**Title: Immune Signatures of SARS-CoV-2 Infection Resolution in Human Lung Tissues**

**Short Title: SARS-CoV-2 infection resolution in human lung tissues**

**One sentence summary:** We comprehensively mapped the human immunological dynamics

associated with successful resolution of SARS-CoV-2 infection in human lung tissues.

Devin Kenney[1,2], Aoife K. O'Connell[1,2,4], Anna E. Tseng[1,2,4], Jacquelyn Turcinovic[1,2,3], Maegan L.

Sheehan[7,12], Adam D. Nitido[7,12], Paige Montanaro[2,4], Hans P. Gertje[2,4], Maria Ericsson[6], John H.

Connor[1,2], Vladimir Vrbanac[7], Nicholas A. Crossland[1,2,4], Christelle Harly[8,9,12]*, Alejandro B.

Balazs[7,12]*, and Florian Douam[1,2,12,13] *

[1]Department of Virology, Immunology, and Microbiology, Boston University Chobanian &

Avedisian School of Medicine, Boston, MA, USA.

[2] National Emerging Infectious Diseases Laboratories, Boston University, Boston, MA, USA.

[3] Bioinformatics Program, Boston University, Boston, MA, USA.

[4] Department of Pathology and Laboratory Medicine, Boston University Chobanian & Avedisian

School of Medicine, Boston, MA, USA.

[6] Electron Microscopy Core Facility, Harvard Medical School, Boston, MA, USA.

[7] Ragon Institute of MGH, MIT and Harvard, Cambridge, MA, USA.

[8] Université de Nantes, INSERM, CNRS, CRCINA, Nantes, France

[9] LabEx IGO 'Immunotherapy, Graft, Oncology', Nantes, France.

[12] These authors contributed equally to the work.

[13] Lead contact

*Correspondence:

Florian Douam, National Emerging Infectious Diseases Laboratories, 620 Albany Street Boston

MA 02118 USA, +1-617-358-9174, fdouam@bu.edu.

Alejandro B. Balazs, Ragon Institute of MGH Harvard and MIT, 400 Technology Square

Cambridge MA 02139 USA, +1-857-268-7197, abalazs@mgh.harvard.edu.

Christelle Harly, CRCINA - Institut de Recherche en Santé de Nantes Université, 8 quai

Moncousu 44 007 Nantes France, +33-2 28 08 02 08, christelle.harly@inserm.fr.

The authors have declared that no conflict of interest exists.

**ABSTRACT**

While human autopsy samples have provided insights into pulmonary immune mechanisms associated with severe viral respiratory diseases, the mechanisms that contribute to a clinically favorable resolution of viral respiratory infections remain unclear due to the lack of proper experimental systems. Using mice co-engrafted with a genetically matched human immune system and fetal lung xenograft (fLX), we mapped the immunological events defining successful resolution of SARS-CoV-2 infection in human lung tissues. Viral infection is rapidly cleared from fLX following a peak of viral replication, histopathological manifestations of lung disease and loss of AT2 program, as reported in human COVID-19 patients. Infection resolution is associated with the activation of a limited number of hematopoietic subsets, including inflammatory monocytes and non-canonical double-negative T-cells with cytotoxic functions, which are highly enriched in viral RNA and dissipate upon infection resolution. Activation of specific human fibroblast and endothelial subsets also elicit robust antiviral and monocyte chemotaxis signatures, respectively. Notably, systemic depletion of human CD4+ cells, but not CD3+ cells, abrogates infection resolution in fLX and induces persistent infection, supporting evidence that peripheral CD4+ monocytes are important contributors to SARS-CoV-2 infection resolution in lung tissues. Collectively, our findings unravel a comprehensive picture of the immunological events defining effective resolution of SARS-CoV-2 infection in human lung tissues, revealing markedly divergent immunological trajectories between resolving and fatal COVID-19 cases.

**INTRODUCTION**

Coronavirus disease 2019 (COVID-19) is a respiratory disease that has swept the world since its emergence in the Wuhan province of China in late 2019. The etiologic agent of COVID-19, the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is a plus-sense, enveloped RNA virus that targets the epithelium of the respiratory tract. Infection results in varying severities of COVID-19, with most cases being mild to asymptomatic. The onset of severe disease is associated with aberrant immune responses (e.g., excessive pulmonary infiltration of myeloid cells, inflammasome-activated monocytic cells, macrophage exacerbated inflammation) and severe lung injury (e.g., lung consolidation, diffuse alveolar damage (DAD), and thrombosis) (*1-9*).

A large number of human studies have been instrumental in unraveling cellular and molecular processes driving severe COVID-19 disease in infected tissues, particularly in the respiratory tract, using autopsy samples (*10-13*). In parallel, human studies of resolving COVID-19, including controlled human challenge studies, have leveraged peripheral blood and nasopharyngeal samples to identify human signatures of effective infection resolution and mild disease (*14-16*). Notably, this includes individuals with specific HLA haplotypes (*15*), evidence of previous coronavirus exposure (*15, 17, 18*), rapid nasopharyngeal immune infiltration (*14*), and non-productive infection of nasopharyngeal T-cells and macrophages (*14*). However, tissue-specific human immunological processes associated with protection, such as extravasation of recruited immune lineages and their differentiation processes, have remained elusive due to the ethical considerations associated with tissue sampling of individuals with mild disease and human challenge models. Although large and small animal models of COVID-19 are available, the high-cost and limited reagent availability associated with non-human primate models, and the large divergence between rodent and human immune systems (*19, 20*) further underscore the need for additional models capable of recapitulating human protective immune responses to SARS-CoV-2 infection.

Mice engrafted with human fetal lung xenograft (fLX) support infection by multiple human

respiratory viruses, including human cytomegalovirus (HCMV), Middle East respiratory syndrome

coronavirus (MERS-CoV) and SARS-CoV-2 (*21-23*). Upon co-engraftment with a human immune

system (HIS), these animals also mount lung-resident human immune responses against these

pathogens (*21, 23-25*). Recently, our group reported that mice engrafted with fLX are highly

susceptible to SARS-CoV-2 infection and lung tissue damage and support persistent viral

infection (*23*). However, co-engraftment of fLX and HIS in a xenorecipient strain supporting

enhanced myelopoiesis (i.e., HNFL mouse model) rapidly blunted viral infection and prevented

widespread acute viral replication across fLX, resulting in protection from histopathology of fLX

(*23*). Our findings unraveled a human macrophage antiviral signature as a correlate of such rapid

protection against SARS-CoV-2 infection. However, how the human immune system mobilizes

and resolves infection upon extensive viral replication within human lung tissues, a context that

more likely describes mild cases of SARS-CoV-2 infection, has remained elusive.

In this study, we leverage previously described immunodeficient mice engrafted with a

human fetal lung xenograft (fLX) and a genetically matched human immune system, fetal liver

and thymus (BLT-L mice) (*21*) to conduct the first comprehensive mapping of human

immunological correlates of resolution of acute SARS-CoV-2 infection in human lung tissues.

Consistent with previous reports (*24, 25*), BLT-L mice are permissive to SARS-CoV-2 infection

following direct viral inoculation into the fLX.  Infection swiftly resolves by 6 days post-infection

(dpi) following an early viral replication peak at 2 dpi. Acute viral infection is associated with lung

histopathological damage and loss of AT2 program, as described in COVID-19 patients (*1*).

Notably, acute infection is also defined by the emergence of a limited set of hyper-activated

hematopoietic subsets, including inflammatory monocytes and T-cells expressing genes

canonically found in macrophages. Of these, inflammatory monocytes appear to mount the most

robust antiviral responses and are highly enriched in viral RNA before phasing out from the tissues

after resolution. No inflammasome activation in these cells was observed despite enrichment in

viral RNA. This contrasts with reports from fatal COVID-19 cases (*26*) and evidence suggesting

that monocyte and macrophage infections in vitro and in vivo are associated with inflammasome

activation (*27, 28*).

133            Specific fibroblast and endothelial cell subsets also contribute to antiviral responses and

hematopoietic chemotaxis, respectively, underscoring how crosstalk between hematopoietic and

non-hematopoietic lineages likely contributes to infection resolution. At 12 dpi, the immune

landscape in fLX is characterized by an increase in CD4+ patrolling monocytes, conventional

dendritic cells and CD206+ interstitial macrophages (IM). Notably, systemic depletion of human

CD4+ cells, but not human CD3+ cells, abrogates SARS-CoV-2 clearance in fLX and causes

persistent infection. These findings underscore CD4+ circulating monocyte infiltration as a likely

feature that defines SARS-CoV-2 infection resolution.

141            Collectively, our work sheds light on a unique set of immunological events associated with

SARS-CoV-2 resolution in human lung tissues which dramatically contrasts with the immune

trajectories reported in fatal COVID-19 cases. This work opens avenues for mechanistically

dissecting the respective contributions of the hematopoietic and non-hematopoietic subsets

identified in this study, which could inform the development of immunotherapies against viral

respiratory diseases.

**RESULTS**

**BLT-L mice effectively clear SARS-CoV-2 infection following acute viral replication**

150            Previous work from our laboratory (*23*) and others (*21, 24, 25*) have shown that

immunodeficient mice can successfully be engrafted with human fetal lung xenograft (fLX) alone

or in combination with a human immune system (HIS). In this study, we leveraged a previously

reported mouse model co-engrafted with fetal liver and thymus as well as with human

hematopoietic stem cells (HSC) and fLX (BLT-L mice) (*21*). Fetal liver and thymus were engrafted

under the renal capsule of adult NOD.Cg-*Prkdc^{scid}^ Il2rg^{tm1Wjl}^*/SzJ (NSG) mice (12-16 weeks old)

prior to intravenous HSC injection. A piece of fetal lung tissue was subcutaneously engrafted on

the flank of each animal, as described previously (*21*) (**Fig. 1A**). To determine the susceptibility

of BLT-L mice to SARS-CoV-2 infection and their ability to effectively clear infection, BLT-L mice

were inoculated with SARS-CoV-2 (2019-nCoV/USA_WA1/2020) via intra-graft injection. We

used a viral dose ($10^6$ PFU) that we previously established to drive robust and persistent infection

in fLX of immunodeficient mice not engrafted with a HIS (*23*) (**Fig. 1B**).

162           Throughout the course of infection, mice did not display any weight loss (**Fig. 1C)** or

clinical signs of disease such as lethargy or lack of responsiveness (data not shown). To assess

lung histopathology and viral titers longitudinally, fLX were collected at 2, 6, and 12 dpi. Plaque

assay was performed on fLX homogenates to determine viral titers. A significant amount of

infectious viral particles could be recovered from fLX at 2 dpi (3.20 ± 2.52 log([PFU/mg of tissue]),

but not at 6dpi (0.281 ± 0.688 log[PFU/mg of tissue]) and 12 dpi (0.111 ± 0.293 log[PFU/mg of

tissue]) (**Fig. 1D**). These data demonstrated a peak of viral infection at 2 dpi, prior to resolution of

infection by 6 dpi. Immunohistochemistry (IHC) for SARS-CoV-2 Spike (S) protein revealed

infection was mainly found in the alveolar epithelium of fLX at 2 dpi (**Fig. 1E,F**). Consistent with

viral quantification, most viral antigen was cleared by 6 dpi and became undetectable by 12 dpi

(**Fig.1G,H**). SARS-CoV-2 S was primarily detected in the alveolar and bronchiole epithelium along

with necrotic cellular debris, consistent with the primary cell targets of SARS-CoV-2 (**Fig.1E**).

These findings were confirmed through IHC for SARS-CoV-2 nucleoprotein (N) at 2, 6 and 12 dpi

(**Fig. 1I-L**) and transmission electron microscopy (TEM) imaging of fLX at 2 dpi (**Fig. S1A-C**).

Notably, TEM substantiated evidence of productive fLX infection, as indicated by the presence of

viral particles in the cytosol of epithelial cells and budding/invagination of virions (**Fig. S1A-C**).

178           Next, we examined histopathological phenotypes associated with active and resolved

infection. Interpretation of hematoxylin and eosin (H&E) staining illustrated denuding of

pneumocytes, neutrophil infiltration, edema, hemorrhage, thrombosis, and pneumocyte necrosis,

which correlated with sites of infection at 2 dpi (**Fig. 2A-E**). No major signs of histopathological

lung damage were observed at 6 and 12 dpi compared to naïve fLX, indicating that fLX can mount

repair mechanisms upon resolution of infection (**Fig. 2F-G**). A previously described semi-

quantitative histopathological scoring system (*23*) provided statistical confirmation for a significant

increase in lung pathology at 2 dpi, which was no longer apparent at 6 (**Fig. 1H**) or 12 dpi. Of

note, minor lung pathology was observed at baseline (naïve), likely reflecting limited graft vs. host

disease. Together, infection of fLX of BLT-L mice recapitulates many important hallmarks of acute

SARS-CoV-2 infection, including viral replication and histopathological manifestations of disease,

prior to effective viral clearance and lung tissue repair.

**Humoral responses do not drive SARS-CoV-2 clearance in BLT-L mice despite evidence of**

**Spike selective pressure.**

We first asked whether SARS-CoV-2 infection resolution was driven by human

neutralizing humoral responses. Consistent with the rapid clearance of infectious viral particles

by 6 dpi and the known caveat that humanized mice mount limited humoral responses (29), there

were no detectable neutralizing antibodies in serum collected at 2 or 12 dpi (**Fig. S2A, B**).

However, interestingly, genomic sequencing of virus isolated from fLX at 2 dpi revealed the

selection of two stable mutations in 75% of fLX (**Fig. S2C**, **Table S1**). Both mutations were located

in the Spike N-terminal domain (NTD): an insertion (216KLRS) and a non-synonymous mutation

(R245H), neither of which were present in the inoculum (**Fig. S2C**, **Table S1**). Interestingly, these

two mutations were found together in 100% of the viral sequences, suggesting potential co-

evolution (**Fig. S2D**). They have also been reported as positively selected in the context of sub-

optimal neutralizing antibody concentration (216KLRS) (*29*) or the context of cross-species

adaptation (216KLRS and R245H) (*30*). Despite lacking humoral responses, these findings reveal

that BLT-L mice can recapitulate host-pathogen interactions that drive SARS-CoV-2 evolution

and host adaptation.

**SARS-CoV-2 infection remodels the human lung cellular environment.**

To comprehensively map the fLX responses associated with resolution of SARS-CoV-2 infection, we performed single-cell RNA sequencing (scRNA-seq) on fLX from naïve BLT-L mice and at 2 and 12 dpi. While most cells detected in fLX by scRNA-seq were human, a minor population of mouse cells was detected, which were excluded from downstream analysis (**Fig. S3A-C**). Initial analysis of the human compartment revealed diverse hematopoietic (T-cells/innate lymphoid cells (ILC), B cells, myeloid cells and mast cells) and non-hematopoietic lineages (ciliated and non-ciliated epithelial, endothelial, mesenchymal cells and chondrocytes) in both naïve and infected fLX (**Fig. 3A-C and Fig. S3D,E**).

T-cells and innate lymphoid cells (ILC) have the same transcriptional programs, express TCR genes, and play similar effector functions (*31*). Their shared features render them very challenging to distinguish by scRNAseq in small datasets and without TCR sequencing information (*32*). We thus considered them together as part of a T-cell/ILC population. T-cell/ILC frequency dramatically decreased in fLX upon infection (naïve, 66.8%; 2 dpi, 35.3%; 12 dpi, 21.8%), which is consistent with evidence that COVID-19 can induce lymphoid depletion(*33, 34*) (**Fig. 3B,C**). Notably, lymphopenia was not observed in the peripheral blood of BLT-L mice, suggesting lymphocyte depletion in fLX was not directly attributed to declining circulating peripheral lymphocytes (**Fig. S3F**). In contrast, myeloid (naïve, 9.94%; 2 dpi, 25.3%; 12 dpi, 22.2%) and B-cell subsets (naïve, 3.64%; 2 dpi, 5.38%; 12 dpi, 5.31%) relatively expanded upon infection (**Fig. 3B,C**). The epithelial, endothelial and mesenchymal cell frequencies increased upon infection, while mast cell, ciliated cell and chondrocyte frequencies decreased (**Fig. 3B,C**). Notably, most human clusters showed temporal segregation between naïve fLX and fLX at 2 and 12 dpi (**Fig. 3B and Fig. S3E**; naïve: black subclusters; 2 dpi: red subclusters and 12 dpi: blue subclusters). This suggests that infection alters the transcriptional state of many cell types and/or drives the emergence of novel cell subsets. Notably, acute infection at 2 dpi led to the emergence of distinct T, myeloid, and mesenchymal cell populations, labeled as subclusters A, B, and C,

respectively (**Fig. 3B**). Resolved infection (12 dpi) was associated with the emergence of

transcriptomically distinct epithelial, endothelial, mesenchymal, B-cell and myeloid subclusters;

while we did not observe the emergence of transcriptomically divergent subclusters within the T-

cell lineages at that time point.  Temporal separation of several human lineages suggests a two-

step tissue remodeling process during infection involving i) an initial antiviral phase mediated by

a limited set of human subpopulations and ii) a tissue repair phase involving a broader range of

human subpopulations.

Next, we wished to utilize scRNAseq to determine the cellular compartments enriched in

viral RNA. We found that viral RNA was mainly within three major lineages: mesenchymal, T-

cell/ILCs and myeloid (**Fig.4A,B**). AT2 and the overall epithelial compartment were not

significantly enriched in viral RNA despite histological evidence of epithelial infection in fLX at 2

dpi (**Fig. 4A and 1F,J**). Previous evidence that SARS-CoV-2 induces significant cytopathic

damage in AT2 cells (*35*) may suggest that infected AT2 cells are lost when undergoing the

scRNAseq pipeline. Within the mesenchymal cluster, viral reads were limited and distributed

sporadically without significant enrichment within a given subcluster (**Fig. 4B,C**). However, viral

RNA was strongly enriched within the 2 dpi-specific populations in the myeloid and T-cell clusters,

previously labeled as clusters A and B (**Fig. 4A,B**), warranting further investigations.

**Human lung epithelium signatures upon SARS-CoV-2 infection recapitulate features of**

**COVID-19.**

Subclustering of the human epithelial compartment revealed subclusters of airway basal

and secretory cells, alveolar type 1 (AT1) and type 2 (AT2) cells, and serous cells (**Fig. 5A**). The

dynamic changes of the epithelial compartment upon infection were consistent with COVID19

human studies (*2, 5, 36*).  We noted a relative reduction of the AT2 compartment at 2 dpi (25.6%)

compared to naïve mice (73.1%). The AT2 compartment was partially restored at 12 dpi (49.1%)

(**Fig. 5B**). The frequency of basal airway cells followed a reversed trend, indirectly reflecting the

disruption of the AT2 compartment. The size of the AT1 compartment showed a relative increase

at 2 dpi (9.89% compared to 8.00% in naïve fLX), consistently with the ability of AT2 cells to

differentiate into AT1 upon lung damage (*37*) before returning to minimal relative levels as the

AT2 compartment re-expanded following infection resolution.

The AT2 compartment displayed the most robust remodeling upon viral infection across

the entire epithelial compartment, with 2 dpi and 12 dpi AT2 subclusters showing distinctive

transcriptomic signatures compared to naïve AT2 cells (**Fig. 5C**). Interestingly, AT2 cells did not

exhibit robust antiviral responses upon acute infection (**Fig. 5C**). However, they displayed gene

signatures previously reported during and after SARS-CoV-2 infection including, downregulation

of MHC genes (*HLA-C, HLA-B, and HLA-DPB1)* and *CCL5* (a cytokine associated with immune

recruitment during respiratory infection) (*6*), upregulation of immunomodulatory genes regulating

inflammation and cell stress (*SFTPA1*, *EGR1*), and elevated expression of genes associated with

mitochondrial dysfunctions and increased oxidative stress responses (i.e., *LARS2*). (**Fig. 5C**).

Activation of tissue repair mechanism (*SNHG8*) was also observed, further underscoring evidence

of physiologically relevant AT2 response to viral infection (**Fig. 5C**). The absence of detectable

viral RNA in AT2 cells was in line with the lack of antiviral responses. Such a lack of infected AT2

cells in our scRNAseq data is likely the reflection of their vulnerability and death during infection,

given histological evidence of epithelial infection (**Fig. 1F, J**) and the marked reduction of the AT2

compartment in fLX upon infection (**Fig. 5B**).

Collectively, the epithelial compartment of fLX of BLT-L mice recapitulates several

previously reported features of COVID-19 in humans, including loss of AT2 program,

downregulation of immune genes and antigen presentation, and activation of tissue repair

mechanisms.

**SARS-CoV-2 infection resolution is associated with the emergence of a viral RNA-enriched**

**double-negative T-cell subcluster with cytotoxic functions.**

We first aimed to characterize the T-cell signatures to SARS-CoV-2 infection resolution,

and with that, the features of our viral RNA-enriched, 2 dpi-specific, T-cell/ILC subcluster A. T-

cell/ILC responses to infection were mainly limited to this specific viral RNA-associated subcluster

(**Fig. 6A-D**), which segregated very distinctively from other CD3+ T-cell lineages, including

canonical CD4+ and CD8+ T-cells (Subcluster 6; **Fig. 6A-C**).  Using doublet discriminators and

assessing viral RNA abundance; we determined that this subcluster was not a doublet and

displayed low levels of *CD4* and *CD8* transcripts, suggesting a double-negative profile, and

elevated levels of mitochondrial transcripts **(Fig. S4A,B)**. Most notably, it was uniquely defined

by the expression of several transcripts known to be enriched in macrophages (*MARCO*, *TIMP1*,

*LYZ*), fibroblasts (*MGP*, *CALD1*, *COL1A*), or both (*A2M*, *IFITM3*) but not in T-cells (**Fig. 6E and**

**Fig. S4C**). This subcluster also exhibited evidence of cytotoxic function through the expression of

*GZMA*, *GZMB*, and *GNLY* transcripts, albeit expression was also detected in CD8+ T-cells and

activated tissue-resident memory T-cells (TRM) (Subclusters 1,3,4, **Fig. 6E**). Notably, a

subpopulation of double-negative T-cells sharing some of these cytotoxic markers has been

previously reported in mouse spleen(*38*). Given the proximity of this subcluster with myeloid

lineages (**Fig. 3A,B**) and evidence of expression of macrophage-enriched transcripts, we referred

to this subcluster as myeloid-like double-negative T-cells (mDNT cells). mDNT cells also exhibited

upregulation of key gene pathways related to COVID-19, SARS-CoV-2 cell-intrinsic immune

responses, cellular cytotoxicity (e.g., degranulation) and cell death, consistently with a role for this

subset to serve as a robust primary responder to viral infection (**Fig. 6D**). Of note, a recent human

challenge study reported that self-resolving SARS-CoV-2 infection is associated with non-

productive infection of human nasopharyngeal T-cells(*14*). In contrast, in lung autopsy samples

of fatal COVID-19 cases, viral RNA was enriched in myeloid cells but was not detected in T-

cells(*39*). Collectively, our findings underscore that the emergence of viral RNA-enriched T cell

populations displaying myeloid-like features in infected lung tissues is associated with effective

SARS-CoV-2 infection resolution.

**Transient antiviral responses by a viral RNA-enriched inflammatory monocyte subset define lung myeloid responses driving SARS-CoV-2 infection resolution.**

Sub-clustering of myeloid lineages unveiled diverse subpopulations, including alveolar and macrophages (AM) and IM, various monocyte subsets, and one conventional DC subset (cDC) (**Fig. 7A and Fig. S5A-C**). Across all time points, monocytes represented the largest sublineage, which was divided into three subgroups: naïve/resting, patrolling intravascular monocytes (PIM), and inflammatory monocytes (iMO). iMO and AM were the only two subclusters increasing in frequencies upon acute infection at the relative expense of cDC and IM (**Fig.7B,C**). An increase in frequencies within these subclusters was associated with the emergence of distinctive iMO and AM cell populations with infection-induced transcriptomic signatures, underscoring a direct response of these subclusters against viral infection. Following infection resolution, most iMO had phased out and frequencies of alveolar macrophages were concomitantly reduced. Conjointly, the relative number of IM, cDCs and PIM increased through the recruitment of cell populations with distinctive transcriptomic identities from naïve and infection-associated cell populations (**Fig. 7B**, blue).

Despite the moderate expansion of AM at 2 dpi, iMO were the most abundant myeloid lineage at 2 dpi (33.5%; **Fig. 7B, C**) and uniquely defined among other monocyte subclusters through elevated expression of *VCAN, S100A8, CD14, CD163* and absence/minimal expression of *CD4, MARCO and CD206 (MRC1)* (**Fig. 7D-F and Fig. S5D**). iMO were also the leading mediators of antiviral responses across all other myeloid subclusters, as exemplified by the robust upregulated expression of interferon-stimulated genes (ISGs) and inflammatory cytokines (*CCL8, CXCL10, CCL2, ISG15, DEFB1, IL1RN, IFIT1, ISG20, IFIT3,* and *MX1*), as well as inflammatory markers such as *CD163* (**Fig. 7E,G**).  While some ISGs (e.g., *ISG15, IFIT3, and MX1*) were expressed in other myeloid lineages at 2 dpi, their expression was markedly lower compared to iMO. Notably, upregulation of *CCL8*, *CXCL11* and *DEFB1* transcripts were the most exclusive to

iMO (**Fig. 7G**). iMO were also highly enriched in viral RNA, and corresponded to the previously

referred 2 dpi-specific viral RNA-enriched subcluster B (**Fig. 3B, 4B, and 7A,B,H**), which also

suggested an association between enrichment in viral RNA and potentiated antiviral responses.

Concomitantly, we examined whether viral RNA-associated iMOs (CoV-iMOs) displayed a

specific transcriptomic signature compared to iMOs (noCoV-iMOs) with undetectable viral RNA.

Notably, only two genes correlated with the presence of viral RNA in iMOs: *FGL2 and C15orf48*

(**Fig. 7I**). While *FGL2* was down-regulated in CoV-iMO compared to noCoV-iMOs, *C15orf48* was

upregulated (**Fig. 7I**). Soluble FGL2 exerts immunosuppressive functions (notably by inhibiting

the NF-kB pathway) (*40*). Conversely, the mitochondrial protein C15orf48 is positively regulated

by NF-kB signaling (*41*) and has previously been implicated in severe COVID-19, acting as a

positive regulator of inflammation (*42*). A recurring feature of lung monocytes in fatal cases of

COVID-19 is the expression of *IL-1β* (*1, 2, 6-8*), and inflammasome activation has been

associated with the non-productive infection of monocytes(*27*). However, no *IL-1β* expression or

inflammasome activation was detected in CoV-iMOs despite enrichment in viral RNA.

In contrast to iMO, *CD4+ CD163- CD206(MRC1)*-PIM were detectable at all time points

(**Sub-cluster 0**; **Fig. 7 A-C**). 2 dpi-specific PIM harbored a distinctive, intermediate transcriptomic

signature bridging naïve PIM and 2 dpi-specific iMO, and that was defined by the upregulation of

specific interferon-stimulated genes (ISGs) such as *IFI27 and XAF1*, and pro-inflammatory genes

(*S100A11*) (**Fig. 7J**). At 12 dpi, PIM was the dominant subset over other monocyte subsets

(50.3%; **Fig. 7C**) and displayed a unique transcriptomic profile associated with anti-inflammation,

tissue repair and cellular debris clearance mechanisms, notably through the upregulation of

*FGL2, DYNLL1* and *CX3CR1*. At 12 dpi, expanded cDC and IM populations also elicited

comparable transcriptomic signatures.

mIHC analysis supported our scRNA seq findings. We observed a significant increase in

CD4- CD3- CD163+ cells in fLX in the extravascular interstitium and alveolar spaces, mirroring

the 2 dpi-specific expansion of CD4- CD163+ iMO and alveolar macrophages uncovered through

scRNA seq (**Fig. 7K**). CD3- CD4+ CD163- at that time point could be associated with the 2 dpi-specific PIM population infiltrating the infected fLX. mIHC also recapitulated the reduction of CD3+ T-cells (**Fig. 7K**) at 2 dpi. At 12 dpi, fLX were still enriched in CD4+ cells, likely reflecting PIM infiltration (**Fig. 7K**). CD163+ cells also persisted in fLX at 12 dpi, which can be explained by the combined expansion of IM (CD4+ CD163+) and presence of AM (CD4- CD163+) at that time point as CD163+ iMO phase out (**Fig. 7K**).

Collectively, our findings underscore a coordinated myeloid mobilization to infection resolution and tissue repair, with iMO concentrating viral materials and robust antiviral responses. As the viral materials are resolved, iMO populations dissipate, opening niches for other myeloid lineages to engage in a coordinated tissue repair process.

**Mesenchymal and endothelial signatures of infection resolution**

We then examined the contribution of the mesenchymal and endothelial compartment in mediating effective infection resolution. Relative expansion of the mesenchymal compartment upon infection was mainly driven by an increase in fibroblast populations (**Fig. 3C; Fig. 8A,B**). An increase in alveolar fibroblasts was observed at 12 dpi (49.6%; naïve=28.1%), which was preceded by the emergence of a 2 dpi-specific cluster of activated fibroblasts (26.5%; naïve=8.9%) (**Fig. 8A-C**), consistent with effective tissue repair in response to lung damage. Activated fibroblasts were the dominant mesenchymal population in mediating antiviral responses (**Fig. 8D**), as displayed by robust upregulation of major pro-inflammatory mediators such as *CXCL10* and *TIMP1* (**Fig. 8F**).

The endothelial compartment is an important modulator of immune recruitment through cytokine/chemokine signaling. Infected fLX showed the emergence of an activated, 2 dpi-specific endothelial cell cluster (**Fig. 8G,H**), which was identified by increased expression of several transcripts involved in myeloid chemotaxis (**Fig. 8I**), including *CXCR4* and *MCAM* (*43, 44*). Notably, venous endothelial cells also displayed upregulation of a panel of interferon-stimulated

gene transcripts (*XAF1, IFIT1, MX1, IFI44L, RSAD2)* at 2 dpi (**Fig. 8J**) and the downregulation of

several transcripts coding for activation markers (*CD69*), MHC genes (*HLA-DPB1, HLA-C*) and

major proteins regulating cellular metabolism and transcription (*HSP90AA1, RPL13A*), which was

reflective of endothelial dysfunction and stress (**Fig. 8J**) consistently with human patient reports

and animal studies (*45-48*). Collectively, our findings support the contribution of the mesenchymal

and endothelium compartments in driving antiviral responses and myeloid chemotaxis,

respectively, in driving infection resolution.

**Systemic depletion of CD4+ cells abrogates viral clearance in fLX.**

Many of our findings underscore a robust association between monocyte recruitment into

fLX and SARS-CoV-2 infection resolution. This includes: 1) the recruitment of CD4+ PIM into

infected fLX, 2) the monocyte nature of iMO and of their dominant antiviral responses, 3) the high

enrichment in viral RNA of iMO during infection and 4) the endothelial-mediated myeloid

chemotaxis signature at 2 dpi. These pieces of evidence also complement recent human findings

which associate effective control of SARS-CoV-2 infection in the nasopharynx with monocyte

recruitment(*14*). Antibody-mediated depletions are commonly used to deplete specific

hematopoietic subsets, and they are particularly amenable to HIS mouse model studies; further

emphasizing the power of these *in vivo* platforms to mechanistically dissect immunological

mechanisms in a human context. However, no anti-human CD14 antibodies have been well

characterized for *in vivo* depletion of human monocytes. In contrast, anti-human CD4 antibodies

have been. To experimentally validate the importance of the recruitment of circulating monocytes

to drive infection resolution in fLX, we therefore performed systemic depletion of CD3+, CD4+,

and CD8+ cells through the administration of OKT3, OKT4, or OKT8 depleting antibodies,

respectively, via intraperitoneal injection of BLTL mice both prior to and after infection (**Fig. 9A**).

Human monocytes, as well as some macrophages and dendritic cells (DC) express CD4 (*49, 50*),

and the use of these three depleting antibodies will allow us to deconvolute the distinctive impact

of T-cells and monocytes in driving infection resolution.

Flow cytometry confirmed effective systemic depletion in the blood (**Fig. S6A**). All animals

were euthanized at 12 dpi to assess for the persistence of SARS-CoV-2 infection in fLX. While

depletion of CD3+, CD4+, and CD8+ cells induced persistent infection in some or most of the

animals, only CD4+ cell depletion (mean Log PFU/mg tissue = 2.37) resulted in a statistically

significant defect in infection resolution compared to all other experimental conditions (isotype,

OKT3, and OKT8-treated mice) (**Fig. 9B**). The rate of productively infected fLX was also higher

in CD4+ cell-depleted animals, with 72% of fLX (8/11) showing infection at 12 dpi compared to

50% (4/8) and 33% (2/6) in CD3- and CD8-depleted mice, respectively (**Fig. 9B**). Additionally,

when examining the average viral titer (Log PFU/mg of tissue) of persistently infected fLX, the

ones from CD4+ cell-depleted animals were significantly higher compared to CD3+ cell- depleted

fLX (**Fig. S6B**), furthering that CD4+ cells have a more consequential impact on driving infection

resolution than CD3+ cells. Anti-SARS-CoV-2 N IHC confirmed these findings and superior defect

of CD4+ cell-depleted animals to resolve infection (**Fig. 9C**). Using multiplex fluorescent IHC

(mIHC), we also validated the reduction in CD3+, CD4+ and CD3+ CD4- cells in fLX from CD3-,

CD4+ and CD8+ cell-depleted animals respectively, compared to isotype-treated animals

(**Fig.9D**).  In the fLX of CD4+ cell-depleted mice, CD4+ cells depletion also associated with

significant reduction of CD163+ cells at 12 dpi, underscoring the association between CD4+

infiltration into fLX and CD163+ cell recruitment and differentiation (**Fig. 9E,F**).

Persistent infection in CD4+ cell-depleted animals was also associated with significant

downregulation of MHC class I (**Fig. 10**), a phenomenon we similarly observed in acutely infected

fLX (2 dpi) (**Fig. 10)** and that has been previously reported in cells with active SARS-CoV-2

replication (*51, 52*). This further emphasizes that depletion of CD4+ cells is associated with

defective viral clearance mechanisms. These findings suggest that circulating CD4-expressing

cells significantly mediate SARS-CoV-2 infection resolution in BLT-L mice.

**Defective monocyte recruitment is associated with systemic and local signatures of chronic infection.**

To further interrogate the impact of CD4+ depletion and monocyte recruitment on SARS-CoV-2 infection, we investigated fLX antiviral responses during acute and persistent infection by quantifying the concentration of 32 cytokines in the peripheral blood of naïve, acutely infected, persistently infected or recovered BLTL mice.  Systemic levels of human CCL2 and CCL3, major monocyte attractants, were elevated in CD4+ cell-depleted mice and comparable to those of acutely infected mice (**Fig. 11A,B**). The maintenance of myeloid chemotaxis signals in persistently infected fLX further underlines the contribution of myeloid recruitment in SARS-CoV-2 infection resolution. Notably, among all cytokines and chemokines analyzed, CXCL10 was the only one displaying significantly increased serum levels in acutely infected mice (2 dpi) prior to returning to undetectable levels upon infection resolution (12 dpi) (**Fig 11C**).  Several human subsets within fLX express *CXCL10* upon acute infection, including myeloid, mesenchymal and endothelial subsets (**Fig. 11D,E**). However, the myeloid compartment was the major source of *CXCL10* among all human clusters at 2 dpi (**Fig. 11D,E**) and this phenotype was dominantly driven by iMO (**Fig. 7G and 11E**). Therefore, our findings suggest an association between circulating human CXCL10 during acute infection and the simultaneous differentiation of iMO. Notably, levels of circulating human CXCL10 in persistently infected CD4+ cell-depleted mice were not statistically different than those of mice that resolved infection (12 dpi) (**Fig. 11C**). These results emphasize a link between persistent infection and lack of effective CXCL10 responses, further strengthening a connection between monocyte recruitment, iMO antiviral responses and SARS-CoV-2 infection resolution.

Persistently-infected mice antiviral responses were also distinguishable from acutely infected (2 dpi) and isotype-treated (12 dpi) mice by elevated levels of CCL19 in serum (**Fig. 11F**). CCL19 is a pro-inflammatory cytokine that has been linked with persistent viral replication and

inflammation such as in the context of HIV-1 infection (*53*), underlining that persistent SARS-CoV-

2 infection in BLT-L mice recapitulate key immunological features of chronic viral infection.

We have previously reported that fLX exhibit significant histopathological manifestations

of disease upon SARS-CoV-2 infection in the absence of an engrafted human immune system

(*3*), highlighting that tissue damage in fLX is virally induced. However, persistent infection did not

result in any significant histopathological manifestations of disease in fLX, which strongly

contrasts with acutely infected fLX (**Fig. 2H, Fig. 11G**). As minimal tissue damage is a hallmark

of chronic viral infection (*54*), these findings further support evidence that CD4+ cell depletion

promotes tissue remodeling processes underlying chronic infection and lasting viral persistence.

**DISCUSSION**

As our appreciation of the immunological differences between mice and humans continues

to grow, humanized mouse models increasingly stand out as robust platforms to understand how

viral pathogens interact with human tissues and the human immune system. These models are

especially valuable when investigating tissue and mucosal immunity since such investigations

remain impractical in human patients.

The SARS-CoV-2 pandemic has emphasized the need to increase our understanding of

immune mechanisms that can drive protection against immunologically novel respiratory viruses.

Mice engrafted with human immune systems and human lung tissues have emerged as valuable

tools for such investigations (*21, 23-25*), bridging the limitations of conventional animal models

and the challenges associated with human studies. We previously reported using the HNFL

mouse model that such models can be leveraged to capture immunological signatures defining

effective control of SARS-CoV-2 infection (*23*). However, the enhanced myeloid reconstitution of

the HNFL model rapidly inhibits viral replication in fLX, precluding our ability to study protective

naive immunological mechanisms at play upon acute and potentially symptomatic infection.  This

study aimed to uncover novel facets of these mechanisms using a humanized mouse model that

exhibits robust susceptibility to acute SARS-CoV-2 infection prior resolving infection in a

hematopoietic-dependent manner.

We found that BTL-L mice are able to effectively clear infectious viral particles following

an early peak of viral replication in fLX.  Infection resolution was associated with rapid mobilization

of the human immune cells into fLX upon viral inoculation, which aligns with previous evidence

that immunodeficient mice only engrafted with fLX are unable to clear infection (*22, 23*). Using

scRNAseq analysis, we identified a comprehensive network of novel factors involved in the

resolution of SARS-CoV-2 infection (**Fig. 12**). Our study identified three major hallmarks of this

process: 1) The recruitment of iMO, which display high level of enrichment in viral RNA and mount

dominant antiviral responses across all myeloid lineages, notably defined by CXCL10 expression;

2) The differentiation of mDNT cells, which are highly enriched in viral RNA and exhibit non-

canonical macrophage features and cytotoxic signatures; And 3) The synergistic contribution of

endothelial and mesenchymal cells in infection resolution via potentiating antiviral responses and

myeloid chemotaxis, respectively. Consistent with monocyte infiltration into infected fLX being a

dominant feature of our infection resolution model, systemic depletion of CD4+ cells, but not T-

cells, abrogated viral clearance. Persistent infection is associated with the lack of systemic

CXCL10 responses, dominantly mediated by iMO, as well as with signatures of chronic infection,

including systemic CCL19 expression and lack of fLX damage.

Our findings empower a recent human study reporting that protection from SARS-CoV-2

infection is associated with a rapid monocyte response in the nasal cavity and a decreased

number of circulating monocytes(*14*). Specifically, our work extends the findings of this human

study beyond the limitations of the human model by exploring tissue-resident events, unraveling

the identity of monocyte populations that extravasate tissues, differentiate and mount tissue-

resident immune responses to clear infection. By providing enhanced resolution on key protective

immunological processes and mediators that human studies alone cannot capture, our works

underscores how human and humanized mouse studies can effectively complement themselves

for improving our understanding of human antiviral immunity.

To the best of our knowledge, this study represents the initial evidence of a direct role

played by human lung extravascular inflammatory monocytes in the resolution of respiratory viral

infections. Extravascular monocytes have been proposed to serve as immune sentinels through

their position at the interface of the lung capillaries and alveoli (*55, 56*). However, although

previous research reported that these cells can promote T-cell resident memory differentiation

following viral infection (*57*), their direct antiviral functions have not been documented until now.

Four major features characterized these cells: 1) a dominant CD163-expressing population that

emerges during acute infection before dissipating, 2) a major source of CXCL10 expression, 3) a

high enrichment in viral RNA, and 4) the induction of robust antiviral responses.

Several lines of evidence support the monocyte nature of these cells. They display a close

transcriptomic relationship with patrolling monocytes, which have been reported to give rise to

transient, non-classical extravascular (including alveolar) monocytes in the mouse lung (*56*). The

iMO gene expression profile is also similar to those of human FCN1-monocytes recovered from

the broncho-alveolar lavage of COVID-19 patients with acute respiratory disease syndrome (*58*).

While further studies are needed, our data collectively suggest CD4+ PIM infiltrating the infected

fLX may differentiate into iMO to promote robust antiviral responses. Furthermore, our data show

that some iMO are negative for viral RNA, suggesting that such differentiation is independent of

an association with viral materials, although the presence of viral RNA could potentiate antiviral

responses.

The fate of iMO following infection resolution also remains unclear and will have to be

further deciphered. By 12 dpi and infection resolution, fLX display residual iMO and are enriched

in CD4+ PIM and CD163+ CD206+ IM. CD206+ IM are involved in response to wounding and

infection recovery (*56*) and CD163+ CD206+ monocyte-derived IM (*58*) have been shown to

directly derive from extravasating, circulating CD14+ monocytes in another human immune

system-engrafted mouse model (*59*). Therefore, one hypothesis would be that the pool of CD163+

CD206+ IM is derived both from iMO and newly recruited circulating monocytes.

Our findings also reveal that different monocytic fates are likely associated with distinct

clinical outcomes of SARS-CoV-2 infection. While our study emphasizes the protective role of

monocytes in preventing severe COVID-19, monocytic lineages have also been associated with

severe COVID-19(*2, 5, 36, 58*). Excessive monocyte infiltration, macrophage inflammation and

fibrotic response can lead to potentially fatal acute respiratory distress syndrome (ARDS) and

fibrosis despite promoting infection resolution. A suspected driver of excessive myeloid

inflammation is the ability of SARS-CoV-2 to trigger abortive infection of both monocytes and

macrophages(*27, 60-62*) through viral RNA replication and protein production, which results in

inflammasome activation(*27, 28, 63*). Consistently, viral RNA is enriched in inflammatory

monocytes and macrophages in lung autopsy samples from fatal COVID-19 cases (*5*). However,

our findings that viral RNA in iMO is associated with infection resolution and tissue protection

suggest that rapid clearance of viral RNA-enriched inflammatory monocytes, as opposed to their

persistence, is associated with favorable clinical outcomes.

Lung extravascular monocytes deriving from PIM also exhibit a dynamic transition state

and can differentiate into CD206- IM(*56*), primary responders to infection and important drivers of

inflammation. However, no CD206- macrophages were observed in infected fLX at any time point.

The transient presence of viral RNA-enriched iMO in fLX, leading to the absence of iMO

differentiation into CD206- IM, could therefore mitigate the risk of uncontrolled inflammation during

infection resolution. This process could prevent severe tissue damage and excessive

extravasation of circulating monocytes, which would otherwise differentiate into pro-fibrotic

CD163+ CD206+ IM, a major driver of ARDS(*58*).

Understanding the cellular and molecular players defining the fate of iMO upon exposure

to SARS-CoV-2 and how key regulatory crossroads in this subset may result in differential clinical

outcomes is of particular interest. Compounding immune dysregulations, such as elevated

inflammatory baseline and/or epigenetic imprinting related to innate immune training (*64*), may

hinder PIM and/or iMO's ability to adequately regulate their inflammatory responses in a timely

manner upon encountering viral materials or inflammatory clues. This could also favor extended

cell survival (NF-κB is both involved in inflammasome activation and cell survival (*65*)) and

subsequent differentiation into inflammatory macrophages, fostering exacerbated inflammation.

Our study also creates a mandate for probing the ubiquitous nature of iMO antiviral responses

during other respiratory viral infections and investigating more comprehensively the antiviral

functions of this novel subset beyond SARS-CoV-2 infection.

In parallel to ExiMO, we also identified mDNT cells, a transient DNT cell population highly

enriched in viral RNA and only observed upon acute infection, and which exhibits expression of

key macrophage-defining genes and cytotoxic markers. This population displays similarities with

a subpopulation of cytotoxic DNT cells previously reported in mouse spleen(*38*) and with innate-

like T-cells(*66*), which also harbor cytotoxic signatures. mDNT also share transcriptomic

similarities with γδ T-cell subsets previously described as bearing myeloid and cytotoxic

functions(*67, 68*). However, mDNT appear unique through a specific blend of lymphoid, and

conventional macrophage and mesenchymal markers and their short-lived nature (suggested by

their complete absence at 12 dpi). While CD8+ T-cells and macrophages isolated from

nasopharyngeal swabs have been identified to harbor SARS-CoV-2 RNA in human challenge

studies(*14*), mDNT did not meet the canonical transcriptomic signatures of these subsets,

suggesting the existence of tissue-resident events driving specific cellular differentiation

processes upon infection.  Additional investigations are required to better understand the identity,

fate and functions of this cell subset in lung antiviral immunity. However, it is tempting to speculate

that mDNT may uniquely synergize CD8+ cytotoxic T-cell functions with viral particle phagocytosis

to effectively control infection prior to undergoing rapid cell death.

Despite our findings related to mDNT, the contribution of T-cells in SARS-CoV-2 infection

resolution remains elusive in our model. CD3+ cell depletion resulted in a partial abrogation of

infection resolution, with 50% of animals still capable of clearing infection. In addition, even when

viral infection persisted in CD3+ cell-depleted animals, viral titers in fLX remained significantly

lower than in the fLX of persistently infected CD4+ cell-depleted animals. One possible

interpretation is that T-cell depletion delays but does not abolish infection resolution. CD8+ cell-

depleted mice may display delayed viral clearance compared to isotype-treated mice (with

resolution occurring between 6dpi and 12 dpi), and CD3+ T-cell-depleted mice may be undergoing

progressive resolution by 12 dpi until complete clearance by a later time. In contrast, the viral titer

observed in fLX of CD4+ cell-depleted mice show no evidence of ongoing infection resolution.

How CD3+ depletion may impact mDNT differentiation, and how T-cells (including mDNT) and

ExiMO functions may synergize to rapidly resolve infection without extensive inflammation-

induced tissue damage will need further investigation.

There are inherent limitations associated with the BLT-L mouse model. First, direct viral

inoculation into fLX potentially bypasses key immune checkpoints in the upper respiratory tract,

which are not recapitulated in our model. Second, the hematopoietic reconstitution and functions

of the BLT-L mouse model remain imperfect, notably through the underrepresentation of specific

hematopoietic subsets, including dendritic cells and granulocytes, and limited B-cell responses –

which may ultimately bias the dynamics of viral clearance. Despite these limitations, BLT-L mice

recapitulated human lung responses to SARS-CoV-2 infection through AT2 loss of programming,

fibroblast and endothelial activation, and effective clearance of infection through robust

hematopoietic responses. This mouse model also supported viral adaptation processes with

public health relevance, and recapitulated signatures of persistent viral replication observed in

patients suffering from chronic infection. Our work demonstrates the potential of the BLT-L mouse

model to uncover naïve immune mechanisms and mediators governing the effective resolution of

lung infection by SARS-CoV-2 and open avenues for a comprehensive examination of such

processes against other viral respiratory infections, which may pave the way toward innovative

immunotherapy strategies against these diseases.

**AUTHOR CONTRIBUTIONS**

D.K., A.B.B. and F.D. conceptualized the overall study.  C.H. conceptualized the computational

component of this study and oversaw the analysis of the single-cell RNA sequencing data. D.K.,

A.B.B., N.A.C. and F.D. designed the experiments. D.K., A.K.O., M.S, A.N, H.P.G., M.E., V.V.,

A.B.B. and F.D. performed experiments. D.K., A.K.O., A.T., J.T., M.S, A.N, P.M., M.E., J.H.C.,

V.V., N.A.C., C.H., A.B.B. and F.D. analyzed the data. J.T and C.H. carried out computational

analysis. M.E. carried out electron microscopy analysis. A.B.B. and V.V. provided access to key

resources. D.K. and F.D. wrote the manuscript with contributions from all authors.

**ACKNOWLEDGMENTS**

This work was supported in part by a start-up fund and Peter Paul Career Development

Professorship from Boston University (to F.D.), grants from the National Institutes of Health (K22

AI144050 to F.D.), Clinical and Translational Science Awards (grant UL1 TR001430) from the

National Center for Advancing Translational Sciences of the National Institutes of Health (to F.D.,

and N.A.C.) and an Evergrande MassCPR award (to NEIDL). A.B.B. was supported by

R01AI174875, R01AI174276, DP2DA040254, as well as CDC subcontract 200-2016-91773-

T.O.2. and a Massachusetts Consortium on Pathogenesis Readiness (MassCPR) grant. We also

thank the Evans Center for Interdisciplinary Biomedical Research at Boston University Chobanian

and Avedisian School of Medicine for their support of the Affinity Research Collaborative on

'Respiratory Viruses: A Focus on COVID19'. This work utilized a Ventana Discovery Ultra and

Akoya PhenoImager HT that were purchased with funding from National Institutes of Health SIG

grants (S10 OD026983 & S10OD030269). We thank Ronald B. Corley, NEIDL director at the time

648 of this study and MassCPR award recipient, for his constant encouragement and support for this

649 study. We thank the Boston University Animal Science Center, the Ragon Institute Human

650 Immune System Mouse core, the single cell sequencing core and microarray and sequencing

651 core, and the flow cytometry core at Boston University Chobanian and Avedisian School of

652 Medicine, and all the NEIDL animal core staff for their outstanding support. We also thank all the

653 Douam, Balazs, Connor, and Crossland lab members, NEIDL members, and members of the

654 Department of Virology, Immunology, and Microbiology and Pathology at Boston University for

655 their constant support and advice. D.K. is supported by a T32 training grant in immunology

656 (T32AI007309).

658 **MATERIALS AND METHODS**

659 Detailed descriptions of the materials and methods used for this study are in the Supplemental

660 Materials.

662 ***Institutional approvals.*** All experiments in this study, including those conducted in BSL-3, were

663 approved by an institutional biosafety committee. Animal experiments described in this study were

664 performed in accordance with protocols that were reviewed and approved by the Institutional

665 Animal Care and Use and Committee of the Ragon Institution and Boston University. All mice

666 were maintained in facilities accredited by the Association for the Assessment and Accreditation

667 of Laboratory Animal Care (AAALAC). All replication-competent SARS-CoV-2 experiments were

668 performed in a biosafety level 3 laboratory (BSL-3) at the Boston University National Emerging

669 Infectious Diseases Laboratories (NEIDL).

671 ***Mouse strains and sex as biological variable.*** Female NOD.Cg.-$Prkdc^{Scid}Il2rg^{tm1Wjl}$/SzJ (NSG)

672 mice were obtained from the Jackson Laboratory, catalog number 005557. NSG mice were

673 maintained by the Ragon Institute Human Immune System Mouse core prior to engraftment and

shipment to the NEIDL, Boston University. In our study, only female mice were engrafted with

human fetal tissues because of their ability to support higher levels of engraftment than males.

As we investigated human tissue responses to infection and only leveraged mice as

xenorecipients, the sex of the animals does represent a critical variable for our study.

*Human fetal tissues.* De-identified human fetal tissues were procured from Advanced Bioscience

Resources (Alameda, CA, USA).

*Generation of BLT-L mice.* BLT mice were generated via irradiation of female NOD.Cg.-

*Prkdc$^{Scid}$Il2rg$^{tm1Wjl}$*/SzJ mice (NSG mice; Jackson Laboratory #005557) prior to implantation of

human fetal thymic and fetal liver tissue (Advanced Bioscience Resources) under the murine

kidney capsule. Two pieces of homologous human fetal lung tissue were implanted into

subcutaneous dorsal pockets of mice. Post-implantation, mice intravenously received $1x10^5$

homologous CD34+ cells. Human immune reconstitution was determined by flow cytometry at

weeks post-implantation. Three distinct human donors were used for this study.

*Mouse inoculation with SARS-CoV-2.* BLT-L mice were anesthetized with 1–3% isoflurane prior

to inoculation via subcutaneous, intra-fetal lung xenograft (intra-fLX) injection with $10^6$ plaque

forming unites (PFU) of SARS-CoV-2 WA-1 isolate in 50 µL of sterile 1X PBS. Mice were

euthanized at 2-, 6-, and 12-days post inoculation.

*In vivo antibody depletion.* BLT-L mice were administered 200 mg of anti-CD3e (OKT3)

(BioxCell; cat. # BE0001-2), anti-CD4 (OKT4) (BioxCell; cat. # BE0003-2), anti-CD8 (OKT8)

(BioxCell; cat. # BE0004-2), or isotype IgG2a (Thermofisher; cat # 02-6200) antibody 3-, 2-, and

1-day prior to inoculation and 4- and 8-days post inoculation with $1x10^6$ PFU SARS-CoV-2 WA-

1.

*Quantification and statistical Analysis.* For histopathological score and viral load/titer comparisons a Kruskla-Wallis, non-parametric one-way ANOVA with Benjamini, Krieger, and Yekutieli correction for multiple comparisons was applied given the non-continuous nature of the data. For cytokine data, an Ordinary one-way ANOVA with uncorrected Fishers LSD was used as the data was collected from different time points, treatment conditions, and cohorts. A Kruskla-Wallis, non-parametric one-way ANOVA with an uncorrected Dunn's test was applied for CD163+ Area quantification (AQ) due to the independent comparisons between the samples. All statistical tests and graphical depictions of results were performed using GraphPad Prism version 9.0.1 software (GraphPad Software, La Jolla, CA). For all tests, $p \leq 0.05$ was considered statistically significant. Statistical significance on figures and supplemental figures is labeled with p-values and non-significant values are labeled with n.s. or left unlabeled.

*Data availability*. The Genome Expression Omnibus (GEO) accession number to access the raw data of our scRNAseq analysis is GSE255200.

**REFERENCES**

1. J. C. Melms *et al.*, A molecular single-cell lung atlas of lethal COVID-19. *Nature* **595**, 114-119 (2021).
2. A. F. Rendeiro *et al.*, The spatial landscape of lung pathology during COVID-19 progression. *Nature* **593**, 564-569 (2021).
3. E. Wauters *et al.*, Discriminating mild from critical COVID-19 by innate and adaptive immune single-cell profiling of bronchoalveolar lavages. *Cell Res* **31**, 272-290 (2021).
4. R. A. Grant *et al.*, Circuits between infected macrophages and T cells in SARS-CoV-2 pneumonia. *Nature* **590**, 635-641 (2021).
5. T. M. Delorey *et al.*, COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature*, (2021).
6. M. Liao *et al.*, Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat Med* **26**, 842-844 (2020).
7. X. Ren *et al.*, COVID-19 immune features revealed by a large-scale single-cell transcriptome atlas. *Cell* **184**, 5838 (2021).
8. P. A. Szabo *et al.*, Longitudinal profiling of respiratory and systemic immune responses reveals myeloid cell-driven lung inflammation in severe COVID-19. *Immunity* **54**, 797-814 e796 (2021).

734    9.     S. S. Batah, A. T. Fabro, Pulmonary pathology of ARDS in COVID-19: A pathological
735           review for clinicians. *Respir Med* **176**, 106239 (2021).
736    10.    C. Fan *et al.*, Animal models for COVID-19: advances, gaps and perspectives. *Signal
737           Transduct Target Ther* **7**, 220 (2022).
738    11.    E. Stephenson *et al.*, Single-cell multi-omics analysis of the immune response in COVID-
739           19. *Nat Med* **27**, 904-916 (2021).
740    12.    X. Ren *et al.*, COVID-19 immune features revealed by a large-scale single-cell
741           transcriptome atlas. *Cell* **184**, 1895-1913 e1819 (2021).
742    13.    R. Edahiro *et al.*, Single-cell analyses and host genetics highlight the role of innate
743           immune cells in COVID-19 severity. *Nat Genet* **55**, 753-767 (2023).
744    14.    R. G. H. Lindeboom *et al.*, Human SARS-CoV-2 challenge uncovers local and systemic
745           response dynamics. *Nature*,  (2024).
746    15.    D. G. Augusto *et al.*, A common allele of HLA is associated with asymptomatic SARS-
747           CoV-2 infection. *Nature* **620**, 128-136 (2023).
748    16.    C. G. K. Ziegler *et al.*, Impaired local intrinsic immunity to SARS-CoV-2 infection in severe
749           COVID-19. *Cell* **184**, 4713-4733 e4722 (2021).
750    17.    M. Dugas *et al.*, Less severe course of COVID-19 is associated with elevated levels of
751           antibodies against seasonal human coronaviruses OC43 and HKU1 (HCoV OC43, HCoV
752           HKU1). *Int J Infect Dis* **105**, 304-306 (2021).
753    18.    M. Sagar *et al.*, Recent endemic coronavirus infection is associated with less-severe
754           COVID-19. *J Clin Invest* **131**,  (2021).
755    19.    K. Medetgul-Ernar, M. M. Davis, Standing on the shoulders of mice. *Immunity* **55**, 1343-
756           1353 (2022).
757    20.    A. Rongvaux *et al.*, Human hemato-lymphoid system mice: current use and future
758           potential for medicine. *Annu Rev Immunol* **31**, 635-674 (2013).
759    21.    A. Wahl *et al.*, Precision mouse models with expanded tropism for human pathogens. *Nat
760           Biotechnol* **37**, 1163-1173 (2019).
761    22.    A. Wahl *et al.*, SARS-CoV-2 infection is effectively treated and prevented by EIDD-2801.
762           *Nature* **591**, 451-457 (2021).
763    23.    D. J. Kenney *et al.*, Humanized mice reveal a macrophage-enriched gene signature
764           defining human lung tissue protection during SARS-CoV-2 infection. *Cell Rep* **39**, 110714
765           (2022).
766    24.    Y. Di *et al.*, SARS-CoV-2 Variant-Specific Infectivity and Immune Profiles Are Detectable
767           in a Humanized Lung Mouse Model. *Viruses* **14**,  (2022).
768    25.    R. Sun *et al.*, Humanized mice for investigating SARS-CoV-2 lung infection and
769           associated human immune responses. *Eur J Immunol* **52**, 1640-1647 (2022).
770    26.    L. Baena Carstens *et al.*, Lung Inflammasome Activation in SARS-CoV-2 Post-Mortem
771           Biopsies. *Int J Mol Sci* **23**,  (2022).
772    27.    C. Junqueira *et al.*, FcgammaR-mediated SARS-CoV-2 infection of monocytes activates
773           inflammation. *Nature* **606**, 576-584 (2022).
774    28.    E. Sefik *et al.*, Inflammasome activation in infected macrophages drives COVID-19
775           pathology. *Nature* **606**, 585-593 (2022).
776    29.    PMDA, Xevudy for Intravenous Infusion 500 mg_GlaxoSmithKline K.K._Report on Special
777           Approval for Emergency; https://www.pmda.go.jp/files/000245006.pdf.  (2021).
778    30.    A. Fagre *et al.*, SARS-CoV-2 infection, neuropathogenesis and transmission among deer
779           mice: Implications for spillback to New World rodents. *PLoS Pathog* **17**, e1009585 (2021).
780    31.    C. Harly, M. Cam, J. Kaye, A. Bhandoola, Development and differentiation of early innate
781           lymphoid progenitors. *J Exp Med* **215**, 249-262 (2018).
782    32.    G. Pizzolato *et al.*, Single-cell RNA sequencing unveils the shared and the distinct
783           cytotoxic hallmarks of human TCRVdelta1 and TCRVdelta2 gammadelta T lymphocytes.
784           *Proc Natl Acad Sci U S A* **116**, 11906-11915 (2019).

33. Q. Zhao *et al.*, Lymphopenia is associated with severe coronavirus disease 2019 (COVID-
19) infections: A systemic review and meta-analysis. *Int J Infect Dis* **96**, 131-135 (2020).
34. S. Tavakolpour, T. Rakhshandehroo, E. X. Wei, M. Rashidian, Lymphopenia during the
COVID-19 infection: What it shows and what can be learned. *Immunol Lett* **225**, 31-32
(2020).
35. J. Huang *et al.*, SARS-CoV-2 Infection of Pluripotent Stem Cell-Derived Human Lung
Alveolar Type 2 Cells Elicits a Rapid Epithelial-Intrinsic Inflammatory Response. *Cell stem*
*cell* **27**, 962-+ (2020).
36. E. Wauters *et al.*, Discriminating mild from critical COVID-19 by innate and adaptive
immune single-cell profiling of bronchoalveolar lavages. *Cell Res* **31**, 272-290 (2021).
37. A. Wu, H. Song, Regulation of alveolar type 2 stem/progenitor cells in lung injury and
regeneration. *Acta Biochim Biophys Sin (Shanghai)* **52**, 716-722 (2020).
38. L. Yang *et al.*, Transcriptome landscape of double negative T cells by single-cell RNA
sequencing. *J Autoimmun* **121**, 102653 (2021).
39. T. M. Delorey *et al.*, A single-cell and spatial atlas of autopsy tissues reveals pathology
and cellular targets of SARS-CoV-2. *bioRxiv*, 2021.2002.2025.430130 (2021).
40. L. Fu, Z. Liu, Y. Liu, Fibrinogen-like protein 2 in inflammatory diseases: A future
therapeutic target. *Int Immunopharmacol* **116**, 109799 (2023).
41. Y. Takakura *et al.*, Mitochondrial protein C15ORF48 is a stress-independent inducer of
autophagy that regulates oxidative stress and autoimmunity. *Nat Commun* **15**, 953 (2024).
42. S. A. Clayton *et al.*, Inflammation causes remodeling of mitochondrial cytochrome c
oxidase mediated by the bifunctional gene C15orf48. *Sci Adv* **7**, eabl5182 (2021).
43. S. K. Gupta, P. G. Lysko, K. Pillarisetti, E. Ohlstein, J. M. Stadel, Chemokine receptors in
human endothelial cells. Functional expression of CXCR4 and its transcriptional regulation
by inflammatory cytokines. *J Biol Chem* **273**, 4282-4287 (1998).
44. A. S. Leroyer *et al.*, CD146 (Cluster of Differentiation 146). *Arterioscler Thromb Vasc Biol*
**39**, 1026-1033 (2019).
45. H. M. Otifi, B. K. Adiga, Endothelial Dysfunction in Covid-19 Infection. *Am J Med Sci* **363**,
281-287 (2022).
46. Z. Qin *et al.*, Endothelial cell infection and dysfunction, immune activation in severe
COVID-19. *Theranostics* **11**, 8076-8091 (2021).
47. T. Kadiyska *et al.*, Role of endothelial dysfunction in the severity of COVID-19 infection
(Review). *Molecular medicine reports* **26**, (2022).
48. S. W. Xu, I. Ilyas, J. P. Weng, Endothelial dysfunction in COVID-19: an overview of
evidence, biomarkers, mechanisms and potential therapies. *Acta Pharmacol Sin* **44**, 695-
709 (2023).
49. A. Zhen *et al.*, CD4 ligation on human blood monocytes triggers macrophage
differentiation and enhances HIV infection. *J Virol* **88**, 9934-9946 (2014).
50. L. Jardine *et al.*, Rapid detection of dendritic cell and monocyte disorders using CD4 as a
lineage marker of the human peripheral blood antigen-presenting cell compartment. *Front*
*Immunol* **4**, 495 (2013).
51. N. Arshad *et al.*, SARS-CoV-2 accessory proteins ORF7a and ORF3a use distinct
mechanisms to down-regulate MHC-I surface expression. *Proc Natl Acad Sci U S A* **120**,
e2208525120 (2023).
52. J. S. Yoo *et al.*, SARS-CoV-2 inhibits induction of the MHC class I pathway by targeting
the STAT1-IRF1-NLRC5 axis. *Nat Commun* **12**, 6602 (2021).
53. J. K. Damas *et al.*, Homeostatic chemokines CCL19 and CCL21 promote inflammation in
human immunodeficiency virus-infected patients with ongoing viral replication. *Clin Exp*
*Immunol* **157**, 400-407 (2009).
54. E. I. Zuniga, M. Macal, G. M. Lewis, J. A. Harker, Innate and Adaptive Immune Regulation
During Chronic Viral Infections. *Annu Rev Virol* **2**, 573-597 (2015).

55.    M. P. Rodero *et al.*, Immune surveillance of the lung by migrating tissue monocytes. *Elife*
837           **4**, e07847 (2015).
838    56.    J. Schyns *et al.*, Non-classical tissue monocytes and two functionally distinct populations
839           of interstitial macrophages populate the mouse lung. *Nat Commun* **10**, 3964 (2019).
840    57.    P. R. Dunbar *et al.*, Pulmonary monocytes interact with effector T cells in the lung tissue
841           to drive T(RM) differentiation following viral infection. *Mucosal Immunol* **13**, 161-171
842           (2020).
843    58.    D. Wendisch *et al.*, SARS-CoV-2 infection triggers profibrotic macrophage responses and
844           lung fibrosis. *Cell* **184**, 6243-6261 e6227 (2021).
845    59.    E. Evren *et al.*, Distinct developmental pathways from blood monocytes generate human
846           lung macrophage diversity. *Immunity* **54**, 259-275 e257 (2021).
847    60.    J. Zheng *et al.*, Severe Acute Respiratory Syndrome Coronavirus 2-Induced Immune
848           Activation and Death of Monocyte-Derived Human Macrophages and Dendritic Cells. *J*
849           *Infect Dis* **223**, 785-795 (2021).
850    61.    A. Boumaza *et al.*, Monocytes and Macrophages, Targets of Severe Acute Respiratory
851           Syndrome Coronavirus 2: The Clue for Coronavirus Disease 2019 Immunoparalysis. *J*
852           *Infect Dis* **224**, 395-406 (2021).
853    62.    D. Yang *et al.*, Attenuated Interferon and Proinflammatory Response in SARS-CoV-2-
854           Infected Human Dendritic Cells Is Associated With Viral Antagonism of STAT1
855           Phosphorylation. *J Infect Dis* **222**, 734-745 (2020).
856    63.    D. Munnur *et al.*, Altered ISGylation drives aberrant macrophage-dependent immune
857           responses during SARS-CoV-2 infection. *Nat Immunol* **22**, 1416-1427 (2021).
858    64.    J. Ochando, W. J. M. Mulder, J. C. Madsen, M. G. Netea, R. Duivenvoorden, Trained
859           immunity - basic concepts and contributions to immunopathology. *Nat Rev Nephrol* **19**,
860           23-37 (2023).
861    65.    M. Mussbacher, M. Derler, J. Basilio, J. A. Schmid, NF-kappaB in monocytes and
862           macrophages - an inflammatory master regulator in multitalented immune cells. *Front*
863           *Immunol* **14**, 1134661 (2023).
864    66.    M. Molgora, M. Colonna, Innate-like T cells: A promising asset in anti-cancer immunity.
865           *Cancer Cell* **40**, 714-716 (2022).
866    67.    M. Lebedev, B. Faburay, J. A. Richt, A. Young, Myeloid-like gammadelta T cell subset in
867           the immune response to an experimental Rift Valley fever vaccine in sheep. *Vet Immunol*
868           *Immunopathol* **233**, 110184 (2021).
869    68.    M. Muto, M. Baghdadi, R. Maekawa, H. Wada, K. Seino, Myeloid molecular characteristics
870           of human gammadelta T cells support their acquisition of tumor antigen-presenting
871           capacity. *Cancer Immunol Immunother* **64**, 941-949 (2015).

872
873

877    **FIGURES**

**Figure 1. BLT-L mice effectively resolve SARS-CoV-2 infection. (A)** Representative

schematic of BLT-L mice. **(B)** Fetal lung xenografts (fLX) of BLT-L mice were infected with 1x10$^6$

PFU of SARS-CoV-2 WA-1 isolate via subcutaneous intra-graft injection. **(C)** BLT-L mice were

monitored for weight change over the course of 12 days post-infection (dpi). **(D)** Viral titer

(log(PFU/mg)) in fLX at 2, 6, and 12 dpi as determined by plaque assay. **(E-H)**

Immunohistochemistry for SARS-CoV-2 S and **(I-L)** N protein were performed on naïve fLX **(E,I)**

and fLX at 2 dpi **(F,J)**, 6dpi **(G,K),** and 12 dpi **(H,L)**. Scale bar = 100 μM. Data are representative

of two or three independent experiments.  n = 6 -10 per timepoint. Error bars represent mean ±

standard error of the mean. *One-way ANOVA analysis was performed. p-values are indicated.*

*n.s. = non-significant.*

**Figure 2. BLT-L mice resolve histopathological damage in fLX. (A-E)** Hematoxylin and eosin staining was performed on naïve fLX **(A)** and fLX at 2 dpi **(B-E)**, 6dpi **(F),** and 12 dpi **(G)**. Scale bar = 100 μM. **(C)** Bronchiolar attenuation and denuding (blue arrows) correlates directly with the presence of SARS-CoV-2 Spike protein. **(D)** Alveolar spaces are multifocally filled with necrotic debris admixed with neutrophils and edema (blue hashes). Type II pneumocytes (ATII) are often denuded or attenuated in areas of inflammation that correlates directly with the presence of SARS-CoV-2 Spike protein. Fibrin thrombi are routinely observed in neighboring parenchyma (black asterisks). **(E)** Coagulative necrosis as evidenced by loss of cellular detail and generalized eosinophilia (above the black hashed line) with numerous regional fibrin thrombi (black asterisks). Although SARS-CoV-2 Spike protein is not observed in the area of coagulative necrosis viral antigen is located within adjacent tissue. **(H)** Histopathological scoring of naïve fLX and fLX at 2, 6, and 12 dpi. Data are representative of two or three independent experiments.  n = 6 -10 per timepoint. Error bars represent mean ± standard error of the mean. *One-way ANOVA analysis was performed. p-values are indicated. n.s. = non-significant.*

**Figure 3**. **Cellular remodeling and viral dynamics in fLX upon SARS-CoV-2 infection**

Single cell RNA sequencing analysis was performed on single cell suspensions of naïve fLX and

fLX at 2 and 12 dpi. Naïve n=3 fLX (9,605 cells), 2 dpi n=2 fLX (4,405 cells), and 12 dpi n = 2 fLX

(5,857 cells). **(A)** UMAP plot clustering of the human cell compartment of naïve fLX and fLX at 2

and 12 dpi. **(B)** Temporal annotation of the human clusters on the UMAP plot: naïve (black), 2 dpi

(red), 12 dpi (blue). Sub-clusters unique to 2 dpi are annotated by A (T cell compartment), B

(myeloid compartment) and C (mesenchymal compartment). **(C)** Frequency of each cell

compartment determined by single cell RNA-sequencing.

**Figure 4. Cellular compartmentalization of viral RNA during infection. (A)** Dot plots

displaying the expression of SARS-CoV-2 viral RNA transcripts identified in scRNA-seq. **(B)**

UMAP plot showing the distribution of SARS-CoV-2 viral RNA transcripts by density across all

human cell clusters and time points. **(C)** UMAP plot showing the distribution of SARS-CoV-2 viral

RNA reads across all human cell clusters and time points. Inset showing distribution of viral RNA

reads in the mesenchymal cluster.

**Figure 5. Remodeling of the epithelial compartment of fLX during infection. (A)** UMAP plot

sub-clustering of the human epithelial compartment in fLX across all time points. **(B)** Temporal

annotation of the human epithelial sub-clusters on the UMAP plot: naïve (black), 2 dpi (red), 12

dpi (blue). The inset graph indicates the change in frequency of each compartment (AT2; black,

Basal; pink, Secretory; teal, and AT1; purple) within the epithelial compartment. Red line indicates

the change in the AT2 compartment over time. **(C)** Heatmap displaying the relative expression

(from -1 to 1) of the top differentially regulated genes over the course of infection within the AT2

compartment.

**Figure 6. Myeloid-like double-negative T-cell emerges upon acute infection and are enriched in viral RNA. (A)** UMAP plot showing the sub-clustering of the human T-cell compartment of fLX across all time points. **(B)** Temporal annotation of the human T-cell sub-clusters on the UMAP plot: naïve (black), 2 dpi (red),2 dpi (blue). **(C)** UMAP plot (all time points) showing the distribution of SARS-CoV-2 viral RNA transcripts in the T-cell compartment. **(D)** Go-term analysis showing the most highly upregulated signaling pathways in the mDNT (sub-cluster 6). **(E)** Relative expression of T cells, macrophage, mesenchymal and antiviral markers within each of the T-cell sub-clusters (From 0 to 7; as labeled in panel A).

**Figure 7. CD163+ extravascular inflammatory monocytes arise during acute viral infection and are predominant antiviral mediators. (A)** UMAP plot showing sub-clustering of the human myeloid compartment of fLX across all time points. **(B)** Temporal annotation of the myeloid sub-clusters on the UMAP plot: naïve (black), 2 dpi (red), 12 dpi (blue). iMO sub-cluster is indicated with a blue circle. **(C)** Frequency of each myeloid sub-cluster by timepoint. **(D-F)** UMAP plot showing the expression of *MRC1* **(D)**, CD*163* **(E)**, and *CD4* **(F)** across myeloid sub-sets. **(G)** Relative expression of highly upregulated ISGs and inflammatory markers in each of the myeloid sub-clusters. **(H)** UMAP plot (all time points) showing the distribution of SARS-CoV-2 viral RNA transcripts in the myeloid compartment. **(I)** Violin plots showing expression level of differentially expressed genes between SARS-CoV-2 positive and negative ExiMO (sub-cluster 3). **(J)** Heatmap displaying the relative expression (from -1 to 1) of the top differentially regulated genes over the course of infection within the patrolling monocyte (0) compartment. **(K)** Multiplex fluorescent immunohistochemistry of naïve fLX and fLX at 2 and 12 dpi. CD163: teal, CD4: yellow, CD3: orange, PPARγ: green, Dapi: gray. Two representative images. Scale bar = 100 μM.

**Figure 8. (A)** UMAP plot sub-clustering of the human mesenchymal compartment in fLX across

all time points. **(B)** Temporal annotation of the mesenchymal sub-clusters on the UMAP plot:

naïve (black), 2 dpi (red), 12 dpi (blue). **(C)** Frequency of mesenchymal sub-clusters. **(D)** Go-term

analysis showing cellular pathways linked to significantly upregulated genes within the activated

fibroblasts sub-cluster. **(E-F)** UMAP plot showing *CXCL10* **(E)** and *TIMP1* **(F)** expression within

all human mesenchymal sub-clusters and across all time points. Activated fibroblasts are

indicated by a red circle in A and B. **(G)** UMAP plot sub-clustering of the human endothelial

compartment of fLX across all time points. **(H)** Temporal annotation of the endothelial sub-clusters

on the UMAP plot: naïve (black), 2 dpi (red), 12 dpi (blue). **(I)** List of upregulated (red) and

downregulated (blue) genes specific to the activated endothelial sub-cluster (Cluster 2) over the

other endothelial sub-clusters (*p.adj*≤0.05). **(J)** Heatmap displaying the relative expression (from

-1 to 1) of the top differentially regulated genes over the course of infection within the venous

endothelial compartment.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

**Figure 9. Systemic depletion of CD4+ cells results in persistent infection of fLX. (A)** BLT-L

mice were administered 200 μg of anti-CD3e (OKT3), anti-CD4 (OKT4), anti-CD8 (OKT8) or

IgG2a isotype. **(B)** Viral titer (log(PFU/mg)) in fLX extracted from BLT-L mice treated with isotype,

OKT3, OKT4, or OKT8 antibody at 12 dpi. **(C)** Immunohistochemistry for SARS-CoV-2 N protein

on fLX of depleted or non-depleted BLT-L mice at 12 dpi. **(D)** Multiplex immunohistochemistry on

fLX of depleted or non-depleted BLT-L mice at 12 dpi. Dapi = gray, CD3 = teal, CD4 = red.  Scale

bar = 50 μm. **(E-F)** Multiplex fluorescent immunohistochemistry (E) and CD163+ area

quantification (AQ) (F) of fLX that resolved infection (12 dpi, isotype treated), or eliciting acute (2

dpi) or persistent infection (12 dpi, OKT4 treated). CD163: teal, CD4: yellow, CD3: orange,

PPARγ: green, Dapi: gray. Scale bar = 50 μM.

**Figure 10.** MHC-I staining of fLX tissue sections (anti-MHC class I (EMR8-5) CST 88274)

extracted from naïve mice, OKT4-treated mice (12 dpi), or from infected fLX at 2 and 12 dpi. Scale

bar = 100 μM

**Figure 11. CD4+ cell depletion results in signatures of chronic infection. (A-C)** Cytokine quantification (**A**: CCL2, **B**: CCL3, **C**: CXCL10) in the serum of naïve, infected (2 dpi and 12 dpi isotype-treated or not) and OKT4-treated BLT-L mice (12 dpi). **(D)** Violin plot showing expression level of *CXCL10* per cell and within each human lineage. **(E)** UMAP plot showing *CXCL10* expression within all human lineages in naive fLX and at 2- and 12 dpi. Locations of the mesenchymal, endothelial, myeloid and ExiMO clusters at 2 dpi are indicated. **(F)** Quantification of CCL19 in the serum of naïve, infected (2 dpi and 12 dpi isotype-treated or not) and OKT4-treated BLT-L mice (12 dpi). **(G)** Histopathological scoring of fLX extracted from isotype and OKT4-depleted BLT-L mice (12 dpi). Error bars indicate mean ± Standard error of the mean. One-way ANOVA, t-test. p-values are indicated on graphs.

1048

**Figure 12. Immune signatures of SARS-CoV-2 infection resolution in human lung tissues.**
During effective viral resolution and tissue repair (left panel), myeloid cells are recruited to the site
of infection and differentiates into CD163+ inflammatory monocytes (iMO). iMO produce antiviral
and inflammatory signals and capture viral RNA/particles. Additionally, myeloid-like double
negative T cells (mDNT), act to enhance the capture of viral RNA/particles and antiviral
responses. In parallel, activated endothelial cells and fibroblasts contribute to the recruitment of
myeloid cells and inflammatory response, respectively. Collectively, these events lead to viral
clearance and infection resolution. In the context of persistent infection (left panel), absence of
effective CD4+ monocyte recruitment prevents infection resolution.