HIV-1 Nef synergizes with APOL1-G1 to induce nephrocyte cell death in a *Drosophila*

model of HIV-related kidney diseases

Jun-yi Zhu[1,2], Yulong Fu[1,3], Joyce van de Leemput[1,2] Jing Yu[4], Jinliang Li[5], Patricio E. Ray[4, *],

and Zhe Han[1,2, *]

1 Center for Precision Disease Modeling, Department of Medicine, University of Maryland

School of Medicine, Baltimore, Maryland

2 Division of Endocrinology, Diabetes and Nutrition, Department of Medicine, University of

Maryland School of Medicine, Baltimore, Maryland

3 Current: Department of Pathology, University of Alabama at Birmingham, Birmingham,

Alabama

4 Child Health Research Center, Department of Pediatrics, University of Virginia School of

Medicine

5 Children's Research Institute, Children's National Hospital, Washington DC

Running title: HIV-1 Nef synergizes with APOL1 in cytotoxicity

Keywords: APOL1, HIV, nephropathy, Nef, nephrocyte, autophagy, HIVAN, *Drosophila*

* CORRESPONDING AUTHORS

Dr. Zhe Han, PhD

Center for Precision Disease Modeling

Department of Medicine, University of Maryland School of Medicine

670 West Baltimore Street, Baltimore, MD 21201, USA

Email: zhan@som.umaryland.edu; ORCID identifier: 0000-0002-5177-7798

Dr. Patricio E. Ray MD

Child Health Research Center,

Department of Pediatrics, University of Virginia School of Medicine

Room 2120, MR4 Building, 409 Lane Road, Charlottesville, VA 22908, USA

E-mail: Pray@virginia.edu; ORCID identifier: 0000-0003-2223-9584

HIGHLIGHTS

- A new transgenic *Drosophila* model to study the pathogenesis of HIV-1-related kidney diseases with nephrocyte-specific expression of HIV-1 *nef* and an *APOL1-G1* risk allele derived from a patient with HIVAN.

- APOL1-G1 caused organelle acidification defects, reduced formation of autophagolysosomes, and reduced autophagy and protein aggregation, which culminated in ER stress.

- HIV-1 Nef induced ER stress through an autophagy-independent pathway. Furthermore, Nef and APOL1-G1 acted synergistically to heighten ER stress, which resulted in nephrocyte dysfunction and cell death.

SIGNIFICANCE STATEMENT

APOL1 risk alleles are strongly linked to HIV-associated nephropathy (HIVAN) in people of African descent, but how HIV-1 and APOL1 interact and which pathways they might converge upon is unclear. A new *Drosophila* model to study HIV-1 Nef and APOL1-G1 (a risk allele) showed that Nef can induce ER stress in nephrocytes by itself, as well as exacerbate the organelle acidification defects and reduced autophagy induced by APOL1-G1, which further stimulates ER stress to a level that could cause nephrocyte cell death. Thus, we identified ER stress as the converging point for the synergy between APOL1-G1 and HIV-1 Nef in kidney cells, providing a potential therapeutic target for HIV-1 and APOL1-associated nephropathies.

ABSTRACT

Background: People carrying two *APOL1* risk alleles (RA) *G1* or *G2* are at greater risk of developing HIV-associated nephropathy (HIVAN). Studies in transgenic mice showed that the expression of HIV-1 genes in podocytes, and *nef* in particular, led to HIVAN. However, it remains unclear whether APOL1-RA and HIV-1 Nef interact to induce podocyte cell death.

Method: We generated transgenic (Tg) flies that express *APOL1-G1* (derived from a child with HIVAN) and HIV-1 *nef* specifically in the nephrocytes, the fly equivalent of mammalian podocytes, and assessed their individual and combined effects on the nephrocyte filtration structure and function.

Results: We found that HIV-1 Nef acts in synergy with APOL1-G1 resulting in nephrocyte structural and functional defects. Specifically, HIV-1 Nef itself can induce endoplasmic reticulum (ER) stress without affecting autophagy. Furthermore, Nef exacerbates the organelle acidification defects and autophagy reduction induced by APOL1-G1. The synergy between HIV-1 Nef and APOL1-G1 is built on their joint effects on elevating ER stress, triggering nephrocyte dysfunction and ultimately cell death.

Conclusions: Using a new *Drosophila* model of HIV-1-related kidney diseases, we identified ER stress as the converging point for the synergy between HIV-1 Nef and APOL1-G1 in inducing nephrocyte cell death. Given the high relevance between *Drosophila* nephrocytes and human podocytes, this finding suggests ER stress as a new therapeutic target for HIV-1 and APOL1-associated nephropathies.

INTRODUCTION

HIV-associated nephropathy (HIVAN) is a kidney disease characterized by heavy proteinuria

and rapid progression to chronic kidney failure [1,2]. HIVAN renal histology shows collapsing

glomerulopathy, de-differentiation and proliferation of podocytes and glomerular parietal

epithelial cells, development of focal and segmental glomerulosclerosis (FSGS), and

microcystic dilatation of renal tubules leading to kidney enlargement [1,2]. People of African

descent are at increased risk of developing HIVAN. Their risk to develop HIVAN or another

chronic kidney disease is strongly associated with two *Apolipoprotein L1* risk alleles

(*APOL1-RA*), *G1* and *G2* [3,4]. People living with HIV-1 that carry two copies of the *APOL1-RA*

have a ~50% lifetime risk of developing HIVAN [3–5], and poorly controlled HIV-1 infection is the

most powerful factor known to contribute to APOL1-kidney diseases [4]. However, to date, the

basic mechanisms through which the APOL1-RA interacts with HIV-1 to cause HIVAN remain

unclear.

Mice and rats do not have an *APOL1* ortholog, nor can they be infected with HIV-1 [6,7].

Nonetheless, transgenic (Tg) rodent models have been used to characterize the molecular

mechanism through which the APOL1-RA and HIV-1 genes induce renal diseases [8,9].

HIV-Tg26 mice and rats carrying a replication defective HIV-1 construct (driven by HIV-LTR

lacking a 3kb sequence with the *gag* and *pol* genes, which are essential for viral replication),

showed that the expression of HIV-genes in podocytes and tubular epithelial cells plays a

critical role in inducing HIVAN in rodents [10–15]. In addition, several HIV-Tg mouse models

showed that the HIV accessory protein negative factor (Nef), is a key determinant of HIVAN

pathogenesis [10–15]. Studies in cultured kidney cells and *APOL1* Tg mice found that the

expression levels and localization of APOL1 are crucial in determining its cytotoxicity [17–20].

Notably, different dual APOL1-HIV-Tg26 mouse models have generated conflicting results,

supporting either toxicity or protection [17,21]. Thus, it remains unclear whether HIV-1 Nef and

APOL1-RA interact directly to induce HIVAN, underlining the need for new animal models.

Drosophila is an established model for kidney diseases and development [18,22–27]. The fly

nephrocyte is equivalent to the mammalian podocyte, with highly conserved filtration structure

called the slit diaphragm [28–31]. They also share many other similarities in endocytosis and

exocytosis for the formation and maintenance of the filtration structure [28–31]. A majority of

genes associated with kidney diseases have functional homologs in flies that are required for

nephrocyte function [23,27]. The *Drosophila* nephrocyte has been well-established to study the

pathogenesis of *APOL1*-associated nephropathies [18,25,26,32,33]. Here, we generated Tg flies

that express HIV-1 *nef* and *APOL1-G1*, the most frequent risk variant, specifically in

nephrocytes. Our findings indicate that Nef and APOL1-G1 act synergistically to induce

nephrocyte cell death through the ER stress pathway, providing a potential therapeutic

target for HIV-1 and APOL1-associated nephropathies.

METHODS

Fly strains

Flies were maintained on standard food (Meidi LLC) at 22°C under standard conditions. The

following *Drosophila* lines were obtained from the Bloomington Drosophila Stock Center

(BDSC): *Dot*-Gal4 (ID_6903), UAS-YFP-*Rab5* (ID_24616), UAS-YFP-*Rab7* (ID_23270),

UAS-mCherry-*Atg8a* (ID_37750), and *w*[1118] (ID_3605). We previously generated the

Hand-GFP flies labeling the nuclei of nephrocytes and cardiomyocytes [34].

The APOL1-G0 and APOL1-G1 (cDNA from a child with HIVAN) *Drosophila* lines were

generated as described before [18]. The HIV-1 Nef cDNA (pGM91) was obtained from the NIH

AIDS Research and Reference Reagent Program (contributed by Dr. John Rossi) [35]. To

generate UAS-*nef* constructs, the *nef* allele cDNA was cloned into pUAST-attB and the

transgene was introduced into a docking site by germline transformation. Flies expressing

Dot-Gal4;UAS-*APOL1*-(*G0*/*G1*) were crossed with UAS-*nef* transgenic flies at 22°C.

10 kD dextran and FITC-albumin uptake assay

Dextran and albumin uptake by nephrocytes was assessed *ex vivo* in (20-day) female adult

flies to expose nephrocytes in artificial hemolymph containing 70 mmol/l NaCl, 5 mmol/l KCl,

1.5 mmol/l $CaCl_2$, 4 mmol/l $MgCl_2$, 10 mmol/l $NaHCO_3$, 5 mmol/l trehalose, 115 mmol/l

sucrose, and 5 mmol/l HEPES (Sigma-Aldrich). Following a 20-minute incubation with 10kD

Texas Red-dextran (0.05 mg/ml, Invitrogen) or FITC-albumin (100 mM, Sigma-Aldrich) and

fixation (10 min) in 4% paraformaldehyde in phosphate buffered saline (4%PFA), the

nephrocytes were mounted with VectaShield mounting medium (Vector Laboratories), and

imaged.

Nephrocyte size and number

Nephrocytes were dissected from 20-day-old female adult flies and kept in artificial

hemolymph, followed by fixation (10 min) in 4%PFA, and imaged. Nephrocyte size was

determined using the area measurement function in ImageJ [36] (version 1.52a). Nephrocyte

numbers were manually counted using images of *Hand-GFP*.

Immunochemistry

Female (20-day-old) flies were dissected and fixed [20 sec heat fix in 100°C artificial

hemolymph for Polychaetoid (Pyd); or in 4%PFA for Ref(2)p, ubiquitin (FK2), and PDI].

Immunochemistry was carried out using established methods [18]. Antibodies used in this study

includes: mouse anti-Pyd antibody (1:100, Developmental Studies Hybridoma Bank); rabbit

anti-Ref(2)P (1:100, ab178440, Abcam); mouse anti-ubiquitin FK2 (1:100, Enzo Life

Sciences); rabbit anti-protein disulfide isomerase (PDI) (1:100, Sigma-Aldrich), and Alexa

Fluor 555 (1:1,000, Thermo-Fisher-Scientific).

LysoTracker assay

Fly nephrocytes were dissected in artificial hemolymph, incubated (20 min) with LysoTracker

(Thermo-Fisher-Scientific), fixed (10 min) with 4%PFA, mounted with VectaShield, and

imaged.

TUNEL assay

Nephrocytes were dissected in artificial hemolymph and heat-fixed (100°C, 20 min) in 4%PFA,

blocked with blocking buffer (50mM Tris-Cl pH 7.4, 0.1% Triton X-100 and 188mM NaCl),

incubated with In Situ Cell Death detection kit (Roche), mounted with VectaShield and

imaged.

Confocal imaging

Confocal images were obtained using a ZEISS LSM900 microscope and ZEN Blue (edition

3.0) acquisition software. For quantitative comparison of intensities, common settings were

chosen to avoid oversaturation and then applied across images for all samples within an

assay. ImageJ [36] (version 1.52a) was used for all image processing.

Statistical analysis

Statistical tests were performed using PAST.exe software (University of Oslo). Data were

tested for normality using the Shapiro-Wilk test ($\alpha=0.05$). Normally distributed data were

analyzed by one-way ANOVA followed by Tukey-Kramer post-test for comparing multiple

groups. Non-normal distributed data were analyzed by Kruskal-Wallis H-test followed by a

Dunn's test for comparisons between multiple groups. The results are presented as mean ±

s.d. Statistical significance was determined as $P<0.05$. Details are provided in Supplementary

File 1.

RESULTS

HIV-1 Nef exacerbates APOL1-G1-induced nephrocyte dysfunction

We previously generated transgenic *Drosophila* lines with nephrocyte-specific expression of *APOL1-G0* or the *APOL1-G1* risk allele derived from a child with HIVAN [18]. Here, we generated transgenic flies with nephrocyte-specific expression of HIV-1 *nef* to explore how it alone or combined with *APOL1* affects the structure and function of nephrocytes.

We determined the capacity of dissected nephrocytes to filter and endocytose 10kD dextran or larger FITC-albumin particles. HIV-1 *nef* nor *APOL1-G0* showed a reduction in uptake (Figure 1A-C). Nephrocytes expressing *APOL1-G0+nef* or *APOL1-G1+nef* showed a significant decreased uptake of dextran and FITC-albumin compared to those expressing *nef, APOL1-G0*, or *APOL1-G1*; with the greatest reduction in *APOL1-G1+nef* nephrocytes (Figure 1A-C), reflecting the higher toxicity of the *APOL1-G1* risk allele. These results show that HIV-1 Nef exacerbates APOL1-induced nephrocyte dysfunction.

Combined HIV-1 Nef and APOL1 cause nephrocyte hypertrophy and cell death

Previously, we showed that *APOL1-G1* induces progressive hypertrophy and accelerated cell death in nephrocytes [18]. Hypertrophy validated here in 20-day-old fly nephrocytes expressing *APOL-G1* (Figure 1D). In contrast, nephrocytes expressing HIV-1 *nef* or *APOL1-G0*, did not show a significant change (Figure 1D), while those expressing *APOL1-G0+nef* or *APOL1-G1+nef* were significantly larger; with *APOL1G1+nef* nephrocytes the biggest (Figure 1D).

We observed reduced nephrocyte numbers in 20-day-old *APOL1-G1* Tg flies (Figure 1E). However, we did not observe significant changes in nephrocyte numbers in flies expressing HIV-1 *nef* or *APOL1-G0* (Figure 1E). In contrast, flies expressing *APOL1-G0+nef* or *APOL1-G1+nef*, showed a significant reduction in nephrocyte number compared to those expressing *APOL1-G0* or *APOL1-G1*; with numbers lowest in *APOL1-G1+nef* (Figure 1E).

These results show that HIV-1 Nef combined with APOL1-G1 exacerbates the damage and

loss of nephrocytes.

HIV-1 Nef and APOL1-G1 combined disrupt the slit diaphragm filtration structure

The *Drosophila* slit diaphragm is a highly organized structure critical for the nephrocyte

filtration function [28]. Thus, we assessed the localization of the slit diaphragm protein

polychaetoid (Pyd - the homolog of human tight junction protein ZO-1) [30]. Control nephrocytes

showed the typical fingerprint-like localization pattern of Pyd on the cell surface (Figure 2A),

and the characteristic continuous circular ring in the optical medial view (Figure 2B). Notably,

nephrocytes expressing *nef* or *APOL1-G0* did not show a visible disruption of Pyd (Figure

2A,B). However, nephrocytes expressing *APOL1-G0+nef* or *APOL1-G1+nef* showed highly

disorganized Pyd localization, more so than either APOL-1 risk variant alone. Moreover,

mislocalized Pyd was most evident in nephrocytes that expressed *APOL1-G1+nef* (Figure

2A,B), consistent with the altered uptake observed in nephrocytes expressing *nef* and

APOL1-G1 (Figure 1) which indicates a disrupted slit diaphragm filtration structure.

Combined HIV-1 Nef and APOL1-G1 exert detrimental effects on nephrocyte

endocytosis

We previously showed that HIV-1 can infect cultured human podocytes via

dynamin-dependent endocytosis in a process facilitated by transmembrane TNF-α [8]. Here, we

assayed the expression of early (Rab5) and late (Rab7) endosomal markers in nephrocytes.

Expression of *nef*, *APOL1-G0,* or *APOL1-G1*, nor combined, induced changes in the

expression level or localization of Rab5 (Figure 3A,B). In contrast, Rab7 protein levels were

significantly increased in nephrocytes expressing *APOL1-G1*, but not in those expressing *nef*

or *APOL1-G0* (Figure 3C,D). Nephrocytes expressing *APOL1-G0+nef* or *APOL1-G1+nef* had

significantly higher Rab7 protein levels; with the highest expression levels in *APOL1-G1+nef*

nephrocytes (Figure 3C,D).

We also showed previously that *APOL1-G1* altered the acidification of organelles and the

lysosomes in nephrocytes, which could impair their function [18]. Here, we used LysoTracker to

examine acidic vacuoles and lysosomes in nephrocytes with *nef* and *APOL1*. LysoTracker

signal was significantly reduced in nephrocytes expressing *APOL1-G1*, but not *nef* or

APOL1-G0 (Figure 3E,F). Nephrocytes expressing *APOL1-G0+nef* or *APOL1-G1+nef*

showed significantly lower LysoTracker signals (Figure 3E,F). Moreover, *APOL1-G1+nef*

nephrocytes had the lowest signal (Figure 3E,F). These findings indicate that HIV-1 Nef

exacerbates the changes induced by APOL1-G1 in key endocytic and acidification pathways

that are essential for nephrocyte function.

HIV-1 Nef enhances APOL1-G1-induced autophagic changes with accumulation of

autophagosomes in fly nephrocytes

APOL1 can modulate autophagy in various experimental model systems [17,19,25,32,37–39]. In

Drosophila, autophagy-related 8a (Atg8a), the equivalent of human LC3, is a widely used

marker for autophagy. Nephrocytes expressing *nef* showed no altered Atg8a-mCherry, but

those expressing *nef* together with *APOL1* showed significantly higher levels of

Atg8a-mCherry, with the highest signal in *APOL1-G1+nef* nephrocytes (Figure 4A,B).

Likewise, we did not find altered refractory to Ref(2)P, a marker for cargo destined to be

degraded by autophagy [40], in nephrocytes with *nef* alone (Figure 4C,D). In contrast, Ref(2)P

was significantly increased in nephrocytes expressing *APOL1-G1*, and these changes were

exacerbated in *APOL-G1+nef* nephrocytes (Figure 4C,D). In addition, we observed increased

accumulation of ubiquitinylated proteins in nephrocytes expressing *APOL1-G1*, but not *nef* or

APOL1-G0 (Figure 4E,F). This accumulation was highest in *APOL1-G1+nef* nephrocytes

(Figure 4E,F). These data demonstrate that HIV-1 Nef markedly increases APOL1-induced

autophagy changes in nephrocytes.

HIV-1 Nef and APOL1-G1 induce ER stress in nephrocytes

APOL1-G1 can elicit the endoplasmic reticulum (ER) stress response [26,32], while HIV-1 Nef

can bind the ER chaperone Calnexin (CNX) to induce ER stress [41–43]. Therefore, we assessed

protein disulfide-isomerase (PDI), a marker of ER stress in nephrocytes. PDI was significantly

increased in nephrocytes expressing *nef* (Figure 5A,B). This increase was significantly more

substantial when *nef* was expressed simultaneously with *APOL1-G1*; even when compared to

APOL1-G1 alone or *APOL1-G0+nef* nephrocytes (Figure 5A,B). Furthermore, nephrocytes

that expressed *APOL1-G1*, but not *nef* or *APOL1-G0*, showed positive results in the TUNEL

assay, a marker of cell death (Figure 5C). TUNEL-positive cells were also observed when

nephrocytes expressed *APOL1-G0+nef* or *APOL1-G1+nef* (Figure 5C). This indicates that

HIV-1 Nef alone can induce ER stress in *Drosophila* nephrocytes in an

autophagy-independent manner, and that Nef can exacerbate the APOL1-G1-induced

autophagy-dependent pathway to further stimulates ER stress, leading to nephrocyte

dysfunction and cell death.

DISCUSSION

We have developed a new *Drosophila* system to explore how HIV-1 Nef and APOL1-G1

interact in nephrocytes. It revealed that HIV-1 Nef in addition to causing ER stress directly

through an autophagy-independent pathway, also exacerbates the pathogenic effects of

APOL1-G1 on the autophagy pathway, disrupting the function of nephrocytes and leading to

their premature death.

APOL1-G1-induced cytotoxicity is not exclusively dependent on the nucleotide substitutions

that define this risk variant. It is also affected by the BH3 domain [26,44], exon 4-encoded

sequences [37], and different haplotypes [45]. Therefore, we generated Tg flies expressing a

common natural *APOL1-G1* derived from culture podocytes from a child with HIVAN [14]. This

APOL1-G1 contains the haplotypes (E150; I228; K255) and only differs at S342 and I384 from

a common natural APOL1-G0 haplotype used to generate the control flies. Notably, this

APOL1-G0 control haplotype is more toxic in cultured human kidney epithelial cells (HEK 293)

than the reference APOL1-G0 haplotype [14,19,25,46–52]. We previously showed that *APOL1-G0*

expression in nephrocytes induced toxicity, albeit less severe compared to *APOL1-G1* [18,32].

When we reduced the expression level of *APOL1-G0* in nephrocytes by lowering the

temperature to 22°C, *APOL1-G0* toxicity was eliminated [33]; hence all current fly experiments

were conducted at 22°C (instead of 25°C used in our previous study) so that only *APOL1-G1*

remains toxic. Nevertheless, we found that Nef could also synergize with APOL1-G0, making

it toxic at 22°C (Figure 1), consistent with clinical studies showing that people of West African

genetic ancestry living with a high HIV viral load can develop HIVAN even if they do not carry

an *APOL1-RA* [4].

A study using *nephrin* promoter driven *APOL1-G0* Tg mice crossbred with HIV-Tg26 mice

showed that fewer renal HIVAN lesions were developed compared to single HIV-Tg26 mice,

suggesting a protective role for APOL1-G0 against HIV-induced nephropathy [21].   However,

studies in *APOL1* Tg mice with a BAC chromosome harboring *G0*, *G1,* or *G2* human *APOL1*

found greater renal pathogenicity in *G2*/*G2* transgenic mice, compared to *G1*/*G1*; and that

APOL1-G0 did not rescue *APOL1-RA*-induced kidney injury when these mice were injected

with Interferon-gamma (INF-γ) [20]. Likewise, we found that *APOL1-G2* Tg flies developed more

significant nephrocyte injury compared to *APOL1-G1* Tg flies [32]. Overall, these studies

indicate that the toxicity of APOL1-RA is mainly a function of its renal expression level [20],

explaining why *APOL1* Tg mice driven by different promoters show different results [9,17,20,53].

Our data is also consistent with a recent study using dual BAC/*APOL1*-HIV-Tg26 mice [54] to

show that *APOL1-G0* did not minimize HIVAN lesions, whereas *APOL1-G1* aggravated them

[54].

Previous studies showed that HIV-1 Nef can induce ER stress by binding a chaperon protein,

Calnexin (CNX) [41–43]. We showed that Nef caused ER stress in nephrocytes through an

autophagy-independent pathway, likely by binding to a similar chaperon. In addition, we

demonstrated that Nef together with APOL1-G1, disrupted the autophagy pathway (Figure 4)

and promoting additional ER stress [26,32] (Figure 5). Defective endosomal trafficking has been

reported in yeast and flies that expressed *APOL1-G1* [25]. Likewise, immunofluorescence

studies in cultured human podocytes revealed increased Rab7 (late endosomes) and LC3II

(autophagosomes) positive organelles in cells expressing the high-risk (G1/G2) compared to

low-risk (G0/G0) alleles [17]. Studies in inducible podocyte-specific *APOL1-RA* transgenic mice

also showed that G1 interferes with intracellular vesicular trafficking by impairing endocytic,

autophagic, and acidification pathways, as well as by disrupting the maturation of

autophagosomes and autophagic flux, which correlated with the development of FSGS [17].

APOL1-RA was also shown to affect autophagy in cultured human podocytes [65,66].

Several studies have shown that HIV-1 can restrict autophagy to promote viral replication by

inhibiting the fusion between autophagosomes and lysosomes [38,55–57]. Nef overexpression in

human astrocytes led to increased ATG8/LC3 [56], and impaired acidification in human

cardiomyocytes, altering Rab7 localization and autophagy pathways [57]. Alternatively, HIV-1

Nef reduced the fusion of autophagosomes to lysosomes by interacting with Beclin1

(BECN1)/Rab7 in cultured human cells [58,59]. Together with our finding here, it appears that

HIV-1 Nef-mediated inhibition of autophagic flux is conserved across cell types and species.

Notably, the minor changes induced by Nef alone in nephrocytes resemble the mild podocyte

changes observed in Tg mice with podocyte-specific *nef* expression [12,60]. Whereas, other

studies showed more severe podocyte injury when HIV-1 *nef* and *vpr* were expressed

simultaneously in mouse podocytes[12], suggesting that other HIV-1 genes and cytokines

released by HIV infected cells are needed to evoke HIVAN phenotype fully. HIV-*nef* Tg mice

driven by the CD4 promoter also developed kidney disease [15,61,62]. Even though podocytes

might not express HIV-1 genes [15,61,62], Nef can still be released by HIV-infected cells

predominately in exosomes [63] and can be taken up by uninfected cells [64].

Our fly model has limitations, including that it is based on the ectopic overexpression of

APOL1 haplotypes, which can affect its intracellular localization and cytotoxicity [7, 8], and

bypasses the infection or renal cells and the effects of other HIV-genes and circulating

cytokines acting as second hits. HIVAN is a complex disease, involving many renal cell types

and structures that cannot be studied in flies. However, our model permits the exploration of

HIV-1 Nef and APOL1 interactions independently of many confounding variables. Most

findings from Tg mice, cultured podocytes, and HIV-infected cells published so far are

consistent with what we observed in nephrocytes. Since most relevant pathogenic pathways

are highly conserved between fly nephrocytes and human podocytes, this new fly model

provides a unique opportunity for rapid and cost-effective genetic or drug screens to better

understand the APOL1 and HIV-1 interactions for inducing renal injuries, as well as to discover

novel therapeutic treatment.

DISCLOSURES

The authors declare no competing interests.

FUNDING

This work was supported by National Institutes of Health grants R01-DK115968 (P.E.R. and

Z.H.), R01-DK103564 (P.E.R.), R01-DK120908 (Z.H.), and R01-DK098410 (Z.H.).

ACKNOWLEDGMENTS

We thank the Bloomington Drosophila Stock Center (BDSC) based at Indiana University

(Bloomington, IN) for the *Drosophila* stocks; and the Developmental Studies Hybridoma Bank

(DSHB) based at the University of Iowa (Iowa City, IA) for providing the antibodies.

AUTHOR CONTRIBUTIONS

J.Z. and Z.H. designed the study; J.Z., Y.F., J.Y. and JL. carried out the experiments; J.Z.,

J.vdL., and Z.H. analyzed and interpreted the data; J.Z. prepared the figures; J.Z, J.vdL., P.R.,

and Z.H. drafted and revised the manuscript; the manuscript has been critically reviewed and

the final version approved by all authors.

REFERENCES

1. Ross MJ: Advances in the pathogenesis of HIV-associated kidney diseases. *Kidney Int.*
86: 266–274, 2014

2. Swanepoel CR, Atta MG, D'Agati VD, Estrella MM, Fogo AB, Naicker S, et al.: Kidney
disease in the setting of HIV infection: conclusions from a Kidney Disease: Improving
Global Outcomes (KDIGO) Controversies Conference. *Kidney Int.* 93: 545–559, 2018

3. Kopp JB, Nelson GW, Sampath K, Johnson RC, Genovese G, An P, et al.: APOL1
genetic variants in focal segmental glomerulosclerosis and HIV-associated nephropathy.
J. Am. Soc. Nephrol. 22: 2129–2137, 2011

4. Kasembeli AN, Duarte R, Ramsay M, Mosiane P, Dickens C, Dix-Peek T, et al.: APOL1
risk variants are strongly associated with HIV-associated nephropathy in black South
Africans. *J. Am. Soc. Nephrol.* 26: 2882–2890, 2015

5. Rednor SJ, Ross MJ: Molecular Mechanisms of Injury in HIV-Associated Nephropathy.
Front. Med. 5: 177, 2018

6. Keppler OT, Welte FJ, Ngo TA, Chin PS, Patton KS, Tsou C-L, et al.: Progress toward a
human CD4/CCR5 transgenic rat model for de novo infection by human
immunodeficiency virus type 1. *J. Exp. Med.* 195: 719–736, 2002

7. Baumann JG, Unutmaz D, Miller MD, Breun SKJ, Grill SM, Mirro J, et al.: Murine T cells
potently restrict human immunodeficiency virus infection. *J. Virol.* 78: 12537–12547, 2004

8. Li J, Das JR, Tang P, Han Z, Jaiswal JK, Ray PE: Transmembrane TNF-α Facilitates
HIV-1 Infection of Podocytes Cultured from Children with HIV-Associated Nephropathy. *J.*
Am. Soc. Nephrol. 28: 862–875, 2017

9. Yoshida T, Latt KZ, Rosenberg AZ, Shrivastav S, Heymann J, Halushka MK, et al.:
Transcriptomic Analysis of Human Podocytes In Vitro: Effects of Differentiation and
APOL1 Genotype. *Kidney Int Rep* 8: 164–178, 2023

10. Kajiyama W, Kopp JB, Marinos NJ, Klotman PE, Dickie P: Glomerulosclerosis and viral
gene expression in HIV-transgenic mice: role of nef. *Kidney Int.* 58: 1148–1159, 2000

11. Ray PE, Liu X-H, Robinson LR, Reid W, Xu L, Owens JW, et al.: A novel HIV-1 transgenic
rat model of childhood HIV-1-associated nephropathy. *Kidney Int.* 63: 2242–2253, 2003

12. Zuo Y, Matsusaka T, Zhong J, Ma J, Ma L-J, Hanna Z, et al.: HIV-1 genes vpr and nef
synergistically damage podocytes, leading to glomerulosclerosis. *J. Am. Soc. Nephrol.*
17: 2832–2843, 2006

13. Lu T-C, He JC, Wang Z-H, Feng X, Fukumi-Tominaga T, Chen N, et al.: HIV-1 Nef
disrupts the podocyte actin cytoskeleton by interacting with diaphanous interacting
protein. *J. Biol. Chem.* 283: 8173–8182, 2008

14. Xie X, Colberg-Poley AM, Das JR, Li J, Zhang A, Tang P, et al.: The basic domain of
HIV-tat transactivating protein is essential for its targeting to lipid rafts and regulating
fibroblast growth factor-2 signaling in podocytes isolated from children with
HIV-1-associated nephropathy. *J. Am. Soc. Nephrol.* 25: 1800–1813, 2014

15. Hu C, Priceputu E, Cool M, Chrobak P, Bouchard N, Forestier C, et al.: NEF-Induced
HIV-Associated Nephropathy Through HCK/LYN Tyrosine Kinases. *Am. J. Pathol.* 193:
702–724, 2023

16. Gharavi AG, Ahmad T, Wong RD, Hooshyar R, Vaughn J, Oller S, et al.: Mapping a locus
for susceptibility to HIV-1-associated nephropathy to mouse chromosome 3. *Proc. Natl.*
Acad. Sci. U. S. A. 101: 2488–2493, 2004

17. Beckerman P, Bi-Karchin J, Park ASD, Qiu C, Dummer PD, Soomro I, et al.: Transgenic
expression of human APOL1 risk variants in podocytes induces kidney disease in mice.
Nat. Med. 23: 429–438, 2017

18. Fu Y, Zhu J-Y, Richman A, Zhang Y, Xie X, Das JR, et al.: APOL1-G1 in Nephrocytes
Induces Hypertrophy and Accelerates Cell Death. *J. Am. Soc. Nephrol.* 28: 1106–1116,
2017

19. O'Toole JF, Schilling W, Kunze D, Madhavan SM, Konieczkowski M, Gu Y, et al.: ApoL1
Overexpression Drives Variant-Independent Cytotoxicity. *J. Am. Soc. Nephrol.* 29:
869–879, 2018

20. McCarthy GM, Blasio A, Donovan OG, Schaller LB, Bock-Hughes A, Magraner JM, et al.:
Recessive, gain-of-function toxicity in an APOL1 BAC transgenic mouse model mirrors
human APOL1 kidney disease. *Dis. Model. Mech.* [Internet] 14: 2021 Available from:
http://dx.doi.org/10.1242/dmm.048952

21. Bruggeman LA, Wu Z, Luo L, Madhavan S, Drawz PE, Thomas DB, et al.: APOL1-G0
protects podocytes in a mouse model of HIV-associated nephropathy. *PLoS One* 14:
e0224408, 2019

22. Zhang F, Zhao Y, Han Z: An in vivo functional analysis system for renal gene discovery in
Drosophila pericardial nephrocytes. *J. Am. Soc. Nephrol.* 24: 191–197, 2013

23. Fu Y, Zhu J-Y, Richman A, Zhao Z, Zhang F, Ray PE, et al.: A Drosophila model system
to assess the function of human monogenic podocyte mutations that cause nephrotic
syndrome. *Hum. Mol. Genet.* 26: 768–780, 2017

24. Hermle T, Braun DA, Helmstädter M, Huber TB, Hildebrandt F: Modeling Monogenic
Human Nephrotic Syndrome in the Drosophila Garland Cell Nephrocyte. *J. Am. Soc.*
Nephrol. 28: 1521–1533, 2017

25. Kruzel-Davila E, Shemer R, Ofir A, Bavli-Kertselli I, Darlyuk-Saadon I, Oren-Giladi P, et
al.: APOL1-Mediated Cell Injury Involves Disruption of Conserved Trafficking Processes.
J. Am. Soc. Nephrol. 28: 1117–1130, 2017

26. Gerstner L, Chen M, Kampf LL, Milosavljevic J, Lang K, Schneider R, et al.: Inhibition of
endoplasmic reticulum stress signaling rescues cytotoxicity of human apolipoprotein-L1
risk variants in Drosophila. *Kidney Int.* [Internet] 2022 Available from:
http://dx.doi.org/10.1016/j.kint.2021.12.031

27. Rani L, Gautam NK: Drosophila Renal Organ as a Model for Identification of Targets and
Screening of Potential Therapeutic Agents for Diabetic Nephropathy. *Curr. Drug Targets*
19: 1980–1990, 2018

28. Weavers H, Prieto-Sánchez S, Grawe F, Garcia-López A, Artero R, Wilsch-Bräuninger M,
et al.: The insect nephrocyte is a podocyte-like cell with a filtration slit diaphragm. *Nature*
457: 322–326, 2009

29. Zhuang S, Shao H, Guo F, Trimble R, Pearce E, Abmayr SM: Sns and Kirre, the
Drosophila orthologs of Nephrin and Neph1, direct adhesion, fusion and formation of a slit
diaphragm-like structure in insect nephrocytes. *Development* 136: 2335–2344, 2009

30. Wang L, Wen P, van de Leemput J, Zhao Z, Han Z: Slit diaphragm maintenance requires
dynamic clathrin-mediated endocytosis facilitated by AP-2, Lap, Aux and Hsc70-4 in
nephrocytes. *Cell Biosci.* 11: 83, 2021

31. Lang K, Milosavljevic J, Heinkele H, Chen M, Gerstner L, Spitz D, et al.: Selective
endocytosis controls slit diaphragm maintenance and dynamics in Drosophila
nephrocytes.   *Elife*   [Internet]   11:   2022   Available   from:
http://dx.doi.org/10.7554/eLife.79037

32. Zhu J-Y, Lee J-G, Fu Y, van de Leemput J, Ray PE, Han Z: APOL1-G2 accelerates
nephrocyte cell death by inhibiting the autophagy pathway. *Dis. Model. Mech.* [Internet]
2023 Available from: http://dx.doi.org/10.1242/dmm.050223

33. Lee J-G, Fu Y, Zhu J-Y, Wen P, van de Leemput J, Ray PE, et al.: A SNARE protective
pool antagonizes APOL1 renal toxicity in Drosophila nephrocytes. *Cell Biosci.* 13: 199,
2023

34. Han Z, Olson EN: Hand is a direct target of Tinman and GATA factors during Drosophila
cardiogenesis and hematopoiesis. *Development* 132: 3525–3536, 2005

35. Murakawa GJ, Zaia JA, Spallone PA, Stephens DA, Kaplan BE, Wallace RB, et al.: Direct
detection of HIV-1 RNA from AIDS and ARC patient samples. *DNA* 7: 287–295, 1988

36. Schneider CA, Rasband WS, Eliceiri KW: NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* 9: 671–675, 2012

37. Khatua AK, Cheatham AM, Kruzel ED, Singhal PC, Skorecki K, Popik W: Exon
4-encoded sequence is a major determinant of cytotoxicity of apolipoprotein L1. *Am. J.*
Physiol. Cell Physiol. 309: C22-37, 2015

38. Chang C, Young LN, Morris KL, von Bülow S, Schöneberg J, Yamamoto-Imoto H, et al.:
Bidirectional Control of Autophagy by BECN1 BARA Domain Dynamics. *Mol. Cell* 73:
339-353.e6, 2019

39. Chun J, Zhang J-Y, Wilkins MS, Subramanian B, Riella C, Magraner JM, et al.:
Recruitment of APOL1 kidney disease risk variants to lipid droplets attenuates cell
toxicity. *Proc. Natl. Acad. Sci. U. S. A.* 116: 3712–3721, 2019

40. Gatica D, Lahiri V, Klionsky DJ: Cargo recognition and degradation by selective
autophagy. *Nat. Cell Biol.* 20: 233–242, 2018

41. Jennelle L, Hunegnaw R, Dubrovsky L, Pushkarsky T, Fitzgerald ML, Sviridov D, et al.:
HIV-1 protein Nef inhibits activity of ATP-binding cassette transporter A1 by targeting
endoplasmic reticulum chaperone calnexin. *J. Biol. Chem.* 289: 28870–28884, 2014

42. Ryan D, Carberry S, Murphy ÁC, Lindner AU, Fay J, Hector S, et al.: Calnexin, an
ER-induced protein, is a prognostic marker and potential therapeutic target in colorectal
cancer. *J. Transl. Med.* [Internet] 14: 2016 Available from:
http://dx.doi.org/10.1186/s12967-016-0948-z

43. Guérin R, Arseneault G, Dumont S, Rokeach LA: Calnexin is involved in apoptosis
induced by endoplasmic reticulum stress in the fission yeast. *Mol. Biol. Cell* 19:
4404–4420, 2008

44. Heneghan JF, Vandorpe DH, Shmukler BE, Giovinazzo JA, Raper J, Friedman DJ, et al.:
BH3 domain-independent apolipoprotein L1 toxicity rescued by BCL2 prosurvival
proteins. *Am. J. Physiol. Cell Physiol.* 309: C332-47, 2015

45. Lannon H, Shah SS, Dias L, Blackler D, Alper SL, Pollak MR, et al.: Apolipoprotein L1
(APOL1) risk variant toxicity depends on the haplotype background. *Kidney Int.* 96:
1303–1307, 2019

46. Lan X, Jhaveri A, Cheng K, Wen H, Saleem MA, Mathieson PW, et al.: APOL1 risk
variants enhance podocyte necrosis through compromising lysosomal membrane
permeability. *Am. J. Physiol. Renal Physiol.* 307: F326-36, 2014

47. Nichols B, Jog P, Lee JH, Blackler D, Wilmot M, D'Agati V, et al.: Innate immunity
pathways regulate the nephropathy gene Apolipoprotein L1. *Kidney Int.* 87: 332–342,
2015

48. Olabisi OA, Zhang J-Y, VerPlank L, Zahler N, DiBartolo S 3rd, Heneghan JF, et al.:
APOL1 kidney disease risk variants cause cytotoxicity by depleting cellular potassium
and inducing stress-activated protein kinases. *Proc. Natl. Acad. Sci. U. S. A.* 113:
830–837, 2016

49. Granado D, Müller D, Krausel V, Kruzel-Davila E, Schuberth C, Eschborn M, et al.:
Intracellular APOL1 Risk Variants Cause Cytotoxicity Accompanied by Energy Depletion.
J. Am. Soc. Nephrol. 28: 3227–3238, 2017

50. Hayek SS, Koh KH, Grams ME, Wei C, Ko Y-A, Li J, et al.: A tripartite complex of suPAR,
APOL1 risk variants and αvβ3 integrin on podocytes mediates chronic kidney disease.
Nat. Med. 23: 945–953, 2017

51. Ma L, Chou JW, Snipes JA, Bharadwaj MS, Craddock AL, Cheng D, et al.: APOL1
Renal-Risk Variants Induce Mitochondrial Dysfunction. *J. Am. Soc. Nephrol.* 28:
1093–1105, 2017

52. Wen H, Kumar V, Lan X, Shoshtari SSM, Eng JM, Zhou X, et al.: APOL1 risk variants
cause podocytes injury through enhancing endoplasmic reticulum stress. *Biosci. Rep.*
[Internet] 38: 2018 Available from: http://dx.doi.org/10.1042/BSR20171713

53. Bruggeman LA, Wu Z, Luo L, Madhavan SM, Konieczkowski M, Drawz PE, et al.:
APOL1-G0 or APOL1-G2 Transgenic Models Develop Preeclampsia but Not Kidney
Disease. *J. Am. Soc. Nephrol.* 27: 3600–3610, 2016

54. Yoshida T, Latt KZ, Santo BA, Shrivastav S, Zhao Y, Fenaroli P, et al.: APOL1 kidney risk
variants in glomerular diseases modeled in transgenic mice. *bioRxiv* [Internet] 2023
Available from: http://dx.doi.org/10.1101/2023.03.27.534273

55. Campbell GR, Rawat P, Bruckman RS, Spector SA: Human Immunodeficiency Virus
Type 1 Nef Inhibits Autophagy through Transcription Factor EB Sequestration. *PLoS*
Pathog. 11: e1005018, 2015

56. Saribas AS, Khalili K, Sariyer IK: Dysregulation of autophagy by HIV-1 Nef in human
astrocytes. *Cell Cycle* 14: 2899–2904, 2015

57. Gupta MK, Kaminski R, Mullen B, Gordon J, Burdo TH, Cheung JY, et al.: HIV-1
Nef-induced cardiotoxicity through dysregulation of autophagy. *Sci. Rep.* 7: 8572, 2017

58. Franco-Juárez B, Coronel-Cruz C, Hernández-Ochoa B, Gómez-Manzo S,
Cárdenas-Rodríguez N, Arreguin-Espinosa R, et al.: TFEB; Beyond Its Role as an
Autophagy and Lysosomes Regulator. *Cells* [Internet] 11: 2022 Available from:
http://dx.doi.org/10.3390/cells11193153

59. Zhang W, Li X, Wang S, Chen Y, Liu H: Regulation of TFEB activity and its potential as a
therapeutic target against kidney diseases. *Cell Death Discov* 6: 32, 2020

60. Husain M, D'Agati VD, He JC, Klotman ME, Klotman PE: HIV-1 Nef induces
dedifferentiation of podocytes in vivo: a characteristic feature of HIVAN. *AIDS* 19:
1975–1980, 2005

61. Hanna Z, Kay DG, Rebai N, Guimond A, Jothy S, Jolicoeur P: Nef harbors a major
determinant of pathogenicity for an AIDS-like disease induced by HIV-1 in transgenic
mice. *Cell* 95: 163–175, 1998

62. Hanna Z, Kay DG, Cool M, Jothy S, Rebai N, Jolicoeur P: Transgenic mice expressing
human immunodeficiency virus type 1 in immune cells develop a severe AIDS-like
disease. *J. Virol.* 72: 121–132, 1998

63. Dubrovsky L, Brichacek B, Prashant NM, Pushkarsky T, Mukhamedova N, Fleetwood AJ,
et al.: Extracellular vesicles carrying HIV-1 Nef induce long-term hyperreactivity of
myeloid cells. *Cell Rep.* 41: 111674, 2022

64. Mukhamedova N, Hoang A, Dragoljevic D, Dubrovsky L, Pushkarsky T, Low H, et al.:
Exosomes containing HIV protein Nef reorganize lipid rafts potentiating inflammatory
response in bystander cells. *PLoS Pathog.* 15: e1007907, 2019

65. Kumar V, Ayasolla K, Jha A, Mishra A, Vashistha H, Lan X, et al.: Disrupted
apolipoprotein L1-miR193a axis dedifferentiates podocytes through autophagy blockade
in an APOL1 risk milieu. *Am. J. Physiol. Cell Physiol.* 317: C209–C225, 2019

66. Ekulu PM, Adebayo OC, Decuypere J-P, Bellucci L, Elmonem MA, Nkoy AB, et al.: Novel
human podocyte cell model carrying G2/G2 APOL1 high-risk genotype. *Cells* 10: 1914,
2021

FIGURES

Figure 1. Expression of HIV-1 protein Nef facilitated nephrocyte function reduction,

hypertrophy, and cell death due to APOL1-G1.

(A-E) Flies used (20-day-old adult females): Control (*Dot>w*[1118]); *nef* (*Dot>nef*-OE);

APOL1-G0 (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE); *APOL1-G0+nef*

(*Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>APOL1-G1*-OE+*nef*-OE). **(A) Top:** 10kD

fluorescent dextran particle uptake (red) by nephrocytes using nephrocyte specific driver

Dot-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C.

Scale bar: 15 µm. **Bottom:** FITC-albumin particle uptake (green) by nephrocytes using

nephrocyte specifically driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0*

and *APOL1-G1* at 22°C. Scale bar: 15 µm. **(B)** Quantitation of 10kD dextran uptake, relative to

uptake in control flies. n=40 nephrocytes from 6 flies, per group. **(C)** Quantitation of

FITC-albumin uptake, relative to uptake in control flies. n=40 nephrocytes from 6 flies, per

group. **(D)** Quantitation of adult nephrocyte size, relative to size in control flies. n=40

nephrocytes from 6 flies, per group. **(E)** Quantitation of adult nephrocyte number, relative to

cell number in control flies. n=10 flies, per group. Results have been presented as mean ± s.d.,

normalized to the control group. Kruskal–Wallis H-test followed by a Dunn's test; statistical

significance: *$P<0.05$, **$P<0.01$, ***$P<0.001$.

Figure 2. The expression of HIV-1 protein Nef in nephrocytes facilitated the disruption of the slit diaphragm due to APOL1-G1.

(A, B) Flies used (20-day-old adult females): Control (*Dot>w*^{1118}); *nef* (*Dot>nef*-OE); *APOL1-G0* (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>APOL1-G1*-OE+*nef*-OE). **(A, B)** Localization of slit diaphragm protein Polychaetoid (Pyd; red) at the surface (A) and medial (B) optical sections in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 µm; Scale bar (magnified): 1 µm.

Figure 3. The expression of HIV-1 protein Nef in nephrocytes facilitated the disruption in endocytic membrane trafficking due to APOL1-G1.

(A, B) Flies used (20-day-old adult females): Control (*Dot>Rab5*-YFP); *nef* (*Dot>Rab5*-YFP +*nef*-OE); *APOL1-G0* (*Dot>Rab5*-YFP+*APOL1-G0*-OE); *APOL1-G1* (*Dot>Rab5*-YFP+*APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>Rab5*-YFP+*APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>Rab5*-YFP+*APOL1-G1*-OE+*nef*-OE). **(C, D)** Flies used (20-day-old adult females): Control (*Dot>Rab7*-YFP); *nef* (*Dot>Rab7*-YFP +*nef*-OE); *APOL1-G0* (*Dot>Rab7*-YFP+*APOL1-G0*-OE); *APOL1-G1* (*Dot>Rab7*-YFP+*APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>Rab7*-YFP+*APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>Rab7*-YFP+*APOL1-G1*-OE+*nef*-OE). **(E, F)** Flies used (20-day-old adult females): Control (*Hand*-GFP, *Dot>*GFP+*w*^{1118}); *nef* (*Hand*-GFP, *Dot>nef*-OE); *APOL1-G0* (*Hand*-GFP, *Dot>APOL1-G0*-OE); *APOL1-G1* (*Hand*-GFP, *Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Hand*-GFP, *Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Hand*-GFP, *Dot>APOL1-G1*-OE+*nef*-OE). **(A)** Expression of endocytosis-related proteins Rab5 (green, early endosome) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 µm; Scale bar (magnified): 1 µm. **(B)** Quantitation

of Rab5 fluorescence intensity, relative to fluorescence in control nephrocytes. n=40

nephrocytes from 6 flies, per group. Note: None of the comparisons between groups were

significant. **(C)** Expression of endocytosis-related proteins Rab7 (green, late endosome) in

nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or

together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to

each image. Scale bar: 5 μm; Scale bar (magnified): 1 μm. **(D)** Quantitation of Rab7

fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from

6 flies, per group. **(E)** LysoTracker dye fluorescence level (red) in nephrocytes using the

nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0*

and *APOL1-G1* at 22°C. *Hand*-GFP transgene expression was visualized as green

fluorescence concentrated in the nuclei of nephrocytes. Scale bar: 15 μm. **(F)** Quantitation of

LysoTracker dye fluorescence intensity, relative to fluorescence in control nephrocytes. n=40

nephrocytes from 6 flies, per group. Results have been presented as mean ± s.d., normalized

to the control group. Kruskal–Wallis H-test followed by a Dunn's test; statistical significance:

*P<0.05, ***P<0.001.

Figure 4. The expression of HIV-1 protein Nef in nephrocytes facilitated autophagy

impairment due to APOL1-G1.

(A, B) Flies used (20-day-old adult females): Control (*Dot>Atg8a*-mCherry); *nef*

(*Dot>Atg8a*-mCherry+*nef*-OE); *APOL1-G0* (*Dot>Atg8a*-mCherry+*APOL1-G0*-OE);

APOL1-G1 (*Dot>* *Atg8a*-mCherry+*APOL1-G1*-OE); *APOL1-G0+nef*

(*Dot>Atg8a*-mCherry+*APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>Atg8a*-mCherry

+*APOL1-G1*-OE+*nef*-OE). **(C-F)** Flies used (20-day-old adult females): Control (*Dot>w*[1118]);

nef (*Dot>nef*-OE); *APOL1-G0* (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE);

APOL1-G0+nef (*Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef*

(*Dot>APOL1-G1*-OE+*nef*-OE). **(A)** Expression of autophagy-related protein 8a (Atg8a; red,

mCherry) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef*

alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified

next to each image. Scale bar: 5 μm; Scale bar (magnified): 1 μm. **(B)** Quantitation of Atg8a

fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from

6 flies, per group. **(C)** Autophagy receptor refractory to sigma P (Ref(2)P; red) in nephrocytes

using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with

APOL1-G0 and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image.

Scale bar: 5 μm; Scale bar (magnified): 1 μm. **(D)** Quantitation of Ref(2)P fluorescence

intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per

group. **(E)** Expression of ubiquitinylated proteins (green) in nephrocytes using the nephrocyte

specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and

APOL1-G1 at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 μm;

Scale bar (magnified): 1 μm. **(D)** Quantitation of ubiquitinylated proteins fluorescence

intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per

group. Results have been presented as mean ± s.d., normalized to the control group.

Kruskal–Wallis H-test followed by a Dunn's test; statistical significance: *P<0.05, **P<0.01,

***P<0.001.

Figure 5. Expression of HIV-1 protein Nef in nephrocyte-induced ER stress.

(A, B) Flies used (20-day-old adult females): Control (*Dot*>w^{1118}); *nef* (*Dot*>*nef*-OE);

APOL1-G0 (*Dot*>*APOL1-G0*-OE); *APOL1-G1* (*Dot*>*APOL1-G1*-OE); *APOL1-G0*+*nef*

(*Dot*>*APOL1-G0*-OE+*nef*-OE); *APOL1-G1*+*nef* (*Dot*>*APOL1-G1*-OE+*nef*-OE). **(C, D)** Flies

used (20-day-old adult females): Control (*Hand*-GFP, *Dot*>GFP+w^{1118}); *nef* (*Hand*-GFP,

Dot>*nef*-OE); *APOL1-G0* (*Hand*-GFP, *Dot*>*APOL1-G0*-OE); *APOL1-G1* (*Hand*-GFP,

Dot>*APOL1-G1*-OE); *APOL1-G0*+*nef* (*Hand*-GFP, *Dot*>*APOL1-G0*-OE+*nef*-OE);

APOL1-G1+*nef* (*Hand*-GFP, *Dot*>*APOL1-G1*-OE+*nef*-OE). **(A)** ER stress marker protein

disulfide isomerase (PDI; red) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to

express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Scale bar: 15

μm. **(B)** Quantitation of PDI fluorescence intensity, relative to fluorescence in control

nephrocytes. n=40 nephrocytes from 6 flies, per group. **(C)** Cell apoptosis marker TUNEL (red)

in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or

together with *APOL1-G0* and *APOL1-G1* at 22°C. *Hand*-GFP transgene expression was

visualized as green fluorescence concentrated in the nuclei of nephrocytes. Scale bar: 5 µm.

Figure 6. Model of HIV-1 Nef acting in synergy with APOL1-G1 through ER stress.

Graphic depiction of the proposed model by which HIV-1 Nef induces ER stress, which in turn

leads to cellular toxicity. Via a separate pathway, Nef exacerbates APOL1-G1-mediated ER

stress. APOL1-G1 disrupts the late endosomes and lysosomes, which affects the fusion of

autophagosomes and lysosomes, the reduced autophagy results in protein accumulation,

leading to ER stress. Thus, when together HIV-1 Nef and APOL1-G1 significantly increase

cellular toxicity through ER stress, which ultimately leads to cell death and tissue damage in

the kidneys. (Green text indicates data showed upregulation, whereas red text indicates data

showed downregulation, yellow text indicates no change.) APOL1-G1, Apolipoprotein L1-risk

allele G1; Atg8a-mCherry, Autophagy-related 8a with mCherry fluorescent tag; ER,

endoplasmic reticulum; PDI, protein disulfide isomerase; Rab5/7-YFP, Rab5 or Rab7 with

yellow fluorescent protein tag; Ref(2)p, refractory to sigma P; TUNEL, terminal

deoxynucleotidyl transferase dUTP nick end labeling assay.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6