HIV-1 Nef acts in synergy with APOL1-G1 to induce nephrocyte cell death in a new

Drosophila **model of HIV-related kidney diseases**

Jun-yi Zhu[1,2], Yulong Fu[1,3], Joyce van de Leemput[1,2] Ying Yu[4], Jinliang Li[5], Patricio E. Ray[4,*],

and Zhe Han[1,2,*]

1 Center for Precision Disease Modeling, Department of Medicine, University of Maryland

School of Medicine, Baltimore, Maryland

2 Division of Endocrinology, Diabetes and Nutrition, Department of Medicine, University of

Maryland School of Medicine, Baltimore, Maryland

3 Current: Department of Pathology, University of Alabama at Birmingham, Birmingham,

Alabama

4 Child Health Research Center, Department of Pediatrics, University of Virginia School of

Medicine

5 Children's Research Institute, Children's National Hospital, Washington DC

Running title: HIV-1 Nef synergizes with APOL1 in cytotoxicity

Keywords: APOL1, HIV, nephropathy, Nef, nephrocyte, autophagy, HIVAN, *Drosophila*

* CORRESPONDING AUTHORS

Dr. Zhe Han, PhD
Center for Precision Disease Modeling
Department of Medicine, University of Maryland School of Medicine
670 West Baltimore Street, Baltimore, MD 21201, USA
Email: zhan@som.umaryland.edu; ORCID identifier: 0000-0002-5177-7798

Dr. Patricio E. Ray MD
Child Health Research Center,
Department of Pediatrics, University of Virginia School of Medicine
Room 2120, MR4 Building, 409 Lane Road, Charlottesville, VA 22908, USA
E-mail: Pray@virginia.edu; ORCID identifier: 0000-0003-2223-9584

HIGHLIGHTS

- Developed a new transgenic *Drosophila* model to study the pathogenesis of HIV-1-related kidney diseases with nephrocyte-specific expression of HIV-1 *nef* and an *APOL1-G1* risk allele derived from a patient with HIVAN.
- APOL1-G1 caused organelle acidification defects, reduced formation of autophagolysosomes, and reduced autophagy and protein aggregation, which culminated in ER stress.
- HIV-1 Nef induced ER stress through an autophagy-independent pathway. In addition, Nef acted through an autophagy-dependent pathway that heightened the ER stress. It did so through APOL1-G1-mediated autophagy pathway disruption, which resulted in nephrocyte dysfunction and cell death.

SIGNIFICANCE STATEMENT

APOL1 risk alleles are strongly linked to HIV-associated nephropathy (HIVAN) in people of African descent, but how HIV-1 and APOL1 interact and which pathways they might converge upon is unclear. A new *Drosophila* model to study HIV-1 Nef and APOL1-G1 (a risk allele) showed that Nef can induce ER stress in nephrocytes by itself, as well as exacerbate the organelle acidification defects and reduced autophagy induced by APOL1-G1, which further stimulates ER stress to a level that could cause nephrocyte cell death. Thus, we identified ER stress as the converging point for the synergy between APOL1-G1 and HIV-1 Nef in kidney cells, providing a potential therapeutic target for HIV-1 and APOL1-associated nephropathies.

ABSTRACT

Background: People carrying two *APOL1* risk alleles (RA) *G1* or *G2* are at greater risk of developing HIV-associated nephropathy (HIVAN). Studies in transgenic mice showed that the expression of HIV-1 genes in podocytes, and *nef* in particular, led to HIVAN. However, it remains unclear whether APOL1-RA and HIV-1 Nef interact to induce podocyte cell death.

Method: We generated transgenic (Tg) flies that express *APOL1-G1* (derived from a child with HIVAN) and HIV-1 *nef* specifically in the nephrocytes, the fly equivalent of mammalian podocytes, and assessed their individual and combined effects on the nephrocyte filtration structure and function.

Results: We found that HIV-1 Nef acts in synergy with APOL1-G1 resulting in nephrocyte structural and functional defects. Specifically, HIV-1 Nef itself can induce endoplasmic reticulum (ER) stress (without affecting autophagy). Through a different pathway, Nef exacerbates the organelle acidification defects and reduced autophagy induced by APOL1-G1. The synergy between HIV-1 Nef and APOL1-G1 is built on their joint effects on elevating ER stress, triggering nephrocyte dysfunction and ultimately cell death.

Conclusions: A new *Drosophila* model of HIV-1-related kidney diseases identified ER stress as the converging point for the synergy between HIV-1 Nef and APOL1-G1 in inducing nephrocyte cell death. Given the high relevance between *Drosophila* nephrocytes and human podocytes, this finding suggests ER stress as a new therapeutic target for HIV-1 and APOL1-associated nephropathies.

INTRODUCTION

HIV-associated nephropathy (HIVAN) is a kidney disease characterized by heavy proteinuria

and rapid progression to chronic kidney failure if proper antiretroviral therapy (ART) is not

available [1,2]. From the renal histological point of view, HIVAN is characterized by collapsing

glomerulopathy, de-differentiation and proliferation of podocytes and glomerular parietal

epithelial cells, development of focal and segmental glomerulosclerosis (FSGS), and

microcystic dilatation of renal tubules leading to kidney enlargement [1,2]. People of African

descents are at increased risk of developing HIVAN. In this regard, previous studies showed

that two *Apolipoprotein L1* risk alleles (*APOL1-RA*), *G1* and *G2*, are strongly associated with

the risk for people of African descent to develop HIVAN, as well as other chronic kidney

diseases (CKD) [3,4]. Poorly controlled HIV infection is the most powerful factor known to

contribute to APOL1- kidney diseases [4]. People living with HIV-1 carrying two copies of the

APOL1 RA have a ~50% lifetime risk of developing HIVAN [3]. However, to date, the basic

mechanisms through which the APOL1-RA interacts with HIV to cause HIVAN are not clearly

understood.

Mice do not have an ortholog to the human *APOL1* gene and cannot be infected with HIV.

Nonetheless, APOL1 and HIV transgenic mouse models have been used to characterize the

molecular mechanism through which the APOL1-RA and HIV-1 genes induce renal diseases.

Functional studies to date exploring how APOL1 interacts with HIV have been limited to cell

culture models and HIV-transgenic (Tg) mice and rats [5,6]. HIV-Tg26 mice and rats carrying a

replication defective HIV construct driven by the HIV-LTR that lacks a 3kb sequence

containing the *gag* and *pol* HIV genes (which are essential for viral replication), showed that

the expression of HIV-genes in podocytes and tubular epithelial cells plays a critical role in

inducing HIVAN in rodents [7–12]. In addition, follow up studies in several HIV-Tg mouse models

showed that the HIV accessory protein negative factor (Nef), is a key determinant of the

pathogenesis of HIVAN [7–12]. Nonetheless, in the absence of APOL1, other genetic factors are

needed to facilitate the development of HIVAN in mice [13]. Previous studies in cultured kidney

cells as well as *APOL1* Tg mice and flies, have shown that the expression levels and

localization of APOL1 are crucial in determining its cytotoxicity [14–17]. Yet, different dual APOL1-

HIV-Tg26 mouse models have generated conflicting results, supporting toxicity or protection

[14,18], and it remains unclear whether HIV-1 Nef interacts with the APOL1-RA directly to induce

HIVAN. Thus, new animal models are needed to determine whether HIV-1 and APOL1 can

interact in a simultaneous and cell type specific manner.

Drosophila is an established model for kidney diseases and development [15,19–24]. Its

nephrocytes, equivalent to mammalian podocytes, contain a filtration structure made of lacuna

channels and a basement membrane[25–28]. Moreover, nephrocytes and podocytes have many

molecular similarities, including the importance of endocytosis and exocytosis for the

formation and maintenance of their slit diaphragm filtration structures [25–28]. Furthermore, many

of the genes associated with kidney diseases not only have functional homologs in *Drosophila*,

but these homologous genes are required for nephrocyte function [20,24]. In this capacity,

Drosophila nephrocytes have been used successfully to study the pathogenesis of *APOL1*-

associated nephropathies [15,22,23,29,30]. Therefore, since APOL1-G1 is the most frequent risk

variant present in African Americans living with HIVAN, we generated Tg flies expressing

HIV-1 *nef* and *APOL1-G1* specifically in nephrocytes. Our findings suggest that Nef and

APOL1-G1 act synergistically to induce nephrocyte cell death through the ER stress

pathway, providing a potential therapeutic target for HIV-1 and APOL1-associated

nephropathies.

METHODS

Fly strains

Flies were crossed, reared, and kept on standard food (Meidi LLC) at 22°C under standard conditions. The following *Drosophila* lines were obtained from the Bloomington Drosophila Stock Center (BDSC; Indiana University, IN): *Dot*-Gal4 (ID 6903; Ugt36A1 driver), UAS-YFP-*Rab5* (ID 24616), UAS-YFP-*Rab7* (ID 23270), UAS-mCherry-*Atg8a* (ID 37750), and w^{1118} (BDSC; ID 3605) which were used in the crosses. The *Hand*-GFP line (express green fluorescent protein in the nuclei of nephrocytes and cardiomyocytes) was previously generated by our team [31].

DNA cloning and generation of transgenic fly strains

The APOL1-G0 and APOL1-G1 *Drosophila* lines were generated as described before [15]. Briefly, the APOL1-G1 cDNA was obtained from a cDNA library generated from podocytes cultured from a child with HIV-associated nephropathy, as described by Xie et al. [11], and differs from the wild-type APOL1-G0 (AAI43040) only at S342 and I384. The HIV-1 Nef cDNA (pGM91) was obtained from the NIH AIDS Research and Reference Reagent Program (contributed by Dr. John Rossi) [32]. To generate UAS-*nef* constructs, the above cDNAs of the *nef* allele were cloned into the pUASTattB vector and the transgene was introduced into a fixed chromosomal docking site by germline transformation.

10 kD dextran and FITC-albumin uptake assay

20-day-old female flies from the *Dot*-Gal4 transgenic lines expressing UAS-*APOL1-G1* were crossed with UAS-*nef* transgenic flies at 22°C. Dextran and albumin uptake by nephrocytes was assessed ex vivo in adult flies by dissecting heart tissues in artificial hemolymph [70 mmol/l NaCl (Carolina), 5 mmol/l KCl (Sigma), 1.5 mmol/l $CaCl_2 \cdot 2H_2O$ (Sigma-Alrich), 4 mmol/l $MgCl_2$ (Sigma-Aldrich), 10 mmol/l $NaHCO_3$ (Sigma-Aldrich), 5 mmol/l trehalose (Sigma), 115 mmol/l sucrose (Sigma-Aldrich), and 5 mmol/l HEPES (Sigma-Aldrich), in H_2O] and examining

cells by fluorescence microscopy after 20 minutes of incubation with Texas Red-dextran (10

kD, 0.05 mg/ml, Invitrogen) or FITC-albumin (100 mM, Sigma-Aldrich) and 10 minutes fixation

in 4% paraformaldehyde in phosphate buffered saline (1xPBS, Thermo Fisher Scientific). The

nephrocytes were mounted with Vectashield mounting medium (H-1000, Vector Laboratories).

For quantification, 40 nephrocytes were analyzed from 6 female flies per genotype.

Nephrocyte size and number

For this in vivo assay, we dissected 20-day-old adult female flies. The flies were from the *Dot-*

Gal4 transgenic lines that expressed UAS-APOL1-G0 or UAS-APOL1-G1 were crossed with

flies from the UAS-nef transgenic line at 22°C. The nephrocytes were dissected and kept in

artificial hemolymph [70 mmol/L NaCl (Carolina), 5 mmol/L KCl (Sigma), 1.5 mmol/L

$CaCl_2.2H_2O$ (Sigma-Aldrich), 4 mmol/L $MgCl_2$ (Sigma-Aldrich), 10 mmol/L $NaHCO_3$ (Sigma-

Aldrich), 5 mmol/L trehalose (Sigma), 115 mmol/L sucrose (Sigma-Aldrich), and 5 mmol/L

HEPES (Sigma-Aldrich), in H_2O], followed by a 10 minutes fixation in 4% paraformaldehyde

in 1XPBS (Thermo scientific). Images were obtained by confocal microscopy (ZEISS LSM

900; see details below). Nephrocyte size was determined using the area measurement

function in ImageJ [33] (version 1.52a), whereas nephrocyte numbers were manually counted in

the same fluorescent images. For quantitation of nephrocyte size, 40 nephrocytes were

analyzed from 6 female flies per genotype. For quantitation of nephrocyte number, 10 female

flies were analyzed per genotype.

Fluorescent immunochemistry

For Polychaetoid (Pyd) immunochemistry, 20-day-old female flies were dissected and heat-

fixed for 20 seconds in 100°C artificial hemolymph [70 mmol/l NaCl (Carolina), 5 mmol/l KCl

(Sigma), 1.5 mmol/l $CaCl_2·2H_2O$ (Sigma-Alrich), 4 mmol/l $MgCl_2$ (Sigma-Aldrich), 10 mmol/l

$NaHCO_3$ (Sigma-Aldrich), 5 mmol/l trehalose (Sigma), 115 mmol/l sucrose (Sigma-Aldrich),

and 5 mmol/l HEPES (Sigma-Aldrich), in H_2O]. Primary mouse monoclonal anti-Pyd antibody

(PYD2) was obtained from Developmental Studies Hybridoma Bank and was used at 1:100

dilutions in 1xPBS with 0.1% Triton X-100 (Sigma) (PBST). The secondary antibody Alexa

Fluor 555 (A-21422, Thermo Fisher Scientific) was used diluted 1:1,000 in PBST. The

nephrocytes were washed with PBST three times, blocked in PBST+2% bovine serum albumin

(BSA; Sigma-Aldrich) for 40 min, incubated with primary antibody overnight at 4°C. Next day,

washed with PBST three times, incubated with secondary antibody for 2 hrs at room

temperature, washed with PBST three times, and mounted with Vectashield mounting medium

(H-1000, Vector Laboratories). 40 nephrocytes were analyzed from 6 female flies per

genotype.

For Ref(2)p, ubiquitin (FK2), and PDI immunochemistry, 20-day-old female flies were

dissected and fixated for 10 min in 4% paraformaldehyde in 1xPBS (Thermo Fisher Scientific).

Primary antibodies were diluted in PBST: Rabbit polyclonal antibody against Ref(2)P (1:100)

(ab178440, Abcam); Monoclonal mouse antibody ubiquitin FK2, raised against human

polyubiquitin-B and polyubiquitin-C (1:100) (BML-PW8810, Enzo Life Sciences); Rabbit

polyclonal antibody against protein disulfide isomerase (1:100) (PDI; P7122, Sigma).

Secondary antibody Alexa Fluor 555 (A-21422, Thermo Fisher Scientific) was used at 1:1,000

dilution in PBST. The nephrocytes were washed with PBST three times, blocked in PBST+2%

BSA (Sigma-Aldrich) for 40 min, incubated with primary antibodies overnight at 4°C. Next day,

washed with PBST three times, incubated with secondary antibody for 2 hrs at room

temperature, washed with 1xPBST three times, and mounted with Vectashield mounting

medium (H-1000, Vector Laboratories). For quantitation, 40 nephrocytes were analyzed from

6 female flies per genotype.

Imaging of tagged protein

For Rab5-GFP, Rab7-GFP, and Atg8a-mCherry imaging, 20-day-old adult flies (female) that

expressed the protein of interest with a fluorescent tag in their nephrocytes were dissected in

artificial hemolymph [70 mmol/L NaCl (Carolina), 5 mmol/L KCl (Sigma), 1.5 mmol/L

$CaCl_2.2H_2O$ (Sigma-Aldrich), 4 mmol/L $MgCl_2$ (Sigma-Aldrich), 10 mmol/L $NaHCO_3$ (Sigma-

Aldrich), 5 mmol/L trehalose (Sigma), 115 mmol/L sucrose (Sigma-Aldrich), and 5 mmol/L

HEPES (Sigma-Aldrich), in H_2O]. The nephrocytes were kept in artificial hemolymph and

imaged live.

LysoTracker assay

20-day-old female flies from the *Hand*-GFP and *Dot*-Gal4 transgenic lines expressing UAS-

APOL1-G1 were crossed with UAS-*nef* transgenic flies at 22°C. LysoTracker intensity by

nephrocytes was assessed ex vivo in adult flies by dissecting heart tissues in artificial

hemolymph [70 mmol/l NaCl (Carolina), 5 mmol/l KCl (Sigma), 1.5 mmol/l $CaCl_2 \cdot 2H_2O$ (Sigma-

Alrich), 4 mmol/l $MgCl_2$ (Sigma-Aldrich), 10 mmol/l $NaHCO_3$ (Sigma-Aldrich), 5 mmol/l

trehalose (Sigma), 115 mmol/l sucrose (Sigma-Aldrich), and 5 mmol/l HEPES (Sigma-Aldrich),

in H_2O] and examining cells by fluorescence microscopy after 20 minutes incubation with

LysoTracker (Red DND-99, Thermo Fisher Scientific) used according to manufacturer

instructions and 10 minutes fixation in 4% paraformaldehyde in 1xPBS (Thermo Fisher

Scientific). For quantification, 40 nephrocytes were analyzed from 6 female flies per genotype.

The nephrocytes were mounted with VectaShield mounting medium (H-1000, Vector

Laboratories). For quantification, 40 nephrocytes were analyzed from 6 female flies per

genotype.

TUNEL assay

Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) assay detects DNA

fragmentation by labeling the 3′-hydroxyl termini in the double-strand DNA breaks generated

during apoptosis. 20-day-old female flies were dissected in 100 °C artificial hemolymph [70

mmol/l NaCl (Carolina), 5 mmol/l KCl (Sigma), 1.5 mmol/l $CaCl_2 \cdot 2H_2O$ (Sigma-Alrich), 4 mmol/l

$MgCl_2$ (Sigma-Aldrich), 10 mmol/l $NaHCO_3$ (Sigma-Aldrich), 5 mmol/l trehalose (Sigma), 115

mmol/l sucrose (Sigma-Aldrich), and 5 mmol/l HEPES (Sigma-Aldrich), in H_2O] and heat-fixed

for 20 min in 4% paraformaldehyde in PBS (Thermo Fisher Scientific). The nephrocytes were

blocked with blocking buffer [50 mM Tris-Cl (pH 7.4) (Sigma-Alrich), 0.1% Triton X-100

257 (Sigma), and 188 mM NaCl (Carolina), in H_2O] and incubated with In Situ Cell Death detection

258 kit (Roche) used according to manufacturer instructions. The nephrocytes were mounted with

259 Vectashield mounting medium (H-1000, Vector Laboratories). 40 nephrocytes were analyzed

260 from 6 female flies per genotype.

262 **Confocal imaging**

263 Images for 10 kD dextran and FITC-albumin uptake and LysoTracker assays were obtained

264 using a ZEISS LSM900 microscope with a 20× Plan-Apochromat 0.8 NA air objective (ZEN

265 Blue, edition 3.0, acquisition software). Fluorescent immunochemistry confocal images for

266 anti-Pyd, anti-Ref(2)P, anti-PDI, anti-ubiquitin (anti-Fk2), and TUNEL were obtained using the

267 ZEISS LSM900 confocal microscope with a 63× Plan-Apochromat 1.4 NA oil objective under

268 AiryScan SR mode (ZEN Blue, edition 3.0, acquisition software). Live-cell fluorescence

269 confocal images for the UAS-YFP-*Rab5*, UAS-YFP-*Rab7*, and UAS-mCherry-*Atg8a* fly lines

270 were obtained with a ZEISS LSM900 confocal microscope with a 63× Plan-Apochromat 1.4

271 NA oil objective under AiryScan SR mode (ZEN Blue, edition 3.0, acquisition software). For

272 quantitative comparison of intensities, common settings were chosen to avoid oversaturation

273 and then applied across images for all samples within an assay. ImageJ [33] software (version

274 1.52a) was used for all image processing.

276 **Statistical analysis**

277 Statistical tests were performed using PAST.exe software [Natural History Museum, University

278 of Oslo (UiO), Oslo, Norway]. Data were first tested for normality using the Shapiro-Wilk test

279 ($\alpha=0.05$). Normally distributed data were analyzed by one-way ANOVA followed by Tukey-

280 Kramer post-test for comparing multiple groups. Non-normal distributed data were analyzed

281 Kruskal-Wallis H-test followed by a Dunn's test for comparisons between multiple groups. The

282 results are presented as mean ± s.d. Statistical significance was determined as $P<0.05$.

283 Details for sample size used for quantitation are provided in the figure legends. The statistical

284 tests applied to determine significance are provided in Supplementary File 1.

RESULTS

APOL1-G1-induced nephrocyte dysfunction was exacerbated by HIV-1 Nef

In a previous study, we described the generation of transgenic *Drosophila* lines carrying the human *APOL1-G0* wild-type allele or the *APOL1-G1* risk-allele (RA) derived from a child with HIV-associated nephropathy [15]. Here, to define how HIV-1 Nef affects the structure and function of nephrocytes, we generated additional transgenic fly lines that express HIV-1 *nef* specifically in the nephrocytes and explored its role alone or in combination with *APOL1-G0* or *APOL1-G1*.

To measure functional changes induced by nephrocyte-specific *APOL1* expression together with HIV-1 *nef*, we used an ex vivo functional assay that measures the capacity of dissected nephrocytes to filter and endocytose 10 kD fluorescent dextran or FITC-albumin particles. HIV-1 *nef* or *APOL1-G0* by itself did not show a reduction in 10 kD dextran or FITC-albumin uptake compared to control nephrocytes (Figure 1A, B, and C). Nephrocytes that expressed *nef* and *APOL1* combined (*APOL1-G0+nef* or *APOL1-G1+nef*), showed a significant decreased uptake of 10 kD dextran and FITC-albumin compared to those that express either *nef, APOL1-G0,* or *APOL1-G1* alone (Figure 1A, B and C). The uptake of either 10 kD dextran or FITC-albumin was significantly reduced in *APOL1-G1+nef* Tg flies, when compared to *APOL1-G0+nef* flies (Figure 1A, B and C), thus reflecting the higher toxicity of the *APOL1-G1* risk allele. In summary, these results show that HIV-1 Nef exacerbates APOL1-induced nephrocyte dysfunction.

Combined HIV-1 Nef and APOL1 exacerbated nephrocyte hypertrophy and cell death

Previously, we showed that the expression of *APOL1-G1* in nephrocytes induced progressive hypertrophy and accelerated their death [15]. Here, we confirmed that the expression of APOL-G1 in aged flies (20-day-old) also increased the size of nephrocytes (Figure 1D). In contrast, nephrocytes expressing HIV-1 *nef* or *APOL1-G0* alone, did not show a significant increase in

cell size (Figure 1D), while those that expressed *nef* combined with either *APOL1-G0* or

APOL1-G1 were significantly larger (*APOL1-G0+nef* or *APOL1G1+nef*; Figure 1D).

Furthermore, nephrocytes expressing *nef* and *APOL1-G1* were larger than those expressing

nef and *APOL1-G0* combined (Figure 1D).

Like before [15], we observed reduced nephrocyte numbers in aged (20-day-old) Tg flies

expressing *APOL1-G1* (Figure 1E). However, we did not observe significant changes in the

number of nephrocytes in flies expressing HIV-1 *nef* or *APOL1-G0* alone (Figure 1E). In

contrast, flies expressing *nef* combined with *APOL1-G0* or *APOL1-G1*, showed a significant

reduction in the nephrocyte number when compared to those expressing *APOL1-G0* or

APOL1-G1 alone (Figure 1E). Moreover, nephrocyte numbers were significantly reduced in

flies expressing *nef* and *APOL1-G1* combined, when compared to those expressing both *nef*

and *APOL1-G0 together* (Figure 1E). These results show that HIV-1 Nef combined with

APOL1-G1 further exacerbates the damage and loss of nephrocytes.

HIV-1 Nef and APOL1-G1 combined disrupt the slit diaphragm filtration structure

The slit diaphragm in *Drosophila* is a highly organized structure that plays a critical role

maintaining the filtration function of nephrocytes [25]. To determine whether the changes

induced by HIV-1 Nef in the uptake of 10 kD dextran were accompanied by an altered

nephrocyte filtration structure, we assessed the localization of the slit diaphragm protein

polychaetoid (Pyd - the homolog of human tight junction protein ZO-1) [27]. As expected, control

nephrocytes showed the typical fingerprint-like localization pattern of Pyd on their cell surface

(Figure 2A). In the optical medial view, control healthy nephrocytes also showed the

characteristic continuous circular ring expression pattern of Pyd (Figure 2B). Notably,

nephrocytes that expressed *nef* or *APOL1-G0* alone did not show a visible disruption of Pyd

(Figure 2A and B). However, nephrocytes that expressed *nef* combined with *APOL1-G0* or

APOL1-G1 showed highly disorganized Pyd localization, more so than either APOL-1 risk

variant alone. Moreover, mislocalized Pyd was more evident in nephrocytes that expressed

both *nef* and *APOL1-G1* together, when compared to those expressing combined *nef* and

APOL1-G0 (Figure 2A and B). In agreement with the altered uptake observed in nephrocytes

expressing *nef* and *APOL1* combined (Figure 1), these data demonstrate that both genes

acting together affect the localization of Pyd and disrupt the slit diaphragm filtration structure

of nephrocytes.

Combined HIV-1 Nef and APOL1-G1 exacerbated the detrimental effects on nephrocyte

endocytosis

Previously, we showed that HIV-1 can infect cultured human podocytes via dynamin-

dependent endocytosis in a process that is facilitated by transmembrane TNF-α [5]. To examine

whether HIV-1 Nef and APOL1 interact to regulate this process in fly nephrocytes, we assayed

the expression level of the early and late endosomal markers, Rab5 and Rab7, respectively.

Expression of *nef*, *APOL1-G0* or *APOL1-G1* alone, or combined did not induce significant

changes in the expression levels or localization of Rab5 (Figure 3A and B). In contrast, Rab7

protein expression levels were significantly increased in nephrocytes that expressed *APOL1-*

G1, but not in those expressing *nef* or *APOL1-G0* alone (Figure 3C and D). Nephrocytes that

expressed *nef* combined with *APOL1-G0* or *APOL1-G1* had significantly higher Rab7 protein

expression levels (Figure 3C and D). Moreover, the highest Rab7 protein expression levels

were observed in flies that expressed *nef* and *APOL1-G1* combined, when compared to all

other flies (Figure 3C and D).

We had also shown previously that *APOL1-G1* altered the acidification of organelles and the

lysosomes in fly nephrocytes, which could impair their function [15]. Thus, here we used the

fluorescent LysoTracker dye to examine acidic vacuoles and lysosomes in nephrocytes

(Figure 3E). LysoTracker fluorescence intensity was significantly reduced in nephrocytes that

expressed *APOL1-G1*, but not *nef* or *APOL1-G0* (Figures 3E and F). Nephrocytes that

expressed *nef* combined with *APOL1-G0* or *APOL1-G1* showed significantly lower

LysoTracker fluorescence intensity levels (Figure 3E and F). Moreover, lower LysoTracker

fluorescence intensity was observed in nephrocytes from dual *APOL1-G1+nef* Tg flies, when

compared to any other fly (Figure 3E and F). Together these findings indicate that HIV-1 Nef

exacerbates the changes induced by APOL1-G1 in key endocytic and acidification pathways

that appear to be essential for the normal function of nephrocytes.

HIV-1 Nef enhances APOL1-G1-induced autophagic changes leading to the

accumulation of autophagosomes in fly nephrocytes

APOL1 has been shown to modulate autophagy in various experimental model systems

[14,16,22,29,34–36]. Autophagy-related 8a (Atg8a), the equivalent of human LC3 (officially known as

MAP1LC3A), is a widely used marker for studying autophagy in *Drosophila* because it allows

the visualization of all autophagic structures, including autophagosomes. Nephrocytes that

expressed *nef* alone did not show higher expression levels of Atg8a-mCherry compared to

controls, although those expressing *nef* together with *APOL1-G0* or with *APOL1-G1* showed

significantly higher expression levels of Atg8a-mCherry, with more significant changes in

APOL1-G1+nef dual Tg flies (Figure 4A and B). We also examined the expression levels of

the refractory to sigma P [Ref(2)P] protein, which is a receptor for cargo destined to be

degraded by autophagy [37]. We did not observe significant changes in Ref(2)P expression

levels in nephrocytes that expressed *nef* alone compared to controls (Figure 4C and D). In

contrast, Ref(2)P was significantly increased in nephrocytes that expressed *APOL1-G0* or

APOL1-G1, and these changes were exacerbated in dual Tg flies that expressed both *nef* and

APOL-G1 (Figure 4C and D). In addition, we observed an increase in the accumulation of

ubiquitinylated proteins in nephrocytes expressing *APOL1-G1*, but not *nef* or *APOL1-G0*

compared to controls (Figure 4E and F). Moreover, more ubiquitinylated proteins were

accumulated in the nephrocytes of *APOL1-G1+nef* dual Tg flies when compared to the

APOL1-G0+nef dual Tg flies (Figure 4E and F). Combined these data demonstrate that HIV-

1 Nef markedly increases APOL1-induced autophagy changes in nephrocytes.

HIV-1 Nef and APOL1-G1 induce ER stress in fly nephrocytes

APOL1-G1 has been shown to elicit the endoplasmic reticulum (ER) stress response [23,29], while HIV-1 Nef bound the ER chaperone Calnexin (CNX) to induce ER stress [38–40]. Therefore, we assessed the expression levels of protein disulfide-isomerase (PDI), a marker of ER stress in nephrocytes. We found a significant increase in PDI in nephrocytes that expressed *nef* alone compared to controls (Figure 5A and B). This increase was significantly more substantial when *APOL1* was expressed simultaneously with *nef*; even when compared to the changes detected in *APOL1-G0+nef* dual Tg flies (Figure 5A and B). Furthermore, nephrocytes that expressed *APOL1-G1*, but not *nef* or *APOL1-G0*, alone showed positive results in the TUNEL assay, an indicator of DNA fragmentation and considered a marker of cell death (Figure 5C). Likewise, TUNEL-positive nephrocytes were observed in nephrocytes that expressed combined *APOL1-G0+nef* or *APOL1-G1+nef* (Figure 5C). These findings indicate that HIV-1 Nef alone can induce ER stress in *Drosophila* nephrocytes in an autophagy-independent pathway, and that Nef can also exacerbate the APOL1-G1-induced autophagy-dependent pathway to further stimulates ER stress, leading to nephrocyte dysfunction and cell death.

DISCUSSION

We have developed a new *Drosophila* model system to explore how HIV-1 Nef and APOL1-G1 interact in fly nephrocytes to affect the structure and function of these cells *in vivo*. In brief, we found that HIV-1 Nef acts in a synergistic manner with APOL1-G1 to impair the function of key endocytic, acidification, and autophagic pathways, thereby affecting the normal structure and function of fly nephrocytes. Previous studies showed that HIV-1 Nef can induce ER stress by binding a chaperon protein called Calnexin (CNX) [38–40]. On the other hand, APOL1 disrupts autophagy at the point of endosome-lysosome transition, leading to the accumulation of proteins in the ER and thus promoting ER stress [23,29]. Here, we showed that in addition to causing ER stress by itself through an autophagy-independent pathway (likely by binding to

Calnexin homologs in nephrocytes), Nef also exacerbates the pathogenic effect of APOL1-G1

on the autophagy pathway which also leads to ER stress. This synergistic interactions

between HIV-1 Nef and APOL1-G1 culminate in cellular toxicity and the premature death of

nephrocytes (Figure 6). We anticipate that this new *Drosophila* model will provide a clinically

relevant experimental tool to identify novel genetic interactions through which HIV-1 Nef and

APOL1-G1 may affect the function and survival of human podocytes precipitating HIV-1 and

APOL1-associated kidney diseases.

It is worth noting that APOL1-G1-induced cytotoxicity is not exclusively dependent on the

nucleotide substitutions or deletions that define the G1 risk variant. Other components of the

APOL1 molecule, including the BH3 domain [23,41], exon 4-encoded sequences [34], and different

haplotypes [42], also affect the toxicity of APOL1 in renal cells. Because introducing *APOL1-G1*

mutations to unnatural haplotype backgrounds, e.g., an *APOL1-G0* reference haplotype, could

affect the toxicity of APOL1, we generated our Tg flies that express a common natural *APOL1-*

G1 derived from podocytes cultured from a child with HIVAN [11]. This APOL1-G1 contains the

haplotypes (E150; I228; K255) and only differs at S342 and I384 from a common natural

APOL1-G0 haplotype used to generate the control flies. Notably, this APOL1-G0 control

haplotype is more toxic in cultured human kidney epithelial cells (HEK 293) than the reference

APOL1-G0 haplotype used in other studies [11,16,22,43–49]. Indeed, we previously found that the

ectopic expression of *APOL1-G0* (reference allele) in fly nephrocytes, also induced

nephrocyte toxicity, albeit less severe compared to the *APOL1-G1* (risk allele) [15,29].

Subsequently, we showed that reducing the expression levels of *APOL1-G0* in fly nephrocytes

by lowering the temperature to 22°C, eliminated *APOL1-G0* toxicity [30], hence in the current

study all fly experiments were conducted at 22°C (instead of the conventional 25°C) so that

APOL1-G1, but not *APOL1-G0*, remains toxic in nephrocytes. In addition, we found that Nef

together with APOL1-G0 also affected the function of nephrocytes, although these changes

were less severe when compared to nephrocytes that expressed Nef combined with APOL1-

G1. Overall, these findings agree with clinical studies that showed that people of West African

genetic ancestry living with a high HIV-1 viral load can develop HIVAN even if they do not

carry an *APOL1-RA*; however, those that carry a risk allele are at much higher risk [4].

Interestingly, one study in *APOL-1* Tg mice, with expression driven by the podocyte-specific

promoter of *nephrin* (*Nphs1*), crossbred with HIV-Tg26 mice, unexpectedly found that the

APOL1 wild-type allele (*G0*) ameliorated the podocyte loss characteristic of HIVAN, while

APOL1-G2 did not induce significant renal changes [18]. This study suggested that APOL1-G0

could play a protective role in the pathogenesis of HIVAN. However, subsequent studies in

APOL1 Tg mice developed with a coisogenic bacterial artificial chromosome (BAC) harboring

either wild-type (*G0*), *G1* or *G2* human *APOL1*, showed that the renal disease phenotype was

greater in *G2/G2* transgenic mice, compared to *G1/G1*, and that *APOL1-G0* did not rescue

APOL1-G1 or *APOL1-G2* induced kidney toxicity after these mice were injected with

Interferon-gamma (INF-γ) [17]. Likewise, in a previous study, we found that *APOL1-G2* Tg flies

developed more significant nephrocyte injury compared to the *APOL1-G1* Tg flies [29]. These

studies also confirmed that the renal toxicity of APOL1-RA is mainly a function of the APOL1

expression level in renal cells [17], explaining why some Tg mice with *APOL1* driven by

podocyte-specific promoters develop more significant renal injury than others [6,14,17,50]. Our

current data from dual HIV-1 *nef* + *APOL1-G0* Tg flies, do not support the hypothesis that G0

offers protection against HIV-1 Nef, since Nef combined with APOL1-G0 induced significant

nephrocyte dysfunction. A recent study on bioRxiv used dual BAC/*APOL1*-HIV-Tg26 mice, in

which the *APOL1* transgene is a 47 kb human cDNA that contains the sequence from

chromosome 22 within a BAC [51]. It likewise showed that *APOL1-G1* aggravated the renal

injury in HIV-Tg26 mice, and that *APOL1-G0* did not prevent or minimize HIVAN in these mice

[51]. Taken together, our findings suggest that HIV-1 Nef per se is an independent risk factor

and potential second hit for developing APOL1 nephropathies.

We found that Nef together with APOL1-G1, but not alone, disrupted the autophagy pathway

and increased the accumulation of Atg8a-positive autophagosomes in fly nephrocytes. In line

with these findings, several studies have shown that HIV-1 can restrict autophagy to promote

viral replication by inhibiting the fusion between autophagosomes and lysosomes [35,52–54]. For

example, Nef overexpression in human astrocytes led to increased ATG8/LC3 [53]. Likewise,

Nef caused a significant acidification defect in human and macaque cardiomyocytes, altering

the subcellular localization of Rab7, and inducing cardiotoxicity through dysregulation of

autophagy [54]. Alternatively, in several human cell lines, HIV-1 Nef reduced the fusion of

autophagosomes to lysosomes by interacting with Beclin1 (BECN1)/Rab7 and dysregulating

the activity of the transcription factor EB (TFEB), which plays a key role in lysosomal

biogenesis [55,56]. Together, these findings show that HIV-1 Nef mediated inhibition of

autophagic flux is conserved across cell types and species. We did not observe visible

changes in the structure and function of nephrocytes that expressed HIV-1 Nef alone. It is

possible that increasing the expression of Nef would lead to a mild phenotype, as previous

studies in podocyte-specific HIV-1 *nef* Tg mice have shown mild podocyte injury or de-

differentiation changes [9,57]. Notably, one study showed much more severe podocyte injury

when HIV-1 *nef* and *vpr* were expressed simultaneously in mouse podocytes, indicating a

synergistic interaction between these HIV-1 genes [9]. These findings suggest that other HIV-1

genes and cytokines are needed to evoke the complete phenotype of HIVAN in human

podocytes. On the other hand, Tg mice with HIV-1 *nef* driven by the CD4 promoter developed

kidney disease [12,58,59]. In this case, the podocytes did not express HIV-1 genes and showed

only mild changes, while renal infiltrating mononuclear cells that expressed HIV-1 *nef* showed

significant renal interstitial injury [12,58,59]. Nef, however, can be released by CD+ HIV-1 infected

cells predominately as a component of exosomes [60] and can also be taken up by uninfected

cells [61]. Therefore, the renal recruitment of CD4+ cells may facilitate the transfer of Nef to renal

epithelial cells.

Defective endosomal trafficking has been reported in yeast and flies that expressed *APOL1-*

G1 [22]. Here, we demonstrated that APOL1-G1 impairs late endosome to lysosome transition

(Rab5 and Rab7) and thus the autophagy pathway [Atg8 and Ref(2)P]. Furthermore, Nef

enhanced the autophagic changes induced by APOL1-G1. Likewise, immunofluorescence

studies in cultured human podocytes derived from low-risk (*G0/G0*) and high-risk (*G1/G2*)

allele subjects revealed increased Rab7 (late endosomes) and LC3II (autophagosomes)

positive intracellular organelles in high-risk compared to low-risk allele podocytes [14]. Studies

in inducible podocyte-specific *APOL1-RA* transgenic mice also showed that G1 interferes with

intracellular vesicular trafficking by impairing endocytic, autophagic, and acidification

pathways, as well as by disrupting the maturation of autophagosomes and autophagic flux,

which correlated with the development of FSGS [14]. Indeed, both *APOL-1* risk alleles have

been shown to affect autophagy in cultured human podocytes [62,63].

Taken together, these studies suggest that the new *APOL1* and HIV-1 *nef* Tg fly models can

generate clinically relevant information for humans. However, the *APOL1+nef* Tg fly models

have several limitations as well. Firstly, the models are based on the ectopic overexpression

of *APOL1* haplotypes, which can affect APOL1 intracellular localization and cytotoxicity [7, 8].

Overexpression also bypasses the infection process as well as the effects of circulating

cytokines released by HIV-1 infected cells (e.g., TNF-α, INF-γ), which can act as a second hit.

On the other hand, our model permits the exploration of HIV-1 Nef and APOL1 interactions

independently of these confounding variables. Secondly, we explored only the interactions

between the APOL1-G1 and HIV-1 Nef, while other HIV-1 proteins (e.g., Vpr or Tat) acting

either independently or in synergy with Nef or APOL1 can play relevant roles the

pathomechanism underlying HIVAN. Thirdly, HIVAN is a complex disease, involving other

renal glomerular and tubular cell types besides podocytes, as well as lesions that cannot be

studied in flies (e.g., collapsing glomerulopathy). Nevertheless, based on the notion that the

relevant pathways are conserved in human podocytes, flies provide a unique opportunity to

perform rapid and cost-effective genetic screenings to uncover new genetic interactions and

pathways that are modulated by HIV-1 and APOL1.

DISCLOSURES

The authors declare no competing interests.

FUNDING

This work was supported by National Institutes of Health grants R01-DK115968 (P.E.R. and Z.H.), R01-DK103564 (P.E.R.), R01-DK120908 (Z.H.), and R01-DK098410 (Z.H.).

ACKNOWLEDGMENTS

We thank the Bloomington Drosophila Stock Center (BDSC) based at Indiana University (Bloomington, IN) for the *Drosophila* stocks; and the Developmental Studies Hybridoma Bank (DSHB) based at the University of Iowa (Iowa City, IA) for providing the antibodies.

AUTHOR CONTRIBUTIONS

J.Z. and Z.H. designed the study; J.Z., Y.F., Y.Y. and JL. carried out the experiments; J.Z., J.vdL., and Z.H. analyzed and interpreted the data; J.Z. prepared the figures; J.Z, J.vdL., P.R., and Z.H. drafted and revised the manuscript; the manuscript has been critically reviewed and the final version approved by all authors.

REFERENCES

1. Ross MJ: Advances in the pathogenesis of HIV-associated kidney diseases. *Kidney Int.* 86: 266–274, 2014

2. Swanepoel CR, Atta MG, D'Agati VD, Estrella MM, Fogo AB, Naicker S, et al.: Kidney disease in the setting of HIV infection: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney Int.* 93: 545–559, 2018

3. Kopp JB, Nelson GW, Sampath K, Johnson RC, Genovese G, An P, et al.: APOL1 genetic variants in focal segmental glomerulosclerosis and HIV-associated nephropathy. *J. Am. Soc. Nephrol.* 22: 2129–2137, 2011

4. Kasembeli AN, Duarte R, Ramsay M, Mosiane P, Dickens C, Dix-Peek T, et al.: APOL1 risk variants are strongly associated with HIV-associated nephropathy in black South Africans. *J. Am. Soc. Nephrol.* 26: 2882–2890, 2015

5. Li J, Das JR, Tang P, Han Z, Jaiswal JK, Ray PE: Transmembrane TNF-α Facilitates HIV-1 Infection of Podocytes Cultured from Children with HIV-Associated Nephropathy. *J. Am. Soc. Nephrol.* 28: 862–875, 2017

6. Yoshida T, Latt KZ, Rosenberg AZ, Shrivastav S, Heymann J, Halushka MK, et al.: Transcriptomic Analysis of Human Podocytes In Vitro: Effects of Differentiation and APOL1 Genotype. *Kidney Int Rep* 8: 164–178, 2023

7. Kajiyama W, Kopp JB, Marinos NJ, Klotman PE, Dickie P: Glomerulosclerosis and viral gene expression in HIV-transgenic mice: role of nef. *Kidney Int.* 58: 1148–1159, 2000

8. Ray PE, Liu X-H, Robinson LR, Reid W, Xu L, Owens JW, et al.: A novel HIV-1 transgenic rat model of childhood HIV-1-associated nephropathy. *Kidney Int.* 63: 2242–2253, 2003

9. Zuo Y, Matsusaka T, Zhong J, Ma J, Ma L-J, Hanna Z, et al.: HIV-1 genes vpr and nef synergistically damage podocytes, leading to glomerulosclerosis. *J. Am. Soc. Nephrol.* 17: 2832–2843, 2006

10. Lu T-C, He JC, Wang Z-H, Feng X, Fukumi-Tominaga T, Chen N, et al.: HIV-1 Nef disrupts the podocyte actin cytoskeleton by interacting with diaphanous interacting protein. *J. Biol. Chem.* 283: 8173–8182, 2008

11. Xie X, Colberg-Poley AM, Das JR, Li J, Zhang A, Tang P, et al.: The basic domain of HIV-tat transactivating protein is essential for its targeting to lipid rafts and regulating fibroblast growth factor-2 signaling in podocytes isolated from children with HIV-1-associated nephropathy. *J. Am. Soc. Nephrol.* 25: 1800–1813, 2014

12. Hu C, Priceputu E, Cool M, Chrobak P, Bouchard N, Forestier C, et al.: NEF-Induced HIV-
Associated Nephropathy Through HCK/LYN Tyrosine Kinases. *Am. J. Pathol.* 193: 702–
724, 2023

13. Gharavi AG, Ahmad T, Wong RD, Hooshyar R, Vaughn J, Oller S, et al.: Mapping a locus
for susceptibility to HIV-1-associated nephropathy to mouse chromosome 3. *Proc. Natl.*
Acad. Sci. U. S. A. 101: 2488–2493, 2004

14. Beckerman P, Bi-Karchin J, Park ASD, Qiu C, Dummer PD, Soomro I, et al.: Transgenic
expression of human APOL1 risk variants in podocytes induces kidney disease in mice.
Nat. Med. 23: 429–438, 2017

15. Fu Y, Zhu J-Y, Richman A, Zhang Y, Xie X, Das JR, et al.: APOL1-G1 in Nephrocytes
Induces Hypertrophy and Accelerates Cell Death. *J. Am. Soc. Nephrol.* 28: 1106–1116,
2017

16. O'Toole JF, Schilling W, Kunze D, Madhavan SM, Konieczkowski M, Gu Y, et al.: ApoL1
Overexpression Drives Variant-Independent Cytotoxicity. *J. Am. Soc. Nephrol.* 29: 869–
879, 2018

17. McCarthy GM, Blasio A, Donovan OG, Schaller LB, Bock-Hughes A, Magraner JM, et al.:
Recessive, gain-of-function toxicity in an APOL1 BAC transgenic mouse model mirrors
human APOL1 kidney disease. *Dis. Model. Mech.* [Internet] 14: 2021 Available from:
http://dx.doi.org/10.1242/dmm.048952

18. Bruggeman LA, Wu Z, Luo L, Madhavan S, Drawz PE, Thomas DB, et al.: APOL1-G0
protects podocytes in a mouse model of HIV-associated nephropathy. *PLoS One* 14:
e0224408, 2019

19. Zhang F, Zhao Y, Han Z: An in vivo functional analysis system for renal gene discovery
in Drosophila pericardial nephrocytes. *J. Am. Soc. Nephrol.* 24: 191–197, 2013

20. Fu Y, Zhu J-Y, Richman A, Zhao Z, Zhang F, Ray PE, et al.: A Drosophila model system
to assess the function of human monogenic podocyte mutations that cause nephrotic
syndrome. *Hum. Mol. Genet.* 26: 768–780, 2017

21. Hermle T, Braun DA, Helmstädter M, Huber TB, Hildebrandt F: Modeling Monogenic
Human Nephrotic Syndrome in the Drosophila Garland Cell Nephrocyte. *J. Am. Soc.*
Nephrol. 28: 1521–1533, 2017

22. Kruzel-Davila E, Shemer R, Ofir A, Bavli-Kertselli I, Darlyuk-Saadon I, Oren-Giladi P, et
al.: APOL1-Mediated Cell Injury Involves Disruption of Conserved Trafficking Processes.
J. Am. Soc. Nephrol. 28: 1117–1130, 2017

23. Gerstner L, Chen M, Kampf LL, Milosavljevic J, Lang K, Schneider R, et al.: Inhibition of
endoplasmic reticulum stress signaling rescues cytotoxicity of human apolipoprotein-L1

risk variants in Drosophila. *Kidney Int.* [Internet] 2022 Available from:
http://dx.doi.org/10.1016/j.kint.2021.12.031

24. Rani L, Gautam NK: Drosophila Renal Organ as a Model for Identification of Targets and
Screening of Potential Therapeutic Agents for Diabetic Nephropathy. *Curr. Drug Targets*
19: 1980–1990, 2018

25. Weavers H, Prieto-Sánchez S, Grawe F, Garcia-López A, Artero R, Wilsch-Bräuninger
M, et al.: The insect nephrocyte is a podocyte-like cell with a filtration slit diaphragm.
Nature 457: 322–326, 2009

26. Zhuang S, Shao H, Guo F, Trimble R, Pearce E, Abmayr SM: Sns and Kirre, the
Drosophila orthologs of Nephrin and Neph1, direct adhesion, fusion and formation of a
slit diaphragm-like structure in insect nephrocytes. *Development* 136: 2335–2344, 2009

27. Wang L, Wen P, van de Leemput J, Zhao Z, Han Z: Slit diaphragm maintenance requires
dynamic clathrin-mediated endocytosis facilitated by AP-2, Lap, Aux and Hsc70-4 in
nephrocytes. *Cell Biosci.* 11: 83, 2021

28. Lang K, Milosavljevic J, Heinkele H, Chen M, Gerstner L, Spitz D, et al.: Selective
endocytosis controls slit diaphragm maintenance and dynamics in Drosophila
nephrocytes. *Elife* [Internet] 11: 2022 Available from:
http://dx.doi.org/10.7554/eLife.79037

29. Zhu J-Y, Lee J-G, Fu Y, van de Leemput J, Ray PE, Han Z: APOL1-G2 accelerates
nephrocyte cell death by inhibiting the autophagy pathway. *Dis. Model. Mech.* [Internet]
2023 Available from: http://dx.doi.org/10.1242/dmm.050223

30. Lee J-G, Fu Y, Zhu J-Y, Wen P, van de Leemput J, Ray PE, et al.: A SNARE protective
pool antagonizes APOL1 renal toxicity in Drosophila nephrocytes. *Cell Biosci.* 13: 199,
2023

31. Han Z, Olson EN: Hand is a direct target of Tinman and GATA factors during Drosophila
cardiogenesis and hematopoiesis. *Development* 132: 3525–3536, 2005

32. Murakawa GJ, Zaia JA, Spallone PA, Stephens DA, Kaplan BE, Wallace RB, et al.: Direct
detection of HIV-1 RNA from AIDS and ARC patient samples. *DNA* 7: 287–295, 1988

33. Schneider CA, Rasband WS, Eliceiri KW: NIH Image to ImageJ: 25 years of image
analysis. *Nat. Methods* 9: 671–675, 2012

34. Khatua AK, Cheatham AM, Kruzel ED, Singhal PC, Skorecki K, Popik W: Exon 4-encoded
sequence is a major determinant of cytotoxicity of apolipoprotein L1. *Am. J. Physiol. Cell*
Physiol. 309: C22-37, 2015

35.   Chang C, Young LN, Morris KL, von Bülow S, Schöneberg J, Yamamoto-Imoto H, et al.:
667   Bidirectional Control of Autophagy by BECN1 BARA Domain Dynamics. *Mol. Cell* 73:
668   339-353.e6, 2019

669   36.   Chun J, Zhang J-Y, Wilkins MS, Subramanian B, Riella C, Magraner JM, et al.:
670   Recruitment of APOL1 kidney disease risk variants to lipid droplets attenuates cell toxicity.
671   *Proc. Natl. Acad. Sci. U. S. A.* 116: 3712–3721, 2019

672   37.   Gatica D, Lahiri V, Klionsky DJ: Cargo recognition and degradation by selective
673   autophagy. *Nat. Cell Biol.* 20: 233–242, 2018

674   38.   Jennelle L, Hunegnaw R, Dubrovsky L, Pushkarsky T, Fitzgerald ML, Sviridov D, et al.:
675   HIV-1 protein Nef inhibits activity of ATP-binding cassette transporter A1 by targeting
676   endoplasmic reticulum chaperone calnexin. *J. Biol. Chem.* 289: 28870–28884, 2014

677   39.   Ryan D, Carberry S, Murphy ÁC, Lindner AU, Fay J, Hector S, et al.: Calnexin, an ER-
678   induced protein, is a prognostic marker and potential therapeutic target in colorectal
679   cancer.   *J.   Transl.   Med.*   [Internet]   14:   2016   Available   from:
680   http://dx.doi.org/10.1186/s12967-016-0948-z

681   40.   Guérin R, Arseneault G, Dumont S, Rokeach LA: Calnexin is involved in apoptosis
682   induced by endoplasmic reticulum stress in the fission yeast. *Mol. Biol. Cell* 19: 4404–
683   4420, 2008

684   41.   Heneghan JF, Vandorpe DH, Shmukler BE, Giovinazzo JA, Raper J, Friedman DJ, et al.:
685   BH3 domain-independent apolipoprotein L1 toxicity rescued by BCL2 prosurvival
686   proteins. *Am. J. Physiol. Cell Physiol.* 309: C332-47, 2015

687   42.   Lannon H, Shah SS, Dias L, Blackler D, Alper SL, Pollak MR, et al.: Apolipoprotein L1
688   (APOL1) risk variant toxicity depends on the haplotype background. *Kidney Int.* 96: 1303–
689   1307, 2019

690   43.   Lan X, Jhaveri A, Cheng K, Wen H, Saleem MA, Mathieson PW, et al.: APOL1 risk
691   variants enhance podocyte necrosis through compromising lysosomal membrane
692   permeability. *Am. J. Physiol. Renal Physiol.* 307: F326-36, 2014

693   44.   Nichols B, Jog P, Lee JH, Blackler D, Wilmot M, D'Agati V, et al.: Innate immunity
694   pathways regulate the nephropathy gene Apolipoprotein L1. *Kidney Int.* 87: 332–342,
695   2015

696   45.   Olabisi OA, Zhang J-Y, VerPlank L, Zahler N, DiBartolo S 3rd, Heneghan JF, et al.:
697   APOL1 kidney disease risk variants cause cytotoxicity by depleting cellular potassium
698   and inducing stress-activated protein kinases. *Proc. Natl. Acad. Sci. U. S. A.* 113: 830–
699   837, 2016

46. Granado D, Müller D, Krausel V, Kruzel-Davila E, Schuberth C, Eschborn M, et al.: Intracellular APOL1 Risk Variants Cause Cytotoxicity Accompanied by Energy Depletion. *J. Am. Soc. Nephrol.* 28: 3227–3238, 2017

47. Hayek SS, Koh KH, Grams ME, Wei C, Ko Y-A, Li J, et al.: A tripartite complex of suPAR, APOL1 risk variants and αvβ3 integrin on podocytes mediates chronic kidney disease. *Nat. Med.* 23: 945–953, 2017

48. Ma L, Chou JW, Snipes JA, Bharadwaj MS, Craddock AL, Cheng D, et al.: APOL1 Renal-Risk Variants Induce Mitochondrial Dysfunction. *J. Am. Soc. Nephrol.* 28: 1093–1105, 2017

49. Wen H, Kumar V, Lan X, Shoshtari SSM, Eng JM, Zhou X, et al.: APOL1 risk variants cause podocytes injury through enhancing endoplasmic reticulum stress. *Biosci. Rep.* [Internet] 38: 2018 Available from: http://dx.doi.org/10.1042/BSR20171713

50. Bruggeman LA, Wu Z, Luo L, Madhavan SM, Konieczkowski M, Drawz PE, et al.: APOL1-G0 or APOL1-G2 Transgenic Models Develop Preeclampsia but Not Kidney Disease. *J. Am. Soc. Nephrol.* 27: 3600–3610, 2016

51. Yoshida T, Latt KZ, Santo BA, Shrivastav S, Zhao Y, Fenaroli P, et al.: APOL1 kidney risk variants in glomerular diseases modeled in transgenic mice. *bioRxiv* [Internet] 2023 Available from: http://dx.doi.org/10.1101/2023.03.27.534273

52. Campbell GR, Rawat P, Bruckman RS, Spector SA: Human Immunodeficiency Virus Type 1 Nef Inhibits Autophagy through Transcription Factor EB Sequestration. *PLoS Pathog.* 11: e1005018, 2015

53. Saribas AS, Khalili K, Sariyer IK: Dysregulation of autophagy by HIV-1 Nef in human astrocytes. *Cell Cycle* 14: 2899–2904, 2015

54. Gupta MK, Kaminski R, Mullen B, Gordon J, Burdo TH, Cheung JY, et al.: HIV-1 Nef-induced cardiotoxicity through dysregulation of autophagy. *Sci. Rep.* 7: 8572, 2017

55. Franco-Juárez B, Coronel-Cruz C, Hernández-Ochoa B, Gómez-Manzo S, Cárdenas-Rodríguez N, Arreguin-Espinosa R, et al.: TFEB; Beyond Its Role as an Autophagy and Lysosomes Regulator. *Cells* [Internet] 11: 2022 Available from: http://dx.doi.org/10.3390/cells11193153

56. Zhang W, Li X, Wang S, Chen Y, Liu H: Regulation of TFEB activity and its potential as a therapeutic target against kidney diseases. *Cell Death Discov* 6: 32, 2020

57. Husain M, D'Agati VD, He JC, Klotman ME, Klotman PE: HIV-1 Nef induces dedifferentiation of podocytes in vivo: a characteristic feature of HIVAN. *AIDS* 19: 1975–1980, 2005

58. Hanna Z, Kay DG, Rebai N, Guimond A, Jothy S, Jolicoeur P: Nef harbors a major
determinant of pathogenicity for an AIDS-like disease induced by HIV-1 in transgenic
mice. *Cell* 95: 163–175, 1998

59. Hanna Z, Kay DG, Cool M, Jothy S, Rebai N, Jolicoeur P: Transgenic mice expressing
human immunodeficiency virus type 1 in immune cells develop a severe AIDS-like
disease. *J. Virol.* 72: 121–132, 1998

60. Dubrovsky L, Brichacek B, Prashant NM, Pushkarsky T, Mukhamedova N, Fleetwood AJ,
et al.: Extracellular vesicles carrying HIV-1 Nef induce long-term hyperreactivity of
myeloid cells. *Cell Rep.* 41: 111674, 2022

61. Mukhamedova N, Hoang A, Dragoljevic D, Dubrovsky L, Pushkarsky T, Low H, et al.:
Exosomes containing HIV protein Nef reorganize lipid rafts potentiating inflammatory
response in bystander cells. *PLoS Pathog.* 15: e1007907, 2019

62. Kumar V, Ayasolla K, Jha A, Mishra A, Vashistha H, Lan X, et al.: Disrupted apolipoprotein
L1-miR193a axis dedifferentiates podocytes through autophagy blockade in an APOL1
risk milieu. *Am. J. Physiol. Cell Physiol.* 317: C209–C225, 2019

63. Ekulu PM, Adebayo OC, Decuypere J-P, Bellucci L, Elmonem MA, Nkoy AB, et al.: Novel
human podocyte cell model carrying G2/G2 APOL1 high-risk genotype. *Cells* 10: 1914,
2021

 FIGURES

Figure 1. Expression of HIV-1 protein Nef facilitated nephrocyte function reduction,

hypertrophy, and cell death due to APOL1-G1.

(A-E) Flies used (20-day-old adult females): Control (*Dot>w[1118]*); *nef* (*Dot>nef*-OE); *APOL1-*

G0 (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>APOL1-*

G0-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>APOL1-G1*-OE+*nef*-OE). **(A) Top:** 10kD fluorescent

dextran particle uptake (red) by nephrocytes using nephrocyte specific driver *Dot*-Gal4 to

express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Scale bar: 15

μm. **Bottom:** FITC-albumin particle uptake (green) by nephrocytes using nephrocyte

specifically driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and

APOL1-G1 at 22°C. Scale bar: 15 μm. **(B)** Quantitation of 10kD dextran uptake, relative to

uptake in control flies. n=40 nephrocytes from 6 flies, per group. **(C)** Quantitation of FITC-

albumin uptake, relative to uptake in control flies. n=40 nephrocytes from 6 flies, per group.

(D) Quantitation of adult nephrocyte size, relative to size in control flies. n=40 nephrocytes

from 6 flies, per group. **(E)** Quantitation of adult nephrocyte number, relative to cell number in

control flies. n=10 flies, per group. Results have been presented as mean ± s.d., normalized

to the control group. Kruskal–Wallis H-test followed by a Dunn's test; statistical significance:

*P<0.05, **P<0.01, ***P<0.001.

Figure 1

Figure 2. The expression of HIV-1 protein Nef in nephrocytes facilitated the disruption of the slit diaphragm due to APOL1-G1.

(A, B) Flies used (20-day-old adult females): Control (*Dot>w[1118]*); *nef* (*Dot>nef*-OE); *APOL1-G0* (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>APOL1-G1*-OE+*nef*-OE). **(A, B)** Localization of slit diaphragm protein Polychaetoid (Pyd; red) at the surface (A) and medial (B) optical sections in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 µm; Scale bar (magnified): 1 µm.

Figure 2

Figure 3. The expression of HIV-1 protein Nef in nephrocytes facilitated the disruption in endocytic membrane trafficking due to APOL1-G1.

(A, B) Flies used (20-day-old adult females): Control (*Dot>Rab5*-YFP); *nef* (*Dot>Rab5*-YFP +*nef*-OE); *APOL1-G0* (*Dot>Rab5*-YFP+*APOL1-G0*-OE); *APOL1-G1* (*Dot>Rab5*-YFP+*APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>Rab5*-YFP+*APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>Rab5*-YFP+*APOL1-G1*-OE+*nef*-OE). **(C, D)** Flies used (20-day-old adult females): Control (*Dot>Rab7*-YFP); *nef* (*Dot>Rab7*-YFP +*nef*-OE); *APOL1-G0* (*Dot>Rab7*-YFP+*APOL1-G0*-OE); *APOL1-G1* (*Dot>Rab7*-YFP+*APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>Rab7*-YFP+*APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>Rab7*-YFP+*APOL1-G1*-OE+*nef*-OE). **(E, F)** Flies used (20-day-old adult females): Control (*Hand*-GFP, *Dot>*GFP+*w*[1118]); *nef* (*Hand*-GFP, *Dot>nef*-OE); *APOL1-G0* (*Hand*-GFP, *Dot>APOL1-G0*-OE); *APOL1-G1* (*Hand*-GFP, *Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Hand*-GFP, *Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Hand*-GFP, *Dot>APOL1-G1*-OE+*nef*-OE). **(A)** Expression of endocytosis-related proteins Rab5 (green, early endosome) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 μm; Scale bar (magnified): 1 μm. **(B)** Quantitation of Rab5 fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per group. Note: None of the comparisons between groups were significant. **(C)** Expression of endocytosis-related proteins Rab7 (green, late endosome) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 μm; Scale bar (magnified): 1 μm. **(D)** Quantitation of Rab7 fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per group. **(E)** LysoTracker dye fluorescence level (red) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. *Hand*-GFP transgene expression was visualized as green fluorescence concentrated in the nuclei of nephrocytes. Scale bar: 15 μm. **(F)** Quantitation of LysoTracker dye fluorescence intensity, relative to fluorescence in control

903   nephrocytes. n=40 nephrocytes from 6 flies, per group. Results have been presented as mean

904   ± s.d., normalized to the control group. Kruskal–Wallis H-test followed by a Dunn's test;

905   statistical significance: *P<0.05, ***P<0.001.

Figure 3

Figure 4. The expression of HIV-1 protein Nef in nephrocytes facilitated autophagy impairment due to APOL1-G1.

(A, B) Flies used (20-day-old adult females): Control (*Dot>Atg8a*-mCherry); *nef* (*Dot>Atg8a*-mCherry+*nef*-OE); *APOL1-G0* (*Dot>Atg8a*-mCherry+*APOL1-G0*-OE); *APOL1-G1* (*Dot>Atg8a*-mCherry+*APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>Atg8a*-mCherry+*APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>Atg8a*-mCherry +*APOL1-G1*-OE+*nef*-OE). **(C-F)** Flies used (20-day-old adult females): Control (*Dot>w*[1118]); *nef* (*Dot>nef*-OE); *APOL1-G0* (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>APOL1-G1*-OE+*nef*-OE). **(A)** Expression of autophagy-related protein 8a (Atg8a; red, mCherry) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 µm; Scale bar (magnified): 1 µm. **(B)** Quantitation of Atg8a fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per group. **(C)** Autophagy receptor refractory to sigma P (Ref(2)P; red) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 µm; Scale bar (magnified): 1 µm. **(D)** Quantitation of Ref(2)P fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per group. **(E)** Expression of ubiquitinylated proteins (green) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Boxed areas are shown magnified next to each image. Scale bar: 5 µm; Scale bar (magnified): 1 µm. **(D)** Quantitation of ubiquitinylated proteins fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per group. Results have been presented as mean ± s.d., normalized to the control group. Kruskal–Wallis H-test followed by a Dunn's test; statistical significance: *P<0.05, **P<0.01, ***P<0.001.

Figure 4

Figure 5. Expression of HIV-1 protein Nef in nephrocyte-induced ER stress.

(A, B) Flies used (20-day-old adult females): Control (*Dot>w[1118]*); *nef* (*Dot>nef*-OE); *APOL1-G0* (*Dot>APOL1-G0*-OE); *APOL1-G1* (*Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Dot>APOL1-G1*-OE+*nef*-OE). **(C, D)** Flies used (20-day-old adult females): Control (*Hand*-GFP, *Dot*>GFP+*w[1118]*); *nef* (*Hand*-GFP*, Dot>nef*-OE); *APOL1-G0* (*Hand*-GFP*, Dot>APOL1-G0*-OE); *APOL1-G1* (*Hand*-GFP*, Dot>APOL1-G1*-OE); *APOL1-G0+nef* (*Hand*-GFP, *Dot>APOL1-G0*-OE+*nef*-OE); *APOL1-G1+nef* (*Hand*-GFP*, Dot>APOL1-G1*-OE+*nef*-OE). **(A)** ER stress marker protein disulfide isomerase (PDI; red) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. Scale bar: 15 µm. **(B)** Quantitation of PDI fluorescence intensity, relative to fluorescence in control nephrocytes. n=40 nephrocytes from 6 flies, per group. **(C)** Cell apoptosis marker TUNEL (red) in nephrocytes using the nephrocyte specific driver *Dot*-Gal4 to express HIV-1 *nef* alone or together with *APOL1-G0* and *APOL1-G1* at 22°C. *Hand*-GFP transgene expression was visualized as green fluorescence concentrated in the nuclei of nephrocytes. Scale bar: 5 µm.

Figure 5

Figure 6. Model of HIV-1 Nef acting in synergy with APOL1-G1 through ER stress.
Graphic depiction of the proposed model by which HIV-1 Nef induces ER stress, which in turn leads to cellular toxicity. Via a separate pathway, Nef exacerbates APOL1-G1-mediated ER stress. APOL1-G1 disrupts the late endosomes and lysosomes, which affects the fusion of autophagosomes and lysosomes, the reduced autophagy results in protein accumulation, leading to ER stress. Thus, when together HIV-1 Nef and APOL1-G1 significantly increase cellular toxicity through ER stress, which ultimately leads to cell death and tissue damage in the kidneys. (Green text indicates data showed upregulation, whereas red text indicates data showed downregulation, yellow text indicates no change.) APOL1-G1, Apolipoprotein L1-risk allele G1; Atg8a-mCherry, Autophagy-related 8a with mCherry fluorescent tag; ER, endoplasmic reticulum; PDI, protein disulfide isomerase; Rab5/7-YFP, Rab5 or Rab7 with yellow fluorescent protein tag; Ref(2)p, refractory to sigma P; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling assay.

Figure 6