The auditory midbrain mediates tactile vibration sensing

Erica L. Huey[1,2], Josef Turecek[1,2], Michelle M. Delisle[1,2], Ofer Mazor[1], Gabriel E. Romero[1],

Malvika Dua[1,2], Zoe K. Sarafis[1,2], Alexis Hobble[1,2], Kevin T. Booth[1], Lisa V. Goodrich[1], David

P. Corey[1], and David D. Ginty[1,2]

[1]Department of Neurobiology, [2]Howard Hughes Medical Institute, Harvard Medical School,

220 Longwood Avenue, Boston, MA 02115

Correspondence: david_ginty@hms.harvard.edu

Summary

Vibrations are ubiquitous in nature, shaping behavior across the animal kingdom. For mammals, sound waves traveling through air are captured by the cochlea and the neural signals they generate are encoded in the auditory system[1]. Mechanical vibrations acting on the body are detected by mechanoreceptors of the skin and deep tissue and processed by the somatosensory system[2]. As such, the neural pathways that underlie perception and reaction to sound waves and mechanical vibrations are believed to be anatomically distinct and functionally independent. Here, we report that mechanical vibrations are prominently encoded by neurons in the lateral cortex of the inferior colliculus (LCIC) of the auditory midbrain. LCIC responses to environmental vibrations are mediated by Aβ rapidly adapting (RA)-LTMRs associated with Pacinian corpuscles, which are exquisitely sensitive and unique in their ability to entrain to high - frequency (40-1000 Hz) environmental vibrations[2-5]. Remarkably, most LCIC neurons are dual somatosensory and auditory environmental vibration detectors. Moreover, the LCIC is required for behavioral responses to high frequency mechanical vibrations, but not other somatosensory modalities. Thus, environmental vibrations captured by Pacinian corpuscles of the body are encoded in the auditory midbrain to mediate behavior.

Main text

Vibration sensing is ubiquitous across the animal kingdom, enabling organisms to interpret their dynamic surroundings, communicate, detect threats, and navigate their environment[6-10]. In mammals, vibrations that propagate through the air in the form of sound waves are transduced into electrical signals in the cochlea and encoded by the auditory system[1]. Analogously, mechanical vibrations acting on the body are detected by fast conducting low-threshold mechanoreceptors (Aβ LTMRs) of the body and encoded by the somatosensory system[2,11]. In many animals, including humans, the vibration frequency detection range of the auditory system (20 Hz-20 kHz) partially overlaps with that of the somatosensory system (1-1000 Hz)[2,12-14]. In some smaller mammals, such as mice, the encoding of vibration frequency in the somatosensory system (1-1000 Hz) and auditory system (~1 khz-100 kHz) is largely non-overlapping[5,12,13,15-17].

Our understanding of LTMRs and their role in detecting mechanical vibrations mainly arises from experiments in which vibratory stimuli are presented directly to the body; however, the somatosensory system can also detect subtle, low amplitude environmental vibrations. In a noteworthy example, elephants can communicate over miles, sensing vocalizations and movements of conspecifics by detecting vibrations of the ground using mechanoreceptors in their feet and trunks[9,18]. Among the LTMRs, Pacinian corpuscle Aβ rapidly adapting type-2 (RA2)-LTMRs are unique in their capacity to encode high frequency vibrations[2]. In humans and primates, Pacinians are located in the deep dermis of glabrous skin and also associated with joints and connective tissue[19,20]. In rodents, Pacinians are enriched in the periosteum of wrist and ankle bones, rendering them responsive to vibrations transmitted through the skeleton[2,5,16,21,22]. Recordings from Pacinian afferents in awake, behaving mice have recently revealed their exquisite sensitivity to self-movements and substrate vibrations under naturalistic conditions[5]. Indeed, Pacinian corpuscle Aβ RA2-LTMRs can fire robustly in response to low-amplitude environmental vibrations, including those initiated by gentle movements on surfaces located meters from where an animal stands[5,23-26].

The canonical view is that the neural impulses encoding mechanical vibrations of the body are conveyed via the brainstem dorsal column nuclei (DCN) to the ventroposterolateral thalamus (VPL) *en route* to primary somatosensory cortex (S1)[11,27]. However, recent work suggests that

high frequency vibration sensitive DCN neurons project to the inferior colliculus, a midbrain

region involved in auditory information processing[1,28]. The inferior colliculus (IC) is comprised

of multiple subregions: the central nucleus of the inferior colliculus (CNIC), which solely

receives auditory input in a tonotopic manner, and the lateral cortex of the inferior colliculus

(LCIC), which receives inputs from multiple sensory modalities, including somatosensory inputs

from the DCN (Fig. 1A, Extended Data Fig. 1)[1,28-39]. Here, we use electrophysiological

recordings, genetic manipulations, and behavioral analyses to show that most LCIC neurons

encode both high frequency mechanical vibrations captured by Pacinian corpuscle LTMRs and

auditory vibrations captured by the cochlea to mediate behavioral responses to environmental

vibrations.

LCIC and VPL neurons are differentially tuned to mechanical vibrations

Distinct populations of DCN neurons send axonal projections to either the VPL or the LCIC

(Fig. 1A, Extended Data Fig. 1)[38]. To compare how mechanical vibrations are represented in the

LCIC and VPL, we performed *in vivo* multi-electrode array (MEA) recordings in these brain

regions in urethane anesthetized animals (Extended Data Fig. 2). We first brushed the animal to

identify neurons containing limb receptive fields. Sinusoidal mechanical vibrations of different

frequencies and amplitudes were then delivered to the limbs using a mechanical stimulator.

While the LCIC and VPL both contain forelimb- and hindlimb-responsive neurons (Extended

Data Fig. 2, Extended Data Fig. 3), we focused primarily on hindlimb responsive neurons.

Vibrations were delivered to the most responsive region, or "hotspot," of a neuron's receptive

field. In the VPL, vibratory receptive fields tended to be focal areas such as an individual pad or

digit (Fig 1C). In contrast, receptive fields of LCIC neurons lacked a focal hotspot and were

spatially extensive (Fig. 1E). In these cases, the mechanical stimulator was positioned on the heel

of the hindlimb. Vibratory stimuli were then delivered at frequencies between 10-900 Hz and

forces between 2-50 mN. The majority of VPL neurons responded robustly to low frequency

vibrations (<200 Hz), but only transiently to high frequency vibrations (>200 Hz; Fig. 1B,

Extended Data Fig. 2C-D, Extended Data Fig. 5A,E). While their pre-synaptic DCN inputs were

able to entrain (phase-lock firing to each cycle of the stimulus) to vibratory frequencies up to 100

Hz[18], no VPL neurons could entrain up to 100 Hz, although some neurons could entrain to

vibratory frequencies up to 50 Hz. Strikingly, LCIC neurons showed an inverse response pattern

compared to VPL neurons, with most neurons responding transiently to low frequency vibrations

(<200 Hz) but firing robustly to high frequency stimuli (200-900 Hz; Fig. 1D, Extended Data

Fig. 2G-H, Extended Data Fig. 5I,M). While many DCN projection neurons targeting the LCIC

entrained to vibratory frequencies from 10-500 Hz[18], LCIC neurons only transiently responded to

low frequency stimuli (Fig. 1D, S2G-H, Extended Data Fig. 5I), suggesting a synaptic filtering

mechanism between the DCN and LCIC that preferentially favors high frequency vibratory

stimuli. Furthermore, although LCIC neurons responded robustly to high frequency vibrations,

they exhibited little or no entrainment (Fig. 1D, Extended Data Fig. 2G-H, Extended Data Fig.

5M).

While high frequency vibratory responses were abundant throughout the LCIC, the sparsity of

high frequency responses detected in the VPL was surprising and could be attributed to missing a

potential "high frequency hotspot" region in the VPL in our recordings. Therefore, considering

that the VPL is an elongated structure extending along the anterior-posterior axis of the brain, we

next used a four shanked MEA to perform a detailed mapping of somatotopy across the VPL to

search for high frequency vibratory responses. We recorded from a minimum of 9 penetrations

per animal using the four-shanked probe, enabling sampling across at least 36 regions of the

VPL, distributed across the medio-lateral and anterio-posterior axes. The resulting somatotopic

map of the VPL aligned well with those previously reported, with receptive fields moving from

forelimb to hindlimb to trunk, when moving from medial to lateral within the VPL (Extended

Data Fig. 3A, C-D)[40-43]. For VPL neurons with brush responses to hindbody regions, high

frequency vibration of the hindlimb activated ~9 percent of neurons, and these high frequency

responsive neurons were sparsely distributed throughout the VPL, not concentrated in a

particular subregion (Fig. 1H, Extended Data Fig. 3E). In contrast, >75 percent of brush

responsive LCIC neurons responded robustly to high frequency vibrations, indicating that high

frequency vibratory information is particularly salient in the LCIC (Fig. 1H).

To assess the relationship between mechanical force thresholds and vibration frequency in the

VPL and LCIC, we next obtained force-frequency tuning curves for vibration sensitive VPL and

LCIC neurons by determining force thresholds for individual neurons across a range of vibratory

frequencies. The force threshold for a given frequency was defined as the minimum force

required to elicit 20 percent of a neuron's maximum firing rate. The tuning curves for VPL and

LCIC neurons were approximately U-shaped, although the most sensitive regions of their

respective tuning curves were non-overlapping: VPL neurons had the lowest force thresholds to

vibrations between 50-200 Hz and LCIC neurons had the lowest force thresholds to vibrations

between 300-500 Hz (Fig. 1F-G, I-K). Thus, most LCIC neurons are activated by high frequency

vibratory stimuli and exhibit expansive receptive fields, while most VPL neurons are

preferentially activated by low frequency stimuli in a spatially restricted manner.

LCIC vibration responses are mediated by Pacinian corpuscle Aβ RA2-LTMRs

The distinct frequency tuning curves and receptive field properties of VPL and LCIC neurons

prompted us to ask whether these brain regions receive inputs from different primary

mechanosensory neuron types. Rapidly adapting Aβ LTMRs exhibit distinct force-frequency

tuning relationships and receptive fields. Most rapidly adapting LTMRs are sensitive to low

frequency vibratory stimuli (40-150 Hz) and exhibit relatively small receptive fields[2,11,15,44-46].

This includes Meissner corpuscle Aβ RA1-LTMRs, enriched in glabrous skin of pedal pads and

digits, hair follicle Aβ RA-LTMRs associated with body hairs, and Krause corpuscle Aβ RA-

LTMRs of the genitalia[2,15,45,47]. In contrast, Pacinian corpuscle Aβ RA2-LTMRs, which in mice

are enriched in the periosteum of ankle and wrist bones, are most sensitive to high frequency

vibratory stimuli (50-1000 Hz) and have extremely large receptive fields[4,5,21,48-52]. To directly

compare the tuning of Meissner corpuscle Aβ RA1-LTMRs and Pacinian corpuscle Aβ RA2-

LTMRs to VPL and LCIC neurons, we performed *in vivo* extracellular recordings of DRG

neurons and measured force-frequency tuning curves of Meissner and Pacinian afferents (Fig.

1L-N). As predicted, the force-frequency tuning curves of Pacinian corpuscle Aβ RA2-LTMRs

resembled those of LCIC neurons, while the tuning curves of Meissner corpuscle Aβ RA1-

LTMRs more closely resembled VPL neurons (Fig. 1L-N). These findings led us to hypothesize

that Pacinian afferents have an outsized role in shaping the response properties of most LCIC

neurons, while Meissner corpuscle afferents and other Aβ RA-LTMRs of the body preferentially

contribute to the response properties of most VPL neurons.

To directly test the relationship between primary sensory neuron activity and vibration tuning in

the LCIC and VPL, we utilized mouse genetic tools to assess the contributions of Pacinian

corpuscles and Meissner corpuscles to LCIC and VPL responses. Ret signaling in somatosensory

neurons is necessary for formation of Pacinian corpuscles, and thus *Avil^Cre; Ret^floxGFP/floxGFP* (Ret

cKO) mice, which lack *Ret* in DRG neurons, lack Pacinian corpuscles (Fig. 2B, Extended Data

Fig. 4F)[53,54]. Conversely, TrkB signaling is necessary for formation of Meissner corpuscles, and

thus *Avil^Cre; TrkB^flox/flox* (TrkB cKO) mice lack Meissner corpuscles (Fig. 2A)[15,45,55]. Importantly,

Ret cKO mice have a normal density of Meissner corpuscles and TrkB cKO mice have a normal

density of Pacinian corpuscles (Extended Data Fig. 4).

Surprisingly, VPL neurons recorded in TrkB cKO and Ret cKO mice showed no notable shifts in

their vibration tuning curves (Fig. 2C, F). Meissner corpuscle and Pacinian corpuscle afferents

are not the only primary sensory neurons responsive to vibrations, and indeed Aβ RA-LTMRs

associated with hair follicles and other end organs and Aβ SA1-LTMRs found in both hairy and

glabrous skin also entrain to vibratory stimuli[2,52,56,57]. Therefore, it is likely that VPL neurons in

TrkB cKO and Ret cKO mice are sensitive to mechanical vibrations via inputs from primary

sensory neuron subtypes other than just Meissner and Pacinian corpuscle LTMRs[55,58]. In stark

contrast, vibration responsiveness of LCIC neurons was nearly absent in animals lacking

Pacinian corpuscles, although LCIC responses were normal in mice lacking Meissner corpuscles

(Fig. 2D, E, G). Taken together, these findings indicate that high frequency mechanical

vibrations acting on the body and detected by Pacinian corpuscle Aβ RA2-LTMRs are

prominently encoded in the LCIC.

LCIC neurons integrate multimodal tactile-auditory stimuli in awake mice

Recent recordings from Pacinian afferents under awake conditions revealed that Pacinian Aβ

RA2-LTMRs are exquisitely sensitive to environmental vibrations encountered during natural

behaviors[5]. Therefore, we next asked whether LCIC neurons are also sensitive to low amplitude

substrate vibrations in awake animals. We performed MEA recordings in awake head-fixed mice,

but instead of applying vibration directly to the paw, we vibrated the acrylic platform the animal

stood on (Fig. 3A). Remarkably, in awake mice, 100 percent of LCIC neurons with hindlimb

receptive fields responded robustly to 500 Hz stimulation of the surface platform, but few

responded to a 50 Hz stimulus, similar to results from direct stimulation of the limb in recordings

from anesthetized mice (Fig. 1D, H, Extended Data Fig. 2G-H, Fig. 3B-C). Thus, as with

Pacinian corpuscle Aβ RA2-LTMRs, LCIC neurons encode low-amplitude, high frequency

environmental vibrations.

The inferior colliculus is primarily an auditory nucleus, and the LCIC receives extensive auditory

input from brainstem auditory nuclei, prompting us to ask whether LCIC neurons responsive to

high frequency tactile vibrations also respond to sound[1]. Prior findings indicate that many LCIC

neurons are broadly tuned to auditory stimuli, and most auditory-sensitive LCIC neurons respond

to short pulses of broadband noise[31,36,59,60]. Therefore, we exposed animals to 100 ms pulses of a

broadband white noise stimulus (0-50 kHz) at a 65 dB sound pressure level (SPL) with the goal

of activating LCIC neurons that receive auditory inputs. We delivered 500 Hz mechanical

vibrations to the platform surface and white noise pulses in isolation, as well as coincident

tactile-auditory stimulation (Fig. 3D) to awake mice while recording in the LCIC. Remarkably,

~75% of LCIC neurons were activated by both high frequency mechanical vibration of the

platform and auditory white noise, with a minority of neurons responding to only platform

vibration or sound (Fig. 3E, H). These findings reveal a high degree of convergence of auditory

and mechanical vibration signals in the LCIC. Interestingly, of the LCIC neurons that received

convergent tactile-auditory input, the majority had a larger response to coincident presentation of

mechanical vibration and sound than to either vibration or sound alone (Fig. 3J, Extended Data

Fig. 6D-I).

We further assessed the multimodal nature of LCIC responses to environmental vibrations by

measuring responses of LCIC neurons in mice lacking either Pacinian corpuscles or cochlear

function. As predicted from the findings in anesthetized mice (Fig. 2D-G), LCIC neurons in

awake Ret cKO mice, which lack Pacinian corpuscles, were unresponsive to high frequency

surface vibrations (Fig. 3F, I-J). Despite their insensitivity to vibration, LCIC neurons in Ret

cKO mice were responsive to brush, suggesting that the LCIC receives inputs from other classes

of cutaneous LTMRs (Extended Data Fig. 7D). It is noteworthy that deletion of Ret does not

impact the auditory system, as the Auditory Brainstem Response (ABR) test in Ret cKO mice

indicated similar auditory thresholds to littermate controls (Extended Data Fig. 7A-C).

Furthermore, LCIC neurons of Ret cKO mice were comparably sensitive to auditory white noise

stimuli compared to controls, indicating that developmental loss of Pacinian corpuscles does not

alter LCIC sensitivity to broadband auditory stimuli (Fig. 3F, I-J). In complementary

experiments, we tested whether disrupting auditory system function alters mechanical vibration

responsiveness in the LCIC. For this, we utilized $Sox10^{Cre}$; $Gjb2^{flox/flox}$ mice (Gjb2 cKO), which

lack connexin 26 expression in epithelial cells of the cochlea[61,62]. Gjb2 cKO mice lack cochlear

hair cell function but have an intact vestibular system, such that this congenital deafness model

exhibits extensive hearing loss but retains normal balance and motor function[61-65]. ABR testing in

Gjb2 cKO mice confirmed their inability to hear, as none of the Gjb2 cKO mice tested had a

discernable brainstem response at the highest sound level tested (Extended Data Fig. 8A-C).

Recordings in anesthetized Gjb2 cKO mice revealed that in the absence of normal hearing, LCIC

neurons exhibit robust responses to mechanical vibration stimulation (Extended Data Fig. 8F-G).

Force-frequency tuning curves in Gjb2 cKO showed no difference from controls (Fig S8F-G).

Furthermore, recordings in awake Gjb2 cKO mice demonstrated that LCIC neurons responded to

substrate vibrations although they showed no responsivity to white noise auditory stimuli (Fig.

3G, I-J). Thus, genetically disrupting either somatosensory/Pacinian corpuscle or auditory

system function does not grossly alter LCIC neuron responses to the other sensory modality.

These findings indicate that most LCIC neurons receive both auditory input emanating from the

cochlea and somatosensory input emanating from the body's Pacinian corpuscles. Therefore, the

LCIC is a site of auditory-mechanical vibration multisensory integration and enhancement (Fig.

3E-J, Extended Data Fig. 6).

The LCIC mediates behavioral avoidance to tactile vibration

The LCIC has been shown to mediate defensive behaviors in response to auditory cues[66]. To

begin investigating whether the LCIC can drive behavioral responses to high frequency

mechanical vibrations, we developed a behavioral preference assay that relies on an animal's

ability to detect and respond to high frequency mechanical vibrations of the environment.

Animals were placed in a two-chamber behavioral setup where the floor of one chamber vibrates

at 500 Hz, a stimulus that robustly activates Pacinian corpuscle afferents and the LCIC of awake

animals (Fig. 3B, E., Fig. 4A, Fig. S9A-D)[5]. Wildtype animals had a strong and reliable

preference for the non-vibrating chamber, spending ~80% of their time there when the vibration

stimulus was on (Fig. 4B-C, Extended Data Fig. 9E-H). Since the production of the 500 Hz

vibration generated a low amplitude 500 Hz auditory tone, we performed a sound-only control.

We decoupled the vibration motor from the chamber floor, such that a comparable sound was

produced with no surface vibration. In the sound-only paradigm, animals showed no aversion

toward the stimulus side of the chamber, demonstrating that sound alone does not drive a

preference (Fig. 4D). To test whether the behavior is dependent on Pacinian corpuscles, the

behavioral preference assay was done using Ret cKO mice, which lack Pacinian corpuscles. Ret

cKO mice demonstrated no preference for the non-vibrating side of the chamber over the

vibrating side, indicating that avoidance of the 500 Hz vibration stimulus is dependent on

Pacinian corpuscles (Fig. 4E-F, Extended Data Fig. 9I-J).

The vibration place preference assay was next used to determine whether avoidance of 500 Hz

mechanical vibration is dependent on the LCIC. Targeted, bilateral injections of a fluorescently-

tagged GABA-A receptor agonist, muscimol, were used to silence neural activity in the inferior

colliculus (IC) (Fig. 4G, Extended Data Fig. 10C). Post-hoc histology confirmed that muscimol

was reliably restricted to the IC and did not extend into neighboring brain regions, including the

superior colliculus, periaqueductal gray, or cerebellum (Fig. 4G, Extended Data Fig. 10A-B)[66-68].

IC-silenced mice displayed normal locomotor activity, traveling equal distances during the assay

as controls (Fig. 4H). Strikingly, IC-silenced mice did not exhibit avoidance of the vibratory

stimulus compared to saline injected controls, revealing that the IC is required for the behavioral

response to high frequency mechanical vibration (Fig. 4I-K, Extended Data Fig. 10D). To

determine whether the IC is required for general behavioral avoidance, or whether the effect of

IC silencing is specific to vibratory stimuli, we also tested whether IC silencing alters avoidance

of other somatosensory stimuli. We first performed a texture preference assay in saline and

muscimol treated mice. In a two-chamber setup, one side of the chamber was covered in a

smooth construction paper, while the other side was covered with a coarse grit sandpaper, which

is aversive (Fig. 4L, Extended Data Fig. 10F-G). Mice injected with either saline or muscimol in

the IC showed strong preference for the smooth side of the chamber, revealing that the IC is not

necessary for aversive texture avoidance (Fig. 4M). A cold temperature avoidance test was also

performed. For this, we used a two-chamber cold avoidance assay in which mice were given a

choice between one chamber, which was held at room temperature (30 ºC), and a second

chamber, which was held at 18 °C (Fig. 4N, Extended Data Fig. 10E). Similar to the texture

preference assay, both control and IC-silenced animals displayed robust avoidance of the cold

chamber (Fig. 4O). These findings indicate that the IC mediates behavioral response to high

frequency mechanical vibrations but is not required for avoidance of an aversive texture or

temperature.

Discussion

The majority of research into the neurobiological basis of discriminative touch of the body has

focused on the ascending pathway through the thalamus (VPL) to somatosensory cortex (S1),

while somatosensory representations in other brain areas have remained largely unexplored.

Here, we report that neurons in the LCIC and VPL exhibit highly distinct vibrotactile tuning

properties that may underlie unique functions of divergent ascending somatosensory system

streams. Most VPL neurons had restrictive receptive fields and were tuned to low frequency

vibrations. These properties are well suited for tactile acuity and discrimination—functions

hypothesized to be mediated by the VPL to S1 pathway[69-72]. Only a small fraction of VPL

neurons (~9%) encoded high frequency vibration (Fig. 1H). In contrast, most LCIC neurons

exhibited expansive receptive fields and were tuned to high frequency vibrations (>75%)

conveyed by Pacinian corpuscle Aβ RA2-LTMRs (Fig. 1H). The salience of low amplitude

environmental vibrations encoded in the LCIC is highlighted by our finding that IC silencing

virtually eliminated animals' avoidance of a 500 Hz vibratory stimulus in a place preference

paradigm (Fig. 4I-J, Extended Data Fig. 10D). It is possible that the divergent ascending streams

through the LCIC and VPL underlie distinct responses to high frequency vibrations in different

contexts. For example, the sparser representation of high frequency vibration in the VPL to S1

stream may underlie operant tasks that use vibratory cues[16,71-73]. Future work will discern the

relative contributions of distinct ascending streams encoding environmental vibrations in

different contexts.

Strikingly, most mechanical vibration sensitive LCIC neurons are also tuned to auditory stimuli.

Moreover, the majority of LCIC neurons that received convergent mechanical vibration and

auditory inputs exhibited multisensory enhancement, responding more strongly to coincident

tactile-auditory stimulation than to either modality alone (Fig. 3H-J. Extended Data Fig. 6E-F).

Thus, the LCIC encodes environmental vibrations, whether they are mechanical vibrations

propagated through the ground, detected by Pacinian corpuscles, and conveyed via the ascending

somatosensory pathway or sound waves traveling through air, captured by the cochlea, and

conveyed via the ascending auditory pathway (Extended Data Fig. 10H). We suspect that the

LCIC enables enhanced detection or reaction to environmental cues through auditory and

vibrotactile coincidence detection. A myriad of natural phenomena generates coincident sounds

and substrate vibrations, including wind, rivers, seismic events or earth movements, objects

moving across or falling to the ground, vocalizations or rumblings of conspecifics, and the

motions of prey or predators. Multisensory audio-vibrotactile integration in the LCIC may

increase saliency of these ethologically relevant environmental cues and may be particularly

instrumental under conditions in which cues are weak and difficult to discern. We propose that,

through their convergent inputs to the LCIC, the somatosensory and auditory systems collaborate

to detect, localize, and respond to environmental vibrations.

  Figures and Legends

Figure 1: LCIC and VPL neurons are differentially tuned to mechanical vibratory stimuli.

A) The brainstem dorsal column nuclei (DCN) receive inputs from dorsal root ganglia (DRG)

primary sensory neurons and spinal cord projection neurons. Non-overlapping subsets of DCN

projection neurons target the ventroposterolateral thalamus (VPL; blue) and the external nucleus of the inferior colliculus (LCIC; red).

B and D) Representative VPL (B) and LCIC (D) neuron spike raster and histogram in response to 50 Hz and 500 Hz vibratory stimuli delivered at a 10 mN force to the hindlimb. Top: spike raster; bottom: average histogram in 10 ms bins.

C and E) Example VPL (C) and LCIC (E) neuron receptive field organization for 50 Hz and 500 Hz stimulation to multiple regions across the hindlimb, delivered at 10 mN and 20 mN intensities. Stimulus locations are superimposed on images of the hindpaw.

F and G) Representative VPL (F) and LCIC (G) neuron responses to vibrational frequencies delivered at different forces to the hindlimb. Heatmap intensity corresponds to the neuron's average change in firing to the sustained portion of vibratory stimuli (200 ms window beginning 50 ms after stimulus onset).

H) Percentage of neurons (mean ± SD) with hindbody brush receptive fields (tested by brushing the hindpaw, thigh, and trunk) that respond to 500 Hz stimulation of the limb at a 10 mN force in the VPL (n= 182 neurons from 4 animals) and LCIC (n= 76 neurons from 4 animals). Each data point corresponds to measurements in one animal. See also Extended Data Fig. 3.

I and J) Force-frequency threshold curves obtained from heatmaps in F and G, where the force threshold for a vibrational frequency is defined as the minimum force required to elicit 20% of the neuron's maximum firing during the sustained portion of the stimulus. Frequencies in which the highest force tested (50 mN) failed to reach threshold are designated NR (non-responsive).

K) Average threshold curves (mean ± 95% confidence interval) for neurons responsive to hindlimb vibration recorded in the VPL (n=15 neurons from 6 animals) and LCIC (n=22 neurons from 6 animals).

L and M) Force-frequency threshold curves for representative Meissner corpuscle innervating Aβ RA1-LTMR afferent (L) and Pacinian corpuscle innervating Aβ RA2-LTMR afferent (M). Threshold is defined as the minimum force necessary for the neuron to reach 50% vibratory entrainment. Frequencies in which the highest force tested (50 mN) failed to reach threshold are designated NR (non-responsive).

N) Average threshold curves (mean ± 95% confidence interval) for hindlimb Meissner Aβ RA1-LTMRs (n=4 neurons from 4 animals) and Pacinian Aβ RA2-LMTRs (n=6 neurons from 5 animals).

Figure 2: LCIC vibration responses are mediated by Pacinian corpuscle Aβ RA2-LTMRs.

A) Meissner corpuscles, located in the plantar pad and digit glabrous skin, are absent in *Avil^{Cre}; TrkB^{flox/flox}* (TrkB cKO) mice, which lack TrkB in DRG sensory neurons. Images of skin histology from a TrkB cKO and littermate control, stained for anti-NFH (labels myelinated axons, green) and anti-S100 (labels glial cells, magenta). Scale bar = 50 µm. See also Extended Data Fig. 4 for quantification.

B) Pacinian corpuscles, located in the periosteum of the fibula, are absent in *Avil^{Cre}; Ret^{flox/flox}* (Ret cKO) mice, which lack Ret in DRG sensory neurons. Images of whole mount histology of the limb of a Ret cKO animal and littermate control, stained for anti-NFH and anti-S100 to visualize Pacinian corpuscles. White dashed line indicates position of the fibula. Scale bar = 100 µm. See Extended Data Fig. 4 for quantification.

C and D) Force-frequency heat maps for representative VPL (C) and LCIC (D) neurons in control, TrkB cKO and Ret cKO animals. Heatmap intensity corresponds to the neuron's average change in firing to the sustained portion of vibratory stimuli.

E) Maximum firing rate for VPL and LCIC (mean ± 95% confidence interval) neurons to vibratory stimuli in control, TrkB cKO and Ret cKO mice. Firing rates of LCIC neurons in Ret cKO and control mice were significantly different (****$p<0.0001$ Mann-Whitney U test). Firing rates of VPL neurons in TrkB cKO, Ret cKO, and controls were not significantly different. Firing rates of LCIC neurons in TrkB cKO and controls were not significantly different (Mann-Whitney U test).

F) Average threshold curves of VPL neurons (mean ± 95% confidence interval) recorded in control (n=15 neurons from 6 animals), TrkB cKO (n=10 neurons from 3 animals), and Ret cKO (n=14 neurons from 4 animals) mice. Thresholds between Ret cKO, TrkB cKO and controls were not significantly different (Mann-Whitney U Test with Bonferroni correction for multiple comparisons).

G) Average threshold curves of LCIC neurons (mean ± 95% confidence interval) recorded in control (n=22 neurons from 6 animals), TrkB cKO (n=9 neurons from 4 animals), and Ret cKO (n=18 neurons from 4 animals) mice. (*$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, Mann-Whitney U test with Bonferroni correction for multiple comparisons). While thresholds between Ret cKO and controls were significantly different, thresholds between TrkB cKO and controls were not (Mann-Whitney U Test with Bonferroni correction for multiple comparisons).

Figure 3: Audio-tactile multimodality of LCIC neurons in awake mice.

A) Recordings of LCIC neurons were conducted in awake head-fixed mice as they stood on an

acrylic platform, which could be stimulated at different vibratory frequencies. An accelerometer

was attached underneath the acrylic platform to measure vibration frequency and amplitude. An

example motor command and accelerometer signal are shown for 500 Hz vibration of the platform. A speaker is positioned on the contralateral side of the head from the recording site.

B) Peristimulus time histogram (PSTH) for a representative LCIC neuron, aligned to 50 Hz (left) and 500 Hz (right) vibration stimulation of the acrylic platform at a vibration amplitude of ~0.2 g (gravitational acceleration). PSTHs are baseline subtracted and reported in 10 ms bins.

C) Percentage of hindlimb sensitive LCIC neurons that responded to 50 Hz and 500 Hz vibratory stimuli (n=24 neurons from 5 animals).

D) 500 Hz substrate vibration (amplitude 0.2 g) and auditory white noise (frequency 0-50 kHz, 100 ms pulses, 65 dB SPL) were delivered in isolation and coincidentally to awake head-fixed mice.

E) Example PSTHs from an LCIC neuron of a wildtype mouse aligned to vibration (v), sound (s) and convergent (vs) stimuli. PSTHs are baseline subtracted and in 10 ms bins. Right: The example neuron's average firing (during 0-0.1 s and 0.5-0.6 s windows) for each stimulus type.

F and G) As in (E), example LCIC neurons in a Ret cKO animal, which lacks Pacinian corpuscles (F), and a Gjb2 cKO animal, which is unable to hear (G).

H) Percentage of neurons in control animals that respond to 500 Hz vibration only, sound only, or convergent stimuli (n=29 neurons from 5 animals).

I) Scatter plot of each neuron's average change in firing rate to vibration and sound stimuli in control (n=29 neurons from 5 animals), Ret cKO (n=14 neurons from 3 animals) and Gjb2 cKO (n=15 neurons from 3 mice) animals.

J) Average change in firing rate to vibration, sound, or combined stimuli (mean ± SD) in control (n=29 neurons from 5 animals), Ret cKO (n=14 neurons from 3 animals) and Gjb2 cKO (n=15 neurons from 3 mice) animals (**$p<.01$, ***$p<0.001$, ****$p<0.0001$, Mann-Whitney U test).

Figure 4: The LCIC mediates behavioral avoidance to platform vibration.

A) Animals were placed in a 2-chamber behavioral setup. The floors of each chamber are

disconnected to prevent propagation of vibration between them, and an exciter speaker is

attached to the floor of one chamber to generate mechanical vibrations. The chamber walls are

partially open to prevent sound resonance from the exciter.

B and C) The percentage of time wildtype CD-1 animals spend on each side of the chamber (mean ± SD) during the 60 s baseline explore period (B) and the vibration periods (C) of the trial. Each dot represents one animal (n=20 animals, ****p<0.0001 paired t-test).

D) The speaker was detached from the floor of the chamber to test whether the sound of the exciter speaker alone can drive a behavioral preference. Time spent on each side of the chamber during the sound on periods was not statistically different (n=10 animals, paired t-test).

E and F) The percentage of time Ret cKO mice (n=14 animals) and littermate controls (n=24 animals) spent on each side of the chamber during the 60 s baseline explore period (E) and the vibration periods (F) of the behavioral trial. (****p<0.0001, paired t-test within group, unpaired t-test between groups).

G) Targeted injections of TMR-X conjugated to the GABA-A receptor agonist, muscimol, were used to silence neural activity in the inferior colliculus. Top-down view of a schematic (left) and representative muscimol injected brain. Dotted lines depict borders between the inferior colliculus and indicated neighboring brain regions.

H) Total distance traveled during the behavioral assay in saline injected controls (n=8) and muscimol injected animals (n=9 animals, not significant, unpaired t-test).

I and J) The proportion of time muscimol injected mice (n=9 animals) and saline injected controls (n=8 animals) spent on each side of the chamber during the 60 s baseline explore period (I) and the vibration periods (J) of the trial. (**p<0.01, ***p<0.001, paired t-test within group, unpaired t-test between groups)

K) Representative activity traces for saline injected (control) and muscimol injected animals during the baseline and vibration periods of the behavioral trial. See also Extended Data Fig. 9.

L) Schematic of texture preference assay.

M) Percentage of time spent on the smooth and rough sides of the chamber during the 10 min assay for saline controls (n=9 animals) and muscimol silenced animals (n=9 animals, ***p<0.001, paired t-test within group, unpaired t-test between groups).

N) Schematic of temperature preference assay.

O) Percentage of time spent on the control (30 °C) and cold (1 8°C) sides of the chamber during the 5 min assay for saline controls (n=8) and muscimol silenced animals (n=8, ***p<0.001, paired t-test within group, unpaired t-test between groups).

none
Extended Data Figures and Legends

Extended Data Figure 1: Anterograde tracing from the dorsal column nuclei.

A) To identify downstream projections from the dorsal column nuclei (DCN), an AAV encoding

Cre-dependent synaptophysin-tdT was injected into the gracile nucleus and an AAV encoding

Cre-dependent synaptophysin-GFP was injected into the cuneate nucleus of Vglut2-Cre mice.

B-C) TdT and GFP positive terminals were observed throughout the ventroposterolateral

thalamus (VPL) and lateral cortex of the inferior colliculus (LCIC). Scale bar = 150μm.

Extended Data Figure 2: MEA recordings in the VPL and LCIC to characterize vibratory responses.

A) Representative probe tract from MEA recording in the VPL. Scale bar = 1000μm.

B) Average waveforms and corresponding autocorrelogram for three example VPL neurons.

C) Example VPL neuron raster and PSTHs in response to vibration of different frequencies

delivered at a 10 mN force to the forelimb. Top: spike raster; bottom: PSTHs, which are baseline

subtracted and in 10ms bins.

D) Same as C for a hindlimb responsive neuron.

E) Representative probe tract from MEA recording in the LCIC. Scale bar = 500 μm.

F) Average waveforms and corresponding autocorrelogram for three example LCIC neurons.

G) Same as C for an example LCIC neuron.

H) Same as D for an example LCIC neuron.

Extended Data Figure 3: Somatotopic and vibratory mapping in the VPL and LCIC.

A) Representative VPL neurons responding to brush of different regions of the body. Each row

corresponds to one neuron. Each body region is stroked four times per trial. Right: Example

raster plot and PSTHs aligned to hindlimb vibration at 50 Hz and 500 Hz at 10 mN,

corresponding to adjacent brush responsive neurons. PSTHs are baseline subtracted and in 10ms

bins.

B) As in A, representative LCIC neurons. Note that, unlike VPL neurons, most LCIC neurons

respond to brush across multiple body locations.

C-E) Example mapping of the VPL in one animal, shown at different coronal planes on the

anterior-posterior axis. The approximate atlas location (C) for recorded neurons is shown for the

corresponding somatotopic map of the VPL (D), where each neuron is colored for the body

region most responsive to brush. The same neurons shown in D are colored by their

responsiveness to 500 Hz mechanical vibration, which is normalized to the neuron's response to

brush (E). Note that most brush responsive VPL neurons do not respond to the 500 Hz vibration

stimulus during the sustained period of the stimulation.

F and G) Percentage of neurons (mean ± SD) with hindbody receptive fields that respond to

brush of the hindpaw, thigh, and trunk in the VPL (F) (n= 182 neurons from 4 animals) and LCIC

(G) (n= 76 neurons from 4 animals). Each data point corresponds to a single animal.

Pacinian corpuscles

A Control — S100 NFH

B TrkB cKO

Meissner corpuscles

C Control — DAPI S100 NFH

D Ret cKO

E Meissner Density

ns

Density (mm²)

Control   Ret cKO
Genotype

F Pacinian Counts

Control / TrkB cKO

ns   ns

Pacinians

Hindlimb   Forelimb

Control / Ret cKO

**   **

Pacinians

Hindlimb   Forelimb

Extended Data Figure 4: Anatomical characterization of peripheral endings in Ret cKO

and TrkB cKO mice.

A-B) Images of Pacinian corpuscles from the limbs of TrkB cKO animals and littermate controls.

Limbs were whole-mount stained with antibodies against NFH (green), which labels large caliber

axons, and S100 (magenta), which labels lamellar cells of the corpuscle and myelinating

Schwann cells. Left scale bar = 250μm. Right scale bar = 50 μm.

C-D) Images of Meissner corpuscles (indicated by white arrows) from Ret cKO animals and

littermate controls. Skin tissue was stained using antibodies against NFH (green) and S100

(magenta). DAPI staining is shown in gray. Left scale bar = 50 μm. Right scale bar =20 μm.

E) Quantification of Meissner density in Ret cKO (n=3 animals) and littermate controls (n=3

animals). Each data point is the Meissner density for a single skin section, and data points of the

same color came from the same animal (no significant difference using unpaired t-test).

F) Quantification of Pacinian corpuscles in TrkB cKO (n=3 animals) and littermate controls (n=3

animals), and Ret cKO (n=3 animals) and littermate controls (n=3 animals). The number of

Pacinians was counted from wholemount staining of hindlimbs and forelimbs (mean ± SD,

**p<0.001, unpaired t-test).

Extended Data Figure 5: Onset and offset responses to vibratory stimuli in Ret cKO and TrkB cKO mice.

A) Top: Heatmap of neurons in the VPL responding to 100 Hz vibration at a 15 mN force (n=14 neurons from 6 animals). Each row is one neuron. Bottom: PSTHs of each neuron in above heatmap (gray), and population average PSTH (red). PSTHs are in 20ms bins.

B-C) Same as A in TrkB cKO (n=10 neurons from 3 animals) and Ret cKO mice (n=14 neurons

from 4 animals).

D) Average firing rate of VPL neurons to the onset (top) and offset (bottom) of the 100 Hz

vibration stimulus in control, TrkB cKO, and Ret cKO mice (mean ± SEM, not significantly

different, Mann-Whitney U test with Bonferroni correction for multiple comparisons). On

period is 20 ms after the stimulus onset, and off period is 20 ms after the end of the stimulus.

E-G) Same as A-C to 500 Hz vibration at 15 mN.

H) Same as D for 500 Hz stimulus (*p<0.05, **p<0.01 , ***p<0.001, ****p<0.0001, Mann-

Whitney U test with Bonferroni correction for multiple comparisons).

I-K) Same as A-C for neurons in the LCIC in control (n=18 neurons from 5 animals), TrkB cKO

(n=8 neurons from 4 animals), and Ret cKO (n=16 neurons from 5 animals) animals.

L) Average firing rate of LCIC neurons to the onset (top) and offset (bottom) of the 100 Hz

vibration stimulus in control, TrkB cKO, and Ret cKO mice (mean ± SEM, not significantly

different, Mann-Whitney U test with Bonferroni correction for multiple comparisons).

M-O) Same as I-K to 500 Hz vibration at 15 mN.

P) Same as L for a 500 Hz stimulus (*p<0.05, **p<0.01, Mann-Whitney U test with Bonferroni

correction for multiple comparisons).

Extended Data Figure 6: Audio-tactile multimodality of LCIC neurons in awake and anesthetized animals.

A) 500 Hz substrate vibration and auditory white noise (frequency 0-50 kHz, 100 ms pulses, 65 dB SPL) were delivered in isolation and coincidentally, as in Fig. 3.

B) Example PSTHs from two wildtype neurons aligned to vibration (v), sound (s) and convergent (vs) stimuli in anesthetized animals. PSTHs are baseline subtracted and in 10ms bins. Right: The example neuron's average firing (during 0-0.1 s and 0.5-0.6 s windows) for each stimulus type.

C) Same as B, except recordings were done in awake animals.

D) Scatter plot of change in firing rate to vibration or sound in neurons from awake (n=29 neurons from 5 animals) and anesthetized mice (n=9 neurons from 3 animals).

E) Scatter plot of firing rates in response to combined vibration and sound (vs) plotted against best unimodal response (vibration only or sound only). Unitary line denotes equal firing between best unimodal stimulus and combined stimulus conditions.

F) Quantification of multisensory enhancement in awake and anesthetized conditions (corresponding to E), calculated using the following equation: $\frac{vs-(Max(v\ or\ s))}{Max(v\ or\ s)}*100$. Neurons above 100% display multisensory enhancement[74,75].

G) Average change in firing rate to vibration, sound, or combined stimuli (mean ± SD) in control (n=29 neurons from 5 animals), and anesthetized (n=9 neurons from 3 animals) conditions (not significant, Mann-Whitney U test).

H) Scatter plot of firing rates in response to combined vibration and sound (vs) plotted against hypothetical summed vibration and sound responses (v+s). Unitary line denotes equal firing to combined stimulation (vs) and the sum of firing to vibration and sound stimuli (v+s).

I) Quantification of multisensory integration in awake and anesthetized conditions (corresponding to H), calculated using the following equation: $\frac{vs-(v+s)}{v+s}*100$. Neurons above 100% display super-additivity and neurons below 100% display sub-additivity[75,76].

Extended Data Figure 7: Ret cKO mice have normal auditory thresholds and brush responses.

A) Auditory brainstem response (ABR) thresholds were measured in Ret cKO mice and littermate controls at 5 different auditory frequencies (mean ± SEM). Auditory thresholds between Ret cKO (n=6 animals) and controls (n=6 animals) were not significantly different (Mann-Whitney U Test with Bonferroni correction for multiple comparisons).

B) ABR thresholds to a broadband frequency "click" were measured in Ret cKO mice (n=6 animals) and littermate controls (n=6 animals). Thresholds were not significantly different (Mann-Whitney U Test).

C) Distortion product otoacoustic emissions (DPOAE) thresholds, which are sounds generated by mechanically active cells in the ear in response to sound stimuli, were measured in Ret cKO mice (n=6 animals) and littermate controls (n=6 animals) for 8 different frequency pairs (mean ± SEM). Thresholds were not significantly different (Mann-Whitney U Test with Bonferroni correction for multiple comparisons), suggesting normal function of the sensory cells responsible for fine-tuning cochlear gain.

574   **D)** Average change in firing to brush stimuli applied to the hindbody of LCIC neurons in awake

575   control (n=10 neurons) and Ret cKO (n=13 neurons) mice. Responses to brush were not

576   significantly different (Mann-Whitney U Test).

ABR and DPOAE thresholds in Gjb2 cKO mice

Auditory responses in LCIC of Gjb2 cKO mice

Vibration tuning in LCIC of Gjb2 cKO mice

Extended Data Figure 8: Characterization of auditory and tactile responses in anesthetized Gjb2 cKO mice.

A) Auditory brainstem response (ABR) thresholds were measured in Gjb2 cKO mice (n=4 animals) and littermate controls (n=6 animals) at 5 different auditory frequencies (mean ± SEM).

Unlike littermate controls, Gjb2 cKO animals had no response at any frequency, even at the

highest sound level tested (80 dB).

B) ABR thresholds to a broadband frequency "click" were measured in Gjb2 cKO mice (n=4

animals) and littermate controls (n=6 animals). Gjb2 cKO animals had no response, even at the

highest sound level tested (80 dB).

C) Distortion product otoacoustic emissions (DPOAE) thresholds, which are sounds generated

by mechanically active cells in the ear in response to sound stimuli, were measured in Gjb2 cKO

mice (n=4 animals) and littermate controls (n=6 animals) for 8 different frequency pairs (mean ±

SEM). Gjb2 cKO animals had no DPOAE at any frequency, even at the highest sound level

tested (75 dB).

D) Example neurons responding to white noise at varying sound levels (dB) in anesthetized

littermate control and Gjb2 cKO mice. Each row corresponds to an individual neuron.

E) Average change in firing rate of neurons (mean ± SE) in littermate control (n=13 neurons

from 2 mice) and Gjb2 cKO mice (n=9 neurons from 2 mice) to white noise at varying sound

intensities.

F) Force-frequency heat maps in response to vibration of differing frequencies and intensities in

two neurons from littermate controls and two neurons from Gjb2 cKO mice.

G) Average force-frequency threshold curves of LCIC neurons (mean ± 95% confidence

interval) recorded in control (n=9 neurons from 2 animals), and Gjb2 cKO mice (n=11 neurons

from 2 animals). Thresholds between Gjb2 cKO and controls were not significantly different

(Mann-Whitney U Test with Bonferroni correction for multiple comparisons).

Extended Data Figure 9: A place preference behavioral assay for measuring behavioral responses to 500 Hz vibration of a platform.

A) Timeline of behavioral trial.

B-D) Accelerometer traces from vibration side (blue) and control side (gray) of the behavioral chamber during the baseline period (B), vibration period (C), and post-vibration period (D) of the

behavioral trial. Amplitude of the platform vibration was ~ 0.3 g (gravitational force) when the

vibration stimulus was on.

E-G) Percent of time spent that mice spent on the control and vibration side of the chamber

during the explore period (E), vibration period (F), and post-vibration period (G) of the

behavioral trial for four different conditions: naïve C57BL/6 mice (n=19 animals), naïve CD-1

mice (n=20 animals), CD-1 mice run on the behavior for the second time, one week later (n=20

animals), and naïve CD-1 mice that had been whisker trimmed (n=10 animals; ***$p<0.001$,

****$p<0.0001$, paired t-test).

H-J) Representative activity traces of two example wildtype CD-1 animals (H), littermate

control animals (I), and Ret cKO animals (J), tracking movement during the baseline and

vibration periods.

Extended Data Figure 10: Behavioral and anatomical characterization of LCIC silenced mice.

A) Post-hoc histology of three different TMR-X conjugated muscimol injected animals, showing restriction of the injection to the inferior colliculus.

B) Coronal sections of a representative TMR-X conjugated muscimol injected brain to show ventral spread of the injection. Scale bar = 500 μm.

C) Timeline of IC silencing experiments.

D) An additional example activity trace from a saline injected control and muscimol injected animal, showing movement during the baseline and vibration period of the behavioral trial.

E) Temperature preference assay done for naïve wildtype CD-1 animals (n=10 animals, ****p<0.0001, unpaired t-test). Note, this is a separate cohort from the group used in silencing experiments, reported in Fig. 4.

F) Texture preference assay done for naïve CD-1 animals (n=10 animals, ****p<0.0001, unpaired t-test). Note, this is a separate cohort from the group used in silencing experiments in Fig. 4.

E) Some types of sandpaper are aversive to mice due to their odor. To ensure our texture avoidance behavioral measure reflects avoidance of an aversive texture and not odor, the sandpaper was covered by a smooth paper with small holes punched in it, such that the odor of the paper could be sensed, but the texture could not be felt. In this condition, mice (n=10 animals) did not demonstrate an aversion to the sandpaper side. (Not significant, paired t-test.)

H) A model in which the LCIC integrates auditory cues detected by the cochlea and vibrotactile cues detected by Pacinian corpuscles to drive behavioral responses to vibration.

Methods

Animals

Avil^Cre, *TrkB^flox*, *Ret^flox(eGFP)*, and *Vglut2^Cre* mouse lines were of mixed background (CD-1 and C57BL/6J). *Gjb2^flox* and *Sox10^Cre* were on a C57BL/6 background. CD-1 animals were imported from Charles River (#022) and C57BL/6 animals were imported from The Jackson Laboratory (#000664). The following mouse lines have been previously described, including: *Sox10^Cre* (Ref: 77, Jax 025807), *Avil^Cre*(Ref: 78, JAX 032536), *TrkB^flox*(Ref:79), *Ret^flox(eGFP)* (JAX 029846), and *Vglut2^Cre* (JAX 016963). The *Gjb2^flox* mouse line, in which loxP sites flank the entire coding sequence in exon 2, was generated by KTB in the laboratory of DPC and will be described elsewhere. Mice were handled and housed in standard cages in accordance with Harvard Medical School and IACUC guidelines. Both male and female mice were used in all experiments. Control littermates of *Ret*, *TrkB,* and *Gjb2* conditional mutants included Cre-negative animals that were negative, heterozygous, or homozygous for the floxed allele. Our data indicated no difference among controls of different genotypes; therefore, they were combined for statistical analyses and collectively referred to as "controls."

Perfusions

Mice were anesthetized under isoflurane and transcardially perfused with 5-10 mL of PBS, followed by 5-10 mL of 4% paraformaldehyde (PFA) in PBS. After perfusion, the skull was removed and post-fixed in 4% PFA in PBS at 4°C overnight. Skin samples were removed and fixed overnight in Zamboni's fixation buffer with gentle agitation at 4°C. Samples were washed 3x10 min in PBS at room temperature after overnight fixation before fine dissection.

Immunohistochemistry of skin cryosections, brain cryosections and whole-mount tissue

The following dilutions and primary antibodies were used: chicken anti-NFH (Aves # NFH, 1:500), rabbit anti-S100 Beta (VWR/Protein Tech #15146-1-AP, 1:300), rabbit anti-dsRed (Takara #632496. 1:1000), chicken anti-GFP (Abcam #13970, 1:1000), goat anti-GFP (US Biological Life Sciences # G8965-01E, 1:500). The secondary antibodies used were Alexa 488, 546 or 647 conjugated donkey or goat anti-rabbit, chicken, or goat (Life Technologies or Jackson ImmunoResearch) and were prepared at 1:500 or 1:1000 dilutions.

For Meissner corpuscle histology, paws were removed from perfused mice. The pads and fingertips were clipped off of the glabrous skin using small spring scissors and cryoprotected in 30% sucrose in PBS at 4 °C for 1-2 days, embedded in OCT (1437365, Fisher), frozen using dry ice, and stored at −80 °C. The tissues were cryosectioned (25 μm) using a cryostat (Leica) and collected on glass slides (12-550-15, Fisher). Sections were washed 3 times for 5 minutes each with PBS containing 0.1% Triton X-100 (0.1% PBST), incubated with blocking solutions (0.1% PBST containing 5% normal goat serum (S-1000, Vector Labs) or normal donkey serum (005-000-121, Jackson Immuno) for 1 hour at room temperature, incubated with primary antibodies diluted in blocking solutions at 4 °C overnight, washed 3 times for 10 minutes each with 0.1% PBST, incubated with secondary antibodies diluted in blocking solutions at 4 °C overnight, washed again 4 times for 10 minutes each with 0.1% PBST (DAPI solution was included in the second wash at 1:5000 dilution), and mounted in Fluoromount-G mounting medium (Fisher 0100-01).

For whole-mount staining of Pacinian corpuscles, animals were anesthetized under isoflurane and euthanized by cervical dislocation. For Pacinian corpuscle dissections, the tibia and fibula of the ankle, and radius and ulna of the wrists were dissected immediately after euthanasia, keeping muscle and surrounding tissue intact, and fixed overnight in Zamboni's fixative at 4 °C. Limbs were washed 3x10 minutes in PBS and then finely dissected to remove muscle on the bone, while leaving the interosseous membrane of the bone intact. Tissues were then washed every 30 minutes for 5-6 hours with 1% PBST (PBS + triton X-100). Primary antibodies were applied in blocking solution (5% normal goat or donkey serum, 75% PBST, 20% DMSO, 0.01% NaN3) and incubated at room temperature for 3-5 nights with gentle rocking. Tissues were rinsed 3x in PBST, then washed every 30 minutes for 5-6 hours in 1% PBST. Secondary antibodies were applied in blocking solution and incubated at room temperature for 3-5 nights, with gentle rocking. Tissues were rinsed 3x in 1% PBST, washed every 30 minutes for 5-6 hours in 1% PBST, then serially dehydrated (10-15 minutes per wash) in 50%, 75%, and 100% ethanol diluted in distilled water. Tissue was stored at -20 °C in 100% ethanol until clearing and imaging. Tissue was cleared in BABB (1 part benzyl alcohol to 2 parts benzyl benzoate) for 2-12 hours at room temperature and mounted in BABB for imaging. For mounting, chambers were

made on slides consisting of a thin layer of vacuum grease to create four walls followed by placement of a coverslip. Tissue was imaged on a Zeiss LSM 900 confocal microscope using the 10x objective.

For brain sections, brains were cryoprotected in 30% sucrose overnight. Tissue was cryosectioned (50 uM sections) and collected in a 12-well plate filled with PBS. Sections were stained free floating using a 12-well plate. Sections were washed 3x5 minutes each with 1x PBS containing 0.1% Triton X-100 (0.1% PBST), incubated with blocking solutions (0.1% PBST containing 5% normal goat serum (S-1000, Vector Labs) or normal donkey serum (005-000-121, Jackson Immuno) for 1-2 hours at room temperature, incubated with primary antibodies diluted in blocking solutions at 4 °C overnight. Subsequently, sections were washed 3x10 minutes each with 0.1% PBST, incubated with secondary antibodies diluted in blocking solutions at room temperature for 2 hours, and washed again 4x10 minutes each with 0.1% PBST (DAPI solution was included in the second wash at 1:5000 dilution). Sections were mounted onto glass slides and coverslipped in Vectashield (Vector Labs).

Quantification of peripheral anatomy

For quantification of Meissner corpuscles in Ret cKO mice and littermate controls, cryosectioned and stained skin from the pads and fingertips were imaged with a widefield Zeiss LSM 700 microscope. Images of sections were analyzed in the Fiji distribution of Image J (www.fiji.sc). In each section, the number of Meissner corpuscles, visualized by S100 antibody staining, were counted. The length of the surface of the skin was traced and measured in ImageJ, and density was calculated by dividing the number of Meissner corpuscles by the length of the skin surface. Quantification was done blind to genotype.

For quantification of Pacinian corpuscles in Ret cKO mice, TrkB cKO mice, and littermate controls, images of stained and intact limbs were taken using a Zeiss LSM 900 confocal microscope using the 10x objective. The number of Pacinian corpuscles, visualized by S100 antibody staining, were counted. Quantification was done blind to genotype.

DRG recordings

Animals were anesthetized with urethane (1.5 g/kg) or isoflurane (1.5-2%). A skin incision was

made over the L3-L5 vertebrae and muscle surrounding the vertebrae was removed. The L3 and

L5 vertebrae were fixed in place using custom-built clamps. The L4 DRG was exposed by

removing overlaying bone. A borosilicate glass electrode (1 MΩ) was lowered into the DRG

using a motorized manipulator (Scientifica) to search for units. Recordings were amplified using

a MultiClamp 700B (Molecular Devices) under the 100x AC differential amplification mode

with additional 20x gain. Signals were collected with a 0.1 kHz high-pass filter and 3 kHz Bessel

filter, and acquired using a Digidata 1550B (Molecular Devices) sampled at 20 kHz. For

Pacinian corpuscle innervating Aβ-RA2-LTMRs, a search stimulus of 300 Hz vibration (40 mN)

was applied to the ankle. Pacinians were identified by their large receptive fields, and their end

organ location on the fibula was identified through electrical stimulation following

measurements of their vibration tuning. For Meissner corpuscle innervating Aβ-RA1-LTMRs,

the hindlimb digits were brushed as the electrode was descended. Aβ-RA1-LTMRs were

identified by their small skin receptive fields, adapting responses to indentation, low mechanical

thresholds (<10 mN), and cutaneous endings that could be excited by electrical stimulation of the

skin. Vibrations (10-900 Hz) of varying intensity (1-50 mN) were delivered to the most sensitive

area of the receptive field for each unit, collecting at least 4 trials per force for each frequency.

Craniotomies for MEA recordings

Mice were injected with sustained release buprenorphine (0.1 mg/kg), anesthetized with

isoflurane, and placed in a small animal stereotaxic frame (David Kopf Instruments). A titanium

headplate was cemented to the mouse's skull (C&B Metabond). A craniotomy was made

centered over the right inferior colliculus (stereotaxic coordinates: 0.5 mm posterior, 1.8 mm

lateral of lambda), or centered over the right VPL (stereotaxic coordinates: 1.7 mm posterior,

1.9 mm lateral of bregma), keeping the dura intact. Kwik-sil adhesive (World Precision

Instruments) was used to cover the craniotomy until the day of recording (within 3 days of the

craniotomy).

Multielectrode array (MEA) recordings

For anesthetized recordings, the mouse was injected intraperitoneally with urethane (0.5-

1.5 mg/g), placed on a heated platform, and head fixed such that the paws on the side

contralateral to the craniotomy were accessible for mechanical stimulation. MEAs were inserted

into the LCIC (H6b-32ch or H7b-32ch, Cambridge Neurotech) or VPL (H6b-32ch, H7b-32ch, or

E4-64ch, Cambridge Neurotech) with a 3-axes micromanipulator (IVM mini, Scientifica). MEA

signals were high-pass filtered at 200 Hz, amplified (RHD2132), and acquired at 20 kHz

(RHD2000, Intantech) for offline processing.

For VPL recordings, hindlimb VPL was targeted using coordinates (1.6 mm posterior, 1.9 mm

lateral to bregma), and the location was verified by brushing the hindlimb. The LCIC was

targeted by positioning the MEA probe posterior to the transverse sinus and ~1.6-1.8 mm lateral,

with a 45-degree angle to insert the probe underneath the transverse sinus. In preliminary

experiments, probe penetrations that contained only auditory responsive neurons were shown to

be in the CNIC, unlike penetrations in the LCIC that contained both mechanical and auditory

responses. Therefore, subsequent penetrations that contained only auditory responses were

assumed to be in the CNIC and were not included in the analysis. For verification of recording

location, after recording was complete, the probe was lifted from the brain, the probe was coated

with DiI (Thermo Fisher) then re-inserted to the same coordinates, and the brain was collected

and post-fixed overnight in 4% PFA post-recording. Recordings were performed and analyzed

blind to genotype.

Force-frequency tuning experiments (anesthetized): The receptive fields for neurons on the

probe were determined by gently brushing the digits or pads using a fine paintbrush. The paw

was fixed to the platform with Easy Mold Silicone Putty (Environmental Technologies). The

mechanical stimulator (with a blunt 1 mm probe) was then targeted to the receptive fields

identified by brush. In the case of LCIC neurons, which had large receptive fields that lacked a

hotspot, the probe was targeted to the heel of the hindlimb. As above, for verification of

recording location, after recording was complete, the probe was lifted from the brain, the probe

was coated with DiI (Thermo Fisher) then re-inserted to the same coordinates, and the brain was

collected and post-fixed overnight in 4% PFA post-recording. Thresholds for each vibratory

frequency were defined as the force required to elicit 20% of a unit's maximum firing during the

sustained portion of the stimulus (200 ms window beginning 50 ms after stimulus onset).

VPL and LCIC mapping experiments (anesthetized): For VPL mapping experiments (Extended Data Fig. 3) a 4 shanked probe (E4-64 ch, Cambridge Neurotech) was positioned with the shanks aligned to the anterio-posterior axis. Mapping began medially (at ~1.7 mm lateral) and moved 100 μm lateral for each successive penetration. For each probe penetration, the contralateral hindlimb, thigh, trunk, forelimb, and face were stroked with a paintbrush. The hindlimb was fixed to the platform with Easy Mold Silicone Putty and stimulation at 50 Hz and 500 Hz was delivered to the hindpaw with a 10 mN intensity using a mechanical stimulator with a wide blunt tip (3 mm diameter). Complementary LCIC experiments used a single shank probe (H6b or H7b, Cambridge Neurotech), with probe penetrations also separated by 100 μm intervals. Before each probe penetration the probe was coated with DiI and recording locations were verified post-experiment. Neurons were considered brush responsive to a body region if their change in firing exceeded 10 Hz during the brush period. Similarly, neurons were considered responsive to 500 Hz vibration if their change in firing exceeded 10 Hz during the sustained portion (200 ms time period beginning 50 ms after stimulus onset) of the stimulus.

Awake LCIC recordings: For awake LCIC recordings, animals were head-fixed, and the LCIC was targeted as described above. Mechanical stimulation was targeted to the platform the animal stood on rather than the limb. The platform was made of thin (1/8") acrylic. Trials where the animal was moving were excluded.

Mechanical and auditory stimuli

For *in vivo* electrophysiology, vibrations were generated using a DC motor with a blunt 1-mm or 3-mm diameter probe, as previously described[5,28]. The motor was driven by a custom-built current supply controlled by Wavesurfer and a multifunction I/O board (USB-6343, National Instruments), and static forces calibrated using a fine scale. For awake recordings, the mechanical stimulator was covered with a sound dampening foam tip, and placed underneath the acrylic platform the animal stands on. The magnitude of vibration was measured using an accelerometer attached to the bottom of the platform and reported in units of gravitational force (g). Brush stimuli were manually delivered with a 4 mm diameter brush at a rate of one stroke per 2 seconds in the rostral to caudal direction.

Auditory stimuli were amplified by an SA1 stereo power amplifier (Tucker-Davis Technologies)

and delivered through an MF1 multi-field magnetic speaker (Tucker-Davis Technologies),

positioned 14 cm from the ear contralateral to recording site. Auditory stimuli were generated in

MATLAB and delivered using WaveSurfer and a multifunction I/O board (USB-6343, National

Instruments). Broadband (white) noise (0-50 kHz) was delivered in 100 ms pulses (5 ms rise-fall

time) at 65 dB SPL. White noise, vibration, and combined trials were interleaved. For awake

recordings, trials where the animal was moving were excluded.

Spike detection and cluster assignment

Data was converted into contiguous binary files (https://github.com/peltonen/kwik-tools) and

processed using JRClust (Janelia). Spikes were detected and assigned to clusters corresponding

to individual or multiples units with JRClust (https://github.com/JaneliaSciComp/JRCLUST).

These initial clusters were manually curated to remove artifacts that did not correspond to spike

waveforms, and clusters were examined and split if their PCA decompositions were not

monodispersed. This collection of clusters was then examined and merge operations performed if

consideration of the average spike waveform and spiking cross-correlation indicated that two

clusters corresponded to the same neuron. Manual curation typically resulted in putative isolated

single units with intra-cluster waveform correlations greater than 0.9, inter-cluster waveform

correlations less than 0.1, and clusters that did not meet these criteria were not considered in the

dataset. Spike event times for each cluster were exported and processed in Python.

Anterograde tracing experiments

For anterograde tracing experiments (Extended Data Fig. 1), brainstem DCN neurons of

Slc17a6^{Cre} (Vglut2) mice were transduced with either an AAV encoding synaptophysin-GFP,

AAV1-CAG-FLEX-Synaptophysin-GFP-WPRE (1.088E+14gc/ml), or synaptophysin-tDT,

AAV1-CAG-FLEX-Synaptophysin-tdTomato-WPRE (1.1482E+14gc/ml). Viruses were

produced and packaged at the Boston Children's Hospital Viral Core Facility. Mice were

injected with sustained release buprenorphine (0.1 mg/kg), anesthetized with isoflurane and

placed in a small animal stereotaxic frame (David Kopf Instruments) with their neck bent at a 45

degree angle. The dorsal column nuclei were exposed between the base of the skull and C1

through retraction of the paraspinal muscles. A small nick in the overlying dural membrane was

made to allow access for the injection pipette. A borosilicate glass pipette was lowered to a depth

of 0.5 mm in either the gracile (0.1 mm lateral to obex) or cuneate (0.5mm anterior to obex, 0.1

mm lateral to fourth ventricle) nucleus. 50-100 nL of AAV virus with fast green was injected at a

rate of 50 nL min−1 using a Microinject system (World Precision Instruments). Three weeks

after injection, mice were perfused, and brain tissue was collected for cryosectioning and

immunohistochemistry.

Auditory brainstem responses (ABR) and distortion product otoacoustic emissions

(DPOAE)

All measurements and analyses were conducted blind to the animal's experimental condition.

For anesthesia, mice were given an IP injection of ketamine (100 mg/kg body weight, BW),

xylazine (10 mg/kg BW) and acepromazine (0.3 mg/kg BW) and placed on their side on a 37°C

metal heating pad (ATC-2000, World Precision Instruments). Experiments were performed in a

sound-attenuating, electrically shielded chamber (IAC Acoustics). After anesthesia induction,

eye lubricant (GenTeal Tears, Alcon) was applied to prevent drying. A ketamine boost ($0.3 - 0.5$

mg/kg BW) was used to maintain anesthesia at the first sign of movement or elevation in the

signal rejection rate (rejection threshold was 15 μV).

An acoustic system consisting of a microphone and dual earphone assembly coupled to a probe

tube was used to deliver and measure acoustic stimuli. This custom system was built by the

Engineering Core of the Eaton-Peabody Laboratories (EPL) at Massachusetts Eye and Ear

(MEE) (Boston). The probe end of the acoustic system was placed ~1 mm over the center of the

ear canal. Stimuli were digitally generated and then amplified by an SA1 stereo power amp

(Tucker Davis Technologies) to drive the acoustic system. The probe tube microphone was

amplified by a custom microphone amplifier (EPL at MEE) and sent to a 24-bit I/O board (PXI-

4461) installed into a PXI chassis (PXIe-1073, National Instruments).

Bioelectrical signals were recorded using three subdermal electrodes: an active electrode at the

vertex, a reference electrode near the pinna, and a ground electrode near the tail. Signals were

passed through a custom pre-amplifier (EPL at MEE) before digitization. Acoustic stimuli

consisted of clicks and 5 ms tone pips with a repetition rate of 30 s^{-1} and a 0.5 ms rise-fall time at

the following frequencies (in kHz): 8, 16, 22.6, and 32. Stimuli were presented in alternating

polarity to cancel out the cochlear microphonic potential. Cochlear Function Test Suite (CFTS)

software (version 2.37.2; MEE, Boston) amplified (x104), filtered (0.3–3 kHz passband), and

averaged 512 responses at each sound pressure level (SPL). ABR thresholds were manually

designated blind to genotype using the ABR Peak Analysis Software from the EPL at MEE.

DPOAEs were recorded in response to primary tones f1 and f2 at a frequency ratio of $f2/f1 = 1.2$,

and a SPL level of $L1 = L2 + 10$ dB. For individual DPOAE trials, f2 consisted of 5.6, 8, 11.3,

16, 22.6, 32, 42.5, or 64 kHz tones that were incremented in 10 dB steps from 10 to 70 dB SPL.

The cubic distortion product 2f1-f2 was extracted by Fourier analysis of the ear-canal sound

pressure after waveform and spectral averaging. DPOAE threshold was defined as the f1 level

required to produce a DPOAE of 5 dB SPL, except at 64 kHz where the threshold was defined as

10 dB SPL due to a higher noise floor.

Muscimol silencing

For muscimol silencing experiments, mice were injected with sustained release buprenorphine

(0.1 mg/kg), anesthetized with isoflurane, and placed in a small animal stereotaxic frame (David

Kopf Instruments). A small section of skin above the rear of the skull was removed with surgical

scissors and the skull was dried. The skin was glued to the skull using Vetbond (3M healthcare),

and a small amount of dental cement (C&B Metabond) was used to cover the exposed region of

the skull, leaving the area where the craniotomy would be drilled clear. A small craniotomy was

drilled approximately 0.5 mm posterior, 1.8 mm lateral to lambda. The craniotomy was covered

with Kwik-sil adhesive.

Animals were given 3-4 days to recover. On the day of the behavioral experiment, animals were

anesthetized with isoflurane and placed in a small animal stereotaxic frame. A borosilicate glass

pipette was used to bilaterally inject into the inferior colliculus (0.5-0.8 mm posterior to lambda,

1.5-1.8 mm lateral, 0.5 mm deep) with TMR-X muscimol dissolved in ACSF (1.5 mM,

Invitrogen M23400). 50 nL injections were made in two locations per side (100 nL total per

side). Control mice received the same injection volume of sterile saline. Injections were

performed at a rate of 50 nL min−1 using a Microinject system (World Precision Instruments)

and the pipette was allowed to sit in the brain for 3 minutes post-injection. Animals were allowed

to recover from isoflurane and habituate to the behavior room for ~1 hour, and the behavioral

assay (vibration, temperature, or texture) was run within 2 hours. Behavioral assays were run

blind to experimental group (saline or muscimol). Post-behavior, animals were perfused.

Muscimol spread was confirmed post-hoc following vibratome sectioning the brain and imaging.

Behavior methods for Openfield, Vibration Platform and Temperature Preference assays

To reduce anxiety and improve handling, animals were enriched with Bio-Serv igloos, tunnels or

chew shacks at least one month prior to all behavior testing. One week prior to behavioral

testing, animals were moved to the testing room and home cage habituated to the behavior room

to acclimate to room lighting (warm white, 2700k), sound (ambient 45-50 dB), odor (cleaned

with ECOS unscented dish soap and 70% EtOH), and temperature (73 °F, 30% humidity).

Vibration, temperature, and texture assays were conducted in 3-4 month old CD-1 animals

(unless otherwise noted). Vibration assays in Ret cKO and littermate controls were performed at

8 weeks of age and were performed blind to genotype.

Vibration Platform and Sound Only Test: Two days prior to testing, animals were moved to the

testing room and habituated to investigator handling. Animals were group habituated to the black

matte test chamber (12 in L x 6 in W x 7 in H x 0.25 in), center divided with an arched doorway

and allowed to freely explore for 5 minutes. During the three-day room/chamber habituation,

animals were also habituated to the vibration motor sound only. The vibration platform was

elevated 6 inches above the base. Two flat surface vibration motors, (PUI Audo, ASX08604-

SW-R 4 ohm, BST) were suspended above the base, resting on spring mounted posts, one motor

on each side of the chamber to act as the suspension base for the two square white matte acrylic

floors, (5.5 in L x 5.5 in W x 0.125 in D) to rest on and provide vibration stimulus to each floor

when activated. The acrylic floors were separated by a thin gap and lightly attached to the

motor. An accelerometer was attached to each floor to ensure vibration floor displacement was

constant for each test animal. The entire floor was covered with a thin stretch white matte cloth

napkin (WypAll L30, 05812) to prevent the animals from investigating the gap in the floor and

edges of the chamber walls as well as maintain a clean odor free test environment. The black

matte acrylic test chamber rested on four pillars (6 inches above the floor) to ensure the chamber

walls snugly surrounded the vibration floors without direct contact to prevent the

chamber from vibrating when the motor was activated during testing. The room and test platform

were lined with odorless 1-inch thick Bonded Logic Natural Fiber Acoustic Sound

Absorbing panels (Model #60600-11212).

Vibration Test: The test chamber was modified on test day by removing the front and side wall

to create a semi-open and closed end (L-shape) to increase animals' awareness of the floor

surface when exploring both sides of the test chamber. The center divider with open doorway

was left in place. The semi-closed side of the chamber was the non-stimulus side and the semi-

open side was the vibration stimulus side. A custom Bonsai script was used to trigger the

vibration motor and camera. The three-minute vibration test sequence was as follows: The

animal was placed in the chamber facing the semi-open side which encouraged the animal to

walk through the doorway and explore the newly semi-closed side of the chamber.

Stage-1: Start, allow one minute baseline explore of test chamber,

Stage-2: 30 second vibration stimulus on the semi-open side,

Stage-3: 30 second interim period with no stimulus,

Stage-4: 30 second vibration stimulus on the semi-open side,

Stage-5: 30 second final period with no stimulus.

For the sound only test, the motor was disconnected from floor contact and the motor sound only

was played for the 30 second stimulus during stage-2 and stage-4. Ambient room and chamber

decibel level was 45-50 dB. During sound only and vibration stimulus periods, the background

sound in the testing room and chamber was set to 70 dB at 500 Hz. A top video recording device

was used to track distance traveled and time exploring the two chambers.

Temperature Preference Test: Two days prior to testing, animals were moved to the testing room

and home cage habituated to the behavior. Animals were group habituated for 5 minutes to the

black matte test chamber (12 in L x 6 in W x 7 in H x 0.25 in), center divided with an arched

doorway, resting on two metal plate floors (5.5 in L x 5.5 in W ea) both set to neutral

temperature 30 °C. On the third day following room habituation, each animal was tested on

contrasting temperatures, 30 °C (warm) vs 18 °C (cold). Each animal was placed in the test

chamber on the warm side facing the doorway to the cold side to encourage the animals to walk

through the doorway and freely explore both temperatures for 5 minutes. A top video recording

device was used to track distance traveled and time exploring the two chambers.

Texture Preference Assay: Two days prior to testing, animals were moved to the testing room

and home cage habituated to the behavior. Animals were group habituated for 10 minutes to the

black matte test chamber (12 in L x 6 in W x 7 in H x 0.25 in) with no divider. On the third day

following room habituation, each animal was tested on contrasting floors, construction paper

(Tru-Ray Heavyweight, Pacon) vs coarse 60-grit sandpaper (ceramic version, 26060PGP-4,

3M). There was no divider between the smooth and rough textured floors. Each animal

was placed in the test chamber on the smooth side and animals were given 10 minutes to freely

explore both chambers. A top video recording was used to track distance traveled and time

exploring the two chambers.

Quantification and statistical analysis

Statistical tests were conducted in Python 3.7.9 using the SciPy stats module or in GraphPad

Prism. Normality of the data was tested using the Shapiro-Wilk test. Comparisons between two

independent groups were performed using the unpaired t-test (in the case of two groups

distributed normally, with no significant difference between variances), Welch's t-test (in the

case of two groups distributed normally, but with unequal variances), or Mann-Whitney test

(non-parametric data). When correction for multiple comparisons was necessary, the Bonferroni

correction was used. An adjusted p value < 0.05 was considered significant. Additional details on

sample sizes and statistical tests for each experiment can be found in figure legends and the main

text.

References

References

1.   Pickles JO. Auditory pathways: anatomy and physiology. *Handb Clin Neurol.*
2015;129:3-25.

2.   Handler A, Ginty DD. The mechanosensory neurons of touch and their mechanisms of
activation. *Nat Rev Neurosci.* 2021;22(9):521-537.

3.   Sato M. Response of Pacinian corpuscles to sinusoidal vibration. *J Physiol.*
1961;159(3):391-409.

4.   Bolanowski SJ, Jr., Zwislocki JJ. Intensity and frequency characteristics of pacinian
corpuscles. II. Receptor potentials. *J Neurophysiol.* 1984;51(4):812-830.

5.   Turecek J, Ginty DD. Coding of self and environment by Pacinian neurons in freely
moving animals. *bioRxiv.* 2023.

6.   Cividini S, Sfenthourakis S, Montesanto G. Are terrestrial isopods able to use stridulation
and vibrational communication as forms of intra and interspecific signaling and defense
strategies as insects do? A preliminary study in Armadillo officinalis.
Naturwissenschaften. 2019;107(1):4.

7.   Narins PM, Reichman OJ, Jarvis JU, Lewis ER. Seismic signal transmission between
burrows of the Cape mole-rat, Georychus capensis. *J Comp Physiol A.* 1992;170(1):13-
21.

8.   Nevo E, Heth G, Pratt H. Seismic communication in a blind subterranean mammal: a
major somatosensory mechanism in adaptive evolution underground. *Proc Natl Acad Sci*
U S A. 1991;88(4):1256-1260.

9.   O'Connell-Rodwell CE. Keeping an "ear" to the ground: seismic communication in
elephants. *Physiology (Bethesda).* 2007;22:287-294.

10.   Virant-Doberlet M, Stritih-Peljhan N, Zunic-Kosi A, Polajnar J. Functional Diversity of
Vibrational Signaling Systems in Insects. *Annu Rev Entomol.* 2023;68:191-210.

11.   Johnson KO. The roles and functions of cutaneous mechanoreceptors. *Curr Opin*
Neurobiol. 2001;11(4):455-461.

12.   Heffner RS, Koay G, Heffner HE. Audiograms of five species of rodents: implications for
the evolution of hearing and the perception of pitch. *Hear Res.* 2001;157(1-2):138-152.

13.   Heffner HE, Heffner RS. Hearing ranges of laboratory animals. *J Am Assoc Lab Anim*
Sci. 2007;46(1):20-22.

14. Yau JM, Olenczak JB, Dammann JF, Bensmaia SJ. Temporal frequency channels are
linked across audition and touch. *Curr Biol.* 2009;19(7):561-566.

15. Neubarth NL, Emanuel AJ, Liu Y, et al. Meissner corpuscles and their spatially
intermingled afferents underlie gentle touch perception. *Science.* 2020;368(6497).

16. Prsa M, Morandell K, Cuenu G, Huber D. Feature-selective encoding of substrate
vibrations in the forelimb somatosensory cortex. *Nature.* 2019;567(7748):384-388.

17. Lee K-S, Wagner DdT, Sanders M, Huber D. Emergence of a somatosensory tonotopic
map for substrate vibration in the brainstem. *bioRxiv.* 2023.

18. Bouley DM, Alarcon CN, Hildebrandt T, O'Connell-Rodwell CE. The distribution,
density and three-dimensional histomorphology of Pacinian corpuscles in the foot of the
Asian elephant (Elephas maximus) and their potential role in seismic communication. *J
Anat.* 2007;211(4):428-435.

19. Germann C, Sutter R, Nanz D. Novel observations of Pacinian corpuscle distribution in
the hands and feet based on high-resolution 7-T MRI in healthy volunteers. *Skeletal
Radiol.* 2021;50(6):1249-1255.

20. Kumamoto K, Senuma H, Ebara S, Matsuura T. Distribution of pacinian corpuscles in the
hand of the monkey, Macaca fuscata. *J Anat.* 1993;183 (Pt 1)(Pt 1):149-154.

21. Silfvenius H. Characteristics of receptors and afferent fibers of the forelimb interosseous
nerve of the cat. *Acta Physiol Scand.* 1970;79(1):6-23.

22. Zelena J. The development of Pacinian corpuscles. *J Neurocytol.* 1978;7(1):71-91.

23. Hulliger M, Nordh E, Thelin AE, Vallbo AB. The responses of afferent fibres from the
glabrous skin of the hand during voluntary finger movements in man. *J Physiol.*
1979;291:233-249.

24. Salinas E, Hernandez A, Zainos A, Romo R. Periodicity and firing rate as candidate
neural codes for the frequency of vibrotactile stimuli. *J Neurosci.* 2000;20(14):5503-
5515.

25. Luna R, Hernandez A, Brody CD, Romo R. Neural codes for perceptual discrimination in
primary somatosensory cortex. *Nat Neurosci.* 2005;8(9):1210-1219.

26. Harvey MA, Saal HP, Dammann JF, 3rd, Bensmaia SJ. Multiplexing stimulus
information through rate and temporal codes in primate somatosensory cortex. *PLoS Biol.*
2013;11(5):e1001558.

27. Abraira VE, Ginty DD. The sensory neurons of touch. *Neuron.* 2013;79(4):618-639.

28. Turecek J, Lehnert BP, Ginty DD. The encoding of touch by somatotopically aligned

dorsal column subdivisions. *Nature.* 2022;612(7939):310-315.

29. Aitkin LM, Dickhaus H, Schult W, Zimmermann M. External nucleus of inferior

colliculus: auditory and spinal somatosensory afferents and their interactions. *J*

Neurophysiol. 1978;41(4):837-847.

30. Aitkin LM, Kenyon CE, Philpott P. The representation of the auditory and somatosensory

systems in the external nucleus of the cat inferior colliculus. *J Comp Neurol.*

1981;196(1):25-40.

31. Aitkin LM, Webster WR, Veale JL, Crosby DC. Inferior colliculus. I. Comparison of

response properties of neurons in central, pericentral, and external nuclei of adult cat. *J*

Neurophysiol. 1975;38(5):1196-1207.

32. Bjorkeland M, Boivie J. An anatomical study of the projections from the dorsal column

nuclei to the midbrain in cat. *Anat Embryol (Berl).* 1984;170(1):29-43.

33. Jain R, Shore S. External inferior colliculus integrates trigeminal and acoustic

information: unit responses to trigeminal nucleus and acoustic stimulation in the guinea

pig. *Neurosci Lett.* 2006;395(1):71-75.

34. Lesicko AM, Hristova TS, Maigler KC, Llano DA. Connectional Modularity of Top-

Down and Bottom-Up Multimodal Inputs to the Lateral Cortex of the Mouse Inferior

Colliculus. *J Neurosci.* 2016;36(43):11037-11050.

35. Oliver DL, Morest DK. The central nucleus of the inferior colliculus in the cat. *J Comp*

Neurol. 1984;222(2):237-264.

36. Syka J, Popelar J, Kvasnak E, Astl J. Response properties of neurons in the central

nucleus and external and dorsal cortices of the inferior colliculus in guinea pig. *Exp Brain*

Res. 2000;133(2):254-266.

37. Zhou J, Shore S. Convergence of spinal trigeminal and cochlear nucleus projections in

the inferior colliculus of the guinea pig. *J Comp Neurol.* 2006;495(1):100-112.

38. Lehnert BP, Santiago C, Huey EL, et al. Mechanoreceptor synapses in the brainstem

shape the central representation of touch. *Cell.* 2021;184(22):5608-5621 e5618.

39. Berkley KJ, Budell RJ, Blomqvist A, Bull M. Output systems of the dorsal column nuclei

in the cat. *Brain Res.* 1986;396(3):199-225.

40.   Francis JT, Xu S, Chapin JK. Proprioceptive and cutaneous representations in the rat
1098   ventral posterolateral thalamus. *J Neurophysiol.* 2008;99(5):2291-2304.

1099   41.   Horne MK, Tracey DJ. The afferents and projections of the ventroposterolateral thalamus
1100   in the monkey. *Exp Brain Res.* 1979;36(1):129-141.

1101   42.   Kaas JH, Nelson RJ, Sur M, Dykes RW, Merzenich MM. The somatotopic organization
1102   of the ventroposterior thalamus of the squirrel monkey, Saimiri sciureus. *J Comp Neurol.*
1103   1984;226(1):111-140.

1104   43.   Mountcastle VB, Henneman E. The representation of tactile sensibility in the thalamus of
1105   the monkey. *J Comp Neurol.* 1952;97(3):409-439.

1106   44.   Handler A, Zhang Q, Pang S, et al. Three-dimensional reconstructions of
1107   mechanosensory end organs suggest a unifying mechanism underlying dynamic, light
1108   touch. *Neuron.* 2023;111(20):3211-3229 e3219.

1109   45.   Qi L, Iskols M, Handler A, Ginty DD. Krause corpuscles of the genitalia are vibrotactile
1110   sensors required for normal sexual behavior. *bioRxiv.* 2023.

1111   46.   Saal HP, Bensmaia SJ. Touch is a team effort: interplay of submodalities in cutaneous
1112   sensibility. *Trends Neurosci.* 2014;37(12):689-697.

1113   47.   Santiago C, Sharma N, Africawala N, et al. Activity-dependent development of the body's
1114   touch receptors. *bioRxiv.* 2023.

1115   48.   Alvarez-Buylla R, Ramirez de Arellano J. Local responses in Pacinian corpuscles. *Am J*
1116   *Physiol.* 1953;172(1):237-244.

1117   49.   Bell J, Bolanowski S, Holmes MH. The structure and function of Pacinian corpuscles: a
1118   review. *Prog Neurobiol.* 1994;42(1):79-128.

1119   50.   Hunt CC, Mc IA. Characteristics of responses from receptors from the flexor longus
1120   digitorum muscle and the adjoining interosseous region of the cat. *J Physiol.*
1121   1960;153(1):74-87.

1122   51.   Mountcastle VB, Talbot WH, Darian-Smith I, Kornhuber HH. Neural basis of the sense
1123   of flutter-vibration. *Science.* 1967;155(3762):597-600.

1124   52.   Talbot WH, Darian-Smith I, Kornhuber HH, Mountcastle VB. The sense of flutter-
1125   vibration: comparison of the human capacity with response patterns of mechanoreceptive
1126   afferents from the monkey hand. *J Neurophysiol.* 1968;31(2):301-334.

1127   53.   Luo W, Enomoto H, Rice FL, Milbrandt J, Ginty DD. Molecular identification of rapidly
1128   adapting mechanoreceptors and their developmental dependence on ret signaling.
1129   *Neuron.* 2009;64(6):841-856.
1130   54.   Luo W, Wickramasinghe SR, Savitt JM, Griffin JW, Dawson TM, Ginty DD. A
1131   hierarchical NGF signaling cascade controls Ret-dependent and Ret-independent events
1132   during development of nonpeptidergic DRG neurons. *Neuron.* 2007;54(5):739-754.
1133   55.   Emanuel AJ, Lehnert BP, Panzeri S, Harvey CD, Ginty DD. Cortical responses to touch
1134   reflect subcortical integration of LTMR signals. *Nature.* 2021;600(7890):680-685.
1135   56.   Gottschaldt KM, Vahle-Hinz C. Merkel cell receptors: structure and transducer function.
1136   *Science.* 1981;214(4517):183-186.
1137   57.   Ribot-Ciscar E, Vedel JP, Roll JP. Vibration sensitivity of slowly and rapidly adapting
1138   cutaneous mechanoreceptors in the human foot and leg. *Neurosci Lett.* 1989;104(1-
1139   2):130-135.
1140   58.   Chirila AM, Rankin G, Tseng SY, et al. Mechanoreceptor signal convergence and
1141   transformation in the dorsal horn flexibly shape a diversity of outputs to the brain. *Cell.*
1142   2022;185(24):4541-4559 e4523.
1143   59.   Duque D, Perez-Gonzalez D, Ayala YA, Palmer AR, Malmierca MS. Topographic
1144   distribution, frequency, and intensity dependence of stimulus-specific adaptation in the
1145   inferior colliculus of the rat. *J Neurosci.* 2012;32(49):17762-17774.
1146   60.   Perez-Gonzalez D, Malmierca MS, Covey E. Novelty detector neurons in the mammalian
1147   auditory midbrain. *Eur J Neurosci.* 2005;22(11):2879-2885.
1148   61.   Takada Y, Beyer LA, Swiderski DL, et al. Connexin 26 null mice exhibit spiral ganglion
1149   degeneration that can be blocked by BDNF gene therapy. *Hear Res.* 2014;309:124-135.
1150   62.   Yu Q, Wang Y, Chang Q, et al. Virally expressed connexin26 restores gap junction
1151   function in the cochlea of conditional Gjb2 knockout mice. *Gene Ther.* 2014;21(1):71-80.
1152   63.   Guo J, Ma X, Skidmore JM, et al. Gjb2 gene therapy and conditional deletion reveal
1153   developmental stage-dependent effects on inner ear structure and function. *Mol Ther
1154   Methods Clin Dev.* 2021;23:319-333.
1155   64.   Lalwani AK. Evaluation of childhood sensorineural hearing loss in the post-genome
1156   world. *Arch Otolaryngol Head Neck Surg.* 2002;128(1):88-89.

65.   Lee MY, Takada T, Takada Y, et al. Mice with conditional deletion of Cx26 exhibit no
vestibular phenotype despite secondary loss of Cx30 in the vestibular end organs. *Hear*
Res. 2015;328:102-112.

66.   Xiong XR, Liang F, Zingg B, et al. Auditory cortex controls sound-driven innate defense
behaviour through corticofugal projections to inferior colliculus. *Nat Commun.*
2015;6:7224.

67.   Allen TA, Narayanan NS, Kholodar-Smith DB, Zhao Y, Laubach M, Brown TH. Imaging
the spread of reversible brain inactivations using fluorescent muscimol. *J Neurosci*
Methods. 2008;171(1):30-38.

68.   Arikan R, Blake NM, Erinjeri JP, Woolsey TA, Giraud L, Highstein SM. A method to
measure the effective spread of focally injected muscimol into the central nervous system
with electrophysiology and light microscopy. *J Neurosci Methods.* 2002;118(1):51-57.

69.   Kwon SE, Yang H, Minamisawa G, O'Connor DH. Sensory and decision-related activity
propagate in a cortical feedback loop during touch perception. *Nat Neurosci.*
2016;19(9):1243-1249.

70.   Sathian K. Tactile sensing of surface features. *Trends Neurosci.* 1989;12(12):513-519.

71.   Park JM, Hong YK, Rodgers CC, et al. Deep and superficial layers of the primary
somatosensory cortex are critical for whisker-based texture discrimination in mice
bioRxiv. 2022.

72.   Stuttgen MC, Schwarz C. Barrel cortex: What is it good for? *Neuroscience.* 2018;368:3-
16.

73.   Hong YK, Lacefield CO, Rodgers CC, Bruno RM. Sensation, movement and learning in
the absence of barrel cortex. *Nature.* 2018;561(7724):542-546.

74.   Meredith MA, Stein BE. Interactions among converging sensory inputs in the superior
colliculus. *Science.* 1983;221(4608):389-391.

75.   Holmes NP, Spence C. Multisensory integration: space, time and superadditivity. *Curr*
Biol. 2005;15(18):R762-764.

76.   King AJ, Palmer AR. Integration of visual and auditory information in bimodal neurones
in the guinea-pig superior colliculus. *Exp Brain Res.* 1985;60(3):492-500.

77.   Matsuoka T, Ahlberg PE, Kessaris N, et al. Neural crest origins of the neck and shoulder.
Nature. 2005;436(7049):347-355.

78.   Hasegawa H, Abbott S, Han BX, Qi Y, Wang F. Analyzing somatosensory axon

projections with the sensory neuron-specific Advillin gene. *J Neurosci.*

2007;27(52):14404-14414.

79.   Liu Y, Rutlin M, Huang S, et al. Sexually dimorphic BDNF signaling directs sensory

innervation of the mammary gland. *Science.* 2012;338(6112):1357-1360.

1193

Acknowledgements

We thank A. Handler, H. Kwak, K. Lezgiyeva, R. Martinez-Garcia, J. Peng, K. Ng, A. Emanuel, and A. Shuster for comments on the manuscript. We thank A. Emanuel for code used for the MEA data analysis. This work was supported by a HHMI Hannah Gray fellowship (GER), NEI P30 Core Grant for Vision Research #EY012196 (OM), NIH grants F31 NS097344 (ELH) and R35 5R35NS097344-05 (DDG), and the Edward R. and Anne G. Lefler Center for Neurodegenerative Disorders (DDG). DDG is an investigator of the Howard Hughes Medical Institute. This article is subject to HHMI's Open Access to Publications policy. HHMI lab heads have previously granted a nonexclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their research articles. Pursuant to those licenses, the author-accepted manuscript of this article can be made freely available under a CC BY 4.0 license immediately upon publication.

Author Contributions

ELH and DDG conceived the study. ELH did the MEA recordings in both anesthetized and awake conditions, with help from JT and MD. JT and ZKS performed primary sensory neuron recordings. ELH, AH, and MD performed anatomy and immunohistochemistry experiments. MMD, OM, and ELH developed the vibration behavioral assay. ELH did the saline and muscimol IC injections, and MMD performed behavioral experiments for vibration, temperature, and texture preference assays, with help from ELH. GER performed auditory brainstem response (ABR) measurements with guidance from LVG. KTB and DPC generated and provided the *Gjb2^{flox/flox} Sox10^{Cre}* mice. ELH and DDG wrote the manuscript with input from all authors.