High-resolution analysis of the treated coeliac disease microbiome reveals increased inter-individual variability

Authors:

Jelle Slager[1], Hanna L. Simpson[1], Ranko Gacesa[1,2], Lianmin Chen[1,3,4,5], Jody Gelderloos[1], Astrid Maatman[1], Cisca Wijmenga[1], Alexandra Zhernakova[1], Jingyuan Fu[1,3], Rinse Weersma[2], Gieneke Gonera[1,6], Iris H. Jonkers[1], Sebo Withoff[1,*]

* Corresponding author: s.withoff@umcg.nl
1. Department of Genetics, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands
2. Department of Gastroenterology and Hepatology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands
3. Department of Pediatrics, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands
4. Department of Cardiology, Nanjing Medical University, The First Affiliated Hospital of Nanjing Medical University, Nanjing, China
5. Cardiovascular Research Center, The Affiliated Suzhou Hospital of Nanjing Medical University, Suzhou Municipal Hospital, Gusu School, Nanjing Medical University, Suzhou, China
6. Department of Pediatrics, Wilhelmina Hospital Assen, Assen, the Netherlands

Highlights:

- Shotgun metagenomic sequencing of the largest coeliac disease (CeD) patient cohort to date provides insights into gut microbiome composition down to the strain level.
- Novel associations between tCeD (treated CeD; adhering to a gluten-free diet) were identified.
- tCeD patients have a less uniform microbiome structure.
- Bacteria display higher mutation frequency in tCeD, compared to controls.
- The European subspecies of *Eubacterium rectale* is almost absent in tCeD patients.

Abstract:

Background: Coeliac disease (CeD) is a prevalent immune-mediated disorder primarily affecting the small intestine, characterised by a breakdown of tolerance to dietary gluten. Its onset results from the multifaceted interplay of genetic and environmental factors. Recent data show that alterations in gut microbiome composition could play an important role in CeD. However, many current studies have been constrained by small sample sizes and limited resolution.

Methods: To address these limitations, faecal gut microbiota from two Dutch patient cohorts, CeDNN (129 treated CeD patients; tCeD and 106 non-CeD controls) and the Lifelines Dutch Microbiome Project (24 self-reported treated CeD and 659 non-CeD controls), was analyzed using shotgun metagenomic sequencing. Controls were matched by age, sex, and sequencing depth. Self-reported IBS and IBD were used as comparative conditions of the gastrointestinal tract. Inter-individual variation among cases and controls was calculated for the whole microbiome and on the strain-level. Finally, species-specific gene repertoires were analysed in tCeD patients and controls.

Results: Alpha diversity was decreased in patients with self-reported IBS and IBD, but not in tCeD. Each condition displayed a unique microbial pattern, with minimal overlap in differentially abundant species. In addition to confirming existing studies, we identified an increase in the abundance of previously unreported species, including *Clostridium sp. CAG:253*, *Roseburia hominis*, and *Clostridium lavalense*. We demonstrate that changes in tCeD gut microbiome can be, at least partly, explained by adherence to a GFD and report an increased inter-individual variation in gut microbiome composition among tCeD patients. More detailed analyses revealed that a higher bacterial mutation frequency in CeD also leads to a higher inter-individual variation on the strain-level. In line with this, we found that the immotile European subspecies of *Eubacterium rectale*, with distinct carbohydrate metabolism potential, was nearly absent in tCeD patients.

Conclusion: Our study sheds light on the complex interplay between the gut microbiome and CeD, revealing unique microbial patterns and increased inter-individual variation among tCeD patients. These findings expand our understanding of the role of the microbiome in intestinal health and disease.

Keywords:

Coeliac disease, gut microbiome, metagenomic sequencing, gluten, gluten-free diet

Introduction:

Coeliac disease (CeD) is a multifactorial immune-mediated disorder characterised by an inflammatory response in the small intestine (1). CeD arises in genetically-susceptible individuals and is triggered by a loss of tolerance to dietary gluten—a storage protein found in grains such as wheat, barley, and rye (1–3). Approximately 1-2% of the global population is affected by CeD, with prevalence varying depending on the carriage of risk human leukocyte antigen (HLA) haplotypes and the extent of gluten consumption (4,5). In CeD patients, an inflammatory immune response is initiated by the translocation of partially digested gluten peptides from the intestinal lumen into the lamina propria. The peptides then undergo deamidation by tissue transglutaminase (tTG2), conferring an increased binding affinity to class II HLA-DQ2 or -DQ8 molecules on antigen-presenting cells which leads to the potent activation of gluten-specific CD4+ T cells upon peptide recognition (6–8). Subsequently, a complex cytokine-mediated cascade ensues, ultimately resulting in the impairment of the intestinal epithelial barrier by intraepithelial lymphocytes (9). Currently, strict life-long adherence to a gluten-free diet (GFD) is the only available treatment option, which requires considerable patient education and monitoring (10).

CeD is a complex disorder influenced by a combination of genetic and environmental factors. While the presence of the HLA-DQ2/DQ8 haplotypes constitutes the primary genetic risk factor, approximately 30% of the general population carries these haplotypes without developing CeD. Moreover, genome-wide association studies (GWAS) have identified an additional 39 non-HLA CeD risk loci, primarily associated with T cell activation and innate immunity (11–13). However, these non-HLA loci contribute to only 14% of the genetic variance in CeD (14). These findings indicate that while genetic susceptibility is necessary, it alone is insufficient for the development of CeD. Thus, other non-genetic factors likely play a role in the propagation of an inflammatory response to gluten and the subsequent development of CeD.

The gut microbiome has been proposed to be one of these factors. Multiple studies on the CeD microbiome have shown alterations in gut microbiome composition in CeD patients, such as reduced levels of beneficial bacteria including *Bifidobacterium* and *Lactobacillus* species, and increased levels of *Proteobacterium* species, compared to healthy controls (15–19). Several mechanisms have been proposed that implicate changes in microbiome composition to CeD pathophysiology including, but not limited to, a) specific microbial proteases could be involved in processing immunogenic gluten peptides (20–22), b) toxins or other microbe-derived metabolites secreted by pathobionts may affect intestinal barrier permeability thereby influencing the passage of gluten peptides into the lamina propria c) microbial proteins may contain sequences similar to immunogenic gluten peptides (mimicry), potentially initiating an inflammation immune response (23). Viral infections have also been implicated in a decrease in tolerance to gluten (24–26). Together, these findings underscore

the intricate interplay among the gut microbiota, gluten, and the immune system in CeD
development.

Although these findings provide interesting insights, further analysis is necessary as many of
the CeD microbiome studies carried out thus far have been constrained by small patient
cohorts and/or relied upon 16S sequencing which limits the statistical power and taxonomic
resolution to the genus level, respectively. Furthermore, most studies do not provide
information on the functionality of the gut microbiota.

Our results, in addition to confirming previous studies (27,28), revealed novel associations
between tCeD and the microbiome, explored the contribution of a GFD to gut microbiome
composition and reported a less uniform microbiome composition, as well as increased strain-
to-strain variation in patients with tCeD. Upon taxonomic analyses, we found both tCeD and
self-reported IBD to present a unique microbiome signature, with limited overlap among
changes in microbial abundance. Lastly, we reported shifts in subspecies distribution of
Eubacterium rectale and *Firmicutes bacterium CAG:83* among tCeD patients, as compared to
controls.

Materials and Methods

Study population

Treated CeD

The Celiac Disease Northern Netherlands cohort (CeDNN; unpublished as of yet) comprises
129 CeD patients and 106 controls that provided faecal samples at the project's onset and had
complete metadata including age and sex. Most CeD patients in CeDNN had already been
clinically diagnosed and were adhering to a GFD at the time of inclusion. Since nearly all
controls were relatives or partners of the included patients, sharing genetic and/or lifestyle
factors, we chose to enhance our study by merging this dataset with a subset of participants
from the Lifelines Dutch Microbiome Project (LL-DMP) (29): 24 self-reported CeD-patients
from LL-DMP were included, as well as 659 controls. Controls were selected in such a way,
that the combined dataset contained 5 matched (age, sex, sequencing depth) controls for
every CeD patient (Table 1). Matching was performed with the R program *MatchIt* (v4.5.5)
(30). Self-reported IBD- and IBS-patients were excluded from the control group.

IBD and IBS

For comparative conditions, 93 self-reported IBD patients (Crohn's disease or ulcerative colitis)
from LL-DMP were matched with 5 controls each (age, sex, sequencing depth; 465 controls
total). 570 self-reported IBS patients from LL-DMP were matched with 3 controls each (age,
sex, sequencing depth; 1710 controls total). Self-reported CeD, IBD- and IBS-patients were
excluded from both control groups (Table 1).

Table 1: Description of the study population

Variable	Cohort	CeD		IBD		IBS	
		CeD	Controls	IBD	Controls	IBS	Controls
Sample size	CeDNN	129	106				
	LL-DMP	24	659	93	465	570	1710
Age (range)		30 (3-82)	30 (2-82)	53 (33-74)	54 (13-81)	49 (19-84)	49 (13-83)
Sex female (%)		98 (64%)	450 (59%)	56 (60%)	267 (57%)	411 (72%)	1233 (72%)
GFD (%)		93%	N/A				

Summary statistics of the data used in this study. The average age is indicated. Adherence to a gluten-free (GFD) diet was largely unknown for controls.

Ethical approval

The LL-DMP study and the CeDNN study were approved by the medical ethical committee of the University Medical Center Groningen (METc numbers 2017/152 and 2013/440, respectively). Additional written consent was signed by all participants, their parents or legal representatives (for participants aged under 18).

Phenotypic data

The CeDNN cohort uses the same questionnaires as Lifelines. Additionally, clinical information is available for many of the participants, including date of diagnosis and adherence to a GFD. The time between the date of diagnosis and the date of faecal sampling was used as a proxy for the duration of GFD.

Faecal sample collection, DNA extraction and sequencing

Participants from both CeDNN and LL-DMP provided stool samples from home. Participants were asked to freeze stool samples within 15 minutes of stool production. Frozen samples were collected by personnel/students of the University Medical Center of Groningen, (CeDNN) and Lifelines (LL-DMP), transported to their respective biorepositories on dry ice and stored at −80 °C until DNA extraction. Microbial DNA was isolated with the QIAamp Fast DNA Stool Mini Kit (Qiagen), according to the manufacturer's instructions, using the QIAcube (Qiagen) automated sample preparation system. Library preparation for samples with total DNA yield lower than 200 ng (as determined by Qubit 4 Fluorometer) was performed using NEBNext Ultra DNA Library Prep Kit for Illumina, while libraries for other samples were prepared using NEBNext Ultra II DNA Library Prep Kit for Illumina. Metagenomic sequencing was performed at Novogene, China using the Illumina HiSeq 2000 platform to generate approximately 8 Gb of 150 bp paired-end reads per sample.

Metagenomic data processing

Trimming and filtering of reads

Metagenomes were profiled as follows, consistent with previous analyses of 1000IBD (31), Lifelines-DEEP (32) and the Lifelines-DMP cohort (29). Illumina adapters and low-quality reads

187   (Phred score < 30) were filtered out using *KneadData* (trimmomatic options: "LEADING:20
188   TRAILING:20 SLIDINGWINDOW:4:20 MINLEN:50"; v0.10.0) (33). Following trimming, the
189   KneadData-integrated *Bowtie2* tool (v2.5.0) (34) was used to remove reads that aligned to the
190   human genome (GRCh37/hg19).
191
Taxonomy and pathway analysis
The taxonomic composition of metagenomes was profiled by *MetaPhlAn3* (v3.0.1) (35) using
the MetaPhlAn database of marker genes *mpa_v30*. None of the included samples had a
eukaryotic or viral abundance >25% of total microbiome content or a total read depth <10
million). MetaCyc pathways profiling was performed with *HUMAnN3* (default settings; v3.0.1)
(35) integrated with the *DIAMOND* alignment tool (v2.0.15) (36), UniRef90 protein database
(v0.1.1) (37) and *ChocoPhlAn* pan-genome database (*ChocoPhlAn_201901*) (35). Since single
genes can be mapped to multiple pathways, pathway abundances were instead normalized to
the sum of gene family abundances per sample.
201
To address the scarcity of microbial data during subsequent analysis, we focused on bacterial
taxa/pathways that were present at a minimum abundance of 0.001% in at least 5% (taxa) or
2.5% (pathways) of participants. This yielded 301 taxa (6 phyla, 14 classes, 20 orders, 33
families, 70 genera and 158 species) and 338 pathways for tCeD vs. controls; 310 taxa (7 phyla,
15 classes, 20 orders, 31 families, 71 genera and 166 species) and 349 pathways for IBD vs.
controls; and 316 taxa (9 phyla, 17 classes, 23 orders, 35 families, 73 genera and 159 species)
and 331 pathways for IBS vs. Controls.
209
210   *UpSet plot*
211   To visualize the extent of overlap among observed differences in the three comparisons (tCeD,
212   IBD and IBS vs. their respective matched controls), the data was plotted using the *UpSetR*
213   package in R (38).
214
215   *Alpha and beta diversity and inter-individual variation*
216   Based on the abundance profiles of the taxa that passed the filtering process, we calculated
217   alpha diversity, as measured by Shannon entropy, at the genus and species level using the
218   *diversity* function in R package *vegan* (v.2.6.4) (39). Beta diversity was assessed through Bray-
219   Curtis dissimilarity indices, as determined with the *vegdist* function from the R package *vegan*.
220
Strain- and gene-level analyses
For the comparison between individuals with tCeD and controls, we focused on studying
differences at a sub-species level. Initially, we conducted strain-level profiling using
StrainPhlAn3 (v3.0) (35), starting with default parameters for every species. For species for
which default parameters were found to be overly stringent, and that were significantly
associated with CeD according to the MetaPhlAn output, we adjusted the StrainPhlAn3 as
follows: a)   "--marker_in_n_samples   50",   and   b)   "--marker_in_n_samples

20 --sample_with_n_markers 10". Among these adjustments, three species
(*Eisenbergiella tayi*, *Haemophilus parainfluenzae* and *Anaerotruncus colihominis*) were
successfully analyzed only with the most lenient parameter set. Only *Enterococcus faecium*
failed with all attempted settings. Distance matrices were constructed from the multiple
alignments, using the command line tool *distmat*. Subsequently, we used *PanPhlAn3* (35) to
analyze the gene repertoire of 9 species with at least a nominally significant increase in
interindividual strain distance among CeD patients. Heatmaps were created with the
ComplexHeatmap package in R (40).
236
237 Analyses were performed using locally installed tools and databases on the high-performance
238 computing infrastructure available at our institution and using the MOLGENIS data platform
239 (41).
240
241 **Statistical analyses**
242 *Associations between alpha diversity and sample group*
243 We conducted a comparison of Shannon diversity between disease groups (tCeD, IBD, IBS)
244 and controls using a linear model. Specifically, when comparing tCeD to controls, we included
245 sequencing batch (CeDNN_batch1, CeDNN_batch2, and LL-DMP) as a covariate.
246
Associations of sample group with bacterial taxa and MetaCyc pathways
We tested for differentially abundant bacterial taxa and MetaCyc pathways by modelling their
abundance using linear models with FDR correction. Differences in prevalence between
sample groups were tested using logistic regression, followed with a c^2-test using the *step*
function from the *stats* package in R (with FDR correction). For the comparisons of tCeD versus
controls, sequencing batch (CeDNN_batch1, CeDNN_batch2 and LL-DMP) was used as a
covariate. The correlation between taxonomic and pathway abundance was quantified using
Kendall's t coefficient.
255
256 *Associations between duration of GFD and bacterial abundance*
257 The duration of GFD was set to 0 for patients, not on a GFD. Using a linear model, effects of
258 age, sex, sequencing depth, and sequencing batch on bacterial abundance were regressed
259 out. Subsequently, Spearman's r tests were used to assess the effect of duration of GFD on
260 the residuals of bacterial abundance.
261
Beta diversity
To evaluate the relationship between microbiome composition and disease groups, we
conducted a PERMANOVA test using the adonis2 function from the vegan package. We utilized
Bray-Curtis distance metrics and performed 999 permutations for each test. When comparing
tCeD to controls, sequencing batch (CeDNN_batch1, CeDNN_batch2, and LL-DMP) was
incorporated as a covariate.
268

To measure the heterogeneity within each sample group (tCeD, IBD, IBS, and their respective

control groups), we computed the distribution of all pairwise Bray-Curtis distances, excluding

sample pairs that were related to each other. Significance was determined using permutation

tests with 1000 permutations.

Strain-level analyses

To investigate whether increased inter-individual differences occur at a strain level, we

measured strain-to-strain variation using the distance matrices generated by distmat. These

matrices served as proxies for the mutational distance between strains of the same species in

different samples within either the control group or tCeD group. Sample pairs that were

related were excluded from this analysis. Significance was assessed using permutation tests

with 1000 permutations.

We then conducted PAM-clustering with the pam function from the cluster package in R. The

prediction.strength function from the *fpc* package (*clustermethod=claraCBI*) was employed to

determine the optimal number of clusters that best described the data. Correlation between

tCeD and strain-clusters was examined using Fisher's exact tests, with FDR-correction for

multiple comparisons.

Gene prevalence and gene enrichment analysis

Associations between the prevalence of UniRef90 gene families, as determined by PanPhlAn,

and tCeD were determined using logistic regression, followed with a c^2-test using the *step*

function from the *stats* package in R (with FDR correction). Age, sex, sequencing depth and

sequencing batch were used as covariates here. Genes that were present in less than 20% or

more than 80% of all samples were excluded from this analysis. Protein family (Pfam) and

Gene Ontology annotations were added to UniRef90 gene families and gene enrichment

analysis was performed using Fisher's exact tests with FDR correction. Nominally significant

gene families were used as input for the enrichment tests.

Correlation between RadC-like JAB domain proteins and inter-individual strain distance

To investigate whether the presence of proteins with the RadC-like JAB domain (PF04002), as

identified by PanPhlAn, influences the mutational load of associated bacterial species

(specifically, Bacteroides species *B. ovatus*, *B. vulgatus*, *B. uniformis*, *B. xylanisolvens)*, we

conducted the following analysis:

For each sample analyzed by both StrainPhlAn and PanPhlAn, we recorded the median

mutational distance (as determined by StrainPhlAn) to all other samples within the same

sample group (either tCeD or control), as well as the number of proteins with a RadC-like JAB

domain identified by PanPhlAn. We assessed trends within sample groups using a linear

model. Additionally, we examined any negative trend irrespective of the sample group using

a linear mixed-effects model (implemented with the *lmer* function from the *lmerTest* package
in R), with the sample group as a random effect.

Results

Alpha diversity is decreased in self-reported IBS and IBD subjects, but not in CeD patients
on GFD
Although contradicting data can be found in literature, most often it is described that within-
sample microbiome diversity (alpha diversity) in untreated CeD is higher than that in controls
(27,42) and that adherence to a GFD restores alpha diversity to normal levels (27,28). Indeed,
using the Shannon index for evenness of the microbiome, we did not observe a significant
difference in alpha diversity on either genus or species level, when comparing tCeD (adhering
to a GFD) patients and matched controls (Table 2). Self-reported IBS and IBD patients display
a small but significant (p < 0.05) decrease in alpha diversity at the genus level when compared
with matched controls. This is in line with studies that observe a marked decrease in diversity,
especially in more inflamed IBD-like samples (43–45). Although tCeD patients show a
comparable alpha diversity to controls, several differences are found when studying
microbiome composition in more detail.

tCeD, IBD and IBS show distinct microbial patterns
Next, we analyzed the levels of faecal bacteria in individuals with tCeD, IBD, and IBS, along
with their matched control groups. Our findings revealed strikingly different trends among
these sample sets (Table 2, Figure 1). While all groups were characterized by an increase of
several *Bacteroides* species, even within that genus several differences can be observed: *B.*
intestinalis, *B. stercoris* and *B. vulgatus* are exclusively altered in IBD, while *B. xylanisolvens* is
only enriched in tCeD. Only *B. ovatus* is enriched in more than one subject group (tCeD and
IBD). All other species with changed abundance in tCeD are unaltered in IBD and IBS. Only
three taxa show significant differential abundance in self-reported IBS-patients, compared to
controls (*Bacteroides* and *Veillonella* genera and *P. merdae*), which are in line with earlier
studies as reviewed by Pittayanon et al. (46). More overlap between the diseases can be seen
when looking at prevalence (Table 3), but several bacteria were identified that are affected
exclusively in tCeD.

Figure 1: Overlap in microbiome changes between tCeD, IBD and IBS. Bacterial species with A. higher abundance or prevalence in disease, and B. lower abundance or prevalence in disease.

Taxonomic analyses reveal known and novel associations with tCeD

Multiple of the observed differences in bacterial abundance and/or prevalence between tCeD and controls (Table 2 and Table 3) are in accordance with earlier reports, such as the enrichment of *H. parainfluenzae* (28)*, R. inulinivorans* (28), *E. coli* (18,47), and *Bacteroides* species (18,27,28), as well as the decreased prevalence of *R. bicirculans* , *C. eutactus* (27), *C. fastidiosus* (28,48), *Roseburia sp. CAG:309* (28)*,* and several *Bifidobacterium* species (18,28,47). In addition, however, several novel species were identified in our analysis, including an increase in *Clostridium sp. CAG:253*, *R. hominis*, and *C. lavalense*.

Table 2: Taxa with significantly altered abundance in tCeD, IBD and IBS

Name	Previously reported	tCeD	IBD	IBS
Shannon index - Genus (control/case)		1.86 / 1.84	1.87 / 1.70 *	1.87 / 1.83 *
Shannon index - Species (control/case)		2.44 / 2.48	2.44 / 2.35	2.45 / 2.44
Bacterial taxa				
Acidaminococcus intestini			3.2	
Anaerotruncus colihominis		2.5[b]		
Bacteroides (genus)			1.0	0.3
Bacteroides fragilis	(27)	1.3		
Bacteroides intestinalis			2.4	
Bacteroides ovatus	(49)	1.5	1.2	
Bacteroides stercoris			1.7	
Bacteroides thetaiotaomicron	(50)	0.9		
Bacteroides uniformis			0.7	
Bacteroides vulgatus			1.4	
Bacteroides xylanisolvens	(51)	1.5		
Bifidobacterium dentium			2.3	
Bifidobacterium pseudocatenulatum			2.5	
Clostridium bolteae			5.2	
Clostridium clostridioforme			5.3	
Clostridium sp. CAG:253		2.3		
Eisenbergiella tayi	(28)	3.2		
Enterococcus faecium		5.1[a,b]		

Name	Previously reported	tCeD	IBD	IBS
Firmicutes bacterium CAG:145			2.9	
Firmicutes bacterium CAG:94			1.2	
Flavonifractor plautii			1.1	
Fusobacterium (genus)			3.4	
Haemophilus parainfluenzae	(28)	1.2		
Parabacteroides distasonis			0.8	
Parabacteroides merdae				0.4
Paraprevotella xylaniphila	(28)	1.0		
Roseburia hominis		1.7		
Roseburia inulinivorans	(28)	1.4		
Roseburia sp. CAG:182		2.7[a,b]		
Ruthenibacterium lactatiformans			1.1	
Veillonella (genus)				3.2

Yellow and blue cells indicate an increase and decrease, respectively, in the Shannon index or bacterial abundance in the respective disease. An asterisk (*) indicates a significant difference in the Shannon index. Numbers in the bacterial taxa section represent $^2\log$(fold change) that are significant after multiple testing correction. Coloured cells without text indicate nominally significant differences. Previously reports of associations with CeD are included in a separate column. [a]Significant batch effect within the CeDNN cohort. [b]Significant difference between observed effect in the CeDNN and Lifelines cohorts.

Table 3: Taxa with significantly altered prevalence in tCeD, IBD and IBS

Name	Previously reported	tCeD	IBD	IBS
Acidaminococcus intestini			4 > 12%	
Akkermansia muciniphila			70 > 49%	
Alistipes finegoldii	(51)	99 > 95%		
Alistipes indistinctus			69 > 51%	
Alistipes shahii			89 > 78%	
Anaerotruncus colihominis		20 > 46%		13 > 18%
Bacteroides dorei	(50)	72 > 84%		
Bacteroides galacturonicus		8 > 5%		
Bacteroides sp. CAG:144		28 > 16%	43 > 26%	
Barnesiella intestinihominis			87 > 73%	
Bifidobacterium adolescentis	(16)	85 > 64%		
Bifidobacterium bifidum		37 > 23%		
Bifidobacterium catenulatum		29 > 6%[a]		
Bifidobacterium dentium				5 > 9%
Bifidobacterium longum	(52)	91 > 76%		
Butyrivibrio crossotus				16 > 10%
Clostridium bolteae			3 > 14%	
Clostridium clostridioforme			2 > 10%	
Clostridium lavalense		6 > 16%		
Clostridium leptum	(53)	23 > 54%[a]		
Clostridium sp. CAG:253		7 > 19%[a]		
Clostridium sp. CAG:58				28 > 36%
Coprobacter fastidiosus	(48)	46 > 29%		
Coprococcus eutactus	(54)	49 > 25%	47 > 25%	
Desulfovibrio piger			43 > 22%	
Desulfovibrionaceae (family)		88 > 76%	89 > 76%	
Dialister invisus			31 > 45%	
Eggerthella lenta		25 > 51%	17 > 37%	
Eisenbergiella massiliensis		3 > 16%		
Eisenbergiella tayi	(28)	4 > 32%[a]		
Enterococcus faecium		1 > 14%[a]		

Species		Cohort 1	Cohort 2	Cohort 3
Escherichia coli	(17)	44 > 72%		
Eubacterium rectale			92 > 83%	
Eubacterium siraeum	(48)	70 > 37%	71 > 54%	
Eubacterium sp. CAG:180				30 > 23%
Eubacterium sp. CAG:251			25 > 13%	
Faecalibacterium prausnitzii			100 > 96%	
Firmicutes bacterium CAG:110			71 > 41%	65 > 55%
Firmicutes bacterium CAG:170		34 > 16%	49 > 19%	
Firmicutes bacterium CAG:238		11 > 5%	22 > 6%	
Firmicutes bacterium CAG:83			87 > 66%	
Firmicutes bacterium CAG:95		27 > 9%	36 > 15%	
Flavonifractor plautii			67 > 85%	
Fusicatenibacter saccharivorans			98 > 85%	
Gemmiger formicilis			93 > 67%	93 > 88%
Gordonibacter pamelaeae				31 > 40%
Haemophilus parainfluenzae	(28)	21 > 45%		
Harryflintia acetispora		6 > 13%		8 > 13%
Hungatella hathewayi		5 > 11%		
Odoribacter splanchnicus		96 > 89%	96 > 87%	
Oscillibacter sp. 57_20		92 > 85%		
Oscillibacter sp. CAG:241			84 > 61%	
Parabacteroides distasonis				92 > 85%
Parabacteroides merdae			77 > 62%	
Paraprevotella clara			28 > 14%	
Paraprevotella xylaniphila			52 > 33%	48 > 38%
Phascolarctobacterium succinatutens			30 > 12%	
Prevotella sp. CAG:279				17 > 10%
Roseburia sp. CAG:182		24 > 43%		
Roseburia sp. CAG:309	(28)	14 > 2%		
Ruminococcaceae (family)	(55)	80 > 91%	71 > 86%	76 > 82%
Ruminococcus bicirculans	(28)	64 > 20%[a]		
Ruminococcus gnavus			6 > 17%	8 > 12%
Ruminococcus torques			51 > 37%	
Selenomonadaceae (family)		8 > 3%	12 > 3%	12 > 6%
Selenomonadales (order)			17 > 5%	
Veillonella atypica		4 > 10%		
Veillonella parvula				5 > 9%
Veillonellaceae (family)	(56)	13 > 24%	8 > 18%	9 > 14%
Victivallis vadensis		40 > 16%	49 > 23%	

Yellow and blue cells indicate an increase and decrease, respectively, in prevalence in the respective disease. Numbers represent shifts in prevalence (control > case) that are significant after multiple testing correction. Coloured cells without text indicate nominally significant differences. [a]Significant interaction with batch and/or cohort observed.

Duration of GFD explains part of the observed differences

Although we are primarily reporting on the microbiome composition of tCeD patients, we attempted to assess the contribution of GFD to the increased abundance of species shown in Table 2. After correction for covariates, we found that the duration of adherence to a GFD positively correlated with the increased abundance of *Paraprevotella xylaniphila* and *Roseburia sp. CAG:182* (Spearman rank correlation, $p_{adj} < 0.05$; Table 4). Moreover, given that the majority of the 13 tested species exhibit a positive correlation, these findings imply that GFD indeed plays a role in several species-level associations observed in this study. Notably,

378 we were unable to replicate the adverse impact of the GFD on the abundance of *H.*
379 *parainfluenzae* as seen in other research (28,57).
380
381 Table 4: Spearman correlation between duration of gluten-free diet and species abundance

Name	Spearman's ρ
Bacteroides fragilis	0.15
Bacteroides ovatus	0.20
Bacteroides thetaiotaomicron	0.13
Bacteroides xylanisolvens	0.05
Paraprevotella xylaniphila	0.28*
Enterococcus faecium	0.04
Clostridium sp. CAG:253	0.12
Eisenbergiella tayi	0.23
Roseburia hominis	-0.11
Roseburia inulinivorans	0.20
Roseburia sp. CAG:182	0.36*
Anaerotruncus colihominis	-0.03
Haemophilus parainfluenzae	0.16

382
383 Spearman correlation coefficients between GFD duration and species abundance, for all species significantly enriched in tCeD
384 patients. *p_{adj} < 0.05.
385

Differences in metabolic pathways are largely driven by specific bacterial taxa
After establishing the taxonomic landscape in each of the subgroups analyzed in this study,
we analyzed the functional potential of the associated microbiota as can be inferred by
mapping of identified gene families to metabolic pathways in the MetaCyc database. Although
several differences were found between tCeD patients and controls, very few of these
differences were unique to tCeD, as there was a striking concordance with differences
observed in IBD and IBS (Table 5). These similarities can probably be explained by the fact that
most differentially abundant metabolic pathways are strongly associated to differentially
abundant taxa, with *Bacteroides* species and *E. coli* explaining most of the observed changes
(Table 5 and Supp. Figure 1). Furthermore, even pathways with lower abundance in tCeD than
controls (Table 5, blue) appear to be primarily driven by an increase in bacteria from the
Bacteroidetes phylum that replaces bacteria carrying this metabolic potential, reflecting the
compositional nature of these data. Since pathway abundance seems to mainly reflect
taxonomic changes, rather than functional enrichment across several taxa, the identified
pathways are, in our assessment, unlikely to be relevant in the context of the disease studies.
401
402 Table 5: Metabolic pathways differentially abundant in tCeD

Pathway	CeD	IBD	IBS	Top correlated taxa ($\tau > 0.3$)	τ
PENTOSE-P-PWY: pentose phosphate pathway	0.3	0.3	0.1	*B. uniformis*	0.47
POLYISOPRENSYN-PWY: polyisoprenoid biosynthesis	0.6	0.7	0.2	*B. uniformis* *B. vulgatus*	0.59
PWY-2941: L-lysine biosynthesis II	0.4[a]	0.3	0.1	*Bacteroidaceae* (family)	0.32

Pathway				Correlated taxa	Kendall's τ
PWY-5083: NAD/NADH (de)phosphorylation	1.4	1.1		*B. ovatus*	0.41
PWY-5838: superpathway of menaquinol-8 biosynthesis I	1.3[a,b]	1.1		*E. coli*	0.54
PWY-5941: glycogen degradation II	0.9[a]	0.8	0.3	*B. ovatus* *B. thetaiotaomicron* *B. uniformis*	0.46
PWY-5971: palmitate biosynthesis II	1.3[b]	1.1		*E. coli*	0.58
PWY-6353: purine nucleotide degradation II (aerobic)	0.7[a,b]	0.5			
PWY-6531: mannitol cycle	1.1	0.8		*B. ovatus*	0.40
PWY-6572: chondroitin sulfate degradation I	0.8[a]	0.7	0.3	*B. thetaiotaomicron*	0.58
PWY-6608: guanosine nucleotide degradation III	0.6	0.4			
PWY-6859: all-trans-farnesol biosynthesis	0.6[a]	0.7	0.2	*B. vulgatus, B. uniformis*	0.55
PWY-6953: dTDP-3-acetamido-3,6-dideoxy-α-D-galactose biosynthesis	3.0			*B. cellulosilyticus*	0.35
PWY-7316: dTDP-N-acetylviosamine biosynthesis	1.7			*B. fragilis*	0.42
RHAMCAT-PWY: L-rhamnose degradation I	0.4	0.3		*B. uniformis*	0.36
ARGSYNBSUB-PWY: L-arginine biosynthesis II (acetyl cycle)	-0.1[a,b]	-0.2	-0.1	*Bacteroidetes* (phylum) / *F. prausnitzii*	-0.44 / 0.32
PWY-7219: adenosine ribonucleotides de novo biosynthesis	-0.1[a]	-0.1	-0.03	*Bacteroidaceae* (family)	-0.39
PWY-7383: anaerobic energy metabolism	-0.3	-0.4		*Bacteroidetes* (phylum) / *A. muciniphila*	-0.40 / 0.31
PWY66-399: gluconeogenesis III	-0.2	-0.3		*Bacteroidetes* (phylum)	-0.43
TRNA-CHARGING-PWY: tRNA charging	-0.1	-0.1		*Bacteroidetes* (phylum)	-0.36

Yellow and blue cells indicate an increase and decrease, respectively, in abundance in the respective disease. Numbers represent ^{2}log(fold change) that are significant after multiple testing correction. Coloured cells without text indicate nominally significant differences. Correlated taxa are reported with associated Kendall's τ coefficient. [a]Significant batch effect within the CeDNN cohort. [b]Significant difference between observed effect in the CeDNN and Lifelines cohorts.

tCeD faecal microbiome is less uniform than that of controls

Since we identified several bacterial species specifically enriched in tCeD, we suspected that tCeD patients' microbiomes would display a certain (t)CeD signature and be more uniform than those of non-CeD controls. To test this hypothesis, we first calculated Bray-Curtis dissimilarity between samples, a measure of beta diversity, and performed principal coordinate analyses (Figure 2A). Not surprisingly, the average overall composition of the tCeD microbiome differed significantly from that of controls (PERMANOVA; R^2=0.003; $p < 0.05$). However, when we tested which group was the most homogeneous, i.e. which group displayed the lowest median within-group dissimilarity (Figure 2B), we found that the microbiome composition of two arbitrary, unrelated tCeD patients tended to be more dissimilar than that of two unrelated controls (permutation test, $p < 0.05$). Interestingly, the same phenomenon, where patients have a less uniform microbiome structure than controls,

421 was observed among self-reported IBD patients (Supp. Figure 2A-B; $p < 0.05$). In contrast, the
422 distribution of Bray-Curtis dissimilarity scores among controls and self-reported IBS patients
423 were indistinguishable (Supp. Figure 2C-D).
424

426 Figure 2: Beta diversity in tCeD vs. controls, using Bray-Curtis dissimilarity as a distance metric. A. Principal
427 coordinate plot of tCeD samples and matched controls. B. Within-group dissimilarity histograms. Top: pairs of
428 unrelated controls. Bottom: pairs of unrelated tCeD patients. Indicated are the number of different permutations
429 of two samples that fall into each group.

tCeD dissimilarity is also present on sub-species level

Next, we wanted to assess whether the increased diversity among tCeD patients was merely
a global phenomenon, acting on the level of the full microbiome, or whether this effect was
present on multiple levels. Using StrainPhlAn to estimate the pairwise evolutionary distance
between bacterial strains of the same species from two different participants, we were able
to analyze strain-level diversity for a total of 67 bacterial species [Supp. Table S9]. For 9 species
(including 4 *Bacteroides* species; Figure 4), the inter-individual evolutionary strain distance
among tCeD patients tended (nominal $p < 0.05$) to be higher than among controls, while only
1 showed the tendency toward more diversity among controls. After multiple testing
correction, *Eubacterium rectale* (Figure 3B) and *Bacteroides* species *B. ovatus, B. uniformis,*
and *B. xylanisolvens* remained significant and were all more diverse among tCeD patients.
441

442
Figure 3: Strain-level differences of *Eubacterium rectale* between tCeD patients and controls. A. Abundance and
prevalence of *E. rectale* in patients and controls. Violin plots represent the distribution of abundance (log-scale)
in the two groups. Samples with absent *E. rectale* are excluded from this plot. Shaded bars on the background
represent prevalence among all samples. B. Distribution of pairwise genetic distances between *E. rectale* strains.
Control vs. control (top) and tCeD vs. tCeD (bottom). Pairs of relatives were excluded from both comparisons. C.
Non-metric multidimensional scaling plots of pairwise genetic distances between *E. rectale* strains, coloured by
cluster-ID (PAM clustering; left) and disease status (right). Ellipses include 95% of the samples in the indicated
group. D. Heatmap of presence/absence of *E. rectale* genes in all analyzed *E. rectale* strains. Each row represents
the *E. rectale* strain from one participant. A phylogenetic tree based on pairwise genetic distances is shown on
the left, as well as the participant's disease status. Each column represents a UniRef90 cluster (gene) that is
differentially prevalent in tCeD patients (nominal *p* < 0.05). Coloured bars in the bottom indicate whether the
genes are predicted to encode glycosyltransferases or flagellum-related proteins.

European subspecies of *Eubacterium rectale* is reduced among tCeD patients

To further investigate strain-to-strain variation, we performed principal coordinate analyses, combined with PAM (partition around medoids) clustering. This approach revealed distinct sub-species clustering for 46 out of the 67 species studied. For *E. rectale* (Figure 3C) and *Firmicutes bacterium CAG:83* (Supp. Figure 3; likely part of the *Oscillibacter* genus (58)), the dominant clusters among controls (61% and 69%, respectively) were significantly reduced among tCeD patients to 33% and 44% (Fisher's exact test, p_{adj} < 0.05).

Finally, we performed gene-level comparison of cases and controls (PanPhlAn), followed by enrichment analysis of protein family (Pfam) and gene ontology (GO) terms among nominally significant (p < 0.05) genes. With this analysis, we aimed to identify gene-level explanations of the strain-level observations (increased inter-individual differences and depletion of the dominant *E. rectale* cluster). The only class enriched among *E. rectale* genes with lower prevalence in tCeD was PF00535 (glycosyl transferase family 2), while over-prevalent genes were enriched for terms related to flagellum assembly and organization, chemotaxis, protein targeting and secretion, and phosphoenolpyruvate-dependent sugar phosphotransferase systems (Figure 3D). Karcher et al. described an *E. rectale* subspecies almost exclusively found in Europe (ErEurope) that is characterized specifically by the presence of several genes from glycosyl transferase family 2 and the absence of motility operons, including chemotaxis and flagellum genes (59). We conclude, therefore, that specifically ErEurope is less prevalent in tCeD patients.

Figure 4: Distribution of pairwise genetic distances between strains of A. *Bacteroides ovatus, Bacteroides uniformis, Bacteroides vulgatus, Bacteroides xylanisolvens*. Control vs. control (top) and tCeD vs. tCeD (bottom). Pairs of relatives were excluded from both comparisons. The significance of increased distance among tCeD patients is indicated by p-values from permutation test (1000 iterations). Asterisks (*) indicate p_{adj} < 0.05.

Gene families related to transmembrane transport and DNA repair are less prevalent in
multiple species in tCeD patients
Besides *E. rectale*, we performed gene-level analyses for 6 other bacterial species. When
analyzed separately, no enriched terms could be identified for any of these species. To assess
whether there were any bacterial gene or protein classes enriched across species, we also
performed the analysis after pooling gene-level data of all 7 bacteria. Among under-prevalent
genes in tCeD, we identified two new enriched classes. Firstly, 23 of 150 genes from
GO:0055085 (transmembrane transport) were under-prevalent in tCeD. Many of the
contributing genes, from 6 different species, were annotated to encode permeases, with some
of them having a predicted specificity for carbohydrates. We suggest that differences in
nutrient availability during active CeD or a gluten-free diet may play a role in the loss of these
genes encoding carbohydrate uptake proteins.
494
495 Secondly, 9 of 21 genes encoding proteins from PF04002 (RadC-like JAB domain) were under-
496 prevalent, across all tested *Bacteroides* species. Proteins from this family have been
497 speculated to have proteolytic and/or nuclease activity and to be involved in DNA repair (60).
498
Presence of DNA repair genes in *Bacteroides* species results in lower mutational drift
Next, sought to assess whether the reduced prevalence in tCeD of DNA repair genes
containing the RadC-like JAB domain could be associated with the increased inter-individual
variation of strains of *Bacteroides* species. Indeed, we found that the number of genes in a
specific strain encoding a RadC-like JAB domain negatively correlated (Figure 5) with the
median distance of that strain to strains of the same species within the same sample group
(tCeD and control). This negative correlation was observed in all species tested (*B. ovatus*, *B.*
uniformis, *B. vulgatus* and *B. xylanisolvens*) and was significant in the two species with most
data points, *B. uniformis* and *B. vulgatus* (linear mixed-effects model, $p < 0.05$). Moreover, for
all four species, there was a negative correlation both in the tCeD group and in controls, in
line with our hypothesis that these DNA repair proteins are limiting the mutation rate in these
Bacteroides species.
511

Figure 5: Correlation between prevalence, in a specific sample, of genes encoding the RadC-like JAB domain ('# Genes') and the median mutational distance (StrainPhlAn) from other strains within the same sample group, for each of A. *Bacteroides ovatus*, B. *Bacteroides uniformis*, C. *Bacteroides vulgatus*, and D. *Bacteroides xylanisolvens*. Significance within sample groups is indicated with p-values inside the plot, while p-values in the plot title across both groups report on the association in both groups combined (linear mixed-effects model).

Individual gene families from *Eubacterium rectale*, *Alistipes putredinis* and *Bifidobacterium adolescentis* are differentially prevalent in tCeD.

Besides gene families that were part of above-mentioned functional classes, 413 more gene families were differentially prevalent in tCeD patients, compared to controls ($p_{adj} < 0.05$; Supp. Table S10). These gene families are derived from *Eubacterium rectale* (38 up, 101 down), *Alistipes putredinis* (168 up, 96 down) and *Bifidobacterium adolescentis* (4 up, 6 down).

Interestingly, *B. adolescentis* strains found in tCeD less frequently encode a LacI/GalR-family transcriptional regulator and galactose-1-phosphate uridyltransferase GalT, part of the Leloir pathway for galactose metabolism. This finding constitutes another potential implication of altered carbohydrate availability during inflammation and/or GFD (61).

Discussion

In this paper, we performed a comprehensive analysis of the largest cohort of CeD microbiota to date, comparing the faecal microbiota of 153 tCeD patients with 765 controls, matched by age, sex, and sequencing depth. Our approach utilises shotgun metagenomic sequencing

535 allowing for microbiome classification at the strain level and providing insights into bacterial
536 genes and pathways.
537
Taxonomic and functional profiling
We aimed to compare taxonomic profiles at the species level with those of previous CeD
microbiome studies. In summary, our data (Table 2 and Table 3) confirms previous findings in
(treated) CeD patients, demonstrating an increase in several *Bacteroides* species, *H.*
parainfluenzae, R. inulinivorans, and *E. coli*, along with a decrease of various bifidobacteria,
R. bicirculans, C. eutactus, and *C. fastidiosus* (18,27,28,47). Despite a lower prevalence of the
major butyrigenic bacterium *Butyrivibrio crossotus* (62) (Table 3), we observed novel
associations with increased abundances of other butyrate-producing bacteria including
Clostridium sp. CAG:253, R. hominis, and *C. lavalense* in tCeD patients in our tCeD cohort.
Interestingly, these bacteria, as well as some of the previously reported bacteria (e.g. *R.*
inulinivorans) all produce butyrate (62,63). Higher production of butyrate by the CeD
microbiome would be in line with observations by Baldi et al. that CeD patients display slightly
higher butyrate fractions (64). However, Tjellström et al. showed that one year of adherence
to a GFD restored butyrate levels in pediatric CeD patients to normal levels (65). We speculate
that the negative effect of a GFD on butyrate production does not necessarily reflect the
restoration of the microbiome to a pre-CeD state, but may rather be mediated by other
butyrigenic species, such that effects of disease and diet largely cancel each other out.

555
Further taxonomic analyses revealed that tCeD, IBD and IBS were all characterised by an
increase in the *Bacteroides* genus, which drives remarkably similar changes in metabolic
pathways observed in each disease group. Moreover, it seems that even all (moderately)
under-abundant pathways are explained by the (moderate) increase of bacteria from the
Bacteroidetes phylum, therefore likely an artefact caused by the compositional nature of the
data (Table 5, Supp. Figure 1).

562
563 Besides the genus *Bacteroides*, however, the different disease groups studied each show
564 highly distinct microbiome profiles (Table 2, Table 3, Figure 1). Although we showed that the
565 GFD contributes to an altered faecal microbiome (Table 4), the current data does not allow us
566 to separate the consequences of CeD and/or GFD from characteristics of the microbiome that
567 contribute to the onset of CeD.

568
Although most metabolic pathways are seemingly altered because the bacteria encoding
them happen to be altered in tCeD, we cannot dismiss the possibility that certain bacteria are
enriched *because* they encode those specific metabolic pathways. Indeed, other studies
report several *Bacteroides* species to be enriched in active (18,42,66) and treated CeD patients
(18,27,42), as well as in at-risk individuals (51,67,68), and in these studies, *B. fragilis* was
speculated to be involved in pathogenesis (49). Although none of the previous studies report

to have performed multiple testing correction, compromising their true discovery rate, taken
together they do suggest that the overabundance of *Bacteroides* species is not solely caused
by GFD and that some of the pathways we have identified might therefore be relevant in CeD
pathogenesis. Additionally, the 1.3-fold enrichment of genes involved in the rhamnose
degradation pathway (Table 5) could instead be related to differences in carbohydrate supply
from the diet. Furthermore, two pathways were uniquely affected in tCeD, also representing
the two largest differences with controls: the dTDP-N-acetylviosamine biosynthesis and dTDP-
3-acetamido-3,6-dideoxy-a-D-galactose biosynthesis pathways are 3.3-fold and 7.8-fold
enriched in tCeD. Both these pathways use α-D-glucopyranose 1-phosphate, a product of
glycogen degradation, as a substrate. Since both these pathways and, as in IBD, that of
glycogen degradation are overabundant in tCeD and associated with various *Bacteroides*
species, it could be that the ability to survive in harsh conditions, with low carbohydrate
availability, allows these bacteria to outcompete other bacteria (69). Only two pathways
enriched in tCeD do not appear to be driven by specific bacterial taxa, PWY-6353 and PWY-
6608 (Table 5), both of which are related to purine nucleotide degradation.
590
Inter-individual variation
Besides species-specific differences between tCeD patients and controls, we found an
increased inter-individual variation among patients as compared to the diversity among
controls (Figure 2). While we expected to find that patients might share a certain microbiome
signature, either related to the disease or GFD, the inverse appears to be true: their
microbiome, more than that of controls, deviates from the average non-CeD microbiome (not
shown), but these changes apparently occur in different directions between patients.
Although we want to refrain from using the term *dysbiosis* (70), this observation could reflect
the loss of certain healthy components of the microbiome. On the other hand, it could be that
a gluten-*containing* diet, rather than disease or GFD, induces a microbiome signature. While
the possibility of a loss of specific 'healthy' bacteria characterizing CeD seems incompatible
with the fact that we exclusively observed over-abundant bacteria in tCeD, presence-absence
analyses did reveal some bacteria with reduced prevalence in tCeD, such as several species of
bifidobacteria, generally regarded as 'healthy' bacteria (71), along with *R. bicirculans*, *E.*
siraeum and *C. eutactus* (Table 3).

606
Interestingly, the increased diversity among patients is also visible in strain-level analyses,
where, for several bacteria, the average evolutionary distance between strains of the same
species in two unrelated tCeD patients is larger than that between strains in two control
individuals (Supp. Table S9). This phenomenon was observed most clearly ($p_{adj} < 0.05$) in
Eubacterium rectale (Figure 3B) and *Bacteroides ovatus, Bacteroides uniformis*, and
Bacteroides xylanisolvens (Figure 4), but appears to be more general, as 48 out of 67 analysed
species (72%) show the same trend. The most plausible explanation, in our opinion, is that
inflammation and oxidative stress accompanying CeD and gluten intake (72) lead to a higher

mutational burden in the gut microbiome. In line with this hypothesis, transcriptome analysis
of intestinal epithelial cells recently revealed the upregulation of pathways related to DNA
double-strand break repair in active CeD, as compared to controls (73). Moreover, we find that
Bacteroides strains (multiple species) that lack the RadC-like JAB domain, involved in DNA
repair (60), are less prevalent in tCeD than in controls (Supp. Table S10 and Supp. Table S11),
and that the presence of these repair proteins in *Bacteroides* strains limits their mutation rate
and thereby inter-strain variation (Figure 5). These observations together suggest it may be
beneficial to these bacteria to increase the genetic diversity of their offspring under stressful
conditions. It is, in fact, an established strategy among bacteria to upregulate their evolution
rate in response to stress (74). Examples of this strategy are the widespread SOS-response,
utilizing an error-prone DNA polymerase (75), and the activation of competence for
transformation during DNA replication stress (76,77). It is unclear whether the observed
increase in mutational load in the tCeD microbiota is an indication of residual inflammation or
merely a result of periods of inflammation in the past.
629
Intriguingly, while *Eubacterium rectale*, too, shows clear evidence of what seems to be random
mutagenesis (Figure 3B, Supp. Figure 4), we also observed that the ErEurope subspecies (59)
was almost absent in tCeD, while nearly half of the controls carried this subspecies.
Importantly, ErEurope lacks many motility genes and has a distinct carbohydrate metabolism
potential. Karcher et al. suggested that ErEurope utilizes complex plant-derived carbohydrates
more efficiently (59). As no interaction with diet was previously observed for ErEurope, most
of the different *E. rectale* strains do not co-colonize (59), and no differences in prevalence or
abundance were observed between tCeD and controls, it is possible that the strain-level
difference in *E. rectale* may exist prior to GFD. It remains unclear whether the lack of motility
limits the ability of ErEurope to access carbohydrates during inflammation, whether their
carbohydrate metabolism potential plays a role in gluten processing, or whether pre-existent
genetic or environmental risk factors for CeD also affect *E. rectale* substrain colonization
success.
643
Limitations
Since the patients with CeD in our study strictly followed a GFD, our findings do not allow us
to definitively ascertain the role of the microbiome in the onset or progression of the disease.
This challenge is common in much of the existing research, where data interpretation is
hindered by diet and the possibility that observed differences stem from studying patients
with active CeD, potentially influenced by associated intestinal inflammation. Furthermore,
our study is limited by the use of faecal samples, which may not accurately reflect conditions
in the small intestine.
652
Despite these limitations, the current findings are a starting point for future work, addressing
questions on which differences do, in fact, reflect changes in the small intestine, which of
these precede the onset of disease, and what mechanisms are responsible for these

differences in microbial composition. Additional future data, ideally obtained from
prospective and longitudinal studies - including those of active CeD patients, healthy
individuals adhering to a GFD, and/or at-risk individuals - will help pull apart the various
influences of genetics, diet, and disease on the gut microbiome of CeD patients. A longitudinal
study, where data is available before CeD onset, was published in 2021 and reported.
However, that study had limited taxonomic resolution (16S sequencing) and statistical power,
with only 10 cases and 10 controls (51). Furthermore, as the significance levels in this study
were, apparently, not corrected for multiple testing, it is difficult to judge whether the
reported differences are beyond what would be expected by chance.
665
Conclusion
We characterized the microbiota of tCeD patients, as compared to controls, leading to both
the confirmation of previously reported CeD-microbiome associations and the identification
of novel associations, that may be linked to either CeD, GFD, or both. Strain- and gene-level
comparisons revealed a higher mutation frequency in several species, which may be induced
by inflammation or oxidative stress. Data on the prevalence of the RadC-like JAB domain in
multiple *Bacteroides* species suggests that the increased mutational load may be beneficial
for bacteria in stressful conditions. Finally, we find that specifically the European subspecies
of *Eubacterium rectale* is virtually absent in tCeD patients, a finding that may, in the future,
help to shed light on some of the mechanisms that shape the microbiome in health and
disease.

References

1. Ludvigsson JF, Leffler DA, Bai JC, *et al.* The Oslo definitions for coeliac disease and related terms. Gut 2013;62:43–52.

2. Lebwohl B, Sanders DS, Green PHR. Coeliac disease. Lancet 2018;391:70–81.

3. Lebwohl B, Ludvigsson JF, Green PHR. Celiac disease and non-celiac gluten sensitivity. BMJ 2015;351:h4347.

4. Gujral N, Freeman HJ, Thomson ABR. Celiac disease: prevalence, diagnosis, pathogenesis and treatment. World J Gastroenterol 2012;18:6036–6059.

5. Rewers M. Epidemiology of celiac disease: what are the prevalence, incidence, and progression of celiac disease? Gastroenterology 2005;128:S47-51.

6. Kårhus LL, Thuesen BH, Skaaby T, *et al.* The distribution of HLA DQ2 and DQ8 haplotypes and their association with health indicators in a general Danish population. United European Gastroenterol J 2018;6:866–878.

7. Bodd M, Ráki M, Tollefsen S, *et al.* HLA-DQ2-restricted gluten-reactive T cells produce IL-21 but not IL-17 or IL-22. Mucosal Immunol 2010;3:594–601.

8. Di Sabatino A, Vanoli A, Giuffrida P, *et al.* The function of tissue transglutaminase in celiac disease. Autoimmun Rev 2012;11:746–753.

9. Troncone R, Jabri B. Coeliac disease and gluten sensitivity. J Intern Med 2011;269:582–590.

10. Al-Toma A, Volta U, Auricchio R, *et al.* European Society for the Study of Coeliac Disease (ESsCD) guideline for coeliac disease and other gluten-related disorders. United European Gastroenterol J 2019;7:583–613.

11. Heel DA van, Franke L, Hunt KA, *et al.* A genome-wide association study for celiac disease identifies risk variants in the region harboring IL2 and IL21. Nat Genet 2007;39:827–829.

12. Hunt KA, Zhernakova A, Turner G, *et al.* Newly identified genetic risk variants for celiac disease related to the immune response. Nat Genet 2008;40:395–402.

13. Dubois PCA, Trynka G, Franke L, *et al.* Multiple common variants for celiac disease influencing immune gene expression. Nat Genet 2010;42:295–302.

14. Kumar V, Wijmenga C, Withoff S. From genome-wide association studies to disease mechanisms: celiac disease as a model for autoimmune diseases. Semin Immunopathol 2012;34:567–580.

15. Collado MC, Calabuig M, Sanz Y. Differences between the fecal microbiota of coeliac infants and healthy controls. Curr Issues Intest Microbiol 2007;8:9–14.

16. Sanz Y, Sánchez E, Marzotto M, *et al.* Differences in faecal bacterial communities in
coeliac and healthy children as detected by PCR and denaturing gradient gel
electrophoresis. FEMS Immunology & Medical Microbiology 2007;51:562–568.

17. Nadal I, Donant E, Ribes-Koninckx C, *et al.* Imbalance in the composition of the duodenal
microbiota of children with coeliac disease. J Med Microbiol 2007;56:1669–1674.

18. Collado MC, Donat E, Ribes-Koninckx C, *et al.* Specific duodenal and faecal bacterial
groups associated with paediatric coeliac disease. Journal of Clinical Pathology
2009;62:264–269.

19. Collado MC, Donat E, Ribes-Koninckx C, *et al.* Imbalances in faecal and duodenal
Bifidobacterium species composition in active and non-active coeliac disease. BMC
Microbiol 2008;8:232.

20. Caminero A, McCarville JL, Galipeau HJ, *et al.* Duodenal bacterial proteolytic activity
determines sensitivity to dietary antigen through protease-activated receptor-2. Nat
Commun 2019;10:1198.

21. Constante M, Libertucci J, Galipeau HJ, *et al.* Biogeographic Variation and Functional
Pathways of the Gut Microbiota in Celiac Disease. Gastroenterology 2022;163:1351-
1363.e15.

22. Caminero A, Galipeau HJ, McCarville JL, *et al.* Duodenal bacteria from patients with
celiac disease and healthy subjects distinctly affect gluten
breakdown and immunogenicity. Gastroenterology 2016;151:670–683.

23. Petersen J, Ciacchi L, Tran MT, *et al.* T cell receptor cross-reactivity between gliadin and
bacterial peptides in celiac disease. Nat Struct Mol Biol 2020;27:49–61.

24. Welch NL, Welch T, Songtanin B. Viral Triggered Celiac Disease: A Case Report. Cureus
15:e40429.

25. Kagnoff MF, Austin RK, Hubert JJ, *et al.* Possible role for a human adenovirus in the
pathogenesis of celiac disease. J. Exp. Med. 1984;160:1544–1557.

26. Stene LC, Honeyman MC, Hoffenberg EJ, *et al.* Rotavirus infection frequency and risk of
celiac disease autoimmunity in early childhood: a longitudinal study. Am J Gastroenterol
2006;101:2333–2340.

27. Palmieri O, Castellana S, Bevilacqua A, *et al.* Adherence to Gluten-Free Diet Restores
Alpha Diversity in Celiac People but the Microbiome Composition Is Different to Healthy
People. Nutrients 2022;14:2452.

28. Francavilla A, Ferrero G, Pardini B, *et al.* Gluten-free diet affects fecal small non-coding
RNA profiles and microbiome composition in celiac disease supporting a host-gut
microbiota crosstalk. Gut Microbes 2023;15:2172955.

29.   Gacesa R, Kurilshikov A, Vich Vila A, *et al.* Environmental factors shaping the gut
746   microbiome in a Dutch population. Nature 2022;604:732–739.

747   30.   Ho D, Imai K, King G, *et al.* MatchIt: Nonparametric Preprocessing for Parametric Causal
748   Inference. Journal of Statistical Software 2011;42:1–28.

749   31.   Imhann F, Van der Velde KJ, Barbieri R, *et al.* The 1000IBD project: multi-omics data of
750   1000 inflammatory bowel disease patients; data release 1. BMC Gastroenterol
751   2019;19:5.

752   32.   Tigchelaar EF, Zhernakova A, Dekens JAM, *et al.* Cohort profile: LifeLines DEEP, a
753   prospective, general population cohort study in the northern Netherlands: study design
754   and baseline characteristics. BMJ Open 2015;5:e006772.

755   33.   McIver LJ, Abu-Ali G, Franzosa EA, *et al.* bioBakery: a meta'omic analysis environment.
756   Bioinformatics 2018;34:1235–1237.

757   34.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat. Methods
758   2012;9:357–359.

759   35.   Beghini F, McIver LJ, Blanco-Míguez A, *et al.* Integrating taxonomic, functional, and
760   strain-level profiling of diverse microbial communities with bioBakery 3. Elife
761   2021;10:e65088.

762   36.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. Nat
763   Methods 2015;12:59–60.

764   37.   The UniProt Consortium. UniProt: the universal protein knowledgebase. Nucleic Acids
765   Res. 2017;45:D158–D169.

766   38.   Lex A, Gehlenborg N, Strobelt H, *et al.* UpSet: Visualization of Intersecting Sets. IEEE
767   Trans Vis Comput Graph 2014;20:1983–1992.

768   39.   Dixon P. VEGAN, a package of R functions for community ecology. Journal of Vegetation
769   Science 2003;14:927–930.

770   40.   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
771   multidimensional genomic data. Bioinformatics 2016;32:2847–2849.

772   41.   Swertz MA, Dijkstra M, Adamusiak T, *et al.* The MOLGENIS toolkit: rapid prototyping of
773   biosoftware at the push of a button. BMC Bioinformatics 2010;11 Suppl 12:S12.

774   42.   Schippa S, Iebba V, Barbato M, *et al.* A distinctive "microbial signature" in celiac pediatric
775   patients. BMC Microbiol 2010;10:175.

776   43.   Pisani A, Rausch P, Bang C, *et al.* Dysbiosis in the Gut Microbiota in Patients with
777   Inflammatory Bowel Disease during Remission. Microbiology Spectrum 2022;10:e00616-
778   22.

44. Bourgonje AR, Roo-Brand G, Lisotto P, *et al.* Patients With Inflammatory Bowel Disease
Show IgG Immune Responses Towards Specific Intestinal Bacterial Genera [Internet].
Front. Immunol. 2022;13[cited 2024 Mar 6] Available from:
https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2022.842911
/full

45. Vila AV, Hu S, Andreu-Sánchez S, *et al.* Original research: Faecal metabolome and its
determinants in inflammatory bowel disease. Gut 2023;72:1472.

46. Pittayanon R, Lau JT, Yuan Y, *et al.* Gut Microbiota in Patients With Irritable Bowel
Syndrome—A Systematic Review. Gastroenterology 2019;157:97–108.

47. Di Cagno R, De Angelis M, De Pasquale I, *et al.* Duodenal and faecal microbiota of celiac
children: molecular, phenotype and metabolome characterization. BMC Microbiol
2011;11:219.

48. El Mouzan M, Al-Hussaini A, Serena G, *et al.* Microbiota profile of new-onset celiac
disease in children in Saudi Arabia. Gut Pathog 2022;14:37.

49. Sánchez E, Laparra JM, Sanz Y. Discerning the Role of Bacteroides fragilis in Celiac
Disease Pathogenesis. Appl Environ Microbiol 2012;78:6507–6515.

50. Sánchez E, Donat E, Ribes-Koninckx C, *et al.* Intestinal Bacteroides species associated
with coeliac disease. Journal of Clinical Pathology 2010;63:1105–1111.

51. Leonard MM, Valitutti F, Karathia H, *et al.* Microbiome signatures of progression toward
celiac disease onset in at-risk children in a longitudinal prospective cohort study.
Proceedings of the National Academy of Sciences 2021;118:e2020322118.

52. De Palma G, Nadal I, Collado MC, *et al.* Effects of a gluten-free diet on gut microbiota
and immune function in healthy adult human subjects. Br. J. Nutr. 2009;102:1154–1160.

53. Valitutti F, Cucchiara S, Fasano A. Celiac Disease and the Microbiome. Nutrients
2019;11:2403.

54. Lionetti E, Dominijanni V, Iasevoli M, *et al.* Effects of the supplementation with a
multispecies probiotic on clinical and laboratory recovery of children with newly
diagnosed celiac disease: A randomized, placebo-controlled trial. Digestive and Liver
Disease 2023;55:1328–1337.

55. Dieterich W, Schuppan D, Schink M, *et al.* Influence of low FODMAP and gluten-free diets
on disease activity and intestinal microbiota in patients with non-celiac gluten sensitivity.
Clin Nutr 2019;38:697–707.

56. Bonder MJ, Tigchelaar EF, Cai X, *et al.* The influence of a short-term gluten-free diet on
the human gut microbiome. Genome Med 2016;8:45.

57. Sample D, Fouhse J, King S, *et al.* Baseline Fecal Microbiota in Pediatric Patients With
Celiac Disease Is Similar to Controls But Dissimilar After 1 Year on the Gluten-Free Diet

[Internet]. JPGN reports 2021;2[cited 2023 Nov 24] Available from:
https://pubmed.ncbi.nlm.nih.gov/37206457/

58. Gophna U, Konikoff T, Nielsen HB. Oscillospira and related bacteria – From metagenomic
species to metabolic features. Environmental Microbiology 2017;19:835–841.

59. Karcher N, Pasolli E, Asnicar F, *et al.* Analysis of 1321 Eubacterium rectale genomes from
metagenomes uncovers complex phylogeographic population structure and subspecies
functional adaptations. Genome Biol 2020;21:138.

60. Iyer LM, Zhang D, Rogozin IB, *et al.* Evolution of the deaminase fold and multiple origins
of eukaryotic editing and mutagenic nucleic acid deaminases from bacterial toxin
systems. Nucleic Acids Res 2011;39:9473–9497.

61. Ruiz L, Bottacini F, Boinett CJ, *et al.* The essential genomic landscape of the commensal
Bifidobacterium breve UCC2003. Sci Rep 2017;7:5648.

62. Amiri P, Hosseini SA, Ghaffari S, *et al.* Role of Butyrate, a Gut Microbiota Derived
Metabolite, in Cardiovascular Diseases: A comprehensive narrative review [Internet].
Frontiers in Pharmacology 2022;12[cited 2024 Jan 10] Available from:
https://www.frontiersin.org/articles/10.3389/fphar.2021.837509

63. Liu H, Wang J, He T, *et al.* Butyrate: A Double-Edged Sword for Health? Adv Nutr
2018;9:21–29.

64. Baldi S, Menicatti M, Nannini G, *et al.* Free Fatty Acids Signature in Human Intestinal
Disorders: Significant Association between Butyric Acid and Celiac Disease. Nutrients
2021;13:742.

65. Tjellström B, Högberg L, Stenhammar L, *et al.* Faecal short-chain fatty acid pattern in
childhood coeliac disease is normalised after more than one year's gluten-free diet.
Microb Ecol Health Dis 2013;24:10.3402/mehd.v24i0.20905.

66. De Palma G, Nadal I, Medina M, *et al.* Intestinal dysbiosis and reduced immunoglobulin-
coated bacteria associated with coeliac disease in children. BMC Microbiol 2010;10:63.

67. Sánchez E, De Palma G, Capilla A, *et al.* Influence of Environmental and Genetic Factors
Linked to Celiac Disease Risk on Infant Gut Colonization by Bacteroides Species ▽ . Appl
Environ Microbiol 2011;77:5316–5323.

68. De Palma G, Capilla A, Nova E, *et al.* Influence of Milk-Feeding Type and Genetic Risk of
Developing Coeliac Disease on Intestinal Microbiota of Infants: The PROFICEL Study. PLoS
One 2012;7:e30791.

69. Esteban-Torres M, Ruiz L, Rossini V, *et al.* Intracellular glycogen accumulation by human
gut commensals as a niche adaptation trait. Gut Microbes 15:2235067.

70. Brüssow H. Problems with the concept of gut microbiota dysbiosis. Microb Biotechnol
2019;13:423–434.

71. Tojo R, Suárez A, Clemente MG, *et al.* Intestinal microbiota in health and disease: role of
bifidobacteria in gut homeostasis. World J Gastroenterol 2014;20:15163–15176.

72. Ferretti G, Bacchetti T, Masciangelo S, *et al.* Celiac Disease, Inflammation and Oxidative
Damage: A Nutrigenetic Approach. Nutrients 2012;4:243–257.

73. Ramírez-Sánchez AD, Zühlke S, Aguirre-Gamboa R, *et al.* Gene expression and eQTL
analysis reflect the heterogeneity in the inflammatory status of the duodenal epithelial
lining in coeliac disease [Internet]. 2024;2024.02.29.582756.[cited 2024 Mar 7] Available
from: https://www.biorxiv.org/content/10.1101/2024.02.29.582756v1

74. Galhardo RS, Hastings PJ, Rosenberg SM. Mutation as a Stress Response and the
Regulation of Evolvability. Critical Reviews in Biochemistry and Molecular Biology
2007;42:399.

75. Radman M. SOS repair hypothesis: phenomenology of an inducible DNA repair which is
accompanied by mutagenesis. Basic Life Sci 1975;5A:355–367.

76. Prudhomme M, Attaiech L, Sanchez G, *et al.* Antibiotic stress induces genetic
transformability in the human pathogen *Streptococcus pneumoniae*. Science
2006;313:89–92.

77. Slager J, Kjos M, Attaiech L, *et al.* Antibiotic-induced replication stress triggers bacterial
competence by increasing gene dosage near the origin. Cell 2014;157:395–406.

869