Repeatability of an extended phenotype: potential causes and consequences of nest variation in the Northern House Wren (Troglodytes aedon aedon)

Running Title: Repeatability of nest construction

Chandler E. G. Carr[1,*], Zoë M. Swanson[1], and Dustin G. Reichard[1]

[1]Ohio Wesleyan University, Department of Biological Sciences, 61 S Sandusky St, Delaware, OH 43015

*Corresponding author: Chandler E. G. Carr

ACKNOWLEDGMENTS

We thank Ohio Wesleyan University, the Big Walnut School System, and the Davis, Koban, and Fink families for access to facilities and field sites. We also like to thank Ali Amer for helping in nest dissection, Ava Swanson for the creation of the House Wren and box drawings used in Figures 3 and 7, and Martin Stoffel for statistical advice. Finally, we would like to thank the House Wrens for making our work possible. This project was funded by the Ohio Wesleyan University Summer Science Research Program.

All procedures were reviewed and approved by the Ohio Wesleyan University Institutional Animal Care and Use Committee (Protocol Number: 05-2021-06), U.S. Federal Bird Banding Lab (Permit Number: 24035), and State of Ohio Department of Natural Resources (Permit Numbers: 23-009, 23-010).

Keywords: Extended Phenotype, Repeatability, Individual Preferences, Construction Behavior, Cavity-nesting, Niche Construction Hypothesis, Nest Building

LAY SUMMARY

- Repeatability of construction behaviors in the wild is understudied
- House Wrens are a cavity-nesting songbird
- House Wren nests are highly variable in their morphology
- Female House Wrens are repeatable in cup composition, but not nest dimensions
- The dimensions of the nest are built to match the dimensions of the cavity
- House Wren nests are unrelated to nestling survival or female condition
- Female preferences for nest building may explain nest variation

ABSTRACT

Construction behavior is an aspect of the extended phenotype that allows organisms to build structures that alter their environments in potentially beneficial ways. Although individuals vary in the expression of this extended phenotype (e.g., structure morphology), the repeatability of construction behaviors remain understudied, especially among free-living populations. Many oviparous taxa construct nests, making them of particular interest because variation in nest architecture may directly affect fitness. Using a free-living, cavity-nesting songbird, the Northern House Wren (*Troglodytes aedon aedon*), as our model, we estimated the contribution of the primary builder (the female) to nest variability by measuring the repeatability of nest morphology between successive clutches. We further examined whether nest morphology was related to the dimensions of the nesting cavity, breeding date, or nest success. We found the composition of the cup lining to be a highly repeatable behavior for the nesting female, although the size and composition of the structural platform appeared more related to the dimensions of the cavity. Nest morphology remained variable throughout the breeding season, showing no significant correlations with breeding date, and it was unrelated to clutch size or offspring survival. Our study suggests that variation in construction behavior is likely the product of multiple factors including the preferences of the builder and physical constraints. The absence of any clear links between construction behavior and fitness indicates that nest morphology is not under strong selection. As a result, diverse female building preferences may explain the extreme among-individual variation in nest structure in this species.

INTRODUCTION

Alfred Russel Wallace (1865) once stated: "…so great is the tendency to vary that it is difficult to find two individuals exactly alike." Phenotypic variation is both ubiquitous and a fundamental requirement for evolution (Darwin & Wallace 1858). As such, a considerable amount of research has been devoted to understanding the causes and consequences of among-individual variation. The interaction between an organism's phenotype and the environment determines its success, and selection can push populations toward one or more optimal phenotypes (van Rijssel et al. 2018, LeGrice et al. 2019). However, the relationship between the phenotype and the environment can be bi-directional, allowing individuals to reshape their environments in ways that affect their fitness (i.e., "niche construction"; Laland et al. 1999). For example, an individual managing a thermoregulatory challenge might rely on costly physiological mechanisms like shivering or increasing adipose tissue, but an alternative behavioral mechanism, such as constructing a burrow, could insulate the animal through the construction of a *new* environment, thereby buffering against this selective pressure (Clark et al. 2020).

Construction behavior is a unique behavioral phenotype where the product of the construction is an extension of the builder (i.e., "extended phenotype"; Dawkins 1982) that modifies the environment to be more beneficial for the builder. Construction behaviors may enhance the builder's fitness by providing protection from predators (Beebe 1953, Hermes, 2013, Sugiura 2016) and environmental exposure (Zeibis et al. 1996, Korb & Linsenmair 2000, Korb 2003, Rosell & Campbell-Palmer 2022), as well as increasing access to resources such as food (Hölldobler & Wilson 1990, Miyashita & Shinkai 1995, Blamires et al. 2010) and mates (Borgia 1985, McKaye et al. 1990). As a result, construction behavior has become

taxonomically widespread across both vertebrate and invertebrate species (Hansell 2005).

Eusocial insects build colonial shelters (e.g. termite mounds), some mammals dig burrows,

most spiders spin webbed traps, and certain species of fishes and birds construct bowers.

However, despite the clear potential for construction behavior to benefit the fitness of an

individual, trade-offs often occur between investment in the construction process and other

behaviors that affect fitness which can result in negative effects on both the builder (Madden

2002, Bailey et al. 2015) and the structure (Ellendula et al. 2021, Parthasarathy et al. 2022).

For example, investing more time in the construction process potentially reduces available

energy for foraging or mate-searching.

Many organisms build multiple versions of the same structure throughout their lifetimes,

which provides an opportunity to evaluate the consistency of this behavior. It is well

established that the products of construction behaviors exhibit among-individual variation, but

the extent and cause of variation *within* an individual's construction behavior is less clear

(DiRienzo & Aonuma 2018). For example, an individual's construction behavior might be

consistent, plastic, or some combination of the two as its experience and environment changes.

Repeatability estimates, calculated as intra-class correlation coefficients, can be used to assess

how much of the variation observed among individuals can be explained by variation within

individuals (Hayes & Jenkins 1997). When among-individual variation is high and

within-individual variation is low, then the repeatability estimate will be high and that trait will

be considered repeatable. Repeatability estimation can be used to understand the extent to

which phenotypic traits are the product of static characteristics, such as heritability ("upper

level of heritability"; Bell et al. 2009, Møller 2006) and individual preferences (Mennerat et al.

2009, Järvinen & Brommer 2020, Silva et al. 2024), or plastic characteristics, such as

experience (DiRienzo et al. 2019, Whittaker et al. 2023). For this reason, the repeatability of a large variety of behaviors have been examined across taxa (Bell et al. 2009), but there are few studies examining the repeatability of construction behaviors. Most of these studies have focused on spider web structure (Fisher et al. 2019, DiRienzo et al. 2020, DiRienzo & Montiglio 2016) or nest morphology (Rushbrook et al. 2008, Muth & Healy 2011, Whittaker et al. 2023), but few have examined the behavior exclusively in the wild (Møller 2006, Järvinen et al. 2017, Silva et al. 2024).

Nest building is a widespread construction behavior among oviparous taxa that serves the primary function of providing a safe, hospitable space for eggs or altricial young to develop (Mainwaring et al. 2014a). With few exceptions (e.g., brood parasites), bird species produce an incredible array of nests that can be structurally complex or simple, but must contain at least one of four zones: an outer nest layer, an attachment to the environment, a structural layer, and a lining (Hansell 2000). For example, scrape nests, built on the ground, may be considered just structural while nests built in trees often require an attachment, structure, and a cup lining. Zones are often discernible from one another by the materials used, with zone composition being largely species specific (Biddle et al. 2018). Nests can be built by the male, female, or both with some division of labor by zone (Hansell 2000). The morphology of the nest, structure, size, and composition, can be important for providing the same benefits of niche construction observed for other structures by protecting the young from predators (Møller 1990, Hays et al. 2022) and environmental exposure (Franklin 1995, Mainwaring et al. 2014b, Cerezo & Deeming 2016, Perez et al. 2020, Medina et al. 2022), as well as potentially playing a role in mate attraction for some species (Ueda 1985, Mainwaring et al. 2008, Jelínek et al. 2016).

Here, we investigated potential causes and consequences of the variation and repeatability of Northern House Wren (*Troglodytes aedon aedon*) nest morphology and composition. House Wrens are secondary cavity nesters that raise two to three clutches a season, often reusing the same nest throughout (Johnson 2020). They will, however, build a new nest when the previous nest is stolen, destroyed, or experimentally removed (Alworth 1996), making them an ideal model for the examination of repeatability in construction behavior. House Wren nests contain 2 to 3 zones: sometimes an outer layer consisting of spider egg sac material, a structural zone made of various types of sticks, henceforth referred to as the "stick platform," and a cup defined as a simple depression, at the top of the stick platform, lined with grasses, feathers, and other material (McCabe 1965). Although the layout and composition of the zones are considered standard, House Wren nests are highly variable in their morphology. For example, in some nests the stick platform may be built so large that it covers the entrance hole of the box (McCabe 1965; Figure S1A) while other nests have a stick platform only a few centimeters in height (Figure S1B). While establishing a territory, unmated males place a variable number of sticks (range: <10 to >400; Johnson 2020) into potential nesting cavities. After pairing, the female completes the stick platform (Alworth & Scheiber 2000, Johnson 2020) and creates a lined cup either before or during egg laying. Considering the extensive effort of the female in both completing the stick platform and constructing the nest cup, we defined the female as the primary nest builder in our study.

Although the architecture of the nests of other bird species have shown to be beneficial to the fitness of the builder (Mainwaring et al. 2014), House Wren nest architecture appears to be unrelated to the mate selection process, the survivability of offspring (Alworth 1996, Eckerle & Thompson 2005), or the limiting of brood parasitism (Pribil & Picman 1997). Thus, the

available evidence suggests that nest architecture in this species has limited effects on fitness, and so it remains unclear why House Wren nests exhibit such extreme variation among individuals. The absence of selective pressure on nest architecture could be one explanation for the amount of variability observed. Alternatively, some research suggests that constraints imposed by the cavity dimensions and the accessibility of building materials may also drive variation in nest architecture (Stanback et al., 2013). However, no study to date has explored the role of the preferences of females during nest construction, such as the types and quantity of material used, as an explanation for the variation. By sampling successive nests built by the same female one can investigate whether nest architecture is repeatable at the individual level. If nest architecture *is* repeatable, then differences between individual females in their building preferences can provide a previously underappreciated explanation for the high intraspecific variation in House Wren nests.

We sampled House Wren nests across an entire breeding season to assess the contribution of the female builder, the box cavity, and breeding date to nest architecture. We also examined whether variation in nest characteristics correlated with one measure of female body condition (residual between the individual's mass and tarsus length), clutch size, or offspring survival, which provided proxies of fitness. Unlike most construction behavior research that often relies on coarse measures of structure like nest weight (Mainwaring et al. 2008, Walsh et al. 2010, Adreani et al. 2022), we used a finer approach of directly quantifying the individual building materials (e.g., exact count of sticks) used in nest construction.

We predicted that nests would have high variability among individuals but that there would be low within-individual variability. Specifically, since the female builds the majority of the structure after pairing, the female identity would be the best predictor of nest dimensions

and composition as measured by a repeatability estimate. If supported, this outcome would

highlight the importance of female preferences for specific nest characteristics and building

materials in determining nest architecture. Furthermore, based on previous research in other

House Wren populations (Alworth 1996, Eckerle & Thompson 2005, Stanback et al., 2013),

we predicted that nest architecture would not relate to offspring survival, but may relate to the

dimensions of the cavity in which it was built or the breeding date when it was measured.

Successive nests that are built in the same box should exhibit repeatability for those

characteristics that are most affected by the box's dimensions. Furthermore, seasonal changes

in other variables such as weather and vegetative cover might favor different nest architecture

as breeding progresses. For example, thinner nest cups could be favored to provide less

insulation as average temperatures increase (Mertens 1977). Alternatively, because clutch size

declines in this species across the breeding season (Johnson 2020), females might build smaller

nests over time due to fewer offspring or a more general decline in reproductive investment

later in the season.

METHODS

Study System

We monitored Northern House Wren (*Troglodytes aedon aedon*) breeding behavior in

160 nest boxes across six sites in Delaware and Union counties in Ohio, U.S.A., between May

9 and August 25 of 2022. Of those 160 boxes, 79 produced at least one nest with a complete

clutch, which was the threshold for inclusion in our study. With the exception of three nest

boxes located in suburban yards, the majority of nest boxes were located in edge habit

separating grass from forest. We measured nest box dimensions with digital calipers (nearest

0.001 mm). Nest boxes were rectangular wooden cuboids with internal dimensions of

approximately 110 x 115 x 195 mm (Figure 1, see Table 1 for variation in box dimensions).

Each box possessed a circular entrance hole with an average diameter of 34 mm and depth of

46 mm, drilled an average of 119 mm from the bottom of the box. Most ($n = 73/79$; 92.4%)

boxes also had a rectangular gap separating the top of the front or side panel from the roof that

allowed the panel to swing open. This gap was the same width as the box with a height ranging

from 4 mm to 30 mm. Some wrens used this gap as an entrance, particularly when their nest

obstructed the primary entrance hole. The variation in box dimensions were due to imperfect

construction and the preferences of landowners for slightly different box designs. All boxes

were suspended about 1.75 meters off the ground by a greased metal conduit pole to limit

predation.

We monitored 108 total nests (some boxes were used multiple times), but were unable to

visit boxes daily, so exact aging of nests was not always possible. However, House Wrens have

predictable rates of egg laying and incubation time, so we were able to estimate first egg dates

by counting eggs and incubation start dates by backdating from hatch date (Johnson 2020). In

cases where we were not present for hatch day, nestling ages and estimated hatch date were

evaluated by measuring nestling weights, wing chord length, and feather tract development as

described by Brown et al. (2011).

As soon as we discovered a complete clutch, the female was caught in a mist net at the

nest box and banded with a USFWS aluminum leg band and a unique combination of plastic

colored bands for future identification. It was essential to capture and band females to sample

repeated nesting attempts, but we chose not to capture males to minimize further disrupting

both our experiment and a separate, unrelated study. The female's weight (nearest 0.1 g; Pesola

spring scale), tarsus length (nearest 1 mm; ruler), and wing length (nearest 1 mm; ruler) were

also measured. While the female was processed, we measured the height of the nest and the

width of the "berm" (Stanback et al. 2013), which was defined as the front wall of the cup

directly against the entrance hole (see Figure 1), using a dial caliper (nearest 0.1 mm). On day

eight of nestling life, we collected measurements and banded each nestling as described above

for adult females. We continued to monitor nests until the clutch fledged or failed, at which

point the nest was collected in a large and resealable plastic bag for dissection in the lab and to

prevent the nest from being reused or built upon for a successive clutch.

Figure 1 - Model of the nest box and nest (in orange). Both the front (left) and side (right) of the box are shown. A: height of the rectangular gap, B: diameter of the entrance hole, C: distance from the bottom of the box to the bottom of the entrance hole, D: internal length of the box and length of the nest, E: depth of the entrance hole, F: width of the berm, G: internal width of the box and width of the nest, H: height of the nest, I: internal height of the box.

234 *Table 1 - Average, standard deviation, and range of the dimensions of the nest boxes used in*

235 *this study. See Figure 1 for a model of an average nest box.*

	Average (mm)	Standard Deviation (mm)	Range (mm)
Internal Box Height	195.19	14.92	172.20 - 219.90
Internal Box Volume	2462.20	434.52	1812.99 - 4133.73
Entrance Diameter	33.60	5.22	27.90 - 44.65
Entrance Depth	44.11	6.78	25.00 - 56.40
Height of Entrance	118.94	21.19	83.10 - 151.60
Gap Height (when present)	14.82	6.11	4.30 - 29.55

Nest Analysis

After collection, we placed nests in a -20 C freezer for 24 - 48 hours to kill any

arthropods. For dissection, nests were placed on a large sheet of white paper and carefully

pulled apart to quantify nest materials and characteristics. Most nests were dissected by the

lead author, but two assistants were also trained for dissection. We made exact counts of

individual sticks, grasses, plumulaceous, and pennaceous feathers. Sticks and grasses needed to

be larger than an arbitrary length of 4 cm in order to be counted. Smaller pieces of vegetation

were not quantified along with other debris not described below. Down, contour, and

semiplume feathers were categorized as plumulaceous, while wing and tail feathers were

categorized as pennaceous. We used an ordinal scale of 0 to 3 to quantify the amount of fecal

material and spider egg sacs present within each nest. Finally, we noted, as a binary yes or no,

the presence of snake skin and anthropogenic objects (e.g., plastic).

Statistical analysis

Although we sampled 108 nests in total, a small number had to be excluded from our

final analysis. One nest was removed during dissection because it contained signs of reuse (i.e.,

two individual grass layers separated by a thick stick layer). Two nests were not collected from

the field due to reuse or extensive nestling decomposition after predation, and so were unable

to be dissected. Due to data loss, the dissection measurements of two additional nests are

unaccounted for. Our final sample size was 103 nests.

R (version 4.2.3) was used for all analyses. We first assessed whether the observed

variation in nest measurements was normally distributed using separate Shapiro-Wilk tests. To

test whether nest measurements were repeatable within females, we calculated repeatability

estimates (packages: rptR, version 0.9.22) with the nest measurement as the dependent

variable, and the identity of the nesting female as the grouping factor. We used the functions

rptGaussian or rptBinomial depending on the nest measurement being explored. Parametric

bootstrapping and permutation, at 10,000 iterations each, were performed to produce

confidence intervals and adjusted alpha values for the repeatability estimates, respectively.

Significant repeatability estimates were indicated by 1) confidence intervals not overlapping

zero (DiRienzo & Aonuma, 2018) and 2) an adjusted alpha value of 0.05. Where nest

measurement was binomial, the link-scale approximation (Stoffel et al. 2017) is reported. We

used the same approach to test whether the box identity predicted nest measurements. In these

models, the nest measurement was the dependent variable, and the identity of the box was the

grouping factor. Although there was some overlap in sampling between models with the same

female nesting more than once in the same box, the sample size (5/24; 21% of females reused

the same box and 6/27; 22% of boxes were used by the same female) was too small to provide

a reliable estimate of the effect of these repeated samples on nest measurements. Thus, we

chose to avoid a more complex model that included both box and female identity. Finally, we

defined repeatability estimates less than 0.20 as low, between 0.20 and 0.40 as moderate, and

greater than 0.4 as high repeatability (Bohn et al. 2017).

Linear regressions (Base R Package) were used to test for a relationship between nest

measurements and box proportions, date of measurement, and age of the nest, which was

defined as the number of days between the first egg and the nest collection date. We used

generalized linear models (Base R Package) to analyze the effect of nest measurements on the

outcome of the clutch. These models assumed a binomial distribution with nest outcome

(fledge/fail) as the dependent variable. Linear models were used to analyze the effect of nest

measurement on clutch size and fledge count, assuming a gaussian distribution. Both the

generalized linear models and linear models had the nest measurement as the fixed effect, and

the date of nest measurement as a covariate to control for seasonal variation in reproductive

output.

RESULTS

Variability in Nest Architecture and Correlations Between Nest Components

House Wren nests were highly variable (Figure 2). The ratio of the nest height to internal

box height ranged from 0.254 to 0.902 (Figure 2A; $\underline{x} \pm$SD: 0.59 ± 0.144). Similarly, the

volume of the nest ranges from 655.71 mm³ to 2776.96 mm³ (Figure 2B; $\underline{x} \pm$SD: 1437.44 ±

409.29). Sticks were the primary contributor to the height and volume of the nest, and we

found that stick count was also variable, ranging from 170 to 1491 sticks in a single nest

(Figure 2C; x ±SD: 676.14 ± 25.01). The width of the nest berm ranged from 7.00 mm to

77.45 mm (Figure 2D; x ±SD: 39.62 ± 1.27).

We found the number of grass strands within the cup of a single nest ranged from zero to

836 (Figure 2E; x ±SD: 210.19 ± 16.49). Feather count within the cup ranged from a singular

pennaceous feather to 271 total feathers (Figure 2F; x ±SD: 83.79 ± 5.70). Plumulaceous

feathers were the most common feather found in nests with an average ratio of 5 plumulaceous

feathers for every 1 pennaceous. Although the cup is primarily lined with grasses and feathers,

other materials such as snake skin ($n = 19/103$; 18.4%), anthropogenic objects (e.g., plastic; $n =$

54/103; 52.4%), and hair, noted anecdotally but not quantified, from a variety of mammalian

species, including human, were observed. We also frequently found varying amounts of spider

egg sac material interspersed throughout the stick platform ($n = 88/103$; 85.4%).

The ratio of the nest height to internal box height (P = 0.74; Figure 2A) and the width of

the nest berm (Figure 2D; P = 0.94) were normally distributed across the population. All other

continuous nest characteristics showed a right-skewed distribution (P < 0.001).

The width of the berm positively correlated with the height ($r^2 = 0.03$, $P = 0.04$), volume

($r^2 = 0.09$, $P < 0.001$, Figure S2A), and stick count ($r^2 = 0.04$, $P = 0.02$, Figure S2B) of the

nest. The stick count positively correlated with the height ($r^2 = 0.37$, $P < 0.001$, Figure S2C)

and volume ($r^2 = 0.41$, $P < 0.001$, Figure S2D) of the nest. Total feather count ($r^2 = 0.03$, $P =$

0.04, Figure S2E) and pennaceous feather count ($r^2 = 0.06$, $P = 0.005$, Figure S2F), but not

plumulaceous feather count ($P = 0.15$) positively correlated with the grass count. No other

correlations were statistically significant ($P > 0.19$, Table S2).

Figure 2 - Histograms showing the number of nests that contained the specified variable.

"Height Ratio" (A; P = 0.74) and "Width of the Nest Berm" (D; P = 0.94) exhibit a normal

distribution, but other factors show a right-skewed distribution (P < 0.001). The blue plots

layered over the histograms represent the outcome of an estimated density distribution.

Variation Explained by Female Builder Identity and Body Condition

Over the course of the nesting season, 24 females renested in our boxes after their initial

clutch. Of those females, 19 renested once and 5 renested twice on our study site, and the

majority of those females renested in a different box from any previous clutch (22/29 total

renests; 75.9%; Table S1). The identity of the nesting female did not explain the variation in the general nest dimensions, such as the ratio of the nest height to internal box height ($R = 0.24$ [95% CI = 0 - 0.56], $P = 0.08$, Figure 3C), nest volume ($R = 0.22$ [95% CI = 0 - 0.53], $P = 0.11$), or the the width of the berm ($R = 0$ [95% CI = 0 - 0.34], $P > 0.99$, Figure 3E). The female identity also did not explain the number of sticks within the stick platform ($R = 0.15$ [95% CI = 0 - 0.49], $P = 0.20$, Figure 3A). However, when considering the cup lining, the grass material count was highly repeatable for the female ($R = 0.56$ [95% CI = 0.23 - 0.77], $P = 0.003$, Figure 3B). While the total count of feathers in the cup was not repeatable ($R = 0.04$ [95% CI = 0 - 0.39], $P = 0.37$), the pennaceous feather count appeared moderately repeatable ($R = 0.33$ [95% CI = 0 - 0.62], $P = 0.05$, Figure 3F), although we do note that the confidence interval overlaps with zero. The ratio of pennaceous feathers to plumulaceous feathers within the cup of the nest was highly repeatable ($R = 0.90$ [95% CI = 0.79 - 0.95], $P = 0.002$, Figure 3D). No other individual measurement was repeatable (Table 2). No significant correlations between the body condition of the female builder and nest morphology were observed ($P > 0.11$, Table S3).

Table 2 - Listed nesting materials were not significantly repeatable for both female and nesting box identity. Material specified with "†" was measured as a simple yes/no presence.

	Box ID		Female ID	
	R & 95% CI	P	R & 95% CI	P
Plumulaceous	0.03 [0 - 0.38]	0.39	0.02 [0 - 0.37]	0.43
Fecal Rate	0.07 [0 - 0.24]	< 0.99	0.07 [0 - 0.25]	< 0.99
Spider Rate	0.07 [0 - 0.24]	< 0.99	0.07 [0 - 0.23]	< 0.99

| † Snake Skin | <0.001 [0 - 0.88] | 0.64 | 0.14 [0 - 0.93] | 0.17 |
| † Plastic | 0.04 [0 - 0.35] | 0.32 | <0.001 [0 - 0.32] | 0.48 |

Figure 3 - The repeatability of nest measurements across successive clutches for individual

females. Characteristics of the stick platform are shown in orange (left), and characteristics of

the cup lining are shown in blue (right). Significant repeatability by female identity is indicated

by the presence of a House Wren drawing. Female House Wrens construct similar nest cups

across multiple nesting attempts with the grass count (B; R = 0.56, P = 0.004), feather ratio

(D; R = 0.90, P = 0.002), and pennaceous feather count (F; R = 0.33, P = 0.05) each

indicating significant repeatability for females. However, none of the characteristics of the

stick platform showed significant repeatability (P > 0.08).

Variation Explained By Nest Box Identity and Dimensions

Twenty-six of our boxes were used more than once over the course of the nesting season. Of the boxes, 23 were used twice while 3 were used three times. Of the boxes used more than once, the majority of the nesting attempts were from a different female than any previous attempt (22/29 total reuses; 75.9%; Table S1). With respect to general nest dimensions, the ratio of the nest height to internal box height ($R = 0.56$ [95% CI = 0.25 - 0.77], $P < 0.001$, Figure 4C), nest volume ($R = 0.65$ [95% CI = 0.37 - 0.82], $P < 0.001$), and the width of the nest's berm ($R = 0.49$ [95% CI = 0.15 - 0.72], $P = 0.002$, Figure 4E) were all highly repeatable within a nest box. The stick platform, measured by the number of sticks in the nest, was also highly repeatable within a nest box ($R = 0.45$ [95% CI = 0.10 - 0.70], $P = 0.008$, Figure 4A). However, the components of the nest cup, the grass count ($R = 0.04$ [95% CI = 0 - 0.38], $P = 0.38$, Figure 4B), total count of feathers in the nest ($R = 0$ [95% CI = 0 - 0.34], $P > 0.99$) or the pennaceous feather count ($R = 0.08$ [95% CI = 0 - 0.42], $P = 0.32$, Figure 4F) were not statistically repeatable. The pennaceous to plumulaceous feather ratio showed moderate repeatability within a nest box ($R = 0.29$ [95% CI = 0 - 0.59], $P = 0.05$, Figure 4D), but the confidence interval overlaps with zero. The box ID did not explain the amount of spider egg sac material in the nest ($R = 0$ [95% CI = 0 - 0.21], $P > 0.99$). No other measurements were statistically repeatable (Table 2).

The height ($r^2 = 0.06$, $P = 0.006$) and volume ($r^2 = 0.16$, $P < 0.001$, Figure 5A) of the nest platform positively correlated with the internal height of the box. Likewise, the number of

sticks within the platform positively correlated with the internal height ($r^2 = 0.12$, $P < 0.001$,

Figure 5B) and volume ($r^2 = 0.14$, $P < 0.001$) of the box. The height of the nest was positively

correlated with the height of the rectangular gap above the entrance hole ($r^2 = 0.13$, $P < 0.001$,

Figure 6A). It is worth noting that within our sample, boxes with taller internal heights tended

to have higher rectangular gaps ($r^2 = 0.13$, $P < 0.001$, Figure 6B). The width of the berm

positively correlated with the diameter of the entrance hole ($r^2 = 0.03$, $P = 0.05$, Figure 7A), but

not the depth of the entrance hole ($P = 0.43$, Figure 7B). The closer the top of the nest was built

to the entrance hole (See Figure 1), the larger the width of the berm ($r^2 = 0.09$, $P = 0.001$). No

other measurement correlated with box dimensions (Table 3).

Figure 4 - The repeatability of nest measurements across successive clutches built in the same nest box. Characteristics of the stick platform are shown in orange (left), and characteristics of the cup lining are shown in blue (right). Significant repeatability by box identity is indicated by the presence of a nest box drawing. The stick count (A; R = 0.45, P = 0.007), height ratio (C; R = 0.65, P < 0.001), berm width (E; R = 0.49, P = 0.002), and feather ratio (D; R = 0.29, P = 0.05) were significantly repeatable traits for the nesting box identity, while grass Count (B; P = 0.39) and pennaceous (F; P = 0.32) were not.

Table 3 - P-values of the correlations between nest characteristics and box dimensions.

Significant correlations are indicated by bold text.

	Internal Box Height	Internal Box Volume	Gap Height	Entrance Diameter	Entrance Depth
Nest Height	**0.006**	0.86	**< 0.001**	0.63	0.58
Nest Volume	**< 0.001**	**< 0.001**	0.12	0.23	0.85
Nest Berm	0.42	0.053	0.54	**0.05**	0.43
Stick	**< 0.001**	**< 0.001**	0.12	0.81	0.85
Grass	0.20	0.37	0.84	0.19	0.65
Total Feather	0.60	0.88	0.81	0.27	0.08
Pennaceous	0.65	0.69	0.88	0.30	0.16
Plumulaceous	0.64	0.97	0.73	0.33	0.10

Figure 5 - Significant positive correlations between the internal height of the nesting box and the total volume of the nest (A; $r^2 = 0.16$, $P < 0.001$) and the stick count (B; $r^2 = 0.12$, $P < 0.001$). Gray shading encompasses the 95% confidence interval for the line of best fit.

Figure 6 - Significant positive correlations between rectangular gap height and the height of the nest (A; $r^2 = 0.13$, P < 0.001) and internal box height (B; $r^2 = 0.13$, P < 0.001). Gray shading encompasses the 95% confidence interval for the line of best fit.

Figure 7 - Relationships between the width of the nest berm and the diameter or depth of the entrance hole. There was a significant positive correlation between the diameter of the entrance hole and the width of the nest berm (A; $r^2 = 0.03$, P = 0.05), but no detectable correlation between the depth of the entrance hole and the width of the nest berm (B; $r^2 = 0.09$, P = 0.43). Gray shading encompasses the 95% confidence interval for the line of best fit.

Effects of Measurement Date and Nest Age

The date of measurement did not significantly correlate with the ratio of the nest height to

internal box height ($P = 0.54$) or the width of the berm ($P = 0.74$). Neither the stick count ($P =$

0.72) or grass count ($P = 0.88$) correlated with the measurement date. A statistical trend was

found with the pennaceous feather count positively correlating with the measurement date ($r^2 =$

0.02, $P = 0.07$, Figure S3C), with the result reaching statistical significance with the removal of

an extreme outlier (Date of year of outlier: 144; $r^2 = 0.09$, $P = 0.001$), but this trend was not

found with plumulaceous feathers ($P = 0.27$) or total feather count ($P = 0.15$). The amount of

spider egg sacs within the nest positively correlated with the measurement date ($t = 4.49$, $P <$

0.001, Figure S3D). No other nest component significantly correlated with the date of

measurement ($P > 0.08$, Table S3)

The pennaceous feather count positively correlated with the age of the nest ($r^2 = 0.04$, $P =$

0.02, Figure S4A), but the plumulaceous ($P = 0.43$) and total feather ($P = 0.18$) counts did not.

The grass count ($r^2 = 0.03$, $P = 0.054$, Figure S4D) and stick count ($r^2 = 0.03$, $P = 0.052$, Figure

S4C) marginally negatively correlated with the age of the nest. As expected, younger nests

were also less likely to contain fecal material, with nests older than 20 days almost always

containing some amount of fecal material ($t = 5.24$, $P < 0.001$, Figure S4B). No other nest

component significantly correlated with nest age ($P > 0.052$, Table S2).

Reproductive Success

435   No correlations between clutch size or offspring success and nest morphology were

436   observed ($P > 0.06$, Table S3). Nests sampled later in the breeding season were more likely to

produce successful clutches than nests built earlier in the season (t = 2.04, *P* = 0.04, Figure S3A) but late season nests also fledged fewer offspring (t = -5.89, *P* < 0.001, Figure S3B). However, this result is likely due to females laying fewer eggs later in the season (t = -10.26, *P* < 0.001).

DISCUSSION

Extended Phenotypes are Complex

The niche construction hypothesis predicts that the selective pressures on construction behaviors within a species should be relatively invariable, both spatially and temporally, because the benefits of the constructed product, such as a nest, should be long term and stable (Clark et al. 2020). This invariable selection should, in turn, create consistency in the expression of the constructed product. However, we found Northern House Wren nests to be highly variable both among individuals and across a single breeding season. Consistent with previous research (Alworth 1996, Eckerle & Thompson 2005), the variation was unrelated to female body condition or the number of offspring fledged, two proxies of the builder's fitness, suggesting that nest architecture is not under strong selection. We asked: in the absence of strong selection for specific nest characteristics, what explains the observed variability?

Extended phenotypes are a complex bidirectional interaction between the builder and their environment, with builders attempting to express their phenotype within the confines of the available space and the limitations of available building materials. We found evidence of repeatability for the female builder in multiple components of the cup composition. We also found that nest morphology was affected by the dimensions of the nesting cavity, supported by repeatability in components of successive nests built in the same box, often by different

females. Our findings suggest a largely unexplored explanation for variation in nest

architecture: consistent differences between individual females in their building preferences for

both the types and quantity of building materials. Together with the ability of the cavity to

constrain or enhance the nest dimensions, these two factors help shape the wren's extended

phenotype. We also employed an uncommon method for measuring construction behavior (i.e.,

directly measuring individual building materials) leading to the discovery of repeatable

characteristics that would have otherwise remained hidden with more coarse measurements

like nest weight. Future research should compare nest variation across populations and

breeding seasons to fully understand the spatial and temporal variation of nest architecture in

this species.

Preferences in Construction behavior

Our prediction that the nesting female would be the best predictor of variation in nest

architecture was only partially supported. The cup composition (i.e., the grass count,

pennaceous feather count, and ratio of pennaceous to plumulaceous feathers) was moderately

to highly repeatable for the female, but female identity did not explain variation in the rest of

the nest morphology. This finding reinforces previous conclusions that the female constructs

the nest cup independent of the male (McCabe 1965), but implicates other factors in the

construction of the stick platform, such as physical constraints of the cavity. The cup appears

largely unaffected by these constraints and similarly unrelated to fitness. We hypothesize that

when constraints and selective pressures favoring a specific nest architecture are reduced, the

effect of diverse female building preferences on nest variation is enhanced. Female Blue Tit

(*Cyanistes caeruleus*) nests, for example, are also highly variable in height and this behavior is

moderately repeatable for females (R = 0.40; Järvinen et al. 2017a) despite nest height in this

species appearing unrelated to nestling survival or female body condition (Järvinen et al.

2017b). Therefore, variation between individuals in their construction preferences is likely a

driving force behind much of the observed variability in nest architecture for cavity nesters.

If much of the variation in nest architecture is explained by diversity in individual

preferences, it is important to understand what causes builders to differ from each other.

Although this study is one of the first to tie the repeatability of an extended phenotype directly

to individuals in a free-living population, lab experiments have implicated differences in

experience as a potential factor that affects individual preferences. For example, Zebra Finch

(*Taeniopygia castanotis*) pairs that had previous experience constructing nests built more

repeatable nests during experimental trials (Whittaker et al. 2023) and juvenile black widow

spiders (*Latrodectus mactans*) exposed to conspecific webs built more repeatable webs in

adulthood (DiRienzo et al. 2019). Overall, having prior experiences with construction

behaviors tends to result in individuals building more repeatable structures, with experiences

being either previous attempts or social interactions.

Some females in our study were more consistent builders than others, but we find it

unlikely that experience from prior breeding seasons could have contributed to these

differences. Long-term demographic data from multiple Northern House Wren populations

indicate that the majority of individuals only breed in a single season (average lifespan is 1.5

years; interannual return rates ~30%; Johnson 2020) and so are unlikely to provide a sufficient

sample size for an analysis of interannual individual differences. The age of individuals could

act as a proxy of experience in some species, but we are unable to present an analysis of its

effect on repeatability in this study. While molt patterns allow many songbirds to be aged as

either in their first breeding season or older, this process is challenging in House Wrens (Pyle

1997), and we are not confident in our age assignments.

Individuals are also affected by trade-offs between the expression of the extended

phenotype and other behaviors important for fitness. Starved colonies of *Stegodyphus*

sarasinorum, for example, invest more in web construction than fed spiders (Ellendula et al.

2021), and immune stressed three-spined stickleback (*Gasterosteus aculeatus*) males construct

less compact nests than healthy conspecifics (Barber et al. 2001). Thus, construction behaviors

can be important indicators of the builder's condition. We calculated the linear residual

between body mass and tarsus length as a measure of female condition, but it did not correlate

with the observed variability in nest cup composition. However, we collected this singular

measure after the completion of egg laying, and it may not have been reflective of the female's

condition during nest building, which mostly occurs before egg laying begins (i.e., 5-7 days

before our measurement). Although not measured in this study, it is likely that female body

condition declined over the course of the breeding season due to the energetic costs of

reproduction (Merilä and Wiggins 1997), which could also have affected nest construction. The

clutch size of female House Wrens declines throughout the season, so we predicted that pairs

would similarly put less effort into nest construction as the season progressed. However, we

found no detectable relationships between breeding date and nest architecture. Instead, female

wrens remained consistent in multiple components of the nest cup throughout the breeding

season despite the fact that clutch size declines over time. Nest cup composition also did not

differ between successful or unsuccessful nests or correlate with the number of eggs laid, and

remained variable throughout the breeding season. Conversely, we found that nests collected

longer after the initiation of construction tended to contain more pennaceous feathers than nests

collected sooner, and even observed one female returning to her complete clutch of eggs with a

feather in her bill, indicating a potential upkeep behavior in cup composition despite the

increasing needs of the offspring. The lack of any apparent reproductive cost of the nest

architecture conflicts with the results of other studies on extended phenotypes. For example,

Zebra Finches modify successive nests when the previous clutch fails (Edwards et al. 2020).

However, this finding remains consistent with previous literature on Northern House Wrens

(Alworth 1996, Pribil & Picman 1997, Eckerle & Thompson 2005). We discuss potential

causes for this lack of fitness-related nest architecture below.

Physical Constraints and Environmental Pressures on Constructions

The box cavity was also a significant predictor of nest morphology and composition,

specifically that of the stick platform. This repeatability supports the prediction that nest

architecture would be related to constraints posed by the cavity dimensions. House Wrens

appeared to be filling the available space and adjusting nest morphology based on differences

between box openings. Stanback et al. (2013) suggested that wrens were modulating the nest

structure to accommodate for perceived vulnerabilities in the cavity, and our data support this

conclusion. For example, the size of the entrance hole is a potential source of vulnerability, as

more accessible nests are more likely to be depredated and parasitized (Paclík et al. 2009,

Hauber et al. 2024). Alternatively, larger entrances could increase heat loss, increasing the

threat of hypothermia for the offspring (Lamprecht and Schmolz 2004). We found that House

Wrens built wider berms in boxes with larger entrance hole diameters, potentially modulating

their nest structure to better protect the contents of the nest from predators or to increase heat

retention. We found a similar correlation between the height of the rectangular gap and the

height of the nest, as larger gaps resulted in taller nests. Previous research has suggested that

the berm could act as a barrier between the nestlings within the cup and any outside threats

(Stanback et al. 2013), however evidence for this relationship is inconsistent (Pribil & Picman,

1997). Despite our support of Stanback et al.'s (2013) finding, the width of the nest berm had

no effect on the number of eggs laid or nestling survival. Variation in the width of the berm

also did not change throughout the breeding season, despite the prediction that nest architecture

that reduces insulation would be favored as temperatures increased. House Wrens may be

modulating their construction behaviors in response to perceived vulnerability, but our results

suggest that this action has little to no effect on nest success. Previous literature has implicated

the climate in affecting the evolution of nest characteristics, but this relationship appears to be

affecting nest shape at the among- rather than within-species level (e.g., domed v. open-cup

nesting species) and still explains only a small portion of the observed variation (Colombo et

al. 2024).

However, evidence from other species still suggests that structures protect inhabitants

from predators and environmental exposure by the construction of a physical barrier. For

example, eurasian beavers (*Castor fiber*) in the Netherlands, which face lower predation rates

and milder climates than their northern counterparts, build lodges with thinner walls (Rosell &

Campbell-Palmer 2022). Cavity-nesters are unique in that they do not need to construct a

physical barrier, as the cavity itself is the barrier between the inhabitants and the outside

environment. Therefore, we hypothesize that the lack of any detectable relationship between

nest architecture and fitness in House Wrens could be due to selection favoring a better quality

cavity rather than specific characteristics of the nest. Observations of platypus

(*Ornithorhynchus anatinus*) nests, which are housed inside burrows, are consistent with this

hypothesis as the dimensions of the burrow explained offspring production, while effort in nest construction was unrelated (Thomas et al. 2018). Support for this hypothesis in House Wrens is more indirect, but males with empty cavities due to experimentally removed nests were just as likely to acquire a mate as control males that initiated nest building (Alworth 1996), suggesting that the cavity rather than the contents are more important in mate choice. Female House Wrens also use cavity quantity before male quality when choosing mates (Eckerle & Thompson, 2006) and male House Wrens selectively choose nesting boxes with smaller entrances that likely provide superior protection from predators and brood parasites (Pribil & Picman 1997). Collectively, the available data suggest that the reproductive success of House Wrens is more related to the cavity itself than the nest.

Limitations

Extended phenotypes become additionally complex when more than one individual contributes to the construction effort. Both sexes contribute to the nest building in House Wrens, so it is important to consider how the male contribution may affect our data and interpretation. Although their contributions are likely not equal, with the female completing the majority of the nest (Alworth & Scheiber 2000), we were unable to directly account for the effect of individual males on construction behavior because the majority of males were unbanded. Despite this limitation, Northern House Wren pairs often change between nesting attempts (Baldwin 1921, Drilling & Thompson 1988), so it is rarely the same pair building successive nests. In 2019, we found that out of 31 banded pairs in our population, only 6 (19.4%) remained with the same mate for the entire breeding season, with the female being more likely to leave a territory than the male. The fact that females were repeatable in cup

composition, despite frequent mate switching, supports our conclusion that the female

constructs some aspects of the nest independently of the male. Box identity could potentially

be a proxy of the male identity, as cavities are controlled by individual males on their own

territories; however, males often control many boxes within the same territory and the rate of

movement between these boxes is unknown. Therefore, we are unable to split apart the identity

of the male from the identity of the box in this study. Males begin bringing sticks to the cavity

before acquiring a mate (Johnson 2020), and so if the box identity is a reliable proxy for male

identity, then the male contribution to building could help explain the observed variation in the

stick platform, but more research is needed.

In most cases, we only measured the construction attempts of individuals twice. House

Wrens only produce two to three clutches a season, so we are limited by this number and the

likelihood the female renested within our nest boxes instead of natural cavities. However, Bell

et al. (2009) found no significant effect of the number of observations per individual on

repeatability estimates. Instead, the number of observations appears to have a greater effect on

the range of uncertainty in the estimate. We provided this range of uncertainty, as a confidence

interval, and required the lower end to remain higher than zero for significance. Furthermore,

all of our estimates are calculated from a sample size ($N = 24$ for female identity; $N = 26$ for

box identity) larger than the median sample size ($N = 22$) of all studies examined by Bell et al.

(2009), and so we do not believe that our sample size is a major limitation.

Conclusions

The repeatability of construction behaviors is greatly understudied, particularly in

free-living populations. We examined the Northern House Wren, a species with incredible

variation in nest architecture (See Figure S1), to estimate the amount of variability in

construction behavior explained by the identity of the female builder and the box identity and

its dimensions. We found female northern House Wrens to be moderately to highly repeatable

in the composition of the nest cup, but this behavior was unrelated to female body condition or

nestling survival. The general dimensions of the nest, as well as the stick platform, correlated

with the dimensions of the box, and these measurements were highly repeatable for box

identity. These results support previous findings that House Wrens modify aspects of the nest

to compensate for vulnerabilities in the cavity. We hypothesize that the observed variability in

nest architecture is due to the nest architecture being under minimal selective pressure, thus

allowing the diversity of female building preferences to have a greater effect on the expression

of the phenotype. Instead, choosing a protective cavity is likely to be more important for the

builder's fitness, with the nest construction process being relatively unimportant for mate

selection or nestling survival. However, much of House Wren nest construction still remains

unclear, including the contribution of the male in constructing the stick platform. Extended

phenotypes are complex, requiring an examination of both the environment *and* the builder.

The cause of variation in House Wren nest morphology has been largely unexplained due to a

narrow focus on potential external factors. Studying the repeatability of extended phenotypes

across taxa has provided new insights into the construction behaviors of animals and provides

an important method for understanding the causes and consequences of individual preferences.

Data Availability Statement

The data and code that support these findings are available in the Mendeley Data

Repository at doi: 10.17632/fnhr5mm99p.1.

REFERENCES

Adreani, N. M., Valcu, M., Scientists, C., & Mentesana, L. (2022). Asymmetric architecture is non-random and repeatable in a bird's nests. *Current Biology*, *32*(9), R412–R413. https://doi.org/10.1016/j.cub.2022.03.075

Alworth, T. (1996). An Experimental Test of the Function of Sticks in the Nests of House Wrens. *The Condor*, *98*(4), 841–844. https://doi.org/10.2307/1369866

Alworth, T., & Scheiber, I. B. R. (2000). NEST BUILDING IN HOUSE WRENS (TROGLODYTES AEDON): A REEXAMINATION OF MALE AND FEMALE ROLES. *Journal of Field Ornithology*, *71*(3), 409–414. https://doi.org/10.1648/0273-8570-71.3.409

Bailey, I., Morgan, K., Oschadleus, D., DeRuiter, S., Meddle, S., & Healy, S. (2015). Nest-building males trade-off material collection costs with territory value. *The Emu: Official Organ of the Australasian Ornithologists' Union*, *116*. https://doi.org/10.1071/MU15022

Baldwin, S. P. (1921). The Marriage Relations of the House Wren (Troglodytes A. Aedon). *The Auk*, *38*(2), 237–244. https://doi.org/10.2307/4073886

Barber, I., Nairn, D., & Huntingford, F. A. (2001). Nests as ornaments: Revealing construction by male sticklebacks. *Behavioral Ecology*, *12*(4), 390–396. https://doi.org/10.1093/beheco/12.4.390

Beebe, W., & Beebe, W. (1953). A contribution to the life history of the euchromid moth, Aethria carnicauda Butler. *Zoologica : Scientific Contributions of the New York Zoological Society*, *38*(13), 155--160. https://doi.org/10.5962/p.203451

Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: A meta-analysis. *Animal Behaviour*, *77*(4), 771–783.

Biddle, L. E., Broughton, R. E., Goodman, A. M., & Deeming, D. C. (2018). Composition of Bird Nests is a Species-Specific Characteristic. *Avian Biology Research*, *11*(2), 132–153. https://doi.org/10.3184/175815618X15222318755467

Blamires, S. J., Lee, Y.-H., Chang, C.-M., Lin, I.-T., Chen, J.-A., Lin, T.-Y., & Tso, I.-M. (2010). Multiple structures interactively influence prey capture efficiency in spider orb webs. *Animal Behaviour*, *80*(6), 947–953. https://doi.org/10.1016/j.anbehav.2010.09.011

Bohn, S. J., Webber, Q. M. R., Florko, K. R. N., Paslawski, K. R., Peterson, A. M., Piche, J. E., Menzies, A. K., & Willis, C. K. R. (2017). Personality predicts ectoparasite abundance in an asocial sciurid. *Ethology*, *123*(10), 761–771. https://doi.org/10.1111/eth.12651

Borgia, G. (1985). Bower quality, number of decorations and mating success of male satin bowerbirds (Ptilonorhynchus violaceus): An experimental analysis. *Animal Behaviour*, *33*(1), 266–271. https://doi.org/10.1016/S0003-3472(85)80140-8

Brown, W., Alexander, A., Alexander, D., Zuefle, M., & Underwood, T. (2024). Estimating Ages of House Wren Nestlings Based on Body Mass, Wing Chord Length, and Feather Tract Development Patterns. *North American Bird Bander*, *36*(3). https://digitalcommons.usf.edu/nabb/vol36/iss3/1

Cerezo, A. T., & Deeming, D. C. (2016). Nest Insulation in the European Robin (Erithacus Rubecula): Effects of Latitude and Construction Materials. *Avian Biology Research*, *9*(2), 96–102. https://doi.org/10.3184/175815516X14551263654405

Clark, A. D., Deffner, D., Laland, K., Odling-Smee, J., & Endler, J. (2020). Niche Construction Affects the Variability and Strength of Natural Selection. *The American Naturalist*, *195*(1), 16–30. https://doi.org/10.1086/706196

Colombo, S., Newman, K. D., Langmore, N. E., Taylor, C. J., & Medina, I. (2024). The role of climatic variables on nest evolution in tanagers. *Ecology and Evolution*, *14*, e11168. https://doi.org/10.1002/ece3.11168

Darwin, C., & Wallace, A. (1858). On the Tendency of Species to form Varieties; and on the Perpetuation of Varieties and Species by Natural Means of Selection. *Journal of the Proceedings of the Linnean Society of London. Zoology*, *3*(9), 45–62. https://doi.org/10.1111/j.1096-3642.1858.tb02500.x

Dawkins, R. (1982). *The Extended Phenotype: The Long Reach of the Gene*. Oxford University Press.

DiRienzo, N., & Aonuma, H. (2018). Plasticity in extended phenotype increases offspring defence despite individual variation in web structure and behaviour. *Animal Behaviour*, *138*, 9–17. https://doi.org/10.1016/j.anbehav.2018.01.022

DiRienzo, N., Johnson, J. C., & Dornhaus, A. (2019). Juvenile social experience generates differences in behavioral variation but not averages. *Behavioral Ecology*, *30*(2), 455–464. https://doi.org/10.1093/beheco/ary185

DiRienzo, N., & Montiglio, P.-O. (2016). Linking consistent individual differences in web structure and behavior in black widow spiders. *Behavioral Ecology*, *27*(5), 1424–1431. https://doi.org/10.1093/beheco/arw048

DiRienzo, N., Schraft, H. A., Montiglio, P. O., Bradley, C. T., & Dornhaus, A. (2020). Foraging behavior and extended phenotype independently affect foraging success in spiders. *Behavioral Ecology*, *31*(5), 1242–1249. https://doi.org/10.1093/beheco/araa080

Drilling, N. E., & Thompson, C. F. (1988). Natal and Breeding Dispersal in House Wrens (Troglodytes aedon). *The Auk*, *105*(3), 480–491.

Eckerle, K. P., & Thompson, C. F. (2005). Addition of arthropod cocoons to House Wren nests is correlated with delayed pairing. *Behavioral Ecology*, *16*(1), 1–7. https://doi.org/10.1093/beheco/arh122

Edwards, S. C., Shoot, T. T., Martin, R. J., Sherry, D. F., & Healy, S. D. (2020). It's not all about temperature: Breeding success also affects nest design. *Behavioral Ecology*, *31*(4), 1065–1072. https://doi.org/10.1093/beheco/araa052

Ellendula, S., Tresa, C., & Uma, D. (2021). Influence of prey availability on web-building in the social spider Stegodyphus sarasinorum (Araneae: Eresidae). *The Journal of Arachnology*, *49*(1), 141–145. https://doi.org/10.1636/JoA-S-19-080

Fisher, D. N., Yeager, J., & Pruitt, J. N. (2019). *Orb-weaving spiders show a correlated syndrome of morphology and web structure in the wild* (p. 767335). bioRxiv. https://doi.org/10.1101/767335

Franklin, D. C. (1995). Helmeted Honeyeaters Build Bulkier Nests in Cold Weather. *The Auk*, *112*(1), 247–248. https://doi.org/10.2307/4088785

Hansell, M. (2000). *Bird Nests and Construction Behaviour*. Cambridge University Press. https://doi.org/10.1017/CBO9781139106788

Hansell, M. H. (2005). *Animal Architecture*. OUP Oxford.

Hauber, M. E., Nagy, J., Sheard, C., Antonson, N. D., Street, S. E., Healy, S. D., Lala, K. N.,
& Mainwaring, M. C. (2024). Nest architecture influences host use by avian brood
parasites and is shaped by coevolutionary dynamics. *Proceedings of the Royal Society
B: Biological Sciences*, *291*(2014), 20231734. https://doi.org/10.1098/rspb.2023.1734

Hayes, J. P., & Jenkins, S. H. (1997). Individual Variation in Mammals. *Journal of
Mammalogy*, *78*(2), 274–293. https://doi.org/10.2307/1382882

Hays, S. C., Cheek, R. G., Mouton, J. C., Sillett, T. S., & Ghalambor, C. K. (2022). Lack of
avian predators is associated with behavioural plasticity in nest construction and height
in an island songbird. *Animal Behaviour*, *194*, 35–42.
https://doi.org/10.1016/j.anbehav.2022.08.017

Hermes, M. G., Somavilla, A., & Garcete-Barrett, B. R. (2013). On the nesting biology of
Pirhosigma Giordani Soika (Hymenoptera, Vespidae, Eumeninae), with special
reference to the use of vegetable matter. *Revista Brasileira de Entomologia*, *57*,
433–436. https://doi.org/10.1590/S0085-56262013005000044

Hölldobler, B., & Wilson, E. O. (1990). *The Ants*. Harvard University Press.

Järvinen, P., & Brommer, J. E. (2020). Nest ornaments and feather composition form an
extended phenotype syndrome in a wild bird. *Behavioral Ecology and Sociobiology*,
74(11), 134. https://doi.org/10.1007/s00265-020-02912-2

Järvinen, P., Kluen, E., & Brommer, J. E. (2017a). Low heritability of nest construction in a
wild bird. *Biology Letters*, *13*(10), 20170246. https://doi.org/10.1098/rsbl.2017.0246

Järvinen, P., Kluen, E., Tiiri, M., & Brommer, J. (2017b). Experimental manipulation of
Blue Tit nest height does not support the thermoregulation hypothesis. *Ornis Fennica*,
94. https://doi.org/10.51812/of.133914

Jelínek, V., Požgayová, M., Honza, M., & Procházka, P. (2016). Nest as an extended phenotype signal of female quality in the great reed warbler. *Journal of Avian Biology*, *47*(3), 428–437. https://doi.org/10.1111/jav.00700

Johnson LS. 2020. House Wren (*Troglodytes aedon*), version 1.0. In: Poole AF, editor. Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA.

Korb, J. (2003). Thermoregulation and ventilation of termite mounds. *Naturwissenschaften*, *90*(5), 212–219. https://doi.org/10.1007/s00114-002-0401-4

Korb, J., & Linsenmair, K. E. (2000). Thermoregulation of termite mounds: What role does ambient temperature and metabolism of the colony play? *Insectes Sociaux*, *47*(4), 357–363. https://doi.org/10.1007/PL00001731

Laland, K. N., Odling-Smee, F. J., & Feldman, M. W. (1999). Evolutionary consequences of niche construction and their implications for ecology. *Proceedings of the National Academy of Sciences of the United States of America*, *96*(18), 10242–10247.

Lamprecht, I., & Schmolz, E. (2004). Thermal investigations of some bird nests. *Thermochimica Acta*, *415(1–2)*, 141–148. https://doi.org/10.1016/j.tca.2003.08.029

LeGrice, R. J., Tezanos-Pinto, G., de Villemereuil, P., Holwell, G. I., & Painting, C. J. (2019). Directional selection on body size but no apparent survival cost to being large in wild New Zealand giraffe weevils. *Evolution*, *73*(4), 762–776. https://doi.org/10.1111/evo.13698

Madden, J. R. (2002). Bower Decorations Attract Females but Provoke Other Male Spotted Bowerbirds: Bower Owners Resolve This Trade-off. *Proceedings: Biological Sciences*, *269*(1498), 1347–1351.

Mainwaring, M., Benskin, C., C.McW.H, & Hartley. (2008). The Weight of Female-Built Nests Correlates with Female but not Male Quality in the Blue Tit Cyanistes caeruleus. *Acta Ornithologica*, *43*, 43–48. https://doi.org/10.3161/000164508X345310

Mainwaring, M. C., Deeming, D. C., Jones, C. I., & Hartley, I. R. (2014b). Adaptive latitudinal variation in Common Blackbird Turdus merula nest characteristics. *Ecology and Evolution*, *4*(6), 851–861. https://doi.org/10.1002/ece3.952

Mainwaring, M. C., Hartley, I. R., Lambrechts, M. M., & Deeming, D. C. (2014a). The design and function of birds' nests. *Ecology and Evolution*, *4*(20), 3909–3928. https://doi.org/10.1002/ece3.1054

McCabe, R. A. (1965). Nest Construction by House Wrens. *The Condor*, *67*(3), 229–234. https://doi.org/10.2307/1365401

McKaye, K. R., Louda, S. M., & Stauffer, J. R. (1990). Bower Size and Male Reproductive Success in a Cichlid Fish Lek. *The American Naturalist*, *135*(5), 597–613.

Medina, I., M. Perez, D., Silva, A. C. A., Cally, J., León, C., Maliet, O., & Quintero, I. (2022). Nest architecture is linked with ecological success in songbirds. *Ecology Letters*, *25*(6), 1365–1375. https://doi.org/10.1111/ele.13998

Mennerat, A., Perret, P., & Lambrechts, M. M. (2009). Local Individual Preferences for Nest Materials in a Passerine Bird. *PLOS ONE*, *4*(4), e5104. https://doi.org/10.1371/journal.pone.0005104

Merilä, J., & Wiggins, D. A. (1997). Mass Loss in Breeding Blue Tits: The Role of Energetic Stress. *Journal of Animal Ecology*, *66*(4), 452–460. https://doi.org/10.2307/5940

Mertens, J. A. (1977). Thermal conditions for successful breeding in great tits (Parus major L.). *Oecologia*, *28(1)*, 1–29. https://doi.org/10.1007/bf00346834

Miyashita, T., & Shinkai, A. (1995). Design and Prey Capture Ability of Webs of the Spiders Nephila clavata and Argiope bruennichii. *Acta Arachnologica*, *44*(1), 3–10. https://doi.org/10.2476/asjaa.44.3

Møller, A. P. (1990). Nest Predation Selects for Small Nest Size in the Blackbird. *Oikos*, *57*(2), 237–240. https://doi.org/10.2307/3565945

Møller, A. P. (2006). Rapid change in nest size of a bird related to change in a secondary sexual character. *Behavioral Ecology*, *17*(1), 108–116. https://doi.org/10.1093/beheco/arj003

Muth, F., & Healy, S. D. (2011). The role of adult experience in nest building in the zebra finch, *Taeniopygia guttata*. *Animal Behaviour*, *82*(2), 185–189. https://doi.org/10.1016/j.anbehav.2011.04.021

Paclík, M., Misík, J., & Weidinger, K. (2009). Nest predation and nest defence in European and North American woodpeckers: A review. *Annales Zoologici Fennici*, *46*(5), 361–379.

Parthasarathy, B., Bouchard, M., & Schneider, J. M. (2022). Extended phenotypes can underlie trade-offs: A case of social spiders. *Die Naturwissenschaften*, *109*(6), 51. https://doi.org/10.1007/s00114-022-01826-5

Perez, D. M., Gardner, J. L., & Medina, I. (2020). Climate as an Evolutionary Driver of Nest Morphology in Birds: A Review. *Frontiers in Ecology and Evolution*, *8*. https://www.frontiersin.org/articles/10.3389/fevo.2020.566018

Pribil, S., & Picman, J. (1997). Parasitism of House Wren nests by Brown-headed Cowbirds: Why is it so rare? *Canadian Journal of Zoology*, *75*(2), 302–307. https://doi.org/10.1139/z97-038

Pyle, P. (1997). Identification Guide To North American Birds: A Compendium Of Information On Identifying, Ageing, And Sexing "Near-Passerines" And Passerines In The Hand. *Books by Alumni*. https://works.swarthmore.edu/alum-books/4499

Rosell, F., & Campbell-Palmer, R. (2022). Habitat use and constructions. In F. Rosell & R. Campbell-Palmer (Eds.), *Beavers: Ecology, Behaviour, Conservation, and Management* (p. 0). Oxford University Press. https://doi.org/10.1093/oso/9780198835042.003.0004

Rushbrook, B. J., Dingemanse, N. J., & Barber, I. (2008). Repeatability in nest construction by male three-spined sticklebacks. *Animal Behaviour*, *75*(2), 547–553. https://doi.org/10.1016/j.anbehav.2007.06.011

Silva, N. J., Benoit, F., Elliott, A., Rault, C., Colençon, P., Covas, R., & Doutrelant, C. (2024). Sex and age differences in the preference for materials for the communal nests of sociable weavers Philetairus socius. *Ethology*, *00*, e13472. https://doi.org/10.1111/eth.13472

Stanback, M. T., Stanbeck, M. T., Bartholomew, J. E., Bergner, L. M., Cline, E. L., Helms, P. I., McGovern, P. G., Millican, D. M., & Roth, J. E. (2013). House Wrens Alter Nest Architecture to Compensate for Cavity Vulnerability. *The Wilson Journal of Ornithology*, *125*(1), 174–178.

Stoffel, M., Schielzeth, H., & Nakagawa, S. (2017). rptR: Repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *8*. https://doi.org/10.1111/2041-210X.12797

Sugiura, S. (2016). Bagworm bags as portable armour against invertebrate predators. *PeerJ*, *4*, e1686. https://doi.org/10.7717/peerj.1686

Thomas, J., Handasyde, K., Parrott, M., & Temple-Smith, P. (2018). The platypus nest: Burrow structure and nesting behaviour in captivity. *Australian Journal of Zoology*, *65*. https://doi.org/10.1071/ZO18007

Ueda, K. (1985). *Nest Building Activity and Characteristics of the Courtship Nest of the Fan-tailed Warbler* Cisticola juncidis. https://www.jstage.jst.go.jp/article/jjo1915/34/2-3/34_2-3_23/_article

van Rijssel, J. C., Moser, F. N., Frei, D., & Seehausen, O. (2018). Prevalence of disruptive selection predicts extent of species differentiation in Lake Victoria cichlids. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1871), 20172630. https://doi.org/10.1098/rspb.2017.2630

Wallace, A. R. (1865). I. On the Phenomena of Variation and Geographical Distribution as illustrated by the Papilionidæ of the Malayan Region. *Transactions of the Linnean Society of London*, *25*(1), 1–71. https://doi.org/10.1111/j.1096-3642.1865.tb00178.x

Walsh, P. T., Hansell, M., Borello, W. D., & Healy, S. D. (2010). Repeatability of nest morphology in African weaver birds. *Biology Letters*, *6*(2), 149–151. https://doi.org/10.1098/rsbl.2009.0664

Whittaker, B. A., Nolet-Mulholland, L., Nevoit, A., Yun, D., Lambert, C. T., Blunk, S. C., & Guillette, L. M. (2023). Zebra finches have style: Nest morphology is repeatable and associated with experience. *IScience*, *26*(11), 108194. https://doi.org/10.1016/j.isci.2023.108194

Ziebis, W., Forster, S., Huettel, M., & Jørgensen, B. B. (1996). Complex burrows of the mud

shrimp Callianassa truncata and their geochemical impact in the sea bed. *Nature*,

382(6592), Article 6592. https://doi.org/10.1038/382619a0

SUPPLEMENTARY

Table S1 - Individuals examined for Female Identity repeatable traits and Box Identity traits.

Females with single IDs were unbanded individuals.

Female Repeatability			Box Repeatability		
Female ID	Clutch	Box ID	Box ID	Clutch	Female ID
288064919	1	3	1	1	288067191
288064919	2	2	1	2	288067191
288067191	1	1	1	3	288067191
288067191	2	1	3	1	288064919
288067191	3	1	3	2	294063102
288064901	1	6	4	1	288064918
288064901	2	4	4	2	288064901
294063102	1	8	9	1	288064920
294063102	2	3	9	2	294063222
288064903	1	19	13	1	288064917
288064903	2	18	13	2	294063155
288067184	1	25	17	1	288064902
288067184	2	24	17	2	294063262
288064906	1	28	19	1	288064903

288064906	2	29	19	2	294063200
288064906	3	29	20	1	288064905
288064947	1	33	20	2	294063294
288064947	2	30	29	1	288064906
288064905	1	20	29	2	288064906
288064905	2	21	30	1	1
288064913	1	46	30	2	288064947
288064913	2	46	33	1	288064947
288064913	3	54	33	2	294063157
288064915	1	41	38	1	294063259
288064915	2	41	38	2	294063453
294063218	1	47	39	1	288064923
294063218	2	49	39	2	288064991
288064991	1	40	41	1	288064915
288064991	2	39	41	2	288064915
288064923	1	39	43	1	288064912
288064923	2	47	43	2	294063461
288064997	1	56	44	1	288064911
288064997	2	56	44	2	294063173
288064939	1	54	46	1	288064913
288064939	2	55	46	2	288064913
288067301	1	80	47	1	4
288067301	2	79	47	2	294063218

288064925	1	68	47	3	288064923
288064925	2	67	48	1	5
288067007	1	70	48	2	288064996
288067007	2	71	49	1	6
288067007	3	65	49	2	294063135
288064916	1	63	49	3	294063218
288064916	2	63	54	1	288064939
288064916	3	62	54	2	288064913
294063101	1	61	56	1	288064997
294063101	2	60	56	2	288064997
294063230	1	76	63	1	288064916
294063230	2	75	63	2	288064916
294063219	1	77	65	1	288067007
294063219	2	66	65	2	7
288064910	1	58	66	1	288064924
288064910	2	59	66	2	294063219
			70	1	288067007
			70	2	294063180

875 *Table S2 - P-values supporting the lack of significant correlations between nest characteristics*

876 *and the nest age.*

	Grass Count	Total Feather	Wing	Body	Age
Nest Height	0.95	0.94	0.69	0.81	0.12

Nest Volume	0.71	0.89	0.49	0.93	0.14
Nest Berm	0.76	0.93	0.37	0.83	0.78
Stick	0.19	0.35	0.32	0.44	*0.052*

Table S3 - Relationships between the dimensions and components of the nest and various

proxies of fitness: Female residual - defined by the correlation between the tarsus length and

the body mass, Fledge or Fail - the simply outcome of the nest, Eggs laid - the number of eggs

originally laid by the nesting female at the completion of the cup, and Offspring fledged - the

estimated number of offspring that fledged the nest (only successful nests examined). Date - the

relationship between the dimensions and components of the nest and the date at which the nest

was measured. †*Specified material measured as a simple yes/no presence.* * $r^2 = 0.09$, P =

0.001 with removal of outlier date 144

	Female residual	Fledge or Fail	Eggs laid	Offspring fledged	Date
Ratio	t = 1.64, P = 0.11	z = 0.10, P = 0.92	t = -0.77, P = 0.45	t = 0.73, P = 0.47	r^2 = -0.006, P = 0.54
Nest Height	t = 0.90, P = 0.37	z = -0.12, P = 0.91	t = -0.81, P = 0.42	t = 0.54, P = 0.59	r^2 = -0.006, P = 0.54
Nest Volume	t = 0.46, P = 0.65	z = -0.13, P = 0.89	t = -0.63, P = 0.53	t = -0.06, P = 0.95	r^2 = -0.007, P = 0.62
Nest Berm	t = -0.64, P = 0.52	z = 0.11, P = 0.91	t = 0.08, P = 0.94	t = -0.35, P = 0.73	r^2 = -0.008, P = 0.74
Stick	t = -0.39, P = 0.70	z = -0.72, P = 0.47	t = -0.03, P = 0.97	t = -0.66, P = 0.51	r^2 = -0.009, P = 0.72

Grass	t = 0.67, P = 0.50	z = -1.56, P = 0.11	t = 0.89, P = 0.37	t = -1.04, P = 0.30	r^2 = -0.01, P = 0.88
Total Feather	t = 0.03, P = 0.97	z = 0.31, P = 0.76	t = 0.13, P = 0.90	t = -1.45, P = 0.15	r^2 = 0.01, P = 0.15
Wing Rate	t = -0.12, P = 0.90	z = 1.32, P = 0.13	t = 0.28, P = 0.78	t = 0.12, P = 0.91	r^2 = 0.02, P = 0.08
Pennaceous	t = -1.07, P = 0.29	z = 1.58, P = 0.08	t = -0.01, P = 0.99	t = -1.91, P = 0.06	r^2 = 0.02, P = 0.07*
Plumulaceous	t = 0.43, P = 0.67	z = -0.23, P = 0.82	t = -0.16, P = 0.88	t = -0.95, P = 0.35	r^2 = 0.002, P = 0.27
Spider Sac Rate	t = -0.59, P = 0.56	z = 0.41, P = 0.68	t = -1.15, P = 0.25	t = -0.26, P = 0.80	t = 4.49, P < 0.001
† Snake Skin	χ^2_1 = 0.08, P = 0.77	z = -0.81, P = 0.42	t = 1.20, P = 0.23	t = 1.27, P = 0.20	z = -1.40, P = 0.16
† Plastic	χ^2_1 = 0.002, P = 0.96	z = -1.57, P = 0.11	t = 1.74, P = 0.08	t = 0.09, P = 0.93	z = 0.02, P = 0.99

Figure S1 - Images of complete nests constructed in a nesting box in 2022 showing that A)

nests are sometimes constructed to the point where they will cover the primary entrance of the

nesting box, while B) some nests are constructed only a few centimeters tall. Arrow 1 points to

the top of the primary entrance; Arrow 2 points to the top of the nest. In cases like that shown

in nest A, the wrens will use the gap entrance (See Figure 1) as their main entrance instead.

Figure S2 - Relationships between different nest attributes and building materials. The width of

the nest berm was positively correlated with the total volume (A; $r^2 = 0.09$, P < 0.001) and stick

count (B; $r^2 = 0.04$, P = 0.02) of the nest. The stick count further highly positively correlated

with the height of the nest (C; r² = 0.37, P < 0.001) and the total volume of the nest (D; r² =

0.41, P < 0.001). The grass count positively correlated with the total count of feathers (E; r² =

0.03, P = 0.04) and pennaceous feather count (F; r² = 0.06, P = 0.005). Gray shading

encompasses the 95% confidence interval for the line of best fit.

Figure S3 - Significant relationships between nest characteristics and the date of nest

measurement (i.e., completion of egg laying). Nests measured earlier in the season were more

likely to fail than nests measured later in the season (A; t = 2.04, P = 0.04), but also fledged

fewer offspring (B; t = -5.89, P = 0.002). The pennaceous feather count positively correlated

with the measurement date after the removal of the outlier on DOY 144 (C; r² = 0.02, P =

0.001). Nests measured later in the season also contained more spider sac material than those

measured earlier in the season (D; t = 4.49, *P < 0.001). Gray shading in C encompasses the*

95% confidence interval for the line of best fit.

Figure S4 - Relationship between three nest materials and fecal material with the age of the

nest at collection. Pennaceous feather count positively correlated with the nest age (A; r^2 *=*

0.04, P = 0.02). Although not statistically significant, the stick count (C; r^2 *= 0.03, P = 0.052)*

and the grass count (D; r^2 *= 0.03, P = 0.054) negatively correlated with the age of the nest.*

Older nests contained more fecal material than younger nests (B; t = 5.24, *P < 0.001). Gray*

shading in A, C, and D encompasses the 95% confidence interval for the line of best fit.