Repeatability of an extended phenotype: causes and consequences of nest variation in a

free-living songbird

Running Title: Repeatability of nest construction

Chandler E. G. Carr[1,*], Zoë M. Swanson[1], and Dustin G. Reichard[1]

[1]Ohio Wesleyan University, Department of Biological Sciences, 61 S Sandusky St, Delaware,

OH 43015

*Corresponding author: Chandler E. G. Carr

Keywords: Extended Phenotype, Niche Construction Hypothesis, Repeatability, Individual

Preferences, Nest Building, House Wren

HIGHLIGHTS

●   Repeatability of construction behaviours in the wild is understudied

●   House wrens are highly variable in their nest morphology

●   Female house wrens are highly repeatable in cup composition

●   House wren nests are unrelated to nestling survival or parental condition

●   Differences in individual preferences likely explains nest variation

ABSTRACT

Construction behaviour is an aspect of the extended phenotype that allows organisms to

build structures that alter their environments in potentially beneficial ways. Although

individuals vary in the expression of this extended phenotype (e.g., structure morphology), the

repeatability of construction behaviour remains understudied, especially among free-living

populations. Many oviparous taxa construct nests, making them of particular interest because

variation in nest architecture may directly affect fitness. Using a free-living, cavity-nesting

songbird, the northern house wren (*Troglodytes aedon aedon*), as our model, we estimated the

contribution of the primary builder (the female) to nest variability by measuring the

repeatability of nest morphology between successive clutches. We further examined whether

nest morphology was related to the dimensions of the nesting cavity, breeding date, or nest

success. We found the composition of the cup lining to be a highly repeatable behaviour for the

nesting female, although the size and composition of the structural platform appeared more

related to the dimensions of the cavity. Despite the expectation that construction effort would

show a seasonal decline, similar to clutch size in this species, nest morphology remained

incredibly variable throughout the breeding season and was unrelated to the survival of the

offspring. Our study suggests that variation in construction behaviours is a product of multiple

factors including the preferences of the builder and physical constraints. The absence of any

clear links between construction behaviour and fitness indicates that nest morphology is not

under strong selection. As a result, diverse female building preferences may explain the

extreme among-individual variation in nest structure in this species.

INTRODUCTION

Alfred Russel Wallace (1865) once stated: "…so great is the tendency to vary that it is

difficult to find two individuals exactly alike." Phenotypic variation is both ubiquitous and a

fundamental requirement for evolution (Darwin & Wallace 1858). As such, a considerable

amount of research has been devoted to understanding the causes and consequences of among-

individual variation. The interaction between an organism's phenotype and the environment

determines its success, and selection can push populations toward one or more optimal

phenotypes (van Rijssel et al. 2018, LeGrice et al. 2019). However, the relationship between

the phenotype and the environment can be bi-directional, allowing individuals to reshape their

environments in ways that affect their fitness (i.e., "niche construction"; Laland et al. 1999).

This bi-directional interaction can bias the direction of selection in favour of the individual,

and often leads to reduced spatial and temporal variation in phenotypes (Clark et al. 2020).

Construction behaviour is a unique behavioural phenotype where the product of the

construction acts as both an extension of the builder's phenotype (i.e., "extended phenotype";

Dawkins 1982) and an artificial environment more beneficial for the builder. Construction

behaviours may enhance the builder's fitness by providing protection from predators (Beebe

1953, Hermes, 2013, Sugiura 2016) and environmental exposure (Zeibis et al. 1996, Korb &

Linsenmair 2000, Korb 2003, Rosell & Campbell-Palmer 2022), as well as increasing access to

resources such as food (Hölldobler & Wilson 1990, Miyashita & Shinkai 1995, Blamires et al.

2010) and mates (Borgia 1985, McKaye et al. 1990). As a result, construction behaviour has

become taxonomically widespread across both vertebrate and invertebrate species (Hansell

2005). Eusocial insects build colonial shelters (e.g. termite mounds), some mammals dig

burrows, most spiders spin webbed traps, and certain species of fishes and birds construct

bowers. However, despite the clear potential for construction behaviour to benefit the fitness of

an individual, trade-offs often occur between investment in the construction process and other

behaviours that affect fitness, such as foraging, which can result in negative effects on both the

builder (Madden 2002, Bailey et al. 2015) and the structure (Ellendula et al. 2021,

Parthasarathy et al. 2022).

It is well established that the products of construction behaviours exhibit among-

individual variation, but the extent and cause of variation *within* an individual's construction

behaviour is less clear (DiRienzo & Aonuma 2018). For example, an individual's construction

behaviour might be static, plastic, or some combination of the two as its experience and

environment changes. Many organisms build multiple versions of the same structure

throughout their lifetimes, which provides an opportunity to evaluate the consistency of this

behaviour. Repeatability estimates, calculated as intra-class correlation coefficients, can be

used to assess how much of the variation observed among individuals can be explained by

variation within individuals (Hayes & Jenkins 1997). When among-individual variation is high

and within-individual variation is low, then the repeatability estimate will be high and that trait

will be considered repeatable. Repeatability estimation can be used to understand the extent to

which phenotypic traits are the product of static characteristics, such as heritability ("upper

level of heritability"; Bell et al. 2009, Møller 2006) and preferences (Mennerat et al. 2009,

Järvinen & Brommer 2020), or plastic characteristics, such as experience (DiRienzo et al.

2019, Whittaker et al. 2023). For this reason, the repeatability of a large variety of behaviours

have been examined across taxa (Bell et al. 2009), but there are few studies examining the

repeatability of construction behaviours. Most of these studies have focused on spider web

structure (Fisher et al. 2019, DiRienzo et al. 2020, DiRienzo & Montiglio 2016) or nest

morphology (Whittaker et al. 2023, Muth & Healy 2011, Rushbrook et al. 2008), but few have

examined the behaviour exclusively in the wild (Møller 2006, Järvinen et al. 2017).

Nest building is a very common construction behaviour among oviparous taxa that serves

the primary function of providing a safe, hospitable space for eggs or altricial young to develop

(Mainwaring et al. 2014a). With few exceptions (e.g., brood parasites), bird species produce an

incredible array of nests that can be structurally complex or simple, but must contain at least

one of four zones: an outer nest layer, an attachment to the environment, a structural layer, and

a lining (Hansell 2000). Zones are often discernible from one another by the materials used,

with zone composition being largely species specific (Biddle et al. 2018). Nests can be built by

the male, female, or both with some division of labour by zone (Hansell 2000). The

morphology of the nest, structure, size, and composition, can be important for providing the

same benefits of niche construction observed for other structures by protecting the young from

predators (Hays et al. 2022, Møller 1990) and environment exposure (Medina et al. 2022,

Perez et al. 2020, Cerezo & Deeming 2016, Mainwaring et al. 2014b, Franklin 1995), as well

as potentially playing a role in mate attraction for some species (Jelínek et al. 2016,

Mainwaring et al. 2008, Ueda 1985).

Here, we investigated the causes and consequences of the variation and repeatability of

northern house wren (*Troglodytes aedon aedon*) nest morphology and composition. House

wrens are secondary cavity nesters that raise two to three clutches a season, often reusing the

same nest throughout (Johnson 2020). They will, however, build a new nest when the previous

nest is stolen, destroyed, or experimentally removed (Alworth 1996), making them an ideal

model for the examination of repeatability in construction behaviour. House wren nests

arguably contain 2 to 3 zones: sometimes an outer layer consisting of spider egg sac material, a

structural zone made of various types of sticks, henceforth referred to as the "stick platform,"

and a cup defined as a simple depression, at the top of the stick platform, lined with grasses

and feathers (McCabe 1965). Although the layout and composition of the zones are standard,

house wren nests are highly variable in their morphology. For example, in some nests the stick

platform may be built so large that it covers the entrance hole of the box (McCabe 1965).

While establishing a territory, unmated males place a variable number of sticks (range: <10 to

>400; Johnson 2020) into potential nesting cavities. After pairing, the female completes the

stick platform (Alworth & Scheiber 2000, Johnson 2020) and creates a lined cup either before

or during egg laying. Considering the collective effort of the female in both completing the

stick platform and constructing the nest cup during the construction phase, we defined the

female as the primary nest builder in our study.

Despite the benefits of nest construction observed among other bird species (Mainwaring

et al. 2014), house wren nest architecture appears to be unrelated to the mate selection process,

the survivability of offspring (Alworth 1996, Eckerle & Thompson 2005), or the limiting of

brood parasitism (Pribil & Picman 1997). Thus, the available evidence suggests that nest

architecture in this species has limited effects on fitness, and so it remains unclear why house

wren nests exhibit such extreme variation among individuals. The absence of selective pressure

on nest construction could be one explanation for the amount of variability observed. Some

research suggests that constraints imposed by the cavity dimensions and the accessibility of

building materials may also drive variation in nest architecture (Stanback et al., 2013).

However, no study to date has explored the role of female building preferences as an

explanation for the variation.

We sampled house wren nests across an entire breeding season to assess the contribution

of the female builder, the box cavity, and seasonal timing to nest architecture. We also

examined how variation in nest dimensions and composition correlated with offspring survival.

Unlike most construction behaviour research that often relies on coarse measures of structure

like nest weight (Mainwaring et al. 2008, Walsh et al. 2010, Adreani et al. 2022), we used a

finer approach of directly quantifying the individual building materials (e.g., exact count of

sticks) used in nest construction. We predicted that nests would have high variability among

individuals but that there would be low within-individual variability, specifically that the

female identity would be the best predictor of nest dimension and composition as measured by

a repeatability estimate. Based on previous data, we predicted that the architecture of the nest

would not relate to the survivability of offspring, but may relate to the dimensions of the cavity

in which it was built.

METHODS

Study System

We monitored northern house wren (*Troglodytes aedon aedon*) breeding behaviour in

160 nest boxes across six sites in Delaware and Union counties in Ohio, U.S.A., between May

9 and August 25 of 2022. Of those 160 boxes, 79 produced at least one nest with a complete

clutch, which was the threshold for inclusion in our study. We measured nest box dimensions

with digital callipers (nearest 0.001 mm). Nest boxes were rectangular wooden cuboids with

internal dimensions of approximately 110 x 115 x 195 mm (Figure 1, see Table 1 for variation

in box dimensions). Each box possessed a circular entrance hole with an average diameter of

34 mm and depth of 46 mm, drilled an average of 119 mm from the bottom of the box. Most (*n*

= 73/79; 92.4%) boxes also had a rectangular gap separating the top of the front panel from the

roof that allowed the panel to swing open. This gap was the same width as the box with a

height ranging from 4 mm to 30 mm. Some wrens used this gap as an entrance, particularly

when their nest obstructed the primary entrance hole. The variation in box dimensions were

due to imperfect construction and the preferences of landowners for slightly different box

designs. All boxes were suspended about 1.75 metres off the ground by a greased metal

conduit pole to limit predation.

We monitored 108 total nests (some boxes were used multiple times), but were unable to

visit boxes daily, so exact ageing of nests was not always possible. However, house wrens have

predictable rates of egg laying and incubation time, so we were able to estimate first egg dates

by counting eggs and incubation start dates by backdating from hatch date (Johnson 2020). In

cases where we were not present for hatch day, nestling ages and estimated hatch date were

evaluated by measuring nestling weights, wing chord length, and feather tract development as

described by Brown et al. (2011).

As soon as we discovered a complete clutch, the female was caught in a mist net and

banded with a USFWS aluminium leg band and a unique combination of plastic coloured

bands for future identification. It was essential to capture and band females to sample repeated

nesting attempts, but we chose not to capture males to minimise further disrupting both our

experiment and a separate, unrelated study. The female's weight (nearest 0.1 g; Pesola spring

scale), tarsus length (nearest 1 mm; ruler), and wing length (nearest 1 mm; ruler) were also

measured. While the female was processed, we measured the height of the nest and the width

of the "berm" (Stanback et al. 2013), which was defined as the front wall of the cup directly

against the entrance hole (see Figure 1), using a dial calliper (nearest 0.1 mm). On day eight of

nestling life, we collected measurements and banded each nestling as described above for adult

females. We continued to monitor nests until the clutch fledged or failed, at which point the

nest was collected in a large and resealable plastic bag for dissection in the lab and to prevent

the nest from being reused or built upon for a successive clutch.

Figure 1 - Model of the nest box and nest (in orange). Both the front (left) and side (right) of

the box are shown. A: height of the rectangular gap, B: diameter of the entrance hole, C:

distance from the bottom of the box to the bottom of the entrance hole, D: internal length of the

box and length of the nest, E: depth of the entrance hole, F: width of the berm, G: internal

width of the box and width of the nest, H: height of the nest, I: internal height of the box.

193   *Table 1 - Average, standard deviation, and range of the dimensions of the nest boxes used in*

194   *this study. See Figure 1 for a model of an average nest box.*

	Average (mm)	Standard Deviation (mm)	Range (mm)
Internal Box Height	195.19	14.92	[172.20, 219.90]
Internal Box Volume	2462.20	434.52	[1812.99, 4133.73]
Entrance Diameter	33.60	5.22	[27.90, 44.65]
Entrance Depth	44.11	6.78	[25.00, 56.40]
Height of Entrance	118.94	21.19	[83.10, 151.60]
Gap Height (when present)	14.82	6.11	[4.30, 29.55]

Nest Analysis

After collection, we placed nests in a -20 C freezer for 24 - 48 hours to kill any

arthropods. For dissection, nests were placed on a large sheet of white paper and carefully

pulled apart to quantify nest materials and characteristics. Most nests were dissected by the

lead author, but two assistants were also trained for dissection. We made exact counts of

individual sticks, grasses, plumulaceous, and pennaceous feathers. An arbitrary threshold of 4

cm in length was chosen to separate individual sticks and grasses from other debris. Down,

contour, and semiplume feathers were categorised as Plumaceous, while wing and tail feathers

were categorised as Pennaceous. We used an ordinal scale of 0 to 3 to quantify the amount of

faecal material and spider egg sacs present within each nest. Finally, we noted, as a binary yes

or no, the presence of snake skin and anthropogenic objects (e.g., plastic).

Statistical analysis

Although we sampled 108 nests in total, a small number had to be excluded from our

final analysis. One nest was removed during dissection because it contained signs of reuse (i.e.,

two individual grass layers separated by a thick stick layer). Two nests were not collected from

the field due to reuse or extensive nestling decomposition after predation, and so were unable

to be dissected. Due to data loss, the dissection measurements of two additional nests are

unaccounted for. Our final sample size was 103 nests.

R (version 4.2.3) was used for all analyses. We first assessed whether the observed

variation in nest measurements was normally distributed using separate Shapiro-Wilk tests. To

test whether nest measurements were repeatable within females, we calculated repeatability

estimates (packages: rptR, version 0.9.22) with the nest measurement as the dependent

variable, and the identity of the nesting female as the grouping factor. Parametric bootstrapping

and permutation, at 10 000 iterations each, were performed to produce confidence intervals and

adjusted alpha values for the repeatability estimates, respectively. Significant repeatability

estimates were indicated by 1) confidence intervals not overlapping zero (DiRienzo &

Aonuma, 2018) and 2) an adjusted alpha value of 0.05. Where nest measurement was binomial,

the link-scale approximation (Stoffel et al. 2017) is reported. We used the same approach to

test whether the box identity predicted nest measurements. In these models, the nest

measurement was the dependent variable, and the identity of the box was the grouping factor.

Although there was some overlap in sampling between models with the same female nesting

more than once in the same box, the sample size (5/24; 21% of females reused the same box

and 6/27; 22% of boxes were used by the same female) was too small to provide a reliable

estimate of the effect of these repeated samples on nest measurements. Thus, we chose to avoid

a more complex model that included both box and female identity.

Linear regressions (Base R Package) were used to test for a relationship between nest

measurements and box proportions, date of measurement, and age of the nest, which was

defined as the number of days between the first egg and the nest collection date. We used

generalised linear models (Base R Package) to analyse the effect of nest measurements on the

outcome of the clutch. These models assumed a binomial distribution with nest outcome

(fledge/fail) as the dependent variable. Linear models were used to analyse the effect of nest

measurement on clutch size and fledge count, assuming a gaussian distribution. Both the

generalised linear models and linear models had the nest measurement as the fixed effect, and

the date of nest measurement as a covariate to control for seasonal variation in reproductive

output.

Ethical Note

All procedures were reviewed and approved by the Ohio Wesleyan University

Institutional Animal Care and Use Committee (Protocol Number: 05-2021-06), U.S. Federal

Bird Banding Lab (Permit Number: 24035), and State of Ohio Department of Natural

Resources (Permit Numbers: 23-009, 23-010).

RESULTS

Observed Variability in Nest Architecture

House wren nests were extremely variable (Figure 2). The ratio of the nest height to

internal box height ranged from 0.254 to 0.902 (Figure 2A; $\underline{x}$ ±SD: 0.59 ± 0.144). Similarly,

the volume of the nest ranges from 655.71 mm³ to 2776.96 mm³ (Figure 2B; $\underline{x}$ ±SD: 1437.44 ±

409.29). Sticks were the primary contributor to the height and volume of the nest, and we

found that stick count exhibited the same variability, ranging from 170 to 1491 sticks in a

single nest (Figure 2C; $\underline{x}$ ±SD: 676.14 ± 25.01). The width of the nest berm ranged from 7.00

mm to 77.45 mm (Figure 2D; $\underline{x}$ ±SD: 39.62 ± 1.27).

We found the number of grass strands within the cup of a single nest ranged from zero to

836 (Figure 2E; $\underline{x}$ ±SD: 210.19 ± 16.49). Feather count within the cup was also quite variable,

ranging from a singular pennaceous feather to 271 total feathers (Figure 2F; $\underline{x}$ ±SD: 83.79 ±

5.70). Plumulaceous feathers were the most common feather found in nests with an average

ratio of 5 plumulaceous feathers for every 1 pennaceous. Although the cup is primarily lined

with grasses and feathers, other materials such as snake skin ($n = 19/103$; 18.4%),

anthropogenic objects (e.g., plastic; $n = 54/103$; 52.4%), and hair, noted anecdotally but not

quantified, from a variety of mammalian species, including human, were observed. We also

consistently found varying amounts of spider egg sac material interspersed throughout the stick

platform ($n = 88/103$; 85.4%).

The ratio of the nest height to internal box height ($P = 0.74$; Figure 2A) and the width of

the nest berm (Figure 2D; $P = 0.94$) were normally distributed across the population. All other

continuous nest characteristics showed a right-skewed distribution ($P < 0.001$).

Figure 2 - Histograms showing the number of nests that contained the specified variable.

"Height Ratio" (A; P = 0.74) and "Width of the Nest Berm" (D; P = 0.94) exhibit a normal

distribution, but other factors show a right-skewed distribution (P < 0.001). The blue plots

layered over the histograms represent the outcome of an estimated density distribution.

Variation Explained by Female Builder Identity

Over the course of the nesting season, 24 females renested in our boxes after their initial

clutch. Of those females, 19 renested once and 5 renested twice on our study site, and the

majority of those females renested in a different box from any previous clutch (22/29 total

renests; 75.9%; Supplementary 1). The identity of the nesting female did not explain the

variation in the ratio of the nest height to internal box height ($R = 0 \leq 0.24 \leq 0.56$, $P = 0.08$,

Figure 3C), nest volume ($R = 0 \leq 0.22 \leq 0.53$, $P = 0.11$), the width of the berm ($R = 0 \leq 0 \leq$

0.34, $P > 0.99$, Figure 3E), or the number of sticks within the the nest ($R = 0 \leq 0.15 \leq 0.49$, $P =$

0.20, Figure 3A). However, the grass material count was significantly repeatable ($R = 0.23 \leq$

$0.56 \leq 0.77$, $P = 0.003$, Figure 3B). The total count of feathers in the cup was not repeatable (R

$= 0 \leq 0.04 \leq 0.39$, $P = 0.37$), but the pennaceous feather count alone appeared repeatable ($R = 0$

$\leq 0.33 \leq 0.62$, $P = 0.05$, Figure 3F), although we do note that the confidence interval overlaps

with zero. The ratio of pennaceous feathers to plumulaceous feathers within the cup of the nest

was significantly repeatable ($R = 0.79 \leq 0.90 \leq 0.95$, $P = 0.002$, Figure 3D). No other

measurement was repeatable (Table 2).

Table 2 - Listed nesting materials were not significantly repeatable for both female and nesting

box identity. Material specified with "✝" was measured as a simple yes/no presence.

	Box ID		Female ID	
	R & 95% CI	P	R & 95% CI	P
Plumulaceous	$0 \leq 0.03 \leq 0.38$	0.39	$0 \leq 0.02 \leq 0.37$	0.43
Faecal Rate	$0 \leq 0.07 \leq 0.24$	< 0.99	$0 \leq 0.07 \leq 0.25$	< 0.99
Spider Rate	$0 \leq 0.07 \leq 0.24$	< 0.99	$0 \leq 0.07 \leq 0.23$	< 0.99
✝ Snake Skin	$0 \leq <0.001 \leq 0.88$	0.64	$0 \leq 0.14 \leq 0.93$	0.17
✝ Plastic	$0 \leq 0.04 \leq 0.35$	0.32	$0 \leq <0.001 \leq 0.32$	0.48

Figure 3 - The repeatability of nest measurements across successive clutches for individual

females. Significant repeatability by female identity is indicated by the presence of a house

wren drawing. Female house wrens construct similar nest cups across multiple nesting

attempts with the grass count (B; P = 0.004), feather ratio (D; P = 0.002), and pennaceous

feather count (F; P = 0.05) each indicating significant repeatability for females. However,

none of the characteristics of the stick platform showed significant repeatability (P > 0.08).

Characteristics of the stick platform are shown in orange (left), and characteristics of the cup

lining are shown in blue (right).

Effects of Nest Box Dimensions

The height ($r^2 = 0.06$, $P = 0.006$) and volume ($r^2 = 0.16$, $P < 0.001$, Figure 4A) of the nest

platform positively correlated with the internal height of the box. Likewise, the number of

sticks within the platform positively correlated with the internal height ($r^2 = 0.12$, $P < 0.001$,

Figure 4B) and volume ($r^2 = 0.14$, $P < 0.001$) of the box. The height of the nest was positively

correlated with the height of the rectangular gap above the entrance hole ($r^2 = 0.13$, $P < 0.001$,

Figure 5A). It is worth noting that within our sample, boxes with taller internal heights tended

to have higher rectangular gaps ($r^2 = 0.13$, $P < 0.001$, Figure 5B). The width of the berm

positively correlated with the diameter of the entrance hole ($r^2 = 0.03$, $P = 0.05$, Figure 6A), but

not the depth of the entrance hole ($P = 0.43$, Figure 6B). The closer the top of the nest was built

to the entrance hole (See Figure 1), the larger the width of the berm ($r^2 = 0.09$, $P = 0.001$). No

other measurement correlated with box dimensions (Table 3).

Table 3 - P-values of the correlations between nest characteristics and box dimensions.

Significant correlations are indicated by bold text.

	Internal Box Height	Internal Box Volume	Gap Height	Entrance Diameter	Entrance Depth
Nest Height	**0.006**	0.86	**< 0.001**	0.63	0.58
Nest Volume	**< 0.001**	**< 0.001**	0.12	0.23	0.85
Nest Berm	0.42	0.053	0.54	**0.05**	0.43
Stick	**< 0.001**	**< 0.001**	0.12	0.81	0.85
Grass	0.20	0.37	0.84	0.19	0.65
Total Feather	0.60	0.88	0.81	0.27	0.08
Pennaceous	0.65	0.69	0.88	0.30	0.16
Plumulaceous	0.64	0.97	0.73	0.33	0.10

Figure 4 - Significant positive correlations between the internal height of the nesting box and the total volume of the nest (A; P < 0.001) and the stick count (B; P < 0.001).

Figure 5 - Significant positive correlations between rectangular gap height and the height of the nest (A; P < 0.001) and internal box height (B; P < 0.001).

Figure 6 - Relationships between the width of the nest berm and the diameter or depth of the

entrance hole. There was a significant positive correlation between the diameter of the

entrance hole and the width of the nest berm (A; P = 0.05), but no detectable correlation

between the depth of the entrance hole and the width of the nest berm (B; P = 0.43).

Variation Explained By Nest Box Identity

Twenty-six of our boxes were used more than once over the course of the nesting season. Of the boxes, 23 were used twice while 3 were used three times. Of the boxes used more than once, the majority of the nesting attempts were from a different female than any previous attempt (22/29 total reuses; 75.9%; Supplementary 1). The box ID significantly explained the ratio of the nest height to internal box height ($R = 0.25 \leq 0.56 \leq 0.77$, $P < 0.001$, Figure 7C), nest volume ($R = 0.37 \leq 0.65 \leq 0.82$, $P < 0.001$), and the width of the nest's berm ($R = 0.15 \leq 0.49 \leq 0.72$, $P = 0.002$, Figure 7E). Box ID significantly explained the number of sticks within the nest platform ($R = 0.10 \leq 0.45 \leq 0.7$, $P = 0.008$, Figure 7A), but not the grass count ($R = 0 \leq 0.04 \leq 0.38$, $P = 0.38$, Figure 7B), total count of feathers in the nest ($R = 0 \leq 0 \leq 0.34$, $P > 0.99$) or the pennaceous feather count ($R = 0 \leq 0.08 \leq 0.42$, $P = 0.32$, Figure 7F). Box ID

potentially further explained part of the variation in the pennaceous to plumulaceous feather

ratio ($R = 0 \le 0.29 \le 0.59$, $P = 0.05$, Figure 7D), although the confidence interval overlaps with

zero. The box ID did not explain the amount of spider egg sac material in the nest ($R = 0 \le 0 \le$

0.21, $P > 0.99$). No other measurements were repeatable (Table 2).

Table 4 - P-values supporting the lack of significant correlations between nest characteristics

and the nest age.

	Grass Count	Total Feather	Wing	Body	Age
Nest Height	0.95	0.94	0.69	0.81	0.12
Nest Volume	0.71	0.89	0.49	0.93	0.14
Nest Berm	0.76	0.93	0.37	0.83	0.78
Stick	0.19	0.35	0.32	0.44	*0.052*

Figure 7 - The repeatability of nest measurements across successive clutches built in the same

nest box. Significant repeatability by box identity is indicated by the presence of a nest box

drawing. The stick count (A; P = 0.007), height ratio (C; P < 0.001), berm width (E; P =

0.002), and feather ratio (D; P = 0.05) were significantly repeatable traits for the nesting box

identity, while grass Count (B; P = 0.39) and pennaceous (F; P = 0.32) were not. Line colours

as in Figure 3.

Nest Dimensions and Age

The width of the berm positively correlated with the height (r^2 = 0.03, P = 0.04), volume

(r^2 = 0.09, P < 0.001, Figure 8A), and stick count (r^2 = 0.04, P = 0.02, Figure 8B) of the nest.

The stick count positively correlated with the height ($r^2 = 0.37$, $P < 0.001$, Figure 8C) and

volume ($r^2 = 0.41$, $P < 0.001$, Figure 8D) of the nest. Total feather count ($r^2 = 0.03$, $P = 0.04$,

Figure 8E) and pennaceous feather count ($r^2 = 0.06$, $P = 0.005$, Figure 8F), but not

plumulaceous feather count ($P = 0.15$) positively correlated with the grass count. No other

correlations were statistically significant (Table 4).

The pennaceous feather count positively correlated with the age of the nest ($r^2 = 0.04$, $P =$

0.02, Figure 9A), but the plumulaceous ($P = 0.43$) and total feather ($P = 0.18$) counts did not.

The grass count ($r^2 = 0.03$, $P = 0.054$, Figure 9D) and stick count ($r^2 = 0.03$, $P = 0.052$, Figure

9C) marginally negatively correlated with the age of the nest. As expected, younger nests were

also less likely to contain faecal material, with nests older than 20 days almost always

containing some amount of faecal material ($t = 5.24$, $P < 0.001$, Figure 9B).

Figure 8 - Relationships between different nest attributes and building materials. The width of the nest berm positively correlated with the total volume (A; P < 0.001) and stick count (B; P = 0.02) of the nest. The stick count further highly positively correlated with the height of the nest (C; P < 0.001) and the total volume of the nest (D; P < 0.001). The grass count positively correlated with the total count of feathers (E; P = 0.04) and pennaceous feather count (F; P = 0.005).

Figure 9 - Relationship between three nest materials and faecal material with the age of the

nest at collection. Pennaceous feather count positively correlated with the nest age (A; P =

0.02). Although not statistically significant, the stick count (C; P = 0.052) and the grass count

(D; P = 0.054) negatively correlated with the age of the nest. Older nests contained more

faecal material than younger nests (B; *P < 0.001*).

Effects of Seasonal Change and Proxies of Fitness

We did not find the ratio of the nest height to internal box height ($P = 0.54$) or the width

of the berm ($P = 0.74$) to correlate with the date of measurement. Neither the stick count ($P =$

0.72) or grass count ($P = 0.88$) correlated with the measurement date. A statistical trend was

found with the pennaceous feather count positively correlating with the measurement date ($r^2 =$

0.02, $P = 0.07$, Figure 10C), with the result reaching statistical significance with the removal of

an extreme outlier (Date of year of outlier: 144; $r^2 = 0.09$, $P = 0.001$), but this trend was not

found with plumulaceous feathers ($P = 0.27$) or total feather count ($P = 0.15$). The amount of

spider egg sacs within the nest positively correlated with the measurement date ($t = 4.485$, $P <$

0.001, Figure 10D). Nests sampled later in the breeding season were more likely to produce

successful clutches than nests built earlier in the season ($t = 2.039$, $P = 0.04$, Figure 10A) but

also fledged fewer offspring ($t = -5.890$, $P < 0.001$, Figure 10B). However, this result is likely

due to females laying fewer eggs later in the season ($t = -10.26$, $P < 0.001$).

No correlations between the body condition of the female builder, clutch size, or

offspring success and nest morphology were observed (Supplementary 2).

Figure 10 - Significant relationships between nest characteristics and the date of nest

measurement (i.e., completion of egg laying). Nests measured earlier in the season were more

likely to fail than nests measured later in the season (A; P = 0.04), but also fledged fewer

offspring (B; P = 0.002). The pennaceous feather count positively correlated with the

measurement date after the removal of the outlier on DOY 144 (C; P = 0.001). Nests measured

later in the season also contained more spider sac material than those measured earlier in the

season (D; P < 0.001).

DISCUSSION

Extended Phenotypes are Complex

The niche-construction hypothesis argues that construction behaviours should be

relatively invariable, both spatially and temporally, because the benefits of the product of the

construction should be long term and stable (Clark et al. 2020). However, we found house

wren nests to be extremely variable both among-individuals and across the breeding season.

Consistent with previous research (Alworth 1996, Eckerle & Thompson 2005), the variation

was unrelated to female body condition or the number of offspring fledged, two important

proxies of the builder's fitness, suggesting that nest architecture is not under strong selection.

We asked: in the absence of strong selection for specific nest characteristics, what explains the

observed variability?

Extended phenotypes are a complex bidirectional interaction between the builder and

their environment, with builders attempting to express their phenotype within the confines of

available space. We found evidence of repeatability for the female builder in multiple

components of the cup composition. We also found that nest morphology was affected by the

availability of space within the nesting cavity, supported by repeatability in the cavity identity.

Our findings suggest a previously unexplored explanation for variation in nest architecture:

differences between individual females in their building preferences. Together with the cavity

space's ability to constrain or enhance the nest dimensions, these two factors contribute to the

complex interplay between the female builder and the environment in the expression of the

wren's extended phenotype. We also employed an uncommon method for measuring

construction behaviour (i.e., directly measuring individual building materials) leading to the

discovery of repeatable characteristics that would have otherwise remained hidden with more

coarse measurements like nest weight. Future research should compare the variation across

populations and breeding seasons to fully understand the spatial and temporal variation of nest

architecture in this species.

Preferences in Construction behaviour

Our prediction that the nesting female would be the best predictor of variation in nest

architecture was only partially supported. The cup composition (i.e., the grass count,

pennaceous feather count, and ratio of pennaceous to plumulaceous feathers) was highly

repeatable for the female, but female identity did not explain variation in the rest of the nest

morphology. This finding reinforces previous conclusions that the female constructs the nest

cup independent of the male (McCabe 1965), but implicates other factors in the construction of

the stick platform, such as physical constraints. The cup appears largely unaffected by these

constraints and similarly unrelated to fitness. We hypothesise that when constraints and

selective pressures favouring a specific nest architecture are reduced, the effect of diverse

female building preferences on nest variation is enhanced. Female blue tit (*Cyanistes*

caeruleus) nests, for example, are also extremely variable in height and this behaviour is

moderately repeatable for females ($R = 0.40$; Järvinen et al. 2017a) despite nest height in this

species appearing unrelated to nestling survival or female body condition (Järvinen et al.

2017b). Therefore, variation between individual construction preferences is likely a driving

force behind much of the observed variability in nest architecture for cavity nesters.

If much of the nest variation is explained by diversity in individual preferences, it is

important to understand what causes builders to differ from each other. Although this study is

one of the first to tie the repeatability of an extended phenotype directly to individuals in a

free-living population, lab experiments have implicated differences in experience as a potential

factor that affects individual preferences. For example, zebra finch (*Taeniopygia castanotis*)

pairs that had previous experience constructing nests built more repeatable nests during

experimental trials (Whittaker et al. 2023) and juvenile black widow spiders (*Latrodectus*

mactans) exposed to conspecific webs built more repeatable webs in adulthood (DiRienzo et

al. 2019). Overall, having prior experiences with construction behaviours tends to result in

individuals building more repeatable structures, with experiences being either previous

attempts or social interactions.

Some females in our study were more consistent builders than others, but we find it

unlikely that experience from prior breeding seasons could have contributed to these

differences. Long-term demographic data from multiple northern house wren populations

indicate that the majority of individuals only breed in a single season (average lifespan is 1.5

years; interannual return rates ~30%; Johnson 2020) and so are unlikely to provide a sufficient

sample size for an analysis of interannual individual differences. The age of individuals could

act as a proxy of experience in some species, but we are unable to present an analysis of its

effect on repeatability in this study. While moult patterns allow many songbirds to be aged as

either in their first breeding season or older, this process is challenging in house wrens (Pyle

1997), and we are not confident in our age assignments.

Individuals are also affected by trade-offs between the construction of the extended

phenotype and other behaviours important for fitness. Starved colonies of *Stegodyphus*

sarasinorum, for example, invest more in web construction than fed spiders (Ellendula et al.

2021), and immune stressed three-spined stickleback (*Gasterosteus aculeatus*) males construct

less compact nests than healthy conspecifics (Barber et al. 2001). Thus, construction

behaviours can be important indicators of the builder's condition. We examined the female

body mass and the linear residual between body mass and tarsus length as two measures of

female condition as an explanation for among-individual variation, but neither correlated with

the observed variability in nest cup composition.

Construction behaviours can also be directly related to reproductive effort and success,

particularly in oviparous species, with this relationship often resulting in low among-individual

variation (Clark et al. 2020). Zebra finches modify successive nests when the previous clutch

fails (Edwards et al. 2020) and male barn swallows (*Hirundo rustica*) who invest more in nest

construction also invest more in the food provisioning (Møller 2006). The reproductive effort

of female house wrens declines throughout the season (e.g., females lay fewer eggs later in the

season), so we expected that pairs would similarly put less effort into nest construction.

However, in our study, nest cup composition did not differ between successful or unsuccessful

nests or correlate with the number of eggs laid, and remained highly variable throughout the

breeding season. Conversely, we found that nests collected longer after the initiation of

construction tended to contain more pennaceous feathers than nest collected sooner, and even

observed one female returning to her complete clutch of eggs with a feather in her bill,

indicating a potential upkeep behaviour in cup composition despite the increasing needs of the

offspring. The lack of any apparent reproductive cost to construction behaviour conflicts with

the results of other studies on extended phenotypes, but remains consistent with previous

literature on northern house wrens (Alworth 1996, Pribil & Picman 1997, Eckerle & Thompson

2005). We discuss potential causes for this lack of fitness-related nest architecture below.

Physical Constraints to Constructions

The box cavity was also a significant predictor of nest morphology and composition. This

repeatability supports our prediction that nest architecture would be related to constraints posed

by the cavity dimensions. House wrens appeared to be filling the available space and adjusting

nest morphology based on differences between box openings. Stanback et al. (2013) suggested

that wrens were modulating the nest structure to accommodate for perceived vulnerabilities in

the cavity, and our data support this conclusion. For example, the size of the entrance hole is a

potential source of vulnerability, as more accessible nests are more likely to be depredated and

parasitized (Paclík et al. 2009, Hauber et al. 2024). We found house wrens to have larger berm

widths in boxes with larger entrance hole diameters, potentially modulating their nest structure

to better protect the contents of the nest. Previous research has suggested that the berm could

act as a barrier between the nestlings within the cup and any outside threats (Stanback et al.

2013), however evidence for this relationship is inconsistent (Pribil & Picman, 1997). Despite

our support of Stanback et al.'s (2013) finding, the width of the nest berm had no effect on the

number of eggs laid or nestling survival. We found a similar correlation between the height of

the rectangular gap and the height of the nest, as larger gaps resulted in taller nests. House

wrens may be modulating their construction behaviours in response to perceived vulnerability,

but our results suggest that this action has little to no effect on nest success.

Evidence from other species suggests that structures can protect inhabitants from

predators and environmental threats by the construction of a physical barrier. For example,

Eurasian beavers (*Castor fiber*) in the Netherlands, which face lower predation rates and

milder climates than their northern counterparts, build lodges with thinner walls (Rosell &

Campbell-Palmer 2022). However, cavity-nesters do not need to construct a physical barrier, as

the cavity itself is the barrier between the inhabitants and the outside environment. We

hypothesise that the lack of any detectable relationship between nest architecture and fitness in

house wrens could be due to selection favouring a better quality cavity rather than specific

characteristics of the nest. Observations of platypus (*Ornithorhynchus anatinus*) nests, which

are housed inside burrows, are consistent with this hypothesis as the dimensions of the burrow

explained offspring production, while effort in nest construction was unrelated (Thomas et al.

2018). Support for this hypothesis in house wrens is more indirect, but males with empty

cavities due to experimentally removed nests were just as likely to acquire a mate as control

males that initiated nest building (Alworth 1996), suggesting that the cavity rather than the

contents are more important in mate choice. Female house wrens also use cavity quantity

before male quality when choosing mates (Eckerle & Thompson, 2006) and male house wrens

selectively choose nesting boxes with smaller entrances that likely provide superior protection

from predators and brood parasites (Pribil & Picman 1997). Collectively, the available data

suggest that the reproductive success of house wrens is more related to the cavity itself than the

nest.

Limitations

Extended phenotypes become additionally complex when more than one individual

contributes to the object of construction behaviour. Both sexes contribute to the nest building

in house wrens, so it is important to consider how the male contribution may affect our data

and interpretation. Although their contributions are likely not equal, with the female

completing the majority of the nest (Alworth & Scheiber 2000), we were unable to directly

account for the effect of individual males on construction behaviour because the majority of

555 males were unbanded. Despite this limitation, northern house wren pairs often change between

556 nesting attempts (Baldwin 1921, Drilling & Thompson 1988), so it is rarely the same pair

557 building successive nests. In 2019, we found that out of 31 banded pairs in our population, only

558 6 (19.4%) remained with the same mate for the entire breeding season, with the female being

559 more likely to leave than the male. The fact that females were repeatable in cup composition,

560 despite frequent mate switching, supports our conclusion that the female constructs some

561 aspects of the nest independently of the male. Box identity could potentially be a proxy of the

562 male identity, as cavities are controlled by individual males on their own territories; however,

563 males often control many boxes within the same territory and the rate of movement between

564 these boxes is unknown. Therefore, we are unable to split apart the identity of the male from

565 the identity of the box in this study. Males begin bringing sticks to the cavity before acquiring

566 a mate (Johnson 2020), and so if the box identity is a reliable proxy for male identity, then the

567 male contribution to building could help explain the observed variation in the stick platform,

568 but more research is needed.

569  In most cases, we only measured the construction attempts of individuals twice. House

570 wrens only produce two to three clutches a season, so we are limited by this number and the

571 likelihood the female renested within our nest boxes instead of natural cavities. However, Bell

572 et al. (2009) found no significant effect of the number of observations per individual on

573 repeatability estimates. Instead, the number of observations appears to have a greater effect on

574 the range of uncertainty in the estimate. We provided this range of uncertainty, as a confidence

575 interval, and required the lower end to remain higher than zero for significance. Furthermore,

576 all of our estimates are calculated from a sample size ($N = 24$ for female identity; $N = 26$ for

box identity) larger than the median sample size ($N = 22$) of all studies examined by Bell et al.

(2009), and so we do not believe that our sample size is a major limitation.

Conclusions

The repeatability of construction behaviours is greatly understudied, particularly in free-

living populations. We examined the northern house wren, a species with incredible variation

in nest architecture, to estimate the amount of variability in construction behaviour explained

by the identity of the female builder and the box identity and its dimensions. We found female

northern house wrens to be highly repeatable in the composition of the nest cup, but this

behaviour was unrelated to female body condition or nestling survival. The variation in the

main structure of the nest, the stick platform, was best explained by the box identity and its

dimensions, supporting previous findings that house wrens modify aspects of the nest to

compensate for vulnerabilities in the cavity. We hypothesise that the observed variability in

nest architecture is due to the nest being under minimal selective pressure, thus allowing the

diversity of female building preferences to have a greater effect on the expression of the

phenotype. Instead, choosing a protective cavity is likely to be more important for the builder's

fitness, with the nest construction process being relatively unimportant for mate selection or

nestling survival. However, much of house wren nest construction still remains unclear,

including the contribution of the male in constructing the stick platform. Extended phenotypes

are complex, requiring an examination of both the environment *and* the builder. The cause of

variation in house wren nest morphology has been largely unexplained due to a narrow focus

on potential external factors. Studying the repeatability of extended phenotypes across taxa has

provided new insights into the construction behaviours of animals and provides an important

method for understanding the causes and consequences of individual preferences.

Data Availability Statement

The data and code that support these findings are available in the Mendeley Data

Repository at doi: 10.17632/fnhr5mm99p.1.

ACKNOWLEDGMENTS

We thank Ohio Wesleyan University, the Big Walnut School System, and the Davis,

Koban, and Fink families for access to facilities and field sites. We also like to thank Ali Amer

for helping in nest dissection, Ava Swanson for the creation of the house wren and box

drawings used in Figures 3 and 7, and Martin Stoffel for statistical advice. Finally, we would

like to thank the house wrens for making our work possible. This project was funded by the

Ohio Wesleyan University Summer Science Research Program.

REFERENCES

Adreani, N. M., Valcu, M., Scientists, C., & Mentesana, L. (2022). Asymmetric architecture

is non-random and repeatable in a bird's nests. *Current Biology*, *32*(9), R412–R413.

https://doi.org/10.1016/j.cub.2022.03.075

Alworth, T. (1996). An Experimental Test of the Function of Sticks in the Nests of House

Wrens. *The Condor*, *98*(4), 841–844. https://doi.org/10.2307/1369866

Alworth, T., & Scheiber, I. B. R. (2000). NEST BUILDING IN HOUSE WRENS

(TROGLODYTES AEDON): A REEXAMINATION OF MALE AND FEMALE

ROLES. *Journal of Field Ornithology*, *71*(3), 409–414. https://doi.org/10.1648/0273-

8570-71.3.409

Bailey, I., Morgan, K., Oschadleus, D., DeRuiter, S., Meddle, S., & Healy, S. (2015). Nest-

building males trade-off material collection costs with territory value. *The Emu: Official*

Organ of the Australasian Ornithologists' Union, *116*.

https://doi.org/10.1071/MU15022

Baldwin, S. P. (1921). The Marriage Relations of the House Wren (Troglodytes A. Aedon).

The Auk, *38*(2), 237–244. https://doi.org/10.2307/4073886

Barber, I., Nairn, D., & Huntingford, F. A. (2001). Nests as ornaments: Revealing

construction by male sticklebacks. *Behavioral Ecology*, *12*(4), 390–396.

https://doi.org/10.1093/beheco/12.4.390

Beebe, W., & Beebe, W. (1953). A contribution to the life history of the euchromid moth,

Aethria carnicauda Butler. *Zoologica : Scientific Contributions of the New York*

Zoological Society, *38*(13), 155--160. https://doi.org/10.5962/p.203451

Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: A

meta-analysis. *Animal Behaviour*, *77*(4), 771–783.

Biddle, L. E., Broughton, R. E., Goodman, A. M., & Deeming, D. C. (2018). Composition

of Bird Nests is a Species-Specific Characteristic. *Avian Biology Research*, *11*(2), 132–

153. https://doi.org/10.3184/175815618X15222318755467

Blamires, S. J., Lee, Y.-H., Chang, C.-M., Lin, I.-T., Chen, J.-A., Lin, T.-Y., & Tso, I.-M.

(2010). Multiple structures interactively influence prey capture efficiency in spider orb

webs. *Animal Behaviour*, *80*(6), 947–953.

https://doi.org/10.1016/j.anbehav.2010.09.011

Borgia, G. (1985). Bower quality, number of decorations and mating success of male satin

bowerbirds (Ptilonorhynchus violaceus): An experimental analysis. *Animal Behaviour*,

33(1), 266–271. https://doi.org/10.1016/S0003-3472(85)80140-8

Brown, W., Alexander, A., Alexander, D., Zuefle, M., & Underwood, T. (2024). Estimating

Ages of House Wren Nestlings Based on Body Mass, Wing Chord Length, and Feather

Tract Development Patterns. *North American Bird Bander*, *36*(3).

https://digitalcommons.usf.edu/nabb/vol36/iss3/1

Cerezo, A. T., & Deeming, D. C. (2016). Nest Insulation in the European Robin (Erithacus

Rubecula): Effects of Latitude and Construction Materials. *Avian Biology Research*,

9(2), 96–102. https://doi.org/10.3184/175815516X14551263654405

Clark, A. D., Deffner, D., Laland, K., Odling-Smee, J., & Endler, J. (2020). Niche

Construction Affects the Variability and Strength of Natural Selection. *The American*

Naturalist, *195*(1), 16–30. https://doi.org/10.1086/706196

Darwin, C., & Wallace, A. (1858). On the Tendency of Species to form Varieties; and on the

Perpetuation of Varieties and Species by Natural Means of Selection. *Journal of the*

Proceedings of the Linnean Society of London. Zoology, *3*(9), 45–62.

https://doi.org/10.1111/j.1096-3642.1858.tb02500.x

Dawkins, R. (1982). *The Extended Phenotype: The Long Reach of the Gene*. Oxford

University Press.

DiRienzo, N., & Aonuma, H. (2018). Plasticity in extended phenotype increases offspring

defence despite individual variation in web structure and behaviour. *Animal Behaviour*,

138, 9–17. https://doi.org/10.1016/j.anbehav.2018.01.022

DiRienzo, N., Johnson, J. C., & Dornhaus, A. (2019). Juvenile social experience generates

differences in behavioral variation but not averages. *Behavioral Ecology*, *30*(2), 455–

464. https://doi.org/10.1093/beheco/ary185

DiRienzo, N., & Montiglio, P.-O. (2016). Linking consistent individual differences in web

structure and behavior in black widow spiders. *Behavioral Ecology*, *27*(5), 1424–1431.

https://doi.org/10.1093/beheco/arw048

DiRienzo, N., Schraft, H. A., Montiglio, P. O., Bradley, C. T., & Dornhaus, A. (2020).

Foraging behavior and extended phenotype independently affect foraging success in

spiders. *Behavioral Ecology*, *31*(5), 1242–1249. https://doi.org/10.1093/beheco/araa080

Drilling, N. E., & Thompson, C. F. (1988). Natal and Breeding Dispersal in House Wrens

(Troglodytes aedon). *The Auk*, *105*(3), 480–491.

Eckerle, K. P., & Thompson, C. F. (2005). Addition of arthropod cocoons to house wren

nests is correlated with delayed pairing. *Behavioral Ecology*, *16*(1), 1–7.

https://doi.org/10.1093/beheco/arh122

Edwards, S. C., Shoot, T. T., Martin, R. J., Sherry, D. F., & Healy, S. D. (2020). It's not all

about temperature: Breeding success also affects nest design. *Behavioral Ecology*,

31(4), 1065–1072. https://doi.org/10.1093/beheco/araa052

Ellendula, S., Tresa, C., & Uma, D. (2021). Influence of prey availability on web-building in

the social spider Stegodyphus sarasinorum (Araneae: Eresidae). *The Journal of*

Arachnology, *49*(1), 141–145. https://doi.org/10.1636/JoA-S-19-080

Fisher, D. N., Yeager, J., & Pruitt, J. N. (2019). *Orb-weaving spiders show a correlated*

syndrome of morphology and web structure in the wild (p. 767335). bioRxiv.

https://doi.org/10.1101/767335

Franklin, D. C. (1995). Helmeted Honeyeaters Build Bulkier Nests in Cold Weather. *The*

Auk, *112*(1), 247–248. https://doi.org/10.2307/4088785

Hansell, M. (2000). *Bird Nests and Construction Behaviour*. Cambridge University Press.

https://doi.org/10.1017/CBO9781139106788

Hansell, M. H. (2005). *Animal Architecture*. OUP Oxford.

Hauber, M. E., Nagy, J., Sheard, C., Antonson, N. D., Street, S. E., Healy, S. D., Lala, K. N.,

& Mainwaring, M. C. (2024). Nest architecture influences host use by avian brood

parasites and is shaped by coevolutionary dynamics. *Proceedings of the Royal Society*

B: Biological Sciences, *291*(2014), 20231734. https://doi.org/10.1098/rspb.2023.1734

Hayes, J. P., & Jenkins, S. H. (1997). Individual Variation in Mammals. *Journal of*

Mammalogy, *78*(2), 274–293. https://doi.org/10.2307/1382882

Hays, S. C., Cheek, R. G., Mouton, J. C., Sillett, T. S., & Ghalambor, C. K. (2022). Lack of

avian predators is associated with behavioural plasticity in nest construction and height

in an island songbird. *Animal Behaviour*, *194*, 35–42.

https://doi.org/10.1016/j.anbehav.2022.08.017

Hermes, M. G., Somavilla, A., & Garcete-Barrett, B. R. (2013). On the nesting biology of

Pirhosigma Giordani Soika (Hymenoptera, Vespidae, Eumeninae), with special

reference to the use of vegetable matter. *Revista Brasileira de Entomologia*, *57*, 433–

436. https://doi.org/10.1590/S0085-56262013005000044

Hölldobler, B., & Wilson, E. O. (1990). *The Ants*. Harvard University Press.

Järvinen, P., & Brommer, J. E. (2020). Nest ornaments and feather composition form an

extended phenotype syndrome in a wild bird. *Behavioral Ecology and Sociobiology*,

74(11), 134. https://doi.org/10.1007/s00265-020-02912-2

Järvinen, P., Kluen, E., & Brommer, J. E. (2017a). Low heritability of nest construction in a

wild bird. *Biology Letters*, *13*(10), 20170246. https://doi.org/10.1098/rsbl.2017.0246

Järvinen, P., Kluen, E., Tiiri, M., & Brommer, J. (2017b). Experimental manipulation of

Blue Tit nest height does not support the thermoregulation hypothesis. *Ornis Fennica*,

94. https://doi.org/10.51812/of.133914

Jelínek, V., Požgayová, M., Honza, M., & Procházka, P. (2016). Nest as an extended

phenotype signal of female quality in the great reed warbler. *Journal of Avian Biology*,

47(3), 428–437. https://doi.org/10.1111/jav.00700

Johnson LS. 2020. House Wren (*Troglodytes aedon*), version 1.0. In: Poole AF, editor.

Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA.

Korb, J. (2003). Thermoregulation and ventilation of termite mounds. *Naturwissenschaften*,

90(5), 212–219. https://doi.org/10.1007/s00114-002-0401-4

Korb, J., & Linsenmair, K. E. (2000). Thermoregulation of termite mounds: What role does

ambient temperature and metabolism of the colony play? *Insectes Sociaux*, *47*(4), 357–

363. https://doi.org/10.1007/PL00001731

Laland, K. N., Odling-Smee, F. J., & Feldman, M. W. (1999). Evolutionary consequences of

niche construction and their implications for ecology. *Proceedings of the National*

Academy of Sciences of the United States of America, *96*(18), 10242–10247.

LeGrice, R. J., Tezanos-Pinto, G., de Villemereuil, P., Holwell, G. I., & Painting, C. J.

(2019). Directional selection on body size but no apparent survival cost to being large

in wild New Zealand giraffe weevils. *Evolution*, *73*(4), 762–776.

https://doi.org/10.1111/evo.13698

Madden, J. R. (2002). Bower Decorations Attract Females but Provoke Other Male Spotted

Bowerbirds: Bower Owners Resolve This Trade-off. *Proceedings: Biological Sciences*,

269(1498), 1347–1351.

Mainwaring, M., Benskin, C., C.McW.H, & Hartley. (2008). The Weight of Female-Built

Nests Correlates with Female but not Male Quality in the Blue Tit Cyanistes caeruleus.

Acta Ornithologica, *43*, 43–48. https://doi.org/10.3161/000164508X345310

Mainwaring, M. C., Deeming, D. C., Jones, C. I., & Hartley, I. R. (2014b). Adaptive

latitudinal variation in Common Blackbird Turdus merula nest characteristics. *Ecology*

and Evolution, *4*(6), 851–861. https://doi.org/10.1002/ece3.952

Mainwaring, M. C., Hartley, I. R., Lambrechts, M. M., & Deeming, D. C. (2014a). The

design and function of birds' nests. *Ecology and Evolution*, *4*(20), 3909–3928.

https://doi.org/10.1002/ece3.1054

McCabe, R. A. (1965). Nest Construction by House Wrens. *The Condor*, *67*(3), 229–234.

https://doi.org/10.2307/1365401

McKaye, K. R., Louda, S. M., & Stauffer, J. R. (1990). Bower Size and Male Reproductive

Success in a Cichlid Fish Lek. *The American Naturalist*, *135*(5), 597–613.

Medina, I., M. Perez, D., Silva, A. C. A., Cally, J., León, C., Maliet, O., & Quintero, I.

(2022). Nest architecture is linked with ecological success in songbirds. *Ecology*

Letters, *25*(6), 1365–1375. https://doi.org/10.1111/ele.13998

Mennerat, A., Perret, P., & Lambrechts, M. M. (2009). Local Individual Preferences for

Nest Materials in a Passerine Bird. *PLOS ONE*, *4*(4), e5104.

https://doi.org/10.1371/journal.pone.0005104

Miyashita, T., & Shinkai, A. (1995). Design and Prey Capture Ability of Webs of the

Spiders Nephila clavata and Argiope bruennichii. *Acta Arachnologica*, *44*(1), 3–10.

https://doi.org/10.2476/asjaa.44.3

Møller, A. P. (1990). Nest Predation Selects for Small Nest Size in the Blackbird. *Oikos*,

57(2), 237–240. https://doi.org/10.2307/3565945

Møller, A. P. (2006). Rapid change in nest size of a bird related to change in a secondary

sexual character. *Behavioral Ecology*, *17*(1), 108–116.

https://doi.org/10.1093/beheco/arj003

Muth, F., & Healy, S. D. (2011). The role of adult experience in nest building in the zebra

finch, *Taeniopygia guttata*. *Animal Behaviour*, *82*(2), 185–189.

https://doi.org/10.1016/j.anbehav.2011.04.021

Paclík, M., Misík, J., & Weidinger, K. (2009). Nest predation and nest defence in European

and North American woodpeckers: A review. *Annales Zoologici Fennici*, *46*(5), 361–

379.

Parthasarathy, B., Bouchard, M., & Schneider, J. M. (2022). Extended phenotypes can

underlie trade-offs: A case of social spiders. *Die Naturwissenschaften*, *109*(6), 51.

https://doi.org/10.1007/s00114-022-01826-5

Perez, D. M., Gardner, J. L., & Medina, I. (2020). Climate as an Evolutionary Driver of Nest

Morphology in Birds: A Review. *Frontiers in Ecology and Evolution*, *8*.

https://www.frontiersin.org/articles/10.3389/fevo.2020.566018

Pribil, S., & Picman, J. (1997). Parasitism of House Wren nests by Brown-headed

Cowbirds: Why is it so rare? *Canadian Journal of Zoology*, *75*(2), 302–307.

https://doi.org/10.1139/z97-038

Pyle, P. (1997). Identification Guide To North American Birds: A Compendium Of

Information On Identifying, Ageing, And Sexing "Near-Passerines" And Passerines In

The Hand. *Books by Alumni*. https://works.swarthmore.edu/alum-books/4499

Rosell, F., & Campbell-Palmer, R. (2022). Habitat use and constructions. In F. Rosell & R.

Campbell-Palmer (Eds.), *Beavers: Ecology, Behaviour, Conservation, and Management*

(p. 0). Oxford University Press. https://doi.org/10.1093/oso/9780198835042.003.0004

Rushbrook, B. J., Dingemanse, N. J., & Barber, I. (2008). Repeatability in nest construction

by male three-spined sticklebacks. *Animal Behaviour*, *75*(2), 547–553.

https://doi.org/10.1016/j.anbehav.2007.06.011

Stanback, M. T., Stanbeck, M. T., Bartholomew, J. E., Bergner, L. M., Cline, E. L., Helms,

P. I., McGovern, P. G., Millican, D. M., & Roth, J. E. (2013). House Wrens Alter Nest

Architecture to Compensate for Cavity Vulnerability. *The Wilson Journal of*

Ornithology, *125*(1), 174–178.

Stoffel, M., Schielzeth, H., & Nakagawa, S. (2017). rptR: Repeatability estimation and

variance decomposition by generalized linear mixed-effects models. *Methods in*

Ecology and Evolution, *8*. https://doi.org/10.1111/2041-210X.12797

Sugiura, S. (2016). Bagworm bags as portable armour against invertebrate predators. *PeerJ*,

4, e1686. https://doi.org/10.7717/peerj.1686

Thomas, J., Handasyde, K., Parrott, M., & Temple-Smith, P. (2018). The platypus nest:

Burrow structure and nesting behaviour in captivity. *Australian Journal of Zoology*, *65*.

https://doi.org/10.1071/ZO18007

Ueda, K. (1985). *Nest Building Activity and Characteristics of the Courtship Nest of the*
Fan-tailed Warbler Cisticola juncidis. https://www.jstage.jst.go.jp/article/jjo1915/34/2-
3/34_2-3_23/_article

van Rijssel, J. C., Moser, F. N., Frei, D., & Seehausen, O. (2018). Prevalence of disruptive
selection predicts extent of species differentiation in Lake Victoria cichlids.
Proceedings of the Royal Society B: Biological Sciences, *285*(1871), 20172630.
https://doi.org/10.1098/rspb.2017.2630

Wallace, A. R. (1865). I. On the Phenomena of Variation and Geographical Distribution as
illustrated by the Papilionidæ of the Malayan Region. *Transactions of the Linnean*
Society of London, *25*(1), 1–71. https://doi.org/10.1111/j.1096-3642.1865.tb00178.x

Walsh, P. T., Hansell, M., Borello, W. D., & Healy, S. D. (2010). Repeatability of nest
morphology in African weaver birds. *Biology Letters*, *6*(2), 149–151.
https://doi.org/10.1098/rsbl.2009.0664

Whittaker, B. A., Nolet-Mulholland, L., Nevoit, A., Yun, D., Lambert, C. T., Blunk, S. C.,
& Guillette, L. M. (2023). Zebra finches have style: Nest morphology is repeatable and
associated with experience. *IScience*, *26*(11), 108194.
https://doi.org/10.1016/j.isci.2023.108194

Ziebis, W., Forster, S., Huettel, M., & Jørgensen, B. B. (1996). Complex burrows of the
mud shrimp Callianassa truncata and their geochemical impact in the sea bed. *Nature*,
382(6592), Article 6592. https://doi.org/10.1038/382619a0

SUPPLEMENTARY

Supplementary 1 - Individuals examined for Female Identity repeatable traits and Box Identity

traits. Females with single IDs were unbanded individuals.

Female Repeatability			Box Repeatability		
Female ID	Clutch	Box ID	Box ID	Clutch	Female ID
288064919	1	3	1	1	288067191
288064919	2	2	1	2	288067191
288067191	1	1	1	3	288067191
288067191	2	1	3	1	288064919
288067191	3	1	3	2	294063102
288064901	1	6	4	1	288064918
288064901	2	4	4	2	288064901
294063102	1	8	9	1	288064920
294063102	2	3	9	2	294063222
288064903	1	19	13	1	288064917
288064903	2	18	13	2	294063155
288067184	1	25	17	1	288064902
288067184	2	24	17	2	294063262
288064906	1	28	19	1	288064903
288064906	2	29	19	2	294063200
288064906	3	29	20	1	288064905
288064947	1	33	20	2	294063294

288064947	2	30	29	1	288064906
288064905	1	20	29	2	288064906
288064905	2	21	30	1	1
288064913	1	46	30	2	288064947
288064913	2	46	33	1	288064947
288064913	3	54	33	2	294063157
288064915	1	41	38	1	294063259
288064915	2	41	38	2	294063453
294063218	1	47	39	1	288064923
294063218	2	49	39	2	288064991
288064991	1	40	41	1	288064915
288064991	2	39	41	2	288064915
288064923	1	39	43	1	288064912
288064923	2	47	43	2	294063461
288064997	1	56	44	1	288064911
288064997	2	56	44	2	294063173
288064939	1	54	46	1	288064913
288064939	2	55	46	2	288064913
288067301	1	80	47	1	4
288067301	2	79	47	2	294063218
288064925	1	68	47	3	288064923
288064925	2	67	48	1	5
288067007	1	70	48	2	288064996

288067007	2	71	49	1	6
288067007	3	65	49	2	294063135
288064916	1	63	49	3	294063218
288064916	2	63	54	1	288064939
288064916	3	62	54	2	288064913
294063101	1	61	56	1	288064997
294063101	2	60	56	2	288064997
294063230	1	76	63	1	288064916
294063230	2	75	63	2	288064916
294063219	1	77	65	1	288067007
294063219	2	66	65	2	7
288064910	1	58	66	1	288064924
288064910	2	59	66	2	294063219
			70	1	288067007
			70	2	294063180

Supplementary 2 - Relationships between the dimensions and components of the nest and

various proxies of fitness: Female residual - defined by the correlation between the tarsus

length and the body mass, Fledge or Fail - the simply outcome of the nest, Eggs laid - the

number of eggs originally laid by the nesting female at the completion of the cup, and

Offspring fledged - the estimated number of offspring that fledged the nest (only successful

nests examined). ✝ *Specified material measured as a simple yes/no presence.*

	Female	Fledge or Fail	Eggs laid	Offspring

	residual			fledged
Ratio	t = 1.64, P = 0.11	z = 0.10, P = 0.92	t = -0.77, P = 0.45	t = 0.73, P = 0.47
Nest Height	t = 0.90, P = 0.37	z = -0.119, P = 0.91	t = -0.81, P = 0.42	t = 0.54, P = 0.59
Nest Volume	t = 0.46, P = 0.65	z = -0.132, P = 0.89	t = -0.63, P = 0.53	t = -0.06, P = 0.95
Nest Berm	t = -0.64, P = 0.52	z = 0.108, P = 0.91	t = 0.08, P = 0.94	t = -0.35, P = 0.73
Stick	t = -0.39, P = 0.70	z = -0.716, P = 0.47	t = -0.03, P = 0.97	t = -0.66, P = 0.51
Grass	t = 0.67, P = 0.50	z = -1.564, P = 0.11	t = 0.89, P = 0.37	t = -1.04, P = 0.30
Total Feather	t = 0.03, P = 0.97	z = 0.310, P = 0.76	t = 0.13, P = 0.90	t = -1.45, P = 0.15
Wing Rate	t = -0.12, P = 0.90	z = 1.324, P = 0.13	t = 0.28, P = 0.78	t = 0.12, P = 0.91
Pennaceous	t = -1.07, P = 0.29	z = 1.584, P = 0.08	t = -0.01, P = 0.99	t = -1.91, P = 0.06
Plumulaceous	t = 0.43, P = 0.67	z = -0.229, P = 0.82	t = -0.16, P = 0.88	t = -0.95, P = 0.35
Spider Sac Rate	t = -0.59, P = 0.56	z = 0.408, P = 0.68	t = -1.15, P = 0.25	t = -0.26, P = 0.80
✝ Snake Skin	$\chi^2_1 = 0.08$, P = 0.77	z = -0.808, P = 0.42	t = 1.202, P = 0.23	t = 1.269, P = 0.20
✝ Plastic	$\chi^2_1 = 0.002$, P = 0.96	z = -1.567, P = 0.11	t = 1.735, P = 0.08	t = 0.090, P = 0.93

833