Social ageing can protect against infectious disease in a group-living primate

Erin R. Siracusa[a*], Melissa A. Pavez-Fox[b], Josué E. Negron-Del Valle[c], Daniel Phillips[c], Michael
L. Platt[d,e,f], Noah Snyder-Mackler[c,g,h], James P. Higham[i], Lauren J. N. Brent[a] and Matthew J.
Silk[j*]

Affiliations:
[a]School of Psychology, Centre for Research in Animal Behaviour, University of Exeter, Exeter,
UK
[b]Department of Psychology and Neuroscience, University of St Andrews, UK
[c]Center for Evolution and Medicine, Arizona State University, Arizona, USA
[d]Department of Neuroscience, University of Pennsylvania, PA, USA
[e]Department of Psychology, University of Pennsylvania, PA, USA
[f]Department of Marketing, University of Pennsylvania, PA, USA
[g]School of Life Sciences, Arizona State University, Arizona, USA
[h]School for Human Evolution and Social Change, Arizona State University, Arizona, USA
[i]Department of Anthropology, New York University, New York, USA
[j]Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh,
Edinburgh, United Kingdom

*corresponding author; email: erinsiracusa@gmail.com, matthewsilk@outlook.com

Abstract

The benefits of social living are well established, but sociality also comes with costs, including

infectious disease risk. This cost-benefit ratio of sociality is expected to change across

individuals' lifespans, which may drive changes in social behaviour with age. To explore this

idea, we combine data from a group-living primate for which social ageing has been described

with epidemiological models to show that having lower social connectedness when older can

protect against the costs of a hypothetical, directly transmitted endemic pathogen. Assuming no

age differences in epidemiological characteristics (susceptibility to, severity, and duration of

infection), older individuals suffered lower infection costs, which was explained largely because

they were less connected in their social networks than younger individuals. This benefit of

'social ageing' depended on epidemiological characteristics and was greatest when infection

severity increased with age. When infection duration increased with age, social ageing was

beneficial only when pathogen transmissibility was low. Older individuals benefited most from

having a lower frequency of interactions (strength) and network embeddedness (closeness) and

benefited less from having fewer social partners (degree). Our study provides a first examination

of the epidemiology of social ageing, demonstrating the potential for pathogens to influence

evolutionary dynamics of social ageing in natural populations.

Key words: ageing, disease ecology, epidemiology, sociality, senescence, network modeling

Introduction

Social interactions have an important influence on individual fitness through their effects on

survival [1] and reproductive success [2–5]. However, sociality is not exclusively beneficial but

comes with important costs ranging from competition for food, mates, and other resources [6–8]

to increased risk of pathogen and parasite transmission [9,10]. Understanding how these costs

and benefits are balanced is central to understanding the evolution of social living [11].

Recent research indicates that social interactions and relationships are not stable over

time but differ across the lifespan [12–15]. There are multiple hypotheses for why individuals

might change their social behaviour as they get older, including physical and cognitive declines

[16,17], demographic changes and shifting kinship dynamics [18,19], or enhanced skill and

experience with age [20]. Fundamentally, the costs and benefits of interacting with others may

change across the lifespan [12], leading individuals to actively adjust their social behaviour in

response. Recent work, both in humans and non-human animals, suggests older individuals tend

to have fewer affiliative social partners than their younger counterparts [13,15,21–23], spend less

time in affiliative interactions [13,22–24], and to be less well embedded in the wider social

network [23,25,26]. There is evidence to suggest that in some species, at least, this is the result of

older individuals being more selective with whom they interact (i.e. 'social selectivity'

[21,27,28]). This evidence is consistent with the hypothesis that age-based reductions in social

connectedness (i.e. 'social ageing') are not exclusively the by-product of declines in bodily

systems but instead reflect beneficial behavioral changes that ameliorate senescence-associated

costs [21].

One major cost of socialising, which has the potential to be exacerbated in old age, is the

increased risk and costs of contracting infectious diseases [29], with individuals that interact

more with others at greater risk of infection [30,31]. Therefore, one appealing hypothesis is that,

because immunosenescence often means that individuals are less able to fight infections as they

age and so suffer greater morbidity to infectious diseases [32–34], age-based reductions in

individual social network connectedness can help mitigate disease costs in older individuals [35].

By being socially selective (i.e., reducing their number of social partners while socialising for

longer with their closest associates [21,27]), older individuals may be able to reduce their risk of

infection while maintaining the benefits of social relationships [36,37]. However, we currently

lack evidence that these declines in sociality with age (hereafter referred to as social ageing) are

sufficient to protect an individual from infectious disease.

One challenge with demonstrating the disease-associated consequences of social ageing

is the fact that an individual's social interactions and relationships are embedded in a wider

group or population wide social network. These social networks are complex, emergent

structures that depend in a non-linear way on the behaviour of every individual [38]. As a result,

it can be difficult to predict both the consequences of individual behaviour change on the group-

level social structure and the knock-on implications for pathogen spread across the network.

While connectivity (or social network density) is an important component of the vulnerability of

a group or population to an infectious disease outbreak [10], other aspects of social network

structure such as the transitivity of social connections (the tendency of an individual's

connections to also be connected to each other) and modularity (division of the network into a set

of social communities) can also shape how pathogens spread [9]. Consequently, testing whether

being more socially selective with age can protect older individuals from infectious disease costs

requires the use of epidemiological network models [39,40].

Rhesus macaques (*Macaca mulatta)* offer an excellent system in which to test this

question. They live in matrilineal multi-male multi-female social groups [41] where the social

relationships that individuals form have important effects on their fitness [8,20,42,43]. A long-

term study of a free-ranging population of rhesus macaques on Cayo Santiago Island, Puerto

Rico, has generated a unique long-term dataset encompassing multiple social groups that has

provided important insights into primate social behaviour [44]. Recent research investigating

patterns of social ageing in female rhesus macaques has demonstrated that as females get older

they reduce their number of social connections and spend more time socializing with important

partners (such as kin and partners with strong and stable connections; [21]). These changes

resulted from females approaching fewer partners as they got older, while continuing to spend

the same amount of time socializing, and were not driven by partners dying or avoiding them.

This study provides one of the few examples in non-human animals where active reductions in
sociality with age have been distinguished from more passive processes such as declines in social
engagement driven by the loss of social partners, declining social motivation, or impaired
physical ability to engage [21]. The results suggest that age-based reductions in an individual's
number of social partners might enable individuals to cope with the physiological alterations or
limitations that come with age, such as immunosenescence. Female macaques also show changes
in their indirect connectedness (connections to partners of their social partners) with age, with
measures of both betweenness (bridging capacity) and closeness (ability to reach others or be
reached in the network) declining as females get older [25]. In addition to showing clear patterns
of social ageing, rhesus macaques are a powerful system to explore the intersection of social
ageing and disease risk because they are popular biomedical models of human health and ageing
[45–47] and are frequently used as a model species in studies of infection biology, thereby
providing information on the types of pathogen they are susceptible to [45] and how costs of
infection vary with age [48–51]. The fact that the observed patterns of social ageing in macaques
closely resemble those observed in human populations [21], combined with the similarity
between macaques and humans in patterns of immunosenescence [45,47,52] and susceptibility to
pathogens (such as SARS-CoV-2 [53,54]) makes this a uniquely relevant system for
understanding the potential implications of age-based changes in sociality for disease risk in
humans.

Here we combined epidemiological network models of pathogen spread with empirical
data on social ageing from a population of free-ranging rhesus macaques. Using susceptible-
infected-susceptible (SIS) models of the spread of a hypothetical directly-transmitted pathogen
(e.g. respiratory virus) across 23 real-world rhesus macaque social networks, we hypothesised
that lower social connectivity of older individuals [21] would protect them from accumulated
infection costs of an endemic (constantly present) pathogen. Specifically, we quantified how age-
based variation in three measures of social centrality - degree (number of partners), strength
(amount of time spent socializing) and closeness (capacity to reach or be reached in the network)
influenced infection costs with age. Our choice of social metrics was based on prior knowledge
that individuals in this system show within-individual declines in both degree and closeness with
age, but not in strength [21,25]. We explored the effects of strength given it is a measure of
social connectivity that is well-known to have important consequences for disease transmission

[30]. We assessed how an individual's accumulated costs of infection in each simulation were

influenced by their age and the aforementioned network measures of social connection.

We predicted that, (1) across all measures of social centrality, the cost of infection would

decrease with reduced social connectedness in an age-dependent manner, such that for older

individuals it would be much less costly to have lower connectedness in the social network

compared with younger individuals where the benefits of having lower connectedness would be

more moderate. Next, we predicted that (2) most of the decrease in infection cost associated with

social ageing would be driven by older individuals having lower degree and closeness since

strength was not expected to change with age in this population. We further predicted that (3)

social ageing would provide the greatest benefit, in terms of reducing infection cost, when

immunosenescence, and therefore infection cost with age, was greatest. To assess this we

explored the extent to which these social variables influenced infection cost across a range of

pathogen transmissibilities and under conditions where we varied different components of

immunosenescence, including how susceptibility to (likelihood of acquiring infection), as well as

severity (per timestep cost of infection) and duration of infection changed with age.

For the sake of clarity and brevity, throughout the manuscript we refer to this reduction

in infection cost that is the result of older individuals having lower social connectivity as the

'protective effect of social ageing' although our analyses are at the population level rather than

being exclusively within-individual, and this interpretation therefore warrants caution. However,

given that our previous results [21,25] have shown that age-based differences in sociality are

driven by within-individual changes rather than between individual differences (at least for

degree and closeness), it is appropriate to assume that population level trends are the result of

within-individual processes [55,56]. Our models are the first to explicitly test whether reduced

social connectivity among older individuals is sufficient to buffer against infectious disease costs

in a free-ranging population.

Methods

Study System

We studied a population of rhesus macaques on the island of Cayo Santiago off the southeast

coast of Puerto Rico. This population was first introduced to the island in 1938 from India and

currently consists of ~ 1800 individuals living in 12 mixed-sex social groups. The population is

maintained by the Caribbean Primate Research Center, which is responsible for monitoring the population daily and collecting data on births, deaths, and social group membership. The animals are food supplemented and provided *ad libitum* access to fresh water. The island is predator free and there is no veterinary intervention for sick or wounded animals, meaning that the majority of deaths on the island are from natural causes such as illness and injury [8].

For this study we focused on adult female rhesus macaques (aged 6 years and older) from six social groups that have been studied intensively between 2010 and 2022 and therefore for which we had detailed behavioural data to build social networks. In total we used behavioural data collected from twenty-three different group years (group F 2010-2017, group HH 2014 & 2016, group KK 2013, 2015 & 2017, group R 2015 & 2016, group S 2011 & 2019, group V 2015-2017, 2019, 2021-2022). For these analyses we excluded data collected in 2018 and 2020 because Hurricane Maria and the COVID-19 pandemic, respectively, precluded use of our typical protocol to collect behavioural data. In total this resulted in 1176 macaque-years of data from 410 unique females whose ages ranged from 6-28 years old (mean = 11.2; see Fig. S1). We collected behavioural data between 07:30 - 14:00, which are the working hours of the field station. Behavioural data were collected using 10-min (20 group years) or 5-min (3 group years) focal animal samples, where all behaviours were recorded continuously. We stratified sampling to ensure balanced data collection on individuals throughout the day and over the course of the year. Given that previous research in this system has shown that there are clear age-based changes in grooming associations among female macaques [21,25], in addition to the fact that behaviours with prolonged contact (such as grooming) are known to be highly relevant for parasite transmission [57–62], we focused specifically on grooming interactions to build our social networks (see below). During focal observations, we recorded the duration of the grooming bout as well as the identity of the monkeys and the direction of the grooming behaviour (give or receive). Grooming bouts had to be at least 5 seconds long to be recorded and a new bout of grooming was recorded if the identity of the monkeys or direction of grooming changed or there was at least a 15 second pause in grooming behaviour. These thresholds are based on long-term expert knowledge of the study system and have been used since 2010 in the collection of grooming interactions. We focused our study on adult females because the age-based changes in sociality previously demonstrated in this system were for female networks that excluded interactions with males and juveniles or subadults of either sex [21,25]. Focusing on

female-female interactions also allowed us to isolate how changes in cooperative interactions

with age influence disease transmission outside of age-based changes in socio-sexual behaviour.

This is because females' interactions with males are likely to capture both cooperative

interactions as well as reproductive behaviours, making it difficult to parse age-based changes in

cooperation from age-based changes in reproduction.

Social network construction

We constructed grooming networks for 23 group years (6 groups across 12 years; mean = 3.8,

range = 2-8 years per group). In these networks, nodes represent individuals and edges represent

the undirected rate of grooming between a pair of individuals (number of grooming bouts/total

number of focals of both individuals). Although we have data on both grooming given and

received, here grooming serves as a proxy for time spent in close contact. We expected that the

total amount of time and number of partners with which an individual spent time in close contact

was likely to be most relevant for infection risk from a hypothetical directly-transmissible

pathogen, regardless of directionality. Because our social networks were constructed from

observational data, which represent a sample of the interactions between individuals, we used the

R package bisonR [63] to estimate uncertainty in the quantified edge weights based on sampling

effort. Explicitly incorporating uncertainty around the observed social network allows us to

account for how well the estimated network represents the true underlying latent network and

therefore confirm the robustness of our modelling results to sampling effects. For example,

typical network approaches would assign an edge weight of 0.5 both for a dyad that had been

seen together once and apart once as well for a dyad that had been seen together 100 times and

apart 100 times, despite the fact that our certainty about the edge weight of the second dyad is

much greater than the first [64]. By generating a distribution of possible networks from the

observed data rather than a single network, BISoN (Bayesian Inference of Social Networks)

allows us to account for this uncertainty, which can drastically affect the performance of

statistical models [64]. Specifically, for each group year we fitted a Bayesian 'edge model' with

a count conjugate prior to our observed network data, which returns a posterior distribution of

edge weights for each dyad in our network rather than a point estimate (for more information see

[64,65]). We extracted 1000 draws from this posterior distribution and used these draws as the

social networks over which we modelled pathogen spread. From each BISoN network we

calculated three different social centrality measures for each individual: strength (weighted sum

of an individual's social connections); closeness (the inverse of the mean weighted path length

from each individual to all others in the network); and degree centrality (the number of social

connections each individual has). All three measures were chosen because they have well-

established and important consequences for pathogen transmission [30,66–68], and degree and

closeness are known to decline with age in this system [25,69]. BISoN model outputs include

only non-zero edge weights because even though a particular dyad may never have been seen

interacting, BISoN naturally accounts for the possibility that those individuals may have

interacted in a future sample and so computes uncertainty for all edge weights [64]. Therefore, in

order to calculate degree centrality, we set a threshold at which individuals were deemed to have

a social connection or not. As females age, they lose their weakest social connections first. To

best capture this change, and ensure that our measure of degree was as independent of strength as

possible, we set our threshold using the minimum empirical observed non-zero edge weight for

each group-year (mean = 0.017; range = 0.008 - 0.027). We therefore calculated degree centrality

by counting all edges in each BISoN network that were equal to or greater than this minimum

edge weight in the observed network for each group-year.

Epidemiological model

Overview

We modelled SIS (susceptible-infected-susceptible) epidemiological dynamics to simulate the

spread of a hypothetical, directly-transmitted endemic pathogen by close contact through our

social networks (e.g., a respiratory virus). In our modelling framework individuals could either

be susceptible (S) or infected (I), thus we assumed that individuals retained no immunity from

previous infections. In a baseline ('control') version of the model, the probability of infection

depended on interaction strength in the grooming network (parameter: si; more grooming =

higher transmission probability), individuals were then infected for 5 time-steps (parameter: di)

and accumulated 1 unit cost per time step that they were infected (i.e., baseline per time step cost

- parameter: ci). While our per time step cost value was arbitrary, we were interested in the

relative differences between individuals rather than the absolute values of this parameter as our

goal was to compare infection cost between individuals of different ages and social centralities.

As a result, the exact value of this cost parameter was not important. In total, the model ran for

500 time-steps allowing individuals to be infected multiple times within each simulation. We

quantified their total cost of infection during a simulation. We then ran additional iterations of

the model where we allowed for immunosenescence in three different individual characteristics

(infection susceptibility, infection duration and infection severity) based on previous evidence

showing immune dysregulation and delayed response to infection in older rhesus macaques

([49,52,70]; see also Supplementary Methods). To examine how immunosenescence may impact

how changes in social centrality with age influence accumulated disease costs, we included

parameters that allowed us to increase susceptibility to infection (likelihood of acquiring

infection; parameter: *ai*), duration of infection (parameter: *adi*) and severity of infection (per

timestep cost of infection; parameter: *aci*) linearly with age. We simulated pathogen spread

across different combinations of these model parameterisations to provide data linking infection

cost to age and social network centrality.

Model details

We used a stochastic, discrete time implementation of the SIS model. Individuals could either be

susceptible (S) or infected (I). (The model code also includes a recovered state (R) but we set its

duration to 0 for this analysis so that individuals transitioned immediately from infected back to

susceptible). Individuals stayed in a susceptible state until they interacted with an infected

individual. For each interaction with an infected individual, a susceptible individual had a

probability of transitioning into the infected state of

$$P(tr)_{j,i} = (si + ai_i) \times A_{i,j}$$

where $P(tr)_{j,i}$ is the probability of transmission from individual *j* to individual *i*, *si* is a baseline

transmission probability, ai_i is the age effect on susceptibility to infection and $A_{i,j}$ is the

(undirected) edge weight between *i* and *j* in the grooming network. The ai_i term allowed us to

either assume that the probability of infection was homogeneous for every individual (when set

to a constant value) or that the probability of infection changed approximately linearly with age

so that immunosenescent individuals were more susceptible to infection. $A_{i,j}$ allowed the

probability of transmission to depend on how often pairs of animals groomed one another, with

this relationship assumed to be linear.

Once an individual is infected, the cost of each infection was calculated as

$$Cost = (di + adi_i) \times (ci + aci_i)$$

where di is the baseline duration of infection (set at 5 time steps for this analysis) before

returning to the susceptible (S) state, adi is the age effect on infection duration that allows a

linear increase in infection duration with age (to a maximum of 15 time steps), ci is the baseline

per time step cost of infection (set at 1 for this analysis) and aci is the age effect on infection

severity that allows a linear increase in infection severity per time step with age (to a maximum

of 3).

Model Parameterisation

We ran in total 24 combinations of parameters (24 models, Table S1). In all versions of the

model, we fixed the parameters $di = 5$ (baseline infection duration; considered representative of

a typical respiratory infection if a time step is considered to be a day [48,71]; see also

Supplementary Methods), $dr = 0$ (baseline duration of recovered period; set to zero to model

SIS epidemiological dynamics), $ci = 1$ (arbitrary). We also used a correction such that

$A(model)_{i,j} = A(data)_{i,j}{}^{0.7}$ to facilitate the calculation of the transmission probability per edge

by slightly increasing the importance of weak connections for transmission dynamics. We varied

the parameters *si* (baseline transmission probability), *ai* (age-based susceptibility to infection),

adi (age-based duration of infection) and *aci* (age-based severity of infection).

● We varied *si* to have low (0.45), medium (0.6) and high (0.75) transmissibility values.

These values were selected so that equilibrium pathogen prevalence equated to a basic

reproductive ratio (R_0) of approximately 1-1.5, 1.5-2 and 2-3 respectively (the R_0 varied

considerably between groups). R_0 was estimated using the approximation $R_0 = 1/(1-$

prevalence), where prevalence corresponded to the proportion of infected individuals in a

group.

● We varied *ai*, *adi* and *aci* to either be independent of age (no age effect) or increase

linearly with age (age effect). *adi* had a maximum value of 10 so that the duration of

infection was between 5 (for the youngest) and 15 (for the oldest) time steps, and *aci* had

a maximum value of 2 so that the per time step cost of infection was between 1 (for the

youngest) and 3 (for the oldest). All values of *adi* were rounded to the nearest whole

number.

Simulations contained all possible combinations of *ai*, *adi* and *aci* being 'on' (linear age effect)

or 'off' (no age effect) resulting in 8 possible parameterisations for each transmission

probability, and 24 parameter combinations in total (see Table S1).

Simulations

Each simulation run consisted of applying the epidemiological model to a given draw from a

BISoN generated posterior (a weighted network) for 500 time steps. We applied each

parametrisation of the model to each network that we generated - 1000 BISoN posterior

networks from each of the 23 group-years (23,000 networks x 24 parameterisations) - resulting

in 552,000 simulation runs in total. From each simulation run we calculated the total cost of

infection across the whole time period for each individual. Total individual infection costs were

collated together with individual data and measures of social network centrality (*see Social*

Network Construction) for subsequent analyses.

Analysis

We used linear mixed effects models (Gaussian error distribution) to summarise the outputs from

our epidemiological models (nb. we focus on effect size estimates rather than p-values here as

our aim is description rather than inference). We fitted three different linear mixed effects

models to understand the relationship between age and sociality on infection cost. For all models

we used the same random effect structure, fitting group-year, individual ID and combined group-

year and simulation number as random intercepts. We first fitted a model with age as the only

continuous fixed effect (model 1) to assess how infection cost changed with age under different

model parameterisations (i.e., combinations of transmission probabilities and

immunosenescence). We then fitted a model with age, degree, strength, and closeness as fixed

effects (model 2) to assess whether the effect of age on infection cost was in part being buffered

by social variation across ages. We standardized (z-scored) these three measures of social

centrality within group-year and included them as categorical variables in the model to improve

fit. We defined 5 categories for each social centrality measure (degree, strength and closeness) as

follows: < -1.5 = very low, -1.5 to -0.5 = low; -0.5 to 0.5 = average; 0.5 to 1.5 = high; >1.5 =

very high. The estimate of age in model 1 represents the effect of age on infection cost that

incorporates any effects of age-related variation in social centrality measures. The estimate of

age in model 2 represents the effect of age that occurs independently of the effects of degree,

strength, and closeness, and therefore represents the effect of age on infection cost in the

"absence" of social effects. This approach allowed us to calculate the overall protective effect of

social ageing on age-based infection cost by subtracting the age estimate in model 2 from the age

estimate in model 1. If this difference was negative, it would indicate that age-based infection

costs were lower when the social effects were not controlled for, indicating that age-based

variation in social centrality buffers the effects of age on infection cost and therefore has a

protective effect. Finally, we fitted a linear mixed effects model that included an interaction

between age (continuous) and each of the social centrality metrics (categorial) to better

understand how each of the different social metrics contributed to this overall protective effect

(model 3). Fitting an interaction term with each social metric allowed us to assess how the

effects of degree, strength, and closeness on infection cost varied with age.

Our first prediction (see Introduction) was that being less socially central at old ages

would be more beneficial in terms of reducing infection cost than at young ages. To test this

idea, we used the results from model 3 to calculate the reduction in infection cost that resulted

from having "average" rather than "high" social centrality when old (18 years old) versus when

young (8 years old). We chose these ages for our "young" and "old" categories as age 8 reflects

the lower quartile of data and age 18 is the median age of death for females that survive to

reproductive age in this population [20,47]. By including all three social metrics in model 3 we

could also determine their relative effect on infection risk and therefore assess prediction 2 - that

most of the decrease in infection cost associated with social ageing would be driven by older

individuals having lower degree and closeness. Finally, by applying models 1-3 across all

parameter combinations, we tested prediction 3 - that these protective effects of social ageing

would be stronger under conditions where there was greater immunosenescence with age. All

analyses were conducted in R version 4.3.1 and modelling was conducted in R version 4.1.1.

Figure 1. Predicted age and social centrality effects on accumulated infectious disease costs in our simulation results. Effect of age on infection cost at A) high transmissibility, C) medium transmissibility and E) low transmissibility. Across all transmissibilities, infection costs decrease slightly with age with no immunosenescence or immunosenescence only in infection susceptibility, but otherwise increase with age. Points in panel A, C, E represent a random sample of simulation data. Infection costs are higher for individuals with B) higher degree, D)

higher strength and F) higher closeness. Transmission probability is represented by point size in B, D, F. In all panels, colour represents the combination of immunosenescent effects included.

Results

Cost of infection increases with increased social centrality

On average, there were 51 females in each network (range: 19-73 females) and 169 grooming bouts per network (range: 33-392 bouts). Average network degree, strength and closeness centrality were 9.60 (range: 5.46 - 14.26), 0.62 (range: 0.40 - 0.88), 0.0008 (range: 0.0003 - 0.0025), respectively. All social centrality metrics were weakly to moderately correlated (see Table S2). Each of the three social centrality measures had independent, positive effects on infection cost under all parameter combinations (Fig. 1B,D,F). The effects of strength and closeness on disease cost were particularly strong (Fig. 1D,F), while the effects of degree were more moderate (Fig. 1B). These results confirmed that more socially central individuals suffer greater costs of infection.

In the absence of immunosenescence, lower infection cost in old age is mediated through lower social connectedness

Under conditions with no immunosenescence (no change in susceptibility (ai), severity (aci), or duration (adi) of infection with age), age had a negative effect on infection cost at all transmissibilities (high: β = -1.09 +/- 0.25; med: β = -0.86 +/- 0.23; low: β = -0.44 +/- 0.17; Fig. 1A,C,E). This means that, for example, at high transmissibility, each year increase in age results in a decrease in 1 cost unit, equating to a reduction of 22 cost units or 4.4 infections between the ages of 6 (our minimum age) and 28 (our maximum age). At the population level, all three sociality measures were negatively correlated with age in our data (degree: r = -0.11; strength: r = -0.13; closeness: r = -0.12; all p < 0.001; Fig. S2). When we included these social centrality measures in the model with age (model 2), the age estimate became less negative (high: β = -0.26 +/- 0.06; med: β = -0.07 +/- 0.05; low: β = 0.20 +/- 0.07) compared to the age estimate in a model with just age (model 1). This indicates that most of the variation that was explained by age in model 1, is now explained by sociality in model 2, suggesting that lower social centrality among older individuals is responsible for a substantial part of the reduction in infection cost with age. For example, at high transmissibility, the protective effect of our three measures of social centrality (i.e., the difference between the age estimate in models 1 and 2) was estimated

to be -0.83 (-1.09 - -0.26), meaning that age-based variation in strength, closeness and degree were expected to account for a reduction in 0.83 cost units per year of age, which translates to a reduction of about 3.7 infections between individuals of the minimum and maximum ages in our study (Fig. 2A). We found that this protective effect of social ageing was greatest at medium and high transmissibility and somewhat less prominent at low transmissibility (Fig. 2A).

Figure 2. The protective effect of social ageing, measured as the difference in the age estimate between model 1 (without social covariates) and model 2 (with social covariates). A more negative protective effect indicates that the effect of age on infection cost was lower when social centrality measures were not included in the model (model 1), compared to when they were included (model 2). Therefore, the stronger the negative effect, the more variation in infection cost was explained by age-related differences in sociality. Results show the protective effect of

428   *social ageing: A) across different transmission probabilities when there is no*

429   *immunosenescence, and B-D) across each (independent) form of immunosenescence (none,*

430   *susceptibility, severity, duration) at each transmission probability (high, medium, low).*

The protective effects of lower social connectedness in old age depend on epidemiological

characteristics

As expected, linear increases in susceptibility, severity, and duration of infection with age

led to infection cost increasing with age, although the strength of this effect depended on the

specific combination of parameters in the model (Fig. 1A,C,E). The one exception was when

there was only an increase in susceptibility with age, changes in infection cost were similar to

those when there was no immunosenescence (Fig. 1A,C,E). In line with prediction 3, the

protective effects associated with social ageing tended to get stronger when there was

immunosenescence, but this was not always true as it depended on the type of

immunosenescence (e.g., whether there were changes in susceptibility, severity, or duration of

infection with age) as well as the transmissibility of the pathogen (Fig. 2 & Fig. S3). The

protective effects of social ageing were greatest under conditions where there was an increase in

infection severity with age, across transmissibilities (Fig. 2B-D). For example, under high

transmissibility, when there was an increase in severity with age, the estimate for age in model 1

(just age) was 20.9 while the estimate for age in model 2 (age + sociality) was 22.1. The

protective effect was therefore -1.2 meaning that social ageing reduced infection cost by 1.2 units

for each year of increase in age. When there were age-based increases in susceptibility, on the

other hand, age-associated differences in sociality provided no additional protective effects

relative to baseline (i.e., no immunosenescence; Fig. 2B-D). When there were changes in

duration of infection with age, the protective effect of social ageing was more complex and

depended on the pathogen transmissibility. Relative to when there was no immunosenescence,

when transmissibility was low and duration of infection increased with age, social ageing

provided more of a protective effect. However, when transmissibility was medium or high and

duration increased with age, social ageing provided no additional protective effect compared to

baseline. Our results therefore suggest that age-based changes in sociality are likely to provide

the greatest benefit when there are increases in the severity or duration of infection with age, or a

combination thereof (for a full breakdown of the protective effects across all parameter

combinations see Fig. S3).

Old individuals benefit most from having lower strength and closeness

When considering interactions between age and each of the social metrics (model 3), the reduction in infection cost associated with lower social centrality in old age came primarily from having lower strength and closeness, which ran somewhat contrary to our expectations (see prediction 2 above). Strength and closeness showed clear sociality-age interactions whereby having "average" strength rather than "high" strength when old (18 years) resulted in a substantially greater decrease in infection cost than having "average" strength rather than "high" strength when young (8 years) (Fig. 3). For example, under high pathogen transmissibility and increased infection susceptibility and severity with age, having average rather than high strength when young decreased infection cost by 21.6 units (equivalent to 4.3 infections for a young individual), while having average rather than high strength when old decreased infection cost by almost double this amount (39.0 units; 7.8 infections for a young individual) (Fig. 3B,H). Similarly, having lower closeness when young reduced infection cost by 10.0 units (2 infections), while having lower closeness when old reduced infection cost by 19.6 units (3.9 infections) (Fig 3C,H). This supported our prediction that lower connectedness would be more beneficial at old than young ages (see prediction 1 above). However, contrary to our second prediction, lower degree had little to no protective effect, meaning that having "average" degree instead of "high" degree was no more beneficial when old than when young. Following on from the example above, lower degree when young decreased infection cost by 3.9 units and when old by 3.8 units (Fig. 3A,H). In fact, in some cases having fewer social partners for a given strength and closeness centrality seemed to reduce infection cost more at young ages than at old ages (Fig. 3F,H). In line with the results described above, the reduction in infection cost associated with having lower strength or closeness when older was dependent on pathogen transmissibility and on the type of immunosenescence. The reduction in infection cost at old ages compared to young ages was most pronounced when there were age-based increases in infection severity (Fig. 3F,H,J,K). This reduction was somewhat dampened when there were also changes in duration of infection with age (Fig. 3J,K), especially at high/medium transmissibility. When there were changes in infection duration with age but no changes in infection severity, lower values of closeness at old ages were no more beneficial than at young ages, except when pathogen transmissibility was low (Fig. 3G,I). Under this parameter combination, having lower strength was actually more beneficial at young ages than at old ages (Fig. 3G,I). Generally, these findings

align well with the results described above, suggesting that having lower values of social

centrality when old is less protective when there are increases in infection duration with age and

transmissibility is high. We explored the robustness of these patterns by also looking at the

difference in infection cost when an individual had "average" social centrality versus "low"

social centrality and comparing this at young and old ages. We found the patterns highly

comparable to those described above (see Fig. S4).

Figure 3. *An illustration of how social network centrality and age interact to determine*

accumulated infection costs. A-C) Predicted age effects on infection costs separated by social

centrality category for A) degree, B) strength and C) closeness centrality. Results shown for a

specific parameter combination where pathogen transmissibility is high and infection

susceptibility and severity show linear increases with age. Points represent a random sample of

simulation data. D-K) The change in infection cost for an individual moving from high social

centrality to average social centrality when old (18 years; blue points) versus when young (8

years old; green points), for each social centrality measure. Different panels (labelled) show

results for different combinations of immunosenescence and point size represents transmission

probability. Similar results for a change from average to low centrality are shown in the

Supplementary Materials.

Discussion

Our model results suggest that social ageing in rhesus macaques (i.e., lower social

centrality in older individuals) is associated with reduced costs accrued from socially transmitted

infections. We found that infectious disease cost was strongly positively associated with the

network centrality of an individual, as would be expected. Specifically, we found that strength

and closeness centrality had strong, independent effects on the overall cost of infection, while the

effects of degree on infection cost were more moderate. Given the probability of infection in our

model depends on edge weight, the importance of strength on overall cost of infection was

expected but nevertheless reflects the widespread importance of strength in explaining infection

risk in free-living populations [30]. Under conditions with no immunosenescence (no increases

in infection susceptibility, severity, or duration with age) older individuals accrued a lower cost

of infection compared to younger individuals across all transmissibilities modelled, with the

benefits peaking for pathogens with intermediate-high transmission probabilities. These lower

costs of infection were driven by age-associated differences in social centrality, as all three social

centrality metrics were negatively correlated with age and including centrality measures within

the same model resulted in the protective effect largely disappearing. When we included (linear)

immunosenescence we were able to show that having lower social centrality had a much greater

benefit for older than younger individuals, with this effect strongest when the severity of

infection increased with age.

The protective effect of lower social centrality on infection risk in older individuals

We found that older individuals accrued lower disease costs than younger individuals in

the absence of immunosenescence. Given previously documented within-individual declines in

closeness centrality and degree with age [21,25], we anticipated that this protective effect would

be driven by age-based differences in social centrality, which our results support. Our analyses,

however, were not explicitly longitudinal, meaning that we cannot fully rule out selective

disappearance of more central individuals partly explaining these results. It is possible that the

high infectious disease risk associated with being more socially central leads these individuals to

die earlier. However, given that social centrality is associated with survival in this population

[8,20,42], it seems likely that longitudinal changes in social network position reduce older

individual's exposure to infection. This is likely to be particularly true for degree and closeness

for which we have clear evidence that population level differences are driven by within-

individual changes with age [21,25]. The fact that strength was negatively correlated with age

and contributed substantially to the protective effect provided by social ageing was surprising

given that previous analyses in this study system have found a non-significant (although weakly

negative) relationship between age and strength both within and between individuals [21].

However, it is possible that, combined, this weakly negative relationship at both the within and

between-individual level results in a more substantial population-level decline in strength with

age, which we detect in our analyses. It is also possible that the inclusion of more data than

previous analyses has enabled us to detect change in strength with age not previously evident.

Overall, we found the protective effect of social ageing to be strongest for pathogens with

intermediate-high transmissibility and weaker for less transmissible pathogens. That is, the

reduction in infection costs with age (in the absence of immunosenescence) was highest when

transmissibility was high and pathogen prevalence (i.e., proportion of infected individuals in the

group) within groups was therefore relatively high (>30% or more), with a weaker reduction

when transmissibility, and therefore pathogen prevalence, in groups was low. There are a variety

of pathogens that are transmitted via direct contact, and which vary in their level of

transmissibility, which could be represented by our model. For example, Shigella (a bacterial

pathogen) is transmissible by close contact in macaques [62] and has spread very rapidly through

the Cayo Santiago macaque population previously [72]. Additionally, macaques are known to be

infected by a range of respiratory viruses (e.g, coronaviruses, influenza viruses), which will

likely fall across the range of transmissibilities considered in our study (see Supplementary

Methods), suggesting our findings are likely to be highly relevant for real-world pathogens.

Immunosenescence enhances the protective effect of lower social centrality in older individuals

When adding immunosenescence to our models, we found that the protective effects of social ageing were generally stronger than under conditions with no immunosenescence. Overall, we found that lower levels of social centrality at old ages led to the most substantial reductions in infection costs when infection severity increased with age. If we instead considered conditions where infection duration increased with age, then the protective effects of social ageing depended on pathogen transmissibility. That is, lower connectivity at older ages was more important for decreasing disease costs at low compared to high transmissibility when infection duration increased with age. These results can be explained intuitively based on the epidemiological dynamics of high versus low pathogen transmissibility within an SIS model. When transmissibility, and therefore (in our model) prevalence is very high, individuals will typically be re-infected very quickly once they recover [73], meaning that the duration of each infection is less important to overall disease cost. However, when transmissibility, and therefore prevalence, is low it may take a while for an individual to be re-infected once it recovers from infection. As a result of this reduced frequency of infection, the fact that each infection lasts longer is disproportionately costly. Therefore, having lower social centrality when old relative to when young under this low transmissibility scenario has a more substantial protective effect. When we looked at conditions where infection susceptibility increased with age, we found that older individuals gained very little protective effect from having lower social connectivity relative to younger individuals.

It should be noted that comparisons of the relative importance of age-related changes in susceptibility, severity and duration for disease cost are limited somewhat by how their relative effects match (e.g., infection severity can be 3 times higher for our oldest compared with our youngest individual, while the difference caused by changes in susceptibility is between 1.33 and 1.56 times depending on the baseline transmission probability). These differences arise because parameters were chosen to be reflective of a reasonable parameter space for many respiratory infections rather than specifically for comparisons of relative effects. Despite this limitation, we can still draw some general conclusions. When pathogen transmissibility is high, most of the age-based differences in accumulated disease cost are driven by increases in the severity of infection with age, while increases in infection susceptibility or duration with age increase age-

based infection cost only minimally. When pathogen transmissibility is lower, age-related

changes in infection duration become relatively more important for age-based infection costs,

although changes to severity still dominate. Generally, this suggests that age-based differences in

social centrality are most likely when there are increases in severity of infection with age. In the

case of a pathogen with low transmissibility, increases in duration of infection with age might

also result in social declines.

Social trade-offs in old age and the role of infectious disease

The measures of social centrality that were most important for cost of infection in older

individuals were not exclusively those known to change within-individuals as they age,

indicating that there may be potentially important social trade-offs. We found that, in general,

older individuals would benefit most from having lower strength followed by lower closeness

centrality, with lower values of degree having smaller and less consistent effects on infection

risk. When we contrast this with observed within-individual changes in social centrality [21,25],

it is noteworthy that individuals in this population show behavioural changes with age that

reduce both their degree and closeness centrality but not their strength. Although here we have

shown that strength is negatively correlated with age it remains unclear whether this negative

relationship is driven by within-individual changes or between-individual differences because of

selective disappearance or differences between cohorts. Strength to top partners has a positive

link to health and survival in rhesus macaques [42] and diverse group-living species [1].

Therefore, it seems possible that while having lower strength may be the most effective way for

older animals to cut infectious disease risk, the costs of doing so may be high, and preserving

strong social relationships may itself be an important form of social buffering that can protect

against infectious disease ([74]; see Limitations section below). By maintaining strong social

connections and avoiding interactions with less familiar or new social partners [21,25]

individuals can instead reduce other aspects of their social centrality (e.g., closeness) that also

reduce infectious disease costs but which are less strongly associated with other aspects of health

and fitness. In this way, older individuals may still reap the benefits of social relationships while

minimizing the risks of infection [35].

While, to date, most research on social ageing has focused on age-based differences in

direct connectedness (cf. [23,25,26]), measures of "flow" through a network, as captured through

indirect metrics, can also be highly relevant for pathogen transmission [30]. Our previous work

has shown that by changing simple behavioural rules (in this case, reassociating with the same

partners and mixing less widely with the broader network) ageing individuals can facilitate

changes in both their direct connectedness (i.e. degree) and their indirect connectedness (i.e.

closeness) [25]. Here we have seen that it is the effects of these age-based behavioural changes

on closeness which are particularly beneficial, relative to the effects on degree, for mitigating

infection risk. Furthering our understanding of the intersection between social ageing and

infectious disease therefore necessitates deepening our understanding of how behavioural

changes with age facilitate changes not only in direct connectedness but also changes in

connectedness to the wider network.

Potential implications for the evolution of social ageing

638   While our results suggest that individuals could benefit from reduced infectious disease

risk by reducing their social networks in old age [21,25], more work is required before we can

say that social ageing is adaptive. While, it is appealing to consider that social ageing might be

an evolved strategy to counteract declines in immunity and associated increases in disease

burden in later life, the strength of selection on a trait will typically decline with age [75,76].

This has two implications for the evolution of social ageing in response to infectious disease risk.

First, it means that when social network centrality influences fitness we would expect social

behaviour to senesce as a result of weakening selection in later life [75], independent of any late-

life benefits of reduced social centrality. This expected decline could even be enhanced in rhesus

macaques where positive effects of social relationships on survival are reduced for older

individuals [20]. Second, under most conditions the declining strength of selection with age

makes it less likely that social ageing could have evolved to reduce the costs attributed to

infection late in life when immunosenescence increases the risk and severity of infections [35].

However, substantial immunosenescence at ages where individuals still have some reproductive

value could cause infectious disease to influence social ageing. It may be in these cases that

observed patterns of social ageing are a plastic response that can be selected for to mitigate rapid

declines in immune performance with age. Additionally, predictions related to the strength and

direction of demographic selection at old ages can depend on assumptions about density-

dependent population regulation [77–79], and it is not immediately clear where mortality related

to infectious disease fits in this context. Therefore, any selection acting on age-related changes in

social behaviour are likely to be highly context dependent and may change depending on which

aspects of sociality predict fitness earlier in life too. Expanding theory to integrate age-related

changes in social behaviour within existing evolutionary models of age-dependent mortality and

senescence will be key to better understanding the role of infectious disease in social ageing.

Limitations

There are some limitations to our model that should be considered when interpreting our

results and that offer fruitful choices for future work. First, we have considered only directly

transmitted pathogens and focused on SIS (susceptible-infected-susceptible) epidemiological

dynamics. These methodological choices make sense. Directly transmitted pathogens are

widespread threats to human and non-human animal health [80] and their epidemiological

dynamics will be most strongly influenced by social interaction patterns [81]. However,

incorporating indirectly transmitted pathogens may be important if co-infection affects morbidity

[82,83]. We focused on SIS dynamics as this provides a convenient way to model endemic

disease without explicitly incorporating demography or immune dynamics. Future work could

build on ours by testing how the choice of disease model can affect results, or by developing

long-term integrated network-demographic models of disease.

We have also assumed a linear relationship between interaction duration and the

probability of infection. However, for some pathogens with high transmissibility almost any

social contact may be sufficient, while for others only prolonged interactions may allow the

pathogen to spread. These differences would likely mean that the impact of social ageing on

pathogen transmission will depend on the shape of the dose-response curve. For example, we

might predict that social ageing is ineffective against mitigating disease risks for pathogens that

can spread easily via even short duration proximity. While social relationships are known to

expose individuals to disease risk, they can simultaneously help individuals cope with infections

once acquired [74,84]. We have not explicitly modeled these 'social support effects', which we

may expect to play a role in macaques [74]. In general, we would expect the greatest social

support effects from an individual's strongest social relationships [42,85,86], which may amplify

any protective effects of social ageing against disease as long these types of relationships are

maintained [21,27]. Existing models have incorporated social support effects into network

models of infectious disease spread [87], and using them explicitly in this context could provide

additional insight.

Further, while we have focused here on how differences in social behaviour can affect

infection risk we have not considered how the spread of infection can also influence social

behaviour and network structure. In some cases, sickness behaviour and social avoidance of

infected individuals can change an individual's social interactions (reviewed in [88,89])

influencing the structure of the network [90]. Given older individuals tend to be more susceptible

to pathogens, lower connectedness with age could be a consequence of, rather than a proactive

response to, infectious disease risk. However, this seems an unlikely explanation of social ageing

in our system given evidence that older females appear to be more selective in their partner

choice rather than arbitrarily decreasing their connectedness as might be expected with sickness

behaviour or being avoided by other individuals [69]). Alternatively, if social selectivity is

selected for by infectious disease costs (see caveats above), then older individuals might actually

be hyper-responsive to cues of infection, leading to greater infection avoidance behaviour with

age. Older individuals' may also be better able to detect infections in the small number of

individuals they know well and this could enhance avoidance strategies [35]. Ultimately, how

social ageing fits into the co-dynamics of social behaviour and infectious disease spread is likely

to be complex and warrants further research.

Conclusions

We used epidemiological models to demonstrate that reduced social connectedness in old age

has the potential to provide a protective effect against the accrued costs of endemic infectious

diseases in a free-ranging population of rhesus macaques. By considering the impacts of different

forms of immunosenescence we showed that the benefits of social ageing could vary

considerably depending on the interaction between pathogen traits (transmissibility) and how

changes to an individual's immune performance manifest in terms of susceptibility, severity and

duration of infection. In addition, the aspects of social centrality that most impacted disease costs

for older individuals were not exclusively those that we know change within-individuals as they

age, highlighting the trade-offs inherent to interacting with others. Although we focused in this

paper on how age-based changes in susceptibility to infectious disease might facilitate social

ageing, the other way that immunosenescence might affect age-based changes in sociality is

through reduced healing ability. Being less able to recover from wounds might impose greater

social costs at old ages leading individuals to alter their social centrality to avoid competitive

interactions [12]. Generally, understanding how immunosenescence intersects with age-based

variation in sociality remains an open and intriguing question. Our results demonstrate the clear

potential for infectious disease to influence social ageing and point towards the value of

developing new theoretical models that consider the evolutionary dynamics involved.

Acknowledgements Thank you to J. Firth, G. Albery, S. Bouwhuis and R. Salguero-Gomez for

organizing this special issue. We are grateful to the Caribbean Primate Research Center for

maintaining the Cayo Santiago population and for access to the study site, and all the field

technicians who have contributed to the long-term behavioural database over the years. Thank

you to S. Ellis, D. Nussey, and J. Moorad for thoughtful discussion during the manuscript's

development. This work was supported by the following grants from the National Institute of

Health (NIH): grant nos R01-AG060931, R00-AG051764, R01-MH096875, R37-MH109728,

R01-MH108627, R01-MH118203, U01MH121260, R01-NS123054 and the Kaufman

Foundation: grant no KA2019-105548. The Cayo Santiago Field Station is supported by the

Office of Research Infrastructure Programs of the NIH (2P40OD012217). MJS is supported by a

Royal Society University Research Fellowship URF\R1\221800.

References

1.   Snyder-Mackler N *et al.* 2020 Social determinants of health and survival in humans and
other animals. *Science* **368**. (doi:10.1126/science.aax9553)

2.   Feldblum JT, Krupenye C, Bray J, Pusey AE, Gilby IC. 2021 Social bonds provide multiple
pathways to reproductive success in wild male chimpanzees. *iScience* **24**, 102864.
(https://doi.org/10.1016/j.isci.2021.102864)

3.   Vander Wal E, Festa-Bianchet M, Réale D, Coltman DW, Pelletier F. 2015 Sex-based
differences in the adaptive value of social behavior contrasted against morphology and
environment. *Ecology* **96**, 631–641.

4.   Siracusa ER, Boutin S, Dantzer B, Lane JE, Coltman DW, McAdam AG. 2021 Familiar
Neighbors, but Not Relatives, Enhance Fitness in a Territorial Mammal. *Curr. Biol.* **31**,
438–445.e3.

5. Cameron EZ, Setsaas TH, Linklater WL. 2009 Social bonds between unrelated females increase reproductive success in feral horses. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 13850–13853.

6. Stockley P, Bro-Jørgensen J. 2011 Female competition and its evolutionary consequences in mammals. *Biol. Rev. Camb. Philos. Soc.* **86**, 341–366.

7. Clutton-Brock T, Huchard E. 2013 Social competition and its consequences in female mammals. *J. Zool.* **289**, 151–171.

8. Pavez-Fox MA *et al.* 2022 Reduced injury risk links sociality to survival in a group-living primate. *iScience*. **12**, 105454. https://doi.org/10.1016/j.isci.2022.105454

9. Sah P, Mann J, Bansal S. 2018 Disease implications of animal social network structure: A synthesis across social systems. *J. Anim. Ecol.* **87**, 546–558.

10. Altizer S *et al.* 2003 Social Organization and Parasite Risk in Mammals: Integrating Theory and Empirical Studies. *Annu. Rev. Ecol. Evol. Syst.* **34**, 517–547.

11. Alexander RD. 1974 The Evolution of Social Behavior. *Annu. Rev. Ecol. Syst.* **5**, 325–383.

12. Siracusa ER, Higham JP, Snyder-Mackler N, Brent LJN. 2022 Social ageing: exploring the drivers of late-life changes in social behaviour in mammals. *Biol. Lett.* **18**, 20210643.

13. Albery GF, Clutton-Brock TH, Morris A, Morris S, Pemberton JM, Nussey DH, Firth JA. 2022 Ageing red deer alter their spatial behaviour and become less social. *Nat Ecol Evol* **6**, 1231–1238.

14. Machanda ZP, Rosati AG. 2020 Shifting sociality during primate ageing. *Philosophical Transactions of the Royal Society B: Biological Sciences*. **375**, 20190620. (doi:10.1098/rstb.2019.0620)

15. Wrzus C, Hänel M, Wagner J, Neyer FJ. 2013 Social network changes and life events across the life span: a meta-analysis. *Psychol. Bull.* **139**, 53–80.

16. Guan X, Dluzen DE. 1994 Age related changes of social memory/recognition in male Fischer 344 rats. *Behav. Brain Res.* **61**, 87–90.

17. Ryu H, Graham KE, Sakamaki T, Furuichi T. 2016 Long-sightedness in old wild bonobos during grooming. *Curr. Biol.* **26**, R1131–R1132.

18. Croft DP, Weiss MN, Nielsen MLK, Grimes C, Cant MA, Ellis S, Franks DW, Johnstone RA. 2021 Kinship dynamics: patterns and consequences of changes in local relatedness. *Proc. Biol. Sci.* **288**, 20211129.

19. Rodrigues AMM. 2018 Demography, life history and the evolution of age-dependent social behaviour. *Journal of Evolutionary Biology*. **31**, 1340–1353. (doi:10.1111/jeb.13308)

20. Brent LJN, Ruiz-Lambides A, Platt ML. 2017 Family network size and survival across the
lifespan of female macaques. *Proc. Biol. Sci.* **284**. (doi:10.1098/rspb.2017.0515)

21. Siracusa ER, Negron-Del Valle JE, Phillips D, Platt ML, Higham JP, Snyder-Mackler N,
Brent LJN. 2022 Within-individual changes reveal increasing social selectivity with age in
rhesus macaques. *bioRxiv.* , 2022.05.31.494118. (doi:10.1101/2022.05.31.494118)

22. Almeling L, Hammerschmidt K, Sennhenn-Reulen H, Freund AM, Fischer J. 2016
Motivational Shifts in Aging Monkeys and the Origins of Social Selectivity. *Curr. Biol.* **26**,
1744–1749.

23. Rathke E-M, Fischer J. 2021 Social aging in male and female Barbary macaques. *Am. J.
Primatol.* , e23272.

24. Schino G, Pinzaglia M. 2018 Age-related changes in the social behavior of tufted capuchin
monkeys. *Am. J. Primatol.* **80**, e22746.

25. Siracusa ER *et al.* 2023 Ageing in a collective: the impact of ageing individuals on social
network structure. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **378**, 20220061.

26. Kroeger SB, Blumstein DT, Martin JGA. 2021 How social behaviour and life-history traits
change with age and in the year prior to death in female yellow-bellied marmots. *Philos.
Trans. R. Soc. Lond. B Biol. Sci.* **376**, 20190745.

27. Lang FR, Carstensen LL. 1994 Close emotional relationships in late life: further support for
proactive aging in the social domain. *Psychol. Aging* **9**, 315–324.

28. Rosati AG, Hagberg L, Enigk DK, Otali E, Emery Thompson M, Muller MN, Wrangham
RW, Machanda ZP. 2020 Social selectivity in aging wild chimpanzees. *Science* **370**, 473–
476.

29. Kappeler PM, Cremer S, Nunn CL. 2015 Sociality and health: impacts of sociality on
disease susceptibility and transmission in animal and human societies. *Philos. Trans. R.
Soc. Lond. B Biol. Sci.* **370**. (doi:10.1098/rstb.2014.0116)

30. Briard L, Ezenwa VO. 2021 Parasitism and host social behaviour: a meta-analysis of
insights derived from social network analysis. *Anim. Behav.* **172**, 171–182.

31. Lucatelli J, Mariano-Neto E, Japyassú HF. 2021 Social interaction, and not group size,
predicts parasite burden in mammals. *Evol. Ecol.* **35**, 115–130.

32. Peters A, Delhey K, Nakagawa S, Aulsebrook A, Verhulst S. 2019 Immunosenescence in
wild animals: meta-analysis and outlook. *Ecol. Lett.* **22**, 1709–1722.

33. Kang SJ, Jung SI. 2020 Age-Related Morbidity and Mortality among Patients with COVID-
19. *Infect Chemother* **52**, 154–164.

34. Palacios MG, Winkler DW, Klasing KC, Hasselquist D, Vleck CM. 2011 Consequences of

immune system aging in nature: a study of immunosenescence costs in free-living Tree
Swallows. *Ecology* **92**, 952–966.

35. Albery GF, Sweeny AR, Webber Q. 2023 How behavioural ageing affects infectious
disease. *Neurosci. Biobehav. Rev.* **155**, 105426.

36. Ezenwa VO, Worsley-Tonks KEL. 2018 Social living simultaneously increases infection
risk and decreases the cost of infection. *Proc. Biol. Sci.* **285**. (doi:10.1098/rspb.2018.2142)

37. Romano V, MacIntosh AJJ, Sueur C. 2020 Stemming the Flow: Information, Infection, and
Social Evolution. *Trends Ecol. Evol.* **35**, 849–853.

38. Hinde RA. 1976 Interactions, Relationships and Social Structure. *Man* **11**, 1–17.

39. Craft ME, Caillaud D. 2011 Network models: an underutilized tool in wildlife
epidemiology? *Interdiscip. Perspect. Infect. Dis.* **2011**, 676949.

40. Keeling MJ, Eames KTD. 2005 Networks and epidemic models. *J. R. Soc. Interface* **2**, 295–
307.

41. Cooper EB, Brent LJN, Snyder-Mackler N, Singh M, Sengupta A, Khatiwada S,
Malaivijitnond S, Qi Hai Z, Higham JP. 2022 The rhesus macaque as a success story of the
Anthropocene. *Elife* **11**. (doi:10.7554/eLife.78169)

42. Ellis S, Snyder-Mackler N, Ruiz-Lambides A, Platt ML, Brent LJN. 2019 Deconstructing
sociality: the types of social connections that predict longevity in a group-living primate.
Proc. Biol. Sci. **286**, 20191991.

43. Brent LJN, Heilbronner SR, Horvath JE, Gonzalez-Martinez J, Ruiz-Lambides A, Robinson
AG, Skene JHP, Platt ML. 2013 Genetic origins of social networks in rhesus macaques. *Sci.*
Rep. **3**, 1042.

44. Rawlins RG, Kessler MJ. 1986 *The Cayo Santiago Macaques: History, Behavior, and*
Biology. SUNY Press.

45. Roth GS, Mattison JA, Ottinger MA, Chachich ME, Lane MA, Ingram DK. 2004 Aging in
rhesus monkeys: relevance to human health interventions. *Science* **305**, 1423–1426.

46. Shively CA, Register TC, Clarkson TB. 2009 Social stress, visceral obesity, and coronary
artery atherosclerosis: product of a primate adaptation. *Am. J. Primatol.* **71**, 742–751.

47. Chiou KL *et al.* 2020 Rhesus macaques as a tractable physiological model of human ageing.
Philos. Trans. R. Soc. Lond. B Biol. Sci. **375**, 20190612.

48. Yu P *et al.* 2020 Age-related rhesus macaque models of COVID-19. *Animal Model Exp*
Med **3**, 93–97.

49. Speranza E *et al.* 2022 Age-related differences in immune dynamics during SARS-CoV-2

infection in rhesus macaques. *Life Sci Alliance* **5**. (doi:10.26508/lsa.202101314)

50. Didier ES, Sugimoto C, Bowers LC, Khan IA, Kuroda MJ. 2012 Immune correlates of
aging in outdoor-housed captive rhesus macaques (Macaca mulatta). *Immun. Ageing* **9**, 25.

51. Messaoudi I, Estep R, Robinson B, Wong SW. 2011 Nonhuman primate models of human
immunology. *Antioxid. Redox Signal.* **14**, 261–273.

52. Rosado MRS *et al.* 2023 Immune cell composition varies by age, sex and exposure to social
adversity in free-ranging Rhesus Macaques. *Geroscience* (doi:10.1007/s11357-023-00962-
8)

53. Salguero FJ *et al.* 2021 Comparison of rhesus and cynomolgus macaques as an infection
model for COVID-19. *Nat. Commun.* **12**, 1260.

54. Munster VJ *et al.* 2020 Respiratory disease in rhesus macaques inoculated with SARS-
CoV-2. *Nature* **585**, 268–272.

55. Pol M van de, van de Pol M, Verhulst S. 2006 Age☐Dependent Traits: A New Statistical
Model to Separate Within☐ and Between☐Individual Effects. *The American Naturalist.*
167, 766–773. (doi:10.1086/503331)

56. Van de Pol M, Wright J. 2009 A simple method for distinguishing within-versus between-
subject effects using mixed models. *Anim. Behav.* **77**, 753.

57. Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in wild
meerkats. *Proc. Biol. Sci.* **277**, 633–642.

58. MacIntosh AJJ, Hernandez AD, Huffman MA. 2010 Host age, sex, and reproductive
seasonality affect nematode parasitism in wild Japanese macaques. *Primates* **51**, 353–364.

59. Müller-Klein N, Heistermann M, Strube C, Franz M, Schülke O, Ostner J. 2019 Exposure
and susceptibility drive reinfection with gastrointestinal parasites in a social primate. *Funct.*
Ecol. **33**, 1088–1098.

60. Rimbach R, Bisanzio D, Galvis N, Link A, Di Fiore A, Gillespie TR. 2015 Brown spider
monkeys (Ateles hybridus): a model for differentiating the role of social networks and
physical contact on parasite transmission dynamics. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*
370. (doi:10.1098/rstb.2014.0110)

61. Habig B, Jansen DAWAM, Akinyi MY, Gesquiere LR, Alberts SC, Archie EA. 2019
Multi-scale predictors of parasite risk in wild male savanna baboons (Papio cynocephalus).
Behav. Ecol. Sociobiol. **73**, 134.

62. Balasubramaniam KN, Beisner BA, Hubbard JA, Vandeleest JJ, Atwill ER, McCowan B.
2019 Affiliation and disease risk: social networks mediate gut microbial transmission
among rhesus macaques. *Anim. Behav.* **151**, 131–143.

63.   Hart J. 2023 *bisonR: Bayesian Inference of Social Networks in R.*
https://jhart96.github.io/bisonR, https://github.com/JHart96/bisonR.

64.   Hart J, Weiss MN, Franks D, Brent L. 2023 BISoN: A Bayesian framework for inference of
social networks. *Methods Ecol. Evol.* **14**, 2411–2420.

65.   Hart JDA, Franks DW, Brent L, Weiss MN. 2022 BisonR - Bayesian inference of social
networks with R. (doi:10.31219/osf.io/ywu7j)

66.   Williams AE, Worsley-Tonks KEL, Ezenwa VO. 2017 Drivers and consequences of
variation in individual social connectivity. *Anim. Behav.* **133**, 1–9.

67.   Grear DA, Luong LT, Hudson PJ. 2013 Network transmission inference: host behavior and
parasite life cycle make social networks meaningful in disease ecology. *Ecol. Appl.* **23**,
1906–1914.

68.   Christley RM, Pinchbeck GL, Bowers RG, Clancy D, French NP, Bennett R, Turner J. 2005
Infection in social networks: using network analysis to identify high-risk individuals. *Am. J.*
Epidemiol. **162**, 1024–1031.

69.   Siracusa ER, Negron-Del Valle JE, Phillips D, Platt ML, Higham JP, Snyder-Mackler N,
Brent LJN. 2022 Within-individual changes reveal increasing social selectivity with age in
rhesus macaques. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2209180119.

70.   Coe CL, Lubach GR, Kinnard J. 2012 Immune senescence in old and very old rhesus
monkeys: reduced antibody response to influenza vaccination. *Age* **34**, 1169–1177.

71.   Nelson CE *et al.* 2022 Mild SARS-CoV-2 infection in rhesus macaques is associated with
viral control prior to antigen-specific T cell responses in tissues. *Sci Immunol* , eabo0535.

72.   Testard C *et al.* 2021 Rhesus macaques build new social connections after a natural disaster.
Curr. Biol. **31**, 2299–2309.e7.

73.   Parshani R, Carmi S, Havlin S. 2010 Epidemic threshold for the susceptible-infectious-
susceptible model on random networks. *Phys. Rev. Lett.* **104**, 258701.

74.   Balasubramaniam K, Beisner B, Vandeleest J, Atwill E, McCowan B. 2016 Social buffering
and contact transmission: network connections have beneficial and detrimental effects on
Shigella infection risk among captive rhesus macaques. *PeerJ* **4**, e2630.

75.   Hamilton WD. 1966 The moulding of senescence by natural selection. *J. Theor. Biol.* **12**,
12–45.

76.   Williams GC. 1957 Pleiotropy, Natural Selection, and the Evolution of Senescence.
Evolution **11**, 398–411.

77.   Abrams PA. 1993 DOES INCREASED MORTALITY FAVOR THE EVOLUTION OF
MORE RAPID SENESCENCE? *Evolution* **47**, 877–887.

78. Moorad J, Promislow D, Silvertown J. 2019 Evolutionary Ecology of Senescence and a
Reassessment of Williams' 'Extrinsic Mortality' Hypothesis. *Trends Ecol. Evol.* **34**, 519–
530.

79. de Vries C, Galipaud M, Kokko H. 2023 Extrinsic mortality and senescence: a guide for the
perplexed. *Peer Community Journal* **3**. (doi:10.24072/pcjournal.253)

80. Daszak P, Cunningham AA, Hyatt AD. 2000 Emerging infectious diseases of wildlife--
threats to biodiversity and human health. *Science* **287**, 443–449.

81. White LA, Forester JD, Craft ME. 2017 Using contact networks to explore mechanisms of
parasite transmission in wildlife. *Biol. Rev. Camb. Philos. Soc.* **92**, 389–409.

82. Griffiths EC, Pedersen AB, Fenton A, Petchey OL. 2011 The nature and consequences of
coinfection in humans. *J. Infect.* **63**, 200–206.

83. Risco D *et al.* 2014 Severity of bovine tuberculosis is associated with co-infection with
common pathogens in wild boar. *PLoS One* **9**, e110123.

84. Akinyi MY, Tung J, Jeneby M, Patel NB, Altmann J, Alberts SC. 2013 Role of Grooming
in Reducing Tick Load in Wild Baboons (Papio cynocephalus). *Anim. Behav.* **85**, 559–568.

85. Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM,
Seyfarth RM, Cheney DL. 2010 Strong and consistent social bonds enhance the longevity
of female baboons. *Curr. Biol.* **20**, 1359–1361.

86. Schülke O, Bhagavatula J, Vigilant L, Ostner J. 2010 Social bonds enhance reproductive
success in male macaques. *Curr. Biol.* **20**, 2207–2210.

87. Chen X, Wang R, Tang M, Cai S, Eugene Stanley H, Braunstein LA. 2018 Suppressing
epidemic spreading in multiplex networks with social-support. *New J. Phys.* **20**, 013007.

88. Stockmaier S, Stroeymeyt N, Shattuck EC, Hawley DM, Meyers LA, Bolnick DI. 2021
Infectious diseases and social distancing in nature. *Science* **371**.
(doi:10.1126/science.abc8881)

89. Townsend AK, Hawley DM, Stephenson JF, Williams KEG. 2020 Emerging infectious
disease and the challenges of social distancing in human and non-human animals. *Proc.*
Biol. Sci. **287**, 20201039.

90. Stroeymeyt N, Grasse AV, Crespi A, Mersch DP, Cremer S, Keller L. 2018 Social network
plasticity decreases disease transmission in a eusocial insect. *Science* **362**, 941–945.