Navigated range expansion promotes migratory culling

Yi Zhang[1], Qingjuan Hu[1], Yingtong Su[1], Pan Chu[1,2], Ting Wei[1], Xuefei Li[1,2], Chenli Liu[1,2], Xiongfei
Fu[1,2*]

[1]Key Laboratory for Quantitative Synthetic Biology, Shenzhen Institute of Synthetic Biology,
Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055,
China

[2]University of Chinese Academy of Sciences, Beijing 100049, China

*Corresponding author. Email: xiongfei.fu@siat.ac.cn

Author Contributions: Yi Zhang: Data curation; investigation; methodology; writing-original
draft; writing-review and editing. **Qingjuan Hu:** Data curation; investigation. **Yingtong Su:** Data
curation; investigation; writing-original draft. **Pan Chu:** Data curation. **Ting Wei:** Methodology;
writing-review and editing. **Xuefei Li:** Writing-review and editing. **Chenli Liu:** Methodology;
writing-review and editing. **Xiongfei Fu:** Conceptualization; supervision; methodology; writing-
original draft; project administration; writing-review and editing.

Competing Interest Statement: The authors declare no conflict of interest.

Keywords: Range expansion; Pattern formation; Synthetic biology; Migratory culling.

Abstract

Motile organisms can expand into new territories and increase their fitness, while nonmotile
viruses usually depend on host migration to spread across long distances. In general, faster host
motility facilitates virus transmission. However, recent ecological studies have also shown that
animal host migration can reduce viral prevalence by removing infected individuals from the
migratory group. Here, we use a bacteria-bacteriophage co-propagation system to investigate
how host motility affects viral spread during range expansion. We find that phage spread during
chemotaxis-driven navigated range expansion decreases as bacterial migration speed increases.
Theoretical and experimental analyses show that the navigated migration leads to a spatial
sorting of infected and uninfected hosts in the co-propagating front of bacteria-bacteriophage,
with implications for the number of cells left behind. The preferential loss of infected cells in the
co-propagating front inhibits viral spread. Further increase in host migration speed leads to a
phase transition that eliminates the phage completely. These results illustrate that navigated
range expansion of the host can promote the migratory culling of infectious diseases in the
migration group.

Significance Statement

Host migration is commonly believed to accelerate the spread of infectious diseases. However,
recent ecological studies suggest that migration may impede this spread. In our study, we
developed a synthetic host-virus co-propagation model to explore the impact of host range
expansion on the interplay between host mobility and virus spatial distribution. Our experimental
and theoretical analysis revealed the spatial sorting of uninfected and infected hosts in the
navigated propagating front leads to faster back diffusion of infected hosts. This self-organized
structure allowed the migrating host population to eradicate the infectious disease, independent
of intricate host-virus dynamics.

44
45
46   **Main Text**

47
48   **Introduction**

49
Range expansion is a process by which living species invade new territories and gain benefits
such as survival(1–3), reproduction(4, 5), and resources(6, 7). Classic theory predicts that motile
populations expand in space and time by growing and moving(8–10). Recent studies have shown
that many organisms can use self-generated cues, known as navigated range expansion, to
increase the expansion rate toward long-term settlement(11–14). However, nonmotile species,
have to use other strategies to expand their territories, such as viruses, which depend on the
infected hosts to spread across long distances(15–17). It is commonly assumed that host motility
enhances virus transmission and viral range expansion(18–21). However, recent ecological
studies have also reported that animal host migrations can reduce viral prevalence(22–25). This
can occur when animals escape from infected regions ('migratory escape')(26, 27) or when
infected animals are removed from migrating populations ('migratory culling')(22, 28). One of the
prominent examples is that the seasonal migration of butterfly monarchs reduces the risk of
infection of parasites(22, 29). These conflicting findings from epidemiological case studies
suggest that the effects of host spatial expansion on viral prevalence depend on the specific
conditions of the system, which require a close and quantitative examination. However, studying
infectious disease dynamics and their mechanisms in field studies is difficult due to the limitations
of observatory technology and the complexity of the natural system. Therefore, we developed a
co-propagation system with motile bacteria, *Escherichia coli* (*E.coli*), and its chronic
bacteriophage M13 which does not kill infected bacteria cells when producing progeny phage
after infection(30), to investigate how host range expansion affects the relationship between host
motility and viral spatial prevalence.

71
72
73 **Results**
74
Phage spread during the bacterial range expansion
We adopted a swimming assay of motile bacteria by inoculating a small droplet of cells at the
center of a semi-solid agar plate (green dot in Fig. 1A, 2 µL containing about 10^6 cells). The
motile bacterial population grew and expanded radially. Ahead of the expanding bacterial
population, we plated a small volume of nonlethal phage M13 (light pink dot in Fig. 1A, containing
about $5{\times}10^6$ phage particles). When the bacteria encountered the phage zone, they became
infected and temporarily reduced their growth (Fig. S1), resulting in a visible low-density region of
the infected zone compared to the uninfected zones (Materials and Methods, Fig. S2). To better
identify the infected region, we integrated a red fluorescent gene Ruby into the M13 phage
genome (M13-Ruby, Fig. S3A). The infection of M13 phage introduced the fluorescent gene into
the bacterial cells, which expressed the red fluorescent protein. Therefore, the infected region can
be detected by the fluorescent signal measurement, which coincides with the visible low-density
region (Fig. 1B, Fig. S3B). Time-lapsed fluorescent images further showed the dynamics of the
fan-shape infected zone, where the boundary between the infected and uninfected zones was
rapidly fixed once it emerged (Supplementary Movies 1&2).
90
To understand the dynamics of bacteria-phage co-propagation, we extended a growth-expansion
model that incorporates bacterial chemotaxis and M13 phage infection kinetics (Fig. 2A and SI
Text). The model simulation provided a comprehensive picture about the formation dynamics of
the fan-shape infected zone (Fig. S4). The infected cells produced new phages that would further
infect more uninfected cells along the front of the migrating population at the same propagating
speed as the host front migration speed. At the same time, the phages infected other neighboring
bacteria, causing the lateral expansion of the infected region with a lateral expanding speed. The
bi-directional propagation results in a fan-shaped infection region (the light pink region in Fig. 1A).
The fan-shaped angle θ defines how fast the viral spread during the co-propagation with host
bacteria migration, and was named the angle of phage spread. Assuming the phage infection
area is approximately a sector, the angle of phage spread θ is derived as the lateral expansion
speed divided by the migration speed (Fig.1A).
103
The dependence of viral spread on the host migration speed during co-propagation
We then examined experimentally how the angle of phage spread θ depended on the host
bacterial motility by varying the agar concentrations. At high agar concentrations, the probability
of bacterial cells being trapped by the agar gel matrix increased, thereby reducing bacterial
motility(31, 32). For the chemotactic bacterial population, which enhances their group migration
by following their self-generated chemoattractant gradient (known as navigated range
expansion(11)), the speed of range expansion decreased as the agar concentration
increased(33). Interestingly, even though host bacterial motility was limited, the fan-shaped
infected zone increased at a fixed distance from the initial infection site (Fig. 1b). We quantified
the angle of phage spread θ and found it showed a negative dependence on co-propagation
migration speed (Fig. 1c). However, when using a chemotaxis deficient strain (Ft-
MG$\Delta cheRcheB$), which performs unguided range expansion following the Fisher–Kolmogorov
dynamics, in this case, the negative relation did not hold anymore (Fig. S6C-E), demonstrating
that faster host motility enhanced viral spread. These results suggested a counterintuitive trend:
during host navigated range expansion, an increase in host group migration speed would limit
phage spread.
120
121 By varying the host diffusive motility, our model can successfully recapture the negative relation
122 between co-propagation migration speed and viral spread (Fig. S7A). The simulation results
123 indicated that while both co-propagation migration speed and lateral expansion speed increased

124   with the host diffusion coefficient, the co-propagation migration speed with host chemotaxis
125   increased at a much faster rate (Fig. S7BC).
126
In contrast, when considering the model without host bacterial chemotaxis (unguided range
expansion, SI text), lateral expansion increased more rapidly as a function of bacteria diffusion
coefficient (Fig. S5), resulting in a positive dependence of viral spread on the host motility (black
line in Fig. S6D). We also confirmed that this relationship was not influenced by the geometric
effect of radial expansion under the plane wave expansion (Fig. S8). Therefore, our modeling
study predicted that, in the presence of host chemotaxis, increased host motility limits viral
spread.
134
The model predicted that the co-propagation group migration speed also depended on the host
chemotactic coefficient, which measures how responsive bacterial cells are to the environmental
chemoattractant gradient(34). We found that the co-propagating migration speed of the host and
phage increased linearly with the host chemotactic coefficient, while the lateral expansion speed
stayed approximately constant (Fig. S9B, C). As a result, the angle of phage spread θ, as well as
the size of the phage-infected zone, decreased significantly when the host chemotactic migration
speed increased (Fig. 2B, C, Fig. S9A).
142
We further tested the model prediction by measuring the angle of phage spread θ with a
chemotactic coefficient titratable strain (ZF1). Specifically, we deleted the native regulation of a
chemoreceptor gene *tar* in the bacterial genome, and introduced a titratable control of *tar*
expression by a small molecule inducer, anhydrotetracycline (aTc) (Materials and Methods, Fig.
3A). The *tar* variant strain affected the receptor gain of the cells in response to aspartate(35), but
not the tumble bias and growth rate(36). The tumble bias defines the average time a cell spends
on the tumbling motion, which further determines the effective diffusion coefficient of bacteria.
Therefore, by titrating different inducer aTc concentrations, we could manipulate the bacterial
chemotactic migration speed (Fig. 3B, Fig. S10A), but not the effective diffusion coefficient of
bacteria(36). Under different host migration speeds, we measured the angle of phage spread θ
and found it decreased as the migration speed increased (Fig. 3B, Fig. S10B&10C). The
experimental results confirmed our model prediction that virus spread is hindered by the host
chemotactic migration.
156
The migratory culling of phages during co-propagation
Another important prediction of our model is that co-propagation of host bacteria and phage is not
always sustained during the host navigated range expansion. As we decreased the phage
production rate in the model, it resulted in a slower lateral expansion of the infected zone, limiting
the angle of phage spread θ and the viral spread. At very low phage production rate, the angle of
phage spread θ became small. If we further increased the group migration speed, the size of the
infected zone shrank until a sudden drop to almost zero, where infection only occurred near the
initial spot of phage (Fig. S11). This indicated that the co-propagation of host bacteria and phage
could not be maintained and the infected cells as well as the phages were culled from the
propagating front. In other words, the system underwent a phase transition from co-propagation
to migratory culling (Fig. 2B&2C).
168
To experimentally test the phase transition of migratory culling, we engineered both the M13
phage and its bacterial host to enable the titration of the phage production rate after infection.
Specifically, we first deleted the *gIII* gene from the M13 phage genome and replaced by a T7 *rnap*
gene (M13SP)(37, 38). The native GIII protein is essential for the M13 phage into a host bacteria,
and deletion of GIII protein prevents full escape from the infected host(38). In other words, the
variant phage M13SP cannot be produced by wild-type infected cells, as it lacks *gIII* gene(39, 40).
We then incorporated the *gIII* gene into bacterial host cells under a T7 promoter (strain FTT7,
Materials and Methods). When the M13SP infects the engineered host FTT7, the bacteria cells
express the T7 RNAP whose gene is brought by the phage, which then activates the expression

of GIII protein and restores the complete production of M13SP phage. Therefore, the variant
phage M13SP infects the engineered strain FTT7 and generates a fan-shaped infection zone
during the co-propagation range expansion, and the dependence of the angle of phage spread θ
on the host migration speed is similar to the wild-type system (Fig. 3B and Fig. 4B).
182
An important advantage of this engineered phage-bacteria system is that we can generate a
series of engineered strains with T7 promoter variants that have different T7 RNAP binding
affinities. Here, we used a library of T7 promoter variants that we had previously characterized in
our study(38)(Table S2, Fig. S12). The weak T7 RNAP binding affinity, e.g. FTG8 strain, reduces
the GIII protein expression(41), resulting in a smaller fan-shaped zone (Fig. 4B). Increasing the
host migration speed further would lead to a much smaller fan-shaped infected zone. However,
for variant strains such as FTF10 and FTD8, whose promoter expression activity is much smaller
than that of wild type T7 promoter, the fan-shaped infected zone only exists at the slow host
migration speed. When the host migration speed becomes large, the infected zone is confined to
a small spot (Fig. 4B). Away from the initial drop of phage, we verified that no infected bacteria or
phage particles were carried, suggesting they were culled from the front propagation. We then
systematically measured the angle of phage spread θ for a series of variant strains at different
host migration speeds (Fig. 4C&D). The phase transition from co-propagation to migratory culling
is clearly observed at low promoter expression activity and fast host migration speed (Fig. 4D),
demonstrating the occurrence of migratory culling.
198
The spatial sorting mechanism of migratory culling
To better understand the underlying mechanism of migratory culling, we examined the simulated
dynamics of the co-propagation front. During navigated range expansion, the leading front of the
host population exhibited a propagating bulge, resulting from the balance between local cell
growth and back diffusion (leakage of cells out of the bulge due to random movement)(3, 11).
Due to phage infection, along the middle radial axis (the black dashed line in Fig. 4e), the
propagating front of the host population rapidly became infected once it crossed the initial phage
spot, with the phage profile generally following that of the infected cells (Fig. S13). However,
simulation results showed the coexistence of uninfected and infected cells near the edge of the
infected zone (along the blue arrow in Fig. 4E, Fig. S14). These cells, traveling along with
phages, formed a co-propagation front (Fig. 4E, Fig. S14). Within this co-propagation front, the
profile of infected cells was determined by three major factors: continuous conversion from
uninfected cells due to phage infection, self-reproduction, and reduction flux caused by back
diffusion. As the immotile phage always lagged behind the motile host, infection was most active
behind the front of the uninfected cells. This resulted in a spatial sorting structure where the
uninfected cells were located at the leading edge of the leading propagation front, while infected
cells were at the rear (Fig. 4E, Fig. S14).
216
To validate this prediction of spatial sorting, we experimentally measured the profiles of
uninfected and infected cells. We labeled the host cells with a red fluorescent gene *rfp* (ZF1-RFP)
and integrated a *gfp* gene into the phage genome (M13-GFP). This allowed us to quantify the
total cell density profile of the co-propagating front using red fluorescent signals (Fig. 4F, black
curve, see Materials and Methods). Phage infection introduced the *gfp* gene into infected cells,
enabling us to quantify the infected profile (the sum of infected and recovered cells) via green
fluorescent signals (Fig. 4F, orange curve). The difference between these two provided the profile
for uninfected (susceptible) cells (Fig. 4F, green curve). The experimental results clearly showed
the spatial difference between uninfected and infected cells at the propagating front near the
edge, which is consistent with the model prediction.
227
228 Consequently, the spatial sorting structure of the two subpopulations in the leading propagation
229 front indicated that the back diffusion of the infected cells was larger than that of the uninfected
230 cells. Without the conversion from uninfected cells by phage infection, the infected cells could not
231 keep up with the uninfected migration front. In the presence of moderate phage infection (e.g. low

phage production by infected cells), the increase in migration speed leads to stronger total back diffusion(11, 13), but the major increment came from the increase in the back diffusion of infected cells. Increasing the migration speed further or reducing the phage infection would make the back diffusion larger than the sum of the self-growth of infected cells and conversion rate from uninfected cells, causing the infected cells to be culled from the migration front. This resulted in a transition from co-propagation to migratory culling of infected bacterial host and phage.

Discussion

The understanding of the interplay of host migration and viral spread has been developed with the advent of global change and advances in genetics, but direct experimental tools are still lacking (13, 22, 27). Here, we developed a synthetic host-viral copropagating system to investigate how host motility affected viral spread during the host's navigated range expansion. We found a counterintuitive phenomenon that faster host chemotaxis-driven range expansion inhibited viral range expansion and resulted in a phase transition from co-propagation to migratory culling of phage.

Although migratory culling has been increasingly recognized in natural migratory animals and its role in regulating viral transmission dynamics, there is still a lack of agreement on how often and how much it may occur(21, 42). It is thought to depend on the extent to which infection affects the host's physiological and behavioral traits(43). Our finding revealed that the spatial sorting of uninfected and infected hosts in the navigated propagation front resulted in a faster back diffusion of infected hosts, enabling their elimination. Even without the temporary growth reduction during infection, this mechanism still facilitated the migratory culling (Fig. S15). This quantitative understanding does not require complex infection-induced changes to host migrations, which has potential implications for controlling the spread of infectious diseases by altering the host motility or the virus production rate.

Range expansion can significantly alter evolutionary dynamics through multiple factors, e.g. genetic drift(44, 45), selection pressures(46, 47), founder effects(48, 49), and cooperation collapse(50, 51). Long-term co-propagation of host and virus can lead to co-evolution, impacting the genetic diversity and adaptation of both host and virus populations(15, 16). In addition, our previous study utilized the bacteria-phage co-propagating system to develop a spatial phage-assisted continuous evolution system, which revealed the evolutionary process during co-propagation was accelerated compared to a fixed niche(38). Therefore, the inhibition of viral spread by host migration discovered in this study would further help to improve the directed evolution method and provide new insights into host-viral co-evolution dynamics.

Materials and Methods

Media and growth conditions
The Luria-Bertani (LB) medium contained 10 g tryptone, 5 g yeast extract, and 5 g NaCl per liter. The defective rich defined medium (D-RDM) used in this study was based on the Neidhardt's lab recipe(52) and modified: 1× MOPS mixture, 0.25× ACGU, 1× defective amino acid mixture (a mixture of 17 amino acid excluding asparagine, aspartic acid and serine, named as 1×AA), 1.32 mM K_2HPO4, 100 μM aspartic acid(K salt, 0 μM only for the host unguided range expansion experiment) and 0.4% (w/v) glucose. Mops salts and ACGU were prepared to be 10× MOPS and 10× ACGU stocks solution. The defective amino acid mixture 1×AA was prepared to be 5×AA stocks solution (in 1 L): 356.4 mg alanine, 5478.2 mg arginine (HCL), 87.8 mg cysteine (HCL), 611.5 mg glutamic acid (K salt), 438.6 mg glutamine, 300.3 mg glycine, 209.6 mg histidine (HCL H_2O), 262.4 mg isoleucine, 524.8 mg leucine, 365.4 mg lysine, 149.2 mg methionine, 330.4 mg phenylalanine, 230.2 mg proline, 238.2 mg threonine, 102.1 mg tryptophane, 181.2 mg tyrosine, 351.6 mg valine. All the medium in this study was buffered to pH 7.0 with 0.1 M HEPES (pH 7.0).

To prepare semi-solid plates, the bacto-agar (BD,214010) was added to the growth medium, and
the agar concentration varied from 0.2% to 0.4% (w/v). If required, inducer aTc was also added to
the medium and its concentration varied from 0 to 20 ng/mL. Then, 10 mL of the above medium
supplemented was poured into a 90-mm Petri dish and allowed to harden at room temperature for
60 min in a light-proof box. In addition, the medium was supplemented with chloramphenicol (25
µg/mL), spectinomycin (50 µg/mL), tetracycline (10 µg/mL), kanamycin (10 µg/mL) and ampicillin
(20 µg/mL). All experiments were executed at 37°C unless otherwise specified.
The culture condition of Fig. 1 and Fig. S8 (FM15 with different agar concentrations): D-RMD
medium + 100 µM aspartate + 0.2%~0.4% agar concentration + 10 µg/mL tetracycline;
The culture condition of Fig. S6 (Ft-MGΔ*cheRcheB* with different agar concentrations): D-RMD
medium + 0.2%~0.4% agar concentration + 10 µg/mL tetracycline;
The culture condition of Fig. 3 and Fig. S10 (ZF1 with different aTc concentrations): D-RMD
medium + 100 µM aspartate +0 ~ 20 ng/mL aTc + 0.2% agar concentration + 10 µg/mL
kanamycin + 20 µg/mL ampicillin;
The culture condition of Fig. 4B,C,D, and Fig. S20 (Strains of the FT series, the chemotactic
titratable strain ZF1 coupling T7 RNAP activity with the expression of *gIII*: FTT7, FTA1, FTD5,
FTD8, FTD9, FTF10, FTG8): D-RMD medium + 100 µM aspartate +0 ~ 20 ng/mLaTc + 0.25%
agar concentration+ 10 µg/mL kanamycin + 50 µg/mL spectinomycin;
The culture condition of Fig. 4E (ZF1-RFP): D-RMD medium + 200 µM aspartate + 20 ng/mL aTc
+ 0.25% agar concentration + 10 µg/mL kanamycin;
306
Strains and phages construction
The strains and plasmids used in this study are listed in Supplementary Table S1. The *E.coli* CLM
strain was provided by Dr Liu(3) and all other strains in this study, except *E. coli* K12 strain
ER2738, were derived from it. The strain FM15 was a conjugation of the *E. coli* CLM and the *E.*
coli K12 ER2738 (NEB, F plasmid with tetracycline-resistant provider) and was provided by Dr
Liu(38). Based on the understanding of the molecular mechanism about chemotactic signal
transduction of *E.coli*, we constructed a chemotaxis deficient *E.coli* strain Ft-MGΔ*cheRcheB* by
the following steps: (i) we constructed a chemotaxis deficient *E.coli* MGΔ*cheRcheB* by knocking
the chemotactic gene *cheR* and *cheB* of *E.coli* CLM; (ii) we transferred the psim5 plasmid with
chloramphenicol gene into the strain MGΔ*cheRcheB* and obtained the strain MGΔ*cheRcheB*-
psim5 (cultured in 30°C); (iii) we conjugated the recipient strain MGΔ*cheRcheB*-psim5 with the
donor strain FM15 (F plasmid provider) and got a strain Ft-MGΔ*cheRcheB*-psim5; (iv) we
removed the psim5 plasmid by culturing in 37°C and obtained the strain Ft-MGΔ*cheRcheB*. The
tar-titratable strain MGT was constructed as follows: (i) the strain CLM(Δ*tar*) was obtained by
knocking the chemotactic receptor protein Tar gene of the strain CLM. (ii) the bla:P$_{tet}$-*tetR-tar*
feedback loop was amplified and inserted into the strain CLM(Δ*tar*) chromosomal attB site
utilizing λRed homologous recombination, then we obtained the strain MGT. The strain FkP was
constructed by replacing the *tetA-PtetA/tetR-tetR* cassette of F plasmid in the strain Er2738 with
the kanamycin-resistance gene. The tar-titratable strain ZF1 was a conjugation of the strain MGT
and the strain Fkp (F plasmid with Kan resistant provider). The strain FTT7 was constructed by
electroporating the plasmid AP-T7 (T7 RNAP-dependent accessory plasmid contain M13 phage
gene *gIII* with a wild type T7 promoter, gift from Dr Chenli Liu(38)) into the strain ZF1, and the
similar strain FTA1/FTD5/FTD8/FTD9/FTF10/FTG8 was constructed by electroporating the
plasmid AP-A1/ AP-D5/ AP-D8/ AP-D9/ AP-F10/ AP-G8 (The variants of plasmid AP-T7 with T7
promoter variants, and the details of the promoter sequence are shown in Appendix Table S1,
gifts from Dr Liu(38)) into the strain ZF1. To better capture the location difference of the infected
and uninfected cells in the co-propagating front, we constructed the strain ZF1-RFP, in which a
red fluorescence plasmid *RFP* with chloramphenicol gene was inserted into the strain ZF1.

The phage M13 used in this study was a gift from Dr Liu(38) and others were its variants. To
better characterize the infectious state of the bacteria, the coding sequence of green/red
fluorescence protein gene was inserted into the genome of M13 and the strain was designated as
M13-GFP, M13-RFP and M13-Ruby. The phage M13SP was also provided by Dr Liu(38), in
which gene *gIII* of the M13 genome was deleted and replaced by a T7 RNP with a downstream
yellow fluorescence protein gene. All phages were verified by sequencing.
341
Conjugation of F plasmid
The conjugation of F plasmid is based on the Barrick laboratory's recipe(53), and the detailed
protocol is as follow: (i) grow overnight cultures of donor and recipient strains in the presence of
diaminopimelic acid (DAP, 0.3 mM) and appropriate antibiotic. (ii) gently spin down 1 mL culture
(~6000 rpm for 3 minutes) and wash donor and recipient cells in PBS, then repeat and resuspend
in 500 µL PBS. (iii) measure cell density and combine 1:1 ratio of donor and recipient cells in
micro centrifuge tube (100 µL:100 µL). (iv) plate 200 µL of mixture onto non-selective LB medium
plate containing DAP, and then incubate conjugation plate overnight. (v) scrape up the
conjugation mixture into a micro centrifuge tube with 1 mL PBS, vortex and gently spin down and
repeat, then resuspend conjugation mixture in 1 mL of PBS and plate 100 µL of this mixture and
100 µL of a 10-fold dilution onto selective plates. (vi) pick single colonies and culture in selective
media and confirm the strain via PCR amplification of the target strains.
354
Growth curve measurement
The growth curves of uninfected and infected bacteria were measured in a 250-mLflask with 50
mL corresponding growth medium at 37°C, 150 rpm. The general procedure was as follows. First,
the isolated bacteria from –80°C stock was streaked onto the agar plate with LB medium and
cultured at 37°C overnight. Second, 3-5 single colonies were picked and inoculated into 14 mL
round-bottom test tubes containing 2 mL LB medium and cultured overnight in a shaker (220 rpm,
Shanghai Zhichu Instrument) as the seed-culture procedure. Third, the overnight seed-cultures
were diluted into 2 mL D-RDM medium by a ratio of 1:100 and grown to log-phase the next
morning; when the diluted cultures reached OD_{600} 0.2~0.3, the diluting step was repeated. Fourth,
5 mL cultures (OD_{600} around 0.2~0.3) were added into a 250-mL flask with 45 mL prewarmed D-
RDM medium and were cultured in a water-bath shaker (150 rpm, Shanghai Zhichu Instrument).
When the diluted cultures reached an OD_{600} of 0.2, 2 mL bacteria culture was added to 48 mL
prewarmed D-RDM medium with or without phage. The medium was then incubated and
measured. The samples were taken every 10 min for measurement of OD_{600} by using
spectrophotometer reader until bacterial cells entered the stationary phase. For phage growth
assay, one-milliliter samples were extracted at the same time points and filtered through a 0.22
µm pore size PES syringe filter to remove bacterium. Aliquots for the time series were then stored
at -20°C until tittering. The growth curve was illustrated in Fig. S1.
373
Phage propagation and tittering
The strain FM15 was served as host bacteria for propagating and tittering M13 phage. Cells used
for phage propagation were cultured in 20 mL LB medium until they reached OD_{600}= 0.3~0.4 and
then infected with 10^9 pfu of phage. The bacteria-phage mixture was incubated overnight at 37°C.
Cell debris was removed by centrifugation at 5000 rpm for 10 min and filtration. The fresh
supernatant containing the revived phage was collected and stored at -20°C for up to several
months. To determine the phage titer, we employed the double agar overlay plaque assay as
described in previous work(54). The bottom agar for plates and soft agar for overlayers were LB
broth containing 1.5% and 0.4% Bacto agar, respectively. Serial ten-fold dilutions of the phage
stock and the underlay agar plates were prepared in advance. We transferred 10 µL of the
selected dilution of phages to a tube of 3 mL warm overlay medium, immediately added 100 µL
culture of the host bacterium (OD_{600} around 0.3~0.4), mixed and poured the contents over the
surface of a dried and labeled underlay plate. The overlays were allowed to harden for 30 min
and then incubated at 37°C overnight. The following day, we counted plaques on plates with 30–

300 plaques and defeminated the titer of the original phage stock by using the following
calculation: Number of plaques×100×reciprocal of counted dilution = pfu/mL.
390
Expansion experiment procedures
First, the isolated bacteria from −80°C stock was streaked onto the agar plate with LB medium
and cultured at 37°C overnight. 3-5 single colonies were picked and inoculated into 14 mL round-
bottom test tubes containing 2 mL LB medium and cultured overnight in a shaker (220 rpm.,
Shanghai Zhichu Instrument) as the seed-culture procedure. Second, the overnight seed-cultures
were diluted into 2 mL D-RDM medium by a ratio of 1:100 and grown to log-phase the next
morning, and then were further diluted in the same way when the diluted cultures reached OD_{600}
0.2~0.3. In the experiment of measuring the host's range expansion speed, bacteria were then
cultured to the mid-log phase (OD_{600} was around 0.2~0.3), which was inoculated at the line 2 cm
away from the center of a semi-solid agar plate using a 75-mm glass slide for the plane-wave
range expansion. Alternatively, 2 μL of the strain was inoculated at the center of a semi-solid agar
plate for the normal range expansion. In the virus infection experiment procedures, 2 μL M13
phages with different concentrations were inoculated 1 cm away from the bacteria position, and
then incubated at 37°C for several hours until the bacteria occupied the whole plate. For the
experiment of measuring bacteria expansion speed, only the bacteria strain was inoculated and
incubated at 37°C.
407
Bacteria expansion speed measurement
The semi-solid agar plate was illuminated from below by a circular white LED array with a lightbox
as described previously(55) and imaged at 1 h or 2 h intervals using a Canon EOS 600D digital
camera. Images were analyzed using ImageJ. For the normal range expansion, a circle was fitted
to the intensity maximum in each image and the area (A) of the fitted circle was determined. The
radius (r) of the colony was calculated as $r = \sqrt{A/\pi}$. The maximum expansion speed was
calculated using a linear fit over a sliding window of at least five time points, with the requirement
that the fit has an R^2 greater than 0.99. For the plane-wave range expansion, an image analysis
script using MATLAB was written to find the front peak position of the images at different times
and then we got the mean migration speed by a linear fitting.
418
Calculation of the angle of phage spread θ
When the semi-solid agar plate was occupied by bacteria, brightfield images of the plates were
captured by the image device in the above Methods and fluorescence images were captured
using the UVP CHEMSTUDIO™ TOUCH 815 IMAGERS (Analytik Jena, US). Virus spread
intensity can be represented by the angle θ of the fan-shaped pattern and can be determined as
follows. Using ImageJ analysis to fluorescence images, the phage infection area (Ar) of the fixed
region (L centimeter away from the initial phage inoculation position) was calculated, and the
angle of phage spread θ was $\theta = \frac{2Ar}{L^2}$. The angle of phage spread θ was still calculated using this
formula even when the sides of expansion zones are curved.
428
Fluorescence intensity assay
To investigate the location difference of the infected and uninfected cells in the propagating front
(Fig. 4E), the plates were scanned by Nikon Ti-E microscopy equipped with a 10× phase contrast
objective (Nikon CFI Plan Fluor DL4X F, N.A. 0.13, W.D. 16.4 mm, PhL), the green and red
fluorescence was taken with a 2-ms and 500-ms exposure time, respectively. The details were as
follows: firstly, as the above-mentioned method, the semi-solid D-RDM medium plate with 0.25%
agar and 20 ng/mL aTc was prepared, and the strain ZF1-RFP was cultured; Second, the strain
ZF1-RFP was inoculated at the line 2 cm away from the center of the plate and 2 μL 10^9 pfu/mL
M13-GFP phage was inoculated 1 cm away from the bacteria position, and the plate was
incubated at 37°C for 10 hours; Third, the Fluorescence intensity of the whole propagating front
in the plate was scanned by the Nikon Ti-E microscopy and the data was exported by the NIS-
Elements AR software (ver. 4.50.00). The red fluorescence intensity represents the host cell ZF1-

RFP including the uninfected and infective cells (infected cells + recovered cells), and the green fluorescence intensity represent the infective cells which express the green fluorescence gene of M13-GFP because of the phage infection. In the back of the infected zone, both fluorescent signals could be normalized by assuming the majority of the cells had been infected (38). Hence, the uninfected cell can be represented by the red fluorescence intensity minus the green fluorescence intensity. This was processed using a custom-written MATLAB code.

Acknowledgments

The authors thank C. Liu for sharing *E. coli* strain, Y. Bai, S. Huang for discussions, comments, and technical support. This work is partially supported by the National Key Research and Development Program of China (2021YFA0911100), Strategic Priority Research Program of the Chinese Academy of Sciences (XDB0480000), the National Natural Science Foundation of China (32301223,32071417, 32025022).

Data availability

Major experimental data supporting the findings of this study are available within the main text and Supplementary Information. Modeling and analysis code have been submitted to GitHub with a link https://github.com/YiZhangsiat/Host-Phage-Expansion.

References

1. H. Xu, J. Dauparas, D. Das, E. Lauga, Y. Wu, Self-organization of swimmers drives long-range fluid transport in bacterial colonies. *Nat Commun* **10**, 1792 (2019).

2. H. H. Mattingly, T. Emonet, Collective behavior and nongenetic inheritance allow bacterial populations to adapt to changing environments. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2117377119 (2022).

3. W. Liu, J. Cremer, D. Li, T. Hwa, C. Liu, An evolutionarily stable strategy to colonize spatially extended habitats. *Nature* **575**, 664–668 (2019).

4. Y. Cao, J. Neu, A. E. Blanchard, T. Lu, L. You, Repulsive expansion dynamics in colony growth and gene expression. *PLoS Comput Biol* **17**, e1008168 (2021).

5. O. Hallatschek, D. R. Nelson, Life at the front of an expanding population. *Evolution* **64**, 193–206 (2010).

6. J. Liu, *et al.*, Coupling between distant biofilms and emergence of nutrient time-sharing. *Science* **356**, 638–642 (2017).

7. W. Liu, T. A. Tokuyasu, X. Fu, C. Liu, The spatial organization of microbial communities during range expansion. *Current Opinion in Microbiology* **63**, 109–116 (2021).

8. J. G. Skellam, Random dispersal in theoretical populations. *Biometrika* **38**, 196 (1951).

9. R. A. Fisher, The wave of advance of advantageous genes. *Annals of Eugenics* **7**, 355–369 (1937).

10. D. A. Andow, P. M. Kareiva, S. A. Levin, A. Okubo, Spread of invading organisms. *Landscape*
Ecol **4**, 177–188 (1990).

11. J. Cremer, *et al.*, Chemotaxis as a navigation strategy to boost range expansion. *Nature* **575**,
658–663 (2019).

12. N. W. Frankel, *et al.*, Adaptability of non-genetic diversity in bacterial chemotaxis. *eLife* **3**,
e03526 (2014).

13. A. V. Narla, J. Cremer, T. Hwa, A traveling-wave solution for bacterial chemotaxis with
growth. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2105138118 (2021).

14. S. Bi, *et al.*, Discovery of novel chemoeffectors and rational design of *Escherichia coli*
chemoreceptor specificity. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 16814–16819 (2013).

15. D. Ping, *et al.*, Hitchhiking, collapse, and contingency in phage infections of migrating
bacterial populations. *The ISME Journal* **14**, 2007–2018 (2020).

16. S. Heilmann, K. Sneppen, S. Krishna, Coexistence of phage and bacteria on the boundary of
self-organized refuges. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 12828–12833 (2012).

17. X. Li, F. Gonzalez, N. Esteves, B. E. Scharf, J. Chen, Formation of phage lysis patterns and
implications on co-propagation of phages and motile host bacteria. *PLoS Comput Biol* **16**,
e1007236 (2020).

18. C. Viboud, *et al.*, Synchrony, Waves, and Spatial Hierarchies in the Spread of Influenza.
Science **312**, 447–451 (2006).

19. S. Krauss, *et al.*, Coincident ruddy turnstone migration and horseshoe crab spawning creates
an ecological 'hot spot' for influenza viruses. *Proc. R. Soc. B.* **277**, 3373–3379 (2010).

20. J. Waldenström, S. Bensch, S. Kiboi, D. Hasselquist, U. Ottosson, Cross-species infection of
blood parasites between resident and migratory songbirds in Africa. *Molecular Ecology* **11**,
1545–1554 (2002).

21. A. Fritzsche McKay, B. J. Hoye, Are Migratory Animals Superspreaders of Infection?: An
Introduction to the Symposium. *Integr. Comp. Biol.* **56**, 260–267 (2016).

22. S. Altizer, R. Bartel, B. A. Han, Animal Migration and Infectious Disease Risk. *Science* **331**,
296–302 (2011).

23. M. Krkošek, *et al.*, Effects of host migration, diversity and aquaculture on sea lice threats to
Pacific salmon populations. *Proc. R. Soc. B.* **274**, 3141–3149 (2007).

24. A. K. Shaw, S. A. Binning, Migratory Recovery from Infection as a Selective Pressure for the
Evolution of Migration. *The American Naturalist* **187**, 491–501 (2016).

25. D. A. Satterfield, J. C. Maerz, S. Altizer, Loss of migratory behaviour increases infection risk
for a butterfly host. *Proc. R. Soc. B.* **282**, 20141734 (2015).

26. R. J. Hall, S. Altizer, R. A. Bartel, Greater migratory propensity in hosts lowers pathogen
transmission and impacts. *Journal of Animal Ecology* **83**, 1068–1077 (2014).

27. R. A. Bartel, K. S. Oberhauser, J. C. De Roode, S. M. Altizer, Monarch butterfly migration and
parasite transmission in eastern North America. *Ecology* **92**, 342–351 (2011).

28. A. Mysterud, L. Qviller, E. L. Meisingset, H. Viljugrein, Parasite load and seasonal migration
in red deer. *Oecologia* **180**, 401–407 (2016).

29. S. Altizer, K. A. Hobson, A. K. Davis, J. C. De Roode, L. I. Wassenaar, Do Healthy Monarchs
Migrate Farther? Tracking Natal Origins of Parasitized vs. Uninfected Monarch Butterflies
Overwintering in Mexico. *PLoS ONE* **10**, e0141371 (2015).

30. Z. Hobbs, S. T. Abedon, Diversity of phage infection types and associated terminology: the
problem with "Lytic or lysogenic." *FEMS Microbiol Lett* **363**, fnw047 (2016).

31. N. A. Licata, B. Mohari, C. Fuqua, S. Setayeshgar, Diffusion of Bacterial Cells in Porous
Media. *Biophysical Journal* **110**, 247–257 (2016).

32. T. Bhattacharjee, S. S. Datta, Bacterial hopping and trapping in porous media. *Nat Commun*
10, 2075 (2019).

33. O. A. Croze, G. P. Ferguson, M. E. Cates, W. C. K. Poon, Migration of Chemotactic Bacteria in
Soft Agar: Role of Gel Concentration. *Biophysical Journal* **101**, 525–534 (2011).

34. X. Fu, *et al.*, Spatial self-organization resolves conflicts between individuality and collective
migration. *Nat Commun* **9**, 2177 (2018).

35. D. Yu, X. Ma, Y. Tu, L. Lai, Both piston-like and rotational motions are present in bacterial
chemoreceptor signaling. *Sci Rep* **5**, 8640 (2015).

36. Y. Bai, *et al.*, Spatial modulation of individual behaviors enables an ordered structure of
diverse phenotypes during bacterial group migration. *eLife* **10**, e67316 (2021).

37. K. M. Esvelt, J. C. Carlson, D. R. Liu, A system for the continuous directed evolution of
biomolecules. *Nature* **472**, 499–503 (2011).

38. T. Wei, *et al.*, Exploiting spatial dimensions to enable parallelized continuous directed
evolution. *Molecular Systems Biology* **18**, e10934 (2022).

39. M. Ploss, A. Kuhn, Kinetics of filamentous phage assembly. *Phys. Biol.* (2010).

40. S. W. Smeal, M. A. Schmitt, R. R. Pereira, A. Prasad, J. D. Fisk, Simulation of the M13 life
cycle I: Assembly of a genetically-structured deterministic chemical kinetic simulation.
Virology **500**, 259–274 (2017).

41. B. W. Thuronyi, *et al.*, Continuous evolution of base editors with expanded target
compatibility and improved activity. *Nat Biotechnol* **37**, 1070–1079 (2019).

42. A. Risely, M. Klaassen, B. J. Hoye, Migratory animals feel the cost of getting sick: A
meta-analysis across species. *Journal of Animal Ecology* **87**, 301–314 (2018).

43. C. A. Bradley, S. Altizer, Parasites hinder monarch butterfly flight: implications for disease
spread in migratory hosts. *Ecology Letters* **8**, 290–300 (2005).

44. B. T. Weinstein, M. O. Lavrentovich, W. Möbius, A. W. Murray, D. R. Nelson, Genetic drift
and selection in many-allele range expansions. *PLOS Computational Biology* **13**, e1005866
(2017).

45. L. Bosshard, S. Peischl, M. Ackermann, L. Excoffier, Dissection of the mutation accumulation
process during bacterial range expansions. *BMC Genomics* **21**, 253 (2020).

46. T. Wein, T. Dagan, The Effect of Population Bottleneck Size and Selective Regime on Genetic
Diversity and Evolvability in Bacteria. *Genome Biology and Evolution* **11**, 3283–3290 (2019).

47. J. M. Horowitz, M. Kardar, Bacterial range expansions on a growing front: Roughness,
fixation, and directed percolation. *Phys. Rev. E* **99**, 042134 (2019).

48. B. M. Peter, M. Slatkin, The effective founder effect in a spatially expanding population.
Evolution **69**, 721–734 (2015).

49. P. Chu, J. Zhu, Z. Ma, X. Fu, Colony pattern development of a synthetic bistable switch.
(2024). https://doi.org/10.1101/2024.06.17.599191.

50. N. Luo, *et al.*, The collapse of cooperation during range expansion of Pseudomonas
aeruginosa. *Nat Microbiol* **9**, 1220–1230 (2024).

51. K. S. Korolev, The Fate of Cooperation during Range Expansions. *PLoS Comput Biol* **9**,
e1002994 (2013).

52. F. C. Neidhardt, P. L. Bloch, D. F. Smith, Culture Medium for Enterobacteria. *J Bacteriol* **119**,
736–747 (1974).

53. S. P. Leonard, *et al.*, Genetic Engineering of Bee Gut Microbiome Bacteria with a Toolkit for
Modular Assembly of Broad-Host-Range Plasmids. *ACS Synth. Biol.* **7**, 1279–1290 (2018).

54. A. Baer, K. Kehn-Hall, Viral Concentration Determination Through Plaque Assays: Using
Traditional and Novel Overlay Systems. *JoVE* 52065 (2014). https://doi.org/10.3791/52065.

55. C. Liu, *et al.*, Sequential Establishment of Stripe Patterns in an Expanding Cell Population.
Science **334**, 238–241 (2011).

56. A. Roy, S. Mitra, Increased Fragility of Escherichia coli After Infection with Bacteriophage
M13. *J Virol* **6**, 333–339 (1970).

580
581

582
Fig.1. Phage spread during the bacterial navigated range expansion. (A) The co-propagation of
bacteria and phage leads to a fan-shaped infected zone. (B) Typical migration speeds and fan-
shaped infected zone (light pink) under different agar concentrations. *E.coli* FM15 was inoculated
at the center of a semisolid agar plate (Materials and Methods) with varying agar concentrations,
and the reporter phage M13-Ruby was inoculated 1 cm away from the center. Infected cells
expressed red fluorescent protein Ruby, introduced by phage during infection. The merged
pseudo-color image combines brightfield and fluorescence images. (C) Dependence of the angle
of phage spread θ on bacterial migration speeds by varying agar concentrations. Strain details
are given in Supplementary Table 1. Data were taken 96 hours after inoculation. Scale bar, 2 cm.
Experiments in (B) were repeated independently three times. For (C), data are mean ± s.d. for
n=5 biological replicates.
594

597
Fig. 2. Kinetic model of the interaction between bacteria and phage. (A) Schematic illustration of population's navigated range expansion. The M13 phage is chronic. During the interaction between bacteria and phage, the infected host cells are not killed; instead, progeny phages are produced and extruded through the cell membrane as infected cells continue to grow at a slower rate (38, 39, 56). This continues until the cells recover. Once recovered, the cells grow as fast as the susceptible ones and produce progeny phages at a much lower level as compared to freshly infected cells(38). In addition, recovered host cells cannot be re-infected by free phage particles (38). In the schematic illustration of the population's navigated range expansion: bacterial populations navigate forward into unoccupied territories depending on the chemoattractant gradients; Bacterial populations are classified into three categories: susceptible, infected, and recovered bacteria. Infected bacteria are converted from susceptible bacteria by phage infection, and eventually become recovered bacteria. These three bacterial populations all proliferate by consuming the nutrients. Motile bacteria expand their range into unoccupied territories by diffusion and chemotaxis. Meanwhile, nonmotile phages are transmitted by their host bacteria, and their titer depends on the level of infectious progeny phage production rate by infected and recovered bacteria (Supplementary text). (B) Simulated phase diagram of bacterial-viral co-propagation. The model predicted that the angle of phage spread θ was negatively correlated with the bacteria migration speed by varying the chemotactic coefficient and positively correlated with the phage production rate. The system experienced a phase transition from co-propagation to migratory culling as further increasing migration speed at low phage production rates (red dashed line). (C) The typical fan-shaped patterns under the different migration speeds and phage production rates.

A

Fig. 3. Phage spread under different co-propagating migration speeds by titration of host bacteria chemotactic abilities. (A) Design of the chemotactic ability titratable strain ZF1 by inducible expression of chemotactic receptor protein Tar (Materials and Methods). (B) Typical migration speeds and fan-shaped patterns under different aTc concentrations (Materials and Methods). Data were taken 30 h after inoculation. Scale bar, 2 cm. Data are presented as mean ± s.d. for n=3 biological replicates.

Fig. 4. The phase transition and the spatial sorting mechanism of migratory culling. (A) Schematic design of the chemotactic titratable strain ZF1 coupling T7 RNAP activity with the expression of *gIII*. The host bacteria carry a motility regulation module and an activity-dependent phage propagation module, which are harbored by chromosome and the accessory plasmid, respectively (Materials and Methods). (B) The typical migration speeds and fan-shaped patterns for the typical strains (FTT7, FTG8, FTF10, and FTD8) under different aTc concentrations. (C) Detailed relationship between the angle of phage spread θ and the migration speed in (B). (D) Phase transition of migratory culling by seven strains with different phage propagation activities (The detailed sequences of T7 promoter variants are shown in Supplementary Table S2) under different aTc concentrations (Materials and Methods). The angle of phage spread θ was

negatively correlated with the bacteria migration speed and positively correlated with the phage
production rate. Migratory culling occurred at higher migration speeds and lower phage
production rates. (E) Illustration of the spatial sorting mechanism for migratory culling. Along the
leading propagating front near the edge between the infected and uninfected zone (as indicated
by the blue arrow in the left panel), the simulated density profiles (Fig. S14) showed a spatial
sorting structure of the infective cells (infected + recovered bacteria, orange line) and the
susceptible cells (green line), leading to smaller back diffusion for the susceptible cell (green
arrow) compared to infective cells (orange arrow). (F) Co-propagation profiles (right panel) of
ZF1-RFP (a plasmid carrying red fluorescent gene rfp was inserted into the strain ZF1, Materials
and Method) and phage M13-GFP (a *gfp* gene was integrated into phage genome, Materials and
Methods) demonstrated the spatial sorting structure experimentally. The profiles of total (black)
and infective bacteria (infected + recovered bacteria, green) were directly quantified the red and
green fluorescent signals respectively, while the profile of susceptible bacteria (orange) was
calculated by the other two profiles. The shaded areas represent the error bar. Data are
presented as mean ± s.d. for n=4 biological replicates.