Title:

Autoimmune antibody-induced neuronal hyperactivity triggers pathological Tau in IgLON5 disease

Authors:

Bilge Askin[1,2], César Cordero Gómez[1,3], Sophie Lan-Linh Duong[1,3,4], Cagla Kilic[1], Alexander Goihl[5], Andrew Newman[6], Jessica Wagner[7,8,9], Valentine Thomas[1], Janne Lamberty[1], Sabrina Huebschmann[1], Ekaterina Semenova[1], Peter Körtvelyessy[3,10], Dirk Reinhold[5], Hans-Christian Kornau[1,11], Paul Turko[12], Jonas J. Neher[7,8,9], Dietmar Schmitz[1,2,11], Benjamin R. Rost[1], Lisa Diez[1], Harald Prüss[1,2,3], Susanne Wegmann[1,2]*

Affiliations:

[1] German Center for Neurodegenerative Diseases (DZNE), Berlin, Germany
[2] Einstein Center for Neurosciences (ECN), Charité - Universitätsmedizin Berlin, Berlin, Germany
[3] Department of Neurology and Experimental Neurology, Charité – Universitätsmedizin Berlin, Berlin, Germany
[4] Berlin Institute of Health (BIH) at Charité – Universitätsmedizin Berlin, Berlin, Germany
[5] Institute of Molecular and Clinical Immunology, Otto-von-Guericke-University, Magdeburg, Germany
[6] Institute of Cell Biology and Neurobiology, Charité– Universitätsmedizin Berlin, Berlin, Germany
[7] German Center for Neurodegenerative Diseases (DZNE), Munich, Germany
[8] Metabolic Biochemistry, Biomedical Center Munich (BMC), Faculty of Medicine, LMU Munich, Germany
[9] Munich Cluster for Systems Neurology (SyNergy), Munich, Germany
[10] German Center for Neurodegenerative Diseases (DZNE), Magdeburg, Germany
[11] Neuroscience Research Center (NWFZ), Charité-Universitätsmedizin Berlin, Germany
[12] Institute of Integrative Neuroanatomy, Charité, Berlin, Germany

* corresponding author: susanne.wegmann@dzne.de

Key words:

Anti-IgLON5 disease, autoantibodies, autoimmune encephalitis, tauopathy, Tau, hyperactivity, neurodegeneration, neuroinflammation

Abbreviations:

AABs - Autoantibodies
AD - Alzheimer's disease
AE - Autoimmune encephalitis
AMPA -α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
Aβ - Amyloid-beta
BSA - Bovine serum albumin
CA1- Hippocampal area CA1

Ca^{2+} - Calcium
CA3 - Hippocampal area CA3
CSF - Cerebrospinal fluid
DG - Dentate gyrus
ELISA - Enzyme-Linked Immunosorbent Assay
EPSC - Excitatory postsynaptic current
FTD - Frontotemporal dementia
GFAP - Glial fibrillary acidic protein
GO - Gene ontology
hNSCs - Human neural stem cells
IF - Immunofluorescence
Ig - Immunoglobulin
IHC - Immunohistochemistry
KEGG - Kyoto encyclopedia of genes and genomes
LDH - Lactate dehydrogenase
LGI1 - Leucine-rich glioma-inactivated 1 protein
mCtrl - Human monoclonal, non-reactive IgG
MF - Mossy fiber
MFG-E8 - Milk fat globule epidermal growth factor 8
NBA - Neurobasal-A
NfL - Neurofilament light chain
NGS - Normal goat serum
NMDA - N-methyl-D-aspartate
p-Tau - Phosphorylated Tau
PBS - Phosphate-buffered saline
PBS-T - PBS containing Tween
pCtrl - Human serum-derived control IgG purified from healthy control serum in-house
pCtrl$_{rec}$ - Commercial normal human IgG
PFA - Paraformaldehyde
PLL - Poly-L-lysine
PPR - Paired-pulse ratios
Pvr - Synaptic vesicle fusion
RT - Room temperature
SP - Signal peptide SP
TBS - Tris-buffered saline
78

Abstract

Anti-IgLON5 disease is an autoimmunity/neurodegeneration overlap disorder in which
autoantibodies (AABs) against the neuronal cell surface protein IgLON5 lead to profound brain
dysfunction. Brains of patients show Tau pathology, neuroinflammation, and
neurodegeneration in multiple brain regions. Through administering patient-derived α-IgLON5
AABs to mice and cultured neurons, we here deciphered the cellular mechanisms of Tau
pathology and neurodegeneration in α-IgLON5 disease, highlighting a central role of neuronal
activity modulation in the disease pathology. Pathogenic human α-IgLON5 AABs induced
acute neuronal hyperactivity, which triggered Tau changes typically found early in Tau-related
neurodegenerative diseases like Alzheimer's disease (AD). α-IgLON5 AAB-induced Tau

phosphorylation and somatodendritic resorting selectively occurred in key hippocampal
connections, involving dentate gyrus granule cells, mossy fiber projections and commissural
fiber tracts. These changes were accompanied by a Tau-specific neuroinflammatory
response, involving the complement pathway, microglial MHC class II proteins, T cell
receptors, and deregulation of synaptic activity and cell-cell interactions. These findings
provide new insights into the origin of autoimmune-triggered α-IgLON5 disease pathology and
highlight that, similar to recent reports in AD patients, neuronal hyperactivity may be a disease-
overarching driver of Tau pathology.

Introduction

Autoimmune antibodies (autoantibodies, AABs) directed against neuronal or glial epitopes in
the brain can cause neuropathological conditions or exacerbate existing diseases [1]. The
trigger for the production of AABs against brain epitopes is not fully understood, but abnormal
immune responses due to genetic predispositions, viral infections, or tumors can contribute to
certain autoimmune encephalitis (AE) cases [2].

α-IgLON5 disease is a recently described, rare but severe neurological disorder characterized
by the presence of AABs targeting the neuronal cell surface protein IgLON5 [3]. Individuals
diagnosed with α-IgLON5 disease demonstrate diverse clinical manifestations and commonly
experience sleep behavior abnormalities, movement disorders, memory deficits, and seizures
[4–6]. Brains of α-IgLON5 patients show intraneuronal accumulations of highly phosphorylated
Tau protein (p-Tau) in several regions, including hippocampus, cerebellum, brain stem,
hypothalamus, and basal ganglia, as well as in the spinal cord [3,7]. Pathological hallmarks in
primary and secondary tauopathies, such as frontotemporal dementia (FTD) variants and
Alzheimer's disease (AD), include high phosphorylation, somatodendritic re-sorting,
aggregation and deposition of the microtubule associated protein Tau [8,9]. Although α-IgLON5
disease differs from other tauopathies because of its unique set of clinical manifestations,
anatomical Tau distribution, and its autoimmune context, it can be categorized as a novel
tauopathy due to the histological similarities to well-known diseases with Tau pathology [3].

The discovery of α-IgLON5 disease poses the fundamental questions of whether and how
AABs against a cell surface protein can trigger intracellular Tau aggregation and associated
pathological changes in the brain. Understanding the mechanisms of how autoimmunity can
induce neurodegeneration could not only lead to specific treatment options for α-IgLON5 and

further autoantibody-mediated brain diseases, but also provide important insights into how
and why pathological Tau changes occur in more than 20 different neurological diseases [10].

The IgLON protein family is a group of immunoglobulin (Ig) domain cell adhesion molecules
comprising five gene members (OPCML (=IgLON1), NTM (=IgLON2), LSAMP (=IgLON3),
NEGR1 (=IgLON4), and IgLON5) [11]. IgLON5 is a GPI-anchored surface protein with three Ig-
domains (Ig1, Ig2 and Ig3) that extend into the extracellular space and mediate homo-dimer
and hetero-dimer formation with other IgLON family members on the cell surface and at the
synaptic cleft [12–14]. In humans and mice, IgLON5 is primarily expressed in neuronal cells in the
brain, including the hippocampal formation, cortex, hypothalamus, brain stem, basal ganglia,
cerebellum and spinal cord (The Human Protein Atlas:
https://www.proteinatlas.org/ENSG00000142549-IGLON5; Expression atlas:
https://www.ebi.ac.uk/gxa/home). The physiological function of IgLON5 is not fully understood,
however, other IgLON family members have been implicated in neuronal cell adhesion, neurite
growth, neural circuit and synapse formation [15–17].

A neurotoxic potential of α-IgLON5 AABs was previously suggested based on the observation
that long-term treatment (1-4 weeks) of cultured neurons with α-IgLON5 AABs induces
common neurodegenerative phenotypes, such as Tau phosphorylation, cytoskeleton
disruption [18], synapse loss, and cell death [19]. Here, using *in vitro* and *in vivo* models, we now
unravelled the origin of these changes through demonstrating that α-IgLON5 AABs trigger
acute neuronal hyperactivity, which activates persistent neuronal Tau phosphorylation and
somatodendritic resorting. These changes occur in selected hippocampal connections,
including mossy fiber (MF) connections between dentate gyrus (DG) and hippocampal region
CA3, and projections onto ventral CA1, and are accompanied by distinct gene expression
signatures indicative of neuroinflammation and synaptic restructuring/activity modulation.
These findings elucidate the cellular mechanism by which α-IgLON5 AABs trigger tauopathy-
like pathology and neurotoxicity in α-IgLON5 disease, and thereby provide the foundation for
the development of novel α-IgLON5 disease treatment approaches. In addition, our data
demonstrate that neuronal hyperexcitability is sufficient to drive Tau pathology and may
therefore represent a central pathomechanism across different neurological diseases
exhibiting Tau pathology.

Results

Case study and isolation of high-affinity α-IgLON5 AABs from α-IgLON5 patient plasma

A 59-year-old male patient presented with a history of sleep apnea, myoclonic jerks, dysarthria, and gait disorder for one year. After ruling out common neurodegenerative diseases by clinical expertise, we detected α-IgLON5 AABs both in cerebrospinal fluid (CSF; at a titer of 1:100) and serum (at a. titer of 1:1000) of the patient, confirming a diagnosis of α-IgLON5 disease. Oral treatment with the corticosteroid immunosuppressant Methylprednisolone did not improve the symptoms. This treatment approach was followed by plasmapheresis to remove AABs from the serum, after which the patient was stable with the application of non-invasive ventilation at night and immunosuppressive therapy (improved sleep apnea and frequency of the myoclonic jerks; remission was not achieved).

To isolate α-IgLON5 AABs from the patient's plasma for fine characterization, recombinant human IgLON5-Fc chimera were immobilized on affinity chromatography purification columns (Fig. 1a). The purified α-IgLON5 AABs (heavy chain ~50 kDa), light chain ~25 kDa; Fig. 1b) had a purity of 84±7% (determined based on signal intensity of [heavy + light chain] relative to total signal in SDS-PAGE). In dot blot assays, the α-IgLON5 AABs were reactive to recombinant IgLON5 protein but not to LGI1 (leucine-rich glioma-inactivated 1) protein, which functioned as a control for unspecific binding (Fig. 1c). Furthermore, isolated α-IgLON5 AABs spotted on a dot blot membrane were recognized by α-human IgG secondary antibodies specific for subclasses IgG1, IgG2, IgG3 and IgG4 (Supplemental Fig. S1a), indicating that the presence of a polyclonal pool of α-IgLON5 AABs in the patient's plasma. An antibody pool derived from healthy subject serum (pCtrl), which does not detect IgLON5 and contained all human IgG subclasses, was used as control.

α-IgLON5 AABs bind cell surface IgLON5 protein on human and mouse neurons

To validate the specificity of the isolated patient-derived α-IgLON5 AABs, we expressed full-length human IgLON5 protein (Fig. 1d) in HEK cells and detected the binding of α-IgLON5 AABs after incubation with fluorescently labeled α-human secondary IgGs (Fig. 1e). We detected binding to full-length IgLON5 as well as all (combinations of) individual GPI-anchored Ig domains (Ig1/2/3), indicating binding of the isolated polyclonal α-IgLON5 AAB pool to every domain of IgLON5. Titrating the isolated α-IgLON5 AABs to cells recombinantly expressing human IgLON5 showed a dose dependent binding and suggested an $EC_{50}=8$ μg/ml ($K_D=56$ μM at $MW_{AAB}=150$ kDa; Fig. 1f); the relative binding strength was determined based on immunofluorescence intensity after fixation and labeling with fluorescently labeled anti-human

IgG. Notably, the molecular binding affinity of α-IgLON5 AABs to IgLON5 protein may be far higher when determined in purified preparations. Furthermore, α-IgLON5 AABs also bound to the surface of human neural stem cells (hNSCs) (Fig. 1g). In fresh frozen mouse brain sections, α-IgLON5 AABs showed immunoreactivity across various brain regions, including the cortex, olfactory bulb, hippocampus, mid brain, hind brain (pons, medulla) and cerebellum (Fig. 1h). Notably, binding of α-IgLON5 AABs aligns with IgLON5 expression in these brain regions in mice (Supplemental Fig. S1b).

To test whether α-IgLON5 AABs bound preferably to pre- or post-synaptic IgLON5, we treated primary hippocampal mouse neuron cultures with α-IgLON5 AABs and determined the co-localization with MAP2 and pre- (Synapsin-1) and postsynaptic (PSD-95) markers. These experiments suggested no preference of α-IgLON5 AABs for dendritic or axonal processes (Fig. 1i), and similar partial co-localization with pre- and post-synaptic structures (Fig. 1j). Immunolabeling was detectable also with detergent permeabilization, hence representing cell surface IgLON5 labeling.

Neurologically active AABs targeting neuronal surface antigens, e.g., LGI1, N-methyl-D-aspartate (NMDA), α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) and glycine receptors, can induce antibody-antigen complexes and receptor internalization upon binding, which contributes to AE pathophysiology [2]. Accordingly, we detected the internalization and endo-lysosomal entry of α-IgLON5 AABs (labeled with pHrodo[TM] Red) in cultured neurons (after 1 day and 2 days) and in cells expressing human IgLON5 (after 4 h), which was less pronounced for a pool of IgGs lacking antibodies against brain cell targets (pCtrl, isolated from healthy control serum; Supplemental Fig. S1e). Previous data indicated that IgG1, but not IgG4 fractions of α-IgLON5 AABs are internalized by neurons [20], suggesting that only a fraction of our α-IgLON5 AABs may be internalized.

Together these data show that the plasma of the α-IgLON5 patient contains a high affinity polyclonal pool of α-IgLON5 AABs that are reactive to different epitopes in different sub-domains of the IgLON5 protein (Fig. 1k).

Patient-derived α-IgLON5 AABs induce motor deficits in mice

Postmortem brains of α-IgLON5 patients are characterized by the intraneuronal deposition of phosphorylated Tau (p-Tau) in various brain regions [3,7]. Like in other tauopathies, the presence of p-Tau appears to correlate with neuroinflammation and neuronal loss in α-IgLON5 affected regions.

To test whether α-IgLON5 AABs are sufficient to induce these changes in the brain, we
administered patient-derived α-IgLON5 AABs into the brain of adult wildtype mice
cerebroventricular infusion into the right cerebral ventricle for 14 consecutive days; a total
amount of 75 μg AABs were infused (~3 μg/g body weight). A serum-derived pool of IgGs from
healthy controls (pCtrl), as well as a non-reactive human monoclonal IgG (mCtrl: mGo53 [21])
and PBS, were used as control groups. After 14 days of infusion, blood serum and brains were
harvested and analyzed for neuroinflammation and p-Tau accumulation (Fig. 2a). Fluorescent
immunolabeling of fixed brain sections showed that the α-IgLON5 AABs diffused from the
infused ventricle through the ipsilateral hemisphere, with strong binding to ventral regions
(e.g., parts of thalamus, hypothalamus, amygdala), and to some extent into the contralateral
hippocampus (Fig. 2b; Supplementary Fig. S2a). Animals treated with pCtrl showed far less
labeling. These results indicated that patient-polyclonal α-IgLON5 AAB are specific to their
target and can bind to IgLON5 *in vivo*.
In addition, animals treated with α-IgLON5 AABs showed phenotypic changes. In particular,
compared to all control groups (pCtrl, mCtrl, and PBS), the body weight of α-IgLON5 AAB
infused animals stagnated for the first 7 days of treatment, leading to a significantly reduced
body weight after 6 to 14 days of treatment (Fig. 2c). Assessment of motor function in the
rotarod test showed that α-IgLON5 AAB animals started to perform worse on Day 7 (non-
significant) and were significantly impaired in motor function on Day 14 (Fig. 2d).
To assess whether the pathological changes observed in α-IgLON5 AAB-treated animals were
caused by neurodegeneration, we determined the levels of neurofilament light chain (NfL;
indicator for axonal damage) and glial fibrillar acidic protein (GFAP; indicator for
neuroinflammation) in the serum of the animals at Day 14. α-IgLON5 animals had enhanced
serum NfL levels compared to all controls (Fig. 2e left), although this did not reach statistical
significance. No difference was detected in GFAP serum levels (Fig. 2e, right). Of note, our
measurements of NfL and GFAP levels in serum may lack the sensitivity achieved by
measuring their CSF levels, for which increased NfL was previously reported in α-IgLON5
patients [18].
We also performed staining for milk fat globule epidermal growth factor 8 (MFG-E8), an
opsonin for apoptotic cells that recognizes phosphatidylserine exposure (early marker of
apoptotic death), which increases with glial activation in mice with amyloid-beta (Aβ)
aggregation [22] and upon neuronal Tau aggregation *in vitro* [23]. Although some MFG-E8 staining
was evident in all groups, its levels were not affected by AAB treatment (Supplementary Fig.
S2b), suggesting that acute administration of AABs did not induce neuronal death at the
chosen AAB dose.
256

257 Together these results indicate a high pathogenic potential of the patient-derived α-IgLON5
258 AABs, resulting in an *in vivo* phenotype with in rapid onset of deficits in physical condition
259 (body weight loss) and motor performance of the treated animals.
260
α-IgLON5 AABs induce neuroinflammation and deregulate neuronal activity related
genes
The rather rapid decline of motor function in α-IgLON5 AAB treated animals suggested that
α-IgLON5 AABs may have caused cellular and/or molecular changes in the brain directly
related to brain function. We tested whether α-IgLON5 AABs induced neuroinflammation,
which could explain the rapid onset and progression of symptoms characteristic for different
types of AE [1], including α-IgLON5 disease.
To this end, we analyzed astrocytosis (GFAP) and microgliosis (Iba1) in brain regions that are
affected in α-IgLON5 disease through fluorescence intensity. α-IgLON5 animals showed
higher GFAP and Iba1 intensities in several brain regions - including hippocampus,
hypothalamus, cerebellum, medial habenula, and periaqueductal gray - compared to PBS and
mCtrl treated-animals (Fig. 3a-d; Supplementary Fig. S2c, d). pCtrl but not mCtrl infusion also
led to increased GFAP and Iba1 immunoreactivity in some brain areas, particularly in cortex
and hypothalamus. This indicated that infusion of polyclonal human antibodies *per se* can
trigger an inflammatory response in mice. Nevertheless, α-IgLON5 AABs seemed to
additionally cause pathogenic effects, particularly in regions with high IgLON5 protein
expression such as the hippocampus.
The impaired motor function observed in α-IgLON5 AAB treated animals (Fig. 2d) could be
caused by inflammation or other pathological changes in the spinal cord. We therefore
investigated glia cell levels in spinal cord gray matter by immunofluorescence, however, we
did not find a difference between treatment groups (Supplementary Fig. S2e, f).
Notably, in a test cohort that received a higher AAB dose (total amount of 200 μg AAB; 8 μg/g
body weight), we also observed weight loss, motor impairments (rotarod at Day 14), and
increased microglia (Iba1) and astrocyte (GFAP) activation in brain and spinal cord, which
were, however, similar in both α-IgLON5 AAB and pCtrl-infused animals (Supplementary Fig.
S4). This suggested that higher doses of patient-derived AABs are *per se* toxic when
administered into mice.
288
289 To further resolve molecular changes induced by α-IgLON5 AABs vs. pCtrl, we performed
290 bulk RNAseq. To this end, we used contralateral cortices animals since cells in ipsilateral
291 cortices are compromised by the surgery. α-IgLON5 treated animals had several genes that

were significantly up- or downregulated compared to pCtrl animals: genes associated with neuroinflammatory response (e.g., *CXCL10, CXCL16, CCL3, C3,* and *GFAP*) were up-regulated, whereas genes for ion channels (e.g., *SLC10A4, KCNA5,* and *SCN5A*), the dopaminergic system (e.g., *DRD1, DRD2,* and *PPP1R1B*), and cell-cell junction proteins and surface receptors (*NEXN, PDLIM3, NTRK1,* and *TACR1*) were down-regulated (Fig. 3e-g; Supplementary Fig. S3; Supplementary Table T1). We confirmed the upregulation of certain neuroinflammation-associated genes on the protein level by immunostaining (Fig. 3h). Interestingly, MHC class II proteins (*CD74, H2-Aa H2-Eb1* and *H2-Ab1*) and CD8 receptor activity related genes (*CD274* and *CCL5*) were also upregulated in α-IgLON5 animals. Immunostaining of hippocampal sections confirmed a small number of CD8-positive (CD8+) T cells in the hippocampus of α-IgLON5 but not pCtrl infused mice. Infiltration of CD8+ T cells was previously suggested to be triggered by Tau pathology and to mediate neurotoxicity in human Tau transgenic mice [24]. Therefore, α-IgLON5 AABs may trigger a similar Tau related pathological cascade.

Together, these data show that, within 2 weeks, α-IgLON5 AABs induce a distinct pro-inflammatory and neuronal activity-modulating signature in the brain. Additionally, we observed a general glia cell response (increase in Iba1 and GFAP) induced by intrathecal infusion of human autoantibodies (pCtrl and α-IgLON5 AABs) into the mouse.

α-IgLON5 AABs induce Tau phosphorylation in selected hippocampal projections

A characteristic pathological change in the brains of α-IgLON5 patients is the accumulation of p-Tau in the somatodendritic compartment of neurons in the hippocampus and other brain regions [3,7]. However, whether AABs against IgLON5 are directly responsible for Tau changes is not clear. We therefore evaluated whether the cerebroventricular infusion of α-IgLON5 AABs into wildtype mice was sufficient to initiate Tau phosphorylation and accumulation in the brain.

Western Blot analysis of Tau in lysates from spinal cord, fimbria, and cerebellum suggested no differences in p-Tau between groups (p-Tau pS202/pT205 (AT8), pS396/pS404 (PHF-1), and pT231 (AT180) epitopes); Supplemental Fig. S5a), possibly because the localized p-Tau signal was diluted in bulk tissue analysis.

To refine our analysis and detect subtle and spatially restricted changes in Tau phosphorylation, we performed immunofluorescence labeling of p-Tau in brain slices of treated mice. These experiments revealed p-Tau pS396/pS404 (PHF-1) immunoreactivity in the brains of α-IgLON5 AAB but not control (PBS, mCtrl, pCtrl) mice, selectively in (i) cells of the dentate gyrus, (ii) in mossy fiber (MF) projections onto CA3, (iii) in bi-directional projections

on ventral CA1 (vCA1) but not dorsal CA1 (dCA1) regions, and (iv) in commissural fibers (CFs)

of corpus callosum (CC) (Fig. 4a,b). In other brain regions we did not observe p-Tau

pS396/pS404. For p-Tau pS202/pT205 (AT8 epitope), we observed no immunoreactivity in

hippocampal brain regions, suggesting that α-IgLON5 AABs may have triggered

phosphorylation of selected epitopes on Tau. Comparing the levels of p-Tau pS396/pS404

and neuroinflammation in the hippocampus of α-IgLON5 animals, we found a correlation

between p-Tau and microglia (Iba1) for α-IgLON5 but not in any of the control groups (Fig. 4c;

Supplemental Fig. S5b). No correlation was detected between p-Tau and astrocytes (GFAP).

Notably, in the spinal cord (ventral horn), motor neuron cell bodies had more p-Tau

pS202/pT205 in α-IgLON5 AAB compared to pCtrl infused mice (not quantified; Supplemental

Fig. S5c).

To better understand the pathogenic potential and cellular mechanism of α-IgLON5 AAB-

induced Tau accumulation, we performed experiments in primary hippocampal mouse

neurons. Neurons treated with α-IgLON5 AABs for 2 days showed a dose-dependent Tau

accumulation in the somatodendritic compartment at 0.1, 1, and 10 μg/ml AABs (Fig. 4d, e),

notably in the absence of overt neuronal toxicity or death in the cultures, reflecting our findings

in vivo. At high concentrations (50 μg/ml), α-IgLON5 AABs induced fragmentation of neuronal

processes, indicative of toxicity.

The levels of p-Tau (AT8, PHF-1, AT180, pS199 epitopes) in neuronal lysates were similar for

α-IgLON5 and Ctrl AAB treated cultures (Supplemental Fig. S5d, e), probably because of a

high Tau phosphorylation level in cultured neurons at baseline, which commonly renders

Western blot analysis of bulk p-Tau levels in neuronal lysates insensitive to changes.

Interestingly, when removing α-IgLON5 AABs after 2 days, the Tau levels in neuronal cell

bodies remained similar to conditions of continuous treatment with AABs (Fig. 4f), suggesting

that Tau – once activated and resorted – may reside in the somatodendritic compartment for

multiple days.

Together these data show that the infusion of α-IgLON5 AABs for two weeks is sufficient to

induce hippocampal glia activation, as well as localized and epitope-specific (pS396/pS404)

phosphorylation and dendritic accumulation/resorting of Tau in dentate gyrus granule cells

and distinct projections in the hippocampus. These pro-pathological Tau changes may

represent the earliest changes occurring in α-IgLON5 disease. Notably, we observed these

Tau changes in wildtype mice, which are normally rather resilient to pathological endogenous

mouse Tau changes demonstrating the strong pathogenic potential of α-IgLON5 AABs across

species.

In vitro, at doses between 0.1 to 1 µg/ml, patient-derived α-IgLON5 AABs directly triggered

prolonged somatodendritic resorting of Tau, a typical reaction in neuronal stress response [25].

Higher doses of α-IgLON5 AABs were neurotoxic. These observations suggest that low

concentrations of α-IgLON5 AABs could trigger continued stress response and

somatodendritic p-Tau resorting in the brains of α-IgLON5 patients.

Acute neuronal hyperactivity drives α-IgLON5 AABs Tau changes

Neuronal hyperactivity, for example during epileptic activity [26], induced by Aβ oligomers [27], or

through chemical activity stimulation (e.g., using glutamate [28]) - is a known driver of Tau

phosphorylation and somatodendritic resorting. Given that we observed similar Tau changes

upon treatment with α-IgLON5 AAB, we asked whether those were linked to neuronal activity

as well. IgLON5 has not previously been associated with modulating neuronal excitability.

Imaging of spontaneous calcium (Ca^{2+}) transients in GCamp6f expressing hippocampal

neurons (DIV13-15; Fig. 5a) treated with a low dose of AABs (0.1 µg/ml) revealed a significant

increase in neuronal activity upon exposure to α-IgLON5 compared to control antibodies (Fig.

5b-e). This was apparent from an increased average spike rate in cultures treated with α-

IgLON5 AABs compared to pCtrl and mCtrl treated cultures after 15 min (average spikes/min

at 15 min (mean±SEM): α-IgLON5=5.0±0.1, pCtrl=4.0±0.1, mCtrl=3.2±0.1 spikes/min;

N=1095 to 1430 neurons per condition) and after 60 min (average spikes/min: α-

Iglon5=5.8±0.1, pCtrl=5.1±0.1, mCtrl=4.8±0.1 spikes/min; N=1199 to 1379 neurons per

condition; Fig. 5d). Therefore, the fraction of hyperactive neurons in the cultures increased

from ~3% before treatment to ~9% at 15 min and ~11% at 60 min after α-IgLON5 AAB

treatment (Fig. 5e; for definition of activity classes see the figure legend). mCtrl and pCtrl

treated neurons showed a less pronounced increase of hyperactive neurons (mCtrl: 5% at

15min, 7% at 60 min; pCtrl: 7% at 15 min, 7% at 60 min).

Notably, this effect was dose-dependent, because at a higher dose of 1 µg/ml AABs, α-

IgLON5 AAB treatment induced a drastic increase in neuronal activity (2.6±0.1 spikes/min

(mean±SEM), ~50% hyperactive neurons, N=309 neurons) compared to pCtrl (1.9±0.1

spikes/min, ~11% hyperactive neurons, N=263 neurons) and untreated control neurons (Fig.

5f,g; Supplemental Fig. S6a). After 2 days, we detected mild toxicity in α-IgLON5 AAB treated

neurons (LDH assay; Supplemental Fig. S6c) and the average activity dropped again to

baseline (=before treatment) (Supplemental Fig. S6b).

Next, we wondered whether the reduction in neuronal activity seen after 2 days of α-IgLON5

AAB treatment was due to the loss of functional synapses. To this end, we performed

electrophysiological recordings in autaptic hippocampal neuron cultures (individual neurons

grown on astrocytic micro-islands, forming synapses only with themselves, Supplemental Fig.

S6d) treated with AABs (0.1 μg/ml) for 3 days, which showed no difference between treatment

groups for all measured parameters (membrane capacitance, resting membrane potential,

excitatory postsynaptic current (EPSC) amplitude, EPSC paired-pulse ratio, readily-releasable

pool size, synaptic vesicle release probability, EPSC release during 10 Hz stimulation;

Supplemental Fig. S6e). That we detected neither a reduction in EPSC amplitude, nor in

RRPM size indicated that no synapse loss occurred upon 3 days treatment with α-IgLON5

AABs.

To test whether α-IgLON5 AAB-induced neuronal activity was the actual trigger for the Tau

resorting observed after α-IgLON5 AAB treatment, we removed Ca^{2+} ions from the medium

by chelating them with EGTA-AM [29]. This treatment suppressed α-IgLON5 AAB-induced

neuronal hyperactivity in a dose dependent manner (Supplementary Fig. S6f-h), and α-

IgLON5 AAB-induced somatodendritic Tau re-sorting was no longer observed upon treatment

with EGTA-AM (10 μM for 20 min) (Fig. 5h). Therefore, neuronal activity and/or activity induced

Ca^{2+} signaling was necessary to activate Tau somatodendritic resorting upon α-IgLON5 AAB

exposure.

Since we identified neuronal hyperactivity as the driver of Tau resorting induced by α-IgLON5

AABs, we hypothesized that other AABs inducing high neuronal activity would induce a similar

effect on Tau resorting. To test this idea, we treated neurons with a human monoclonal AAB

targeting LGI1 (α-LGI1 AAB), an epilepsy-related protein that regulates excitatory synapses

and stabilizes axonal voltage-gated potassium channels [30–32]. We previously showed that α-

LGI1 AABs increase cellular excitability and excitatory synaptic transmission of hippocampal

CA3 neurons [33]. Here we found that the treatment of neuronal cultures with 1 μg/ml

monoclonal α-LGI1 AAB induced Tau resorting similar to α-IgLON5 AABs (Fig. 5i), suggesting

that α-LGI1 induced neuronal hyperactivity may also induced Tau activation.

Together, these observations show that the binding of α-IgLON5 AABs induces acute

hyperexcitation (within minutes) that triggers lasting somatodendritic Tau resorting (for days)

and a rebalancing of neuronal physiology after transient hyperactivity. When becoming

chronic, for example upon extended or repeated exposure of the brain to α-IgLON5 AABs,

these adaptations to α-IgLON5 AAB-induced acute hyperexcitation could lead to synaptic loss

and neurotoxicity, resulting in a reduction in the overall neuronal network activity (Fig. 5j). In

fact, long-term treatment of cultured neurons with α-IgLON5 AABs for multiple weeks has

been shown to result in a decrease of synapse numbers [19]. In summary, our data identify acute

neuronal hyperactivity as the driver of pro-pathological Tau changes in α-IgLON5 disease –
and possibly other AAB-induced neurological conditions.
435

Discussion

Autoimmune encephalitis (AEs) disorders are characterized by neuroinflammation and
neurodegeneration that are induced by AABs targeting brain antigens [2]. AE patients show a
range of neurological symptoms, including seizures, psychiatric manifestations, and cognitive
deficits. Several AABs against neuronal receptors and cell surface proteins - e.g., NMDA
receptors, GABA$_A$ receptors, and LGI1 - lead to AE diseases by altering neuronal physiology
and inducing neuroinflammation [1,33–35].

Our data now reveal that α-IgLON5 encephalitis – a unique tauopathy – is caused by a similar
neuronal activity-related disease mechanism: plasma from an α-IgLON5 encephalitis patient
containing a polyclonal pool of α-IgLON5 AABs was able to trigger acute neuronal
hyperactivity. At high AAB doses or longer exposure, α-IgLON5 AABs led to neurotoxicity *in*
vitro. *In vivo*, these AABs induced neuroinflammation, motor deficits, and body weight loss.
Both neuronal silencing and inflammation were also affected on the gene expression level,
whereby animals that received α-IgLON5 AABs showed a distinct activation of complement 3,
MHC class II, and T cell receptor pathways. Similar responses of the innate immune system
were previously reported in transgenic mouse models of Tau pathology [24,36,37], indicating that
α-IgLON5 AABs induce an innate immune response that is typical for pro-pathological Tau
changes in the brain. Notably, this neuroinflammatory Tau signature occurred in parallel to p-
Tau accumulation, but not aggregation, in selected neuronal projections. Therefore, early Tau
changes – as observed in α-IgLON5 AAB treated animals - seem sufficient to induce a distinct
glia and immune cell response, which in turn may further accelerate Tau pathology and
neuronal damage.
Notably, the cerebroventricular infusion of AABs in the brain – as it was used in this study - is
currently limited to 2 weeks. Therefore, effects of long-term (or repeated) treatment with α-
IgLON5 AABs on Tau, neuroinflammation, and neurodegeneration are limited and new *in vivo*
models mirroring extended immune and activity stimulation in α-IgLON5 disease are required.
462
We identified α-IgLON5 AAB-induced neuronal hyperactivity as the driver of pro-pathological
Tau changes (somatodendritic p-Tau), a hallmark in α-IgLON5 disease. A similar
hyperactivity-induced Tau "activation", including Tau phosphorylation and resorting, may
occur in other AAB-induced neurological conditions as well – in this study we show similar

effects for α-LGI1 AABs-, suggesting a common route to Tau pathology in AEs. Indeed,

somatodendritic accumulation of p-Tau, accompanied by hippocampal atrophy, was found in

LGI1 patients [38]. It should be investigated whether Tau pathology in other immune-related

secondary tauopathies, e.g., subacute sclerosing panencephalitis (SSPE [39]; measles late

reaction) and head-nodding syndrome [40] (involving DJ-1 antibodies), occurs in response to

antibody-triggered neuronal hyperactivity as well.

Neuronal hyperactivity was previously suggested to also trigger Tau pathology in AD [41–43] and

in epilepsy [44]. The herein discovered route of action for α-IgLON5 AABs now adds another

example for activity-dependent Tau pathology, and, thereby, emphasizes the importance to

consider the modulation of neuronal activity for therapeutic approaches towards preventing or

lowering Tau pathology and toxicity in the brain. Next, it will be important to identify the

molecular mechanisms of how AAB binding to the surface protein IgLON5 can induce neuronal

activity, e.g., receptors and channels involved.

To which extent activity-induced Tau changes and excitotoxicity contribute to neurotoxicity in

α-IgLON5 disease and other AEs is unclear. Our data show that the effects of α-IgLON5 AABs

on neuronal function and Tau depends on AAB concentration and correlates with the amount

of hyperexcitation *in vitro*. Mice that received higher AAB doses showed stronger

neuroinflammation and motor symptoms. We propose that speed and extension of neuronal

dysfunction in AEs that are triggered by hyperexcitation-promoting AABs may depend on the

interplay between acute (minutes to hours) excitotoxicity and mid-term (days to weeks) Tau

toxicity, i.e., extreme, toxic hyperexcitation resulting in fast neuronal death versus moderate,

non-toxic hyperexcitation leading to Tau pathology and (slower) neuronal dysfunction.

Following this idea, hyperexcitation is apparently not immediately toxic in α-IgLON5 disease

and allows for the gradual development of pathological Tau changes. This would also mirror

the clinical observation that immunotherapy can be beneficial when initiated early in α-IgLON5

disease [5]. The extended presence of p-Tau in the somatodendritic compartment, however,

can exhaust the neuronal homeostasis network [45], induce synapse loss [46], or result in the

impairment of other key neuronal functions [47–49]. These Tau effects can ultimately induce

neurotoxicity also in α-IgLON5 disease.

In contrast, in α-LGI1 or α-GABA receptor AEs, extreme hyperexcitation may lead to

immediate neurotoxicity, so that a manifestation of later Tau pathology is less obvious or rare

in these diseases. For the treatment of patients with hyperexcitation-inducing AABs, like α-

IgLON5 and α-LGI1 AABs, determining the neuromodulatory activity of their AAB pool could

help in the choice of treatment, whether to target neuronal hyperexcitation, Tau
phosphorylation and aggregation, or both.

Pathological Tau changes, including high phosphorylation, somatodendritic resorting, and
later aggregation, occur in multiple neurological conditions. In neurodegenerative disease like
AD and FTD, the location and extent of Tau pathology correlates with neuronal loss in the
brain [8]. However, changes in Tau, although commonly classified as pathological, do not
necessarily lead to immediate neuronal dysfunction and death [50,51]. They are, however,
correlated with a progressive decay in synaptic homeostasis [46,52–54], ultimately leading to
connectivity deficits and neuronal loss in AD and tauopathies. For α-IgLON5 AAB-induced
Tau changes, a similar course of Tau action appears to apply. Long-term exposure to α-
IgLON5 AABs (for weeks) was shown to result in neuronal p-Tau accumulation, disrupted
synaptic protein homeostasis, decay in neuronal activity, and cell death, and caused cognitive
deficits and anxiety-like behavior in mice [19,55,56]. These pathological changes can be
interpreted as consequences of prolonged Tau activation, as they occur similarly in
progressive neurodegenerative diseases like AD. Interestingly, we found that the activation of
Tau (in form of somatodendritic re-sorting) lasted for multiple days even after removal of α-
IgLON5 AABs, suggesting that activated Tau may play a role in a prolonged neuronal activity
stress response. Whether a "deactivation" of Tau is facilitated by sorting it back to the axon or
relies on Tau removal through degradation remains to be clarified. It also remains to be
determined whether the persistence of the Tau response could be responsible for poor clinical
response of the patient to the immunosuppression therapy, and whether Tau activation
persists similarly in other AE disorders, e.g., LGI1 disease.

Our study shows that α-IgLON5 AABs activate Tau in selected hippocampal neural
projections. p-Tau accumulation occurred in somata of DG granule cells and their MF
projections onto CA3, which are important structures for learning and memory, and in
commissural fibres in corpus callosum which are output structures of CA3. Similar
hippocampal connections were recently suggested to be affected by α-IgLON5 AABs [55,56].
Additionally, we identified projections onto ventral CA1 (vCA1) and commissural fibres –
probably coming from CA3 neurons [57,58] - as the main responders to α-IgLON5 AAB induced
Tau phosphorylation. In mice, vCA1 neurons project to the entorhinal cortex, olfactory cortex,
and amygdala, and are involved in anxiety, social, and motivational (e.g., feeding) behavior [57].
One may speculate that the accumulation of p-Tau in projections onto vCA1 could therefore
correlate with anxiety-like behavior previously reported for mice treated with α-IgLON5 AABs
[55], and could also be related to the rapid body weight loss we observed in such animals.

Notably, the hippocampal connections affected by α-IgLON5 AABs are similar to the ones

affected by Tau in AD, where Tau pathology starts in the entorhinal cortex and spreads

through the perforant pathway into DG granule cells and further through MFs into CA3 [59–61].

Furthermore, the anterior human hippocampus, which corresponds to the ventral

hippocampus in mice, shows metabolic changes and neuronal atrophy early in AD [62–64]. In

mice with α-IgLON5 AAB treatment, only projections onto CA1 of the ventral hippocampus

(vCA1) accumulated p-Tau. Considering that α-IgLON5 AABs activate Tau via neuronal

hyperactivity, this could be explained by the generally higher excitability of ventral compared

to dorsal CA1 hippocampal neurons [65]. Extending the link between neuronal hyperexcitation

and pro-pathological Tau activation, susceptibility of neurons to hyperexcitation could encode

the selective vulnerability of neuronal circuits to Tau pathology not just in α-IgLON5 disease

but also in AD and other tauopathies. In fact, hippocampal hyperactivity has been described

in patients with preclinical AD and mild cognitive impairment, stages of the disease where p-

Tau starts to accumulate in this region [66]. Similarly, recent MRI and PET imaging identified

that Aβ induces hyperactivity of the default mode network which propagates to the medial

temporal lobe and predicts Tau accumulation in early stages of AD [43,66].

Later in the disease progression, upon prolonged hippocampal Tau phosphorylation and

aggregation, synapse loss and a decline in neuronal activity can be observed. The herein

revealed mechanism, whereby α-IgLON5 AAB-induced neuronal hyperactivity is responsible

for pro-pathological Tau changes, therefore has important implications for the understanding

and treatment of Tau-related neurological conditions, far beyond α-IgLON5 disease. Future

work will need to identify how α-IgLON5 AAB binding induces neuronal hyperactivity on the

molecular level, whether these mechanisms are unique for α-IgLON5 AABs or can be applied

more general to different autoimmune antibodies, whether reversibility of the hyperexcitable

state can interrupt the Tau pathology cascade, and whether and when immunotherapy can

reverse progression of both autoimmunity and neurodegeneration.

Acknowledgement

565   We thank Britta Eickholt for providing us with access to instruments and equipment of the

566   Charité Biochemistry Department, the Advanced Medical Bioimaging Core Facility (AMBIO)

567   of the Charité, which enabled us to perform most microscopy on brain sections and neuron/cell

568   cultures, the Viral Core Facility (VCF) of the Charité for virus production, and the BIH

569   sequencing core for determining the RNA content of mouse brain tissue.

Funding

The work was funded by the German Center for Neurodegenerative diseases (DZNE) in the Helmholtz Society (to S.W. and H.P.), the Einstein Foundation Berlin (to B.A.), and the Deutsche Forschungsgemeinschaft (DFG; KFO 5023 BECAUSE-Y; Project 504745852 (to S.W. and H.P.). PR1274/5-1, PR1274/9-1, and PR1274/13-1 (to H.P.), the Helmholtz Association (HIL-A03 BaoBab to H.P.), and by the German Federal Ministry of Education and Research (Connect-Generate 01GM1908D, to H.P.). B.R.R. was supported by the Deutsche Forschungsgesellschaft (DFG; project 327654276 - SFB1315 and project 273915538 - SPP1926). H.-C.K. was supported by the German Research Council (DFG, FOR3004, KO2290/3-2).

Author contributions

Experiments/data. analysis of *in vitro* and *in vivo* α-IgLON5 AAB binding, Tau changes, neuroinflammation, RNAseq, Calcium imaging: Bilge Askin
Purification of IgLON5 AABs & Clinical experts: Alexander Goihl, Peter Körtvelyessy, Dirk Reinhold
Injection of mice and behavioral assessments: César Cordero Gómez
RNAseq analysis: Andrew Newman
Tissue processing, cryosectioning, immunohistochemistry (IHC) and IHC analysis: Cagla Kilic, Janne Lamberty, César Cordero Gómez, Sophie Lan-Linh Duong
NfL and GFAP ELISA: César Cordero Gómez
FACS for binding analysis: César Cordero Gómez
MFGE8 staining: Jessica Wagner, Jonas Neher
Epitope mapping and internalization experiments: Sophie Lan-Linh Duong
Establishing co-loc analysis pipelines and co-loc experiments: Valentine Thomas
Providing inputs for GCaMP experiments: Paul Turko
Autaptic neuron culture recordings: Benjamin R. Rost, Dietmar Schmitz
Supply of LGI1 AABs: Hans-Christian Kornau
Technical & neuron prep support: Sabrina Huebschmann, Ekaterina Semenova
General help, support, MS revision, discussion: Lisa Diez
Funding acquisition: Susanne Wegmann, Harald Prüss, Dietmar Schmitz, Benjamin R. Rost, Hans-Christian Kornau
Figure preparation: Bilge Askin, Susanne Wegmann, Cagla Kilic
Manuscript writing: Bilge Askin, Susanne Wegmann
Manuscript revision: all authors
Scientific concept: Susanne Wegmann, Harald Prüss, Bilge Askin, Peter Körtvelyessy,

References:

1. Prüss, H. Autoantibodies in neurological disease. *Nat Rev Immunol* **21**, 798–813 (2021).
2. Dalmau, J., Geis, C. & Graus, F. Autoantibodies to synaptic receptors and neuronal cell surface proteins in autoimmune diseases of the central nervous system. *Physiol Rev* **97**, 839–887 (2017).

3. Sabater, L. *et al.* A novel non-rapid-eye movement and rapid-eye-movement parasomnia with sleep breathing disorder associated with antibodies to IgLON5: A case series, characterisation of the antigen, and post-mortem study. *Lancet Neurol* **13**, 575–586 (2014).

4. Fu, Y., Zou, X. & Liu, L. Epileptic Seizures and Right-Sided Hippocampal Swelling as Presenting Symptoms of Anti-IgLON5 Disease: A Case Report and Systematic Review of the Literature. *Front Neurol* **13**, 1–9 (2022).

5. Grüter, T. *et al.* Clinical, serological and genetic predictors of response to immunotherapy in anti-IgLON5 disease. *Brain* **146**, 600–611 (2023).

6. Madetko, N. *et al.* Anti-IgLON5 Disease – The Current State of Knowledge and Further Perspectives. *Front Immunol* **13**, 1–8 (2022).

7. Gelpi, E. *et al.* Neuropathological criteria of anti-IgLON5-related tauopathy. *Acta Neuropathol* **132**, 531–543 (2016).

8. Spillantini, M. G. & Goedert, M. Tau pathology and neurodegeneration. *Lancet Neurol* **12**, 609–622 (2013).

9. Iqbal, K. *et al.* Tau pathology in Alzheimer disease and other tauopathies. *Biochim Biophys Acta Mol Basis Dis* **1739**, 198–210 (2005).

10. Zempel, H. Genetic and sporadic forms of tauopathies—TAU as a disease driver for the majority of patients but the minority of tauopathies. *Cytoskeleton* 1–5 (2023) doi:10.1002/cm.21793.

11. Vanaveski, T. *et al.* Promoter-Specific expression and genomic structure of Iglon family genes in mouse. *Front Neurosci* **11**, 1–15 (2017).

12. Ranaivoson, F. M. *et al.* A Proteomic Screen of Neuronal Cell-Surface Molecules Reveals IgLONs as Structurally Conserved Interaction Modules at the Synapse. *Structure* **27**, 893-906.e9 (2019).

13. Venkannagari, H. *et al.* Highly Conserved Molecular Features in IgLONs Contrast Their Distinct Structural and Biological Outcomes. *J Mol Biol* **432**, 5287–5303 (2020).

14. Landa, J. *et al.* Patients' IgLON5 autoantibodies interfere with IgLON5-protein interactions. *Front Immunol* **14**, 1–7 (2023).

15. Hashimoto, T., Maekawa, S. & Miyata, S. IgLON cell adhesion molecules regulate synaptogenesis in hippocampal neurons. *Cell Biochem Funct* **27**, 496–498 (2009).

16. Karis, K. *et al.* Altered expression profile of igLON family of neural cell adhesion molecules in the dorsolateral prefrontal cortex of schizophrenic patients. *Front Mol Neurosci* **11**, 1–12 (2018).

17. Sanz, R., Ferraro, G. B. & Fournier, A. E. IgLON Cell Adhesion Molecules Are Shed from the Cell Surface of Cortical Neurons to Promote Neuronal Growth. *J Biol Chem* **290**, 4330 (2015).

18. Landa, J. *et al.* Effects of IgLON5 Antibodies on Neuronal Cytoskeleton: A Link between
Autoimmunity and Neurodegeneration. *Ann Neurol* **88**, 1023–1027 (2020).

19. Ryding, M. *et al.* Neurodegeneration Induced by Anti-IgLON5 Antibodies Studied in
Induced Pluripotent Stem Cell-Derived Human Neurons. *Cells* **10**, (2021).

20. Sabater, L., Planagumà, J., Dalmau, J. & Graus, F. Cellular investigations with human
antibodies associated with the anti-IgLON5 syndrome. *J Neuroinflammation* **13**, 1–12
(2016).

21. Wenke, N. K. *et al.* N-methyl-D-aspartate receptor dysfunction by unmutated human
antibodies against the NR1 subunit. *Ann Neurol* **85**, 771–776 (2019).

22. Wagner, J. *et al.* Medin co-aggregates with vascular amyloid-β in Alzheimer's disease.
Nature **612**, 123–131 (2022).

23. Brelstaff, J., Tolkovsky, A. M., Ghetti, B., Goedert, M. & Spillantini, M. G. Living Neurons
with Tau Filaments Aberrantly Expose Phosphatidylserine and Are Phagocytosed by
Microglia. *Cell Rep* **24**, 1939-1948.e4 (2018).

24. Chen, X. *et al.* Microglia-mediated T cell infiltration drives neurodegeneration in tauopathy.
Nature 2023 615:7953 **615**, 668–677 (2023).

25. Zempel, H. & Mandelkow, E. Lost after translation: Missorting of Tau protein and
consequences for Alzheimer disease. *Trends in Neurosciences* vol. 37 721–732 Preprint
at https://doi.org/10.1016/j.tins.2014.08.004 (2014).

26. Gao, Y. *et al.* Targeted Reducing of Tauopathy Alleviates Epileptic Seizures and Spatial
Memory Impairment in an Optogenetically Inducible Mouse Model of Epilepsy. *Front Cell
Dev Biol* **8**, 1–10 (2021).

27. Zempel, H., Thies, E., Mandelkow, E. & Mandelkow, E. M. Aβ oligomers cause localized
Ca2+ elevation, missorting of endogenous Tau into dendrites, Tau phosphorylation, and
destruction of microtubules and spines. *Journal of Neuroscience* **30**, 11938–11950 (2010).

28. Kobayashi, S., Tanaka, T., Soeda, Y. & Takashima, A. Enhanced Tau Protein Translation
by Hyper-Excitation. *Front Aging Neurosci* **11**, (2019).

29. Hardingham, G. E., Arnold, F. J. L. & Bading, H. A calcium microdomain near NMDA
receptors: On switch for ERK-dependent synapse-to-nucleus communication. *Nat
Neurosci* **4**, 565–566 (2001).

30. Seagar, M. *et al.* LGI1 tunes intrinsic excitability by regulating the density of axonal Kv1
channels. *Proc Natl Acad Sci U S A* **114**, 7719–7724 (2017).

31. Fukata, Y. *et al.* Epilepsy-related ligand/receptor complex LGI1 and ADAM22 regulate
synaptic transmission. *Science* **313**, 1792–1795 (2006).

32. Schulte, U. *et al.* The epilepsy-linked Lgi1 protein assembles into presynaptic Kv1
channels and inhibits inactivation by Kvβ1. *Neuron* **49**, 697–706 (2006).

33. Kornau, H. C. *et al.* Human Cerebrospinal Fluid Monoclonal LGI1 Autoantibodies Increase
Neuronal Excitability. *Ann Neurol* **87**, 405–418 (2020).

34. Kreye, J. *et al.* Encephalitis patient-derived monoclonal GABAA receptor antibodies cause
epileptic seizures. *J Exp Med* **218**, (2021).

35. Ceanga, M. *et al.* Human NMDAR autoantibodies disrupt excitatory-inhibitory balance,
leading to hippocampal network hypersynchrony. *Cell Rep* **42**, (2023).

36. Wu, T. *et al.* Complement C3 Is Activated in Human AD Brain and Is Required for
Neurodegeneration in Mouse Models of Amyloidosis and Tauopathy. *Cell Rep* **28**, 2111-
2123.e6 (2019).

37. Dejanovic, B. *et al.* Complement C1q-dependent excitatory and inhibitory synapse
elimination by astrocytes and microglia in Alzheimer's disease mouse models. *Nature
Aging 2022 2:9* **2**, 837–850 (2022).

38. Day, G. S. *et al.* Flortaucipir (tau) PET in LGI1 antibody encephalitis. *Ann Clin Transl
Neurol* **8**, 491–497 (2021).

39. Bancher, C. *et al.* On the relationship between measles virus and Alzheimer neurofibrillary
tangles in subacute sclerosing panencephalitis. *Neurobiol Aging* **17**, 527–533 (1996).

40. Pollanen, M. S. *et al.* Nodding syndrome in Uganda is a tauopathy. *Acta Neuropathol* **136**,
691 (2018).

41. Styr, B. & Slutsky, I. Imbalance between firing homeostasis and synaptic plasticity drives
early-phase Alzheimer's disease. *Nat Neurosci* **21**, 463–473 (2018).

42. Lee, W. J. *et al.* Regional Aβ-tau interactions promote onset and acceleration of
Alzheimer's disease tau spreading. *Neuron* **110**, 1932-1943.e5 (2022).

43. Giorgio, J., Adams, J. N., Maass, A., Jagust, W. J. & Breakspear, M. Amyloid induced
hyperexcitability in default mode network drives medial temporal hyperactivity and early
tau accumulation. *Neuron* (2023) doi:10.1016/J.NEURON.2023.11.014.

44. Tai, X. Y. *et al.* Hyperphosphorylated tau in patients with refractory epilepsy correlates with
cognitive decline: a study of temporal lobe resections. *Brain* **139**, 2441–2455 (2016).

45. Papanikolopoulou, K. & Skoulakis, E. M. C. Altered Proteostasis in Neurodegenerative
Tauopathies. *Adv Exp Med Biol* **1233**, 177–194 (2020).

46. Yoshiyama, Y. *et al.* Synapse Loss and Microglial Activation Precede Tangles in a P301S
Tauopathy Mouse Model. *Neuron* **53**, 337–351 (2007).

47. Meier, S. *et al.* Pathological tau promotes neuronal damage by impairing ribosomal
function and decreasing protein synthesis. *Journal of Neuroscience* **36**, 957–962 (2016).

48. Vanderweyde, T. *et al.* Interaction of tau with the RNA-Binding Protein TIA1 Regulates tau
Pathophysiology and Toxicity. *Cell Rep* **15**, 1455–1466 (2016).

49. Eftekharzadeh, B. *et al.* Tau Protein Disrupts Nucleocytoplasmic Transport in Alzheimer's
Disease. *Neuron* **99**, 925-940.e7 (2018).

50. Kuchibhotla, K. V. *et al.* Neurofibrillary tangle-bearing neurons are functionally integrated
in cortical circuits in vivo. *Proc Natl Acad Sci U S A* **111**, 510–514 (2014).

51. Arendt, T. *et al.* Reversible paired helical filament-like phosphorylation of tau is an adaptive
process associated with neuronal plasticity in hibernating animals. *Journal of*
Neuroscience **23**, 6972–6981 (2003).

52. Henstridge, C. M., Pickett, E. & Spires-Jones, T. L. Synaptic pathology: A shared
mechanism in neurological disease. *Ageing Res Rev* **28**, 72–84 (2016).

53. Pickett, E. K. *et al.* Spread of tau down neural circuits precedes synapse and neuronal
loss in the rTgTauEC mouse model of early Alzheimer's disease. *Synapse* **71**, (2017).

54. Dejanovic, B. *et al.* Changes in the Synaptic Proteome in Tauopathy and Rescue of Tau-
Induced Synapse Loss by C1q Antibodies. *Neuron* **100**, 1322-1336.e7 (2018).

55. Ni, Y. *et al.* Anti-IgLON5 antibodies cause progressive behavioral and neuropathological
changes in mice. *J Neuroinflammation* **19**, 1–18 (2022).

56. Alvente, S. *et al.* Pilot Study of the Effects of Chronic Intracerebroventricular Infusion of
Human Anti-IgLON5 Disease Antibodies in Mice. *Cells* **11**, (2022).

57. Masurkar, A. V. Towards a circuit-level understanding of hippocampal CA1 dysfunction in
Alzheimer's disease across anatomical axes. *J Alzheimers Dis Parkinsonism* **8**, 412–412
(2018).

58. Dong, H. W., Swanson, L. W., Chen, L., Fanselow, M. S. & Toga, A. W. Genomic-anatomic
evidence for distinct functional domains in hippocampal field CA1. *Proc Natl Acad Sci U S*
A **106**, 11794–11799 (2009).

59. Hyman, B. T., Kromer, L. J. & Van Hoesen, G. W. A direct demonstration of the perforant
pathway terminal zone in Alzheimer's disease using the monoclonal antibody Alz-50. *Brain*
Res **450**, 392–397 (1988).

60. Thal, D. R. *et al.* Alzheimer-related tau-pathology in the perforant path target zone and in
the hippocampal stratum oriens and radiatum correlates with onset and degree of
dementia. *Exp Neurol* **163**, 98–110 (2000).

61. De Calignon, A. *et al.* Propagation of Tau Pathology in a Model of Early Alzheimer's
Disease. *Neuron* **73**, 685–697 (2012).

62. Ouchi, Y. *et al.* Altered glucose metabolism in the hippocampal head in memory
impairment. *Neurology* **51**, 136–142 (1998).

63. Wang, L. *et al.* Changes in hippocampal volume and shape across time distinguish
dementia of the Alzheimer type from healthy aging. *Neuroimage* **20**, 667–682 (2003).

64. Greene, S. J., Killiany, R. J., Alzheimer, T., Disease Neuroimaging Initiative, S. & Lilly, E.
Hippocampal Subregions are Differentially Affected in the Progression to Alzheimer's
Disease. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary*
Biology **295**, 132–140 (2012).

65. Dougherty, K. A., Islam, T. & Johnston, D. Intrinsic excitability of CA1 pyramidal neurones
from the rat dorsal and ventral hippocampus. *J Physiol* **590**, 5707–5722 (2012).

66. Targa Dias Anastacio, H., Matosin, N. & Ooi, L. Neuronal hyperexcitability in Alzheimer's
disease: what are the drivers behind this aberrant phenotype? *Translational Psychiatry*
2022 12:1 **12**, 1–14 (2022).

67. Ly, L. T. *et al.* Affinities of human NMDA receptor autoantibodies: implications for disease
mechanisms and clinical diagnostics. *J Neurol* **265**, 2625–2632 (2018).

68. Jin, Y., Tam, O. H., Paniagua, E. & Hammell, M. TEtranscripts: a package for including
transposable elements in differential expression analysis of RNA-seq datasets.
Bioinformatics **31**, 3593–3599 (2015).

69. Patel, T. P., Man, K., Firestein, B. L. & Meaney, D. F. Automated quantification of neuronal
networks and single-cell calcium dynamics using calcium imaging. *J Neurosci Methods*
243, 26–38 (2015).

70. Rost, B. R. *et al.* Autaptic cultures of single hippocampal granule cells of mice and rats.
European Journal of Neuroscience **32**, 939–947 (2010).

Figures

776
777

Figure 1. Purification and verification of α-IgLON5 AABs from patient plasma. a, Principle of α-IgLON5 AAB isolation from clinically verified IgLON5 patient plasma using affinity purification on immobilized recombinant human IgLON5 Fc-chimera proteins. **b,** SDS-PAGE of isolated antibodies. **c,** Dot blot of recombinant IgLON5 and LGI protein (negative control) shows specific reactivity of the

isolated α-IgLON5 AABs against IgLON5. **d,** Illustration of human IgLON5 protein (336 amino acids)
consisting of a signal peptide for secretion (SP), three extracellular Ig domains (Ig1/2/3), and a GPI-
anchor (GPI) in the C-terminus, tethering IgLON5 to the plasma membrane. **e,** Binding of α-IgLON5
AABs in HEK293T cells overexpressing full-length IgLON5 protein or (combinations of) individual Ig
domains. Scale bar = 20 μm. **f,** FACS binding affinity characterization (EC$_{50}$ = 8 μg/ml; K$_D$=55.9 μM) of
α-IgLON5 AABs to HEK cells expressing full-length human IgLON5 (IgLON5+). Control patient IgGs
(pCtrl) did not show binding. Data shown as mean±SEM, N=3 independent experiments. **g,** α-IgLON5
AABs recognize endogenous IgLON5 in human neuronal stem cells (hNSCs). **h,** Immunoreactivity of
α-IgLON5 AABs compared to pCtrl in fresh frozen sagittal brain sections from an adult wildtype mouse.
i, Primary hippocampal mouse neurons (DIV12-15) treated with α-IgLON5 AABs show similar α-IgLON5
AABs binding to dendritic (MAP2+; white arrow heads) and axonal regions (MAP2-; pink arrow heads).
Data were obtained from three independent experiments shown as mean±SEM. Student's T-test. **j,**
Immunofluorescence shows similar partial co-localization of α-IgLON5 AAB with pre-synaptic
(Synapsin-1) and post-synaptic (PSD-95) markers, both in (i) non-permeabilized (~15%) and (ii)
permeabilized (~22%) primary mouse neurons. Data shown as mean±SEM, N=3 independent
experiments, Student T-test. **k,** Model: Polyclonal α-IgLON5 AABs from patient serum bind to different
Ig domains of IgLON5 on the neuronal surface.

Figure 2. Intraventricular infusion of α-IgLON5 and control AABs into wildtype mice. a,
Experimental design for the *in vivo* application of α-IgLON5 AABs. Wildtype mice were infused in the
right ventricle with α-IgLON5 AABs, control antibodies (pCtrl, mCtrl), or PBS. 75 μg AABs were infused
for 14 consecutive days with a flow rate of 0.25 μl/h. Body weight was monitored daily starting on Day
0. At Day 7 and Day 14, rotarod experiments were conducted and serum samples were collected before
animals were sacrificed. Brain and spinal cord tissues were dissected for later investigation by
immunofluorescence, and for protein and RNA analysis. **b,** Immunofluorescence images of coronal
mouse brain sections show the distribution of infused α-IgLON5 AAB and pCtrl in the brain after 14
days (fluorescence anti-human IgG secondary antibody). Inset shows labeling of cells in the
contralateral hippocampus in α-IgLON5 AAB infused brain. **c,** Body weight relative to Day 0 measured
during 14 days of treatment. Data shown as mean±SEM, N=3-6 animals per group, Multiple t-test
comparing IgLON5 and pCtrl at each time point. **d,** Rotarod performance (time spent on rod) of α-
IgLON5 AAB and pCtrl animals determined at Day 7 and Day 14. Data shown as mean±SEM, N=3-4
animals per group. Student T-test. **e,** Serum levels of NfL and GFAP determined by ELISA. Data shown
as mean±SEM, N=3-6 animals, with 3 technical replicates measured per animal.

a GFAP immunolabeling in coronal brain sections of AAB infused animals

b Quantification of GFAP immunolabeling by brain region

Cerebral Cortex | Hippocampus | Hypothalamus | Periaqueductal Gray | Medial Habenula | Cerebellum

c Iba1 immunolabeling in coronal brain sections of AAB infused animals

d Quantification of Iba1 immunolabeling by brain region

Cerebral Cortex | Hippocampus | Hypothalamus | Periaqueductal Gray | Medial Habenula | Cerebellum

e Cortical RNAseq

f Gene deregulation in α-IgLON5 AABs vs. pCtrl treated animals
(bulk RNAseq of cortical tissue)

Down: 83 genes   Up: 167 genes

g Selected deregulated pathways

h Immunofluorescence of selected upregulated neuroinflammation markers

Figure 3. Neuroinflammatory changes in brain upon α-IgLON5 AAB infusion. a, Representative
brain sections immunolabeled for GFAP (astrocytes). **b,** Quantification of GFAP intensity in different
regions of α-IgLON5 AAB and control brains. Data shown as mean±SEM, N=3-6 animals, 3-5 brain
sections per animal. One-way ANOVA with Tukey post-test. Location of brain regions is shown in
images below graphs (adapted from 3D-brain Explorer, Allen Mouse Brain Atlas). **c,** Representative
brain sections immunolabeled for Iba1 (microglia). **d,** Quantification of Iba1 intensity in different regions
of α-IgLON5 AAB and control brains. Data shown as mean±SEM, N=3-6 animals, 3-5 brain sections
per animal. One-way ANOVA with Tukey post-test. **e,** Heat map showing differential gene expression
in contralateral cortical tissue in individual animals from α-IgLON5, pCtrl and PBS groups **f,** Volcano
plots of differentially expressed genes in the cortex of α-IgLON5 AAB infused animals. **g,** Gene ontology
(GO) enrichment analysis of the differentially expressed genes showing deregulated pathways in α-
IgLON5 AAB infused animals. **h**, Immunolabeling of neuroinflammatory markers (C3, C1q, CD68 and
CD8) upregulated in α-IgLON5 vs. mCtrl AAB treated animals. White arrow heads indicate C3-positive
astrocytes (left panel) and CD8-positive T cells (right panel). For quantification of C1q fluorescence,
data shown as mean±SD, N=3-5 hippocampal brain sections per group. One-way ANOVA with Tukey
post-test. Scale bars as indicated.
847

Figure 4. α-IgLON5 AABs induce phosphorylation and somatodendritic resorting of Tau in hippocampal neurons. a, Representative images of p-Tau pS396/pS404 (PHF-1) immunoreactivity in α-IgLON5 AAB and control brains. Top: p-Tau accumulation in dorsal hippocampus in dentate gyrus (DG) granule cells (left; white arrow heads), mossy fiber (MF) projections onto CA3 (center; white arrowhead), and in commissural fiber (CF) tracks of the corpus callosum (CC; right, white arrowhead). No p-Tau is found in dorsal CA1 (dCA1). Bottom: p-Tau accumulation in ventral hippocampus in projections onto ventral CA1 (vCA1; white arrow head) and MF projections onto CA3 (white arrow head). Scale bars as indicated. **b,** Quantification of mean p-Tau pS396/pS404 (PHF-1) intensity in

projections onto CA3. Data presented as mean±SEM, N=3-6 animals. One-way ANOVA with Tukey

post-test. **c,** Correlation between hippocampal Iba1 (top) or GFAP (middle) intensities and p-Tau

pS396/pS404 in projections onto CA3 for α-IgLON5 animals (orange data points and linear fit) vs.

combined control groups (black-white data points and black line fit). Bottom: triple correlation with p-

Tau levels color-coded. Note that IgLON5 animals cluster in the top right of the graph with red/yellow

color coding, indicating elevated hippocampal GFAP and Iba1 levels and above-average p-Tau levels

in projection onto CA3 in animals that received α-IgLON5 AABs. Linear fit values are indicated. **d,**

Representative images of primary hippocampal neurons treated with different concentrations (1, 10,

and 50 $\mu g/\mu l$) of α-IgLON5 AABs and immunolabeled for MAP2 (blue), Tau (magenta), and DNA (DAPI,

white). White solid line and shaded ROI in non-treated neurons (NT) indicate ROIs manually defined

for measuring the intensity of Tau in the soma. At 50 $\mu g/\mu l$**,** fragmentation of neuronal processes (white

arrow heads) indicated neurotoxicity. **e,** Quantification of Tau re-sorting into the soma in primary

neurons after treatment with 0.1, 1, or 10 $\mu g/ml$ α-IgLON5 AABs and controls for 2 days. N=3

independent experiments, data presented as mean±SEM, One-way ANOVA with Tukey post-test **f,**

Quantification of Tau re-sorting into the soma in primary neurons after treatment with 1 $\mu g/ml$ α-IgLON5

AAB and control antibodies for 5 days (bars without pattern fill) or after 2 days antibody treatment and

3 days recovery (bars with pattern fill). Data collected from 2 independent experiments and presented

as mean±SEM. One-way ANOVA with Tukey post-test.

Figure 5. α-IgLON5 AAB-induced neuronal hyperactivity. a, Construct design for genetically encoded Ca²⁺ indicator (GCamp6f). **b** Representative z-projection (sum) of Ca²⁺ recordings, individual activity traces and raster plots for detected calcium spikes after 15 min 0.1 μg/ml α-IgLON5 and **c,** pCtrl treatment. **d,** Frequency (transients/min) of spontaneous Ca²⁺ transients in neurons before (0 min) and after 0.1 μg/ml AAB treatment for 15 min and 60 min. NT = untreated neurons Violin plot (left) shows data distribution in treatment groups. Line plot (right) shows mean±SEM for each group and time point. N=3 independent experiments with 1094 to 1379 neurons per condition and time point. One-way ANOVA with Tukey post-test. **e,** Percentage (%) of inactive (blue), active (green), and hyperactive (pink)

neurons in cultures treated with 0.1 μg/ml α-IgLON5 AABs or controls for 0, 15, or 60 min. Hyperactive

criteria: Number of transients/min > mean+2*SD of pCtrl at each recording time point. Active: 0 <

number of transients/min < mean+2*SD of pCtrl at each recording time point. Silent: no transients. **f,**

Frequency (transients/min) of spontaneous Ca^{2+} transients in neurons before (0 min) and after 1 μg/ml

α-IgLON5 AAB treatment for 60 min or 2 days. N=3 independent experiments with 71 to 994 neurons

per condition and time point. One-way ANOVA with Tukey post-test. **g,** Percentage (%) of inactive

(blue), active (green), and hyperactive (pink) neurons in cultures treated with 1 μg/ml antibody or

controls for 0 or 60 min or 2 days. **h,** Quantification of Tau re-sorting into the soma in neurons treated

with 1 μg/ml α-IgLON5 after suppressing their activity with 10 μM EGTA-AM pre-treatment for 20 min.

Data collected from two independent experiments with 47 to 60 neurons per condition and presented

as mean±SEM. One-way ANOVA with Tukey post-test **i,** Quantification of Tau re-sorting after treatment

with monoclonal 1 μg/ml α-LGI1 AAB, α-IgLON5 AAAB, and controls (mCtrl and pCtrl) for 2 days. Data

collected from two independent experiments and presented as mean±SEM, One-way ANOVA with

Tukey post-test. **j,** Model: Binding of α-IgLON5 AABs triggers acute neuronal hyperactivity (in minutes),

which induces prolonged Tau phosphorylation and somatodendritic resorting (=Tau activation) for days.

Excessive or repeated α-IgLON5 AABs exposure leads to acute excitotoxicity and/or permanent Tau

activation resulting in neuronal death.

904
905
906

Supplementary Figures

Supplementary Figure S1. α-IgLON5 AABs IgG subclass analysis and expression in adult mouse brain. a, Dot blot of patient-derived α-IgLON5 AABs and a control IgG pool isolated from serum of healthy control individuals (pCtrl) detected with IgG subclass-specific secondary antibodies. **b,** Spatial distribution of IgLON5 protein expression in the adult mouse brain (Allen mouse brain atlas: http://mouse.brain-map.org/experiment/show?id=71380538). Scale bar = 2000 μm. **c,** Additional control antibodies tested did not bind to cultured hippocampal mouse neurons: monoclonal human non-reactive AAB (mCtrl) [21] and commercially available normal human IgG (pCtrl_rec). Scale bars = 20 μm. **d,** Pre- (Synapsin-1) and post-synaptic (PSD95) cluster numbers and sizes did not change upon acute α-IgLON5 AAB exposure (1 μg/ml for 60 min). Data collected from three independent experiments and shown as mean±SEM, Student T-test. **e,** Representative images showing internalization and lysosomal acidification (=fluorescence) of pHrodo™-labelled α-IgLON5 or pCtrl in HEK293 cells overexpressing full-length IgLON5 after 4 h (left) and in primary hippocampal mouse neurons after 1 day and 2 days (right). Scale bars = 10 μm.

a Distribution of infused α-IgLON5 AABs in the brain

anterior → posterior

α-IgLON5 AABs
DAPI

2500 μm

b MFG-E8 immunolabeling in brain sections

Coronal brain section of α-IgLON5 treated animal

Cortex
CA3
Dentate gyrus

MFG-E8
Congo red

α-IgLON5   pCtrl   PBS   Positive Ctrl (APP23)

Cortex
Dentate gyrus
50 μm
CA3
50 μm

Positive Control (APP23)

Hippocampus

MFG-E8
Congo red

c GFAP immunolabeling in anterior and posterior hippocampus

Anterior hippocampus
Posterior hippocampus

Anterior HPC
Posterior HPC

Norm. mean intensity
PBS mCtrl pCtrl α-IgLON5

d Iba1 immunolabeling in hippocampus

Anterior HPC
Posterior HPC

e GFAP immunolabeling in spinal cord

α-IgLON5
mCtrl
1000 μm
GFAP
GFAP

f Iba1 immunolabeling in spinal cord

α-IgLON5
mCtrl
1000 μm
Iba1
Iba1

925
926

Supplemental Figure S2. MFEG-8 immunolabeling in brain sections of AAB treated mice. a,
Images of coronal mouse brain sections (from anterior to posterior) show distribution of infused α-
IgLON5 AABs in the brain after 14 days (fluorescence anti-human IgG secondary antibody). **b,** Location
of anterior and posterior hippocampal regions shown in images on the left (adapted from 3D-brain
Explorer, Allen Mouse Brain Atlas). Quantification of GFAP (astrocytes) and **c,** Iba1 (microglia) intensity
in anterior and posterior hippocampus of α-IgLON5 AAB and control brains. Data shown as mean±SEM,
N=3-6 animals, 2-3 brain sections per animal. One-way ANOVA with Tukey post-test. **d,**
Immunolabelling of MFGE-8 and co-staining with congo red in α-IgLON5 AAB and control brains, in
cortex, dentate gyrus and CA3 (Schaffer collaterals). APP23 mouse brain section was used as a
positive control **e**, Representative images, and quantification of GFAP and **f,** Iba1 immunolabeling in
spinal cord sections of α-IgLON5 AAB and control animals. Data shown as mean±SEM, N=3-4 animals,
5-8 spinal cord sections per animal. One-way ANOVA with Tukey post-test.

Supplementary Figure S3. **Gene and pathway deregulation detected by bulk RNAseq of cortical tissue. a,** Heat map showing differential gene expression in contralateral cortical tissue in individual animals from α-IgLON5, pCtrl, and PBS groups. **b,** Gene ontology (GO) enrichment analysis for cellular components and KEGG pathways in α-IgLON5 AAB infused animals. **c,** Immunolabeling of inflammatory markers (C3, C1q, CD68) in hippocampal CA3 region, upregulated in α-IgLON5 vs. pCtrl AAB treated mice.

Supplementary Figure S4. **Changes observed upon infusion of high dose (200 μg AABs). a,** Body weight relative to Day 0 measured during 14 days of AAB treatment. Data shown as mean±SEM, N=4 animals per group. **d,** Rotarod performance (time spent on rod) of α-IgLON5 AAB and pCtrl animals determined at Day 14. Data shown as mean±SEM, N=4 animals per group. Student T-test. **c,** Serum levels of NfL and GFAP determined by ELISA. Data shown as mean±SEM, N=4 animals, with three technical replicates measured per animal. Student T-test. **d,** Representative coronal brain sections (anterior to posterior) immunolabeled for GFAP (astrocytes) and **e,** Iba1 (microglia) in α-IgLON5 AAB and pCtrl animals. Quantification of GFAP and Iba1 intensity in cerebral cortex, hippocampus, brainstem and cerebellum in α-IgLON5 AAB and control brains. Data shown as mean±SEM, N=4 animals, 3-5 brain sections per animal. Student T-test. **f,** Representative images, and quantification of GFAP and **g,** Iba1 immunolabeling in spinal cord sections. Data shown as mean±SEM, N=4 animals, 5-8 spinal cord sections per animal. Student T-test.

a p-Tau in brain lysates of AAB infused mice

Fimbria

Cerebellum

Spinal cord

b Correlation of hippocampal p-Tau and glia cell markers [individual groups]

Iba1 vs. p-Tau

α-IgLON5 Y = 0.6561*X + 0.3388 R² = 0.1405
pCtrl Y = -0.03854*X + 0.5733 R² = 0.0009167
mCtrl Y = -1.130*X + 1.524 R² = 0.8164
PBS Y = 0.01332*X + 0.2713 R² = 0.002310

GFAP vs. p-Tau

α-IgLON5 Y = -0.3932*X + 1.376 R² = 0.1476
pCtrl Y = -0.5178*X + 1.313 R² = 0.1832
mCtrl Y = -1.464*X + 1.844 R² = 0.7609
PBS Y = 0.2078*X + 0.5378 R² = 0.07903

c p-Tau in spinal cord motor neurons of AAB infused mice

p-Tau (AT8) in spinal cord

d p-Tau in cultured neurons after 2 days of AAB treatment (1 μg/ml)

e p-Tau in cultured neurons (2 days AABs at diff. conc.)

963
964

Supplementary Figure S5. **p-Tau levels in brains and cultured neurons upon AAB treatment. a,** Western Blot membranes probed for p-Tau (p-Tau pS202/pT205 (AT8), pS396/pS404 (PHF-1), and pT231 (AT180) epitopes) and total Tau in lysates from fimbria, and cerebellum, and spinal cord tissues of α-IgLON5 AAB and control animals. N=3-4 animals. Lysates from control and AD human brain were included as controls. **b,** Correlation between normalized fluorescence intensities of hippocampal microglia (Iba1, top) and astrocytes (GFAP, bottom) and p-Tau in projections onto CA3 for all treatment groups, with linear fits for all individual control groups. A positive correlation only exists for Iba1 vs. p-Tau in α-IgLON5 animals. **c,** Representative images of p-Tau (pS202/pT205 (AT8)) immunolabeling in spinal cord sections of α-IgLON5 AAB and control mice, and quantification of mean intensity in the spinal cord ventral horn. Data shown as mean±SEM, N=3-6 animals, 5-8 spinal cord sections per animal. One-way ANOVA with Tukey post-test. **d**, Western Blot membranes probed for p-Tau (p-Tau pS202/pT205 (AT8), pS396/pS404 (PHF-1), pS199 and pT231 (AT180) epitopes) and total Tau in lysates from primary neurons treated with 1 μg/ml α-IgLON5 AABs and control antibodies for 2 days. Two independent experiments with three technical replicates per treatment, data shown as mean±SEM. One-way ANOVA with Tukey post-test. **e,** Western Blot membranes probed for p-Tau (p-Tau pS202/pT205 (AT8), pS396/pS404 (PHF-1), pS199 and pT231 (AT180) epitopes) and total Tau in lysates from cultured neurons treated with 1, 10 and 20 μg/μl α-IgLON5 AABs for 2 days. Two independent experiment per treatment, data presented as mean±SEM. One-way ANOVA with Tukey post-test.

Supplementary Figure S6. α-IgLON5 AAB-induced acute neuronal hyperactivity and long-term silencing. a+b, Representative z-projection (sum) of Ca²⁺ recordings, individual activity traces, and raster plots for detected Ca²⁺ spikes after (**a**) 60 min and (**b**) 2 days exposure in neurons treated with 1 μg/ml α-IgLON5 or untreated (NT). **c,** LDH activity in culture media after 60 min and 2 days treatment with 1 μg/ml α-IgLON5 or control AABs. N=3 independent experiments with 3 technical replicates each, data presented as mean±SEM, One-way ANOVA with Tukey post-test **d,** Representative image of autaptic cultures immunolabeled for MAP2 (blue), PSD95 (cyan), Synapsin-1 (magenta) and DNA

(DAPI, white). **e,** Electrophysiological recordings in autaptic cultures treated with 0.1 μg/ml α-IgLON5

AABs and controls for 3 days. Measured parameters are (i) membrane capacitance (pC) and (i) resting

membrane potential, (iii) excitatory postsynaptic current (EPSC) amplitude, (iv) EPSC paired-pulse ratio

(PPR), (v) readily releasable pool size (sucrose charge), (vi) synaptic vesicle release probability (Pvr),

and (vii) EPSC release during 10 Hz stimulation. N=3 independent autaptic cultures. Data are presented

as mean±SEM, One-way ANOVA with Tukey post-test. **f,** Representative z-projection (sum) of Ca^{2+}

recordings and GCamp6f fluorescent intensity measurement in neurons treated with 100 μM glutamate

for 30 min after 0.1, 10, 100 μM EGTA-AM treatment for 20 min. **g,** Frequency (transients/min) analysis

in neurons pre-treated with 0.1, 10, or 100 μM EGTA-AM for 20 min and then exposed to 1 μg/ml α-

IgLON5 or control AABs for 60 min. Data were collected from two primary neurons cultures and are

shown as mean±SEM. One-way ANOVA with Tukey post-test. **h,** Primary neurons treated with 0.1, 10,

or 100 μM EGTA-AM for 1 day and 2 days and imaged by bright field microscope.

1005

1006

1007

1008
1009

1010

Methods

Ethics

All analyses of patient material were approved by the Charité - Universitätsmedizin Berlin Ethics Board (#EA1/258/18).

Primary mouse neuron preparations and experiments involving adult animals (C57/B6, male) were carried out according to the guidelines stated in the directive 2010/63/EU of the European Parliament on the protection of animals used for scientific purposes and were approved by local authorities in Berlin and the animal welfare committee of the Charité university in Berlin, Germany.

Antibody Detection in patient CSF and serum.

Specificity of the purified α-IgLON5 AABs was confirmed by indirect immunofluorescence (IF) performed on HEK293 cells transfected with commercially available IgLON5 expression vector (EUROIMMUN, # 1151-1005-50) and cells with control transfection. α-IgLON5 AAB binding on cells was confirmed at dilutions of 1:100 for CSF and 1:1000 for serum.

Purification of patient-derived α-IgLON5 AABs

Recombinant human IgLON5-Fc chimera (R&D Systems) was bound to a 1 ml HiTrap® NHS-activated HP column (GE Healthcare) according to manufacturer's guide. For antibody isolation, eluates of plasmapheresis were loaded overnight onto the column. Then, the column was washed with 15 ml 20 mM Tris (pH 7.5) and 15 ml 0.5 M NaCl in 20 mM Tris (pH 7.5). α-IgLON5 AABs were eluted with 100 mM glycine (pH 2.2) with subsequent neutralization using 1 M Tris (pH 8.8). Finally, the samples were re-buffered into phosphate-buffered saline (PBS) using Vivaspin® 15R ultrafiltration spin columns (Sartorius). For detection of specificity, recombinant IgLON5 or LG1 protein was spotted onto a nitrocellulose membrane (GE Healthcare), membranes were blocked with 1% BSA in PBS, incubated with α-IgLON5 AABs over night at 4°C, washed and incubated with HRP-coupled anti-human IgG. For detection of IgG subclasses, 2 μg of patient-derived AABs were dotted on a nitrocellulose membrane (GE Healthcare), membranes were probed with mouse anti-human IgG1-POD, anti-human IgG2-POD, anti-human IgG3-POD, and IgG4-POD ab (ThermoFisher Scientific), and sheep-anti-human IgG (Seramun).

For the isolation of control serum IgGs (pCtrl), 750 μl serum from healthy controls were mixed with 450 μl protein A/G agarose (SantaCruz). IgG fractions were isolated according to manufacturer's instructions. Purity was checked by Coomassie staining. Protein concentration

1045 was determined using BCA protein assay Kit (ThermoFisher Scientific). The non-reactive

1046 human monoclonal control AAB (mCtrl; mGO53) was purchased from InVivoBiotech.

1047

Relative binding strength quantification using flow cytometry

Binding strength was determined following previously published protocols [67]. In brief,

HEK293T cells were transfected with cDNA plasmids coding for full human IgLON5 protein

which co-expressed a myc-tag for transfection efficiency control. Three days post-transfection,

cells were harvested and stained with serial dilutions of AABs and c-myc AABs. From live cells

with top 30% protein expression (evaluated by c-myc signal), the mean fluorescence intensity

(MFI) of Alexa488-coupled goat anti-human IgG (1:500; Life Technologies) was evaluated,

and nonlinear regression models [$MFI = MFI_{max} * IgG$ conc./ ($0.5 * MFI_{max} + IgG$ conc.)] under

settings for one site specific binding were generated using GraphPad Prism 8 (GraphPad

Software Inc.)

1058

Tissue reactivity screening

Sagittal mouse brain sections were obtained from unfixed tissue, cut on a cryostat (Leica) at

20 μm thickness and mounted on glass slides. Sections were rinsed with PBS, then blocked

with PBS containing with 2% bovine serum albumin (BSA) and 5% normal goat serum (NGS)

for 1 h at room temperature (RT), and incubated with α-IgLON5 AABs and pCtrl overnight at

4°C. After three washes with PBS, goat anti-human IgG-AlexaFluor488 was applied for 2 h at

RT, followed by staining with DAPI (1:1000 in PBS) for 5 min. Sections were washed with PBS

before mounting with Immo-Mount (Epredia) and imaging using an inverted epifluorescence

microscope (Leica SPE).

1068

Epitope mapping in HEK293T cells expressing IgLON5 constructs

To map the epitopes of patient-derived α-IgLON5 AABs, we cloned human IgLON5 deletion

constructs from full-length Myc-DKK-tagged human IgLON5 plasmid (Origene, #225495)

using a Q5® Site-Directed Mutagenesis kit (New England Biolabs). Individual Ig domains of

IgLON5 (Ig1, Ig2, and Ig3), or combinations thereof (Ig1+Ig2, Ig1+Ig3, and Ig2+Ig3), were

expressed in HEK293T cells for 48 h. HEK293T cells were seeded on poly-L-lysine (PLL)

coated coverslips and transiently transfected with either full-length IgLON5 or its deletional

constructs. After fixation with 4 % paraformaldehyde (PFA) at RT for 10 min, blocking with 3%

NGS in PBS for 1 h, cells were incubated with α-IgLON5 AABs or pCtrl overnight at 4°C. After

washing and incubation with the AlexaFluor488-conjugated anti-human IgG for 1 h at RT.

Binding of α-IgLON5 AABs was evaluated based on epifluorescence microscope (Leica SPE).

1080

Antibody internalization experiments

α-IgLON5 AABs and pCtrl were conjugated with endosomal-pH sensitive red-fluorescent dye (pHrodo^{TM} iFL Red STP ester, ThermoFisher Scientific) following the manufacturer's instructions. pHrodo^{TM}-conjugated antibodies were applied at 5 μg/ml to HEK293T cells overexpressing human full-length IgLON5 for 4 h at 37°C, or to primary mouse hippocampal neurons (DIV12) for 1 day or 2 days at 37°C. Images were captured on a spinning disc confocal microscope (Nikon, SoRa CSU-W1).

Intrathecal osmotic pump infusion

Eight weeks-old male C57Bl/6J wild-type mice were randomized for the different treatment groups by an independent investigator. α-IgLON5 AABs and non-reactive human monoclonal IgG (mCtrl) [21], IgG from healthy control individual serum (pCtrl), and PBS as controls - were delivered continuously into the right lateral ventricle over the course of 14 days, with a flow rate of 0.25 μl/h. Antibody cerebroventricular infusion was performed unilaterally using osmotic pumps (model 1002, Alzet, Cupertino, CA), which were loaded 24 h prior to surgical implantation. For pump implantation, mice were placed in a stereotaxic frame and a cannula was inserted into the right ventricle (Coordinates: 0.2 mm posterior and ± 1.00 mm lateral from bregma, depth 2.2 mm). The cannula was connected to a pump, which was subcutaneously implanted on the interscapular space of the animals. After surgery, mice were monitored daily to assess symptoms and body weight. Mice were sacrificed after 14 days and brain, spinal cord tissues, and serum were harvested.

Two cohorts were infused with different amounts of antibodies: 1) Low dose cohort receiving 75 μg antibody (~3 μg/g b.w.) consisted of α-IgLON5 (N=6), mCtrl (N=3), pCtrl (N=4), and PBS (N=4) animals. 2) High dose cohort receiving 200 μg antibody (8 μg/g b.w.) consisted of α-IgLON5 (N=4) and pCtrl (N=4) animals.

Tissue processing

For protein and RNA analysis, brain and spinal cord tissues were promptly dissected on ice, flash-frozen by placing on dry ice, and stored at -80°C until further use.

For protein extraction, brain and spinal cord tissues were homogenized in ice-cold radioimmunoprecipitation assay (RIPA) buffer (Sigma-Aldrich) containing phosphatase and protease inhibitor cocktail (ThermoFisher Scientific) using a probe homogenizer. After incubating the homogenates on ice for 20 min, samples were centrifuged at 10000xg for 20 min at 4°C and the protein concentration in supernatant was measured with BCA protein assay Kit (Pierce). Concentration of each sample was adjusted to 2 μg/μl and then mixed with 6 X

SDS-containing Laemmli buffer (ThermoFisher Scientific). Subsequently, they were boiled at
95 °C for 5 min and stored at -20°C until the Western blot analysis.
<u>For Immunohistochemistry (IHC),</u> brain and spinal cord tissues were drop-fixed in 4% PFA for
2 days. For cryoprotection, fixed tissues were sequentially transferred to 10%, 20% and 30%
sucrose in PBS with 0.02% sodium azide at 4°C. Tissue was cut into 30 µm thick sections
using cryostat (Leica) and serially collected in 50% glycerol in PBS to be stored at -20°C until
used for IHC.
1123

Western blot

To analyze total and p-Tau levels by Western blot, 10 µg of total protein per sample was
loaded onto 4 to 12% bis-tris SDS- polyacrylamide gel (Invitrogen). Following electrophoresis,
the separated proteins were transferred to a nitrocellulose membrane. The membrane was
blocked with 3% BSA in PBS containing 0.05% of Tween (PBS-T) at RT for 1 h. Subsequently,
the membranes were incubated with primary antibodies for total Tau, p-Tau and loading
control, diluted in the blocking solution, overnight at 4°C. The following day, membranes were
washed 3-times for 5 min in PBS-T and then incubated with fluorescent-dye conjugated
secondary antibodies, diluted in blocking solution, for 1 h at RT. Membranes were then imaged
using a Licor imaging system (Odyssey DLx). A full list of antibodies used in Western blot can
be found in SupplementaryTable T2.
1135

Histology

For immunodetection, 30 µm-thick PFA-fixed brain and spinal cord sections were washed free-
floating in tris-buffered saline (TBS), permeabilized with 0.3% Triton X-100 in TBS for 20 min
at RT, and washed 2-times for 5 min in TBS. All incubation steps were performed on an orbital
shaker (180 rpm). Sections were blocked in blocking solution (3% NGS in TBS) for 1 h at RT.
Sections were incubated with primary antibodies diluted in blocking solution at 4°C overnight.
Next day, the sections were washed 3-times for 10 min in TBS and incubated with Alexa Fluor
conjugated secondary antibody diluted in blocking solution for 2 h at RT. Sections were
washed 3-times for 10 min in TBS and incubated with DAPI (1:1000 in PBS) for 15 min at RT.
Sections were then mounted onto microscope slides and covered with coverslip using
Fluoroshield™ mounting medium (Sigma-Aldrich). A full list of antibodies used IHC can be
found in SupplementaryTable T2. Sections were imaged on a wide field fluorescence
microscope (Eclipse-Ti, Nikon) using 10x objective with tiling function. Same imaging
parameters were used while imaging the sections across different groups. Mean fluorescence
intensity was measured using ImageJ from manually drawn regions of interest (ROIs) in the
brain (on contralateral site to avoid injection confound) and spinal cord sections. Mean

intensity was averaged over 3-5 sections for brain tissue or 5-8 sections for spinal cord tissue

per animal.

Detection of the spread of infused AABs was done in free-floating brain sections with goat

anti-human HRP-conjugated secondary antibody in 3% NGS in TBS for 2 h at RT. Sections

were washed 3-times for 5 min in TBS and the signal was amplified with a CF® 594 tyramide

signal amplification kit (ThermoFisher Scientific).

1158

1159 ***Evaluation of serum***

1160 Levels of neurofilament light chain (NfL) and glial fibrillary acidic protein (GFAP) were

1161 measured from undiluted serum samples using commercial enzyme-linked immunosorbent

1162 (ELISA) kits according to manufacturer's instructions. (GFAP: Abbexa, NfL:Cloud-Clone).

1163

RNA sequencing

RNA was extracted using High Pure RNA Tissue Kit (Roche) according to manufacturer's

instructions. The quality of the purified total RNA was assessed with high sensitivity RNA

screenTape with RIN number >7.0 and the RNA integrity was confirmed using Agilent

Tapestation. Following ribosomal rRNA depletion, total RNAseq was sequenced in a paired

end configuration using Illumina Novaseq 6000. Alignment and analysis scripts can be found

at https://github.com/qoldt/Iglon5_RNAseq. Computation has been performed on the HPC for

Research cluster of the Berlin Institute of Health. Briefly, raw reads were aligned to GRCm39

with STAR using gencode basic gene annotation (vM31) and the prepared

GRCm39_Ensembl_rmsk_TE.gtf (Hammel laboratory). Count tables were generated using

TEcount [68]. Upon investigation, sample IgG_2 (Animal 649) was found to have excessive

gliosis and Iglon5_3 (Animal 660) had incomplete perfusion and were excluded from analysis.

Differential expression was determined using the quasi-likelihood F test in edgeR by

comparing Iglon5 treated vs PBS treated samples, and taking the top 250 differentially

expressed genes using the TopTags function with corrected P values <0.05. Gene ontology

(GO) enrichment analysis was performed using ShinyGO v0.77 (South Dakota State

University, Brookings, SD, USA) to generate the list of pathways affected using gene ontology

molecular function, cellular components and Kyoto Encyclopedia of Genes and Genomes

(KEGG) databases.

1183

Preparation of mouse primary neurons

Primary neurons were prepared from hippocampi dissected from P0-P1 wildtype mice of either

sex. Hippocampi were dissected in ice-cold HBSS (Millipore) containing 1% P/S. Then,

hippocampal tissue was digested in enzyme solution containing DMEM (ThermoFisher

Scientific), (in mM) 3.3 cysteine, 2 CaCl2, 1 EDTA, and papain (20U/ml, Worthington) at 37°C

for 30 min. Papain reaction was inhibited by incubating digested hippocampal tissue in DMEM
containing 10% FBS (ThermoFisher Scientific), 1% P/S, 38 mM BSA, and 95 mM trypsin
inhibitor at 37°C for 5 min. Cells then were triturated in complete Neurobasal-A (NBA) medium
containing 10% FBS, 2% B-27, 1% Glutamax, 1% penicillin/streptomycin (P/S) and seeded in
PLL (0.1 mg/mL) coated glass bottom μ-slide 8-well imaging dishes (ibidi) at a seeding density
of ~30000 cells/cm^2 for IF experiments. For calcium (Ca^{2+}) imaging, neurons were seeded in
96-well clear bottom black microplates (Corning) at a seeding density of 125000 cells/cm^2 and
maintained at 37°C and 5% CO2 until used for experiments. After 3 h seeding the neurons,
the media was completely changed to FBS and phenol-red free complete NBA medium. Every
second day, one-fifth of the media was replaced by fresh complete NBA medium.
1199

Differentiation of human neurons
Human neuronal stem cells (hNSCs) were differentiated using an inducible Neurog2 hNSCs
through doxycycline induction. hNSCs were received from BIH Core Unit pluripotent Stem
Cells and Organoids (CUSCO), Charité-Berlin. Cells were seeded in laminin 521 (Biolamina)-
coated 8-well plates. The composition of the differentiation medium was: NBA, Neural
Induction Supplement, Advanced DMEM/F12 (Gibco), supplemented by antibiotic (Gibco) and
doxycycline (Sigma-Aldrich). On Day 4, the medium change was performed by the medium
supplemented with AraC (Sigma-Aldrich), to reduce the number of non-neuronal cells. The
differentiation medium was refreshed every day, the cells were cultivated for 43 days in 37°C
and 5% CO_2.
1210

IF in cultured neurons
Live staining (surface antigen binding): Primary mouse hippocampal neurons (DIV12) were
treated with 1 μg/ml α-IgLON5 AABs or control antibodies and kept at 37°C and 5% CO_2 for
60 min. Then, neurons were washed with PBS before fixation with 4% PFA in PBS for 15 min,
and then washed with TBS for 10 min at RT. After fixation, neurons were blocked with 3%
NGS for 1 h at RT and incubated with anti-human 488 secondary antibodies for 2 h at RT.
After 3-times 10 min PBS washes, nuclei were stained with DAPI (1:1000 in PBS) for 10 min.
Dishes were imaged with laser scanning confocal microscope (Nikon, A1Rsi+) with a x60 oil
objective.
Co-localization with synaptic markers: After the surface antigen labelling as described above,
neurons were permeabilized with PBS containing 0.3% Triton X-100 for 20 min at RT. After
two consecutive washing steps with excess PBS, neurons were blocked again with 3% NGS
for 1 h. Neurons then were incubated with primary antibodies, diluted in blocking solution, for
microtubule (MAP2) and pre- (Synapsin1), or post-(PSD95) synaptic markers overnight at

4°C. Secondary and DAPI stainings were performed as described above. Dishes were imaged

with laser scanning confocal microscope (Nikon, A1Rsi+) with a x60 oil objective using the z-

stack function (z-step: 0,5 μm). Same imaging settings were used between each biological

replicate (N=3) to allow the comparison for co-localization analysis which was performed using

ImageJ.

<u>Tau missorting:</u> Primary neurons were treated with different concentrations of α-IgLON5 AABs

or control antibodies for either 2 days or 5 days. For Ca^{2+} depletion experiments, neurons

were treated with 10 μM cell permeable Ca^{2+} chelator, EGTA-AM, (AAT Bioquest) 20 min prior

to the antibody treatment. After the treatments, neurons were fixed at DIV9-12 with 4% PFA

for 15 min and immunolabeled for total Tau and MAP2 by following the IF protocol described

above. Antibody specifications are listed in SupplementaryTable T2. Neurons were imaged

with spinning disc confocal microscope (Nikon, CSU-X), using 40X oil objective. Identical

imaging settings were used between each biological replicate (N=3-4). Tau missorting

analysis was done with ImageJ by calculating the mean intensity in somatodendritic

compartment in Tau channel from manually drawn ROIs, as depicted in Fig. 4d.

<u>Cytotoxicy assay:</u> Before fixing the cells, media was collected and centrifuged at 1400 rpm for

1 min to remove the cell debris. A commercially available lactate dehydrogenase (LDH) kit

was used to measure the release of LDH to the culture media (CyQUANT-ThermoFisher

Scientific) according to the manufacturer's description.

1244

Ca^{2+} imaging in cultured neurons

For Ca^{2+} imaging in cultured hippocampal neurons, AAV2/9 serotype was used for neuron

specific expression of GCamp6f under *Syn* promoter. AAVs (ruby-P2A-GCamp6f) were

ordered from Viral Core Facility of Charite Universitätsmedizin, Berlin (Catalog number: BA-

026e). In cultured hippocampal neurons, 0.16 μl of 1,04 x 10^{13} v.g./ml of viruses were added

to medium on DIV5. Ca^{2+} recordings were performed at DIV13-15.

<u>Ca^{2+} transient recordings:</u> Neurons in 96-well plate expressing GCamp6f were recorded in a

wide field fluorescence microscope (Eclipse-Ti, Nikon) with a 10x objective. Imaging dishes

were placed in the live-cell imaging chamber of the microscope to maintain an environment of

37°C, 5% CO_2 and ~95% humidity. α-IgLON5 AABs or control antibodies were added to the

wells by direct application. Ca^{2+} imaging was done in the green channel and mRuby

expressing neurons were visualized in the red channel. Ca^{2+} transients were recorded for 180

sec at ~8 Hz. Two to three wells in three independent cultures were analyzed per condition.

<u>Analysis:</u> ROIs were created for each soma in red channel using a custom written script in

Cell Profiler. Created ROIs were placed over each frame of the video recorded in green

channel and mean intensity over time was measured using ImageJ. The .csv files containing

mean fluorescent intensity from each image of the time-lapse videos were processed using

the FluoroSNNAP application for ΔF/F0 conversion and spike detection [69]. In short, .csv files

were uploaded to FluoroSNNAP and the function "Convert raw fluorescence data to ΔF/F0"

was used to compute baseline fluorescent by taking the average of 50th percentile of the

signal across a 60 s time window. Ca^{2+} transients from individual ROIs were detected with a

template-based approach. This method identifies events by comparing a moving window of

the Ca^{2+} signal to a predefined library of Ca^{2+} waveform templates. Events were detected with

a similarity threshold of 0.75 and a minimum amplitude of 0.01. Hyperactivity criteria was

calculated for each time point independently by taking the mean±SD of pCtrl group as a

reference value for normal activity: Spike frequencies larger than [mean+2*SD] of pCtrl were

considered as hyperactivity.

1272

1273 *Electrophysiology in primary autaptic cultures*

Isolated primary neurons on micro islands of glial cells ("autaptic cultures") were prepared

from wildtype mice as recently described [70], with slight modifications. Briefly, 300 µm spots of

a growth-permissive substrate mix of 0.7 mg/ml collagen and 0.1 mg/ml poly-D-lysine were

printed on glass coverslips coated with agarose. Astrocytes were seeded onto these

coverslips in DMEM (ThermoFisher Scientific), supplemented with 10% fetal calf serum and

0.2% P/S (Invitrogen). After formation of glia micro-islands, DMEM was replaced with NBA

supplemented with 2% B27 and 0.2% P/S. Hippocampal neurons prepared from P0 mice were

added at a density of 370 cells/cm^2. Prior to electrophysiological recordings, autaptic cultures

were treated with antibodies at a final concentration of 0.1 µg/ml for 3 days.

Electrophysiological recordings: Neurons were recorded at DIV13-17 at RT on an IX73

inverted microscope (Olympus) using a Multiclamp 700B amplifier under the control of a

Digidata 1550 AD board and Clampex 10 software (all Molecular Devices). Data were acquired

at 10 kHz and filtered at 3 kHz, and series resistance was compensated at 70%. The

extracellular solution contained (in mM) 140 NaCl, 2.4 KCl, 10 HEPES, 10 glucose, 2 $CaCl_2$,

and 4 $MgCl_2$ (pH adjusted to 7.3 with NaOH, 300 mOsm). The intracellular solution contained

(in mM) 136 KCl, 17.8 HEPES, 1 EGTA, 4.6 $MgCl_2$, 4 Na_2ATP, 0.3 NaGTP, 12 disodium

phosphocreatine, 50 U/ml creatine phosphokinase, pH adjusted to 7.3 with KOH, 300 mOsm.

Autaptic neurons were recorded in whole-cell voltage clamp mode using thick-walled

borosilicate pipettes with a tip resistance of 3-4 MΩ. Membrane potential was set to −70 mV.

Paired EPSCs were evoked every 5 s by triggering two unclamped action potentials with 40

ms interstimulus interval using 1 ms depolarizations of the soma to 0 mV. The readily-

releasable pool of synaptic vesicles was determined by application of a hypertonic 500 mM

sucrose solution for 10 sec. Electrophysiological recordings were analyzed using AxoGraph.

The vesicular release probability was calculated as the ration of the charge of the sucrose
evoked response by the average charge of six EPSCs prior to the sucrose application. The
paired-pulse ratio was calculated as the ratio from the second and first EPSC amplitude.
1300

1301 ***Statistical analysis***
1302 All data plotting and statistical analyses were performed in GraphPad Prism 9. Comparison of
1303 two groups was done by Student T-test and multiple group comparisons were performed by
1304 one-way ANOVA with Tukey or Holmes-Sidak, or as indicated in the figure legends.
1305 ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$.
1306

1307 ***Data and code availability***
1308 Lists of differentially expressed gene can be found in Supplementary Table T1. Scripts for
1309 RNAseq analysis can be found at https://github.com/qoldt/lglon5_RNAseq and the used
1310 code is provided at
1311 file:///Users/wegmanns/Downloads/IgLON5_code%20for%20RNAseq%20analysis.html.
1312