High-throughput Screening of Compounds against Autoluminescent Nonreplicating *Mycobacterium tuberculosis* under Diverse Conditions

Xirong Tian [a,b,c,d*], Wanli Ma [a,b,c,d*], Buhari Yusuf [a,b,c,d], Chunyu Li [a,b,c,d], H.M. Adnan Hameed [a,b,c,d], Xinyue Wang [e,f], Nanshan Zhong [e], Jinxing Hu [f#], Tianyu Zhang [a,b,c,d,f#]

[a] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese Academy of Sciences (CAS), Guangzhou 510530, China

[b] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious Diseases, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese Academy of Sciences (CAS), Guangzhou 510530, China

[c] University of Chinese Academy of Sciences (UCAS), Beijing 100049, China

[d] China-New Zealand Joint Laboratory on Biomedicine and Health, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[e] Guangzhou National Laboratory, Guangzhou 510005, China

[f] Guangzhou Chest Hospital, Guangzhou 510005, China

[*] These authors contributed equally to this work.

[#] Corresponding author:

Tianyu Zhang, Tel: +86-20-32015270, E-mail: zhang_tianyu@gibh.ac.cn

Jinxing Hu, Tel: +86-20-83594895, E-mail: hujinxing2000@163.com

KEYWORDS

Autoluminescence; Low-oxygen; Persistence; *Mycobacterium tuberculosis*; High-

throughput screening

ABSTRACT

The screening of new anti-mycobacterial chemicals is primarily focused on inhibiting

the active growing bacteria. However, a major challenge in tuberculosis control is the

ability of *Mycobacterium tuberculosis* to enter a nonreplicating state for extended

periods, rendering it resistant to many clinical drugs and complicating eradication

efforts. Existing low-oxygen-recovery assays designed for screening compounds

targeting nonreplicating *M. tuberculosis* have limitations, including the colony-

forming unit counting for non-luminous *M. tuberculosis*, and the instability and

exogenous substrate requirements of luminescent *M. tuberculosis* strains carrying free

plasmid with *luxAB* genes. Moreover, these assays fail to accurately replicate the

growth conditions of nonreplicating *M. tuberculosis in vitro*, thus resulting in less

convincing results. To address these challenges, we have developed an

autoluminescence-based, cholesterol-enriched culture evaluation model to assess 17

anti-tuberculosis drugs of different classes against nonreplicating *M. tuberculosis*. Our

findings indicate that the relative light units count, measured in real-time, serves as a

reliable surrogate marker for colony-forming unit count, which typically becomes

available one month later. This suggests the utility of our model for the rapid

determination of drug susceptibility in a dynamic manner. The autoluminescent *M.*

tuberculosis, harbouring *luxCDABE* gene cluster within its genome, can emit blue-

green light stably and autonomously without requiring an external substrate

supplement. The minimal inhibitory concentrations of all the drugs tested under

anaerobic conditions are significantly different from that detected in aerobic

environment. Our model allows for rapid, precise, and efficient assessment of drug

activity under anaerobic conditions, thereby enabling a more comprehensive

evaluation of anti-mycobacterial efficacy. Overall, our model represents a significant

advancement in anti-tuberculosis drug discovery and pharmaceutical development.

INTRODUCTION

The highly infectious pathogen *Mycobacterium tuberculosis* is the causative agent of tuberculosis (TB), and was responsible for 10.6 million TB individuals globally in 2022 (1). The emergence of drug-resistant *M. tuberculosis* strains, coupled with lethal coinfections with human immunodeficiency virus, has significantly exacerbated the global healthcare crisis, necessitating the exploration of novel drugs or treatment regimens against *M. tuberculosis* (2, 3). Notably, after four decades of persistent efforts, bedaquiline (BDQ), the first new anti-TB medication, received approval from the United States Food and Drug Administration in 2012 (4-6).

The persistence of nonreplicating *M. tuberculosis* within lung lesions, capable of enduring for decades and adapting to hypoxic conditions, poses a formidable challenge to TB eradication (7-9). A comprehensive understanding of the complex host environment in which *M. tuberculosis* resides is crucial for developing effective control strategies (10). Previous studies have revealed *M. tuberculosis*'s ability to exploit host-derived cholesterol as a primary carbon source during infection (11). By incorporating cholesterol into growth media, it is more accurately replicate the growth environment of nonreplicating *M. tuberculosis* compared to traditional glycerol-enriched media.

Despite the existence of well-established *in vitro* low-oxygen-recovery assays developed to evaluate drug efficacy against nonreplicating *M. tuberculosis*, these methods have certain limitations (12-14). Some methods rely on enumerating colony forming units (CFUs) or visible bacterial growth, requiring several months for the completion of testing. Others have employed a luminescent *M. tuberculosis* strain with a free plasmid carrying *luxAB* genes to screen for chemical agents against nonreplicating mycobacteria (6). However, this approach is not favored due to its

dependence on additional substrates (6). Our research introduces an

autoluminescence-based, low-oxygen, and cholesterol-enriched evaluation model that

utilizes autoluminescent *M. tuberculosis* H37Ra (AlRa), emitting stable and efficient

blue-green light dynamically during a long period without requiring substrate

supplementation. Bacterial growth is quantified by measuring relative light units

(RLUs) at various time points in a dynamic form at the recovery process. In

conclusion, our model is designed to be straightforward, time-efficient, labor- and

resource-effective, while also facilitating high-throughput screening of drugs and drug

combinations against nonreplicating *M. tuberculosis*.

RESULTS

MICs determination using glycerol- and cholesterol-enriched media under

aerobic conditions. Seventeen anti-TB drugs from distinct classes were selected for

this study. Cholesterol serves as a primary carbon source during *M. tuberculosis*

infection, mimicking the nutritional conditions encountered during the course of *M.*

tuberculosis infection. Moreover, under aerobic conditions, the minimal inhibitory

concentrations (MICs) of the drugs against *M. tuberculosis* remained largely

consistent, irrespective of whether glycerol or cholesterol was used as the carbon

source. An exception to this was observed with para-aminosalicylic acid (PAS), as

detailed in Table 1 and Figure 1. In the glycerol-enriched medium, the MIC of PAS

ranged from 0.25 to 1 μg/mL, however, it increased dramatically to over 64 μg/mL in

the cholesterol-enriched medium (Table 1). The growth curves of actively growing

AlRa treated with nine drugs from different categories are shown in Figure 1, and

those treated with the remaining eight drugs are shown in Figure S1.

MICs determination using glycerol- and cholesterol-enriched media under

anaerobic conditions. We systematically evaluated 17 anti-TB drugs targeting

various crucial aspects of *M. tuberculosis* metabolism, including the cell wall, RNA

polymerase, ribosome and respiration. Our findings revealed that the MICs of these

drugs against *M. tuberculosis* were generally higher under anaerobic conditions

compared to aerobic conditions with the exception of rifampin (RIF) and rifabutin

(RFB), as depicted in Table 1, Figures 2 and Figure S2. Additionally, the MICs of the

drugs remained relatively consistent under anaerobic conditions, regardless of

whether the medium was supplemented with glycerol or cholesterol, with the

exception of moxifloxacin (MOX). The MICs of MOX against nonreplicating *M.*

tuberculosis were found to be in the range of 0.125-2 μg/mL in the presence of

glycerol, whereas in the presence of cholesterol, the MICs spanned from 8-32 μg/mL

(Table 1). Both MOX and levofloxacin (LEV), the two fluoroquinolones, exhibited

increased MICs under anaerobic conditions, regardless of the presence of cholesterol

or glycerol in the media. Remarkably, MOX displayed a greater efficacy compared to

LEV under both anaerobic and aerobic conditions.

Under anaerobic conditions, drugs targeting the cell wall, such as vancomycin (VAN),

isoniazid (INH), ethambutol (EMB), and ethionamide (ETH), did not display

significant antibacterial activity against nonreplicating *M. tuberculosis* as the MICs

were over 64 μg/mL (Table 1). In contrast, these drugs exhibited notable efficacy

against actively growthing *M. tuberculosis* under aerobic conditions.

Compared to other anti-TB drugs, drugs targeting RNA polymerase showed lower

MICs with notably effective anti-*M. tuberculosis* activity in both conditions,

particularly under anaerobic conditions. The MICs of RIF ranged from 0.0625 to

0.125 μg/mL and 0.125 to 2 μg/mL under aerobic and anaerobic conditions

respectively (Table 1). RFB displayed enhanced efficacy compared to RIF against

nonreplicating *M. tuberculosis*, with the MICs as low as 0.03125 to 0.125 μg/mL and

0.03125 to 0.0625 μg/mL in glycerol- and cholesterol-enriched media, respectively

(Table 1). These findings emphasized the promising efficacy of RIF and RFB as

candidates for eradicating nonreplicating *M. tuberculosis*, thus establishing a robust

foundation for their clinical application.

The ribosome-targeting drugs, amikacin (AMK) and streptomycin (STR),

demonstrated anti-TB activity against both actively growing and nonreplicating *M.*

tuberculosis. However, the MICs against nonreplicating *M. tuberculosis* were

significantly higher in comparison to those observed against actively growing *M.*

tuberculosis. Pyrazinoic acid (POA), the active form of pyrazinamide, showed no

anti-*M. tuberculosis* activity under anaerobic conditions, with the MIC 800 μg/mL.

Under anaerobic conditions, drugs targeting folic acid synthesis exhibited no evident

anti-*M. tuberculosis* activity. The MIC of PAS ranged from 0.25-1 μg/mL in glycerol-

enriched 7H9 medium under aerobic conditions, which significantly increased to over

64 μg/mL in cholesterol-enriched 7H9 medium under both aerobic and anaerobic

conditions (Table 1).

Pretomanid (PA-824) demonstrated pronounced anti-*M. tuberculosis* activity,

particularly under anaerobic conditions and with cholesterol as the primary carbon

source, with a MIC range of 1-4 μg/mL (Table 1). TB47 exhibited remarkable anti-*M.*

tuberculosis activity at an exceptionally low concentration of 0.004 μg/mL under

aerobic conditions (Table 1). However, its efficacy against *M. tuberculosis* became

evident only when the concentration was increased to 0.0625-0.25 μg/mL under

anaerobic conditions (Table 1). The MICs of BDQ were 0.5-1 μg/mL and 1-2 μg/mL

against actively growing *M. tuberculosis* when cultivated with glycerol or cholesterol,

respectively (Table 1). Nevertheless, under anaerobic conditions, these MICs

increased to 4-8 μg/mL and 4-16 μg/mL with glycerol or cholesterol, respectively

(Table 1). Clofazimine (CLO) exhibited MICs against nonreplicating *M. tuberculosis*

ranging from 2-16 μg/mL, significantly higher than those observed against actively

growing *M. tuberculosis*, which had an MIC range of 0.5-2 μg/mL (Table 1).

The combined effect of TB47 and CLO against nonreplicating AlRa. Our results

indicated that the combination of TB47 and CLO exhibited partial synergy in terms of

sterilizing activity against nonreplicating *M. tuberculosis* under anaerobic conditions,

both with glycerol and cholesterol-enriched media with a Fractional Inhibitory

Concentration Index (FICI) of 0.75 (Table 2). Notably, the MIC of CLO decreased

from 8 to 4 μg/mL in the presence of TB47 at one-fourth of its MIC. These results

highlighted the potential of this drug combination for clinical evaluation against

nonreplicating *M. tuberculosis*, which has been proved to have strong sterilizing

activities against *M. tuberculosis* in a well-established mouse model in two studies

carried out by us (15, 16), and the manuscript of another study is under revision in

Antimicrob Agents Chemother. Moreover, the assessment model established in this

study for evaluating the efficacy of drug combinations against nonreplicating *M.*

tuberculosis showed significant promise for informing future research and testing

endeavors in this field.

DISCUSSION

During TB infection, *M. tuberculosis* transitions into a nonreplicating state, becoming

resistant to many clinical antibiotics and significantly prolonging therapy (17). *M.*

tuberculosis can acquire and metabolize essential host-derived lipids, particularly

cholesterol, which facilitates disease initiation and progression (18). This necessitates

the establishment of a representative model that faithfully replicates the growth

conditions of nonreplicating *M. tuberculosis* within the host environment. This model

is characterized by oxygen deprivation and cholesterol enrichment of growth medium,

with the goal of screening for effective drugs against nonreplicating *M. tuberculosis*.

To fulfill this objective, we employed AlRa, a strain preserved in our laboratory, to

establish an autoluminescence-based drug efficacy evaluation model for high-

throughput screening of compounds against nonreplicating *M. tuberculosis*. The RLUs

emitted by AlRa, producing blue-green light, exhibit a strong correlation with CFUs,

thus serving as a surrogate marker rather than CFUs in quantifying *M. tuberculosis*

numbers in culture (19).

During the establishment of our model, a notable increase in the growth rate of *M.*

tuberculosis was observed in cholesterol-enriched medium compared to glycerol-

enriched medium, both under aerobic continuously and recovered after cultivation

under anaerobic conditions. This observation supports previous studies suggesting

that cholesterol serves as the primary carbon source during *M. tuberculosis* infection

(11). Our findings suggested that RIF, RFB, PA-824, and TB47 demonstrated

relatively significant potential for inhibiting nonreplicating *M. tuberculosis* under

anaerobic conditions, as previously documented (20-22). Conversely, drugs targeting

the mycobacterial cell wall, such as INH, EMB, ETH, and VAN, showed negligible

anti-mycobacterial activity under anaerobic conditions, consistent with prior

published studies (12, 23). The resistance of nonreplicating *M. tuberculosis* to these

drugs was attributed to the slowed process of cellular division under anaerobic

condition. Drugs that target folic acid synthesis, such as sulfamethoxazole (SMX) and

PAS, manifested minimal bactericidal activities against nonreplicating *M. tuberculosis*

(Table 1). PAS also showed limited activity against actively growing *M. tuberculosis*

with a MIC exceeding 64 µg/mL, when applying the media supplemented with

cholesterol under aerobic conditions. As a prodrug activated through the folate

synthesis pathway, PAS can be converted into folate intermediate analogs, thereby

disrupting *M. tuberculosis*'s folate metabolism (24, 25). It has been hypothesized that

the addition of cholesterol, which prevents PAS from entering *M. tuberculosis*, or the

unique mechanism of action of PAS, contributes to its ineffectiveness against *M.*

tuberculosis under cholesterol-enriched media. Ribosome-targeting drugs, AMK and

STR, exhibited superior activity against nonreplicating *M. tuberculosis* compared to

cell wall-targeting drugs (Table 1). The results are consistent with the observations of

some Chinese clinicians, who have noted good treatment efficacy in multi-drug

resistant tuberculosis patients treated with AMK. This suggests that AMK possesses

relatively potent activity in controlling *M. tuberculosis* persisters, although injection is

one of its disadvantages. Generally, MICs of some drugs against nonreplicating *M.*

tuberculosis were generally higher than those against actively growing *M.*

tuberculosis, indicating that some drugs clinically effective against actively growing

M. tuberculosis may not effectively treat nonreplicating *M. tuberculosis* in lung

lesions, which is demonstrated clearly in animal work (16, 26). It is noteworthy that

fluoroquinolones (either LEV or MOX) have not shown significant effects on

persisting *M. tuberculosis* in previous animal studies, which is consistent with the

predictions made in this study (16, 26). The combination of TB47 and CLO

demonstrated promising partial synergistic activity with a FICI as low as 0.75 under

anaerobic conditions (Table 2), which supports the finding observed in the animal

studies and is promising for clinical trial (15, 16).

The model developed in this study for assessing drug activity against nonreplicating

M. tuberculosis possesses several notable advantages. Firstly, this method stimulates

the growth conditions of nonreplicating *M. tuberculosis* within the human host by

utilizing media supplemented with cholesterol, thereby ensuring maximal fidelity to

the natural environment of the pathogen. Secondly, the model offers simple

manipulation as AlRa consistently emits blue-green light without requiring additional

substrate aldehyde, thereby guaranteeing stable and consistent results. Thirdly, the

results are easily visualized through the consistent detection and comparison of RLUs

emitted by AlRa at various time intervals. Further exploration should prioritize high-

throughput screening of prospective compounds and treatment regimens against

nonreplicating *M. tuberculosis* using the current model. In summary, this model holds

significant application value and possesses high potential to accelerate the discovery

and development of new drugs and treatment strategies targeting nonreplicating *M.*

tuberculosis, which may closely mirror the real state of *M. tuberculosis* persisters *in*

vivo.

MATERIALS AND METHODS

Mycobacterial strains and culture conditions. The mycobacterial strain utilized in

this study is AlRa, an autoluminescent *M. tuberculosis* H37Ra, which has been

genetically engineered to incorporate the *luxCDABE* gene cluster into its genome (19).

AlRa persistently continuously emits blue-green light catalyzed by *luxAB*, facilitated

by *luxCDE*, which consistently supply the necessary substrate aldehyde in a recycling

form, thereby obviating the need for external substrate supplementation (19). Notably,

the drug susceptibility and growth rate of AlRa do not significantly differ from those

of the wild type *M. tuberculosis* H37Ra (19). AlRa was cultured at 37□ in

Middlebrook 7H9 broth (Difco, Detroit, MI, USA) supplemented with 10% oleic

acid-albumin-dextrose-catalase enrichment (BBL, Sparks, MD, USA), 0.2% glycerol,

and 0.05% Tween 80.

Antimicrobials. All pharmaceutical drugs, including ETH, INH, VAN, RIF, RFB,

AMK, STR, MOX, LEV, PA-824, SMX, PAS, POA, BDQ, and CLO, were purchased

from Meilun (Dalian, China) with a minimum purity threshold of ≥ 95%. Additionally,

TB47 was synthesized and sent to us as a gift by Boji (Guangzhou, China). AMK,

INH, and STR were solubilized in sterilized water, while the remaining drugs were

dissolved in dimethyl sulfoxide (DMSO) obtained from Xilong (Shanghai, China). All

drug solutions were freshly prepared and stored at 4□ until use.

Chemotherapy under aerobic condition. The AlRa were cultured in 7H9 medium

supplemented with Tween80 at 37□ until the optical density, measured at a

wavelength of 600 nm, reached to 0.6-0.8 and the RLUs/mL reached 5 million. The

various drug solutions (4 μL each) were combined with AlRa culture (196 μL, 10-fold

diluted using 7H9 without Tween80 and supplemented with either glycerol or

cholesterol) in a sterilized tube and mixed thoroughly. Negative controls were

prepared using DMSO and sterilized water. Each drug concentration and negative

control were assayed simultaneously in three 1.5 mL EP tubes. The AlRa-drug

mixtures underwent a five-day incubation period, during which the RLUs were

consistently assessed at identical intervals each day. The MIC_{lux} (MIC based on the

detection of autoluminescence) of drugs against the actively growing AlRa is defined

as the lowest concentration at which the RLUs in the experimental group are equal to

or less than 10% of the RLUs in the control group after 5 days.

Culture of nonreplicating AlRa. The AlRa strains were cultured until they reached

an optical density of 0.6-0.8, measured at a wavelength of 600 nm, and a state of

5×10^6 RLUs/mL. Subsequently, the AlRa culture was then transferred to a low-

oxygen cabinet (HUAPO, Guangzhou, China) that had already achieved a steady state.

This cabinet was programmed to maintain specific conditions, including a temperature

of 37□, an oxygen content of 0.8%, and a carbon dioxide concentration of 5%. The

methylene blue solution was then added to the AlRa culture to achieve a final

concentration of 6 μg/mL (27, 28). The cultures were incubated until the blue color

disappeared, typically observed within 7-10 days after the addition of methylene blue.

Drug susceptibility tested under anaerobic condition. Under hypoxic conditions,

various drug solutions (4 µL each) were combined with nonreplicating AlRa culture

(196 µL, 10-fold diluted using 7H9 without Tween 80 and supplemented with glycerol

or cholesterol) in a sterile tube and mixed thoroughly. The AlRa-drug mixture was

then incubated for seven days. To remove any potential carryover effects of residual

drugs during the subsequent recovery process, activated carbon (in a volume ratio of

1:5, 50 µL into 200 µL) was added (15). The tubes were subsequently transferred to a

standard aerobic incubator. The RLUs in the co-incubated 1.5 mL EP tubes were

measured and recorded at 7-hour intervals over a period of 28 hours. The

experimental procedure was repeated three times.

Combined activity assay of TB47 and CLO against nonreplicating AlRa. Previous

research from our group has demonstrated that the combined administration of TB47

and CLO significantly exhibits anti-replicating *M. tuberculosis* activity of both *in*

vitro and *in vivo* (16). In this study, we aim to determine whether this combination

would also display notable efficacy against nonreplicating *M. tuberculosis* using the

checkerboard method. Specifically, 2 µL of TB47, 2 µL of CLO, and 196 µL of

diluted AlRa culture in a nonreplicating state were simultaneously added to the same

1.5 mL tube. To eliminate any potential carryover effects of the drugs following seven

days of coincubation, activated carbon (at a volume ratio of 1:5, 50 µL into 200 µL)

was utilized (26). Subsequently, the tubes containing the co-culture were transferred

to aerobic conditions, and RLUs were then detected at 7-hour intervals for the

following 28 hours. The FICI was calculated as the ratio of the MIC of TB47 in

combination to the MIC of TB47 alone, plus the MIC of CLO in combination to the

MIC of CLO alone (29). The effects were categorized based on the FICI into five

groups: synergistic (FICI ≤ 0.5), partially synergistic (0.5 < FICI < 1.0), additive (FICI = 1.0), irrelevant (1.0 < FICI ≤ 4.0), and antagonistic (FICI > 4.0) (29).

Statistical analysis. Before performing statistical analysis, the RLU values were transformed into their logarithmic counts. The time-kill curves were graphically depicted using GraphPad Prism version 8.3.0.

ACKNOWLEDGMENTS

This work received support from the National Key R&D Program of China (2021YFA1300904), the National Natural Science Foundation of China (21920102003) and by the Guangzhou Science and Technology Planning Project (2023A03J0992). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

AUTHOR CONTRIBUTIONS

Conceived the project: Tianyu Zhang, Xirong Tian, Wanli Ma, Nanshan Zhong; Designed the research: Tianyu Zhang, Xirong Tian, Wanli Ma, Xinyue Wang, Jinxing Hu, Nanshan Zhong; Performed the studies: Xirong Tian, Wanli Ma, Chunyu Li, Xinyue Wang; Interpreted the results: Tianyu Zhang, Xirong Tian, Wanli Ma, Jinxing Hu; Drafted the manuscript: Tianyu Zhang, Xirong Tian, Wanli Ma, Buhari Yusuf, H.M. Adnan Hameed; Final approval of the manuscript: all authors.

REFERENCES

1. World Health Organization (WHO). 2023. Global Tuberculosis Report. Available at: https://iris.who.int/bitstream/handle/10665/373828/97892400838 51-eng.pdf?sequence=1

2. Sala C, Dhar N, Hartkoorn RC, Zhang M, Ha YH, Schneider P, Cole ST. 2010. Simple model for testing drugs against nonreplicating *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 54:4150-4158.

305   3.   Koch A, Mizrahi V. 2018. *Mycobacterium tuberculosis. Trends Microbiol*
306   26:555-556.

307   4.   Volynets GP, Tukalo MA, Bdzhola VG, Derkach NM, Gumeniuk MI,
308   Tarnavskiy SS, Starosyla SA, Yarmoluk SM. 2019. Benzaldehyde
309   thiosemicarbazone derivatives against replicating and nonreplicating
310   *Mycobacterium tuberculosis. J Antibiot (Tokyo)* 72:218-224.

311   5.   Kadura S, King N, Nakhoul M, Zhu H, Theron G, Köser CU, Farhat M. 2020.
312   Systematic review of mutations associated with resistance to the new and
313   repurposed *Mycobacterium tuberculosis* drugs bedaquiline, clofazimine,
314   linezolid, delamanid and pretomanid. *J Antimicrob Chemother* 75:2031-2043.

315   6.   He W, Liu C, Liu D, Ma A, Song Y, He P, Bao J, Li Y, Zhao B, Fan J, Cheng Q,
316   Zhao Y. 2021. Prevalence of *Mycobacterium tuberculosis* resistant to
317   bedaquiline and delamanid in China. *J Glob Antimicrob Resist* 26:241-248.

318   7.   Srivastava S, Cirrincione KN, Deshpande D, Gumbo T. 2020. Tedizolid,
319   faropenem, and moxifloxacin combination with potential activity against
320   nonreplicating *Mycobacterium tuberculosis. Front Pharmacol* 11:616294.

321   8.   Wayne LG, Sohaskey CD. 2001. Nonreplicating persistence of *mycobacterium*
322   *tuberculosis. Annu Rev Microbiol* 55:139-163.

323   9.   Iacobino A, Piccaro G, Giannoni F, Mustazzolu A, Fattorini L. 2017. Fighting
324   tuberculosis by drugs targeting nonreplicating *Mycobacterium tuberculosis*
325   bacilli. *Int J Mycobacteriol* 6:213-221.

326   10.   Wipperman MF, Sampson NS, Thomas ST. 2014. Pathogen roid rage:
327   cholesterol utilization by *Mycobacterium tuberculosis. Crit Rev Biochem Mol*
328   *Biol* 49:269-293.

11.   Moopanar K, Nyide ANG, Senzani S, Mvubu NE. 2023. Clinical strains of

Mycobacterium tuberculosis exhibit differential lipid metabolism-associated

transcriptome changes in *in vitro* cholesterol and infection models. *Pathog Dis*

81:ftac046..

12. Cho SH, Warit S, Wan B, Hwang CH, Pauli GF, Franzblau SG. 2007. Low-

oxygen-recovery assay for high-throughput screening of compounds against

nonreplicating *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother*

51:1380-1385.

13. Cho S, Lee HS, Franzblau S. 2015. Microplate Alamar blue assay (MABA)

and low oxygen recovery assay (LORA) for *Mycobacterium tuberculosis*.

Methods Mol Biol 1285:281-292.

14. Macabeo APG, Vidar WS, Chen X, Decker M, Heilmann J, Wan B, Franzblau

SG, Galvez EV, Aguinaldo MAM, Cordell GA. 2011. *Mycobacterium*

tuberculosis and cholinesterase inhibitors from Voacanga globosa. *Eur J Med*

Chem 46:3118-3123.

15. Yu W, Yusuf B, Wang S, Tian X, Hameed HMA, Lu Z, Chiwala G, Alam MS,

Cook GM, Maslov DA, Zhong N, Zhang T. 2021. Sterilizing effects of novel

regimens containing TB47, clofazimine, and linezolid in a murine model of

tuberculosis. *Antimicrob Agents Chemother* 65:e0070621.

16. Yu W, Chiwala G, Gao Y, Liu Z, Sapkota S, Lu Z, Guo L, Khan SA, Zhong N,

Zhang T. 2020. TB47 and clofazimine form a highly synergistic sterilizing

block in a second-line regimen for tuberculosis in mice. *Biomed Pharmacother*

131:110782.

17. Grant SS, Kawate T, Nag PP, Silvis MR, Gordon K, Stanley SA, Kazyanskaya

E, Nietupski R, Golas A, Fitzgerald M, Cho S, Franzblau SG, Hung DT. 2013.

Identification of novel inhibitors of nonreplicating *Mycobacterium*

tuberculosis using a carbon starvation model. *ACS Chem Biol* 8:2224-2234.

18.   Wilburn KM, Fieweger RA, VanderVen BC. 2018. Cholesterol and fatty acids

grease the wheels of *Mycobacterium tuberculosis* pathogenesis. *Pathog Dis*

76:fty021.

19.   Yang F, Njire MM, Liu J, Wu T, Wang B, Liu T, Cao Y, Liu Z, Wan J, Tu Z,

Tan Y, Tan S, Zhang T. 2015. Engineering more stable, selectable marker-free

autoluminescent mycobacteria by one step. *PLoS One* 10:e0119341.

20.   Hu Y, Coates ARM, Mitchison DA. 2003. Sterilizing activities of

fluoroquinolones against rifampin-tolerant populations of *Mycobacterium*

tuberculosis. *Antimicrob Agents Chemother* 47:653-657.

21.   Stover CK, Warrener P, VanDevanter DR, Sherman DR, Arain TM, Langhorne

MH, Anderson SW, Towell JA, Yuan Y, McMurray DN, Kreiswirth BN, Barry

CE, Baker WR. 2000. A small-molecule nitroimidazopyran drug candidate for

the treatment of tuberculosis. *Nature* 405:962-966.

22.   Wayne LG, Sramek HA. 1994. Metronidazole is bactericidal to dormant cells

of *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 38:2054-2058.

23.   Commandeur S, Iakobachvili N, Sparrius M, Nur MM, Mukamolova GV,

Bitter W. 2020. Zebrafish embryo model for assessment of drug efficacy on

mycobacterial persisters. *Antimicrob Agents Chemother* 64:e00801-20.

24.   Zhang Y, Wang S, Chen X, Cui P, Chen J, Zhang W. 2024. Mutations in the

promoter region of methionine transporter gene metM (Rv3253c) confer para-

aminosalicylic acid (PAS) resistance in *Mycobacterium tuberculosis*. *mBio*

15:e0207323.

25.   Wei W, Yan H, Zhao J, Li H, Li Z, Guo H, Wang X, Zhou Y, Zhang X, Zeng J,

Chen T, Zhou L. 2019. Multi-omics comparisons of p-aminosalicylic acid

(PAS) resistance in folC mutated and un-mutated *Mycobacterium tuberculosis*

strains. *Emerg Microbes Infect* 8:248-261.

26.   Grosset JH, Tyagi S, Almeida DV, Converse PJ, Li S-Y, Ammerman NC,

383   Bishai WR, Enarson D, Trébucq A. 2013. Assessment of clofazimine activity

384   in a second-line regimen for tuberculosis in mice. *Am J Respir Crit Care Med*

385   18:608-612.

386   27.   Sumitani M, Takagi S, Tanamura Y, Inoue H. 2004. Oxygen indicator

387   composed of an organic/inorganic hybrid compound of methylene blue,

388   reductant, surfactant and saponite. *Anal Sci* 20:1153-1157.

389   28.   Klementiev AD, Whiteley M. 2022. Development of a versatile, low-cost

390   electrochemical system to study biofilm redox activity at the micron scale.

391   *Appl Environ Microbiol* 88:e0043422.

392   29.   Liu P, Yang Y, Tang Y, Yang T, Sang Z, Liu Z, Zhang T, Luo Y. 2019. Design

393   and synthesis of novel pyrimidine derivatives as potent antitubercular agents.

394   *Eur J Med Chem* 163:169-182.

396 **Table 1. The MICs of clinical drugs against AlRa under aerobic and anaerobic**

397 **conditions using various media.**

Class/Targets	Drugs*	MIC (μg/mL)/aerobic conditions		MIC (μg/mL)/anaerobic conditions	
		Glycerol	Cholesterol	Glycerol	Cholesterol
Cell Wall	EMB	2	4	>64	>64
	INH	0.125	0.125	>64	>64
	VAN	16-32	16-32	>64	>64
	ETH	64	64	>64	>64
RNA polymerase	RIF	0.0625	0.0625-0.125	0.125-0.5	0.25-2
	RFB	0.008	0.008	0.03125-0.125	0.03125-0.0625
Ribosome	AMK	0.25-1	0.25-0.5	1-8	2-16
	STR	0.25-0.5	0.25-0.5	1-8	2-16
Unknown	POA	400	400	>800	>800
DNA gyrases	MOX	0.0625-0.25	0.125	0.125-2	8-32
	LEV	0.125-0.5	0.25-1	>64	>64
Respiration/ Production of NO	PA-824	0.0625	0.0625-0.125	1-4	1-4
Folic acid synthesis	SMX	> 64	> 64	>64	>64
	PAS	0.25-1	> 64	>64	>64
Respiration/ Electron transport chain	TB47	0.004-0.008	0.004-0.008	0.125-0.25	0.0625-0.25
Respiration/ ATP synthesis	BDQ	0.5-1	1-2	4-8	4-16
Unknown	CLO	0.5-1	1-2	2-8	8-16

*EMB, ethambutol; INH, isoniazid; VAN, vancomycin; ETH, ethionamide; RIF,

rifampin; RFB, rifabutin; AMK, amikacin; STR, streptomycin; MOX, moxifloxacin;

LEV, levofloxacin; PA-824, pretomanid; SMX, sulfamethoxazole; PAS, para-

aminosalicylic; BDQ, bedaquiline; CLO, clofazimine.

Figure 1. The time-killing curves of actively growing *M. tuberculosis* treated with nine distinct pharmacological categories of drugs

under aerobic condition using various media.

Solvent, DMSO or distilled water; G, 7H9 supplied with glycerol; C, 7H9 enriched with cholesterol; INH, isoniazid; RIF, rifampin; AMK,

amikacin; MOX, moxifloxacin; PA-824, pretomanid; PAS, para-aminosalicylic; BDQ, bedaquiline; CLO, clofazimine.

Figure 2. The growth curves of nonreplicating *M. tuberculosis* after treatment with nine distinct pharmacological categories of drugs,

using various media under anaerobic condition.

Solvent, DMSO or distilled water; G, 7H9 supplied with glycerol; C, 7H9 enriched with cholesterol; INH, isoniazid; RIF, rifampin; AMK,

amikacin; MOX, moxifloxacin; PA-824, pretomanid; PAS, para-aminosalicylic; BDQ, bedaquiline; CLO, clofazimine.

Table 2. MICs of CLO and TB47 against AlRa along with the corresponding interaction profiles of CLO in conjunction with TB47 assessed by checkerboard method under anaerobic condition.

Drugs[*]	7H9 + Glycerol		7H9 + Cholesterol	
	MIC_a[#]	MIC_c[#]	MIC_a[#]	MIC_c[#]
TB47	0.125	0.03125	0.125	0.03125
CLO	8	4	8	4
FICI[□] / Effect	0.75 / partially synergistic		0.75 / partially synergistic	

[*]CLO, clofazimine.

[□]FICI, Fractional inhibitory concentration index; The FICI was calculated as the ratio of the MIC of TB47 in combination to the MIC of TB47 alone, plus the MIC of CLO in combination to the MIC of CLO alone. The FICI was calculated to classify the interactions as one of five kinds: synergistic (FICI ≤ 0.5), partially synergistic (0.5 < FICI < 1.0), additive (FICI = 1.0), indifferent (1.0 < FICI ≤ 4.0) or antagonistic (FICI > 4.0) (29).

[#]MIC_a, MIC alone; MIC_c, MIC in combination.