High-throughput Screening of Compounds against Autoluminescent Nonreplicating *Mycobacterium tuberculosis* under Diverse Conditions

Xirong Tian [a,b,c,d*], Wanli Ma [a,b,c,d*], Buhari Yusuf [a,b,c,d], Chunyu Li [a,b,c,d], H.M. Adnan Hameed [a,b,c,d], Xinyue Wang[e,f], Nanshan Zhong [e], Jinxing Hu[f#], Tianyu Zhang [a,b,c,d,f#]

[a] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese Academy of Sciences (CAS), Guangzhou 510530, China

[b] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious Diseases, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese Academy of Sciences (CAS), Guangzhou 510530, China

[c] University of Chinese Academy of Sciences (UCAS), Beijing 100049, China

[d] China-New Zealand Joint Laboratory on Biomedicine and Health, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[e] Guangzhou National Laboratory, Guangzhou 510005, China

[f] Guangzhou Chest Hospital, Guangzhou 510005, China

[*] These authors contributed equally to this work.

[#] Corresponding author:

Tianyu Zhang, Tel: +86-20-32015270, E-mail: zhang_tianyu@gibh.ac.cn

Jinxing Hu, Tel: +86-20-83594895, E-mail: hujinxing2000@163.com

KEYWORDS

Autoluminescence; Low-oxygen; Persistence; *Mycobacterium tuberculosis*; High-

throughput screening

ABSTRACT

The screening of new anti-mycobacterial chemicals is primarily focused on inhibiting

the active growing bacteria. However, a major challenge in tuberculosis control is the

ability of *Mycobacterium tuberculosis* to enter a nonreplicating state for extended

periods, rendering it resistant to many clinical drugs and complicating eradication

efforts. Existing low-oxygen-recovery assays designed for screening compounds

targeting nonreplicating *M. tuberculosis* have limitations, including the colony-

forming unit counting for non-luminous *M. tuberculosis* and the instability of the free

plasmid carrying *luxAB* genes in luminescent *M. tuberculosis*, along with exogenous

substrate requirements for light producing. Moreover, these assays fail to accurately

replicate the growth conditions of nonreplicating *M. tuberculosis in vitro*, thus

resulting in less convincing results. To address these challenges, we have developed

an autoluminescence-based, cholesterol-enriched culture evaluation model to assess

17 anti-tuberculosis drugs of different classes against nonreplicating *M. tuberculosis*.

Our findings indicate that the relative light unit, measured in real-time, serves as a

reliable surrogate marker for colony-forming unit, which typically becomes available

one month later. This suggests the utility of our model for the rapid determination of

drug susceptibility dynamically. The autoluminescent *M. tuberculosis*, harbouring

luxCDABE gene cluster within its genome, can emit blue-green light stably and

autonomously without requiring an external substrate supplement. The minimal

inhibitory concentrations of all the drugs tested under anaerobic conditions are

significantly different from that detected in aerobic environment. Our model allows

for rapid, precise, and efficient assessment of drug activity under anaerobic conditions,

thereby enabling a more comprehensive evaluation of anti-mycobacterial efficacy.

Overall, our model represents a significant advancement in anti-tuberculosis drug

discovery and pharmaceutical development.

INTRODUCTION

The highly infectious pathogen *Mycobacterium tuberculosis* is the causative agent of tuberculosis (TB), and was responsible for 10.6 million TB individuals globally in 2022 (1). The emergence of drug-resistant *M. tuberculosis* strains, coupled with lethal coinfections with human immunodeficiency virus, has significantly exacerbated the global healthcare crisis, necessitating the exploration of novel drugs or treatment regimens against *M. tuberculosis* (2, 3). Notably, after four decades of persistent efforts, bedaquiline (BDQ), the first new anti-TB medication, received approval from the United States Food and Drug Administration in 2012 (4-6).

The persistence of nonreplicating *M. tuberculosis* within lung lesions, capable of surviving for decades and adapting to hypoxic conditions, poses a formidable challenge to TB eradication (7-9). A comprehensive understanding of the complex host environment in which *M. tuberculosis* resides is crucial for developing effective control strategies (10). Previous studies have revealed *M. tuberculosis*'s ability to exploit host-derived cholesterol as a primary carbon source during infection (11). By incorporating cholesterol into growth media, it is more accurately replicate the growth environment of nonreplicating *M. tuberculosis* compared to traditional glycerol-enriched media.

Despite the availability of well-established *in vitro* low-oxygen-recovery assays for assessing evaluate drug efficacy against nonreplicating *M. tuberculosis*, these methods still have certain limitations (12-14). Some methods rely on enumerating colony forming units (CFUs) or visible bacterial growth, requiring several months for the completion of testing. Others have employed a luminescent *M. tuberculosis* strain with a free plasmid carrying *luxAB* genes to screen for chemical agents against nonreplicating mycobacteria (6). However, this approach is not favored due to its

dependence on additional substrates and the instability of the plasmid (6). Our

research introduces an autoluminescence-based, low-oxygen, and cholesterol-enriched

evaluation model that utilizes autoluminescent *M. tuberculosis* H37Ra (AlRa),

emitting stable and efficient blue-green light dynamically over an extended period

without requiring substrate supplementation. Bacterial growth is quantified

dynamically by measuring relative light units (RLUs) at various time points during

the recovery process. In conclusion, our model is designed to be straightforward,

time-efficient, labor- and resource-effective, while also facilitating high-throughput

screening of drugs and drug combinations against nonreplicating *M. tuberculosis*.

RESULTS

MICs determination using glycerol- and cholesterol-enriched media under

aerobic conditions. Seventeen anti-TB drugs from distinct classes were selected for

this study. Cholesterol serves as a primary carbon source during *M. tuberculosis*

infection, mimicking the nutritional conditions encountered during the course of *M.*

tuberculosis infection. Moreover, under aerobic conditions, the minimal inhibitory

concentrations (MICs) of the drugs against *M. tuberculosis* remained largely

consistent, irrespective of whether glycerol or cholesterol was used as the carbon

source. An exception to this was observed with para-aminosalicylic acid (PAS), as

detailed in Table 1 and Figure 1. In the glycerol-enriched medium, the MIC of PAS

ranged from 0.25 to 1 μg/mL, however, it increased dramatically to over 64 μg/mL in

the cholesterol-enriched medium (Table 1). The growth curves of actively growing

AlRa treated with nine drugs from different categories are shown in Figure 1, and

those treated with the remaining eight drugs are shown in Figure S1.

MICs determination using glycerol- and cholesterol-enriched media under

anaerobic conditions. We systematically evaluated 17 anti-TB drugs targeting

various crucial aspects of *M. tuberculosis* metabolism, including the cell wall, RNA

polymerase, ribosome and respiration. Our findings revealed that the MICs of these

drugs against *M. tuberculosis* were generally higher under anaerobic conditions

compared to aerobic conditions with the exception of rifampin (RIF) and rifabutin

(RFB), as depicted in Table 1, Figures 2 and Figure S2. Additionally, the MICs of the

drugs remained relatively consistent under anaerobic conditions, regardless of

whether the medium was supplemented with glycerol or cholesterol, with the

exception of moxifloxacin (MOX). The MICs of MOX against nonreplicating *M.*

tuberculosis were found to be in the range of 0.125-2 μg/mL in the presence of

glycerol, whereas in the presence of cholesterol, the MICs spanned from 8-32 μg/mL

(Table 1). Both MOX and levofloxacin (LEV), the two fluoroquinolones, exhibited

increased MICs under anaerobic conditions, regardless of the presence of cholesterol

or glycerol in the media. Remarkably, MOX displayed a greater efficacy compared to

LEV under both anaerobic and aerobic conditions.

Under anaerobic conditions, drugs targeting the cell wall, such as vancomycin (VAN),

isoniazid (INH), ethambutol (EMB), and ethionamide (ETH), did not display

significant antibacterial activity against nonreplicating *M. tuberculosis* as the MICs

were over 64 μg/mL (Table 1). In contrast, these drugs exhibited notable efficacy

against actively growthing *M. tuberculosis* under aerobic conditions.

Compared to other anti-TB drugs, drugs targeting RNA polymerase showed lower

MICs with notably effective anti-*M. tuberculosis* activity in both conditions,

particularly under anaerobic conditions. The MICs of RIF ranged from 0.0625 to

0.125 μg/mL and 0.125 to 2 μg/mL under aerobic and anaerobic conditions

respectively (Table 1). RFB displayed enhanced efficacy compared to RIF against

nonreplicating *M. tuberculosis*, with the MICs as low as 0.03125 to 0.125 μg/mL and

0.03125 to 0.0625 µg/mL in glycerol- and cholesterol-enriched media, respectively

(Table 1). These findings emphasized the promising efficacy of RIF and RFB as

candidates for eradicating nonreplicating *M. tuberculosis*, thus establishing a robust

foundation for their clinical application.

The ribosome-targeting drugs, amikacin (AMK) and streptomycin (STR),

demonstrated anti-TB activity against both actively growing and nonreplicating *M.*

tuberculosis. However, the MICs against nonreplicating *M. tuberculosis* were

significantly higher in comparison to those observed against actively growing *M.*

tuberculosis. Pyrazinoic acid (POA), the active form of pyrazinamide, showed no

anti-*M. tuberculosis* activity under anaerobic conditions, with the MIC 800 µg/mL.

Under anaerobic conditions, drugs targeting folic acid synthesis exhibited no evident

anti-*M. tuberculosis* activity. The MIC of PAS ranged from 0.25-1 µg/mL in glycerol-

enriched 7H9 medium under aerobic conditions, which significantly increased to over

64 µg/mL in cholesterol-enriched 7H9 medium under both aerobic and anaerobic

conditions (Table 1).

Pretomanid (PA-824) demonstrated pronounced anti-*M. tuberculosis* activity,

particularly under anaerobic conditions and with cholesterol as the primary carbon

source, with a MIC range of 1-4 µg/mL (Table 1). TB47 exhibited remarkable anti-*M.*

tuberculosis activity at an exceptionally low concentration of 0.004 µg/mL under

aerobic conditions (Table 1). However, its efficacy against *M. tuberculosis* became

evident only when the concentration was increased to 0.0625-0.25 µg/mL under

anaerobic conditions (Table 1). The MICs of BDQ were 0.5-1 µg/mL and 1-2 µg/mL

against actively growing *M. tuberculosis* when cultivated with glycerol or cholesterol,

respectively (Table 1). Nevertheless, under anaerobic conditions, these MICs

increased to 4-8 µg/mL and 4-16 µg/mL with glycerol or cholesterol, respectively

(Table 1). Clofazimine (CLO) exhibited MICs against nonreplicating *M. tuberculosis*

ranging from 2-16 μg/mL, significantly higher than those observed against actively

growing *M. tuberculosis*, which had an MIC range of 0.5-2 μg/mL (Table 1).

The combined effect of TB47 and CLO against nonreplicating AlRa. Our results

indicated that the combination of TB47 and CLO exhibited partial synergy in terms of

sterilizing activity against nonreplicating *M. tuberculosis* under anaerobic conditions,

both with glycerol and cholesterol-enriched media with a Fractional Inhibitory

Concentration Index (FICI) of 0.75 (Table 2). Notably, the MIC of CLO decreased

from 8 to 4 μg/mL in the presence of TB47 at one-fourth of its MIC. These results

highlighted the potential of this drug combination for clinical evaluation against

nonreplicating *M. tuberculosis*, which has been proved to have strong sterilizing

activities against *M. tuberculosis* in a well-established mouse model in two studies

carried out by us, and the manuscript of another study is in revision in Antimicrobial

Agents and Chemotherapy (15-17). Moreover, the assessment model established in

this study for evaluating the efficacy of drug combinations against nonreplicating *M.*

tuberculosis showed significant promise for informing future research and testing

endeavors in this field.

DISCUSSION

During TB infection, *M. tuberculosis* transitions into a nonreplicating state, becoming

resistant to many clinical antibiotics and significantly prolonging therapy (18). *M.*

tuberculosis can acquire and metabolize essential host-derived lipids, particularly

cholesterol, which facilitates disease initiation and progression (19). This necessitates

the establishment of a representative model that faithfully replicates the growth

conditions of nonreplicating *M. tuberculosis* within the host environment. This model

is characterized by oxygen deprivation and cholesterol enrichment of growth medium,

with the goal of screening for effective drugs against nonreplicating *M. tuberculosis*.

To fulfill this objective, we employed AlRa, a strain preserved in our laboratory, to

establish an autoluminescence-based drug efficacy evaluation model for high-

throughput screening of compounds against nonreplicating *M. tuberculosis*. The RLUs

emitted by AlRa, producing blue-green light, exhibit a strong correlation with CFUs,

thus serving as a surrogate marker rather than CFUs for quantifying *M. tuberculosis*

numbers in culture (20).

During the establishment of our model, a notable increase in the growth rate of *M.*

tuberculosis was observed in cholesterol-enriched medium compared to glycerol-

enriched medium, both continuous cultivation under aerobic conditions and recovery

after cultivation under anaerobic conditions. This observation supports previous

studies suggesting that cholesterol serves as the primary carbon source during *M.*

tuberculosis infection (11). Our findings suggested that RIF, RFB, PA-824, and TB47

demonstrated relatively significant potential for inhibiting nonreplicating *M.*

tuberculosis under anaerobic conditions, as previously documented (21-23).

Conversely, drugs targeting the mycobacterial cell wall, such as INH, EMB, ETH, and

VAN, showed negligible anti-mycobacterial activity under anaerobic conditions,

consistent with prior published studies (12, 24). The resistance of nonreplicating *M.*

tuberculosis to these drugs was attributed to the slowed process of cellular division

under anaerobic condition. Drugs that target folic acid synthesis, such as

sulfamethoxazole (SMX) and PAS, manifested minimal bactericidal activities against

nonreplicating *M. tuberculosis* (Table 1). PAS also showed limited activity against

actively growing *M. tuberculosis* with a MIC exceeding 64 μg/mL, when applying the

media supplemented with cholesterol. As a prodrug activated through the folate

synthesis pathway, PAS can be converted into folate intermediate analogs, thereby

disrupting *M. tuberculosis*'s folate metabolism (25, 26). It has been hypothesized that

the addition of cholesterol, which prevents PAS from entering *M. tuberculosis*, or the

unique mechanism of action of PAS, contributes to its ineffectiveness against *M.*

tuberculosis under cholesterol-enriched media. Ribosome-targeting drugs, AMK and

STR, exhibited superior activity against nonreplicating *M. tuberculosis* compared to

cell wall-targeting drugs (Table 1). The results are consistent with the observations of

some Chinese clinicians, who have noted good treatment efficacy in multi-drug

resistant TB patients treated with AMK. This suggests that AMK possesses relatively

potent activity in controlling *M. tuberculosis* persisters, despite injection is one of its

disadvantages. The MIC of POA against nonreplicating *M. tuberculosis* exceeded 800

μg/mL. We speculated that anaerobic environment might influence the membrane

potential of *M. tuberculosis*, which may contribute to its resistance to POA (27).

Generally, MICs of some drugs against nonreplicating *M. tuberculosis* were generally

higher than those against actively growing *M. tuberculosis*, indicating that some drugs

clinically effective against actively growing *M. tuberculosis* may not effectively treat

nonreplicating *M. tuberculosis* in lung lesions, which is demonstrated clearly in

animal work (16, 28). It is noteworthy that fluoroquinolones (either LEV or MOX)

have not shown significant effects on persisting *M. tuberculosis* in previous animal

studies, which is consistent with the predictions made in this study (16, 28). The

combination of TB47 and CLO demonstrated promising partial synergistic activity

with a FICI as low as 0.75 under anaerobic conditions (Table 2), which supports the

finding observed in the animal studies and is promising for clinical trial (15, 16).

The model developed in this study for assessing drug activity against nonreplicating

M. tuberculosis possesses several notable advantages. Firstly, this method stimulates

the growth conditions of nonreplicating *M. tuberculosis* within the human host by

utilizing media supplemented with cholesterol, thereby ensuring maximal fidelity to

the natural environment of the pathogen. Secondly, the model offers simple

manipulation as AlRa consistently emits blue-green light without requiring additional

substrate aldehyde, thereby guaranteeing stable and consistent results. Thirdly, the

results are easily visualized through the consistent detection and comparison of RLUs

emitted by AlRa at various time intervals. Further exploration should prioritize high-

throughput screening of prospective compounds and treatment regimens against

nonreplicating *M. tuberculosis* using the current model. In summary, this model holds

significant application value and possesses high potential to accelerate the discovery

and development of new drugs and treatment strategies targeting nonreplicating *M.*

tuberculosis, which may closely mirror the real state of *M. tuberculosis* persisters *in*

vivo.

MATERIALS AND METHODS

Mycobacterial strains and culture conditions. The mycobacterial strain utilized in

this study is AlRa, an autoluminescent *M. tuberculosis* H37Ra, which has been

genetically engineered to incorporate the *luxCDABE* gene cluster into its genome (20).

AlRa emits blue-green light continuously catalyzed by LuxAB (bacterial luciferase)

and facilitated by LuxCDE (fatty acid reductase complex), which consistently supply

the necessary substrate aldehyde in a recycling form, thereby obviating the need for

external substrate supplementation (20). Notably, the drug susceptibility and growth

rate of AlRa do not significantly differ from those of the wild type *M. tuberculosis*

H37Ra (20). AlRa was cultured at 37□ in Middlebrook 7H9 broth (Difco, Detroit, MI,

USA) supplemented with 10% oleic acid-albumin-dextrose-catalase enrichment (BBL,

Sparks, MD, USA), 0.2% glycerol, and 0.05% Tween 80.

Antimicrobials. All pharmaceutical drugs, including ETH, INH, VAN, RIF, RFB,

AMK, STR, MOX, LEV, PA-824, SMX, PAS, POA, BDQ, and CLO, were purchased

from Meilun (Dalian, China) with a minimum purity threshold of ≥ 95%. Additionally,

TB47 was synthesized and sent to us as a gift by Boji (Guangzhou, China). AMK,

INH, and STR were solubilized in sterilized water, while the remaining drugs were

dissolved in dimethyl sulfoxide (DMSO) obtained from Xilong (Shanghai, China). All

drug solutions were freshly prepared and stored at 4□ until use.

Chemotherapy under aerobic condition. The AlRa were cultured in 7H9 medium

supplemented with Tween80 at 37□ until the optical density, measured at a

wavelength of 600 nm, reached to 0.6-0.8 and the RLUs/mL reached 5 million. The

various drug solutions (4 μL each) were combined with AlRa culture (196 μL, 10-fold

diluted using 7H9 without Tween80 and supplemented with either glycerol or

cholesterol) in a sterilized tube and mixed thoroughly. Negative controls were

prepared using DMSO and sterilized water. Each drug concentration and negative

control were assayed simultaneously in three 1.5 mL EP tubes. The AlRa-drug

mixtures underwent a five-day incubation period, during which the RLUs were

consistently assessed at identical intervals each day. The MIC_{lux} (MIC based on the

detection of autoluminescence) of drugs against the actively growing AlRa is defined

as the lowest concentration at which the RLUs in the experimental group are equal to

or less than 10% of the RLUs in the control group after 5 days.

Culture of nonreplicating AlRa. The AlRa strains were cultured until they reached

an optical density of 0.6-0.8, measured at a wavelength of 600 nm, and a state of

5×10^6 RLUs/mL. Subsequently, the AlRa culture was then transferred to a low-

oxygen cabinet (HUAPO, Guangzhou, China) that had already achieved a steady state.

This cabinet was programmed to maintain specific conditions, including a temperature

of 37□, an oxygen content of 0.8%, and a carbon dioxide concentration of 5%. The

methylene blue solution was then added to the AlRa culture to achieve a final

concentration of 6 μg/mL (29, 30). The cultures were incubated until the blue color

disappeared, typically observed within 7-10 days after the addition of methylene blue.

Drug susceptibility tested under anaerobic condition. Under hypoxic conditions,

various drug solutions (4 μL each) were combined with nonreplicating AlRa culture

(196 μL, 10-fold diluted using 7H9 without Tween80 and supplemented with glycerol

or cholesterol) in a sterile tube and mixed thoroughly. The AlRa-drug mixture was

then incubated for seven days. To remove any potential carryover effects of residual

drugs during the subsequent recovery process, activated carbon (in a volume ratio of

1:5, 50 μL into 200 μL) was added (15). The tubes were subsequently transferred to a

standard aerobic incubator. The RLUs in the co-incubated 1.5 mL EP tubes were

measured and recorded at 7-hour intervals over a period of 28 hours. The

experimental procedure was repeated three times.

Combined activity assay of TB47 and CLO against nonreplicating AlRa. Previous

research from our group has demonstrated that the combined administration of TB47

and CLO significantly exhibits anti-replicating *M. tuberculosis* activity both *in vitro*

and *in vivo* (16). In this study, we aim to determine whether this combination would

also display notable efficacy against nonreplicating *M. tuberculosis* using the

checkerboard method. Specifically, 2 μL of TB47, 2 μL of CLO, and 196 μL of

diluted AlRa culture in a nonreplicating state were simultaneously added to the same

1.5 mL tube. To eliminate any potential carryover effects of the drugs following seven

days of coincubation, activated carbon (at a volume ratio of 1:5, 50 μL into 200 μL)

was utilized (26). Subsequently, the tubes containing the co-culture were transferred

to aerobic conditions, and RLUs were then detected at 7-hour intervals for the

following 28 hours. The FICI was calculated as the ratio of the MIC of TB47 in

combination to the MIC of TB47 alone, plus the MIC of CLO in combination to the

MIC of CLO alone (31). The effects were categorized based on the FICI into five

groups: synergistic (FICI ≤ 0.5), partially synergistic (0.5 < FICI < 1.0), additive

(FICI = 1.0), irrelevant (1.0 < FICI ≤ 4.0), and antagonistic (FICI > 4.0) (31).

Statistical analysis. Before performing statistical analysis, the RLU values were

transformed into their logarithmic counts. The time-kill curves were graphically

depicted using GraphPad Prism version 8.3.0.

ACKNOWLEDGMENTS

This work received support from the National Key R&D Program of China

(2021YFA1300904), the National Natural Science Foundation of China

(21920102003) and by the Guangzhou Science and Technology Planning Project

(2023A03J0992). The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

AUTHOR CONTRIBUTIONS

Conceived the project: Tianyu Zhang, Xirong Tian, Wanli Ma, Nanshan Zhong;

Designed the research: Tianyu Zhang, Xirong Tian, Wanli Ma, Xinyue Wang, Jinxing

Hu, Nanshan Zhong; Performed the studies: Xirong Tian, Wanli Ma, Chunyu Li,

Xinyue Wang; Interpreted the results: Tianyu Zhang, Xirong Tian, Wanli Ma, Jinxing

Hu; Drafted the manuscript: Tianyu Zhang, Xirong Tian, Wanli Ma, Buhari Yusuf,

H.M. Adnan Hameed; Final approval of the manuscript: all authors.

REFERENCES

1. World Health Organization (WHO). 2023. Global Tuberculosis Report.

Available at: https://iris.who.int/bitstream/handle/10665/373828/97892400838

51-eng.pdf?sequence=1

2. Sala C, Dhar N, Hartkoorn RC, Zhang M, Ha YH, Schneider P, Cole ST. 2010.

Simple model for testing drugs against nonreplicating *Mycobacterium*

tuberculosis. *Antimicrob Agents Chemother* 54:4150-4158.

3.   Koch A, Mizrahi V. 2018. *Mycobacterium tuberculosis*. *Trends Microbiol*

26:555-556.

4.   Volynets GP, Tukalo MA, Bdzhola VG, Derkach NM, Gumeniuk MI,

Tarnavskiy SS, Starosyla SA, Yarmoluk SM. 2019. Benzaldehyde

thiosemicarbazone derivatives against replicating and nonreplicating

Mycobacterium tuberculosis. *J Antibiot (Tokyo)* 72:218-224.

5.   Kadura S, King N, Nakhoul M, Zhu H, Theron G, Köser CU, Farhat M. 2020.

Systematic review of mutations associated with resistance to the new and

repurposed *Mycobacterium tuberculosis* drugs bedaquiline, clofazimine,

linezolid, delamanid and pretomanid. *J Antimicrob Chemother* 75:2031-2043.

6.   He W, Liu C, Liu D, Ma A, Song Y, He P, Bao J, Li Y, Zhao B, Fan J, Cheng Q,

Zhao Y. 2021. Prevalence of *Mycobacterium tuberculosis* resistant to

bedaquiline and delamanid in China. *J Glob Antimicrob Resist* 26:241-248.

7.   Srivastava S, Cirrincione KN, Deshpande D, Gumbo T. 2020. Tedizolid,

faropenem, and moxifloxacin combination with potential activity against

nonreplicating *Mycobacterium tuberculosis*. *Front Pharmacol* 11:616294.

8.   Wayne LG, Sohaskey CD. 2001. Nonreplicating persistence of *mycobacterium*

tuberculosis. *Annu Rev Microbiol* 55:139-163.

9.   Iacobino A, Piccaro G, Giannoni F, Mustazzolu A, Fattorini L. 2017. Fighting

tuberculosis by drugs targeting nonreplicating *Mycobacterium tuberculosis*

bacilli. *Int J Mycobacteriol* 6:213-221.

10.   Wipperman MF, Sampson NS, Thomas ST. 2014. Pathogen roid rage:

cholesterol utilization by *Mycobacterium tuberculosis*. *Crit Rev Biochem Mol*

330   *Biol* 49:269-293.

331   11.   Moopanar K, Nyide ANG, Senzani S, Mvubu NE. 2023. Clinical strains of

332   *Mycobacterium tuberculosis* exhibit differential lipid metabolism-associated

333   transcriptome changes in *in vitro* cholesterol and infection models. *Pathog Dis*

334   81:ftac046.

335   12.   Cho SH, Warit S, Wan B, Hwang CH, Pauli GF, Franzblau SG. 2007. Low-

336   oxygen-recovery assay for high-throughput screening of compounds against

337   nonreplicating *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother*

338   51:1380-1385.

339   13.   Cho S, Lee HS, Franzblau S. 2015. Microplate Alamar blue assay (MABA)

340   and low oxygen recovery assay (LORA) for *Mycobacterium tuberculosis*.

341   *Methods Mol Biol* 1285:281-292.

342   14.   Macabeo APG, Vidar WS, Chen X, Decker M, Heilmann J, Wan B, Franzblau

343   SG, Galvez EV, Aguinaldo MAM, Cordell GA. 2011. *Mycobacterium*

344   *tuberculosis* and cholinesterase inhibitors from Voacanga globosa. *Eur J Med*

345   *Chem* 46:3118-3123.

346   15.   Yu W, Yusuf B, Wang S, Tian X, Hameed HMA, Lu Z, Chiwala G, Alam MS,

347   Cook GM, Maslov DA, Zhong N, Zhang T. 2021. Sterilizing effects of novel

348   regimens containing TB47, clofazimine, and linezolid in a murine model of

349   tuberculosis. *Antimicrob Agents Chemother* 65:e0070621.

350   16.   Yu W, Chiwala G, Gao Y, Liu Z, Sapkota S, Lu Z, Guo L, Khan SA, Zhong N,

351   Zhang T. 2020. TB47 and clofazimine form a highly synergistic sterilizing

352   block in a second-line regimen for tuberculosis in mice. *Biomed Pharmacother*

353   131:110782.

17.   Yu W, Ju Y,Han X, Tian X, Ding J, Wang S, Gao Y, Li L, Li Y, Zhong N,

Zhang T. Bactericidal and sterilizing activity of sudapyridine-clofazimine-
TB47 combined with linezolid or pyrazinamide in a murine model of
tuberculosis. *Antimicrob Agents and Chemother* (Revised and submitted,
AAC00124-24R2).

18.   Grant SS, Kawate T, Nag PP, Silvis MR, Gordon K, Stanley SA, Kazyanskaya
E, Nietupski R, Golas A, Fitzgerald M, Cho S, Franzblau SG, Hung DT. 2013.
Identification of novel inhibitors of nonreplicating *Mycobacterium*
tuberculosis using a carbon starvation model. *ACS Chem Biol* 8:2224-2234.

19.   Wilburn KM, Fieweger RA, VanderVen BC. 2018. Cholesterol and fatty acids
grease the wheels of *Mycobacterium tuberculosis* pathogenesis. *Pathog Dis*
76:fty021.

20.   Yang F, Njire MM, Liu J, Wu T, Wang B, Liu T, Cao Y, Liu Z, Wan J, Tu Z,
Tan Y, Tan S, Zhang T. 2015. Engineering more stable, selectable marker-free
autoluminescent mycobacteria by one step. *PLoS One* 10:e0119341.

21.   Hu Y, Coates ARM, Mitchison DA. 2003. Sterilizing activities of
fluoroquinolones against rifampin-tolerant populations of *Mycobacterium*
tuberculosis. *Antimicrob Agents Chemother* 47:653-657.

22.   Stover CK, Warrener P, VanDevanter DR, Sherman DR, Arain TM, Langhorne
MH, Anderson SW, Towell JA, Yuan Y, McMurray DN, Kreiswirth BN, Barry
CE, Baker WR. 2000. A small-molecule nitroimidazopyran drug candidate for
the treatment of tuberculosis. *Nature* 405:962-966.

23.   Wayne LG, Sramek HA. 1994. Metronidazole is bactericidal to dormant cells
of *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 38:2054-2058.

24.   Commandeur S, Iakobachvili N, Sparrius M, Nur MM, Mukamolova GV,
Bitter W. 2020. Zebrafish embryo model for assessment of drug efficacy on

mycobacterial persisters. *Antimicrob Agents Chemother* 64:e00801-20.

25.   Zhang Y, Wang S, Chen X, Cui P, Chen J, Zhang W. 2024. Mutations in the

382   promoter region of methionine transporter gene metM (Rv3253c) confer para-

383   aminosalicylic acid (PAS) resistance in *Mycobacterium tuberculosis*. *mBio*

384   15:e0207323.

385   26.   Wei W, Yan H, Zhao J, Li H, Li Z, Guo H, Wang X, Zhou Y, Zhang X, Zeng J,

386   Chen T, Zhou L. 2019. Multi-omics comparisons of p-aminosalicylic acid

387   (PAS) resistance in folC mutated and un-mutated *Mycobacterium tuberculosis*

388   strains. *Emerg Microbes Infect* 8:248-261.

389   27.   Zhang Y, Shi W, Zhang W, Mitchison D. 2014. Mechanisms of pyrazinamide

390   action and resistance. *Microbiol Spectr* 2:MGM2-0023-2013.

391   28.   Grosset JH, Tyagi S, Almeida DV, Converse PJ, Li S-Y, Ammerman NC,

392   Bishai WR, Enarson D, Trébucq A. 2013. Assessment of clofazimine activity

393   in a second-line regimen for tuberculosis in mice. American Journal of

394   Respiratory and Critical Care Medicine 188:608-612.

395   29.   Sumitani M, Takagi S, Tanamura Y, Inoue H. 2004. Oxygen indicator

396   composed of an organic/inorganic hybrid compound of methylene blue,

397   reductant, surfactant and saponite. Analytical Sciences : the International

398   Journal of the Japan Society For Analytical Chemistry 20:1153-1157.

399   30.   Klementiev AD, Whiteley M. 2022. Development of a Versatile, Low-Cost

400   Electrochemical System to Study Biofilm Redox Activity at the Micron Scale.

401   Applied and Environmental Microbiology 88:e0043422.

402   31.   Liu P, Yang Y, Tang Y, Yang T, Sang Z, Liu Z, Zhang T, Luo Y. 2019. Design

403   and synthesis of novel pyrimidine derivatives as potent antitubercular agents.

404   European Journal of Medicinal Chemistry 163:169-182.

Table 1. The MICs of clinical drugs against AlRa under aerobic and anaerobic conditions using various media.

Class/Targets	Drugs[*]	MIC (µg/mL)/aerobic conditions		MIC (µg/mL)/anaerobic conditions	
		Glycerol	Cholesterol	Glycerol	Cholesterol
Cell Wall	EMB	2	4	>64	>64
	INH	0.125	0.125	>64	>64
	VAN	16-32	16-32	>64	>64
	ETH	64	64	>64	>64
RNA polymerase	RIF	0.0625	0.0625-0.125	0.125-0.5	0.25-2
	RFB	0.008	0.008	0.03125-0.125	0.03125-0.0625
Ribosome	AMK	0.25-1	0.25-0.5	1-8	2-16
	STR	0.25-0.5	0.25-0.5	1-8	2-16
Unknown	POA	400	400	>800	>800
DNA gyrases	MOX	0.0625-0.25	0.125	0.125-2	8-32
	LEV	0.125-0.5	0.25-1	>64	>64
Respiration/ Production of NO	PA-824	0.0625	0.0625-0.125	1-4	1-4
Folic acid synthesis	SMX	> 64	> 64	>64	>64
	PAS	0.25-1	> 64	>64	>64
Respiration/ Electron transport chain	TB47	0.004-0.008	0.004-0.008	0.125-0.25	0.0625-0.25
Respiration/ ATP synthesis	BDQ	0.5-1	1-2	4-8	4-16
Unknown	CLO	0.5-1	1-2	2-8	8-16

[*]EMB, ethambutol; INH, isoniazid; VAN, vancomycin; ETH, ethionamide; RIF, rifampin; RFB, rifabutin; AMK, amikacin; STR, streptomycin; MOX, moxifloxacin; LEV, levofloxacin; PA-824, pretomanid; SMX, sulfamethoxazole; PAS, para-aminosalicylic; BDQ, bedaquiline; CLO, clofazimine.

Figure 1. The time-killing curves of actively growing *M. tuberculosis* treated with nine distinct pharmacological categories of drugs

under aerobic condition using various media.

Solvent, DMSO or distilled water; G, 7H9 supplied with glycerol; C, 7H9 enriched with cholesterol; INH, isoniazid; RIF, rifampin; AMK,

amikacin; MOX, moxifloxacin; PA-824, pretomanid; PAS, para-aminosalicylic; BDQ, bedaquiline; CLO, clofazimine.

Figure 2. The growth curves of nonreplicating *M. tuberculosis* after treatment with nine distinct pharmacological categories of drugs,

using various media under anaerobic condition.

Solvent, DMSO or distilled water; G, 7H9 supplied with glycerol; C, 7H9 enriched with cholesterol; INH, isoniazid; RIF, rifampin; AMK,

amikacin; MOX, moxifloxacin; PA-824, pretomanid; PAS, para-aminosalicylic; BDQ, bedaquiline; CLO, clofazimine.

Table 2. MICs of CLO and TB47 against AlRa along with the corresponding

interaction profiles of CLO in conjunction with TB47 assessed by checkerboard

method under anaerobic condition.

Drugs[*]	7H9 + Glycerol		7H9 + Cholesterol	
	MIC_a[#]	MIC_c[#]	MIC_a[#]	MIC_c[#]
TB47	0.125	0.03125	0.125	0.03125
CLO	8	4	8	4
FICI[□] / Effect	0.75 / partially synergistic		0.75 / partially synergistic	

[*]CLO, clofazimine.

[□]FICI, Fractional inhibitory concentration index; The FICI was calculated as the ratio

of the MIC of TB47 in combination to the MIC of TB47 alone, plus the MIC of CLO

in combination to the MIC of CLO alone. The FICI was calculated to classify the

interactions as one of five kinds: synergistic (FICI ≤ 0.5), partially synergistic (0.5 <

FICI < 1.0), additive (FICI = 1.0), indifferent (1.0 < FICI ≤ 4.0) or antagonistic (FICI >

4.0) (31).

[#]MIC_a, MIC alone; MIC_c, MIC in combination.