A consensus motif in *ASH1* and further transcripts unifies several RNA motifs required for

interaction with the She2p/She3p transport machinery and mRNA localization in yeast

Markus Seiler[1#*], Annika Niedner[2#], Simone Heber[2,3], Michael Feldbrügge[4], Ralf-Peter Jansen[5], Dierk

Niessing[2,3*], Kathi Zarnack[1*]

[1]Buchmann Institute for Molecular Life Sciences (BMLS), Goethe University Frankfurt, Max-von-Laue-

Str. 15, 60438 Frankfurt a. M., Germany

[2]Institute of Structural Biology, Helmholtz Zentrum München, Ingolstädter Landstrasse. 1, 85764

Neuherberg, Germany

[3]Institute of Pharmaceutical Biotechnology, Ulm University, James Franck Ring N27, 89081 Ulm,

Germany

[4]Institute of Microbiology, Heinrich-Heine University Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf,

Germany

[5]Interfaculty Institute of Biochemistry, Tübingen University, Auf der Morgenstelle 34, 72076 Tübingen,

Germany

[#] Equal contribution

[*] Corresponding authors

Keywords: She2p, She3p, localization machinery, RNA localization, RNA zip-code, modeling, consensus

motif, surface plasmon resonance, computational screening

Abstract

Asymmetric localization of the *ASH1* transcript is a central step in the regulation of mating type

switching in *Saccharomyces cerevisae* and a paradigm for localized mRNAs specifically recognized by the

She2p/3p transport machinery. Four RNA elements in *ASH1* are known to mediate bud localization, but it

remained unclear what a consensus motif of all four She2p/3p recognition sites (SRS) might look like. In

this study, we derive the SRS consensus motif, which is characterized by an L-shaped, double-stranded

RNA conformation with two cytosine-containing sequence motifs at conserved positions and a central

asymmetric bulge. It is noteworthy that the spatial arrangement of these features can be retained, despite

their altered order within the nucleotide sequence. We termed these variations "configurations" and

confirmed in biochemical studies that SRS function in *ASH1* elements is preserved when changing

configurations. We tested other known and predicted SRS elements in target transcripts of She2p/3p by

biochemical and biophysical studies and confirmed the conserved SRS function even in switched

configuration. Consistently, we present the *SRS instance collection*, which includes further predictions of

SRS instances of transcript targets of the She2p/3p translocation pathway.

Introduction

The active intracellular transport of effector molecules contributes to the high degree of organization in life and asymmetric mRNA distribution is one way to ensure correct spatiotemporal expression of encoded proteins[1]. In recent years, it has been studied in various polarized cells, in *D. melanogaster* and *Xenopus laevis* oocytes and embryos[2–4], and also in mammalian fibroblasts[5] and neurons[6]. mRNA transport is involved in embryonic patterning, cellular polarity, and asymmetric cell division. Failure of mRNAs to be properly transferred has been shown to cause cellular dysfunction and disease[7]. In these pathways, mRNAs are recognized by the cellular translocation machinery via *cis*-acting elements in their sequences, called zip-codes or localization elements that may be determined by short RNA sequence motifs, conformational folds[8–10] and spatial restraints between these features[11]. These zip-codes are recognized by RNA-binding proteins that target the mRNAs to their final destination[12]. Within the mRNAs, the zip-codes often reside in the 3' untranslated regions (3' UTRs) of mRNAs, but also in 5' UTRs and coding regions[13–18].

Budding yeast (*S. cerevisiae*) has evolved as a model organism for mechanistic analyses on specific targeting of mRNAs to the bud of emerging daughter cells[19,20]. The current model of how the RNA transport machinery is built suggests a step-wise assembly of its core compounds: She2p interacts specifically with its target mRNA and recruits the adapter protein She3p together with the type V myosin Myo4p[21–24]. This mRNA ribonucleoprotein (mRNP) complex is then transported along the actin cytoskeleton. The paradigm, the yeast *ASH1* mRNA, is localized to the bud tip enabling the asymmetric distribution of its encoded transcription factor Ash1p into the daughter cell nucleus where it represses mating-type switching[25–28]. To date, over twenty transcripts have been reported to be targeted via the She2p/3p machinery. Besides the well-characterized She2p/3p target mRNAs *ASH1*, *EAR1*, and *WSC2*, further candidates were identified using combined RNA immunoprecipitation and microarray techniques[24,29]. In some target mRNAs, the regions containing putative zip-codes had been narrowed down by yeast three-hybrid and *in vivo* localization studies[30]. However, besides of *ASH1*, precise positions of the She2p/3p recognition sites (SRS) remained unknown in most cases. Within the *ASH1* mRNA, four localization elements with intrinsic SRS

function were mapped in accurate fashion[13,14,19]. Three of these (E1, E2A and E2B) were shown to reside

within the coding region, whereas E3 overlapped the stop codon and extended into the 3' UTR

(**Supplementary Fig. 1A, Tab. 1**). Each *ASH1* SRS element was sufficient for triggering the assembly of

the She2p/3p complex and for localizing a reporter RNA to the bud and was predicted to exhibit RNA folds

with extended stem-loop structures. An *in-vivo* screen with mutated *ASH1* fragments supported the notion

that a three-dimensional fold was involved in zip-code function. The fold was described as a stem with

conserved nucleotides on either side: a singular cytosine and a CGA triplet, with a distance of six

nucleotides between the cytosines. Searching for similar RNA folds identified functional SRS elements in

IST2 and *EAR1*[10]. However, these folds lacked an apparent consistency with some of the four *ASH1* SRS

elements and a coherent model for specific SRS recognition by the She2p/3p transport machinery has

remained elusive. Recently, the interaction interface between E3 and She2p/3p was resolved using X-ray

crystallography: a symmetric tetramer of She2p exhibited two interfaces interacting with E3 on opposite

sides of the complex[8]. As suggested before[31], the affinity was increased by She3p in a synergistic manner.

Intriguingly, the ternary complex formation was accompanied by an induced fit in the SRS element: For

the unbound E3 RNA, SHAPE experiments suggested an elongated conformation[8]. When interacting with

She2p/3p, E3 displayed a sharp turn in one of its bulge regions, resulting in a kinked global conformation

that was also confirmed by NMR structures.

In this study, we define a coherent SRS consensus motif that unifies the features of E1, E2A, E2B

and E3 in *ASH1* and that also classifies SRS elements of further bud-localizing transcripts (*WSC2, EAR1,*

IST2). It exhibits a kinked stem-asymmetric bulge-stem-bulge structure, with the recurrent singular cytosine

in the asymmetric bulge and the CGA motif in a second bulge. One characteristic of the consensus motif is

its arrangement of features, conserved in the overall spatial arrangement but present in two variations

("configurations") in linear sequences. Electrophoretic mobility shift assays and surface plasmon resonance

studies showed that E3 and other SRS elements could be switched between configurations without

impairing She2/3p binding. Using computational screening, we identified potential SRS instances in further

twelve transcripts known to interact with She2/3p. Finally, we introduce the *SRS instance collection*,

describing in detail predicted SRS elements and their positions in further RNA targets of the She2p/3p

translocation pathway.

Results

Bioinformatics predictions reveal a consensus capturing all She2/3p recognition sites of *ASH1*

We wondered if a general concept may exist to describe all four localization elements of the *ASH1*

mRNA that act as She2p/3p recognition sites (SRS). For this study, we could rely on a recently published

crystal structure that showed in detail how the *ASH1* localization element E3 interacts in a kinked stem

structure with She2p/3p[8]. Of note, this interaction involved a molecularly induced fit upon binding that

resulted in an asymmetric bulge in the RNA stem with one of its functional cytosines deeply buried inside

the protein interface. The asymmetric bulge of E3 displayed a large stretch ("Lbulge") and a small one

("Sbulge"), with a length ratio of 6 nt : 3 nt (**Fig. 1A, right**) or 4 nt : 1 nt, respectively, if considering

atypical base pairing inside the bulge as further base pairs. We hypothesized that the ability to form such

an asymmetric bulge could be a pre-requisite of the E3 conformation, whose composition would require

different lengths to be kinked. In a next step, we performed a systematic survey of such arrangements in

further SRS elements of *ASH1*. However, common RNA conformation predictors do not consider precise

structural rearrangements due to RNA-protein interaction very well. In a step-wise strategy

(Supplementary Fig. 2, Material and Methods), we first evaluated which of the commonly used

prediction tools would predict the best approximation of the kinked E3 conformation in complex with

She2p/She3p[8] just from the single E3 nucleotide sequence. Of note, we found *Kinefold*[32] to predict the

asymmetric bulge in close approximation (5 nt : 2 nt) that could even be closer approximated when

considering atypical base pairing as found in the asymmetric bulge of the crystal structure, resulting in (6

nt : 3 nt) (**Fig.1**, **Supplementary Fig. 3A**).

For E2A, we found the predicted conformation with an asymmetric bulge with exactly the same

length ratio as found in E3 (5 nt : 2 nt) using *Kinefold* (**Fig. 1B left**). Of note, after 180° rotation, the

predicted E2A conformation harmonized perfectly with the E3 conformation (**Fig. 1A right, 1B right**).

Furthermore, this 180° rotation caused the previously diverging sequence motifs of E3 to fit now to the

single cytosine and CGA motifs of E2A. Surprisingly, sequence comparison revealed 89% sequence

identity between E3 and E2A in the asymmetric bulge and flanking regions (**Supplementary Fig. 1B**). For

E1, the 180° rotation of its predicted conformation resulted in an arrangement comparable with E3 (**Fig.**

1D), but with different length proportions of the asymmetric bulge and a lower degree of sequence identity.

These findings suggested one blueprint, with a common overall spatial arrangement of its

(conformational and sequence) features in E3, E2A and E1 that we referred to as "SRS consensus motif"

(**Fig. 1E,F**). Of note, despite of this conserved spatial arrangement of features, the linear order of features

was interchangeable in the nucleotide sequences (**Fig. 1F**). This was due to the fact that the conformations

of E3, E2A and E1 are harmonized by a 180 degree rotational step of E3. Subsequently, the 5' start of E3

was at the opposite site of the SRS consensus motif than for E2A and E1, resulting in different feature

orders in the nucleotide sequence (**Fig. 1E**). We termed these variations "configurations", with "config-I"

observed in E3 and "config-II" in E2A and E1. Further, we could harmonize the annotation of the sequence

motifs in E3 by labeling the single cytosine motif in the asymmetric bulge as C_1 and the cytosine in the

CGA motif as C_2 (**Fig. 1A, right, 1E**), analogously to E2A and E1. For E2B, at a first glance, the

conformational prediction of E2B displayed the hypothesized "Lbulge" even shorter than the "Sbulge" (**Fig.**

1C, left). Remarkably, Kinefold predicted an alternative conformation with an asymmetric bulge in

agreement with the SRS consensus motif (6 nt : 3 nt), but with a higher content of energy (calculated free

energy = -10.3 kcal/mol versus -7.4 kcal/mol), which could be rearranged by sliding between sequence

repeats in one inverted repeat (**Fig. 1C, right**). Altogether, we had derived a unified model of the SRS

consensus motif (see **Info-Box**). In a subsequent step, we could confirm that the model was sufficient to

summarize previous functional studies with mutationally-derived decreases in RNA localization in all four

ASH1 SRS elements[10,13,14,25] (**Material and Methods)**. For an overview, the distinct modeling steps

(initially characterized elements, Kinefold prediction, rotation, and mutual feature alignments) are

displayed in (**Fig. 2**).

146 **The two configurations are interchangeable in E3 without loss-of-function *in-vitro***

147   One unexpected characteristic of the SRS consensus motif was the spatial conservation of its

148 features in its overall conformation while the order of features in the nucleotide sequences was found to be

149 swapped for the different configurations. To validate if we had grasped the essence of the consensus, we

150 investigated if an artificial switch between the different sequence orders (and the configurations) would

151 conserve SRS function. For this purpose, we used the well-characterized construct $E3_{51nt}wt^8$ and switched

152 its feature order in the nucleotide sequence (config-I) to design a config-II construct ($E3_{51nt}$switch, **Fig. 3A**).

153 Based on the observation that the E3 crystal structure exhibited an interaction of She2p/3p with C_2G but

154 not with C_2GA, and for comparability reasons with previous studies, we limited the construct design to

155 C_2G. Using an electrophoretic mobility shift assay (EMSA), we examined $E3_{51nt}$switch for interaction with

156 She2p and She3p in a titration approach. We used the fusion protein She2p-She3p, which comprised both,

157 She2p and a She3p fragment and was demonstrated to be sufficient to mediate the synergistic binding of

158 both proteins[8]. For the wildtype construct $E3_{51nt}wt$, we observed an interaction strength at the expected

159 nanomolar protein concentration (**Fig. 3A**)[8]. As a control, we tested the construct $E3_{51nt}M6$ characterized

160 by reduced interaction[8] due to a $C_{1813}G$ substitution of C_2. In agreement of our model, $E3_{51nt}wt$ and

161 $E3_{51nt}$switch, interacted with She2p-She3p at comparable nanomolar protein concentration (**Fig. 3A**). These

162 similar binding affinities contrasted with $E3_{51nt}M6$, which displayed severely impaired interaction.

163   Since the $E3_{51nt}M6$ mutant showed impaired but still considerable binding to She2p-She3p-fusion

164 protein, we used the full-length proteins to assess the binding features of E3 elements. Incubation with

165 constant She3p concentration of 0.56 μM and titration of She2p (0.02 - 5 μM) confirmed that both

166 constructs, $E3_{51nt}wt$ and $E3_{51nt}$switch, supported binding even at the lowest She2p concentration used in this

167 experiment (0.02 μM). In contrast, binding of the full-length proteins to $E3_{51nt}M6$ was abolished at

168 concentrations up to 5 μM (**Supplementary Fig. 4B**). The same was true for a reciprocal approach, in

169 which increasing concentrations of She3p (0.02 - 5 μM) were tested in the presence of high amounts of

170 She2p (5 μM) (**Supplementary Fig. 4C**). Moreover, both experiments showed that She3p efficiently binds

171 E3 in the absence of She2p. Surprisingly, this binding was sensitive to mutation of C_2. In conclusion, these

studies successfully confirmed that E3 can be switched from config-I to config-II without compromising

its functionality to interact with She2p/3p.

When we analogously examined E2A and E1, we reversed the direction of switching (now from

config-II to config-I), resulting in the constructs $E1_{27nt}$switch and $E2A_{32nt}$switch (**Supplementary Fig. 5**).

Again, in EMSA validations, we could demonstrate that both constructs largely exhibited the binding

properties to She2p-She3p-fusion. In summary, the "config-switch" approach supported our model, which

suggested that SRS features could be rearranged in the RNA sequence while retaining function. We could

demonstrate that E1, E2A, and E3 recognized She2p/3p regardless of configuration, demonstrating the

essential characteristics of RNA-She2p/3p recognition were reflected in the SRS consensus motif. As a

conclusion, these findings indicate that SRS function requires very few conserved bases in defined positions

of a stem-loop, which must accommodate a bulge that is flexible enough to open up upon binding by She2p-

She3p to adopt a kinked L-shaped conformation in the co-complex.

In a fine-grained approach, we used the analytical power of surface plasmon resonance (SPR), to

investigate the influence of the functional cytosines C_1 and C_2 on SRS function. Previous studies had

determined a preferred distance of six nucleotides between these cytosines[10]. Using the functionally

minimal[8] construct $E3_{29nt}$, we introduced varying distances between C_1 and C_2 and performed titration series

in a surface plasmon resonance experiment with a She2p-She3p-fusion construct (125 nM) (**Fig. 3B**). As a

result, for the wildtype construct $E3_{29nt}$wt, we observed a binding affinity of approx. K_D = 0.89 nM (**Fig.**

3B, Supplementary Fig. 6). In contrast, constructs with C_1-C_2 distances of five and seven nucleotides led

to a reduction in binding of at least one order of magnitude (**Fig. 3B**, **Supplementary Tab. 1**), confirming

the findings of 6 nt as optimal C_1-C_2 distance.

A computational algorithm to screen for the consensus in known She2p/3p substrates

We performed computational screening (details, see **Material and Methods**) to identify further

instances of the SRS consensus motif in further target sequences of the She2p/3p transport machinery. For

this purpose, we screened a list of 24 transcripts reported to show *in-vitro* binding by She2p/3p or bud

localization *in-vivo*[24,29,30]. The identified sequence regions represented potential SRS instances and were

evaluated for their potential to form the correct conformation of one of the configurations using *Kinefold*.

We intended to extend this conformational evaluation by integrating data describing the degree of unpaired

state of nucleotides (obtained in a remarkable genome-wide study using parallel analysis of RNA structure

(PARS)[33]). This strategy was promising, but for our purposes the data were not detailed enough, due to

competing conformations that could not be differentiated in the averaged values per nucleotide position,

even for some of the strong *ASH1* SRS elements (**Material and Methods, Supplemental Fig. 7**).

For the three transcripts *EAR1, IST2* and *WSC2*, their SRS elements had been identified already,

with their sequence regions narrowed down to fragments[10,30]. Now, with knowledge of the SRS consensus

motif, these SRS elements could be classified as a config II arrangements (**Fig. 4C, Supplementary Fig.**

8). Remarkably, for *EAR1* and *IST2*, the asymmetric bulge was even more eccentric than observed in *ASH1*

(*EAR1:* 7 nt : 0 nt; *IST2*: 9 nt : 1 nt). In addition, we identified an additional upstream SRS instance in each

transcript, *EAR1* and *IST2* (**Tab. 2**). In contrast, for *WCS2*-C, the initially identified SRS fragment turned

out to comprise possibly two SRS elements, a canonical one in the center of this fragment, and a potential

SRS instance in the flanking regions, however, with a C_1-C_2 distance of 7 nt (**Supplemental Fig. 8D**).

However, we could not classify the second published instance in *WSC2*, *WSC2*-N, in agreement with our

model (see **Discussion**). As a first outcome of the screening we had identified seven SRS instances (**Fig.**

3C). When we studied the seven canonical SRS elements in detail, we noticed a recurring di-adenosine in

the L-bulge ("AA_L") as a common feature in all elements (**Fig. 3C**), possibly indicating relevance in SRS

function. This was surprising due to the fact that the crystal structure of E3 showed the interaction between

She2p/3p and the AA_L nucleotides occurs exclusively via the backbone atoms of the nucleotides, and thus

unspecifically[8], with their nucleobases having intramolecular contact within the RNA structure. This

observation makes relevance in the context of SRS recognition less likely.

As mentioned before, for most known SRS instances (in 23 of 30 instances in our set), functionality

had been exclusively studied for large RNA fragments or complete transcripts. For 13 of these transcripts,

we could identify now the positions of SRS instances. For 12 transcripts, we found one or two SRS instances

in each sequence, and we found four instances in *TPO1* (**Fig. 4A**). Altogether, in 17 of 24 sequences (*ASH1*,

WSC2, *EAR1*, *IST2* included), we predicted 30 SRS instances, with nine instances (30%) in config-I and 21

instances (70%) in config-II, respectively (**Fig. 4A**). Of notice, more than two thirds of the SRS instances

contained the AA_L di-nucleotide in the Lbulge (**Fig. 4B**).

Validation of SRS elements in *EAR1*, *IST2*, *KSS1* and *TCB3*

As described before, in *IST2* and *EAR1*, sequence fragments with SRS function had been identified

already, and these sequences were also identified in our computational screening (termed *EAR1*-SRS2,

IST2-SRS2; nomenclature see **Tab. 2**). We could classify their functional features, in agreement with the

SRS consensus motif, provided that the consensus was valid for these instances. In such a view, the SRS

elements would resemble the very first characterization (Olivier et al., 2005), however in the larger context

of the SRS consensus motif. To address this aspect, we studied specifically designed SRS constructs,

analogous to our previous studies. In accordance with the consensus, the newly designed constructs

EAR1$_{26nt}$switch and *IST2*$_{29nt}$switch were strictly reduced to those regions assumed to be essential for SRS

function and also switched in configuration (each to config-I) (**Fig. 5A**). As we could show in EMSA

studies, *EAR1*$_{26nt}$switch was bound with affinities comparable to the unmodified wild-type element

EAR1$_{50nt}$wt (**Fig. 5A**). Although for *IST2*$_{29nt}$switch binding was clearly retained, it was considerably weaker

when compared to *IST2*$_{62nt}$wt. In addition, we tested the predicted *KSS1*-SRS1 (config-II, **Tab. 2**) and

TCB3-SRS1 (config-I) for which no detailed position in their transcripts had previously been determined.

As a result, we observed strong binding *in-vitro* for both, *KSS1*$_{72nt}$wt and for the minimal construct

KSS1$_{29nt}$switch (switched to config-I), confirming *KSS1*-SRS1 as a potential SRS element (**Fig. 5B**). In an

analogous manner, we observed for the minimal construct TCB3$_{38nt}$wt (config-I) strong interaction with the

fusion protein suggesting *TCB3*-SRS1 as further SRS element (**Fig. 5B**). In summary, we were able to

demonstrate the identification and *in-vitro* confirmation of new SRS instances. Furthermore, for some of

these instances we were able to prove that the newly designed, size-reduced constructs retain the SRS

function even in switched configuration. In conclusion, the fact that our model, derived exclusively from

ASH1 SRS elements, enabled the identification, size reduction, and config-switch of designed SRS

constructs in previously unexplored sequences while preserving SRS function confirmed its applicability

as an SRS consensus motif.

Validation of further features in the SRS of *EAR1*

In *EAR1* (*YMR171C*), it had been shown for its SRS element (now termed *EAR1*-SRS2) that

substitutions in the sequence motifs decreased function[10]. The substitution of both, C_1 and C_2 (updated

nomenclatures) in the constructs "M1" and "M2"[10] had shown severe effects *in-vivo*. To examine the SRS

consensus motif and *EAR1*-SRS2 in more detail, we chose a fine-grained approach using the $EAR1_{31nt}$switch

construct (switch to config-II) as the base construct for further changes (**Fig. 6A**) and determined its binding

affinity using SPR analysis as described before. As result, we obtained an affinity of the original construct

$EAR1_{31nt}$switch of $K_D = 11$ nM (**Fig. 6D, Supplementary Fig. 9**). Remarkably, compared to the E3 minimal

element, the affinity with the She2p/3p fusion protein was decreased by one order of magnitude ($E3_{29nt}$wt:

$K_D = 0.89$ nM). When we modified $EAR1_{31nt}$switch in the inverted repeat IR1, by inversions or additionally

introduced base pairs, the modified constructs showed affinities in the range of the original construct

suggesting no direct impact on SRS function (**Fig. 6D**). Also, when we analyzed the relevance of the di-

nucleotide AA_L in the Lbulge by deletions of different size (**Fig. 6C**), we observed binding affinities in the

order of magnitude of the basic construct (**Fig. 6D**). The analysis of the functional cytosines C_1 and C_2 was

complex since two cytosines (C_{1601} and C_{1602}) precluded an unambiguous assignment of the C_1 position

(**Fig. 6B**). The substitution of either cytosine (constructs $C_{1-1}U$, $C_{1-2}U$) resulted in a considerable binding

decrease suggesting that both positions contribute to She2p/3p binding (**Fig. 6D**). Unexpectedly, the

deletion of one of the C_1 positions (ΔC_1) or a simultaneous substitution of both cytosines (C_{1-1} U & C_{1-2} U)

restored She2p/3p binding (see **Discussion**). In contrast, substituting C_{1590} in the three-nucleotide C_{1590}GA

motif (C_2U) returned unequivocal results manifesting in a dramatic loss of binding. Altogether, these results

confirmed that neither sequence composition of IR1 nor the AA_L motif in the Lbulge were relevant for

She2p/3p binding *in vitro*. Alterations of the C_1-C_2 distance were tolerated in some SRS contexts but not

others, with a strong bias for a C_1-C_2 distance of six nucleotides. We therefore kept a C_1-C_2 distance

constraint of six nucleotides for subsequent computational SRS screening.

A collection of predicted *SRS* instances in known target transcripts of She2p/3p

Computational screening for the SRS consensus motif had identified SRS instances in 17 of 24

sequences (**Fig. 4A**). After successful *in-vitro* validation of several of the new SRS instances (in *EAR1*,

IST2, *KSS1*, *TCB3*), we finally compiled a collection of these screening results to present to the community

(*SRS instance collection*, **Tab. 2**). This collection contains information on 30 SRS instances in 17 transcripts

that have been reported to show She2p/3p binding *in-vitro* or bud localization *in-vivo* [24,29,30]. In addition, a

standardized nomenclature for SRS elements is introduced and further information is provided on positional

data, classification and other details of SRS instances, several of them are promising new candidates for

future experimental investigations.

Discussion

In this work, we have derived the unified model of a three-dimensional RNA motif that functions as She2p/3p recognition motif (SRS) with the She2p/She3p protein transport machinery, a prerequisite for active transport of transcripts to the yeast bud. We used a computer-aided and conceptual modeling approach that based on the hypothesis that an asymmetric bulge in the RNA conformation might be a prerequisite of SRS function. This approach led to the identification of a common three-dimensional arrangement of SRS functional features (conformational features and sequence motifs) in the *ASH1* localization elements E3, E2A and E1 (and with limitations also for E2B) in a blueprint, thus demonstrating the long-missing consensus motif of all four SRS elements. Unexpectedly, although this arrangement of features was conserved in its overall three-dimensional fold, , there were two variations with respect to the linear arrangement in the nucleotide sequences that we termed "configurations" ("config-I" in E3, "config-II" in E2A, E1 and E2B). The consensus model proved to be correct based on our consistent results from electric mobility shift assays, in which we analyzed the SRS function of RNA-constructs with re-designed feature arrangement. According to our model, this rearrangement of the features affected their linear order in the nucleotide sequences, but allowed the preservation of their overall three-dimensional arrangement, hence representing a switch between their original configuration and the respective counterpart configuration ("config-switch"). We were able to consistently demonstrate that the config-switch retained SRS function in all experiments we performed indicating that the consensus motif comprises the main requirements of SRS function. The fact that also size-reduced constructs proved to keep SRS function was a further hint that we had grasped the essential features of SRS function. It can be described in all *ASH1* SRS elements as follows: an asymmetric bulge, flanked by inverted repeats, forms a kinked, L-shaped RNA conformation. The larger side of the bulge contains the single cytosine motif C_1 inside of the first three positions. The C_2GA motif resides on the opposite strand of C_1, separated by an inverted repeat in downstream direction on this strand. The two configurations are marked by the two different 5' entry sites into this kinked conformation. These findings were further confirmed, when we compared previous thorough mutagenesis studies of the *ASH1* SRS elements and could assign these results well according to

their relevance in the SRS consensus model. Further, by studying the binding kinetics of $E3_{29nt}$wt in surface

plasmon resonance analysis, we could confirm the optimal distance between C_1 and C_2 with six nucleotides

(ca. 28 Å) as it was suggested before[10]. It is noteworthy that the existence of two configurations with double-

stranded folds that can be converted into each other by swapping their features in the nucleotide sequence

finds an analogy in studies on the beta-actin RNA zip code, where the order of its single-stranded 5' and 3'

elements is interchangeable within the target transcripts bound to ZBP1[11].

Due to its universal character to describe all four *ASH1* SRS elements, the exploitation of the SRS

consensus motif went beyond previously computational screens. Hence, we created a computational screen

that searched for both configurations, followed by a conformational check of the overall RNA fold

performed by *Kinefold*. For some of the further transcripts that had been found to localize at the bud tip of

yeast cells, sequence fragments had already been narrowed down with retained SRS function. In this set of

sequence fragments, we could classify SRS elements of *EAR1*, *IST2* and *WSC2* as further instances of the

SRS consensus motif. *EAR1* was first identified as localized in the bud in the study of [29] with its SRS

element characterized in further studies[10] and classified as a config-II SRS element in this work (*EAR1*-

SRS2, **Tab. 2**). For the minimal construct $EAR1_{31nt}$switch, using SRP analysis, we could confirm previous

observations that the nucleotide sequences of the inverted repeats did not contribute to SRS function[8,13]. In

further consistence with previous *in-vivo* studies[10], the loss of C_2 showed and individual substitutions of

each of the two potential C_1 positions (in C_1-C_2-distance of five or six nucleotides) affected SRS function.

Unexpectedly, a single deletion of C_{1601} or a simultaneous substitution of C_{1601} and C_{1602}, both restored SRS

function. When studying the potential di-nucleotide AA_L sequence motif by deletion experiments we could

not detect any impact on SRS function, even if the Lbulge region was almost completely deleted. So far,

we can neither explain the apparent discrepancy in the C_1 mutation effects in the *EAR1* constructs nor the

retained SRS function with deleted asymmetric bulge. It may reflect both complex sequence composition

and inherent redundancy in this particular SRS instance.

So far, we had observed only the occurrence of one config-I (E3) whereas we had identified six

instances of config-II (**Fig. 4C**). This dominance of config-II SRS elements continued when we applied the

computational screen to our complete sequence set and predicted 30 SRS instances in 17 transcripts, with

22 of them (73%) exhibiting config-II. So far, the reason for the unequal frequency of the two configurations

remains unclear. From this outcome, we used EMSA approaches to analyze two instances of each subset:

SRS instances with relatively well-known positions in their transcripts or with unknown SRS positions. For

the first subset, we could confirm SRS function in both SRS elements *EAR1*-SRS2 and *IST2*-SRS2[10]. For

the second subset, we could verify *KSS1*-SRS1 and *TCB3*-SRS1 *in-vitro*. Remarkably, when we switched

configuration in *EAR1*-SRS2, *IST2*-SRS2 and *KSS1*-SRS1 (config-II to config-I), we could demonstrate

again conserved SRS function for all switches, strongly supporting the SRS consensus model. Of note, as

a first successful validation *in-vivo*, we had been able to identify and confirm the nucleotide positions of a

SRS element in the *CLB2* transcript[34]: In the transcript, only a few but functionally crucial positions of the

SRS instance *CLB2*-SRS1 (config-II, 1111 nt - 1145 nt) were altered in non-canonical mutations to reveal

a significant loss-of-function in bud-localization. It is also interesting to note that the regulatory effect of

well-defined asymmetric bulges seems not to be limited to SRS function. As described in a study[35], the

structure of the pre-mRNA autoregulatory complex of the L30 ribosomal protein in yeast displays an

asymmetric bulge (length-ratio 5 nt : 2 nt) with a cytosine positioned in an extension of the Lbulge, and

separated from another cytosine (in a further bulge) by an IR in a distance of 6 nt. Remarkably, it is this

asymmetric bulge region that is part of a mutual induced fit, suggesting analogies with the observed induced

fit mechanisms in the crystal structure of E3 and She2p/3p.

Further aspects of SRS characteristics remain to be investigated. For example, a further extension

of the description of the SRS consensus motif is thinkable: It is reasonable to ask if the SRS function would

be conserved if Lbulge and Sbulge were joined together without IR2 as intervening feature. Although

experimental analysis of such a setting is beyond the scope of this study, we would like to put to discussion

such a theoretical construct, to possibly take a step forward to complete the picture (**Supplementary Fig.**

10A). The SRS element *WSC2*-N[36] that could be not classified, might be such an example (**Supplementary**

Figure 10B, left). Such an extended perspective might have further implications also for the understanding

of some of the SRP results of *EAR1* described before. Both elements, *EAR1*-SRS2 and *IST2*-SRS2 displayed

two CGA motifs with the following consequences: Whereas we considered *EAR1*-SRS2 as config-II with

C_2GA residing in the terminal bulge, this terminal bulge also may be considered as a joint of Lbulge and

Sbulge, hence offering a new perspective: the cytosine motif in this terminal bulge may be considered as

C_1 while its counterpart inside of the asymmetric bulge may be extended and labelled to C_2GA

(**Supplementary Figure 10B, right**). In such a view, the functionality of *EAR1*-SRS2 would exhibit

redundant features of two superimposed and oppositely rotated SRS elements in the bigger context of an

extended SRS consensus motif. This perspective may also explain partially the ambiguous results in the

SPR experiments of our study. Secondly, a further extension of the description of the SRS consensus might

be formed by considering a range in the restraints of the C_1-C_2 distance. In consistency with previous

studies[10] and with our SPR results we focused on the optimal parameters of six nucleotides in this study to

minimize false positive matches. However, we could not exclude a minimal functionality for instances with

(C_1-C_2 distance $\neq$ 6). An example might be a second, potential SRS instance that we identified in the

flanking regions of *WSC2*-SRS2 (*WSC2*-C) but with a C_1-C_2 distance of seven nucleotides (**Supplementary**

Fig 7C). In such a case, *WSC2*-SRS2 may compensate the suboptimal C_1-C_2 distance of its neighboring

SRS element by inducing SRS conformation in a cooperative manner when both SRS instances interact

with the binding interfaces on the opposite sites of the She2p tetramer[37].

As a second aspect, localization elements that mediate bud localization require the interaction with

the She2p/3p protein transport machinery, but further motifs may be required as binding platforms of

auxiliary factors. For example, for E2B, we speculated that its energetically optimal conformation had to

be converted to an alternative conformation to represent the SRS consensus motif. Even strong SRS

elements such as E2A[12] can be subject to major reorganizations, as shown by the PARS data which indicate

competing conformations for E2A[33] (**Supplementary Fig. 7**). Therefore, the She2p/3p interaction can be

seen as a central step in the zip-code function, but it might be framed by further rearrangements in the

chronological order. In such a scenario, transient binding of auxiliary cofactors could stabilize the

asymmetric bulge and increase SRS function. AA_L might be such a co-factor binding motif. Despite the

fact that AA_L was present in each case of our seven well-characterized, bud-localizing SRS elements,

neither the crystal structure[8] nor the results of SPR studies suggest involvement in She2p/3p binding

specificity. Future research will be required to clarify the possible function of AA_L *in-vivo*, also with regard

to transient interactions. In addition, for E3 and E2A, it was tempting to speculate if their extraordinary

sequence identity was a product of convergent evolution, due to further requirements in affecting zip-code

function. Maybe we have to face a trade-off between canonical SRS consensus motif and stability of the

individual RNA conformation, with the weakening of one aspect may be compensated by strengthening of

the other. This would complicate the identification of SRS instances, especially if auxiliary co-factors

would alter the probabilities of formation of SRS conformations *in-vivo*. Summarizing, all these aspects

stress two aspects: (i) The new SRS consensus motif delivers a conceptual framework of how to classify

further aspects of new SRS instances, and (ii) It remains necessary to combine experimental screenings

(identification of bud-localizing transcripts, final validation of instances) and computational screenings

(prediction of positional information in transcripts to narrow down experimental analysis and suggest

crucial positions in mutations studies). Consequently, the *SRS instance collection* contains only such SRS

instances where bud localization had been confirmed already, at least at the transcript level.

In conclusion, the SRS consensus motif enabled the description of the four SRS elements of *ASH1*

in one model, hence providing the long-missing consensus. It further unified the described functional

features of the known SRS elements in *IST2*, *EAR1* and *WSC2* and enabled the identification of new SRS

elements in further bud-localizing transcripts. The consensus model proved to be consistent with previous

mutagenesis studies of these elements and we could demonstrate that a switch between the configurations

retained function in SRS constructs of several transcripts.

Materials and Methods

Overview on the theoretical data flow to derive the SRS consensus model

The theoretical part of deriving the SRS consensus model comprised the following steps

(**Supplementary Fig. 2**): (1) Hypothesis (asymmetric bulge as hallmark of SRS) by analysis of crystal

structure of *ASH1*-E3 (2) Selection of an appropriate conformational prediction tool (3) Conformational

prediction of instances and inspection to derive a consensus (**Fig. 2**) (4) Cross-check of the consensus with

published functional data (5) Creation of a computational screening pipeline (6) Check for further data

sources to filter results (**Supplementary Fig. 7**) (7) Screening for instances in selected sequence sets (8)

Inspection of instances for further patterns

Identification of an appropriate conformational prediction tool

The inspection of the crystal structure of *ASH1*-E3 suggested an asymmetric bulge as a pre-requisite

of kinked RNA conformation. The occurrence of such asymmetric bulges may be favorized in mRNPs by

the energy release due to molecular interaction. However, its occurrence is less likely in an individual RNA

and is less likely to be predicted as plausible conformation by computational prediction tools that commonly

deal with individual RNA sequences. In a survey, when investigating which of common tools[32,38–40] was

most likely to predict the asymmetric bulge of E3, we found Kinefold[32] as the one that best tolerated

asymmetric bulges (**Supplementary Fig. 3**). It predicted the asymmetric bulge consisting of a 5 nt Lbulge

and a 2 nt Sbulge, which closely approximates the length ratio of the asymmetric bulge observed in the E3-

She2p/3p crystal structure[8] that displayed a length ratio of 6 nt : 3 nt (considering only Watson-Crick base

pairs) or 4 nt : 1 nt (considering also atypical base pairings as observed in the crystal structure), respectively

(**Fig. 1A**).

Cross-check of the consensus with published functional data

We used published data of functional studies that involved both, mutagenesis and deletion of RNA

regions to cross-check for consistency with the SRS consensus model[10,13,25,30]. We characterized such

mutations in *ASH1* SRS elements that could clearly be assigned to destroy or to keep SRS function

according to the consensus model. All these constructs were in accordance with the consensus model (data

not shown).

Development of a computational screening approach

For the purpose of identifying further instances of the SRS consensus motif, we transformed the

consensus features into "patterns" to screen for "matches" in RNA sequences. We created a computational

screening pipeline consisting of in-house computational scripts and an external conformational prediction

algorithm. For the screening, three aspects were required: (1) Screening the sequence for conformational

aspects. For each RNA, the sequence was screened for two nested pairs of inverted repeats (each with a

minimal length of four nucleotides) that framed an asymmetric bulge region. We had to consider both

configurations in different screening patterns. The obtained constituted potential SRS instances. (2)

Screening the sequence for functional motifs. For each SRS instance, the Lbulge region was screened for

C_1, the opposite strand for C_2GA (positioned beyond IR1), with a C_1-C_2 distance of 6 nt between. The

screening strategy based on the approach of modular feature screening of mosaic-like patterns[41]. This

modular approach allowed to introduce further features to be screened (e.g., AA_L). It also tolerated length

variations between the distinct (sequence and conformational) features (e.g., one version tolerated C_1

positioned in the last nucleotide of IR1). As a consequence, permutated sets of varying patterns could be

summarized into two screening patterns (one for each configuration). (3) Validation by conformational

prediction. For each SRS instance, it was calculated if it would express a stable conformational fold that

fitted to the conformational arrangement of the SRS consensus. This analysis was performed using *Kinefold*

as energy-minimization approach that had proven best for large asymmetric bulges. On each site , the SRS

stretches were extended by 10 nt of the flanking sequences of their RNA sequences, to tolerate some degree

of freedom during calculation. In a series of tests using known SRS elements, a flanking region extension

by 10 nt had proven to be best in a range of 0 nt to 50 nt (data not shown). Of note, natural limitations of

common RNA conformational predictions occur due to the fact that such calculations are limited to predict

the conformation of distinct RNA stretches but less reliable regarding statements of the complete transcript

and second, they would ignore any energetic benefits by binding of molecular interactors (e.g., She2p/3p

proteins). As a consequence, we analyzed the 25th best conformational predictions of *Kinefold* (maximum

value = 25), to consider slight structural dynamics in each RNA molecule.

Check for further data sources to filter for the consensus

For the purpose to possibly extend the validation step of new SRS instances by conformational

predictions. we intended to cross-check the predicted conformations with experimental data to get deeper

insights on actual RNA conformations *in-vivo*. For our purposes, we tested the remarkable genome-wide

study based on PARS (Parallel Analysis of RNA Structure)[33]. For each nucleotide in transcripts, PARS-

scores described the percentage in occurring in single-strand ("S1-reads" of RNase S1) or double-strand

("V1-reads" of RNase V1) regions of the RNA conformations. We extracted the S1- and V1-data and tested

the strong localization element E3[13] as positive control. As a result the PARS data was well suited to

describe the SRS conformation as observed in the E3 crystal structure[8] (**Supplemental Fig. 7**). However,

for further controls, the data comparison of paired/ unpaired occurrence of nucleotides between PARS data

and SRS consensus model failed (**Supplemental Fig. 7**). Commonly, RNA molecules display not one

distinct conformation in-vivo but are subject to conformational dynamics resulting in collections of

competing conformations. PARS data per nucleotide averaged the binding state (paired/unpaired) over all

competing conformations. It seemed that even the two SRS elements in *ASH1*-E1 and *ASH1*-E2A (proven

to be efficient localization elements[13]) were not sufficiently stable in the consensus conformations

according to their PARS data to distinguish them from alternative conformations

Further bioinformatics resources

For the investigation of molecular structures, we used Pymol[42] as 3D- viewer. Initial protein

characterization of She2p and She3p was performed using the ELM server[43]. RNA sequence alignments

were performed using Jalview[44]. RNA sequences were extracted of the Saccharomyces Genome Database[45].

For the actual screening, the mRNA sequences of *ASH1*, *IST2*, *EAR1* (aka *YMR171C*), *WSC2* and 20 further

mRNA sequences known to She2p/3p interact[10,13,25,29,30,36] were chosen. As sequence data, we used the

CDRs, extended by 200 nt downstream by default, due to the fact that for several mRNAs, the actual 3'

UTR lengths were still unknown. Functional annotation of Supplementary Table S1 followed [46]. Graphical

visualization of RNA conformations was performed using VARNA[47].

Protein expression

Expression in *E. coli* and purification of She2p, She3p, their fusion protein, and variants thereof

were performed as described in [8,23]. Purity and solubility were confirmed by size-exclusion chromatography

and SDS page.

RNA preparation and isotope labeling

For small scale *in vitro* transcription the MEGAshortscript T7 kit (life technologies) was used

according to the manufacturer's instructions. Synthetic DNA oligonucleotides containing a T7 promoter

and the respective sequence were used as a template. After *in vitro* transcription, the template was digested

by DNAse and the RNA was extracted using phenol:chloroform:isoamyl alcohol (25:24:1). Homogeneity

of RNA was assessed by native and denaturing TBE agarose gel electrophoresis.

Electrophoretic mobility shift assay

In a total volume of 20 µL 5 nM radioactively labeled RNA, 100 µg/mL yeast tRNA competitor,

and protein at a given concentration were mixed in buffer (20 mM Hepes, pH 7.8, 200 mM NaCl, 2 mM

$MgCl_2$, 2 mM DTT) with 4 % (v/v) glycerol. After 25 min at room temperature, the samples were resolved

by native TBE PAGE (6 % polyacrylamide) and gels incubated for 15 min in fixing solution (10 % (v/v)

acetic acid, 30 % (v/v) methanol) and subsequently vacuum dried. Gels were analyzed with radiograph

films or by a using Phosphorimager (FujifilmFLA5100).

Surface plasmon resonance analysis (SPR)

SPR experiments were performed with a BIACORE 3000 system (GE Healthcare). For protein-RNA interactions, biotinylated RNA was streptavidin-captured on an SA-Chip (GE Healthcare) surface according to manufacturers instructions. Measurements were performed at a flow rate of 30 µL/min in standard buffer. Data were analyzed using the BIAevaluation software (GE Healthcare). Sample sensograms were referenced against the signal in a ligand-free reference channel, and averaged values at the equilibrium calculated. Steady state binding curves were obtained by plotting the response units at equilibrium against protein concentration with the Origin Pro 9.0 (OriginLab) software.

Computational quality check of the folds of the experimental constructs

Part of the design of experimental constructs (minimal, config-switch) for the *in-vitro* and *in-vivo* approaches was a quality check with *Kinefold*: The conformations of their sequences were predicted to ensure that their expected conformation would be in agreement with the SRS consensus model

Acknowledgements

We would like to thank Herve Isambert for helpful discussions regarding Kinefold and Toby Gibson for further support. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) to R.-P.J. (DFG-FOR2333, Project Number 270067186), to M.F. (CEPLAS EXC 1028, CRC1208, DFG-FOR2333, DFG Fe448/8, DFG Fe448/9), to D.N. (DFG-FOR2333, Project Number 270067186), and to Z.K. (DFG-FOR2333, Project Number 270067186).

Declaration of Interest

The authors declare no potential conflicts of interest.

References

(1) Eliscovich, C.; Singer, R. H. RNP Transport in Cell Biology: The Long and Winding Road. *Curr. Opin. Cell Biol.* **2017**, *45*, 38–46. https://doi.org/10.1016/j.ceb.2017.02.008.

(2) Bashirullah, A.; Cooperstock, R. L.; Lipshitz, H. D. RNA Localization in Development. *Annu. Rev. Biochem.* **1998**, *67*, 335–394. https://doi.org/10.1146/annurev.biochem.67.1.335.

(3) Johnstone, O.; Lasko, P. Translational Regulation and RNA Localization in Drosophila Oocytes and Embryos. *Annu. Rev. Genet.* **2001**, *35*, 365–406. https://doi.org/10.1146/annurev.genet.35.102401.090756.

(4) Kloc, M.; Bilinski, S.; Chan, A. P.; Allen, L. H.; Zearfoss, N. R.; Etkin, L. D. RNA Localization and Germ Cell Determination in Xenopus. *Int. Rev. Cytol.* **2001**, *203*, 63–91. https://doi.org/10.1016/s0074-7696(01)03004-2.

(5) Palacios, I. M.; St Johnston, D. Getting the Message across: The Intracellular Localization of mRNAs in Higher Eukaryotes. *Annu. Rev. Cell Dev. Biol.* **2001**, *17*, 569–614. https://doi.org/10.1146/annurev.cellbio.17.1.569.

(6) Dahm, R.; Kiebler, M.; Macchi, P. RNA Localisation in the Nervous System. *Semin. Cell Dev. Biol.* **2007**, *18* (2), 216–223. https://doi.org/10.1016/j.semcdb.2007.01.009.

(7) Abouward, R.; Schiavo, G. Walking the Line: Mechanisms Underlying Directional mRNA Transport and Localisation in Neurons and Beyond. *Cell. Mol. Life Sci. CMLS* **2021**, *78* (6), 2665–2681. https://doi.org/10.1007/s00018-020-03724-3.

(8) Edelmann, F. T.; Schlundt, A.; Heym, R. G.; Jenner, A.; Niedner-Boblenz, A.; Syed, M. I.; Paillart, J.-C.; Stehle, R.; Janowski, R.; Sattler, M.; Jansen, R.-P.; Niessing, D. Molecular Architecture and Dynamics of ASH1 mRNA Recognition by Its mRNA-Transport Complex. *Nat. Publ. Group* **2017**, *24* (2), 152–161.

(9) Niedner, A.; Edelmann, F. T.; Niessing, D. Of Social Molecules: The Interactive Assembly of ASH1 mRNA-Transport Complexes in Yeast. *RNA Biol.* **2014**, *11* (8), 998–1009. https://doi.org/10.4161/rna.29946.

565 (10) Olivier, C.; Poirier, G.; Gendron, P.; Boisgontier, A.; Major, F.; Chartrand, P. Identification of a

566    Conserved RNA Motif Essential for She2p Recognition and mRNA Localization to the Yeast Bud.

567    *Mol. Cell. Biol.* **2005**, *25* (11), 4752–4766. https://doi.org/10.1128/MCB.25.11.4752-4766.2005.

568 (11) Patel, V. L.; Mitra, S.; Harris, R.; Buxbaum, A. R.; Lionnet, T.; Brenowitz, M.; Girvin, M.; Levy,

569    M.; Almo, S. C.; Singer, R. H.; Chao, J. A. Spatial Arrangement of an RNA Zipcode Identifies

570    mRNAs under Post-Transcriptional Control. *Genes Dev.* **2012**, *26* (1), 43–53.

571    https://doi.org/10.1101/gad.177428.111.

572 (12) Chartrand, P.; Singer, R. H.; Long, R. M. RNP Localization and Transport in Yeast. *Annu. Rev. Cell*

573    *Dev. Biol.* **2001**, *17*, 297–310. https://doi.org/10.1146/annurev.cellbio.17.1.297.

574 (13) Chartrand, P.; Meng, X. H.; Singer, R. H.; Long, R. M. Structural Elements Required for the

575    Localization of ASH1 mRNA and of a Green Fluorescent Protein Reporter Particle in Vivo. *Curr.*

576    *Biol. CB* **1999**, *9* (6), 333–336. https://doi.org/10.1016/s0960-9822(99)80144-4.

577 (14) Gonzalez, I.; Buonomo, S. B. C.; Nasmyth, K.; von Ahsen, U. ASH1 mRNA Localization in Yeast

578    Involves Multiple Secondary Structural Elementsand Ash1 Protein Translation. *Curr. Biol. CB*

579    **1999**, *9* (6), 337–340.

580 (15) Kanke, M.; Jambor, H.; Reich, J.; Marches, B.; Gstir, R.; Ryu, Y. H.; Ephrussi, A.; Macdonald, P.

581    M. Oskar RNA Plays Multiple Noncoding Roles to Support Oogenesis and Maintain Integrity of the

582    Germline/Soma Distinction. *RNA N. Y. N* **2015**, *21* (6), 1096–1109.

583    https://doi.org/10.1261/rna.048298.114.

584 (16) Mori, Y.; Imaizumi, K.; Katayama, T.; Yoneda, T.; Tohyama, M. Two Cis-Acting Elements in the

585    3' Untranslated Region of Alpha-CaMKII Regulate Its Dendritic Targeting. *Nat. Neurosci.* **2000**, *3*

586    (11), 1079–1084. https://doi.org/10.1038/80591.

587 (17) Serano, J.; Rubin, G. M. The Drosophila Synaptotagmin-like Protein Bitesize Is Required for

588    Growth and Has mRNA Localization Sequences within Its Open Reading Frame. *Proc. Natl. Acad.*

589    *Sci. U. S. A.* **2003**, *100* (23), 13368–13373. https://doi.org/10.1073/pnas.1835727100.

590   (18)   Thio, G. L.; Ray, R. P.; Barcelo, G.; Schüpbach, T. Localization of Gurken RNA in Drosophila

591   Oogenesis Requires Elements in the 5' and 3' Regions of the Transcript. *Dev. Biol.* **2000**, *221* (2),

592   435–446. https://doi.org/10.1006/dbio.2000.9690.

593   (19)   Long, R. M.; Gu, W.; Lorimer, E.; Singer, R. H.; Chartrand, P. She2p Is a Novel RNA-Binding

594   Protein That Recruits the Myo4p-She3p Complex to ASH1 mRNA. *EMBO J.* **2000**, *19* (23), 6592–

595   6601. https://doi.org/10.1093/emboj/19.23.6592.

596   (20)   Münchow, S.; Sauter, C.; Jansen, R. P. Association of the Class V Myosin Myo4p with a Localised

597   Messenger RNA in Budding Yeast Depends on She Proteins. *J. Cell Sci.* **1999**, *112 (Pt 10)*, 1511–

598   1518.

599   (21)   Bertrand, E.; Chartrand, P.; Schaefer, M.; Shenoy, S. M.; Singer, R. H.; Long, R. M. Localization of

600   ASH1 mRNA Particles in Living Yeast. *Mol. Cell* **1998**, *2* (4), 437–445.

601   https://doi.org/10.1016/s1097-2765(00)80143-4.

602   (22)   Böhl, F.; Kruse, C.; Frank, A.; Ferring, D.; Jansen, R. P. She2p, a Novel RNA-Binding Protein

603   Tethers ASH1 mRNA to the Myo4p Myosin Motor via She3p. *EMBO J.* **2000**, *19* (20), 5514–5524.

604   https://doi.org/10.1093/emboj/19.20.5514.

605   (23)   Heym, R. G.; Zimmermann, D.; Edelmann, F. T.; Israel, L.; Ökten, Z.; Kovar, D. R.; Niessing, D. In

606   Vitro Reconstitution of an mRNA-Transport Complex Reveals Mechanisms of Assembly and Motor

607   Activation. *J. Cell Biol.* **2013**, *203* (6), 971–984. https://doi.org/10.1083/jcb.201302095.

608   (24)   Takizawa, P. A.; Vale, R. D. The Myosin Motor, Myo4p, Binds Ash1 mRNA via the Adapter

609   Protein, She3p. *Proc. Natl. Acad. Sci.* **2000**, *97* (10), 5273–5278.

610   (25)   Chartrand, P.; Meng, X. H.; Huttelmaier, S.; Donato, D.; Singer, R. H. Asymmetric Sorting of

611   Ash1p in Yeast Results from Inhibition of Translation by Localization Elements in the mRNA. *Mol.*

612   *Cell* **2002**, *10* (6), 1319–1330. https://doi.org/10.1016/s1097-2765(02)00694-9.

613   (26)   Long, R. M.; Singer, R. H.; Meng, X.; Gonzalez, I.; Nasmyth, K.; Jansen, R. P. Mating Type

614   Switching in Yeast Controlled by Asymmetric Localization of ASH1 mRNA. *Science* **1997**, *277*

615   (5324), 383–387. https://doi.org/10.1126/science.277.5324.383.

(27) Sil, A.; Herskowitz, I. Identification of Asymmetrically Localized Determinant, Ash1p, Required

for Lineage-Specific Transcription of the Yeast HO Gene. *Cell* **1996**, *84* (5), 711–722.

https://doi.org/10.1016/s0092-8674(00)81049-1.

(28) Takizawa, P. A.; Sil, A.; Swedlow, J. R.; Herskowitz, I.; Vale, R. D. Actin-Dependent Localization

of an RNA Encoding a Cell-Fate Determinant in Yeast. *Nature* **1997**, *389* (6646), 90–93.

https://doi.org/10.1038/38015.

(29) Shepard, K. A.; Gerber, A. P.; Jambhekar, A.; Takizawa, P. A.; Brown, P. O.; Herschlag, D.;

DeRisi, J. L.; Vale, R. D. Widespread Cytoplasmic mRNA Transport in Yeast: Identification of 22

Bud-Localized Transcripts Using DNA Microarray Analysis. *Proc. Natl. Acad. Sci. U. S. A.* **2003**,

100 (20), 11429–11434. https://doi.org/10.1073/pnas.2033246100.

(30) Jambhekar, A.; McDermott, K.; Sorber, K.; Shepard, K. A.; Vale, R. D.; Takizawa, P. A.; DeRisi, J.

L. Unbiased Selection of Localization Elements Reveals Cis-Acting Determinants of mRNA Bud

Localization in Saccharomyces Cerevisiae. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102* (50), 18005–

18010. https://doi.org/10.1073/pnas.0509229102.

(31) Singh, N.; Blobel, G.; Shi, H. Hooking She3p onto She2p for Myosin-Mediated Cytoplasmic

mRNA Transport. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112* (1), 142–147.

https://doi.org/10.1073/pnas.1423194112.

(32) Xayaphoummine, A.; Bucher, T.; Isambert, H. Kinefold Web Server for RNA/DNA Folding Path

and Structure Prediction Including Pseudoknots and Knots. *Nucleic Acids Res.* **2005**, *33* (Web

Server issue), W605-610. https://doi.org/10.1093/nar/gki447.

(33) Kertesz, M.; Wan, Y.; Mazor, E.; Rinn, J. L.; Nutter, R. C.; Chang, H. Y.; Segal, E. Genome-Wide

Measurement of RNA Secondary Structure in Yeast. *Nature* **2010**, *467* (7311), 103–107.

https://doi.org/10.1038/nature09322.

(34) Tutucci, E.; Maekiniemi, A.; Snoep, J. L.; Seiler, M.; van Rossum, K.; van Niekerk, D. D.; Savakis,

P.; Zarnack, K.; Singer, R. H. *Cyclin CLB2 mRNA Localization Determines Efficient Protein*

Synthesis to Orchestrate Bud Growth and Cell Cycle Progression; preprint; Cell Biology, 2022.

https://doi.org/10.1101/2022.03.01.481833.

(35) Frankel, A. D. If the Loop Fits. *Nat. Struct. Biol.* **1999**, *6* (12), 1081–1083.

https://doi.org/10.1038/70002.

(36) Takizawa, P. A.; DeRisi, J. L.; Wilhelm, J. E.; Vale, R. D. Plasma Membrane Compartmentalization

in Yeast by Messenger RNA Transport and a Septin Diffusion Barrier. *Science* **2000**, *290* (5490),

341–344. https://doi.org/10.1126/science.290.5490.341.

(37) Müller, M.; Richter, K.; Heuck, A.; Kremmer, E.; Buchner, J.; Jansen, R.-P.; Niessing, D.

Formation of She2p Tetramers Is Required for mRNA Binding, mRNP Assembly, and Localization.

RNA N. Y. N **2009**, *15* (11), 2002–2012.

(38) Lorenz, R.; Bernhart, S. H.; Höner Zu Siederdissen, C.; Tafer, H.; Flamm, C.; Stadler, P. F.;

Hofacker, I. L. ViennaRNA Package 2.0. *Algorithms Mol. Biol. AMB* **2011**, *6*, 26.

https://doi.org/10.1186/1748-7188-6-26.

(39) Reuter, J. S.; Mathews, D. H. RNAstructure: Software for RNA Secondary Structure Prediction and

Analysis. *BMC Bioinformatics* **2010**, *11*, 129. https://doi.org/10.1186/1471-2105-11-129.

(40) Zuker, M. Mfold Web Server for Nucleic Acid Folding and Hybridization Prediction. *Nucleic Acids*

Res. **2003**, *31* (13), 3406–3415. https://doi.org/10.1093/nar/gkg595.

(41) Seiler, M.; Mehrle, A.; Poustka, A.; Wiemann, S. The 3of5 Web Application for Complex and

Comprehensive Pattern Matching in Protein Sequences. *BMC Bioinformatics* **2006**, *7*, 144.

https://doi.org/10.1186/1471-2105-7-144.

(42) Schrödinger, L.; DeLano, W. PyMOL. **2020**.

(43) Kumar, M.; Michael, S.; Alvarado-Valverde, J.; Mészáros, B.; Sámano-Sánchez, H.; Zeke, A.;

Dobson, L.; Lazar, T.; Örd, M.; Nagpal, A.; Farahi, N.; Käser, M.; Kraleti, R.; Davey, N. E.;

Pancsa, R.; Chemes, L. B.; Gibson, T. J. The Eukaryotic Linear Motif Resource: 2022 Release.

Nucleic Acids Res. **2022**, *50* (D1), D497–D508. https://doi.org/10.1093/nar/gkab975.

(44) Waterhouse, A. M.; Procter, J. B.; Martin, D. M. A.; Clamp, M.; Barton, G. J. Jalview Version 2--a

Multiple Sequence Alignment Editor and Analysis Workbench. *Bioinforma. Oxf. Engl.* **2009**, *25* (9),

1189–1191. https://doi.org/10.1093/bioinformatics/btp033.

(45) Engel, S. R.; Dietrich, F. S.; Fisk, D. G.; Binkley, G.; Balakrishnan, R.; Costanzo, M. C.; Dwight, S.

S.; Hitz, B. C.; Karra, K.; Nash, R. S.; Weng, S.; Wong, E. D.; Lloyd, P.; Skrzypek, M. S.;

Miyasato, S. R.; Simison, M.; Cherry, J. M. The Reference Genome Sequence of Saccharomyces

Cerevisiae: Then and Now. *G3 Bethesda Md* **2014**, *4* (3), 389–398.

https://doi.org/10.1534/g3.113.008995.

(46) Engel, S. R.; Wong, E. D.; Nash, R. S.; Aleksander, S.; Alexander, M.; Douglass, E.; Karra, K.;

Miyasato, S. R.; Simison, M.; Skrzypek, M. S.; Weng, S.; Cherry, J. M. New Data and

Collaborations at the Saccharomyces Genome Database: Updated Reference Genome, Alleles, and

the Alliance of Genome Resources. *Genetics* **2022**, *220* (4), iyab224.

https://doi.org/10.1093/genetics/iyab224.

(47) Darty, K.; Denise, A.; Ponty, Y. VARNA: Interactive Drawing and Editing of the RNA Secondary

Structure. *Bioinforma. Oxf. Engl.* **2009**, *25* (15), 1974–1975.

https://doi.org/10.1093/bioinformatics/btp250.

 Figures Legends

Figure 1. The SRS consensus model describes all four *ASH1* SRS elements. A Left, conformational
prediction of E3$_{1771-1821}$ by Kinefold displays the asymmetric bulge as a strand of 5 nt ('Lbulge') juxtaposed
with a shorter strand ('Sbulge') both indicated in red, previously known functional motifs in dark green.
Right, the conformation of E3 referred to its crystal structure bound to She2p/3p (**PDB: 5NEW**) displayed
a rearranged the asymmetric bulge with atypical base-pairing, with a length ratio of 6 nt : 3 nt (with atypical
base-pairs dashed in blue, but ignored in the count), and with further positions of sequence motifs adapted
to analogous functional residues as studied in E2A, E2B and E3 (indicated in light green). **B** Left, Kinefold
prediction and 180° rotation of E2A$_{1109-1185}$ revealed an asymmetric bulge with the same 5 nt : 2 nt ratio as
in E3. Right, the high degree of sequence similarity (**Supplementary Fig. 1**) enabled a modeling in tight
accordance to the E3 conformation of the crystal structure that even comprised the uncommon base-pairing
C$_{1122}$-C$_{1174}$ and the A$_{1123}$-A$_{1169}$ base-stacking (dashed in blue). **C** Left, the optimal conformation of E2B$_{1263-}$
$_{1320}$ after 180 ° rotation showed internal repeats (left, boxed in orange) that could be re-modeled according
to Kinefold predictions (right) but with increased energy content (-10.3 kcal/mol vs.-7.4 kcal/mol). **D** SRS
element E1$_{623-701}$ after Kinefold prediction and 180° rotation (left) and with adapted conformation (right).
E Scheme of the features of all four modeled SRS elements, projected on E3$_{1771-1821}$. An asymmetric bulge
(red) is framed by two inverted repeats (yellow), with sequence motifs indicated in green. **F** Schematic
presentation of the SRS consensus motif with its features displayed in both configurations (IR1 boxed in
grey, IR2 boxed in blue, Lbulge and Sbulge in red, sequence motifs indicated by green labeling). Regarding
the two configurations, their different order of features (left) results in a overall spatial arrangement that is
conserved but with varying positioning of 5' start and 3' end in its conformations (middle). Right,
conformational view on the unified model of the SRS consensus motif.

**Figure 2. The modeling of the SRS consensus involves choice of appropriate conformation prediction
tool, rotation and mutual feature alignment of the four SRS elements in *ASH1*.** The overview
demonstrates the effects of the three steps of modeling on the extended sizes of the *ASH1* SRS elements as
they were characterized before. Whereas the initially known conformations displayed a huge degree of
heterogeneity (**A, D, H, L**), the conformation prediction tool Kinefold tolerated asymmetric bulges in each
SRS element (**B, E, I, M**). For E3, using the published crystal structure as a blue print, its conformational
prediction (**B**) was adapted, with altered conformation of the asymmetric bulge (resulting in atypical base
pairings, marked in red, **C**). Its final asymmetric feature is indicated by a red triangle, with the numbers of
nucleotides displayed for both, Lbulge and Sbulge. By later analyses, we could assign one of its functional
cytosines as being analogous to already known CGA sequence motifs with other SRS elements (indicated

in red). For E2A, E1 and E2B, it was the rotation of the initial predictions that harmonized the respective

conformations with E3 (**F, J, N**). For E2A, due to its remarkable sequence identity with E3, we could also

introduce the same atypical base pairings (marked in red, **G**). As mentioned before, E2B represented an

outlier: it was a alternative conformation prediction that could be harmonized with E3, suggesting the need

of a further intramolecular rearrangement to align the inverted boxes (marked in yellow, **N, O**).

Figure 3. The configuration switch in E3 displays conserved function and surface plasmon resonance

analysis confirms the optimal C_1-C_2 distance of 6 nt in minimal constructs of E3. A Representative

results of electrophoretic mobility shift assays (EMSA) of triplicates to analyze complex formation of wild-

type and variant constructs of radioactive ASH1 $E3_{51nt}$ RNA with the She2p-She3p fusion protein ($She2p_{6-}$

$_{246}$-$(GGSGG)_2$-$She3p_{331-405}$) in increasing concentration (0.02 - 5 µM): Wildtype $E3_{1771-1821}$ (left), construct

with switch of $E3_{1771-1821}$ from config-I to config-II (middle) and control M6 mutational variant of $E3_{1771-}$

$_{1821}$ with decreased SRS function[8]. **B** Results of surface plasmon resonance analysis (SRP), performed in

triplicates to analyze the minimal E3 constructs and alterations in their C_1-C_2 distance. The minimal

construct $E3_{28nt}wt$ was re-designed to create constructs with differing C1-C2 distances (5 and 7 nt) and was

incubated with She2p-She3p fusion protein (125 nM).

Figure 4. Computational screen for the SRS consensus motif across 24 known She2p/3p interacting

transcripts results in 30 predicted instances in 17 transcripts. A Computational screen utilized both,

feature screening of the SRS consensus motif and conformational evaluation (**Materials and Methods**)

and was applied on 24 sequences known for She2p/3p interaction but with varying experimental confidence

(**Supplementary Tab. 1**). It identified 30 SRS predictions in 17 sequences, here visualized as bar charts,

with their coding sequences (rectangles) and 3' UTR extensions (200 nt, black lines). SRS predictions were

indicated by their configurations (config-I: red, config-II: blue). **B** For each configuration, the occurrence

of the AA_L dinucleotide was specified. **C** Schematic overview on the high-confidential set of SRS elements.

Similar to the SRS elements in *ASH1*, predictions in *IST2*, *EAR1* and *WSC2* matched published, well-

characterized SRS elements (nomenclature, see **Tab. 2**) and could be arranged accordant to the model.

Color scheme is adopted of **Figure 1**, with visualization of AA_L dinucleotides, boxed in orange).

Uncommon base pairings had detected in the E3-She2p/She3p crystal structure were consistently mapped

to E2A that displayed high sequence similarity with E3 after rotation (blue-dashed). Of note, similar

uncommon base-pairings may also arise in E1 and *IST2*-SRS-2.

Figure 5. The configuration switch in the minimal SRS constructs of *EAR*-SRS2, *IST*-SRS2, *KSS1*-
SRS1 and *TCB3*-SRS1 displays conserved SRS function in electrophoretic mobility shift assays.
Representative results of electrophoretic mobility shift assays (EMSA) to analyze interaction of radioactive
wild-type and designed RNA constructs with the She2p-She3p fusion protein (She2p$_{6-246}$-(GGSGG)$_2$-
She3p$_{331-405}$) in increasing concentration (0.02 - 5 μM). All minimal constructs were designed switched in
configuration (from config-II to config-I). Their final loop sequence GAAA referred to the artificial loop
sequence of the E3 constructs in [8] that was shown not to modify SRS function. **A** EMSA experiments of
well-known SRS elements. Left, *EAR1*-SRS2$_{1572-1621}$ (config-II) was studied as wildtype construct *EAR1*-
SRS2$_{50nt}$wt and as config-switched, minimal construct *EAR1*-SRS2$_{26nt}$-switch (config-I). Right, *IST2*-
SRS2$_{2716-2777}$ (config-II) was studied as wildtype construct *IST2*-SRS2$_{62nt}$wt and as config-switched,
minimal construct *IST2*-SRS2$_{29nt}$-switch (config-I). **B** EMSA experiments of new, predicted SRS elements.
Left, *KSS1*-SRS1 (config-II) was studied as wildtype construct *KSS1*-SRS1$_{72nt}$wt and as config-switched,
minimal construct *KSS1*-SRS1$_{29nt}$-switch (config-I). Right, *TCB3*-SRS1 (config-II) was studied as config-
switched, minimal construct *TCB3*-SRS1$_{38nt}$-switch (config-I).

Figure 6. *EAR1*-SRS2$_{28nt}$switch is partly sensitive to alterations of its functional motifs but tolerates
alterations in its conformational features and in the AA$_L$ dinucleotide of the Lbulge as showed by
surface plasmon resonance analysis. A Schemes of the minimal constructs of *EAR1*-SRS2$_{28nt}$switch that
were re-designed to create constructs with altered conformational features (indicated in red in
conformations). The final loop sequence GAAA referred to the artificial loop sequence of the E3 constructs
in [8] that was shown not to alter SRS function **B** Substitution scheme to analyze the relevance of cytosines
in the sequence motifs of the SRS consensus model. Alterations are marked in red in substitution tables that
refer to both, sequence positions and terminologies of resulting constructs. Since two cytosines (C$_{1601}$ and
C$_{1602}$) precluded an unambiguous assignment of the C$_1$ position, both had to be considered. A third cytosine
C$_{1600}$, marked by a star, was not considered in alterations due to its uncommonly short distance to C$_2$. **C**
Substitution scheme to analyze the relevance of the AA$_L$ dinucleotide in the Lbulge. For each alterations,
construct name (blue), conserved sequence in the Lbulge (black) and alterations (red) are displayed. **D**
Results of surface plasmon resonance analysis (SRP), performed in triplicates to analyze the minimal
EAR1-SRS2 constructs and alterations in their C$_1$-C$_2$ distance. E3 RNA was incubated with She2p-She3p
fusion protein (125 nM).

781
782

Figure 1

Consensus modeling by mutual feature alignment using Kinefold

A

<u>ASH1–E3</u>

B

Figure 4

A

B

C

Figure 5

A

B

Figure 6

EAR1-SRS2 constructs for SRP analysis

A

EAR1_{31nt}sw — FlexA –> stem — plus 1bp stem

CG inversion — inversion 3bp — ΔC1ΔflexA

B

C

D

SRP: EAR1-SRS2 constructs vs. 125nM She2p/3p

788
789
790
791

INFO-BOX: Summary of the SRS consensus motif

1. Two **inverted repeats** (IR-1, IR-2) flank an **asymmetric bulge** (consisting of **Lbulge** and **Sbulge**)

2. The conserved **singular cytosine C_1** resides in one of the first three positions of the Lbulge

3. The conserved **C_2GA triplet** resides in a further bulge, at the opposite strand of C_1, separated by an inverted repeat in downstream direction on this strand in a **C_1-C_2 distance of six positions**

4. The two **configurations** are marked by the two different 5' entry sites into the kinked SRS element conformation.

793
794
795   **TABLE 1 Overview on the four localization elements of *ASH1*.** Positions
796   and configuration classification are listed according to the SRS consensus motif
798

SRS name	SRS config I	SRS config II	SRS AA motif in Lbulge	SRS start	SRS end
E2		II	yes	631	686
E2A		II	yes	1118	1178
E2B		II	yes	1275	1312
E3	I		yes	1774	1817

799
800
801
802

803
804 **TABLE 2 Overview on the *SRS instance collection*.** The columns list info at transcripts
805 and SRS instance level; SRS instance are classified according to the SRS consensus motif
806 (including positions and configuration); the last two columns contain information to the
807 identification of SRS elements in fragment size or as a detailed description in previous
808
809 publications and this study

mRNA #	mRNA name	mRNA gene ID	SRS instance also known as	SRS instance collection name (*)	SRS start	SRS end	SRS config I	SRS config II	SRS as terminal element (***)	SRS with AAt motif	SRS fragment reference	SRS element reference
1	ASH1	YKL185W	E1	ASH1-E1	631	686		II		AA	several	several
			E2A	ASH1-E2A	1118	1178		II		AA	several	several
			E2B	ASH1-E2B	1275	1312		II	terminal	AA	several	several
			E3, Umin	ASH1-E3	1774	1817	I			AA	several	several
2	CLB2	YPR119W		CLB2-SRS1	1111	1145		II		AA		Tutucci et al
3	CPS1	YJL172W		CPS1-SRS1	599	636		II		AA		
				CPS1-SRS2	1684	1738		II		AA		
4	DNM1	YLL001W		DBM1-SRS1	262	302	I					
				DBM1-SRS2	1155	1189		II				
5	EAR1	YMR171C		EAR1-SRS1	110	167		II		AA		
				EAR1-SRS2	1578	1614		II	terminal	AA	Olivier et al	Olivier et al this study
6	EGT2	YNL327W		EGT2-SRS1	2194	2235		II		AA		
7	ERG2	YMR202W		ERG2-SRS1	261	314		II				
				ERG2-SRS2	338	380	I			AA		
8	IRC8	YJL051W		IRC8-SRS1	826	870		II		AA		
9	IST2	YBR086C		IST2-SRS1	2597	2626		II		AA		
				IST2-SRS2	2727	2768		II	terminal	AA	Olivier et al	Olivier et al this study
10	KSS1	YGR040W		KSS1-SRS1	730	781		II		AA		this study
11	MET4	YNL103W		MET4-SRS1	991	1051		II				
12	MMR1	YLR190W		MMR1-SRS1	548	590		II				
13	MTL1	YGR023W		MTL1-SRS1	1581	1651	I					
14	TCB2	YNL087W		TCB2-SRS1	3123	3175	I			AA		
				TCB2-SRS2	3556	3592		II		AA		
15	TCB3	YML072C		TCB3-SRS1	1603	1642	I			AA		this study
				TCB3-SRS2	2134	2173		II		AA		
16	TPO1	YLL028W		TPO1-SRS1	175	240		II		AA		
				TPO1-SRS2	461	527	I					
				TPO1-SRS3	774	822		II				
				TPO1-SRS4	1818	1862	I					
17	WSC2	YNL283C	WSC2C	WSC2-C	1326	1373		II		AA		Jambhekar et al

Remarks

SRS instances studies in this work are indicated in **red**.

(*) <u>SRS names in SRS collection</u>   previously existing SRS terminologies are kept
in case of non-existing previous terminologies, the nomenclature contains "-SRS" suffixes (**)

(**) <u>Numbering with suffixes</u>   the numbering follows their positions in 5'-3' RNA sequence.
SRS elements that were discovered later, are numbered with subsequent numbers

(***) <u>SRS as terminal element</u>   indicates such SRS instances that reside at the terminal region of a RNA stem fold

811