A consensus motif in *ASH1* and further transcripts unifies several RNA motifs required for

interaction with the She2p/She3p transport machinery and mRNA localization in yeast

Markus Seiler[1#*], Annika Niedner[2#], Simone Heber[2,3], Michael Feldbrügge[4], Ralf-Peter Jansen[5], Dierk

Niessing[2,3*], Kathi Zarnack[1*]

[1]Buchmann Institute for Molecular Life Sciences (BMLS), Goethe University Frankfurt, Max-von-Laue-

Str. 15, 60438 Frankfurt a. M., Germany

[2]Institute of Structural Biology, Helmholtz Zentrum München, Ingolstädter Landstrasse. 1, 85764

Neuherberg, Germany

[3]Institute of Pharmaceutical Biotechnology, Ulm University, James Franck Ring N27, 89081 Ulm,

Germany

[4]Institute of Microbiology, Heinrich-Heine University Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf,

Germany

[5]Interfaculty Institute of Biochemistry, Tübingen University, Auf der Morgenstelle 34, 72076 Tübingen,

Germany

[#] Equal contribution

[*] Corresponding authors

Keywords: She2p, She3p, localization machinery, RNA localization, RNA zip-code, modeling, consensus

motif, surface plasmon resonance, computational screening

Abstract

Asymmetric localization of the *ASH1* transcript is a central step in the regulation of mating type

switching in *Saccharomyces cerevisae* and a paradigm for localized mRNAs specifically recognized by the

She2p/3p transport machinery. Four RNA elements in *ASH1* are known to mediate bud localization, but it

remained unclear what a consensus motif of all four She2p/3p recognition sites (SRS) might look like. In

this study, we derive the SRS consensus motif, which is characterized by an L-shaped, double-stranded

RNA conformation with two cytosine-containing sequence motifs at conserved positions and a central

asymmetric bulge. It is noteworthy that the spatial arrangement of these features can be retained, despite

their altered order within the nucleotide sequence. We termed these variations "configurations" and

confirmed in biochemical studies that SRS function in *ASH1* elements is preserved when changing

configurations. We tested other known and predicted SRS elements in target transcripts of She2p/3p by

biochemical and biophysical studies and confirmed the conserved SRS function even in switched

configuration. Consistently, we present the *SRS instance collection*, which includes further predictions of

SRS instances of transcript targets of the She2p/3p translocation pathway.

Introduction

The active intracellular transport of effector molecules contributes to the high degree of organization in life and asymmetric mRNA distribution is one way to ensure correct spatiotemporal expression of encoded proteins [1]. In recent years, it has been studied in various polarized cells, in *D. melanogaster* and *Xenopus laevis* oocytes and embryos [2–4], and also in mammalian fibroblasts [5] and neurons [6]. mRNA transport is involved in embryonic patterning, cellular polarity, and asymmetric cell division. Failure of mRNAs to be properly transferred has been shown to cause cellular dysfunction and disease [7]. In these pathways, mRNAs are recognized by the cellular translocation machinery via *cis*-acting elements in their sequences, called zip-codes or localization elements that may be determined by both, short RNA sequence motifs, conformational folds [8–10] and spatial restraints between these features [11]. These zip-codes are recognized by RNA-binding proteins that target the mRNAs to their final destination [12]. Within the mRNAs, the zip-codes often reside in the 3' untranslated regions (3' UTRs) of mRNAs, but also in 5' UTRs and coding regions [13–18].

Budding yeast (*S. cerevisiae*) has evolved as a model organism for mechanistic analyses on specific targeting of mRNAs to the bud of emerging daughter cells [19,20]. The current model of how the RNA transport machinery is built suggests a step-wise assembly of its core compounds: She2p interacts specifically with its target mRNA and recruits the adapter protein She3p together with the type V myosin Myo4p [21–24]. This mRNA ribonucleoprotein (mRNP) complex is then transported along the actin cytoskeleton. The paradigm, the yeast *ASH1* mRNA, is localized to the bud tip enabling the asymmetric distribution of its encoded transcription factor Ash1p into the daughter cell nucleus where it represses mating-type switching [25–28]. To date, over twenty transcripts have been reported to be targeted via the She2p/3p machinery. Besides the well-characterized She2p/3p target mRNAs *ASH1*, *EAR1*, and *WSC2*, further candidates were identified using combined RNA immunoprecipitation and microarray techniques [24,29]. In some target mRNAs, the regions containing putative zip-codes had been narrowed down by yeast three-hybrid and *in vivo* localization studies [30]. However, besides of *ASH1*, precise positions of the She2p/3p recognition sites (SRS) remained unknown in most cases. Within the *ASH1* mRNA, four localization elements with intrinsic SRS

function were mapped in accurate fashion [13,14,19]. Three of these (E1, E2A and E2B) were shown to reside

within the coding region, whereas E3 overlapped the stop codon and extended into the 3' UTR

(**Supplementary Fig. 1A, Tab. 1**). Each *ASH1* SRS element was sufficient for triggering the assembly of

the She2p/3p complex and for localizing a reporter RNA to the bud and was predicted to exhibit RNA folds

with extended stem-loop structures [13,14]. An *in-vivo* screen with mutated *ASH1* fragments supported the

notion that a three-dimensional fold was involved in zip-code function. The fold was described as a stem

with conserved nucleotides on either side: a singular cytosine and a CGA triplet, with a distance of six

nucleotides between the cytosines. Searching for similar RNA folds identified functional SRS elements in

IST2 and *EAR1* [10]. However, these folds lacked an apparent consistency with some of the four *ASH1* SRS

elements and a coherent model for specific SRS recognition by the She2p/3p transport machinery has

remained elusive.

Recently, the interaction interface between E3 and She2p/3p was resolved using X-ray

crystallography: a symmetric tetramer of She2p exhibited two interfaces interacting with E3 on opposite

sides of the complex [8]. As suggested before [31], the affinity was increased by She3p in a synergistic manner.

Intriguingly, the ternary complex formation was accompanied by an induced fit in the SRS element: For

the unbound E3 RNA, SHAPE experiments suggested an elongated conformation [8]. When interacting with

She2p/3p, E3 displayed a sharp turn in one of its bulge regions, resulting in a kinked global conformation

that was also confirmed by NMR structures.

In this study, we define a coherent SRS consensus motif that unifies the features of E1, E2A, E2B

and E3 in *ASH1* and that also classifies SRS elements of further bud-localizing transcripts (*WSC2, EAR1,*

IST2). It exhibits a kinked stem-asymmetric bulge-stem-bulge structure, with the recurrent singular cytosine

in the asymmetric bulge and the CGA motif in a second bulge. One characteristic of the consensus motif is

its arrangement of features, conserved in the overall spatial arrangement but present in two variations

("configurations") in linear sequences. Electrophoretic mobility shift assays and surface plasmon resonance

studies showed that E3 and other SRS elements could be switched between configurations without

impairing She2/3p binding. Using computational screening, we identified potential SRS instances in further

twelve transcripts known to interact with She2/3p. Finally, we introduce the *SRS instance collection*,

describing in detail predicted SRS elements and their positions in further RNA targets of the She2p/3p

translocation pathway.

Results

Bioinformatics predictions reveal a consensus capturing all She2/3p recognition sites of *ASH1*

We wondered if a general concept may exist to describe all four localization elements of the *ASH1* mRNA that act as She2p/3p recognition sites (SRS). For this study, we could rely on a recently published crystal structure that showed in detail how the *ASH1* localization element E3 interacts in a kinked stem structure with She2p/3p [8]. Of note, this interaction involved a molecularly induced fit upon binding that resulted in an asymmetric bulge in the RNA stem with one of its functional cytosines deeply buried inside the protein interface. We hypothesized that the ability to form such an asymmetric bulge could be a pre-requisite of the E3 conformation, whose composition would require different lengths to be kinked. The asymmetric bulge of E3 displayed a large stretch ("Lbulge") and a small one ("Sbulge"), with a length ratio of 6 nt: 3 nt (**Fig. 1A**) (or 4 nt :1 nt, respectively, if considering atypical base pairing inside the bulge as further base pairs). In a next step, we performed a systematic survey of such an arrangements in further SRS elements of *ASH1*. However, common RNA conformation predictors do not consider precise structural rearrangements due to RNA-protein interaction very well. In a step-wise strategy (**Supplementary Fig. 2, Material and Methods),** we first evaluated which of the commonly used prediction tools would predict the best approximation of the kinked E3 conformation in complex with She2p/She3p [8] just from the (single) E3 nucleotide sequence. Of note, we found *Kinefold* [32] to predict the asymmetric bulge in close approximation (5 nt : 2 nt) that could even be closer approximated if considering further details of the crystal structure, resulting in (6 nt : 3 nt) (**Fig.1**, **Supplementary Fig. 3A**).

For E2A, we found the predicted conformation with an asymmetric bulge with exactly the same length ratio as found in E3 (5 nt: 2 nt) using *Kinefold* (**Fig. 1B left)**. Surprisingly, after 180° rotation, the arrangement of the predicted E2A conformation harmonized perfectly with the E3 conformation (**Fig. 1A right, 1B right**). Furthermore, this 180° rotation caused the previously diverging sequence motifs of E3 to fit now to the single and triplet motifs of E2A. Sequence comparison revealed 89% sequence identity between E3 and E2A in the asymmetric bulge and flanking regions (**Supplementary Fig. 1B**). For E1, the

180° rotation of its predicted conformation resulted in an arrangement comparable with E3 (**Fig. 1D**), but

with different length proportions of the asymmetric bulge and a lower degree of sequence identity.

These findings suggested one blueprint, with a common overall spatial arrangement of its

(conformational and sequence) features in E3, E2A and E1 that we referred to as "SRS consensus motif"

(**Fig. 1E,F**). Of note, despite of this conserved spatial arrangement of features, their linear order was

interchangeable in the nucleotide sequences (**Fig. 1F**). This was due to the fact that we had to harmonize

conformations of E3, E2A and E1 by a 180 degree rotational step of E3. Subsequently, the 5' start of E3 is

at the opposite site of the SRS consensus motif than for E2A and E1, resulting again in different feature

orders in the nuclear sequence (**Fig. 1E**). We termed these variations "configurations", with "config-I"

observed in E3 and "config-II" in E2A and E1. In a next step, we refined the annotation of sequence motifs

in E3. We referred to the single cytosine motif in the asymmetric bulge as C_1 and to the cytosine in the

CGA motif as C_2 (**Fig. 1A, right, 1E**), analogously to E2A and E1. At first glance, the conformational

prediction of E2B displayed the hypothesized "Lbulge" even shorter than the "Sbulge" (**Fig. 1C, left**).

Remarkably, Kinefold predicted an alternative conformation with an asymmetric bulge in agreement with

the SRS consensus motif (6 nt: 3 nt), but with a 30% higher content of energy (calculated free energy = -

10.3 kcal/mol versus -7.4 kcal/mol), which could be rearranged by sliding the sequence repeats of the SRS

element (**Fig. 1C, right**). Altogether, we had derived a model of the SRS consensus motif (see **Info-Box**).

In a subsequent step, we could confirm that the model was sufficient to summarize previous functional

studies with mutationally-derived decreases in RNA localization in all four *ASH1* SRS elements [10,13,14,25]

(**Material and Methods**). For a better overview, the distinct modeling steps (initially characterized

elements, Kinefold prediction, rotation, and mutual feature alignments) are displayed in (**Fig. 2**).

The two configurations are interchangeable in E3 without loss-of-function *in-vitro*

One unexpected characteristic of the SRS consensus motif was the spatial conservation of its

features in its overall conformation while the order of features in the nucleotide sequences was found to be

swapped for the different configurations. To validate if we had grasped the essence of the consensus, we

investigated if an artificial switch between the different sequence orders (and the configurations) would

conserve SRS function. For this purpose, we used the well-characterized construct $E3_{51nt}$wt [8] and switched

its feature order in the nucleotide sequence (config-I) to design a config-II construct ($E3_{51nt}$switch, **Fig. 3A**).

Based on the observation that the E3 crystal structure exhibited an interaction of She2p/3p with C_2G but

not with C_2GA, and for comparability reasons with previous studies, we limited the construct design to

C_2G. Using an electrophoretic mobility shift assay (EMSA), we examined $E3_{51nt}$switch for interaction with

She2p and She3p in a titration approach. We used the fusion protein She2p-She3p, which comprised both,

She2p and a She3p fragment and was demonstrated to be sufficient to mediate the synergistic binding of

both proteins [8]. For the wildtype construct $E3_{51nt}$wt, we observed an interaction strength at the expected

nanomolar protein concentration (**Fig. 3A**) [8]. As a control, we tested the construct $E3_{51nt}$M6 characterized

by reduced interaction due to a $C_{1813}G$ substitution of C_2 [8]. In agreement of our model, $E3_{51nt}$wt and

$E3_{51nt}$switch, interacted with She2p-She3p at comparable nanomolar protein concentration (**Fig. 3A**). These

similar binding affinities contrasted with $E3_{51nt}$M6, which displayed severely impaired interaction.

Since the $E3_{51nt}$M6 mutant showed impaired but still considerable binding to She2p-She3p-fusion

protein, we used the full-length proteins to assess the binding features of E3 elements. Incubation with

constant She3p concentration of 0.56 µM and titration of She2p (0.02 - 5 µM) confirmed that both

constructs, $E3_{51nt}$wt and $E3_{51nt}$switch, supported binding even at the lowest She2p concentration used in this

experiment (0.02 µM). In contrast, binding of the full-length proteins to $E3_{51nt}$M6 was abolished at

concentrations up to 5 µM (**Supplementary Fig. 4B**) x. The same was true for a reciprocal approach, in

which increasing concentrations of She3p (0.02 - 5 µM) were tested in the presence of high amounts of

She2p (5 µM) (**Supplementary Fig. 4C**). Moreover, both experiments showed that She3p efficiently binds

E3 in the absence of She2p. Surprisingly, this binding was sensitive to mutation of C_2. In conclusion, these

studies successfully confirmed that E3 can be switched from config-I to config-II without compromising

its functionality to interact with She2p/3p.

When we analogously examined E2A and E1, we reversed the direction of switching (now from

config-II to config-I), resulting in the constructs $E1_{27nt}$switch and $E2A_{32nt}$switch (**Supplementary Fig. 5**).

Again, in EMSA validations, we could demonstrate that both constructs largely exhibited the binding

properties to She2p-She3p-fusion. In summary, the "config-switch" approach supported our model, which

suggested that SRS features could be rearranged in the RNA sequence while retaining function. We could

demonstrate that E1, E2A, and E3 recognized She2p/3p regardless of configuration, demonstrating the

essential characteristics of RNA-She2p/3p recognition were reflected in the SRS consensus motif. As a

conclusion, these findings indicate that SRS function requires very few conserved bases in defined positions

of a stem-loop, which must accommodate a bulge that is flexible enough to open up upon binding by She2p-

She3p to adopt a kinked L-shaped conformation in the co-complex.

In a fine-grained approach, we used the analytical power of surface plasmon resonance (SPR), to

investigate the influence of the functional cytosines C_1 and C_2 on SRS function. Previous studies had

determined a preferred distance of six nucleotides between these cytosines [10]. Using the functionally

minimal construct $E3_{29nt}$ [8], we introduced varying distances between C_1 and C_2 and performed titration

series in a surface plasmon resonance experiment with a She2p-She3p-fusion construct (125 nM) (**Fig. 3B**).

As a result, for the wildtype construct $E3_{29nt}$wt, we observed a binding affinity of approx. $K_D = 0.89$ nM

(**Fig. 3B, Supplementary Fig. 6**). In contrast, constructs with C_1-C_2 distances of five and seven nucleotides

led to a reduction in binding of at least one order of magnitude (**Fig. 3B**, **Supplementary Tab. 1**),

confirming the findings of 6 nt as optimal C_1-C_2 distance.

A computational algorithm to screen for the consensus in known She2p/3p substrates

We performed computational screening (details, see **Material and Methods**) to identify further

instances of the SRS consensus motif in further target sequences of the She2p/3p transport machinery. For

this purpose, we screened a list of 24 transcripts reported to show *in-vitro* binding by She2p/3p or bud

localization *in-vivo* [24,29,30]. The identified sequence regions represented potential SRS instances and were

evaluated for their potential to form the correct conformation of one of the configurations, again using

Kinefold for this step We intended to extend this conformational evaluation by integrating data describing

the degree of unpaired state of nucleotides (obtained in a remarkable genome-wide study using parallel

analysis of RNA structure (PARS), [33]). This strategy was promising, but for our purposes the data were not

detailed enough, as the averaged values for each nucleotide position suggested competing RNA

conformations even for some of the strong *ASH1* SRS elements (**Material and Methods, Supplemental**

Fig. 7).

For the three transcripts *EAR1, IST2* and *WSC2*, their SRS elements had been identified already,

with their sequence regions narrowed down to fragments [10,30]. Now, with knowledge of the SRS consensus

motif, these SRS elements could be classified as a config II arrangement (**Fig. 4C, Supplementary Fig. 8**).

Remarkably, for *EAR1* and *IST2*, the asymmetric bulge was even more eccentric than observed in *ASH1*

(*EAR1:* 7 nt :0 nt; *IST2*: 9 nt : 1 nt). In addition, we identified an additional upstream SRS instance in each

transcript, *EAR1* and *IST2* (**Tab. 2**). In contrast, for *WCS2*-C, the initially identified SRS fragment turned

out to comprise possibly two SRS elements, a canonical one in the center of this fragment, and a potential

SRS instance in the flanking regions, however, with a C_1-C_2 distance of 7 nt (**Supplemental Fig. 8D**).

However, we could not classify the second published instance in *WSC2*, *WSC2*-N, in agreement with our

model (see **Discussion**). As a first outcome of the screening we had identified seven SRS instances (**Fig.**

3C). When we studied the seven canonical SRS elements in detail, we noticed a recurring di-adenosine in

the L-bulge ("AA_L") as a common feature in all elements (**Fig. 3C**), possibly indicating relevance in SRS

function. This was surprising due to the fact that the crystal structure of E3 showed the interaction between

She2p/3p and the AA_L nucleotides occurs exclusively via the backbone atoms of the nucleotides, and thus

unspecifically [8], with their nucleobases having intramolecular contact within the RNA structure. This

observation makes relevance in the context of SRS recognition less likely. As mentioned before, for most

known SRS instances (20 of 24 instances in our set), functionality had been exclusively studied for large

RNA fragments or complete transcripts. For 13 of these transcripts, we could identify now the positions of

SRS instances. For 12 transcripts, we found one or two instances in each sequence, and we found four

instances in *TPO1* (**Fig. 4A**). Altogether, in 17 of 24 sequences (*ASH1*, *WSC2*, *EAR1*, *IST2* included), we

predicted 31 SRS instances, with ten instances (32%) in config-I and 21 instances (68%) in config-II,

respectively (**Fig. 4A**). Of notice, almost two thirds of the SRS instances contained the AA_L di-nucleotide

in the Lbulge (**Fig. 4B**).

Validation of SRS elements in *EAR1*, *IST2*, *KSS1* and *TCB3*

As described before, in *IST2* and *EAR1*, sequence fragments with SRS function had been identified

already, and these sequences were also identified in our computational screening (termed *EAR1*-SRS2,

IST2-SRS2; referring to the nomenclature, see **Tab. 2**). We could classify their functional features, in

agreement with the SRS consensus motif, provided that the consensus was valid also for all these instances.

In such a view, the SRS elements would resemble the very first characterization (Olivier et al., 2005),

however in the larger context of a SRS consensus motif. To address this aspect, we studied the interaction

between specifically designed SRS constructs and She2p-She3p fusion proteins, analogous to our previous

studies. In accordance with the consensus, the newly designed constructs $EAR1_{26nt}$switch and $IST2_{29nt}$switch

were strictly reduced to those regions assumed to be essential for SRS function and also switched in

configuration (each to config-I) (**Fig. 5A**). As we could show in EMSA studies, $EAR1_{26nt}$switch was bound

with affinities comparable to the unmodified wild-type element $EAR1_{50nt}$wt (**Fig. 5A**). Although for

$IST2_{29nt}$switch binding was clearly retained, it was considerably weaker when compared to $IST2_{62nt}$wt. In

addition, the *SRS instance collection* contained instances in transcripts for which no detailed position

information on SRS instances in transcripts had previously been determined. From these instances, we

chose *KSS1*-SRS1 (config-II, **Tab. 2**) and *TCB3*-SRS1 (config-I, **Tab.2**) for further functional analysis. As

a result, we were able to show that $KSS1_{72nt}$wt interacts strongly with the She2p/3p fusion protein *in-vitro*

(**Fig. 5B**) and observed similar strong binding for the minimal construct switched in configuration

$KSS1_{29nt}$switch (to config-I), confirming *KSS1*-SRS1 as a potential SRS element. In an analogous manner,

we tested the minimal construct $TCB3_{38nt}$wt (config-I) and observed also strong interaction with the fusion

protein (**Fig. 5B**) confirming also *TCB3*-SRS1 as potential SRS element. In summary, we were able to

demonstrate the identification and *in-vitro* confirmation of new SRS instances. Furthermore, for some of

these instances we were able to prove that the newly designed, size-reduced constructs retain the SRS

function even in switched configuration. In conclusion, the fact that our model, derived exclusively from

ASH1 SRS elements, enabled the identification, size reduction, and config-switch of designed SRS

constructs in previously unexplored sequences while preserving SRS function confirmed its applicability

as an SRS consensus motif.

Validation of further features in the SRS of *EAR1*

In *EAR1* (*YMR171C*), it had been shown for its SRS element (now termed *EAR1*-SRS2) that

substitutions in the sequence motifs decreased function [10]. The substitution of both, C_1 (updated

nomenclature) (construct "M2" in [10]) and C_2 (construct "M1") had shown severe effects *in-vivo*. To examine

the SRS consensus motif and *EAR1*-SRS2 in more detail, we chose a fine-grained approach using the

$EAR1_{31nt}$switch construct (switch to config-II) as the base construct for further changes (**Fig. 6A**) and

determined its binding affinity using SPR analysis as described before. As result, we obtained an affinity

of the original construct $EAR1_{31nt}$switch of $K_D = 11$ nM (**Fig. 6D, Supplementary Fig. 9**). Remarkably,

compared to the E3 minimal element, the affinity with the She2p/3p fusion protein was decreased by one

order of magnitude ($E3_{29nt}$wt: $K_D = 0.89$ nM). When we examined the original construct $EAR1_{31nt}$switch for

modifications in the inverted repeat IR1, by inversions or additionally introduced base pairs, the modified

constructs showed affinities in the range of the original construct suggesting no direct impact on SRS

function (**Fig. 6D**). Second, we examined the function of the di-nucleotide AA_L in the Lbulge by deletions

of varying extent (**Fig. 6C**), analogously observing binding affinities in the magnitude of the basic construct

(**Fig. 6D**). The analysis of the functional cytosines C_1 and C_2 was complex since two cytosines (C_{1601} and

C_{1602}) precluded an unambiguous assignment of the C_1 position (**Fig. 6B**). The substitution of either cytosine

(constructs $C_{1-1}U$, $C_{1-2}U$) resulted in a considerable binding decrease suggesting that both positions

contribute to She2p/3p binding (**Fig. 6D**). Unexpectedly, a deletion of one of the C_1 positions (ΔC_1) or a

simultaneous substitution of both cytosines ($C_{1-1}U$ & $C_{1-2}U$) restored She2p/3p binding (see **Discussion**).

In contrast, substituting C_{1590} in the three-nucleotide $C_{1590}GA$ motif (C_2U) returned unequivocal results

manifesting in a dramatic loss of binding. Altogether, these results confirmed that neither sequence

composition of IR1 nor the AA_L motif in the Lbulge were relevant for She2p/3p binding *in vitro*. Alterations

of the C_1-C_2 distance were tolerated in some SRS contexts but not others, with a strong bias for a C_1-C_2

distance of six nucleotides. We therefore kept a C_1-C_2 distance constraint of six nucleotides for subsequent

computational SRS screening.

A collection of predicted *SRS* instances in known target transcripts of She2p/3p

Computational screening for the SRS consensus motif had identified SRS instances in 17 of 24

sequences (**Fig. 4A**). After positive *in-vitro* validation of several of the new SRS instances (in *EAR1*, *IST2*,

KSS1, *TCB3*), we finally created a collection of these screening results to present to the community (*SRS*

instance collection, **Tab. 2**). This collection contains information on 32 SRS instances in 17 transcripts that

have been reported to show She2p/3p binding *in-vitro* or bud localization *in-vivo* [24,29,30]. In addition, a

standardized nomenclature for SRS elements is introduced and further information is provided on positional

data, classification and other details of SRS instances, several of them are promising new candidates for

future experimental investigations.

Discussion

In this work, we have derived the unified model of a three-dimensional RNA motif that functions as She2p/3p recognition motif (SRS) with the She2p/She3p protein transport machinery, a prerequisite for active transport of transcripts to the yeast bud. We used a computer-aided and conceptual modeling approach that based on the hypothesis that an asymmetric bulge in the RNA conformation might be a prerequisite of SRS function. This approach led to the identification of a common three-dimensional arrangement of SRS functional features (conformational features and sequence motifs) in the *ASH1* localization elements E3, E2A and E1 (and with limitations also for E2B) in a blueprint, thus demonstrating the long-missing consensus motif of all four SRS elements. Unexpectedly, although this arrangement of features was conserved in its overall three-dimensional fold, , there were two variations with respect to the linear arrangement in the nucleotide sequences. We termed these variations "configurations" ("config-I" in E3, "config-II" in E2A, E1 and E2B). The consensus model proved to be correct based on our consistent results from electric mobility shift assays, in which we analyzed the SRS function of RNA-constructs with re-designed feature arrangement. According to our model, this rearrangement of the features affected their linear order in the nucleotide sequences, but allowed the preservation of their overall three-dimensional arrangement. This re-design therefore represents a switch between their original configuration and the respective counterpart configuration ("config-switch"). We were able to consistently demonstrate that the config-switch retained SRS function in all experiments we performed indicating that the consensus motif comprises the main requirements of SRS function. The fact that also size-reduced constructs proved to keep SRS function was a further hint that we had grasped the essential features of SRS function. It can be described in all *ASH1* SRS elements as follows: an asymmetric bulge, flanked by inverted repeats, forms a kinked, L-shaped RNA conformation. The larger side of the bulge contains the single cytosine motif C_1 inside of the first three positions. The C_2GA motif resides on the opposite strand of C_1, separated by an inverted repeat in downstream direction on this strand. The two configurations are marked by the two different 5' entry sites into this kinked conformation. These findings were further confirmed, when we compared previous thorough mutagenesis studies of the *ASH1* SRS elements and could assign these results

well according to their relevance in the SRS consensus model. Further, by studying the binding kinetics of

$E3_{29nt}$wt in surface plasmon resonance analysis, we could confirm the optimal distance between C_1 and C_2

with six nucleotides as it was suggested before (ca. 28 Å) [10]. It is noteworthy that the existence of two

(double-stranded) configurations that can be converted into each other by swapping their features in the

nucleotide sequence finds an analogy in studies on the beta-actin RNA zip code, where the order of its

(single-stranded) 5' and 3' elements is interchangeable within the target transcripts bound to ZBP1 [11].

Due to its universal character to describe all four *ASH1* SRS elements, the exploitation of the SRS

consensus motif went beyond previously proposed computational screening strategies. Hence, we created

a computational screen that searched for both configurations, followed by a conformational check of the

overall RNA fold performed by *Kinefold*. For some of the further transcripts that had been found to localize

at the bud tip of yeast cells, sequence fragments had already been narrowed down with retained SRS

function. In this set of sequence fragments, we could classify SRS elements of *EAR1*, *IST2* and *WSC2* as

further instances of the SRS consensus motif.

EAR1 was first identified as localized in the bud by [29] with its SRS element characterized in further

studies [10] and classified as a config-II SRS element in this work (*EAR1*-SRS2, **Tab.1**). We were able to

confirm previous studies that the specific nucleotide sequences of the inverted repeats did not contribute to

SRS function [8,13]) by applying SRP to different modifications of the minimal construct $EAR1_{31nt}$switch. In

contrast, the loss of C_2 showed a strong impact on SRS function that was also consistent with previous *in-*

vivo studies [10]. Analogously, individual substitutions of each of the two potential C_1 positions (in C_1-C_2-

distance of five or six nucleotides) affected SRS function. Unexpectedly, a single deletion of C_{1601} or a

simultaneous substitution of C_{1601} and C_{1602}, both restored SRS function. When studying the potential di-

nucleotide AA_L sequence motif by deletion experiments we could not detect any direct impact on SRS

function, even if the Lbulge region was almost completely deleted. So far, we can neither explain the

apparent discrepancy in the C_1 mutation effects in the *EAR1* constructs nor the retained SRS function with

deleted asymmetric bulge. It may reflect both complex sequence composition and inherent redundancy in

this particular SRS instance.

So far, we had observed a config-I arrangement only in E3 whereas we had identified six SRS

elements with config-II (**Fig. 4C**). This dominance of config-II SRS elements continued when we applied

the computational screen to our complete sequence set and predicted 31 instances in 17 sequences, with 22

predicted elements (71%) exhibited config-II. So far, the reason for the unequal frequency of the two

configurations remains unclear. For each of the two sets of SRS instances (with relatively well-known

positions in their transcripts or with unknown SRS positions), we experimentally analyzed two instances,

utilizing EMSA approaches as described before. For the first subset, we could confirm SRS function of

previous studies in both SRS elements *EAR1*-SRS2 and *IST2*-SRS2 [10]. For the second subset, we could

verify both new identified SRS elements *KSS1*-SRS1 and *TCB3*-SRS1 *in-vitro.* Remarkably, when we

switched configuration in *EAR1*-SRS2, *IST2*-SRS2 and *KSS1*-SRS1 (config-II to config-I), we could

demonstrate again conserved SRS function for <u>all</u> switches, strongly supporting the SRS consensus model.

Of note, as a first successful validation *in-vivo*, we had been able to identify and confirm the nucleotide

positions of a SRS element in the *CLB2* transcript [34]: In the transcript, only a few but functionally crucial

positions of the SRS instance *CLB2*-SRS1 (config-II, 1111 nt - 1145 nt) were altered in non-canonical

mutations to reveal a significant loss-of-function effect in bud-localization and a shift in protein distribution

from the buds towards mother cells. It is also interesting to note that the regulatory effect of well-defined

asymmetric bulges seems not to be limited to SRS function. As described in [35], the structure of the pre-

mRNA autoregulatory complex of the L30 ribosomal protein in yeast displays an asymmetric bulge (length-

ratio 5 nt : 2 nt) with a cytosine positioned in an extension of the Lbulge, and separated from another

cytosine (in a further bulge) by an IR in a distance of 6 nt. Remarkably, it is this asymmetric bulge region

that is part of a mutual induced fit, suggesting analogies with the observed induced fit mechanisms in the

crystal structure of E3 and She2p/3p.

Further aspects of SRS characteristics remain to be investigated. For example, a further extension

of the description of the SRS consensus motif is thinkable: It is reasonable to ask if the SRS function would

be conserved if Lbulge and Sbulge were joined together without IR2 as intervening feature. Although

experimental analysis of such a setting is beyond the scope of this study, we would like to put to discussion

such a theoretical construct, to possibly take a step forward to complete the picture (**Supplementary Fig.**

10A). The SRS element *WSC2*-N [36] that could be not classified, might be such an example (**Supplementary**

Figure 10B, left). Such an extended perspective might have further implications also for the understanding

of some of the SRP results of *EAR1* described before. Both elements, *EAR1*-SRS2 and *IST2*-SRS2 displayed

<u>two</u> CGA motifs with the following consequences: Whereas we considered *EAR1*-SRS2 as config-II with

C_2GA residing in the terminal bulge, this terminal bulge also may be considered as a joint of Lbulge and

Sbulge, hence offering a new perspective: the cytosine motif in this terminal bulge may be considered as

C_1 while its counterpart (initially referred to as C_1 inside of the asymmetric bulge) may be extended to

C_2GA (**Supplementary Figure 10B, right**). In such a view, the functionality of *EAR1*-SRS2 would exhibit

redundant features of two superimposed and oppositely rotated SRS elements in the bigger context of an

(extended) consensus motif. This perspective may also explain partially the ambiguous results in the SPR

experiments of our study. Secondly, a further extension of the SRS consensus could be formed by a range

in the C_1-C_2 distance. In consistency with previous studies [10] and with our SPR results we focused on the

optimal parameters of six nucleotides in this study to minimize false positive matches. However, we could

not exclude a minimal functionality for instances with (C_1-C_2 distance $\neq 6$). An example might be a second,

potential SRS instance that we identified in the flanking regions of *WSC2*-SRS2 (*WSC2*-C) but with a C_1-

C_2 distance of seven nucleotides (**Supplementary Fig 7C**). In such a case, *WSC2*-SRS2 may compensate

the suboptimal C_1-C_2 distance of its neighboring SRS element by inducing SRS conformation in a

cooperative manner when both SRS instances interact with the binding interfaces on the opposite sites of

the She2p tetramer [37].

As a second aspect, localization elements that mediate bud localization require the interaction with

the She2p/3p protein transport machinery, but further motifs may be required as binding platforms of

auxiliary factors. For E2B, we have speculated that its energetically optimized conformation had to be

rearranged from an alternative conformation to display the SRS consensus motif. Even strong SRS elements

such as E2A [12] can be subject to major reorganizations, as shown by the PARS data which indicate

competing conformations for E2A [33] (**Supplementary Fig. 7**). Therefore, the She2p/3p interaction can be

seen as a central step in the zip-code function, but it might be framed by further rearrangements in the

chronological order. In such a scenario, transient binding of auxiliary cofactors could stabilize the

asymmetric bulge and increase SRS function. AA_L might be such a co-factor binding motif. Despite the

fact that AA_L was present in each case of our seven well-characterized, bud-localizing SRS elements,

neither the crystal structure of [8] nor the results of SPR studies suggest involvement in She2p/3p binding

specificity. Future research must clarify the possible function of AA_L *in-vivo*, also with regard to transient

interactions. In addition, for E3 and E2A, it was tempting to speculate if their extraordinary sequence

identity was a product of convergent evolution, due to further requirements in affecting zip-code function.

Maybe we have to face a trade-off between canonical SRS consensus motif and stability of the individual

RNA conformation, with the weakening of one aspect may be compensated by strengthening of the other.

This would complicate the identification of SRS instances, especially if auxiliary co-factors would alter the

probabilities of formation of SRS conformations *in-vivo*. Summarizing, all these aspects stress two aspects:

(i) The new SRS consensus motif delivers a conceptual framework of how to classify further aspects of

new SRS instances, and (ii) It remains necessary to combine experimental screenings (identification of bud-

localizing transcripts, final validation of instances) and computational screenings (prediction of positional

information in transcripts to narrow down experimental analysis and suggest crucial positions in mutations

studies). Consequently, the *SRS instance collection* contains only such SRS instances where bud

localization had been confirmed already, at least at the transcript level.

In conclusion, the SRS consensus motif enabled the description of the four SRS elements of *ASH1*

in one model, hence providing the long-missing consensus. It further unified the described functional

features of the known SRS elements in *IST2*, *EAR1* and *WSC2* and enabled the identification of new SRS

elements in further bud-localizing transcripts. The consensus model proved to be consistent with previous

mutagenesis studies of these elements and we could demonstrate that a switch between the configurations

retained function in SRS constructs of several transcripts.

Materials and Methods

Overview on the theoretical data flow to derive the SRS consensus model

The theoretical part of deriving the SRS consensus model comprised the following steps

(**Supplementary Fig. 2**): (1) Hypothesis (asymmetric bulge as hallmark of SRS) by analysis of crystal

structure of *ASH1*-E3 (2) Selection of an appropriate conformational prediction tool (3) Conformational

prediction of instances and inspection to derive a consensus (**Fig. 2**) (4) Cross-check of the consensus with

published functional data (5) Creation of a computational screening pipeline (6) Check for further data

sources to filter results (**Supplementary Fig. 7**) (7) Screening for instances in selected sequence sets (8)

Inspection of instances for further patterns

Identification of an appropriate conformational prediction tool

The inspection of the crystal structure of *ASH1*-E3 suggested an asymmetric bulge as a pre-requisite

of kinked RNA conformation. The occurrence of such asymmetric bulges may be favorized in mRNPs by

the energy release due to molecular interaction. However, its occurrence is less likely in an individual RNA

and is less likely to be predicted as plausible conformation by computational prediction tools that commonly

deal with individual RNA sequences. In a survey, when investigating which of common tools [32,38–40] was

most likely to predict the asymmetric bulge of E3, we found Kinefold [32] as the one that best tolerated

asymmetric bulges (**Supplementary Fig. 3**). It predicted the asymmetric bulge consisting of a 5 nt Lbulge

and a 2 nt Sbulge, which closely approximates the length ratio of the asymmetric bulge observed in the E3-

She2p/3p crystal structure [8] that displayed a length ratio of 6 nt : 3 nt (considering only Watson-Crick base

pairs) or 4 nt : 1 nt (considering also non-canonical base pairings as found in the crystal structure),

respectively (**Fig. 1A**).

Cross-check of the consensus with published functional data

We used published data of functional studies that involved both, mutagenesis and deletion of RNA

regions to cross-check for consistency with the SRS consensus model [10,13,25,30]. We characterized such

mutations in *ASH1* SRS elements that could clearly be assigned to destroy or to keep SRS function

according to the consensus model. All these constructs were in accordance with the consensus model (data

not shown).

Development of a computational screening approach

For the purpose of identifying further instances of the SRS consensus motif, we transformed the

consensus features into "patterns" to screen for "matches" in RNA sequences. We created a computational

screening pipeline consisting of in-house computational scripts and an external conformational prediction

algorithm. For the screening, three aspects were required: (1) Screening the sequence for conformational

aspects. For each RNA, the sequence was screened for two nested pairs of inverted repeats (each with a

minimal length of four nucleotides) that framed an asymmetric bulge region. We had to consider both

configurations in different screening patterns. The obtained constituted potential SRS instances. (2)

Screening the sequence for functional motifs. For each SRS instance, the Lbulge region was screened for

C_1, the opposite strand for C_2GA (positioned beyond IR1), with a C_1-C_2 distance of 6 nt between. The

screening strategy based on the approach of modular feature screening of mosaic-like patterns [41]. This

modular approach allowed to introduce further features to be screened (e.g., AA_L). It also tolerated length

variations between the distinct (sequence and conformational) features (e.g., one version tolerated C_1

positioned in the last nucleotide of IR1). As a consequence, permutated sets of varying patterns could be

summarized into two screening patterns (one for each configuration). (3) Validation by conformational

prediction. For each SRS instance, it was calculated if it would express a stable conformational fold that

fitted to the conformational arrangement of the SRS consensus. This analysis was performed using *Kinefold*

as energy-minimization approach that had proven best for large asymmetric bulges. On each site , the SRS

stretches were extended by 10 nt of the flanking sequences of their RNA sequences, to tolerate some degree

of freedom during calculation. In a series of tests using known SRS elements, a flanking region extension

by 10 nt had proven to be best in a range of 0 nt to 50 nt (data not shown). Of note, natural limitations of

common RNA conformational predictions occur due to the fact that such calculations are limited to predict

the conformation of distinct RNA stretches but less reliable regarding statements of the complete transcript

and second, they would ignore any energetic benefits by binding of molecular interactors (e.g., She2p/3p

proteins). As a consequence, we analyzed the 25th best conformational predictions of *Kinefold* (maximum

value = 25), to consider slight structural dynamics in each RNA molecule.

Check for further data sources to filter for the consensus

For the purpose to possibly extend the validation step of new SRS instances by conformational

predictions. we intended to cross-check the predicted conformations with experimental data to get deeper

insights on actual RNA conformations *in-vivo*. For our purposes, we tested the remarkable genome-wide

study based on PARS (Parallel Analysis of RNA Structure) [33]. For each nucleotide in transcripts, PARS-

scores described the percentage in occurring in single-strand ("S1-reads" of RNase S1) or double-strand

("V1-reads" of RNase V1) regions of the RNA conformations. We extracted the S1- and V1-data and tested

the strong localization element E3 [13] as positive control. As a result the PARS data was well suited to

describe the SRS conformation as observed in the E3 crystal structure [8] (**Supplemental Fig. 7**). However,

for further controls, the data comparison of paired/ unpaired occurrence of nucleotides between PARS data

and SRS consensus model failed (**Supplemental Fig. 7**). Commonly, RNA molecules display not one

distinct conformation in-vivo but are subject to conformational dynamics resulting in collections of

competing conformations. PARS data per nucleotide averaged the binding state (paired/unpaired) over all

competing conformations. It seemed that even the two SRS elements in *ASH1*-E1 and *ASH1*-E2A that were

shown to function as efficient localization elements [13] were not so stable in their consensus conformations

that these would be so conspicuous that they could be distinguished from alternative conformations in the

PARS data.

Further bioinformatics resources

For the investigation of molecular structures, we used Pymol [42] as 3D- viewer. Initial protein

characterization of She2p and She3p was performed using the ELM server [43]. RNA sequence alignments

were performed using Jalview [44]. RNA sequences were extracted of the Saccharomyces Genome Database

[45]. For the actual screening, the mRNA sequences of *ASH1*, *IST2*, *EAR1* (aka *YMR171C*), *WSC2* and 20

further mRNA sequences known to She2p/3p interact [10,13,25,29,30,36] (Chartrand1999; Chartrand2002;

Olivier2005; Takizawa2000; Shepard2003; Jambhekar2005) were chosen. As sequence data, we used the

CDRs, extended by 200 nt downstream by default, due to the fact that for several mRNAs, the actual 3'UTR

lengths were still unknown. Functional annotation of Supplementary Table S1 followed [46]. Graphical

visualization of RNA conformations was performed using VARNA [47].

Protein expression

Expression in *E. coli* and purification of She2p, She3p, their fusion protein, and variants thereof

were performed as described in [8,23]. Purity and solubility were confirmed by size-exclusion chromatography

and SDS page.

RNA preparation and isotope labeling

For small scale *in vitro* transcription the MEGAshortscript T7 kit (life technologies) was used

according to the manufacturer's instructions. Synthetic DNA oligonucleotides containing a T7 promoter

and the respective sequence were used as a template. After *in vitro* transcription, the template was digested

by DNAse and the RNA was extracted using phenol:chloroform:isoamyl alcohol (25:24:1). Homogeneity

of RNA was assessed by native and denaturing TBE agarose gel electrophoresis.

Electrophoretic mobility shift assay

In a total volume of 20 µL 5 nM radioactively labeled RNA, 100 µg/mL yeast tRNA competitor,

and protein at a given concentration were mixed in buffer (20 mM Hepes, pH 7.8, 200 mM NaCl, 2 mM

$MgCl_2$, 2 mM DTT) with 4 % (v/v) glycerol. After 25 min at room temperature, the samples were resolved

by native TBE PAGE (6 % polyacrylamide) and gels incubated for 15 min in fixing solution (10 % (v/v)

acetic acid, 30 % (v/v) methanol) and subsequently vacuum dried. Gels were analyzed with radiograph

films or by a using Phosphorimager (FujifilmFLA5100).

Surface plasmon resonance analysis (SPR)

SPR experiments were performed with a BIACORE 3000 system (GE Healthcare). For protein-

RNA interactions, biotinylated RNA was streptavidin-captured on an SA-Chip (GE Healthcare) surface

according to manufacturers instructions. Measurements were performed at a flow rate of 30 µL/min in

standard buffer. Data were analyzed using the BIAevaluation software (GE Healthcare). Sample

sensograms were referenced against the signal in a ligand-free reference channel, and averaged values at

the equilibrium calculated. Steady state binding curves were obtained by plotting the response units at

equilibrium against protein concentration with the Origin Pro 9.0 (OriginLab) software.

Computational quality check of the folds of the experimental constructs

Part of the design of experimental constructs (minimal, config-switch) for the *in-vitro* and *in-vivo*

approaches was a quality check with *Kinefold*: The conformations of their sequences were predicted to

ensure that their expected conformation would be in agreement with the SRS consensus model

Acknowledgements

We would like to thank Herve Isambert for helpful discussions regarding Kinefold and Toby

Gibson for further support of M.S. This work was supported by the Deutsche Forschungs–gemeinschaft

(DFG, German Research Foundation) to R.-P.J. (DFG-FOR2333, Project Number 270067186), to M.F.

(CEPLAS EXC 1028, CRC1208, DFG-FOR2333, DFG Fe448/8, DFG Fe448/9), to D.N. (DFG-FOR2333,

Project Number 270067186), and to Z.K. (DFG-FOR2333, Project Number 270067186).

Declaration of Interest

The authors declare no potential conflicts of interest.

References

(1) Eliscovich, C.; Singer, R. H. RNP Transport in Cell Biology: The Long and Winding Road. *Curr.*

Opin. Cell Biol. **2017**, *45*, 38–46. https://doi.org/10.1016/j.ceb.2017.02.008.

(2) Bashirullah, A.; Cooperstock, R. L.; Lipshitz, H. D. RNA Localization in Development. *Annu. Rev.*

Biochem. **1998**, *67*, 335–394. https://doi.org/10.1146/annurev.biochem.67.1.335.

(3) Johnstone, O.; Lasko, P. Translational Regulation and RNA Localization in Drosophila Oocytes and

Embryos. *Annu. Rev. Genet.* **2001**, *35*, 365–406.

https://doi.org/10.1146/annurev.genet.35.102401.090756.

(4) Kloc, M.; Bilinski, S.; Chan, A. P.; Allen, L. H.; Zearfoss, N. R.; Etkin, L. D. RNA Localization and

Germ Cell Determination in Xenopus. *Int. Rev. Cytol.* **2001**, *203*, 63–91.

https://doi.org/10.1016/s0074-7696(01)03004-2.

(5) Palacios, I. M.; St Johnston, D. Getting the Message across: The Intracellular Localization of mRNAs

in Higher Eukaryotes. *Annu. Rev. Cell Dev. Biol.* **2001**, *17*, 569–614.

https://doi.org/10.1146/annurev.cellbio.17.1.569.

(6) Dahm, R.; Kiebler, M.; Macchi, P. RNA Localisation in the Nervous System. *Semin. Cell Dev. Biol.*

2007, *18* (2), 216–223. https://doi.org/10.1016/j.semcdb.2007.01.009.

(7) Abouward, R.; Schiavo, G. Walking the Line: Mechanisms Underlying Directional mRNA Transport

and Localisation in Neurons and Beyond. *Cell. Mol. Life Sci. CMLS* **2021**, *78* (6), 2665–2681.

https://doi.org/10.1007/s00018-020-03724-3.

(8) Edelmann, F. T.; Schlundt, A.; Heym, R. G.; Jenner, A.; Niedner-Boblenz, A.; Syed, M. I.; Paillart, J.-

C.; Stehle, R.; Janowski, R.; Sattler, M.; Jansen, R.-P.; Niessing, D. Molecular Architecture and

Dynamics of ASH1 mRNA Recognition by Its mRNA-Transport Complex. *Nat. Publ. Group* **2017**,

24 (2), 152–161.

(9) Niedner, A.; Edelmann, F. T.; Niessing, D. Of Social Molecules: The Interactive Assembly of ASH1

mRNA-Transport Complexes in Yeast. *RNA Biol.* **2014**, *11* (8), 998–1009.

https://doi.org/10.4161/rna.29946.

(10) Olivier, C.; Poirier, G.; Gendron, P.; Boisgontier, A.; Major, F.; Chartrand, P. Identification of a Conserved RNA Motif Essential for She2p Recognition and mRNA Localization to the Yeast Bud. *Mol. Cell. Biol.* **2005**, *25* (11), 4752–4766. https://doi.org/10.1128/MCB.25.11.4752-4766.2005.

(11) Patel, V. L.; Mitra, S.; Harris, R.; Buxbaum, A. R.; Lionnet, T.; Brenowitz, M.; Girvin, M.; Levy, M.; Almo, S. C.; Singer, R. H.; Chao, J. A. Spatial Arrangement of an RNA Zipcode Identifies mRNAs under Post-Transcriptional Control. *Genes Dev.* **2012**, *26* (1), 43–53. https://doi.org/10.1101/gad.177428.111.

(12) Chartrand, P.; Singer, R. H.; Long, R. M. RNP Localization and Transport in Yeast. *Annu. Rev. Cell Dev. Biol.* **2001**, *17*, 297–310. https://doi.org/10.1146/annurev.cellbio.17.1.297.

(13) Chartrand, P.; Meng, X. H.; Singer, R. H.; Long, R. M. Structural Elements Required for the Localization of ASH1 mRNA and of a Green Fluorescent Protein Reporter Particle in Vivo. *Curr. Biol. CB* **1999**, *9* (6), 333–336. https://doi.org/10.1016/s0960-9822(99)80144-4.

(14) Gonzalez, I.; Buonomo, S. B. C.; Nasmyth, K.; von Ahsen, U. ASH1 mRNA Localization in Yeast Involves Multiple Secondary Structural Elementsand Ash1 Protein Translation. *Curr. Biol. CB* **1999**, *9* (6), 337–340.

(15) Kanke, M.; Jambor, H.; Reich, J.; Marches, B.; Gstir, R.; Ryu, Y. H.; Ephrussi, A.; Macdonald, P. M. Oskar RNA Plays Multiple Noncoding Roles to Support Oogenesis and Maintain Integrity of the Germline/Soma Distinction. *RNA N. Y. N* **2015**, *21* (6), 1096–1109. https://doi.org/10.1261/rna.048298.114.

(16) Mori, Y.; Imaizumi, K.; Katayama, T.; Yoneda, T.; Tohyama, M. Two Cis-Acting Elements in the 3' Untranslated Region of Alpha-CaMKII Regulate Its Dendritic Targeting. *Nat. Neurosci.* **2000**, *3* (11), 1079–1084. https://doi.org/10.1038/80591.

(17) Serano, J.; Rubin, G. M. The Drosophila Synaptotagmin-like Protein Bitesize Is Required for Growth and Has mRNA Localization Sequences within Its Open Reading Frame. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, *100* (23), 13368–13373. https://doi.org/10.1073/pnas.1835727100.

(18) Thio, G. L.; Ray, R. P.; Barcelo, G.; Schüpbach, T. Localization of Gurken RNA in Drosophila

Oogenesis Requires Elements in the 5' and 3' Regions of the Transcript. *Dev. Biol.* **2000**, *221* (2),

435–446. https://doi.org/10.1006/dbio.2000.9690.

(19) Long, R. M.; Gu, W.; Lorimer, E.; Singer, R. H.; Chartrand, P. She2p Is a Novel RNA-Binding

Protein That Recruits the Myo4p-She3p Complex to ASH1 mRNA. *EMBO J.* **2000**, *19* (23), 6592–

6601. https://doi.org/10.1093/emboj/19.23.6592.

(20) Münchow, S.; Sauter, C.; Jansen, R. P. Association of the Class V Myosin Myo4p with a Localised

Messenger RNA in Budding Yeast Depends on She Proteins. *J. Cell Sci.* **1999**, *112 (Pt 10)*, 1511–

1518.

(21) Bertrand, E.; Chartrand, P.; Schaefer, M.; Shenoy, S. M.; Singer, R. H.; Long, R. M. Localization of

ASH1 mRNA Particles in Living Yeast. *Mol. Cell* **1998**, *2* (4), 437–445.

https://doi.org/10.1016/s1097-2765(00)80143-4.

(22) Böhl, F.; Kruse, C.; Frank, A.; Ferring, D.; Jansen, R. P. She2p, a Novel RNA-Binding Protein

Tethers ASH1 mRNA to the Myo4p Myosin Motor via She3p. *EMBO J.* **2000**, *19* (20), 5514–5524.

https://doi.org/10.1093/emboj/19.20.5514.

(23) Heym, R. G.; Zimmermann, D.; Edelmann, F. T.; Israel, L.; Ökten, Z.; Kovar, D. R.; Niessing, D. In

Vitro Reconstitution of an mRNA-Transport Complex Reveals Mechanisms of Assembly and Motor

Activation. *J. Cell Biol.* **2013**, *203* (6), 971–984. https://doi.org/10.1083/jcb.201302095.

(24) Takizawa, P. A.; Vale, R. D. The Myosin Motor, Myo4p, Binds Ash1 mRNA via the Adapter

Protein, She3p. *Proc. Natl. Acad. Sci.* **2000**, *97* (10), 5273–5278.

(25) Chartrand, P.; Meng, X. H.; Huttelmaier, S.; Donato, D.; Singer, R. H. Asymmetric Sorting of

Ash1p in Yeast Results from Inhibition of Translation by Localization Elements in the mRNA. *Mol.*

Cell **2002**, *10* (6), 1319–1330. https://doi.org/10.1016/s1097-2765(02)00694-9.

(26) Long, R. M.; Singer, R. H.; Meng, X.; Gonzalez, I.; Nasmyth, K.; Jansen, R. P. Mating Type

Switching in Yeast Controlled by Asymmetric Localization of ASH1 mRNA. *Science* **1997**, *277*

(5324), 383–387. https://doi.org/10.1126/science.277.5324.383.

(27) Sil, A.; Herskowitz, I. Identification of Asymmetrically Localized Determinant, Ash1p, Required

for Lineage-Specific Transcription of the Yeast HO Gene. *Cell* **1996**, *84* (5), 711–722.

https://doi.org/10.1016/s0092-8674(00)81049-1.

(28) Takizawa, P. A.; Sil, A.; Swedlow, J. R.; Herskowitz, I.; Vale, R. D. Actin-Dependent Localization

of an RNA Encoding a Cell-Fate Determinant in Yeast. *Nature* **1997**, *389* (6646), 90–93.

https://doi.org/10.1038/38015.

(29) Shepard, K. A.; Gerber, A. P.; Jambhekar, A.; Takizawa, P. A.; Brown, P. O.; Herschlag, D.;

DeRisi, J. L.; Vale, R. D. Widespread Cytoplasmic mRNA Transport in Yeast: Identification of 22

Bud-Localized Transcripts Using DNA Microarray Analysis. *Proc. Natl. Acad. Sci. U. S. A.* **2003**,

100 (20), 11429–11434. https://doi.org/10.1073/pnas.2033246100.

(30) Jambhekar, A.; McDermott, K.; Sorber, K.; Shepard, K. A.; Vale, R. D.; Takizawa, P. A.; DeRisi, J.

L. Unbiased Selection of Localization Elements Reveals Cis-Acting Determinants of mRNA Bud

Localization in Saccharomyces Cerevisiae. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102* (50), 18005–

18010. https://doi.org/10.1073/pnas.0509229102.

(31) Singh, N.; Blobel, G.; Shi, H. Hooking She3p onto She2p for Myosin-Mediated Cytoplasmic

mRNA Transport. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112* (1), 142–147.

https://doi.org/10.1073/pnas.1423194112.

(32) Xayaphoummine, A.; Bucher, T.; Isambert, H. Kinefold Web Server for RNA/DNA Folding Path

and Structure Prediction Including Pseudoknots and Knots. *Nucleic Acids Res.* **2005**, *33* (Web

Server issue), W605-610. https://doi.org/10.1093/nar/gki447.

(33) Kertesz, M.; Wan, Y.; Mazor, E.; Rinn, J. L.; Nutter, R. C.; Chang, H. Y.; Segal, E. Genome-Wide

Measurement of RNA Secondary Structure in Yeast. *Nature* **2010**, *467* (7311), 103–107.

https://doi.org/10.1038/nature09322.

(34) Tutucci, E.; Maekiniemi, A.; Snoep, J. L.; Seiler, M.; van Rossum, K.; van Niekerk, D. D.; Savakis,

P.; Zarnack, K.; Singer, R. H. *Cyclin CLB2 mRNA Localization Determines Efficient Protein*

Synthesis to Orchestrate Bud Growth and Cell Cycle Progression; preprint; Cell Biology, 2022.

https://doi.org/10.1101/2022.03.01.481833.

(35) Frankel, A. D. If the Loop Fits. *Nat. Struct. Biol.* **1999**, *6* (12), 1081–1083.

https://doi.org/10.1038/70002.

(36) Takizawa, P. A.; DeRisi, J. L.; Wilhelm, J. E.; Vale, R. D. Plasma Membrane Compartmentalization

in Yeast by Messenger RNA Transport and a Septin Diffusion Barrier. *Science* **2000**, *290* (5490),

341–344. https://doi.org/10.1126/science.290.5490.341.

(37) Müller, M.; Richter, K.; Heuck, A.; Kremmer, E.; Buchner, J.; Jansen, R.-P.; Niessing, D.

Formation of She2p Tetramers Is Required for mRNA Binding, mRNP Assembly, and Localization.

RNA N. Y. N **2009**, *15* (11), 2002–2012.

(38) Lorenz, R.; Bernhart, S. H.; Höner Zu Siederdissen, C.; Tafer, H.; Flamm, C.; Stadler, P. F.;

Hofacker, I. L. ViennaRNA Package 2.0. *Algorithms Mol. Biol. AMB* **2011**, *6*, 26.

https://doi.org/10.1186/1748-7188-6-26.

(39) Reuter, J. S.; Mathews, D. H. RNAstructure: Software for RNA Secondary Structure Prediction and

Analysis. *BMC Bioinformatics* **2010**, *11*, 129. https://doi.org/10.1186/1471-2105-11-129.

(40) Zuker, M. Mfold Web Server for Nucleic Acid Folding and Hybridization Prediction. *Nucleic Acids*

Res. **2003**, *31* (13), 3406–3415. https://doi.org/10.1093/nar/gkg595.

(41) Seiler, M.; Mehrle, A.; Poustka, A.; Wiemann, S. The 3of5 Web Application for Complex and

Comprehensive Pattern Matching in Protein Sequences. *BMC Bioinformatics* **2006**, *7*, 144.

https://doi.org/10.1186/1471-2105-7-144.

(42) Schrödinger, L.; DeLano, W. PyMOL. **2020**.

(43) Kumar, M.; Michael, S.; Alvarado-Valverde, J.; Mészáros, B.; Sámano-Sánchez, H.; Zeke, A.;

Dobson, L.; Lazar, T.; Örd, M.; Nagpal, A.; Farahi, N.; Käser, M.; Kraleti, R.; Davey, N. E.;

Pancsa, R.; Chemes, L. B.; Gibson, T. J. The Eukaryotic Linear Motif Resource: 2022 Release.

Nucleic Acids Res. **2022**, *50* (D1), D497–D508. https://doi.org/10.1093/nar/gkab975.

(44) Waterhouse, A. M.; Procter, J. B.; Martin, D. M. A.; Clamp, M.; Barton, G. J. Jalview Version 2--a

Multiple Sequence Alignment Editor and Analysis Workbench. *Bioinforma. Oxf. Engl.* **2009**, *25* (9),

1189–1191. https://doi.org/10.1093/bioinformatics/btp033.

(45) Engel, S. R.; Dietrich, F. S.; Fisk, D. G.; Binkley, G.; Balakrishnan, R.; Costanzo, M. C.; Dwight, S.

S.; Hitz, B. C.; Karra, K.; Nash, R. S.; Weng, S.; Wong, E. D.; Lloyd, P.; Skrzypek, M. S.;

Miyasato, S. R.; Simison, M.; Cherry, J. M. The Reference Genome Sequence of Saccharomyces

Cerevisiae: Then and Now. *G3 Bethesda Md* **2014**, *4* (3), 389–398.

https://doi.org/10.1534/g3.113.008995.

(46) Engel, S. R.; Wong, E. D.; Nash, R. S.; Aleksander, S.; Alexander, M.; Douglass, E.; Karra, K.;

Miyasato, S. R.; Simison, M.; Skrzypek, M. S.; Weng, S.; Cherry, J. M. New Data and

Collaborations at the Saccharomyces Genome Database: Updated Reference Genome, Alleles, and

the Alliance of Genome Resources. *Genetics* **2022**, *220* (4), iyab224.

https://doi.org/10.1093/genetics/iyab224.

(47) Darty, K.; Denise, A.; Ponty, Y. VARNA: Interactive Drawing and Editing of the RNA Secondary

Structure. *Bioinforma. Oxf. Engl.* **2009**, *25* (15), 1974–1975.

https://doi.org/10.1093/bioinformatics/btp250.

Figures Legends

Figure 1. The SRS consensus model describes all four *ASH1* SRS elements. A Left, conformational
prediction of $E3_{1771-1821}$ by Kinefold displays the asymmetric bulge as a strand of 5 nt ('Lbulge') juxtaposed
with a shorter strand ('Sbulge')(both marked in red). Functionally relevant cytosines are boxed in green.
The asymmetric bulge and the lengths of Lbulge and Sbulge are marked in red, previously known functional
motifs in dark green. Right, the conformation of E3 referred to its crystal structure bound to She2p/3p
(**PDB: 5NEW**) displayed an rearrangement of the asymmetric bulge, in a ratio (Lbulge/Sbulge of
6:3)(uncommon base-pairs ignored in counting but dashed in blue), with sequence motifs adapted to
analogous positions in E2A, E2B and E3 (light green). **B** Left, Kinefold prediction and 180° rotation of
$E2A_{1109-1185}$ revealed an asymmetric bulge with the same 5:2 ratio as in E3. Right, the high degree of
sequence similarity (**Supplementary Fig. 1**) enabled a modeling in tight accordance to the E3 conformation
of the crystal structure that even comprised the uncommon base-pairing C_{1122}-C_{1174} and the A_{1123}-A_{1169} base-
stacking (dashed in blue). **C** Left, the predicted conformation of $E2B_{1263-1320}$ after 180°rotation showed
internal repeats (left, boxed in orange) that had to be aligned to adapt the consensus conformation (right)
that Kinefold also predicted but with 28% gain of energy content (-10.3 kcal/mol vs.-7.4 kcal/mol). **D** SRS
element $E2B_{623-701}$ after Kinefold prediction and 180°rotation (left) and after adaption (right). **E** Scheme of
the common features of all four modeled SRS elements, projected on $E3_{1771-1821}$. An asymmetric bulge(red)
is framed by two inverted repeats (yellow), with sequence motifs indicated in green. **F** Schematic
presentation of the SRS consensus model with its features displayed in the two configurations (IR1 boxed
in grey, IR2 boxed in blue, Lbulge and Sbulge in red, sequence motifs in green). The SRS consensus model
is characterized by a conformational feature arrangement that is identical in both configurations (middle),
but with differing linear order of features at sequence level (left). As a result, the positioning of sequence
start and end differ between the configurations in the conformational view (middle). Right, derived scheme
of the SRS consensus model that unifies the conformational arrangement of both configurations.

Figure 2. The modeling of the SRS consensus involves choice of appropriate conformation prediction
tool, rotation and mutual feature alignment of the four SRS elements in *ASH1*. The overview
demonstrates the effects of the three steps of modeling on the extended sizes of the *ASH1* SRS elements as
they were characterized before. Whereas the initially known conformations displayed a huge degree of
heterogeneity (**A, D, H, L**), the conformation prediction tool Kinefold tolerated asymmetric bulges in each
SRS element (**B, E, I, M**). For E3, using the published crystal structure as a blue print, its conformational
prediction (**B**) was adapted, with altered conformation of the asymmetric bulge (resulting in atypical base
pairings, marked in red, **C**). Its final asymmetric feature is indicated by a red triangle, with the numbers of

nucleotides displayed for both, Lbulge and Sbulge. By later analyses, we could assign one of its functional

cytosines as being analogous to already known CGA sequence motifs with other SRS elements (indicated

in red). For E2A, E1 and E2B, it was the rotation of the initial predictions that harmonized the respective

conformations with E3 (**F, J, N**). For E2A, due to its remarkable sequence identity with E3, we could also

introduce the same atypical base pairings (marked in red, **G**). As mentioned before, E2B represented an

outlier: it was a alternative conformation prediction that could be harmonized with E3, suggesting the need

of a further intramolecular rearrangement to align the inverted boxes (marked in yellow, **N** vs. **O**).

Figure 3. The configuration switch in E3 displays conserved function and surface plasmon resonance

analysis confirms the optimal C_1-C_2 distance of 6 nt in minimal constructs of E3. A Representative

results of electrophoretic mobility shift assays (EMSA) of triplicates to analyze complex formation of wild-

type and variant constructs of radioactive ASH1 $E3_{51nt}$ RNA with the She2p-She3p fusion protein (She2p$_{6-}$

$_{246}$-$(GGSGG)_2$-She3p$_{331-405}$) in increasing concentration (0.02 - 5 µM): Wildtype $E3_{1771-1821}$ (left), construct

with switch of $E3_{1771-1821}$ from config-I to config-II (middle) and control M6 mutational variant of $E3_{1771-}$

$_{1821}$ with decreased SRS function [8]. **B** Results of surface plasmon resonance analysis (SRP), performed in

triplicates to analyze the minimal E3 constructs and alterations in their C_1-C_2 distance. The minimal

construct $E3_{28nt}$wt was re-designed to create constructs with differing C1-C2 distances (5 and 7 nt) and was

incubated with She2p-She3p fusion protein (125 nM).

Figure 4. Computational screen for the SRS consensus model across 24 known She2p/3p interactors

results in 31 predictions of both configurations in 17 sequences. A Computational prediction of SRS

instances by feature screening of the SRS consensus model and evaluation of the resulting, predicted RNA

conformations (**Materials and Methods**). The screen was applied on 24 sequences for She2p/3p interaction

with varying experimental confidence (**Supplementary Tab. 1**). It identified 31 SRS predictions in 17

sequences, visualized as bar charts, with their coding sequences (rectangles) and 3' UTR extensions (200

nt, black lines). SRS predictions were indicated by their configurations (config-I: red, config-II: blue). **B**

For the SRS predictions of each configuration, the number of occurrences of the AA_L dinucleotide was

specified. **C** Schematic overview on the high-confidential set of SRS elements. Similar to the SRS elements

in *ASH1*, predictions in *IST2*, *EAR1* and *WSC2* matched published, well-characterized SRS elements (SRS

nomenclature, see *SRS collection*, **Tab. 2**) and could be arranged in the scheme of the SRS consensus model.

Color scheme is according to **Figure 1**, with further visualization of AA_L dinucleotides in the Lbulge and

adjacent regions, boxed in orange). Uncommon base pairings had been shown for the E3-She2p/She3p

crystal structure and were mapped on E2A that displayed high sequence similarity with E3 after rotation

(blue-dashed). Of note, similar uncommon base-pairings may also arise in E1 and *IST2*-SRS-2.

Figure 5. The configuration switch in the minimal SRS constructs of *EAR*-SRS2, *IST*-SRS2, *KSS1*-SRS1 and *TCB3*-SRS1 displays conserved SRS function in electrophoretic mobility shift assays. Representative results of electrophoretic mobility shift assays (EMSA) to analyze interaction of radioactive wild-type and designed RNA constructs with the She2p-She3p fusion protein (She2p$_{6-246}$-(GGSGG)$_2$-She3p$_{331-405}$) in increasing concentration (0.02 - 5 µM). All minimal constructs were designed switched in configuration (from config-II to config-I). Their final loop sequence GAAA referred to the artificial loop sequence of the E3 constructs in [8] that was shown not to modify SRS function. **A** EMSA experiments of well-known SRS elements. Left, *EAR1*-SRS2$_{1572-1621}$ (config-II) was studied as wildtype construct *EAR1*-SRS2$_{50nt}$wt and as config-switched, minimal construct *EAR1*-SRS2$_{26nt}$-switch (config-I). Right, *IST2*-SRS2$_{2716-2777}$ (config-II) was studied as wildtype construct *IST2*-SRS2$_{62nt}$wt and as config-switched, minimal construct *IST2*-SRS2$_{29nt}$-switch (config-I). **B** EMSA experiments of new, predicted SRS elements. Left, *KSS1*-SRS1 (config-II) was studied as wildtype construct *KSS1*-SRS1$_{72nt}$wt and as config-switched, minimal construct *KSS1*-SRS1$_{29nt}$-switch (config-I). Right, *TCB3*-SRS1 (config-II) was studied as config-switched, minimal construct *TCB3*-SRS1$_{38nt}$-switch (config-I).

Figure 6 *EAR1*-SRS2 is partly sensitive to alterations of its functional motifs but tolerates alterations in its conformational features and in the AA$_L$ dinucleotide of the Lbulge as confirmed for its minimal, config-switched construct by surface plasmon resonance analysis. A Schemes of the minimal constructs of *EAR1*-SRS2$_{28nt}$switch that were re-designed to create constructs with altered conformational features (indicated in red in conformations). The final loop sequence GAAA referred to the artificial loop sequence of the E3 constructs in [8] that was shown not to modify SRS function **B** Substitution scheme to analyze the relevance of cytosines in the sequence motifs of the SRS consensus model. Alterations are marked in red in substitution tables that refer to both, sequence positions and terminologies of resulting constructs. Since two cytosines (C$_{1601}$ and C$_{1602}$) precluded an unambiguous assignment of the C$_1$ position, both had to be considered. A third cytosine C$_{1600}$, marked by a star, was not considered in alterations due to its uncommonly short distance to C$_2$. **C** Substitution scheme to analyze the relevance of the AA$_L$ dinucleotide in the Lbulge. For each alterations, construct name (blue), conserved sequence in the Lbulge (black) and alterations (red) are displayed. **D** Results of surface plasmon resonance analysis (SRP), performed in triplicates to analyze the minimal EAR1-SRS2 constructs and alterations in their C$_1$-C$_2$ distance. E3 RNA was incubated with She2p-She3p fusion protein (125 nM).

794
795

Figure 1

Consensus modeling by mutual feature alignment using Kinefold

Start | Kinefold | Rotation | Feature alignment

A ASH1 E3

D ASH1 E2A

H ASH1 E1

L ASH1 E2B

Figure 3

A

B

Figure 4

A

B

C

Figure 5

A

B

Figure 6

EAR1-SRS2 constructs for SRP analysis

A

EAR1₃₁ₙₜsw

FlexA -> stem

plus 1bp stem

CG inversion

inversion 3bp

ΔC1ΔflexA

B

C

D

SRP: EAR1-SRS2 constructs vs. 125nM She2p/3p

801
802
803
804

TABLE 1 Overview on the four localization elements of *ASH1*. Positions and configuration classification are listed according to the SRS consensus motif

SRS name	SRS config I	SRS config II	SRS AA motif in Lbulge	SRS start	SRS end
E2		II	yes	631	686
E2A		II	yes	1118	1178
E2B		II	yes	1275	1312
E3	I		yes	1774	1817

812
813
814
815

TABLE 2 Overview on the *SRS instance collection.* The columns list info at transcripts and SRS instance level; SRS instance are classified according to the SRS consensus motif (including positions and configuration); the last two columns contain information to the identification of SRS elements in fragment size or as a detailed description in previous publications and this study

mRNA #	mRNA name	mRNA gene ID	SRS instance also known as	SRS instance collection name (*)	SRS start	SRS end	SRS config I	SRS config II	SRS as terminal element (***)	SRS with AA$_L$ motif	SRS fragment reference	SRS element reference
1	ASH1	YKL185W	E1	ASH1-E1	631	686		II		AA	several	several
			E2A	ASH1-E2A	1118	1178		II		AA	several	several
			E2B	ASH1-E2B	1275	1312		II	terminal	AA	several	several
			E3, U$_{min}$	ASH1-E3	1774	1817	I			AA	several	several
2	CLB2	YPR119W		CLB2-SRS1	1111	1145		II		AA		Tutucci et al
3	CPS1	YJL172W		CPS1-SRS1	599	636		II		AA		
				CPS1-SRS2	1684	1738		II		AA		
4	DNM1	YLL001W		DBM1-SRS1	262	302	I					
				DBM1-SRS2	1155	1189		II				
5	EAR1	YMR171C		EAR1-SRS1	110	167		II		AA		
				EAR1-SRS2	1578	1614		II	terminal	AA	Olivier et al	Olivier et al this study
6	EGT2	YNL327W		EGT2-SRS1	2194	2235		II		AA		
7	ERG2	YMR202W		ERG2-SRS1	261	314		II				
				ERG2-SRS2	338	380	I			AA		
8	IRC8	YJL051W		IRC8-SRS1	826	870		II		AA		
9	IST2	YBR086C		IST2-SRS1	2597	2626		II		AA		
				IST2-SRS2	2727	2768		II	terminal	AA	Olivier et al	Olivier et al this study
10	KSS1	YGR040W		KSS1-SRS1	730	781		II		AA		this study
11	MET4	YNL103W		MET4-SRS1	991	1051		II				
12	MMR1	YLR190W		MMR1-SRS1	548	590		II				
13	MTL1	YGR023W		MTL1-SRS1	1581	1651	I					
14	TCB2	YNL087W		TCB2-SRS1	3123	3175	I			AA		
				TCB2-SRS2	3556	3592		II		AA		
15	TCB3	YML072C		TCB3-SRS1	1603	1642	I			AA		this study
				TCB3-SRS2	2134	2173		II		AA		
16	TPO1	YLL028W		TPO1-SRS1	175	240		II		AA		
				TPO1-SRS2	461	527	I					
				TPO1-SRS3	774	822		II				
				TPO1-SRS4	1818	1862	I					
17	WSC2	YNL283C	WSC2N	WSC2-N	1277	1327	I			AA		Jambhekar et al
			WSC2C	WSC2-C	1326	1373		II		AA		Jambhekar et al

Remarks

SRS instances studies in this work are indicated in **red**.

(*) <u>SRS names in SRS collection</u> previously existing SRS terminologies are kept
in case of non-existing previous terminologies, the nomenclature contains "-SRS" suffixes (**)

(**) <u>Numbering with suffixes</u> the numbering follows their positions in 5'-3' RNA sequence.
SRS elements that were discovered later, are numbered with subsequent numbers

(***) <u>SRS as terminal element</u> indicates such SRS instances that reside at the terminal region of a RNA stem fold

824