Title: Negative impact of mild arid conditions on a rodent revealed using a physiological

approach in natura

Authors : Hamilcar Keilani[1*], Nico Avenant[2], Pierre Caminade[1], Neville Pillay[3], Guila

Ganem[1,3]

1.   ISEM, Univ Montpellier, CNRS, IRD, Montpellier, Occitanie, France

2.   Department of Mammalogy, National Museum and Centre for Environmental

Management, University of the Free State, Bloemfontein, Free State, South Africa

3.   School of Animal, Plant and Environmental Science, University of the

Witwatersrand, Wits, Gauteng, South Africa

* Corresponding author; e-mail : hamilcar.keilani@umontpellier.fr

Open research statement

Data will be available from the Dryad repository (TBA)

Keywords

Aridity, blood metabolites, ecophysiology, liver, local adaptation, nutrition, *Rhabdomys*

Abstract

1.   Understanding how organisms respond to seasonal variations in their

environment can be a window to their potential adaptability, a classical problem in

evolutionary ecology. In the context of climate change, inducing increased aridity and

disruption of seasonality, it is crucial to identify the potential for species responses.

2.   Here, the physiological response to food and water shortage during seasonally

dry conditions were investigated. We studied populations of two rodent species of the

genus *Rhabdomys*, one arid and one mesic, in a semi-arid zone where their range overlap

in South Africa. We measured blood concentrations of physiologically relevant

metabolites at the onset and the end of the dry season.

3.   We found similar shifts in blood metabolite levels in the semi-arid populations of

the two species, indicating malnutrition consistent with the observed degradation of

habitat quality between the start and the end of the dry season. Furthermore, regardless

of the period, differences between the two species in blood metabolite concentrations (ex:

amylase, sodium, alkaline phosphatase) were observed, suggesting contrasting diets and

water conservation abilities.

4.   Overall, we show that, as seasonal dry conditions worsen, organisms are

increasingly affected by reduced food availability, and local adaptation to arid

conditions may provide the arid species with an advantage to cope with semi-arid

conditions. Our study suggests that even mild arid conditions could have a negative

impact and questions resilience of animals to harsher arid conditions.

Introduction

Organisms are constantly exposed to a wide range of environmental variations, including short-term changes during their lifetime and longer-term changes across generations (Lopez-Maury et al., 2008). In the coming decades, extreme and more frequent droughts are predicted by most projections, resulting from changes in precipitation patterns, increased temperature, and evaporation (Naumann et al., 2018). These changes will induce great challenges to organisms. Under such circumstances, divergent selection can lead to the evolution of local adaptations where, in a given environment, resident phenotypes outperform nonresident ones (Kawecki & Ebert, 2004).

Local ecological adaptation can be described in several different and complementary ways, such as through its genetic basis, transcriptomic, behavioral, or physiological aspects. Adaptive evolution *sensu stricto* occurs when the genetic constitution of a population changes because of natural selection (Merilä & Hendry, 2013). In contrast, local adaptation *sensu lato*, also considers concepts such as phenotypic plasticity. It is usually assessed using comparative population or species fitness estimates under novel environmental conditions (Webster & Reusch, 2017). The response to local environmental conditions is a major factor in the generation and maintenance of biodiversity, impacting population dynamics, biogeographical ranges, and species interactions (Blanquart et al., 2013; Post et al., 2009). Specifically, physiological limits can drive and be driven by evolution, shape species distributions, and define species response capacities to future climate change, directly impacting risks of extinction (Somero, 2012). For example, adaptive variation in lethal temperature (LT_{50}) has been established in marine invertebrates inhabiting different latitudinal niches (Somero, 2010).

Studies of different physiological systems, such as the cardiovascular (Seebacher et al., 2005,

Berkel & Cacan, 2022) or digestive systems (Naya et al., 2011), in response to different

abiotic conditions, can improve our understanding of the causal mechanisms of responses of

organisms to climate variation (Seebacher & Franklin, 2012). Biotic factors, such as sex, age,

diet, strain, or breeding status also influence individual responses (McClure, 1999). With a

focus on influence of air temperature, water availability and energy resources on an

organism's physiological state, research in physiological ecology has addressed how variation

in physiology evolves and is maintained in relation to environmental conditions (Feder &

Block, 1991).

As a result of climate change, organisms can be faced with warmer and drier environmental

conditions (Parmesan et al., 2000), likely to impact their capacity to maintain homeostasis

(Fuller et al., 2016). In dry environments, food and water can be scarce, generating strong

selection on physiological attributes maximizing energy and water availability. Habitat

generalist and specialist taxa may respond differently to environmental changes. Desert

specialists exhibit specific physiological and/or behavioral traits, resulting from genetic

adaptations to the environment, allowing them to continuously endure or evade periods of

resource restriction (Rocha et al., 2021). In contrast, generalist species may seek to escape the

effects of aridity through temporary avoidance (Abraham et al., 2019) or plastic adjustments

during unfavorable seasons (Kobbe et al., 2011). Compared to mesic species, arid species

have lower basal metabolic rates and evaporative water loss (Muñoz-Garcia et al., 2022),

adopt a more flexible diet (Tshikae et al., 2013), and cope better with prolonged droughts

(Boyers et al., 2021).

Analysis of blood metabolites has successfully been used to investigate kidney and liver

physiology of wild populations (e.g. Al-Eissa et al., 2012). The kidney and the liver both

contribute to homeostasis in periods of food and water scarcity; indeed, the kidney plays a

crucial role in maintaining osmotic balance and the liver is involved in dietary metabolism.

Water loss can also mechanically induce high concentrations of some blood markers and

impact organism's capacity to evacuate products of metabolic waste, such as urea, from the

blood (Ostrowski et al., 2006). Hence, dry conditions can put a strain on these organs and be

detrimental to survival and reproduction.

In our study, potential impact of increased aridity was addressed by investigating the

physiological consequences of seasonal variation in dry conditions, in two species of the

diurnal African striped mouse genus *Rhabdomys*. The two study species, *R.bechuanae* and

R.dilectus dilectus, have different environmental niches (du Toit et al. 2012, Meynard et al.

2012). Throughout most of its distribution, *R.d.dilectus* occurs in mesic habitats with

vegetation cover and nests in dense grass, while *R.bechuanae* is found in semi-arid and arid

habitats, thriving predominantly in sparsely vegetated areas and nesting in bushes (Dufour et

al., 2015; Dufour et al., 2019). Moreover, differences in morphology or behavior suggest

adaptation to dry conditions in *R.bechuanae* (Ganem et al., 2020; Dufour et al., 2019). Here

we address both physiological response to seasonal dry conditions and divergence in

populations of the two species in the same bioclimatic region.

Indeed, at the edge of their distributions in central South Africa, the two species inhabit a

semi-arid region within which they occur either as parapatric or sympatric populations. This

semi-arid zone, the expansion and contraction of which depends on land use and precipitation

(Lian et al., 2021), has experienced a recent trend towards desiccation (Jury, 2021). Such

drying conditions could generate selective pressures, additional to those already experienced

by the semi-arid populations of the two species.

In the Succulent Karoo of South Africa, where the average daily precipitation during the dry

season is 0.26 mm *versus* 0.78 mm during the wet season, the probability of survival of adult

R. pumilio were related to their physiological response at the start of the dry season (Schoepf

et al., 2017a). Schoepf and collaborators found higher serum concentrations of albumin,

glucose, potassium, and lower concentrations of globulin, in animals that survived compared

to those that did not survive the dry season. At the peak of the dry season, *R.pumilio* had

lower concentrations of glucose and phosphorus and higher concentrations in globulin and

urea nitrogen if they survived the dry season. These results strongly suggest that such markers

could be good indicators of individual fitness.

While many studies have harnessed physiology, genomics, and transcriptomics to highlight

potential adaptive interspecific or interpopulational contrasts, studies under common natural

environmental conditions are lacking (Rocha et al., 2021). These approaches provide a mean

to disentangle the roles of interpopulational or interspecific variation from extrinsic

environmental factors in shaping phenotypic variation, taking advantage of the natural

conditions experienced in the field, including their complexity, with little human involvement.

This study took place in a natural semi-arid environment and asked how seasonal variation in

dry conditions influenced the physiological responses of the two species. We compared body

condition, blood concentrations of markers of kidney and liver functions and habitat

characteristics of parapatric populations of *R.bechuanae* and *R.d.dilectus* at the start *versus*

the end of the dry season. First, we expected the habitat available for the mice would be drier

at the end compared to the onset of the dry season, inducing a reduction of food and water,

impairing liver and kidney function, and impacting body condition. Second, assuming local

adaptive plasticity, we hypothesized that both species would be able to adjust their

physiological responses to seasonal changes in dry conditions. Third, we expected

R.bechuanae, having evolved in arid environments, would perform better in semi-arid

conditions and be closer to the expected local optimal response to increased dry conditions

than *R.d.dilectus* (having evolved in mesic areas), as found for *Mus musculus* (Bittner et al.,

2021). Alternatively, if both species evolved specific adaptations to their semi-arid

environment, we expected no species differences (**Table 1**).

Materials and Methods

Ethics statement

Permits to sample and handle animals in the field were obtained from the DESTEA of the

Free State, the North West DEDECT and Northern Cape DENC (respectively,

n°202110000007994, NW 38956/04/2022, FAUNA 0389/2022). Animal handling was

performed under ethical clearance from the Languedoc-Roussillon ethical committee for

animal experimentation (n°2022041512256467 v5).

Choice of sampling periods and sites

We selected six study sites based on their geographical position and their aridity index (AI).

The AI calculation was based on Thornthwaite method (Thornthwaite, 1948; see

Supplementary Material for full formula) using rainfall and temperature data collected from

2010-2021 (South African Weather Service).

We selected parapatric populations of the two species in a semi-arid region of South Africa

(0.2 < AI < 0.5). In this region, a hot and wet season occurs from roughly December to

February, and a cool dry season from May to September. We sampled six sites at the onset

(May 2022) of the dry season: Barberspan Bird Sanctuary, Benfontein Nature Reserve,

Bloemhof Dam Nature Reserve, Gariep Dam Nature Reserve, Kalkfontein Dam Nature

Reserve and Wolwespruit Nature Reserve (**Figures 1&3, Table 2**). We then resampled four

of these sites at the end of the dry season (September 2022). Two sites (Barberspan and

Benfontein) were excluded from the September sampling due to extreme weather conditions.

Capture and field protocol

We used small mammal PVC traps (LxHxW= 29.6x7.5x7.5 cm), baited with a mixture of

oats, peanut butter, salt and sunflower seeds, and added a piece of cotton wool. They were

placed approximately every 15 m along roughly 150 to 300 m transects. Since *Rhabdomys* is

mostly diurnal, the traps were checked twice a day (8 AM and 3 PM). On the field, all trapped

striped mice were measured (body length and mass), their sex and breeding status (breeding

or non-breeding) sassessed based on external morphological metrics, and a 0.5 mm piece of

tail was collected and kept in 98 % ethanol for species identification. All striped mice were

marked with a unique ear-tag before their release at the trapping location unless they were

kept for the physiological study (i.e. kept in their trap with cotton and food).

Overall, trapping effort was 9688 trap nights (**Supplementary Table 1**). Out of 903 trapped

small mammals, 694 were *Rhabdomys*. For the physiological study, we selected adults,

avoiding related individuals as much as possible, by selecting mice trapped at least 100 m

apart, except for breeding pairs (a male and a female) that could be trapped in the same nest.

This distance was based on data obtained for *R. pumilio* indicating minimal relatedness (R=0

between males and R=0.06 between females) at this distance (Solmsen et al. 2012). Trapping

lasted 3 to 6 days per site, until we reached our target of 20 adult individuals.

Habitat

Vegetation composition

Earlier studies have shown inter-species differences in vegetation cover and structure

requirements (Dufour et al., 2015). Since habitat characteristics can vary locally, we

characterized the vegetation structure (i.e. grass *versus* woody vegetation). Around traps in

which a *Rhabdomys* was trapped, the percentage of the surface composed of dry grass, green

grass, dry bush, green bush, succulent plants, holes and uncovered surfaces was measured

within 2mx2m (4m²) and 10x10m (100m²) quadrats centered on the trap position. For each

4m² quadrat we considered four 1x1m subunits assessed using a 1x1m metal frame; the results

obtained for the four subunits were then averaged. Each 100m² quadrat was assessed

following a visual inspection of a 100m² zone around each trap. We characterized 236

quadrats of each type (100m² and 4m²).

Normalised Differential Vegetation Index

NDVI, the satellite imagery-based index informative of ground vegetation greenness, was

retrieved from the Copernicus Open Access Data Hub (Copernicus Sentinel-2 data, 2023). For

each site and sampling session, we used NDVI data available for the closest day to the

beginning of a sampling session, which was expected to represent the conditions experienced

by the mice at the time of capture **(Supplementary Table 3).**

**Blood extraction and assessment of blood metabolites concentrations**

A total of 273 adult mice were euthanized by means of cervical dislocation on their day of

capture, and a blood volume of at least 100 µL was collected in lithium-heparinized tubes by

cardiac puncture. The breeding status was confirmed for all individuals during dissection. The

whole liver, the left kidney, and the skull of every individual were also collected for a

complementary study. Because the two species could not be distinguished visually, the spleen

was also harvested from each mouse and subsequently analyzed for post-hoc species

identification using Cytochrome Oxidase I genotyping (as described in Ganem et al. 2020).

Levels of albumin, alkaline phosphatase, alanine aminotransferase, amylase, total bilirubin,

blood urea nitrogen, total protein, globulin, glucose, calcium, phosphorus, potassium, and

sodium were successfully measured in the whole blood of 257 individuals immediately after

blood collection (Abaxis VETSCAN V2 technology with the Comprehensive diagnostic

profile cartridge), following the manufacturer's instructions. The coefficient of variation

(CV%) of measured levels of these markers was determined from data obtained by analyzing

a given blood sample twice. Samples from six adult striped mice (5 *R.bechuanae*, 1

R.d.dilectus) were used for these calculations. For these six samples, the calculated CV

equaled 5.6 ± 1.9 %, which is consistent with the intra-test CV% provided in the

manufacturer's instruction manual for other mammal species (**Supplementary Table 4**).

Data preparation and analysis

All statistical tests were carried out using the R software (version 4.2.1).

Habitat

<u>Vegetation composition</u>

Principal Component Analysis (package FactoMineR version 2.9) was performed, including

the 7 variables measured within the $4m^2$ quadrats and within the $100m^2$ quadrats for each

successful trap (traps in which *Rhabdomys* was captured). In each case, over 80 % of the

variance was explained by the first five Principal Components (**Supplementary Tables 7 &**

9). As Multivariate normality was violated in our dataset, the coordinates of all traps on the

first 5 PCs were used as response variables of a *PERMANOVA* (package *vegan*, version 2.6-

4). *PERMANOVA* is a multivariate statistical inference tool using permutational algorithms

(Anderson, 2001). This equivalent to *MANOVA* operates in a distribution-free setting and is

robust to non-normality of residuals as well as dispersion heterogeneity, even in slightly

unbalanced designs (Anderson, 2013). We tested differences between sites nested within

sessions (start and end of dry season), with 10 000 permutations, to assess whether the sites

had similar enough habitat characteristics to be considered as suitable replicates in other

analyses.

<u>NDVI</u>

NDVI was averaged over a 60 m radius circular buffer for each trap and then compared across

sessions and sites using a GLM Analysis of Variance, with the site nested within session. To

meet *ANOVA* prerequisites, the response variable was log-transformed; the analysis was

followed by Tukey multiple comparisons of means post-hoc tests.

Body condition

We calculated an index of body condition (log mass/log length), to assess the state of each

individual's fat reserves (Schulte-Hostedde et al., 2005). To test whether breeding status, age,

habitat quality, increased dryness, and/or interspecific differences influence body condition of

individuals trapped in semi-arid locations, an *ANOVA* was used with body condition as a

response variable, site as a random factor and session, species, sex, session*species*sex,

breeding status (nested within sex), age class and NDVI as explanatory variables.

Assumptions of normality and homoscedasticity of residuals were checked using diagnostic

plots (*graphics* package v.3.6.2.). A Tukey multiple comparisons of means post-hoc test was

applied to determine which inter-level differences were driving the significance of each

factorial effect having more than two modalities.

Age classes

Trapped *Rhabdomys* were assigned to age categories: juveniles, subadults and adults based on

their general appearance, size and/or breeding status. Body length of trapped individuals

varied between 5 and 12 cm. The body length of adult individuals involved in the

physiological study varied between 7.7 and 12 cm. To distinguish between older and younger

adults, mass was plotted against body length (**Supplementary figure 8**) showing a curve

defined by an allometric growth equation with a steeper slope at the highest and lowest body

length values. Since bone growth can significantly alter the concentration of some

physiological markers, and growth rate varies with age/body length, we considered 4

length/age classes (Class A: [7.7:9 cm[; Class B : [9:10 cm[; Class C : [10:11 cm[; Class D

[11 cm:]).

Physiology

We performed a Principal Component Analysis to identify outliers in the data set that could

be potential measure errors. We identified twelve outliers that we removed from further

analyses. These outliers were animals from which we collected smaller volume of blood

samples than recommended for the Vetscan or were heavily hemolyzed and flagged as such in

the Vetscan analysis output. In addition, as total protein values were used by the Vetscan

analyzer to infer the Globulin (GLOB) fraction of blood protein (with GLOB=TP-ALB), we

considered this marker as not informative and removed this variable from our analyses.

Finally, for 199 out of 257 samples, blood concentrations of creatinine (CRE) were flagged as

being below the instrument's sensitivity threshold; we hence excluded CRE from subsequent

analyses, as this imprecision would have hindered analytical performance.

Due to covariance in variables such as blood physiological markers, multivariate statistical

techniques provide a unique insight into the main patterns and effects driving variation in the

data. Thus, we performed a multivariate analysis with 236 blood samples, 12 physiological

parameters as response variables, site as a random factor, and session, species, sex, breeding

status (nested within sex), body condition, age class, and NDVI as explanatory variables. As

with habitat, multivariate normality was violated in our physiological dataset, so

permutational multivariate analysis of variance (*PERMANOVA*) was performed, using 10 000

permutations. A backwards stepwise model selection was performed to retain the most

parsimonious model. The best-fitting model was determined by comparing all previous

models, from least to most complex, using the Corrected Akaike's Information Criterion

(AICc) for *PERMANOVA* (package *AICcPermanova* version 0.0.2). We then sought to

identify which of the 12 physiological markers drove significantly the between-group

differences evidenced with the *PERMANOVA*. Twelve univariate *ANOVA* tests were

performed (followed by backwards stepwise model selection), initially using the parameters

of the most parsimonious *PERMANOVA* model. Assumptions of normality and

homoscedasticity of residuals were checked using diagnostic plots. In the case of total

bilirubin, a permutation test had to be performed instead of the *ANOVA*. Through a backwards

stepwise model selection process, the final model retained was based on the Akaike's

Information Criterion (AIC). For all tests, the significance level (α) was set at 0.05.

Results

Variation in habitat characteristics between species and sessions

Vegetation composition

We compared the vegetation characteristics surrounding each successful trap with a

PERMANOVA, using the first 5 Principal Component coordinates (**Supplementary Tables 7**

& 9) of each trap as response variable, and site and session as explanatory variables. We

detected significant differences in overall habitat characteristics between sessions (4m² and

100m²: p<0.001), and sites nested within sessions (2x2m and 10x10m: p<0.001), indicating

clear site differences and seasonal effects on habitat structure (**Supplementary Figures 6 &**

7). Although a permutational dispersion test showed these patterns could be attributable in

part to differences in dispersion between sessions or sites (see **Supplementary Material**),

some general trends could be inferred from the data. Indeed, some differences between sites

amounted to differences between species. Quadrats from *R.d.dilectus* sites (Barberspan,

Bloemhof and Wolwespruit) tended to feature higher proportions of grass (mean 4m²: 83.99 ±

1.12 %, mean 100m² : 71.42 ± 1.40 %) than *R.bechuanae* sites (mean 4m²: 54.30 ± 2.57 %,

mean 100m²: 53.12 ± 1.84 %). Conversely, *R.bechuanae* sites (Benfontein, Gariep and

Kalkfontein) tended to feature higher proportions of bushes (mean 4m²: : 31.07 ± 2.37 %,

mean 100m²: : 27.35 ± 1.42%) than *R.d.dilectus* sites (mean 4m²: 7.33 ± 1.00 %, mean 100m²

: 10.49 ± 1.03 %). Furthermore, differences between sessions could be observed; quadrats

surveyed at the end of the dry season tended to feature more dry grass (mean 4m²: 60.82 ±

2.33 %, mean 100m²: 51.99 ± 1.87 %) than at its onset (mean 4m² : 50.75 ± 2.04 %, mean

100m²: 40.85 ± 1.39 %).

NDVI

The log-transformed NDVI of sites sampled at the two sessions (2 sites per species), extracted

from satellite data, was also analyzed between sites and sessions using *ANOVA*. Site, session

and the interaction between these two variables (all $p<0.001$) contributed significantly to

explain the NDVI variation (**Table 3**). A post-hoc Tukey multiple comparisons of means test

showed that all sites had a significantly lower NDVI in September compared to May (**Figure**

4), indicating a poorer state of the vegetation at the end of the dry season. Furthermore, while

these study locations were chosen in a region with similar semi-arid bioclimatic features,

Gariep Dam and Kalkfontein (where *R.bechuanae* was targeted) seemed to stand out as

having a lower NDVI than all other sites sampled in both May and in September, with

Wolwespruit and Bloemhof (where *R.d.dilectus* was targeted) having the highest NDVI

(**Figure 4**). Therefore, to take into account the impact of variations in NDVI between the two

species trapping locations we included site as a random factor in analyses addressing body

and physiological conditions.

Variation of Body condition

An *ANOVA* assessed the influence of session, species, breeding status, age class and NDVI on

body condition, including site as a random factor. The most parsimonious model included the

significant effects of session (p<0.05), breeding status (p<0.001) and the interaction between

the two (p<0.05), as well as the effect of NDVI verging on significance (p=0.066) and sex

(p=0.286, **Table 4**). Body condition was slightly higher at the end of the dry season than at its

onset, and non-breeding females had lower body condition than breeding females and non-

breeding males (post-hoc pairwise t-tests all p<0.05). The correlation between body condition

and NDVI was negative.

Variation in physiological responses

A *PERMANOVA* was performed using the distance matrix of 12 physiological markers as

response variable and session, species, sex, breeding status, body condition, age class, NDVI

and site (random factor) as explanatory variables (**Supplementary table 12**).

The most parsimonious model **(Table 5)** indicated an overall difference in physiological

parameter concentrations between *R.bechuanae* and *R.d.dilectus* (p=0.002), as well as an

influence of body condition (p=0.019), breeding status (nested within sex) (p=0.028), that

differed between the two species (interaction effect, p=0.022), and the interaction between sex

and species (p=0.020). Session (p=0.065), and sex (p=0.084) verged on overall significance

and were also included in the model, indicating that physiological markers overall tend to

shift between the start and end of the dry season, and to vary between sexes. Importantly, both

species showed the same pattern of variation between sessions (no significant species*session

interaction effect, p=0.498, **Supplementary Table 12**).

Six physiological markers showed significant differences between the start and the end of the

dry season (**Figure 5**). Blood concentrations of markers of nutrition and liver function

(albumin, potassium, alanine aminotransferase, alkaline phosphatase, and total bilirubin) were

lower at the end compared to the onset of the dry season (**Table 4**). Glucose also showed that

trend, but not significantly (p=0.0577). In contrast, globulin levels varied in the opposite

direction (higher at the end compared to the onset of the dry season).

The levels of four markers were significantly lower in *R.bechuanae* than in *R.d.dilectus* : total

bilirubin, alkaline phosphatase (ALP), sodium and potassium (**Table 6, Figure 6**). However,

amylase showed the opposite pattern between the species.

Regardless of species, ALP levels were negatively correlated with body condition, while the

levels of phosphorus exhibited the opposite pattern. Finally, the levels of glucose, ALP and

alanine aminotransferase (ALT) varied significantly with sex, tending to be lower in females

than in males (**Supplementary figure 9**).

Discussion

We asked how seasonal variation of dry conditions influenced the physiological response of

semi-arid parapatric populations of two sister species that evolved under distinct

environmental conditions (du Toit et al., 2012).

Our study took place in a relatively wet year (La Niña). During the El Niño Southern

Oscillation phenomenon, there is a general association between regional wetness and sea

surface temperatures in the neighbouring Atlantic and Indian Oceans, leading to wetter

episodes throughout Southern Africa during La Niña years (Nicholson & Selato, 2000).

Vegetation growth and senescence usually follow seasonal rainfall patterns; above-average

rainfall widens the temporal window for vegetation growth, improving habitat conditions at

the start of the typical dry season (Anyamba et al., 2002). Despite this, and consistent with our

predictions, milder dry conditions did alter habitat quality of our study sites (as attested by

shifts in NDVI and habitat structure).

Consistent with our prediction, we found strong evidence for physiological divergence

between the species. We also observed a tendency for seasonal variation of the physiological

response in both species related to energetic resource depletion at the end of the dry season,

supported by the shifts in the levels of 6 blood parameters between the onset and the end of

the dry season.

Water and nutritional stress are expected to impact the body condition of individuals (Fuller et

al., 2021). Interestingly, the body condition of sampled individuals was higher at the end of

the dry season. As expected, body condition varied with breeding status, and several

physiological markers varied significantly with body condition or sex, indicating their

association with physiological processes involved in body mass regulation and reproduction.

Reproduction in mammals has direct physiological costs, associated with increased energy,

protein and calcium demands and indirectly through compensatory effects such as reduction

in thermogenesis, immune function and physical activity (Speakman, 2008; Rintoul &

Bringham, 2014; Stawski & Rojas, 2016; Schoepf et al., 2017b). It follows that sex-specific

differences in reproductive investments are associated with contrasts in body weight, fat

distribution and energy metabolism (Chen et al., 2012). In support, our results indicate lower

levels of blood glucose in females compared with males. Moreover, albumin (ALB) and

alkaline phosphatase (ALP) were significantly elevated in males compared to females, also

attributable to the pleiotropic effects of sex hormones (Havill et al., 2004; Sullivan et al.,

2007).

Variation in ALP levels is linked to variation in soft tissue and/or bone growth (Walker et al.,

1990). As expected, ALP was also negatively correlated with body condition. Individuals with

the lowest body condition had lower phosphorus levels. Generally, high phosphorus levels can

be observed in dehydrated individuals in which osmolality is high; indeed, low phosphorus

levels related to survival in *R.pumilio* at the peak of the dry season, where water balance was

hardest to maintain (Schoepf et al., 2017). However, blood levels of phosphorus also depend

on its consumption and hence availability in the food (Berndt & Kumar, 2009). Phosphorus

reserve depletion in smaller-sized adults might therefore be consistent with the role of

phosphorus intake as a promoter of bone and soft tissue growth (Shapiro & Heaney, 2003).

Furthermore, higher needs for phosphorus in smaller adults might be hindered by reduced

access to food. Indeed, foraging activity tends to vary between individuals according to size

and breeding status, possibly resulting in larger breeding individuals gaining access to higher

quality food (Ebensperger & Hurtado, 2005).

Seasonal variation in physiology

Despite our study year being relatively wet, mild dry conditions impacted blood marker

concentrations in *R.bechuanae* and *R.d.dilectus*. These changes involved 6 out of 12 tested

physiological markers, and included markers expected to vary in periods of dietary restriction.

Contrary to our prediction, we did not find a significant interactive effect between session and

species, suggesting that the semi-arid populations of the two species were similarly affected

by seasonal dry conditions. However, a follow up study during an El Niño year may be

necessary before one can conclude.

Blood protein levels are generally good indicators of physiological condition (Tothova et al.,

2016). They comprise two distinct and major components: albumin (ALB) and globulins

(GLOB). In our study, ALB was significantly lower at the end than at the start of the dry

season. ALB is synthesized in the liver and involved in the transport of bilirubin and several

hormones. Blood ALB levels are less prevalent in malnourished individuals (Walker et al.,

1990), supporting the hypothesis of reduction of quality and/or availability of food at the end

of the dry season. The Nubian ibex *Capra nubiana*, a desert mammal, also displays low ALB

during the dry season (Al-Eissa et al., 2012), which could reflect poor nutritional condition,

kidney dysfunction leading to poor water balance and high osmolality, or liver dysfunction

causing low ALB synthesis. We found that GLOB was significantly higher at the end than at

the start of the dry season. Higher GLOB levels were also reported for *R.pumilio* individuals

that did not survive the dry season (Schoepf et al., 2017). GLOB is a family of molecules

including immunity proteins and blood protein carriers (including transferrin). An increase in

GLOB levels usually indicates elevation of immunoglobulins and hence could suggest

heightened immune activity. Greater antibody production occurs in response to a higher

prevalence of respiratory infections during the dry season, e.g. in the Nubian goat (Abdelatif

et al., 2009), and/or reduced protein nutritional status in young birds (Lochmiller et al., 1993).

An increase in blood GLOB concentrations may also reflect high levels of transferrin (Walker

et al., 1990), a major GLOB that may accumulate in the blood in cases of severe iron

deficiency (Kasvosve & Delanghe, 2002). Hence, elevated blood GLOB levels at the end of

the dry season may indicate either or both depletion in iron resource (transferrin) and higher

vulnerability to infection (immunoglobulin), both resulting from or being aggravated by

reduced access to food. Lower levels of alkaline phosphatase (ALP), a liver enzyme, at the

end of the dry season in *Rhabdomys*, is also evidence of malnutrition and deficiency in

essential nutrients (proteins, magnesium, zinc) (Yousef et al., 2002; Saraç & Saygili, 2007;

Ray et al., 2017). Alanine aminotransferase (ALT), another marker of liver function, also

occurred at significantly lower concentrations at the end compared to the start of the dry

season, indicating a reduction in liver activity (Walker et al., 1990) or malnutrition (Le

Couteur et al., 2010). Indeed, as environmental resources become scarce, essential nutrients,

such as pyridoxine (or vitamin B6), that constitute some ALT coenzymes, may be less

available, leading to a reduction of ALT blood levels (Vespasiani-Gentilucci et al., 2018).

Blood glucose, another major physiological marker of nutrition state, was also slightly

diminished at the end of the dry season, albeit not significantly. Fasting or low-energy feeding

results in lower blood glucose levels (Jensen et al., 2013).

Acute starvation also typically causes bilirubin (TBIL) to accumulate in the blood, as the

enzymatic process for conversion of heme into TBIL is stimulated (Thaler et al., 1972). Yet,

TBIL was lower at the end compared to the onset of the dry season, rejecting the starvation

hypothesis. Lower TBIL concentrations at the end of the dry season in *Rhabdomys* could stem

from low hemoglobin counts because the body produces fewer red blood cells than usual; low

hemoglobin can be induced by several factors, among which dietary factors such as iron

deficiencies (Clark, 2008). Iron deficient anemia is also supported by higher GLOB

concentrations observed at the end of the dry season. Infection can likewise be a cause of both

decreased TBIL levels (Zhao et al., 2019) and higher GLOB concentrations at the end of the

dry season. Finally, malnutrition and/or lack of potassium intake are among the main potential

causes of the reduction of its levels in the blood (Their, 1986), which could explain why they

were the lowest at the end of the dry season, when the available resources were lowest, in our

study.

Overall, our results indicate that seasonal variation of some physiological markers could

reflect reduced primary productivity throughout the dry season, which is coherent with our

observation of habitat degradation (lower NDVI) at the end of the dry season. Blood

concentrations of various markers seem to indicate difficulties in maintaining nutritional

functions and the necessity of a higher immune output, caused or exacerbated by food

restriction. Indeed, individuals undergoing malnutrition during dry conditions are expected to

show low blood glucose concentrations, and deficiencies in essential nutrients, leading to

pathologies such as anemia (Gordon et al., 1988). Indeed, animals foraging on soils

potentially poor in nutrients such as iron, zinc, and magnesium are more prone to malnutrition

(Graham, 1991; Gupta et al., 2008). Higher GLOB, lower TBIL and lower ALP blood

concentrations in our study might relate to such deficiencies.

We did not observe seasonal variation of sodium or blood urea nitrogen, blood markers of

kidney function or osmoregulation. Indeed, reduced environmental moisture at the end of the

dry season could have caused an overall increase in blood osmolality, increasing blood

concentrations of most biomarkers. Still, most markers exhibited a significant reduction in

blood concentrations. This suggests that, compared to food availability, seasonal patterns in

water availability may impose a milder physiological cost to the striped mice throughout the

dry season in this semi-arid environment during a La Niña episode.

The two species responded similarly to increased dryness, increasing body condition, a proxy

of fat reserves. Similarly, *R.pumilio* individuals tended to maintain their body condition stable

by reducing their energy expenditure and physical activity during periods of limited food

availability (Rimbach, Blanc et al., 2018; Rimbach, Jäger et al., 2018). The semi-arid

populations of *R.bechuanae* and *R.d.dilectus* could be displaying a similar behavioral strategy.

Interspecific differences

R.bechuanae is thought to have evolved in drier environments than *R.d.dilectus*. Moreover, in

the semi-arid region where they co-occur, we showed that *R.bechuanae* populations occupy

slightly drier habitats than *R.d.dilectus*. While body condition did not differ significantly

between species, some interspecific contrasts support our hypothesis that *R.bechuanae* may

have developed a better physiological capacity to cope with dry conditions than *R.d.dilectus*.

Among the markers exhibiting a significant difference between *R.bechuanae* and *R.d.dilectus*, all but amylase showed higher blood concentrations in *R.d.dilectus*. Amylase is positively correlated with digestive activity and reflects differences in feeding status or diet composition between the species (Hidalgo et al., 1999). Given the role of alpha-amylase in digestion, lower blood amylase levels in *R.d.dilectus* than *R.bechuanae* may indicate lower digestive activity due to reduced access to food, or lower starch content in the diet. Pajic et al. (2019) found a correlation between dietary starch content and the number of amylase gene copies in several mammalian genomes, even in different habitats and with different diets. Since amylase activity and amylase gene expression are directly correlated with amylase gene copy number (Arendt et al., 2014), further studies could elucidate the potential genetic basis of the different levels of amylase in our two study species.

Sodium (NA) and potassium (K) concentrations, which depend on both dietary and water intake, were higher in *R.d.dilectus* than in *R.bechuanae*. These differences, coupled with higher ALP and TBIL concentrations in *R.d.dilectus* compared to *R.bechuanae*, could suggest a better nutrition state for *R.d.dilectus* than *R.bechuanae*. However, higher levels of amylase in *R.bechuanae* cast doubt on this interpretation. Instead, lower levels of blood NA, K, ALP, and TBIL, would be more parsimoniously attributed to lower overall osmolality, perhaps due to better water balance, in *R.bechuanae*. A better overall ability of *R.bechuanae* to conserve body moisture could rely on physiological mechanisms such as more efficient excretion or osmoregulation, or through feeding on a diet optimizing water intake. We hypothesize that during the evolutionary history of *R.bechuanae*, selection pressures may have favored specific strategies allowing an overall more efficient water regulation compared to *R.d.dilectus*. These results, combined with the reduction in the levels of K, ALP and TBIL at the end of the dry season suggest the species are exposed to both water and nutrition stress.

We found relatively little evidence that the two species suffer differentially from lack of food

in their semi-arid environment during a dry season impacted by La Niña phenomenon.

Overall, energy levels, as indicated by blood glucose, and body condition, did not vary

between the species. Instead, it seems that the overall differences between the two species can

be accounted for by lower blood concentrations in *R.bechuanae*, and on average higher blood

concentrations in key metabolites in *R.d.dilectus,* possibly due to overall poorer water

regulation. High average blood amylase levels in *R.bechuanae* during the dry season might

indicate an interspecific difference in diet composition or food intake, although not protein-

energy malnutrition. Little is known about *R.bechuanae* and *R.d.dilectus* diets; the genus was

characterised as a generalist feeding on seeds and other plant material, and insects (Curtis &

Perrin, 1979). Some studies showed variation in diet composition within *Rhabdomys* genus

associated with variation in local resources (Taylor & Green, 1976). The two species could

have adjusted their diet to the respective local availabilities of different food resources within

their home ranges. For example, during the dry season and in the semi-arid zone, arthropod

biomass and diversity is affected by seasonal patterns in rainfall. Dalerum et al. (2017) found

lower arthropod trapping success and diversity during the dry season, in shrub habitats

compared to any other biomes, in Benfontein Game Reserve, one of our study sites. Thus,

compared to seeds, the reliability of arthropods as a food source during dry conditions could

be lower in shrublands, inhabited by *R.bechuanae* across its range and by both species in

parapatry, than in grasslands inhabited by *R.d.dilectus* in allopatry. Variation in seed

composition could also explain this interspecific difference in amylase concentrations. Indeed,

granivores prefer seeds that have higher protein content and lower secondary metabolites but

may be forced to take a wider range of seeds in dry areas than in mesic ones (Wilmer et al.,

2009). If the hypothesis of a relatively ancient adaptation of *R.bechuanae* to dry habitats is

true, it could be expected that it would also have adjusted its diet to the specific conditions

encountered in dry habitats. Within this ecosystem, *R.bechuanae* could forage more

efficiently compared with *R.d.dilectus*, which evolved in a more mesic environment; its diet

could also be richer in starch (such as containing more seeds/grains than leaves or fibre).

Higher starch intake could help maintaining water balance; starch is the best substrate for

metabolic water production in a dry environment (Adolph, 1964). This is consistent with the

other interspecific differences revealed in this study, as diet can also significantly affect

osmoregulation (Sabat et al., 2009).

Due to climate change, populations of many species experience changes in their habitats

affecting their fitness and inducing shifts in their geographical distribution (Parmesan et al.,

2006; Fuller et al., 2010). In some species, range shifts are expected to swell edge

populations, as marginal habitats grow more favourable. For instance, edge populations of

Wilsonia citrina hooded warblers have repeatedly expanded their ranges due to climate

change (Melles et al., 2011). In other species, the quality of marginal habitats occupied by

populations at the range edge decline in food availability or nutrient richness, making these

populations particularly vulnerable to climate change and inducing range contraction. For

example, the koala *Phascolarctos cinereus* experiences declines in population sizes and

habitat contraction due to droughts (Seabrook et al., 2011). As such, depending on their

phenotypic plasticity, edge populations may hold the key to the survival of a species in a

changing environment (Wu & Seebacher, 2022, Usui et al., 2023). *R.d.dilectus* edge

populations may experience the same situation in the semi-arid zone, an hypothesis that could

be tested addressing these populations dynamics, compared to *R.bechuanae* semi-arid

populations.

Conclusions

We found that two closely related species with different ecological characteristics displayed

similar seasonal negative shifts in blood concentrations of markers of nutrition and liver

function in spite of a relatively mild seasonal drying of their habitats. We hence expect the

physiological impact of harsher dry conditions, to affect the individuals' fitness even more

dramatically. The fact that the habitat of *R.bechuanae* studied populations was the driest at

both time points, combined with our findings concerning their overall better water balance

and nutrition than *R.d.dilectus*, point towards *R.bechuanae* coping better in the semi-arid

environment during the dry season. Though adaptation *per se* cannot be inferred from our

study, this comparative study considering populations of the two species in the same

biogeographic zone adds to the growing evidence that these cryptic species of striped mice

have diverged phenotypically. These findings show the ecological relevance of physiological

approaches in investigating the effect of climatic variation on organisms and question

resilience of core populations of arid species and edge populations of mesic species in the face

of predicted enhanced aridification.

Acknowledgements

We thank R.Nokha who provided substantial field work contribution to this project.

Constructive and insightful comments were provided by C.Smadja and C.Schradin helped

improve a previous version of this paper. Technical support by D.Greuet (INM), L.Paradis

(Geomatics, ISEM) and GenSeq platform. We are grateful to the rangers and personnel of

Barberspan Bird Sanctuary, Benfontein Nature Reserve, Bloemhof Dam Nature Reserve,

Gariep Dam Nature Reserve, Kalkfontein Dam Nature Reserve and Wolwespruit Nature

Reserve who facilitated our work in these locations. Funding: CNRS/NRF, OSU-OREME

Author contributions

G.G. conceived the idea and designed methodology with input from H.K; G.G., H.K., P.C.,

collected the data; H.K. analyzed the data with input from G.G.; H.K. wrote the first draft of

the manuscript with input from G.G.; N.P. and N.A. contributed critically to the manuscript,

secured funding for the project with G.G. and helped with field logistics. All authors approved

the final version of the manuscript.

Conflict of Interest statement

The authors declare they have no conflict of interest.

References

Abaxis, Inc. "VS2 Operator's Manual." (2009) Retrieved from https://www.zoetis.es/_locale-

assets/spc/vetscan-vs2-analizador.pdf

Abdelatif, Abdallah M., Mariam Y. Ibrahim, and Yahia Y. Hassan. "Seasonal variation in

erythrocytic and leukocytic indices and serum proteins of female Nubian goats." Middle-

Eastern Journal of Scientific Research, 4, no.3 (2009): 168-174. Retrieved from

https://idosi.org/mejsr/mejsr4(3)/9.pdf

Abraham, Joel O., Gareth P. Hempson, and A. Carla Staver. "Drought-response Strategies of

Savanna Herbivores." Ecology and Evolution 9, no. 12 (May 22, 2019): 7047–56.

https://doi.org/10.1002/ece3.5270.

Adolph, Edward F. "Desert Animals: Physiological Problems of Heat and Water. Knut

Schmidt-Nielsen." Physiological Zoology 37, no. 3 (July 1964): 338–39.

https://doi.org/10.1086/physzool.37.3.30152404.

AL Eissa, M S, Alkahtani Saad, S A Al Farraj, A Alarifi Saud, B Al Dahmash, and Al Yahya

Hamad. "Seasonal Variation Effects on the Composition of Blood in Nubian Ibex (Capra

Nubiana) in Saudi Arabia." African Journal of Biotechnology 11, no. 5 (January 16, 2012): 1283–86. https://doi.org/10.5897/ajb11.2004.

Anyamba, Assaf, Compton J. Tucker, and Robert Mahoney. "From El Niño to La Niña: Vegetation Response Patterns over East and Southern Africa during the 1997–2000 Period." Journal of Climate 15, no. 21 (November 1, 2002): 3096–3103. https://doi.org/10.1175/1520-0442(2002)015<3096:FENOTL>2.0.CO;2.

Arendt, Maja, Tove Fall, Kerstin Lindblad-Toh, and Erik Axelsson. "Amylase Activity Is Associated with AMY2B Copy Numbers in Dog: Implications for Dog Domestication, Diet and Diabetes." Animal Genetics 45, no. 5 (June 28, 2014): 716–22. https://doi.org/10.1111/age.12179.

Berkel, Caglar, and Ercan Cacan. "The Differential Influence of Ambient Temperature on Heart Rate in Male and Female Wild Pale-Throated Sloths (Bradypus Tridactylus)." Research Square Platform LLC, August 12, 2022. http://dx.doi.org/10.21203/rs.3.rs-1908943/v1.

Berndt, Theresa, and Rajiv Kumar. "Novel Mechanisms in the Regulation of Phosphorus Homeostasis." Physiology 24, no. 1 (February 2009): 17–25. https://doi.org/10.1152/physiol.00034.2008.

Bittner, Noëlle K. J., Katya L. Mack, and Michael W. Nachman. "Gene Expression Plasticity and Desert Adaptation in House Mice*." Evolution 75, no. 6 (January 23, 2021): 1477–91. https://doi.org/10.1111/evo.14172.

Blanquart, François, Oliver Kaltz, Scott L. Nuismer, and Sylvain Gandon. "A Practical Guide to Measuring Local Adaptation." Ecology Letters 16, no. 9 (July 15, 2013): 1195–1205. https://doi.org/10.1111/ele.12150.

Boyers, Melinda, Francesca Parrini, Norman Owen-Smith, Barend F. N. Erasmus, and Robyn

S. Hetem. "Contrasting Capabilities of Two Ungulate Species to Cope with Extremes of

Aridity." Scientific Reports 11, no. 1 (February 18, 2021). https://doi.org/10.1038/s41598-

021-83732-w.

Chen, Xuqi, Rebecca McClusky, Jenny Chen, Simon W. Beaven, Peter Tontonoz, Arthur P.

Arnold, and Karen Reue. "The Number of X Chromosomes Causes Sex Differences in

Adiposity in Mice." PLoS Genetics 8, no. 5 (May 10, 2012): e1002709.

https://doi.org/10.1371/journal.pgen.1002709.

Clark, Susan F. "Iron Deficiency Anemia." Nutrition in Clinical Practice 23, no. 2 (April

2008): 128–41. https://doi.org/10.1177/0884533608314536.

Curtis, B.A., and M.R. Perrin. "Food Preferences of the Vlei Rat (Otomys Irroratus) and the

Four-Striped Mouse (Rhabdomys Pumilio)." South African Journal of Zoology 14, no. 4

(January 1979): 224–29. https://doi.org/10.1080/02541858.1979.11447675.

Dalerum, Fredrik, J. Low de Vries, Christian W.W. Pirk, and Elissa Z. Cameron. "Spatial and

Temporal Dimensions to the Taxonomic Diversity of Arthropods in an Arid Grassland

Savannah." Journal of Arid Environments 144 (September 2017): 21–30.

https://doi.org/10.1016/j.jaridenv.2017.04.002.

Toit, Nina du, Bettine Jansen van Vuuren, Sonja Matthee, and Conrad A. Matthee. "Biome

Specificity of Distinct Genetic Lineages within the Four-Striped Mouse Rhabdomys Pumilio

(Rodentia: Muridae) from Southern Africa with Implications for Taxonomy." Molecular

Phylogenetics and Evolution 65, no. 1 (October 2012): 75–86.

https://doi.org/10.1016/j.ympev.2012.05.036.

Dufour, Claire M. S., Christine Meynard, Johan Watson, Camille Rioux, Simon Benhamou,

Julie Perez, Jurie J. du Plessis, Nico Avenant, Neville Pillay, and Guila Ganem. "Space Use

Variation in Co-Occurring Sister Species: Response to Environmental Variation or

Competition?" PLOS ONE 10, no. 2 (February 18, 2015): e0117750.

https://doi.org/10.1371/journal.pone.0117750.

Dufour, Claire M. S., Neville Pillay, Nico Avenant, Johan Watson, Etienne Loire, and Guila

Ganem. "Habitat Characteristics and Species Interference Influence Space Use and Nest-site

Occupancy: Implications for Social Variation in Two Sister Species." Oikos 128, no. 4

(November 15, 2018): 503–16. https://doi.org/10.1111/oik.05357.

Ebensperger, Luis A., and María J. Hurtado. "On the Relationship between Herbaceous Cover

and Vigilance Activity of Degus (Octodon Degus)." Ethology 111, no. 6 (May 31, 2005):

593–608. https://doi.org/10.1111/j.1439-0310.2005.01084.x.

Feder, M. E., and B. A. Block. "On the Future of Animal Physiological Ecology." Functional

Ecology 5, no. 2 (1991): 136. https://doi.org/10.2307/2389251.

Fuller, Andrea, Duncan Mitchell, Shane K. Maloney, and Robyn S. Hetem. "Towards a

Mechanistic Understanding of the Responses of Large Terrestrial Mammals to Heat and

Aridity Associated with Climate Change." Climate Change Responses 3, no. 1 (November 28,

2016). https://doi.org/10.1186/s40665-016-0024-1.

Fuller, Andrea, Duncan Mitchell, Shane K. Maloney, Robyn S. Hetem, Vinicius F. C.

Fonsêca, Leith C. R. Meyer, Tanja M. F. N. van de Ven, and Edward P. Snelling. "How

Dryland Mammals Will Respond to Climate Change: The Effects of Body Size, Heat Load

and a Lack of Food and Water." Journal of Experimental Biology 224, no. Suppl_1 (February

15, 2021). https://doi.org/10.1242/jeb.238113.

Fuller, Andrea, Terence Dawson, Brian Helmuth, Robyn S. Hetem, Duncan Mitchell, and

Shane K. Maloney. "Physiological Mechanisms in Coping with Climate Change."

Physiological and Biochemical Zoology 83, no. 5 (September 2010): 713–20.

https://doi.org/10.1086/652242.

Ganem, Guila, Claire Marie Soleil Dufour, Nico L. Avenant, Pierre Caminade, Seth J. Eiseb,

Christelle Tougard, and Neville Pillay. "An Update on the Distribution and Diversification of

Rhabdomys Sp. (Muridae, Rodentia)." Journal of Vertebrate Biology 69, no. 2 (July 7, 2020):

1. https://doi.org/10.25225/jvb.20013.

Gordon, G., A. S. Brown, and T. Pulsford. "A Koala (Phascolarctos Cinereus Goldfuss)

Population Crash during Drought and Heatwave Conditions in South-western Queensland."

Australian Journal of Ecology 13, no. 4 (December 1988): 451–61.

https://doi.org/10.1111/j.1442-9993.1988.tb00993.x.

Graham, Thomas W. "Trace Element Deficiencies in Cattle." Veterinary Clinics of North

America: Food Animal Practice 7, no. 1 (March 1991): 153–215.

https://doi.org/10.1016/s0749-0720(15)30816-1.

Gupta, Umesh C., Kening Wu, and Siyuan Liang. "Micronutrients in Soils, Crops, and

Livestock." Earth Science Frontiers 15, no. 5 (September 2008): 110–25.

https://doi.org/10.1016/s1872-5791(09)60003-8.

Havill, L.M, M.C Mahaney, and J Rogers. "Genotype-by-Sex and Environment-by-Sex

Interactions Influence Variation in Serum Levels of Bone-Specific Alkaline Phosphatase in

Adult Baboons (Papio Hamadryas)." Bone 35, no. 1 (July 2004): 198–203.

https://doi.org/10.1016/j.bone.2004.02.022.

Hidalgo, M.C, E Urea, and A Sanz. "Comparative Study of Digestive Enzymes in Fish with

Different Nutritional Habits. Proteolytic and Amylase Activities." Aquaculture 170, no. 3–4

(January 1999): 267–83. https://doi.org/10.1016/s0044-8486(98)00413-x.

Havill, L.M, M.C Mahaney, and J Rogers. "Genotype-by-Sex and Environment-by-Sex

Interactions Influence Variation in Serum Levels of Bone-Specific Alkaline Phosphatase in

Adult Baboons (Papio Hamadryas)." Bone 35, no. 1 (July 2004): 198–203.

https://doi.org/10.1016/j.bone.2004.02.022.

Jury, Mark R. "Spreading of the Semi-Arid Climate across South Africa." Journal of Water

and Climate Change 12, no. 8 (October 1, 2021): 3734–49.

https://doi.org/10.2166/wcc.2021.187.

Kasvosve, Ishmael, and Joris Delanghe. "Total Iron Binding Capacity and Transferrin

Concentration in the Assessment of Iron Status." Clinical Chemistry and Laboratory Medicine

40, no. 10 (January 29, 2002). https://doi.org/10.1515/cclm.2002.176.

Kawecki, Tadeusz J., and Dieter Ebert. "Conceptual Issues in Local Adaptation." Ecology

Letters 7, no. 12 (November 12, 2004): 1225–41. https://doi.org/10.1111/j.1461-

0248.2004.00684.x.

Kobbe, Susanne, Jörg U. Ganzhorn, and Kathrin H. Dausmann. "Extreme Individual

Flexibility of Heterothermy in Free-Ranging Malagasy Mouse Lemurs (Microcebus

Griseorufus)." Journal of Comparative Physiology B 181, no. 1 (August 18, 2010): 165–73.

https://doi.org/10.1007/s00360-010-0507-5.

Le Couteur, D. G., F. M. Blyth, H. M. Creasey, D. J. Handelsman, V. Naganathan, P. N.

Sambrook, M. J. Seibel, L. M. Waite, and R. G. Cumming. "The Association of Alanine

725 Transaminase With Aging, Frailty, and Mortality." The Journals of Gerontology Series A:

726 Biological Sciences and Medical Sciences 65A, no. 7 (May 24, 2010): 712–17.

727 https://doi.org/10.1093/gerona/glq082.

728 Lian, Xu, Shilong Piao, Anping Chen, Chris Huntingford, Bojie Fu, Laurent Z. X. Li,

729 Jianping Huang, et al. "Multifaceted Characteristics of Dryland Aridity Changes in a

730 Warming World." Nature Reviews Earth & Environment 2, no. 4 (March 9, 2021): 232–

731 50. https://doi.org/10.1038/s43017-021-00144-0.

732 Lochmiller, Robert L., Michelle R. Vestey, and Jon C. Boren. "Relationship between Protein

733 Nutritional Status and Immunocompetence in Northern Bobwhite Chicks." The Auk 110, no.

734 3 (July 1993): 503–10. https://doi.org/10.2307/4088414.

735 López-Maury, Luis, Samuel Marguerat, and Jürg Bähler. "Tuning Gene Expression to

736 Changing Environments: From Rapid Responses to Evolutionary Adaptation." Nature

737 Reviews Genetics 9, no. 8 (August 2008): 583–93. https://doi.org/10.1038/nrg2398.

738 Melles, S. J., M.-J. Fortin, K. Lindsay, and D. Badzinski. "Expanding Northward: Influence

739 of Climate Change, Forest Connectivity, and Population Processes on a Threatened Species'

740 Range Shift." Global Change Biology 17, no. 1 (December 1, 2010): 17–31.

741 https://doi.org/10.1111/j.1365-2486.2010.02214.x.

742 McClure, Diane E. "Clinical Pathology and Sample Collection in the Laboratory Rodent."

743 Veterinary Clinics of North America: Exotic Animal Practice 2, no. 3 (September 1999): 565–

744 90. https://doi.org/10.1016/s1094-9194(17)30111-1.

Merilä, Juha, and Andrew P. Hendry. "Climate Change, Adaptation, and Phenotypic

Plasticity: The Problem and the Evidence." Evolutionary Applications 7, no. 1 (January

2014): 1–14. https://doi.org/10.1111/eva.12137.

Meynard, Christine N., Neville Pillay, Manon Perrigault, Pierre Caminade, and Guila Ganem.

"Evidence of Environmental Niche Differentiation in the Striped Mouse (Rhabdomys Sp.):

Inference from Its Current Distribution in Southern Africa." Ecology and Evolution 2, no. 5

(April 12, 2012): 1008–23. https://doi.org/10.1002/ece3.219.

Muñoz-Garcia, Agustí, Miriam Ben-Hamo, Shai Pilosof, Joseph B. Williams, and Carmi

Korine. "Habitat Aridity as a Determinant of the Trade-off between Water Conservation and

Evaporative Heat Loss in Bats." Journal of Comparative Physiology B 192, no. 2 (January 17,

2022): 325–33. https://doi.org/10.1007/s00360-021-01425-2.

Naumann, G., L. Alfieri, K. Wyser, L. Mentaschi, R. A. Betts, H. Carrao, J. Spinoni, J. Vogt,

and L. Feyen. "Global Changes in Drought Conditions Under Different Levels of Warming."

Geophysical Research Letters 45, no. 7 (April 13, 2018): 3285–96.

https://doi.org/10.1002/2017gl076521.

Naya, Daniel E., Claudio Veloso, Pablo Sabat, and Francisco Bozinovic. "Physiological

Flexibility and Climate Change: The Case of Digestive Function Regulation in Lizards."

Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology

159, no. 1 (May 2011): 100–104. https://doi.org/10.1016/j.cbpa.2011.02.005.

Nicholson, S.E., and J.C. Selato. "The Influence of La Nina on African Rainfall."

International Journal of Climatology 20, no. 14 (n.d.): 1761–76. https://doi.org/10.1002/1097-

0088(20001130)20:14<1761::AID-JOC580>3.0.CO;2-W.

Ostrowski, Stéphane, Joseph B. Williams, Pascal Mésochina, and Helga Sauerwein. "Physiological Acclimation of a Desert Antelope, Arabian Oryx (Oryx Leucoryx), to Long-Term Food and Water Restriction." Journal of Comparative Physiology B 176, no. 3 (November 9, 2005): 191–201. https://doi.org/10.1007/s00360-005-0040-0.

Pajic, Petar, Pavlos Pavlidis, Kirsten Dean, Lubov Neznanova, Rose-Anne Romano, Danielle Garneau, Erin Daugherity, Anja Globig, Stefan Ruhl, and Omer Gokcumen. "Independent Amylase Gene Copy Number Bursts Correlate with Dietary Preferences in Mammals." eLife 8 (May 14, 2019). https://doi.org/10.7554/elife.44628.

Parmesan, Camille. "Ecological and Evolutionary Responses to Recent Climate Change." Annual Review of Ecology, Evolution, and Systematics 37, no. 1 (December 1, 2006): 637–69. https://doi.org/10.1146/annurev.ecolsys.37.091305.110100.

Parmesan, Camille, Terry L. Root, and Michael R. Willig. "Impacts of Extreme Weather and Climate on Terrestrial Biota." Bulletin of the American Meteorological Society 81, no. 3 (March 1, 2000): 443–50. https://doi.org/10.1175/1520-0477(2000)081<0443:IOEWAC>2.3.CO;2.

Post, Eric, Jedediah Brodie, Mark Hebblewhite, Angela D. Anders, Julie A. K. Maier, and Christopher C. Wilmers. "Global Population Dynamics and Hot Spots of Response to Climate Change." BioScience 59, no. 6 (June 2009): 489–97. https://doi.org/10.1525/bio.2009.59.6.7.

Rimbach, Rebecca, Stéphane Blanc, Alexandre Zahariev, Maria Gatta, Neville Pillay, and Carsten Schradin. "Seasonal Variation in Energy Expenditure in a Rodent Inhabiting a Winter-Rainfall Desert." Journal of Comparative Physiology B 188, no. 5 (June 8, 2018): 877–88. https://doi.org/10.1007/s00360-018-1168-z.

Rimbach, Rebecca, Jörg Jäger, Neville Pillay, and Carsten Schradin. "Food Availability Is the

Main Driver of Seasonal Changes in Resting Metabolic Rate in African Striped Mice

(Rhabdomys Pumilio)." Physiological and Biochemical Zoology 91, no. 2 (March 2018):

826–33. https://doi.org/10.1086/696828.

Rintoul, Jody L. P., and R. Mark Brigham. "The Influence of Reproductive Condition and

Concurrent Environmental Factors on Torpor and Foraging Patterns in Female Big Brown

Bats (Eptesicus Fuscus)." Journal of Comparative Physiology B 184, no. 6 (June 28, 2014):

777–87. https://doi.org/10.1007/s00360-014-0837-9.

Rocha, Joana L., Raquel Godinho, José C. Brito, and Rasmus Nielsen. "Life in Deserts: The

Genetic Basis of Mammalian Desert Adaptation." Trends in Ecology & Evolution 36, no.

7 (July 2021): 637–50. https://doi.org/10.1016/j.tree.2021.03.007.

Sabat, Pablo, Sandra Gonzalez-Vejares, and Karin Maldonado. "Diet and Habitat Aridity

Affect Osmoregulatory Physiology: An Intraspecific Field Study along Environmental

Gradients in the Rufous-Collared Sparrow." Comparative Biochemistry and Physiology Part

A: Molecular & Integrative Physiology 152, no. 3 (March 2009): 322–26.

https://doi.org/10.1016/j.cbpa.2008.11.003.

Saraç, F., and F. Saygılı. "Causes of High Bone Alkaline Phosphatase." Biotechnology

& Biotechnological Equipment 21, no. 2 (January 2007): 194–97.

https://doi.org/10.1080/13102818.2007.10817444.

Seabrook, Leonie, Clive McAlpine, Greg Baxter, Jonathan Rhodes, Adrian Bradley, and

Daniel Lunney. "Drought-Driven Change in Wildlife Distribution and Numbers: A Case

Study of Koalas in South West Queensland." Wildlife Research 38, no. 6 (2011): 509.

https://doi.org/10.1071/wr11064.

Schoepf, I., N. Pillay, and C. Schradin. "The Pathophysiology of Survival in Harsh

Environments." Journal of Comparative Physiology B 187, no. 1 (July 15, 2016): 183–201.

https://doi.org/10.1007/s00360-016-1020-2.

Schoepf, I., N. Pillay, and C. Schradin. "Trade-Offs between Reproduction and Health in

Free-Ranging African Striped Mice." Journal of Comparative Physiology B 187, no. 4

(February 4, 2017): 625–37. https://doi.org/10.1007/s00360-016-1054-5.

Schulte-Hostedde, Albrecht I., Bertram Zinner, John S. Millar, and Graham J. Hickling.

"RESTITUTION OF MASS–SIZE RESIDUALS: VALIDATING BODY CONDITION

INDICES." Ecology 86, no. 1 (January 2005): 155–63. https://doi.org/10.1890/04-0232.

Seebacher, Frank, and Craig E. Franklin. "Determining Environmental Causes of Biological

Effects: The Need for a Mechanistic Physiological Dimension in Conservation Biology."

Philosophical Transactions of the Royal Society B: Biological Sciences 367, no. 1596 (June

19, 2012): 1607–14. https://doi.org/10.1098/rstb.2012.0036.

Seebacher, Frank, William Davison, Cara J Lowe, and Craig E Franklin. "A Falsification of

the Thermal Specialization Paradigm: Compensation for Elevated Temperatures in Antarctic

Fishes." Biology Letters 1, no. 2 (April 28, 2005): 151–54.

https://doi.org/10.1098/rsbl.2004.0280.

Shapiro, R, and R.P Heaney. "Co-Dependence of Calcium and Phosphorus for Growth and

Bone Development under Conditions of Varying Deficiency." Bone 32, no. 5 (May 2003):

532–40. https://doi.org/10.1016/s8756-3282(03)00060-7.

Solmsen, N., J. Johannesen, and C. Schradin. "Highly Asymmetric Fine-Scale Genetic

Structure between Sexes of African Striped Mice and Indication for Condition Dependent

Alternative Male Dispersal Tactics." Molecular Ecology 20, no. 8 (March 3, 2011): 1624–34.

https://doi.org/10.1111/j.1365-294x.2011.05042.x.

Somero, G. N. "The Physiology of Climate Change: How Potentials for Acclimatization and

Genetic Adaptation Will Determine 'Winners' and 'Losers.'" Journal of Experimental

Biology 213, no. 6 (March 15, 2010): 912–20. https://doi.org/10.1242/jeb.037473.

Somero, George N. "The Physiology of Global Change: Linking Patterns to Mechanisms."

Annual Review of Marine Science 4, no. 1 (January 15, 2012): 39–61.

https://doi.org/10.1146/annurev-marine-120710-100935.

Speakman, John R. "The Physiological Costs of Reproduction in Small Mammals."

Philosophical Transactions of the Royal Society B: Biological Sciences 363, no. 1490

(August 8, 2007): 375–98. https://doi.org/10.1098/rstb.2007.2145.

Stawski, Clare, and A Daniella Rojas. "Thermal Physiology of a Reproductive Female

Marsupial, Antechinus Flavipes." Mammal Research 61, no. 4 (July 21, 2016): 417–21.

https://doi.org/10.1007/s13364-016-0287-8.

Sullivan, Jennifer C., Laura Semprun-Prieto, Erika I. Boesen, David M. Pollock, and Jennifer

S. Pollock. "Sex and Sex Hormones Influence the Development of Albuminuria and Renal

Macrophage Infiltration in Spontaneously Hypertensive Rats." American Journal of

Physiology-Regulatory, Integrative and Comparative Physiology 293, no. 4 (October 2007):

R1573–79. https://doi.org/10.1152/ajpregu.00429.2007.

Sundar Ray, Chinmaya, Bratati Singh, Itishri Jena, Sudeshna Behera, and Subhashree Ray.

"Low Alkaline Phosphatase (ALP) In Adult Population an Indicator of Zinc (Zn) and

Magnesium (Mg) Deficiency." Current Research in Nutrition and Food Science Journal 5, no. 3 (December 19, 2017): 347–52. https://doi.org/10.12944/crnfsj.5.3.20.

Taylor, K. D., and M. G. Green. "The Influence of Rainfall on Diet and Reproduction in Four African Rodent Species." Journal of Zoology 180, no. 3 (November 1976): 367–89. https://doi.org/10.1111/j.1469-7998.1976.tb04683.x.

Thaler, M Michael, David L Gemes, and Arne F Bakken. "Enzymatic Conversion of Heme to Bilirubin in Normal and Starved Fetuses and Newborn Rats." Pediatric Research 6, no. 3 (March 1972): 197–201. https://doi.org/10.1203/00006450-197203000-00008.

Thornthwaite, C. W. "An Approach toward a Rational Classification of Climate." Geographical Review 38, no. 1 (January 1948): 55. https://doi.org/10.2307/210739.

Tothova, C., O. Nagy, and G. Kovac. "Serum Proteins and Their Diagnostic Utility in Veterinary Medicine: A Review." Veterinární Medicína 61, no. 9 (September 30, 2016): 475–96. https://doi.org/10.17221/19/2016-vetmed.

Tshikae, Balatlhane P., Adrian L. V. Davis, and Clarke H. Scholtz. "Does an Aridity and Trophic Resource Gradient Drive Patterns of Dung Beetle Food Selection across the Botswana Kalahari?" Ecological Entomology 38, no. 1 (January 22, 2013): 83–95. https://doi.org/10.1111/j.1365-2311.2012.01409.x.

Usui, Takuji, David Lerner, Isaac Eckert, Amy L. Angert, Colin J. Garroway, Anna Hargreaves, Lesley T. Lancaster, et al. "The Evolution of Plasticity at Geographic Range Edges." Trends in Ecology & Evolution 38, no. 9 (September 2023): 831–42. https://doi.org/10.1016/j.tree.2023.04.004.

Vespasiani-Gentilucci, Umberto, Antonio De Vincentis, Luigi Ferrucci, Stefania Bandinelli, Raffaele Antonelli Incalzi, and Antonio Picardi. "Low Alanine Aminotransferase Levels in the Elderly Population: Frailty, Disability, Sarcopenia, and Reduced Survival." The Journals of Gerontology: Series A 73, no. 7 (June 19, 2017): 925–30. https://doi.org/10.1093/gerona/glx126.

"Clinical Methods: The History, Physical, and Laboratory Examinations." Annals of Internal Medicine 113, no. 7 (October 1, 1990): 563. https://doi.org/10.7326/0003-4819-113-7-563_2.

Webster, Nicole S, and Thorsten B H Reusch. "Microbial Contributions to the Persistence of Coral Reefs." The ISME Journal 11, no. 10 (May 16, 2017): 2167–74. https://doi.org/10.1038/ismej.2017.66.

Willmer, Pat, Graham Stone, and Ian Johnston. Environmental Physiology of Animals. John Wiley & Sons, 2009.

Wu, Nicholas C., and Frank Seebacher. "Physiology Can Predict Animal Activity, Exploration, and Dispersal." Communications Biology 5, no. 1 (February 3, 2022). https://doi.org/10.1038/s42003-022-03055-y.

Yousef, M.I., H.A. El Hendy, F.M. El-Demerdash, and E.I. Elagamy. "Dietary Zinc Deficiency Induced-Changes in the Activity of Enzymes and the Levels of Free Radicals, Lipids and Protein Electrophoretic Behavior in Growing Rats." Toxicology 175, no. 1–3 (June 2002): 223–34. https://doi.org/10.1016/s0300-483x(02)00049-5.

Zhang, Jianfen, Na Zhang, Shufang Liu, Songming Du, and Guansheng Ma. "Young Adults with Higher Salt Intake Have Inferior Hydration Status: A Cross-Sectional Study." Nutrients 14, no. 2 (January 11, 2022): 287. https://doi.org/10.3390/nu14020287.

Zhao, Miao-Miao, Jeremy Krebs, Xi Cao, Jing Cui, Dong-Ning Chen, Yu Li, Lin Hua, Jim

Mann, and Jin-Kui Yang. "Helicobacter Pylori Infection as a Risk Factor for Serum Bilirubin

Change and Less Favourable Lipid Profiles: A Hospital-Based Health Examination Survey."

BMC Infectious Diseases 19, no. 1 (February 14, 2019). https://doi.org/10.1186/s12879-019-

3787-8.

Zomer, Robert J., Jianchu Xu, and Antonio Trabucco. "Version 3 of the Global Aridity Index

and Potential Evapotranspiration Database." Scientific Data 9, no. 1 (July 15, 2022).

https://doi.org/10.1038/s41597-022-01493-1.

Tables

<u>***Table 1: Summary of main ecological characteristics of R.bechuanae and R.d.dilectus and***</u>

<u>***our predictions on their physiological response during the dry season***</u>

ECOLOGICAL CHARACTERISTICS	*R.BECHUANAE*	*R.D.DILECTUS*
ENVIRONMENTAL NICHE	Arid to semi-arid (Ganem et al., 2020)	Mesic to semi-arid (Ganem et al., 2020)
PREFERRED HABITAT	Bushy patches (Dufour et al., 2015)	Continuous cover / grassland (Dufour et al., 2015)
SOCIAL STRUCTURE	Group-living (Dufour et al., 2018)	Solitary / opportunistic group-living (Dufour et al., 2018)
SPECIES-SPECIFIC RESPONSE TO SEASONAL VARIATIONS OF DRY CONDITIONS: PREDICTIONS		
BODY CONDITION	↘↘ or ↘*	↘↘
NUTRITIONAL METABOLISM	↘↘ or ↘*	↘↘
LIVER ACTIVITY	↘↘ or ↘*	↘↘
WATER BALANCE	↘↘ or ↘*	↘↘
SENSITIVITY TO INFECTION	↗↗ or ↗*	↗↗
ASEASONAL DIFFERENCES: PREDICTIONS		
BODY CONDITION	+	-
LIVER ACTIVITY	+	-
WATER BALANCE	+	-
SENSITIVITY TO INFECTION	-	+

910   *Adaptation to harsher arid conditions could have enabled *R.bechuanae* to withstand growing aridity without

911   suffering as many adverse effects as *R.d.dilectus* in the semi-arid zone. Alternatively, semi-arid populations of

912   both species could both have developed local adaptation.

914   <u>***Table 2: Geographical position and Aridity Index of sampled sites for each species***</u>

SPECIES	SAMPLED SITES	GEOGRAPHICAL COORDINATES	ARIDITY INDEX[*]
R.bechuanae	Gariep Dam Nature Reserve	E25.55537; S30.57589	0.35-0.49
	Kalkfontein Dam Nature Reserve	E25.28031; S29.52058	0.47-0.61
	Benfontein Nature Reserve**	E24.81824; S28.82747	0.32-0.46
R.d.dilectus	Bloemhof Dam Nature Reserve	E25.65144; S27.66063	0.29-0.41
	Barberspan Bird Sanctuary**	E25.58354; S26.56538	0.34-0.44
	Wolwespruit Nature Reserve	E26.26326; S27.41418	0.43-0.63

915   *Average Aridity index (see **Supplementary material** for formula) ± Standard Error of the

916   Mean, based on climate data collected from 2010 to 2021 (South African Weather Service)

917   ** Sampled only in May 2022 (dry season onset).

919 *Table 3:* **Results of the Analysis of Variance (response variable: NDVI)**

Variable	Df	R²	F	P
Session	1	0.445	442.757	**< 2.2*10^{-16}**
Site	3	0.348	115.385	**< 2.2*10^{-16}**
Session*Site	3	0.021	7.113	**1.512*10^{-4}**
Residual	229	0.185		

922 *Table 4:* **Results of the Analysis of Variance (response variable : body condition)**

Variable	Df	R²	F	P
Session	1	0.017	4.489	**0.035**
Breeding status	2	0.079	10.353	**$4.978*10^{-5}$**
Session*Breeding status	2	0.024	3.142	**0.045**
NDVI	1	0.013	3.425	0.066
Sex	1	0.004	1.144	0.286
Session*Sex	1	0.003	0.703	0.403
Residual	229	0.861		

925 ***Table 5: Summary of the results of the best fitting Permutational multivariate analysis of***

926 ***variance model, addressing variation in physiological response (site= random factor)***

Variable	Df	R^2	Pseudo-F	P
Session	1	0.011	2.923	0.065
Species	1	0.042	11.227	**0.002**
Sex	1	0.010	2.786	0.084
Breeding status	2	0.025	3.282	**0.028**
Body condition	1	0.018	4.936	**0.019**
Species * Sex	1	0.018	4.762	**0.020**
Species * Breeding status	2	0.030	3.964	**0.022**
Residual	226	0.846		

Table 6: Summary of the results of univariate analyses of variance addressing variation in

931 ***responses of each of the 12 studied physiological variables (best fitting models).***

Physiological marker	Variable(s) kept in Best Model and their significance	adjusted R²	F	df Model	df Residual	P (model)
Albumin	**Session*****, **Sex***	0.098	13.72	2	233	$2.338*10^{-6}$
Alkaline phosphatase	**Session*****, **Species***, **Sex*****, **Body condition*****	0.310	18.52	6	228	$< 2.2*10^{-16}$
Alanine aminotransferase	**Session***, Species, Sex, Breeding status, Species*Sex, Species*Breeding Status	0.060	2.884	8	227	0.004
Amylase	**Species*****, Sex, Breeding status, body condition, Species*Sex, Species*Breeding Status	0.195	8.104	8	227	$1.277*10^{-9}$
Calcium	Species	0.009	3.095	1	234	0.080
Globulin	**Session*****, Body condition	0.296	50.4	2	233	$< 2.2*10^{-16}$
Glucose	Session\+·, **Sex****	0.079	11.14	2	233	$2.397 *10^{-5}$
Potassium	**Session***, **Species***	0.068	9.569	2	233	$1.014*10^{-4}$
Sodium	Session, **Species*****, Sex	0.110	10.65	3	232	$1.39*10^{-6}$
Blood Urea Nitrogen	*Null Random effect model*	0.072	/	/	/	/
Phosphorus	Species, Sex, **Body condition*****, Breeding status + random	0.282	/	/	/	(χ^2 test vs. null model) $4.603*10^{-9}$

Note: The full models, before backwards elimination, included session, species, body condition, sex, breeding

935 *status, sex*species and breeding status*species as variables, with breeding status nested within sex and site as a*

936 *partially-crossed random factor nested within species.*

937 ***Bold:*** *significant effects (p < 0.05) following Bonferroni correction with:*

938 ****: p < 0.001; **: 0.001 < p < 0.01 ; *: 0.01 < p < 0.05; \+ : 0.05<p<0.1*

Figure captions

Figure 1: Map of known occurrences of 6 main clades of Rhabdomys (translucent dots) and locations sampled in this study (full colour dots) in southern Africa. Based on published data and unpublished data (origin details are available in doi: 0d1b3414-7e2a-11ea-a38d-00163e26bfb0). Base map: World Topographic Map Esri Standard, Aridity Index layer was computed from a 0.5° global grid, using data from the Version 3 of the Global Aridity Index and Potential Evapotranspiration Database (Zomer et al.,2022).

Figure 2: Graphical summary of organization and composition of data used for analysis in this study. Figures between brackets indicate number of modalities or types of categories included in the factor.

Figure 3: Total monthly precipitation for all sampled sites as recorded in 2022.

Figure 4: Average Normalized Differential Vegetation Index calculated within a buffer circle of 60m radius around each successful trap per site and session. NDVI was retrieved from the Copernicus Open Access Data Hub (Copernicus Sentinel-2 data [2023]). For each site and session, we retained NDVI data available for the closest day to the beginning of a sampling session, expected to represent the conditions experienced by the mice at the time of capture.

Figure 5: Significant shifts in blood physiological marker concentrations between the onset and the end of the dry season (mean ± SE)

Figure 6: Significant interspecific differences involving four blood physiological marker concentrations (mean ± SE)

Figures

<u>Figure 1: Map of known occurrences of 6 main clades of Rhabdomys (translucent dots)</u>

<u>and locations sampled in this study (full colour dots) in southern Africa</u>

963   *Based on published data and unpublished data (origin details are available in doi:*

964   *0d1b3414-7e2a-11ea-a38d-00163e26bfb0). Base map: World Topographic Map Esri*

965   *Standard, Aridity Index layer was computed from a 0.5° global grid, using data from*

966   *the Version 3 of the Global Aridity Index and Potential Evapotranspiration Database*

967   *(Zomer et al.,2022).*

Figure 2: Graphical summary of organization and composition of data used for

analysis in this study. Figures between brackets indicate number of modalities or types

of categories included in the factor.

977 ***<u>Figure 3: Total monthly precipitation for all sampled sites as recorded in 2022.</u>***

Figure 4: Average Normalized Differential Vegetation Index calculated within a

buffer circle of 60m radius around each successful trap per site and session.

982 _NDVI was retrieved from the Copernicus Open Access Data Hub (Copernicus_

983 _Sentinel-2 data [2023]). For each site and session, we retained NDVI data available_

984 _for the closest day to the beginning of a sampling session, expected to represent the_

985 _conditions experienced by the mice at the time of capture._

991 ***Figure 5: Significant shifts in blood physiological marker concentrations***

992 ***between the onset and the end of the dry season (mean ± SE)***

996 ***Figure 6: Significant interspecific differences involving four blood***

997 ***physiological marker concentrations (mean ± SE)***

998