Do sister species respond similarly to dry conditions? An ecophysiological approach *in natura*

Authors: Hamilcar Keilani[1], Nico Avenant[2], Pierre Caminade[1], Neville Pillay[3], Guila Ganem[1,3]

1. **ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France ; e-mail : hamilcar.keilani@umontpellier.fr, pierre.caminade@umontpellier.fr, guila.ganem@umontpellier.fr**
2. **Department of Mammalogy, National Museum and Centre for Environmental Management, University of the Free State, Bloemfontein, South Africa; e-mail: navenant@nasmus.co.za**
3. **School of Animal, Plant and Environmental Science, University of the Witwatersrand, Wits, South Africa; e-mail: neville.pillay@wits.ac.za, guila.ganem@wits.ac.za**

Acknowledgements

We thank R.Nokha who provided substantial field work contribution to this project. Constructive and insightful comments were provided by C.Smadja and C.Schradin helped improve a previous version of this paper. Technical support by D.Greuet (INM), L.Paradis (Geomatics, ISEM) and GenSeq platform.

We are grateful to the rangers and personnel of Barberspan Bird Sanctuary, Benfontein Nature Reserve, Bloemhof Dam Nature Reserve, Gariep Dam Nature Reserve, Kalkfontein Dam Nature Reserve and Wolwespruit Nature Reserve who facilitated our work in these locations.

24   Funding: CNRS/NRF, OSU-OREME

25   Conflict of Interest

26   The authors declare they have no conflict of interest.

27   Author contributions

G.G. conceived the idea and designed methodology with input from H.K; G.G.,

H.K., P.C., collected the data; H.K. analyzed the data with input from G.G.; H.K.

wrote the first draft of the manuscript with input from G.G.; N.P. and N.A.

contributed critically to the manuscript, secured funding for the project with G.G.

and helped with field logistics. All authors approved the final version of the

manuscript.

34   Statement on inclusion

35   Our study brings together authors from France and South Africa. All authors were

36   engaged early on with the research and study design to ensure that the diverse sets

37   of perspectives they represent was considered from the onset. Whenever relevant,

38   literature published by scientists from the study region was cited.

39   Data availability statement

40   Data will be available from the Dryad repository (TBA)

Abstract

1. **Understanding the proximate mechanisms of organisms' adaptations to the environment is a classical problem in evolutionary ecology. In the context of climate change, inducing increased aridity and disruption of seasonality, it is crucial to identify the potential for species responses.**

2. **Here, the physiological response to food and water limitation during seasonally dry conditions were investigated. We studied two rodent species of the genus *Rhabdomys*, one arid and one mesic, the ranges of which overlap in a semi-arid zone of South Africa. We measured blood concentrations of physiologically relevant metabolites in semi-arid populations of the two species at the onset and the end of the dry season.**

3. **We found similar shifts in blood metabolite levels in the two species, indicating malnutrition consistent with the observed degradation of habitat quality between the start and the end of the dry season. Furthermore, regardless of season, differences between the two species in blood metabolite concentrations (amylase, sodium, potassium, bilirubin, and alkaline phosphatase) were observed, that could indicate contrasting diets and water conservation abilities.**

4. **Overall, we show that, as dry conditions worsen, organisms are increasingly affected by reduced food availability, and local adaptation to arid conditions may provide the arid species with an advantage to cope with semi-arid conditions. Our study shows that even mild arid conditions could have a negative impact and questions resilience of animals to harsher arid conditions.**

Keywords

Aridity, blood metabolites, ecophysiology, liver, local adaptation, nutrition, *Rhabdomys*

Introduction

Organisms are constantly exposed to a wide range of environmental variations, including short-term changes during their lifetime and longer-term changes across generations (Lopez-Maury et al., 2008). In the coming decades, extreme and more frequent droughts are predicted by most projections, resulting from changes in precipitation patterns, increased temperature, and evaporation (Naumann et al., 2018). These changes will induce great challenges to organisms. Under such circumstances, divergent selection can lead to the evolution of local adaptations where, in a given environment, resident phenotypes outperform nonresident ones (Kawecki & Ebert, 2004).

Local ecological adaptation can be described in several different and complementary ways, such as through its genetic basis, transcriptomic, behavioral, or physiological aspects. Adaptive evolution *sensu stricto* occurs when the genetic constitution of a population changes because of natural selection (Merilä & Hendry, 2013). In contrast, local adaptation *sensu lato*, also considers concepts such as phenotypic plasticity. It is usually assessed using comparative population or species fitness estimates under novel environmental conditions (Webster & Reusch, 2017). The response to local environmental conditions is a major factor in the generation and maintenance of biodiversity, impacting population dynamics, biogeographical ranges, and species interactions (Blanquart et al., 2013; Post et al., 2009). Specifically, physiological limits can drive and be driven by evolution, shape species distributions, and define species

response capacities to future climate change, directly impacting risks of extinction

(Somero, 2012). For example, adaptive variation in lethal temperature (LT_{50}) has been

established in marine invertebrates inhabiting different latitudinal niches (Somero,

2010).

Studies of different physiological systems, such as the cardiovascular (Seebacher et

al., 2005, Berkel & Cacan, 2022) or digestive systems (Naya et al., 2011), in response

to different abiotic conditions, can improve our understanding of the causal

mechanisms of responses of organisms to climate variation (Seebacher & Franklin,

2012). Biotic factors, such as sex, age, diet, strain, or breeding status also influence

individual responses (McClure, 1999). With a focus on influence of air temperature,

water availability and energy resources on an organism's physiological state, research

in physiological ecology has addressed how variation in physiology evolves and is

maintained in relation to environment conditions (Feder & Block, 1991).

As a result of climate change, organisms can be faced with warmer and drier

environmental conditions (Parmesan et al., 2000), likely to impact their capacity to

maintain homeostasis (Fuller et al., 2016). In dry environments, food and water can be

scarce, generating strong selection on physiological attributes maximizing energy and

water availability. Habitat generalist and specialist taxa may respond differently to

environmental changes. Desert specialists exhibit specific physiological and/or

behavioral traits, resulting from genetic adaptations to the environment, allowing them

to continuously endure or evade periods of resource restriction (Rocha et al., 2021). In

contrast, generalist species may seek to escape the effects of aridity through

temporary avoidance (Abraham et al., 2019), plastic adjustments during unfavorable

seasons (Christian et al., 1996; Kobbe et al., 2011)Compared to mesic species, arid

species have lower basal metabolic rates and evaporative water loss (Tieleman et al.,

2003; Muñoz-Garcia et al., 2022), adopt a more flexible diet (Tshikae et al., 2013), and

cope better with prolonged droughts (Boyers et al., 2021).

Analysis of blood metabolites has successfully been used to investigate kidney and

liver physiology of wild populations (e.g. Al-Eissa et al., 2012). The kidney and the liver

both contribute to homeostasis in periods of food and water scarcity; indeed the kidney

plays a crucial role in maintaining osmotic balance and the liver is involved in dietary

metabolism. Hence, dry conditions can put a strain on these organs and contribute to

the emergence of pathologies detrimental to survival and reproduction. Water loss can

also mechanically induce high concentrations of some blood markers and impact

organism's capacity to evacuate products of metabolic waste, such as urea, from the

blood (Ostrowski et al., 2006).

In our study, potential impact of increased aridity was addressed by investigating the

physiological consequences of seasonal variation in dry conditions, in two species of

the diurnal African striped mouse genus *Rhabdomys*. The two study species,

R.bechuanae and *R.dilectus dilectus*, are characterized by different environmental

niches (du Toit et al. 2012, Meynard et al. 2012). Throughout most of its distribution,

R.d.dilectus occurs in mesic habitats with vegetation cover and nests in dense grass,

while *R.bechuanae* is found in semi-arid and arid habitats, thriving predominantly in

sparsely vegetated areas and nesting in bushes (Dufour et al., 2015; Dufour et al.,

2019). Moreover, differences in morphology or behavior suggest adaptation to dry

conditions in *R.bechuanae* (Ganem et al., 2020; Dufour et al., 2019). Here we address

both physiological response to seasonal dry conditions and divergence in the two

species in the same bioclimatic region.

Indeed, at the edge of their distributions in central South Africa, the two species inhabit

a semi-arid region within which they occur either as parapatric or sympatric

populations. This semi-arid zone, the expansion and contraction of which depends on

land use and precipitation (Kruger, 2006; Masubele et al., 2014; Department of

Environmental Affairs, 2018; Lian et al., 2021), has experienced a recent trend towards

desiccation (Jury, 2021). Such dry conditions could generate additional selective

pressures on the semi-arid populations of the two species.

In the Succulent Karoo of South Africa, where the average daily precipitation during

the dry season is 0.26 mm *versus* 0.78 mm during the wet season, the probability of

survival of adult *Rhabdomys pumilio* could be predicted from their physiological

response at the start of the dry season (Schoepf et al., 2017a). Indeed, Schoepf and

collaborators found higher serum concentrations of albumin, glucose, potassium, and

lower concentrations of globulin, in animals that survived compared to those that did

not survive the dry season. At the peak of the dry season, *R.pumilio* had lower

concentrations of glucose and phosphorus and higher concentrations in globulin and

urea nitrogen if they survived the dry season. These results strongly suggest that such

markers could be good indicators of individual fitness.

While many studies have harnessed physiology, genomics, and transcriptomics to

highlight potential adaptive interspecific or interpopulational contrasts, studies under

common natural environmental conditions are lacking (Rocha et al., 2021). This study

took place in a natural semi-arid environment and asked how seasonal variation in dry

conditions influenced the physiological responses of the two species. We compared

body condition, blood concentrations of markers of kidney and liver functions and

habitat characteristics of semi-arid parapatric populations of *R.bechuanae* and

R.dilectus dilectus at the start *versus* the end of the dry season. First, we expected the

habitat available for the mice would be drier at the end compared to the onset of the

dry season, inducing a reduction of food and water, impairing liver and kidney function,

and impacting body condition. Second, assuming local adaptive plasticity, we

hypothesized that both species would be able to adjust their physiological responses

to seasonal changes in dry conditions. Third, we expected *R.bechuanae*, having

evolved in arid environments, would perform better in semi-arid conditions than

R.d.dilectus (having evolved in mesic areas) and be closer to the expected local

optimal response to increased dry conditions, as found for *Mus musculus* (Bittner et

al., 2021). Alternatively, if both species evolved specific adaptations to their semi-arid

environment, we expected no species differences (**Table 1**).

Materials and Methods

Ethics statement

177 Permits to work and handle animals in the field were obtained from the DESTEA of the

178 Free State, the North West DEDECT and Northern Cape DENC (respectively,

179 n°202110000007994, NW 38956/04/2022, FAUNA 0389/2022). Animal handling was

180 performed under ethical clearance from the Languedoc Roussillon ethical committee

181 for animal experimentation (n°2022041512256467 v5).

Choice of sampling periods and sites

We selected six sampling sites based on their geographical position and their aridity

index (AI). The AI calculation was based on Thornthwaite method (Thornthwaite, 1948;

see **Supplementary Material** for full formula) using rainfall and temperature data

collected from 2010-2021 (South African Weather Service).

We selected parapatric populations of the two species in a semi-arid region of South

Africa (0.2 < AI < 0.5). In this region, a hot and wet season occurs from roughly

December to February, and a cool dry season from May to September. We sampled

six sites at the onset (May 2022) of the dry season: Barberspan Bird Sanctuary,

Benfontein Nature Reserve, Bloemhof Dam Nature Reserve, Gariep Dam Nature

Reserve, Kalkfontein Dam Nature Reserve and Wolwespruit Nature Reserve (**Figures**

1&3, Table 2). We then resampled four of these sites at the end of the dry season

(September 2022). Two sites (Barberspan and Benfontein) were excluded from the

September sampling due to extreme weather conditions.

Capture and field protocol

We used small mammal PVC traps (LxHxW= 29.6x7.5x7.5 cm), baited with a mixture

of oats, peanut butter, salt and sunflower seeds, and added a piece of cotton wool.

They were placed approximately every 15 m along roughly 150 to 300 m transects.

Since *Rhabdomys* is mostly diurnal, the traps were checked twice a day (8 AM and 3

PM). On the field, all trapped striped mice were measured (body length and mass),

their sex and breeding status assessed based on external morphological metrics, and

a 0.5 mm piece of tail was collected and kept in 98 % ethanol for species identification.

One of four possible breeding statuses was recorded: breeding males; breeding

females; non-breeding males; and non-breeding females. All striped mice were marked

with a unique ear-tag before their release at the trapping location unless they were

kept for the physiological study (i.e. kept in their trap with cotton and food).

Overall, trapping effort was 9688 trap nights (**Supplementary Table 1**). Out of 903

trapped small mammals, 694 were *Rhabdomys*. For the physiological study, we

selected adults, avoiding related individuals as much as possible, by selecting pairs of

mice (a male and a female) trapped at least 100 m apart. This distance was based on

data obtained for *R. pumilio* indicating minimal relatedness (R=0 between males and

R=0.06 between females) at this distance (Solmsen et al. 2012). Trapping took place

between 3 and 6 days per site, until we reached our target of 20 individuals.

Habitat

Vegetation composition

Earlier studies have shown inter-species differences in vegetation cover and structure

requirements (Dufour et al., 2015). Since habitat characteristics can vary locally, we

characterized the vegetation structure (i.e. grass *versus* woody vegetation) around

traps in which a *Rhabdomys* was trapped by measuring the percentage of grass,

bushes, and bare soil within 2mx2m and 10x10m quadrats centered on the trap

position. Each 2mx2m quadrat comprised four 1x1m subunits assessed using a 1x1m

metal frame; the results obtained for the four subunits were then averaged. Each

10mx10m quadrat was assessed following a visual inspection of a 10x10m zone

around each trap. We characterized 236 quadrats of each type (10x10m and 2mx2m).

Normalised Difference Vegetation Index

NDVI, a satellite imagery-based index informative of ground vegetation greenness,

was retrieved from the Copernicus Open Access Data Hub (Copernicus Sentinel-2

data, 2023). For each site and sampling session, we used NDVI data available for the

closest day to the beginning of a sampling session, which was expected to represent

the conditions experienced by the mice at the time of capture **(Supplementary Table**

3).

Blood extraction and assessment of blood metabolites concentrations

A total of 273 animals were euthanized by means of cervical dislocation on their day of capture, and a blood volume of at least 100 μL was collected in lithium-heparinized tubes by cardiac puncture. The breeding status was confirmed for all individuals during dissection. The whole liver, the left kidney, and the skull of every individual were also collected for a complementary study. Because the two species could not be distinguished visually, the spleen was also harvested from each mouse and subsequently analyzed for post-hoc species identification using Cytochrome Oxidase I genotyping (as described in Ganem et al. 2012).

Levels of albumin, alkaline phosphatase, alanine aminotransferase, amylase, total bilirubin, blood urea nitrogen, total protein, globulin, glucose, calcium, phosphorus, potassium, and sodium were successfully measured in the whole blood of 257 individuals immediately after blood collection (Abaxis VETSCAN V2 technology with the Comprehensive diagnostic profile cartridge), following the manufacturer's instructions. The coefficient of variation (CV%) of measured levels of these markers was determined from data obtained by analysing a given blood sample twice. Samples from six adult striped mice (5 *R.bechuanae*, 1 *R.d.dilectus*) were used for these calculations. For these six samples, the calculated CV equaled 5.6 ± 1.9 %, which is consistent with the intra-test CV% provided in the manufacturer's instruction manual for other mammal species (**Supplementary Table 4**).

Replication statement

Inferences for the habitat analyses were made at the scale of successful traps (**3**).

Inferences for the body condition analysis were made at the individual scale

(depending on the factor of interest, **1** or **2**).

Inferences for the physiology analyses were made at the individual scale (depending

on the factor of interest, **1** or **2**).

Scale of inference	Scale at which the factor of interest is applied	Number of replicates at the appropriate scale
(1) Individuals	Populations	3 *R.bechuanae* onset of the dry season, 3 *R.d.dilectus* onset of the dry season, 2 *R.bechuanae* end of the dry season, 2 *R.d.dilectus* end of the dry season
(2) Individuals	Individuals	236
(3) Successful traps	Sites	6 onset of the dry season, 4 end of the dry season

Data preparation and analysis

All statistical tests were carried out using the R software (version 4.2.1).

Habitat

Vegetation composition

Principal Component Analysis (package FactoMineR version 2.9) was performed, including the 7 variables measured within the $4m^2$ quadrats and within the $100m^2$ quadrats for each successful trap (traps in which *Rhabdomys* was captured). In each case, over 80 % of the variance was explained by the first five Principal Components (**Supplementary Tables 7 & 9**). As Multivariate normality was violated in our dataset, the coordinates of all traps on the first 5 PCs were used as response variables of a *PERMANOVA* (package *vegan*, version 2.6-4). *PERMANOVA* is a multivariate statistical inference tool using permutational algorithms (Anderson, 2001). This equivalent to MANOVA operates in a distribution-free setting and is robust to non-normality of residuals as well as dispersion heterogeneity, even in slightly unbalanced designs (Anderson, 2013). We tested differences between sites nested within sessions (start and end of dry season), with 10 000 permutations, to assess whether the sites had similar enough habitat characteristics to be considered as suitable replicates in other analyses.

NDVI

NDVI was averaged over a 60 m radius circular buffer for each trap and then compared across sessions and sites using a GLM Analysis of Variance, with the site nested within session. To meet *ANOVA* prerequisites, the response variable was log-transformed; the analysis was followed by Tukey multiple comparisons of means post-hoc tests.

Body condition

We calculated an index of body condition (mass/length), to assess the state of each

individual's fat reserves (Schulte-Hostedde et al., 2005). To test whether breeding

status, age, habitat quality, increased dryness, and/or interspecific differences

influence body condition of individuals trapped in semi-arid locations, an *ANOVA* was

used with body condition as a response variable, site as a random factor and session,

species, session*species, breeding status, age class and NDVI as explanatory

variables. Assumptions of normality and homoscedasticity of residuals were checked

using diagnostic plots (*graphics* package v.3.6.2.). A Tukey multiple comparisons of

means post-hoc test was applied to determine which inter-level differences were

driving the significance of each factorial effect having more than two modalities.

Age classes

Trapped *Rhabdomys* were assigned to age categories: juveniles, subadults and adults

based on their general appearance, size and/or breeding status. Body length of

trapped individuals varied between 5 and 12 cm. The body length of adult individuals

involved in the physiological study varied between 7.7 and 12 cm. To distinguish

between older and younger adults, adult mass was plotted against body length

(**Supplementary figure 8**), showing a curve defined by an allometric growth equation

with a steeper slope at the highest and lowest body length values. Since bone growth

can significantly alter the concentration of some physiological markers, and growth rate

varies with age/body length, we considered 4 length/age classes (Class A: [7.7:9 cm[;

Class B : [9:10 cm[; Class C : [10:11 cm[; Class D [11 cm:]).

Physiology

We performed a Principal Component Analysis to identify outliers in the data set that

could be potential measure errors. We identified twelve outliers that we removed from

further analyses. These outliers were animals from which we collected smaller volume

of blood samples than recommended for the Vetscan or were heavily hemolyzed and

flagged as such in the Vetscan analysis output. In addition, as total protein values were

used by the Vetscan analyzer to infer the Globulin (GLOB) fraction of blood protein

(with GLOB=TP-ALB), we considered this marker as not informative and removed this

variable from our analyses. Finally, for 199 out of 257 samples, blood concentrations

of creatinine (CRE) were flagged as being below the instrument's sensitivity threshold;

we hence excluded CRE from subsequent analyses, as this imprecision would have

hindered analytical performance.

Due to covariance in variables such as blood physiological markers, multivariate

statistical techniques provide a unique insight into the main patterns and effects driving

variation in the data. Thus, we performed a multivariate analysis with 236 blood

samples, 12 physiological parameters as response variables, site as a random factor,

and session, species, body condition, age class, and NDVI as explanatory variables.

As with habitat, multivariate normality was violated in our physiological dataset, so

permutational multivariate analysis of variance (*PERMANOVA*) was performed, using

10 000 permutations. A backwards stepwise model selection was performed to retain

the most parsimonious model. The best-fitting model was determined by comparing all

previous models, from least to most complex, using the Corrected Akaike's Information

Criterion (AICc) for *PERMANOVA* (package *AICcPermanova* version 0.0.2). We then

sought to identify which of the 12 physiological markers drove significantly the

between-group differences evidenced with the *PERMANOVA*. Twelve univariate

ANOVA tests were performed, initially using the best *PERMANOVA* model followed by

backwards stepwise model selection. Assumptions of normality and homoscedasticity

of residuals were checked using diagnostic plots. In the case of total bilirubin, a

permutation test had to be performed instead of the *ANOVA*. Through a backwards

stepwise model selection process, the final model retained was based on the Akaike's

Information Criterion (AIC). For all tests, the significance level (α) was set at 0.05.

Results

Variation in habitat characteristics between species and sessions

Vegetation composition

We compared the vegetation characteristics surrounding each successful trap with a

PERMANOVA, using the first 5 Principal Component coordinates (**Supplementary**

Tables 7 & 9) of each trap as response variable, and site and session as explanatory

variable. For 2x2m quadrats, PC1 was most explained by variance in dry grass (39.85

%), green bushes (35.63 %) and uncovered surfaces (20.59 %), PC2 was most

explained by variance in the presence of green grass (62.04 %), PC3 by presence of

holes in the ground (60.13 %), PC4 by presence of dry bushes (74.2 %) and PC5 by

presence of succulent plants (76.56 %). For 10x10m quadrats, PC1 was most

explained by variance in dry grass (32.14 %), green bushes (26.16 %), uncovered

surfaces (15.91 %) and dry bushes (10.96 %); PC2 was most explained by green grass

(47.24 %), dry grass (20.85 %), holes (13.59 %), and uncovered surfaces (11.77 %);

PC3 by succulent plants (53.34 %), PC4 by dry bushes (34.28 %), uncovered surfaces

(28.32 %) and succulent plants (23.96 %); and PC5 by holes (43.92 %), green bushes

(25.85 %) and succulent plants (18.69 %). We detected significant differences in

overall habitat characteristics between sessions (2x2m and 10x10m: p<0.001), and

sites nested within sessions (2x2m and 10x10m: p<0.001), indicating clear site differences and seasonal effects on habitat structure (**Supplementary Figures 6 & 7**). Although a permutational dispersion test showed these patterns could be attributable in part to differences in dispersion between sessions or sites (see **Supplementary Material**), some general trends could be inferred from the data. Indeed, *R.d.dilectus* quadrats tended to feature higher proportions of grass (2x2 : 83.99 %, 10x10 : 71.46 %) than *R.bechuanae* quadrats (2x2 : 54.30 %, 10x10 : 53.16 %). Conversely, *R.bechuanae* quadrats tended to feature higher proportions of bushes (2x2 : 31.07 %, 10x10 : 27.35 %) than *R.d.dilectus* quadrats (2x2 : 7.33 %, 10x10 : 10.49 %). Furthermore, quadrats surveyed at the end of the dry season featured more dry grass (2x2 : 74.65 %, 10x10 : 50.15 %) than at its onset (2x2 : 50.75 %, 10x10 : 40.92 %).

NDVI

The log-transformed NDVI of sites sampled at the two sessions (2 sites per species), extracted from satellite data, was also analysed between sites and sessions using *ANOVA*. Site (F=115.385, df=3, p<0.001) and session (F=442.757, df=1, p<0.001), and the interaction between these two variables (F=7.113, df=3, p<0.001) contributed significantly to explain the NDVI variation (**Table 3**). A post-hoc Tukey multiple comparisons of means test showed that all sites had a significantly lower NDVI in September compared to May (**Figure 4**), indicating a poorer state of the vegetation at the end of the dry season. Furthermore, while these study locations were chosen in a region with similar semi-arid bioclimatic features, Gariep Dam and Kalkfontein (where *R.bechuanae* was targeted) seemed to stand out as having a lower NDVI than all other sites sampled in both May and in September, with Wolwespruit and Bloemhof (where *R.d.dilectus* was targeted) having the highest NDVI (**Figure 4**). Therefore, to take into account the impact of variations in NDVI between the two species trapping locations

we included site as a random factor in analyses addressing body and physiological conditions.

Variation of Body condition

An *ANOVA* assessed the influence of session, species, breeding status, age class and NDVI on body condition, including site as a random factor. The most parsimonious model included the significant effects of breeding status and age class (both p<0.001, **Table 4**). Non-breeding females had lower body condition than all other categories (post-hoc pairwise t-tests all p<0.05); body condition was also significantly higher for bigger/older individuals (**Supplementary table 11**).

Variation in physiological responses

A *PERMANOVA* was performed using the distance matrix of 12 physiological markers as response variable and session, species, breeding status, body condition, age class, NDVI and site (random factor) as explanatory variables (**Supplementary table 12**).

The most parsimonious model **(Table 5)** indicated an overall difference in physiological parameter concentrations between *R.bechuanae* and *R.d.dilectus* (p=0.001), as well as an influence of breeding status (p=0.008), that differed between the two species (interaction effect, p=0.005). Session (p=0.062), and body condition (p=0.074) verged on overall significance and were also included in the model, indicating that physiological markers overall tend to shift between the start and end of the dry season, and to vary with body condition. Importantly, both species showed the same pattern of variation between sessions (no significant species*session interaction effect, **Supplementary Table 12**).

Univariate *ANOVAs* performed for each of the 12 physiological markers, with the initial

model being the one retained in the *PERMANOVA* analysis, showed that maximum

parsimony was achieved for 9 of the physiological markers without the site random

effect, suggesting little impact of inter-site variations on these markers.

Six physiological markers revealed differences between the start and the end of the

dry season (**Figure 5**). Blood concentrations of markers of nutrition and liver function

(albumin, potassium, alanine aminotransferase, alkaline phosphatase, and total

bilirubin) were significantly lower at the end compared to the onset of the dry season

(**Table 4**). In contrast, globulin levels varied in the opposite direction (higher at the end

compared to the onset of the dry season).

The levels of five markers were significantly lower in *R.bechuanae* and *R.d.dilectus*

(R^2=0.12, p<0.001), including total bilirubin, alkaline phosphatase (ALP), sodium and

potassium (**Figure 6**). However, amylase showed the opposite pattern between the

species for all breeding statuses except breeding males (**Figure 7**).

Regardless of species, ALP levels were negatively correlated with body condition,

while the levels of phosphorus exhibited the opposite pattern. Finally, the levels of

glucose varied significantly with breeding statuses, being significantly lower in breeding

females compared with non-breeding males ($p=2*10^{-4}$) and tended to be lower than in

non-breeding males and breeding females (**Supplementary figure 9**).

Discussion

We asked how seasonal variation of dry conditions influenced the physiological

response of semi-arid parapatric populations of two sister species that evolved under

distinct environmental conditions (du Toit et al., 2012).

Our study took place in a relatively wet year (La Niña). During the El Niño Southern

Oscillation phenomenon, there is a general association between regional wetness and

sea surface temperatures in the neighbouring Atlantic and Indian Oceans, leading to

wetter episodes throughout Southern Africa during La Niña years (Nicholson & Selato,

2000). Vegetation growth and senescence usually follow seasonal rainfall patterns;

above-average rainfall widens the temporal window for vegetation growth, improving

habitat conditions at the start of the typical dry season (Anyamba et al., 2002). Despite

this, and consistent with our predictions, milder dry conditions did alter habitat quality

of our study sites (as attested by shifts in NDVI and habitat structure).

Consistent with our prediction, we observed seasonal variation of the physiological

response in both species related to energetic resource depletion at the end of the dry

season. Additionally, as predicted, irrespective of season, we found differences in the

physiological responses between species.

Water and nutritional stress are expected to impact the body condition of individuals

(Fuller et al., 2021). However, the body condition of the species did not vary between

the two study sessions, which may relate to the wetter conditions the sampling year.

As expected, body condition varied with breeding status, and several physiological

markers varied significantly with body condition or breeding status, indicating their

association with physiological processes involved in reproduction and body mass

regulation. Reproduction in mammals has direct physiological costs, associated with

increased energy, protein and calcium demands and indirectly through compensatory

effects such as reduction in thermogenesis, immune function and physical activity

(Speakman, 2008; Rintoul & Bringham, 2014; Stawski & Rojas, 2016; Schoepf et al.,

2017b). In support, our results indicate lower levels of blood glucose in breeding

females compared with all other individuals. Moreover, alkaline phosphatase (ALP)

was significantly elevated in non-breeders compared to breeders. Since non-breeders

were mostly the smallest adults, greater ALP levels might relate to soft tissue and/or

bone growth in the latter (Walker et al., 1990). Body condition was generally better in

larger individuals (**Supplementary Table 11**), and ALP was as expected negatively

correlated with body condition. Individuals with the lowest body condition, mostly

smaller individuals, had lower phosphorus levels. High phosphorus levels can be

observed in dehydrated individuals in which osmolality is high; indeed, low phosphorus

levels related to survival in *R.pumilio* at the peak of the dry season, where water

balance was hardest to maintain (Schoepf et al., 2017). However, blood levels of

phosphorus also depend on its consumption and hence availability in the food (Berndt

& Kumar, 2009). Phosphorus reserve depletion in smaller-sized/younger adults might

therefore be consistent with the role of phosphorus intake as a promoter of bone and

soft tissue growth (Shapiro & Heaney, 2003). Furthermore, higher needs for

phosphorus in smaller adults might be hindered by reduced access to food. Indeed,

foraging activity tends to vary between individuals according to size and breeding

status, possibly resulting in larger breeding individuals gaining access to higher quality

food (Ebensperger & Hurtado, 2005).

Seasonal variation in physiology

Despite our study year being relatively wet, milder dry conditions impacted kidney and

liver blood marker concentrations in *R.bechuanae* and *R.d.dilectus*. These changes

involved 6 out of 12 tested physiological markers, and included markers expected to

vary in periods of dietary restriction. Contrary to our prediction, we did not find a

significant interactive effect between session and species, suggesting that the semi-

arid populations of the two species were similarly affected by seasonal dry conditions.

However, a follow up study during an El Niño year may be necessary before one can

conclude.

Blood protein levels are generally good indicators of physiological condition (Tothova

et al., 2016). They comprise two distinct and major components: albumin (ALB) and

globulins (GLOB). In our study, ALB was significantly lower at the end than at the start

of the dry season. ALB is synthesized in the liver and involved in the transport of

bilirubin and several hormones. Blood ALB levels are less prevalent in malnourished

individuals (Walker et al., 1990), supporting the hypothesis of reduction of quality

and/or availability of food at the end of the dry season. The Nubian ibex *Capra nubiana*,

a desert mammal, also displays low ALB during the dry season (Al-Eissa et al., 2012),

which could reflect poor nutritional condition, kidney dysfunction leading to poor water

balance and high osmolality, or liver dysfunction causing low ALB synthesis. We found

that GLOB was significantly higher at the end than at the start of the dry season. GLOB

is a family of molecules including enzymes, immunity proteins and blood proteins

carriers (including transferrin). An increase in GLOB levels usually indicates elevation

of immunoglobulins and hence could suggest heightened immune activity. Greater

antibody production occurs in response to a higher prevalence of respiratory infections

during the dry season, e.g. in the Nubian goat (Abdelatif et al., 2009), and/or reduced

protein nutritional status in young birds (Lochmiller et al., 1993). Higher GLOB levels,

indicating a strained immune system, were also reported for *R.pumilio* individuals that

did not survive the dry season (Schoepf et al., 2017). An increase in blood GLOB

concentrations may also reflect high levels of transferrin (Walker et al., 1990), a major

GLOB that may accumulate in the blood in cases of severe iron deficiency (Kasvosve

& Delanghe, 2002). Hence, elevated blood GLOB levels at the end of the dry season

may indicate either or both depletion in iron resource (transferrin) and higher

vulnerability to infection (immunoglobulin), both resulting from or being aggravated by reduced access to protein. Lower levels of alkaline phosphatase (ALP), a liver enzyme, at the end of the dry season in *Rhabdomys*, is also evidence of malnutrition and deficiency in essential nutrients (proteins, magnesium, zinc) (Yousef et al., 2002; Saraç & Saygili, 2007; Ray et al., 2017). Alanine aminotransferase (ALT), another marker of liver function, also occurred at significantly lower concentrations at the end compared to the start of the dry season, indicating a reduction in liver activity (Walker et al., 1990). During malnutrition, ALT levels could be low (Le Couteur et al., 2010). Indeed, as environmental resources become scarce, essential nutrients, such as pyridoxine (or vitamin B6), that constitute some ALT coenzymes, may be less available, leading to a reduction of ALT blood levels (Vespasiani-Gentilucci et al., 2018).

Acute starvation typically causes bilirubin (TBIL) to accumulate in the blood, as the enzymatic process for conversion of heme into TBIL is stimulated (Thaler et al., 1972). Yet, TBIL was lower at the end compared to the onset of the dry season, rejecting the starvation hypothesis.

Lower TBIL concentrations at the end of the dry season in *Rhabdomys* could stem from low hemoglobin counts because the body produces fewer red blood cells than usual; low hemoglobin can be induced by several factors, among which dietary factors such as iron deficiencies (Clark, 2008). Iron deficient anemia is also supported by higher GLOB concentrations observed at the end of the dry season. Infection can likewise be a cause of both decreased TBIL levels (Zhao et al., 2019) and higher GLOB concentrations at the end of the dry season. Finally, malnutrition and/or lack of potassium intake are among the main potential causes of the reduction of its levels in

the blood (Their, 1986), which could explain why they were the lowest at the end of the

dry season, when the available resources were lowest, in our study.

Overall, our results indicate that seasonal variation of physiological markers could

reflect reduced primary productivity throughout the dry season, which is coherent with

our observation of habitat degradation (lower NDVI) at the end of the dry season. Blood

concentrations of various markers seem to indicate difficulties in maintaining nutritional

functions and the necessity of a higher immune output, caused or exacerbated by food

restriction. However, since glucose levels were maintained during the dry season,

shifts in six physiological markers related to nutrition and immunity during the milder

than usual dry season suggest a reduction of liver function due to malnutrition, rather

than starvation.

Individuals undergoing malnutrition during dry conditions are expected to show low

blood glucose concentrations, and deficiencies in essential nutrients, leading to

pathologies such as anemia (Gordon et al., 1988). Indeed, animals foraging on soils

potentially poor in nutrients such as iron, zinc, and magnesium are more prone to

malnutrition (Graham, 1991; Gupta et al., 2008). Higher GLOB, lower TBIL and lower

ALP blood concentrations in our study might relate to such deficiencies.

We did not observe seasonal variation of sodium or blood urea nitrogen, blood markers

of kidney function or osmoregulation. Even if such effects may have been mitigated by

the relatively wet conditions, reduced environmental moisture at the end of the dry

season could have caused an overall increase in blood osmolality, increasing blood

concentrations of most biomarkers. In contrast, most markers exhibited a significant

reduction in blood concentrations. This tends to indicate that water availability does not

worsen as strongly as food availability for striped mice throughout the dry season in

this semi-arid environment.

The two species responded similarly to increased dryness, while maintaining stable

body conditions and levels of energy (glucose), a proxy of fat reserves. *R.pumilio*

individuals tended to reduce their energy expenditure and physical activity during

periods of limited food availability (Rimbach, Blanc et al., 2018; Rimbach, Jäger et al.,

2018). The semi-arid populations of *R.bechuanae* and *R.d.dilectus* could display a

similar behavioral strategy, which could explain their relatively stable levels of glucose

during the dry season.

Interspecific differences

R.bechuanae is thought to have evolved in drier environments than *R.d.dilectus*.

Moreover, in the semi-arid region where they co-occur, *R.bechuanae* populations

occupy drier habitats than *R.d.dilectus*. While body condition did not differ significantly

between species, some interspecific contrasts support our hypothesis that

R.bechuanae may have developed a better physiological capacity to cope with semi-

arid conditions than *R.d.dilectus*.

Among the markers exhibiting a significant difference between *R.bechuanae* and

R.d.dilectus, all but amylase showed higher blood concentrations in *R.d.dilectus*.

Amylase is positively correlated with digestive activity and reflects differences in

feeding status or diet composition between the species (Hidalgo et al., 1999). Given

the role of alpha-amylase in digestion, lower blood amylase levels in *R.d.dilectus* than

R.bechuanae may indicate lower digestive activity due to reduced access to food, or

lower starch content in the diet. Pajic et al. (2019) found a correlation between dietary

starch content and the number of amylase gene copies in several mammalian

genomes, even in different habitats and with different diets. Since amylase activity and amylase gene expression are directly correlated with amylase gene copy number (Arendt et al., 2014), further studies could elucidate the potential genetic basis of the different levels of amylase in our two study species.

Sodium (NA) and potassium (K) concentrations, which depend on both dietary and water intake, were higher in *R.d.dilectus* than in *R.bechuanae*. These differences, coupled with higher ALP and TBIL concentrations in *R.d.dilectus* compared to *R.bechuanae*, could have been induced by nutritional deficiencies, which would suggest a better nutrition state for *R.d.dilectus* than *R.bechuanae*. However, higher levels of amylase in *R.bechuanae* cast doubt on this hypothesis. Instead, lower levels of blood NA, K, ALP and TBIL, would be more parsimoniously attributed to lower osmolality due to better water balance in *R.bechuanae*. Indeed, detection of water balance shifts may be muddled by the opposing effects of malnutrition and water constraints, as both influence osmolality (Zhang et al., 2022). A better overall ability of *R.bechuanae* to conserve body moisture could rely on physiological mechanisms such as more efficient excretion or osmoregulation, or through feeding on a diet optimizing water intake. We hypothesize that during the evolutionary history of *R.bechuanae*, selection pressure may have favored specific strategies allowing an overall more efficient water regulation compared to *R.d.dilectus*. These results, combined with the reduction in the levels of K, ALP and TBIL at the end of the dry season suggest the species are exposed to both water and nutrition stress.

We found relatively little evidence that the two species suffer differentially from lack of food in their semi-arid environment during a La Niña phenomenon. Overall, energy levels, as indicated by blood glucose, and body condition, did not vary between the

species. Instead, it seems that the overall differences between the two species can be accounted for by lower blood concentrations in *R.bechuanae*, and on average higher blood concentrations in key metabolites in *R.d.dilectus,* possibly due to overall poorer water regulation. High average blood amylase levels in *R.bechuanae* in the dry season might indicate an interspecific difference in diet composition or food intake, but not of protein-energy malnutrition. Little is known about the *R.bechuanae* and *R.d.dilectus* diets, although the genus was characterised as a generalist feeding on seeds and other plant material, and insects (Curtis & Perrin, 1979). Some studies show variation in diet composition within *Rhabdomys* genus associated with variation in local resources (Taylor & Green, 1976). The two species could have adjusted their diet to the respective local availabilities of different food resources within their home ranges. For example, in the semi-arid zone, arthropod biomass and diversity is affected by seasonal patterns in rainfall. Dalerum et al. (2017) found lower arthropod trapping success and diversity during the dry season, in shrub habitats compared to any other biomes, in Benfontein Game Reserve, one of our study sites. Thus, compared to seeds, the reliability of arthropods as a food source during dry conditions could be lower in shrublands, inhabited by *R.bechuanae* and by both species in parapatry, than in grasslands inhabited by *R.d.dilectus* in allopatry. Variation in seed composition could also explain this interspecific difference in amylase concentrations. Indeed, granivores prefer seeds that have higher protein content and lower secondary metabolites but may be forced to take a wider range of seeds in dry areas than in mesic ones (Wilmer et al., 2009). If the hypothesis of a relatively ancient adaptation of *R.bechuanae* to dry habitats is true, it could be expected that it would also have adjusted its diet to the specific conditions encountered in dry habitats. Within this ecosystem, *R.bechuanae* could forage more efficiently compared with *R.d.dilectus*, which evolved in a more

mesic environment; its diet could also be richer in starch (such as containing more

seeds/grains than leaves or fibre). Higher starch intake could help maintaining water

balance; starch is the best substrate for metabolic water production in a dry

environment (Adolph, 1964). This is consistent with the other interspecific differences

revealed in this study, as diet can also significantly affect osmoregulation (Sabat et al.,

2009).

Differences in blood amylase concentrations between the two species were observed

in all breeding statuses, except for breeding males. This is consistent with findings in

R.pumilio (Scantlebury et al., 2006), a species with similar levels of sociality as

R.bechuanae, where daily energy expenditure is reduced in mice living in groups. In

the solitary *R.d.dilectus*, higher thermoregulatory energy expenditure might reduce

foraging time, allowing only the largest individuals (breeding males) to meet their

foraging needs as efficiently as their *R.bechuanae* counterparts.

Conclusions

We found that two closely related species with different ecological characteristics

displayed similar negative shifts in blood concentrations of markers of nutrition and

liver function in spite of a relatively mild seasonal drying of their habitats. We hence

expect the physiological impact of harsher dry conditions, to affect the individuals'

fitness even more dramatically. The fact that the habitat of *R.bechuanae* studied

populations was the driest at both time points, combined with our findings concerning

their overall better water balance and nutrition than *R.d.dilectus*, point towards

R.bechuanae coping better in the semi-arid environment. If, as projected, the

frequency and length of drought episodes increase in this region, we predict that

R.bechuanae will expand its range at the expense of *R.d.dilectus*. Though adaptation

per se cannot be inferred from our study, this comparative study considering

populations of the two species in the same biogeographic zone adds to the growing

evidence that these cryptic species of striped mice have diverged phenotypically.

These findings support the hypothesis of adaptation to distinct ecological conditions

associated with their divergence. Further studies might seek to identify signatures of

selection in genes linked to crucial physiological functions in dry conditions to

understand the underlying mechanisms of divergence. Moreover, investigation of

seasonal variation in gene expression in the liver and kidney should help identify the

physiological pathways, key to survival, affected by seasonal variation in dry

conditions.

Main tables & figures

Figure 1: Map of known occurrences of 6 main clades of Rhabdomys (translucent dots)

and locations sampled in this study (full colour dots) in southern Africa

Based on published data and unpublished data (origin details are available in doi:

0d1b3414-7e2a-11ea-a38d-00163e26bfb0). Base map: World Topographic Map Esri

Standard, Aridity Index layer was computed from a 0.5° global grid, using data from

the Version 3 of the Global Aridity Index and Potential Evapotranspiration Database

(Zomer et al.,2022).

671 **_Table 1: Summary of main ecological characteristics of R.bechuanae and_**

672 **_R.d.dilectus and our predictions on their physiological response during the_**

673 **_dry season_**

ECOLOGICAL CHARACTERISTICS	R.BECHUANAE	R.D.DILECTUS
ENVIRONMENTAL NICHE	Arid to semi-arid (Ganem et al., 2020)	Mesic to semi-arid (Ganem et al., 2020)
PREFERRED HABITAT	Bushy patches (Dufour et al., 2015)	Continuous cover / grassland (Dufour et al., 2015)
SOCIAL STRUCTURE	Group-living (Dufour et al., 2018)	Solitary / opportunistic group-living (Dufour et al., 2018)
SPECIES-SPECIFIC RESPONSE TO SEASONAL VARIATIONS OF DRY CONDITIONS: PREDICTIONS		
BODY CONDITION	↘↘ or ↘*	↘↘
NUTRITIONAL METABOLISM	↘↘ or ↘*	↘↘
LIVER ACTIVITY	↘↘ or ↘*	↘↘
WATER BALANCE	↘↘ or ↘*	↘↘
SENSITIVITY TO INFECTION	↗↗ or ↗*	↗↗
ASEASONAL DIFFERENCES: PREDICTIONS		
BODY CONDITION	+	-
LIVER ACTIVITY	+	-
WATER BALANCE	+	-
SENSITIVITY TO INFECTION	-	+

674

675 *Adaptation to harsher arid conditions could have enabled _R.bechuanae_ to withstand

676 growing aridity without suffering as many adverse effects as _R.d.dilectus_ in the semi-

677   arid zone. Alternatively, semi-arid populations of both species could both have

678   developed local adaptation.

681   ***Figure 2: Graphical summary of organization and composition of data used for***

682   ***analysis in this study.*** Figures between brackets indicate number of modalities or types

683   of categories included in the factor.

Figure 3: Total monthly precipitation for all sampled sites as recorded in 2022.

Table 2: Geographical position and Aridity Index of sampled sites for each species

SPECIES	SAMPLED SITES	GEOGRAPHICAL COORDINATES	ARIDITY INDEX[*]
R.bechuanae	Gariep Dam Nature Reserve	E25.55537; S30.57589	0.35-0.49
	Kalkfontein Dam Nature Reserve	E25.28031; S29.52058	0.47-0.61
	Benfontein Nature Reserve**	E24.81824; S28.82747	0.32-0.46
R.d.dilectus	Bloemhof Dam Nature Reserve	E25.65144; S27.66063	0.29-0.41
	Barberspan Bird Sanctuary**	E25.58354; S26.56538	0.34-0.44
	Wolwespruit Nature Reserve	E26.26326; S27.41418	0.43-0.63

690 *Average Aridity index (see **Supplementary material** for formula) ± Standard

691 Error of the Mean, based on climate data collected from 2010 to 2021 (South

692 African Weather Service)

693 ** Sampled only in May 2022 (dry season onset).

Figure 4: Average Normalized Differential Vegetation Index calculated within a

buffer circle of 60m radius around each successful trap per site and session.

698 *NDVI was retrieved from the Copernicus Open Access Data Hub (Copernicus*

699 *Sentinel-2 data [2023]). For each site and session, we retained NDVI data available*

700 *for the closest day to the beginning of a sampling session, expected to represent the*

701 *conditions experienced by the mice at the time of capture.*

702 *Table 3:* **Results of the Analysis of Variance (response variable: NDVI)**

Variable	Df	R²	F	P
Session	1	0.445	442.757	**< 2.2*10⁻¹⁶**
Site	3	0.348	115.385	**< 2.2*10⁻¹⁶**
Session*Site	3	0.021	7.113	**1.512*10⁻⁴**
Residual	229	0.185		

704 *Table 4:* **Results of the Analysis of Variance (response variable : body**
705 **condition)**

Variable	Df	R^2	F	P
Age class	3	0.151	14.665	**$9.081*10^{-9}$**
Breeding status	3	0.065	6.309	**$3.952*10^{-4}$**
Residual	229	0.784		

706

707 *Table 5: Summary of the results of the best fitting Permutational multivariate*

708 *analysis of variance model, addressing variation in physiological response*

709 *(site= random factor)*

710

Variable	Df	R^2	Pseudo-F	P
Session	1	0.011	3.000	0.062
Species	1	0.042	11.108	**0.001**
Breeding status	3	0.044	3.874	**0.008**
Body condition	1	0.010	2.710	0.074
Species * Breeding status	3	0.047	4.202	**0.005**
Residual	226	0.846		

711

712

713

714

715

716

717

Table 6: summary of the results of univariate analyses of variance addressing

variation in responses of each of the 12 studied physiological variables (best

fitting models).

Physiological marker	Variable(s) kept in Best Model and their significance	adjusted R²	F	df Model	df Residual	P (model)
Albumin	**Session**[***], Reproductive status, body condition	0.097	6.048	5	230	2.786×10^{-5}
Alkaline phosphatase	**Session**[***]**, Species**[*]**, Reproductive status**[***]**, Body condition**[***]	0.321	19.390	6	228	$< 2.2 \times 10^{-16}$
Alanine aminotransferase	**Session**[*], Species, Reproductive status, Species*Reproductive Status	0.060	2.884	8	227	0.005
Amylase	**Species**[***], Reproductive status, body condition, **Species*Reproductive Status**[*]	0.196	8.164	8	227	1.079×10^{-9}
Calcium	Species	0.009	3.095	1	234	0.08
Globulin	**Session**[***], Body condition	0.300	51.450	2	233	$< 2.2 \times 10^{-16}$
Glucose	Session[·], **Reproductive status**[**]	0.085	6.481	4	231	5.849×10^{-5}
Potassium	**Session**[*]**, Species**[*]	0.068	9.569	2	233	0.0001
Sodium	Session, **Species**[***], Reproductive status, Body condition	0.125	6.614	6	229	1.83×10^{-6}
Blood Urea Nitrogen	*Null Random effect model*	0.072				/
Phosphorus	Session, Species, Reproductive status, **Body condition**[***] + random	0.294				(χ^2 test vs. null model) 9.838×10^{-9}

724   *Note: The full models, before backwards elimination, included session, species, body condition,*

725   *reproductive status and reproductive status*species as variables, and site as a partially-crossed*

726   *random factor nested within species.*

727   **Bold:** *significant effects (p < 0.05) following Bonferroni correction with:*

728   ****: p < 0.001; **: 0.001 < p < 0.01 ; *: 0.01 < p < 0.05*

Figure 5: Significant shifts in blood physiological marker concentrations

between the onset and the end of the dry season (mean ± SE)

Figure 6: Significant interspecific differences involving four blood physiological marker concentrations (mean ± SE)

744   ***Figure 7: Variation of blood amylase concentration between species and***

745   ***breeding status (mean ± SE)***

References

Abaxis, Inc. (2009). VS2 Operator's Manual. Retrieved from https://www.zoetis.es/_locale-assets/spc/vetscan-vs2-analizador.pdf

Abdelatif, A. A., Ibrahim, M. Y., & Hassan, Y. Y. (2009). Seasonal variation in erythrocytic and leukocytic indices and serum proteins of female Nubian goats. Middle-East J. Sci. Res, 4(3), 168-174. Retrieved from https://idosi.org/mejsr/mejsr4(3)/9.pdf

Abraham, J. O., Hempson, G. P., & Staver, A. C. (2019). Drought-response strategies of savanna herbivores. Ecology and Evolution, 9(12), 7047–7056. https://doi.org/10.1002/ece3.5270

Adolph, E. F. (1964). Desert Animals: Physiological Problems of Heat and Water. Knut Schmidt-Nielsen. Physiological Zoology, 37(3), 338–339. https://doi.org/10.1086/physzool.37.3.30152404

AL Eissa, M. S., Saad, A., Al Farraj, S. A., Saud, A. A., Al Dahmash, B., & Hamad, A. Y. (2012). Seasonal variation effects on the composition of blood in Nubian ibex (Capra nubiana) in Saudi Arabia. African Journal of Biotechnology, 11(5), 1283–1286. https://doi.org/10.5897/ajb11.2004

Anyamba, A., Tucker, C. J., & Mahoney, R. (2002). From El Niño to La Niña: Vegetation Response Patterns over East and Southern Africa during the 1997–2000 Period. Journal of Climate, 15(21), 3096–3103. https://doi.org/10.1175/1520-0442(2002)015<3096:fenotl>2.0.co;2

Arendt, M., Fall, T., Lindblad-Toh, K., & Axelsson, E. (2014). Amylase activity is associated with AMY2B copy numbers in dog: implications for dog domestication, diet and diabetes. Animal Genetics, 45(5), 716–722. https://doi.org/10.1111/age.12179

Berkel, C., & Cacan, E. (2022). The differential influence of ambient temperature on

heart rate in male and female wild pale-throated sloths (Bradypus tridactylus).

Research Square Platform LLC. http://dx.doi.org/10.21203/rs.3.rs-1908943/v1

Berndt, T., & Kumar, R. (2009). Novel Mechanisms in the Regulation of Phosphorus

Homeostasis. Physiology, 24(1), 17–25. https://doi.org/10.1152/physiol.00034.2008

Bittner, N. K. J., Mack, K. L., & Nachman, M. W. (2021). Gene expression plasticity

and desert adaptation in house mice*. Evolution, 75(6), 1477–1491.

https://doi.org/10.1111/evo.14172

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to

measuring local adaptation. Ecology Letters, 16(9), 1195–1205.

https://doi.org/10.1111/ele.12150

Boyers, M., Parrini, F., Owen-Smith, N., Erasmus, B. F. N., & Hetem, R. S. (2021).

Contrasting capabilities of two ungulate species to cope with extremes of aridity.

Scientific Reports, 11(1). https://doi.org/10.1038/s41598-021-83732-w

Christian, K. A., Griffiths, A. D., & Bedford, G. S. (1996). Physiological ecology of

frillneck lizards in a seasonal tropical environment. Oecologia, 106(1), 49–56.

https://doi.org/10.1007/bf00334406

Clark, S. F. (2008). Iron Deficiency Anemia. Nutrition in Clinical Practice, 23(2), 128–

141. https://doi.org/10.1177/0884533608314536

Curtis, B. A., & Perrin, M. R. (1979). Food preferences of the vlei rat (Otomys Irroratus)

and the four-striped mouse (Rhabdomys pumilio). South African Journal of Zoology,

14(4), 224–229. https://doi.org/10.1080/02541858.1979.11447675

Dalerum, F., de Vries, J. L., Pirk, C. W. W., & Cameron, E. Z. (2017). Spatial and

temporal dimensions to the taxonomic diversity of arthropods in an arid grassland

savannah. Journal of Arid Environments, 144, 21–30.

https://doi.org/10.1016/j.jaridenv.2017.04.002

DEA (2018). South Africa's Third National Communication under the United Nations

Framework Convention on Climate Change. Retrieved from

https://unfccc.int/sites/default/files/resource/South%20African%20TNC%20Report%2

0%20to%20the%20UNFCCC_31%20Aug.pdf

du Toit, N., Jansen van Vuuren, B., Matthee, S., & Matthee, C. A. (2012). Biome

specificity of distinct genetic lineages within the four-striped mouse Rhabdomys

pumilio (Rodentia: Muridae) from southern Africa with implications for taxonomy.

Molecular Phylogenetics and Evolution, 65(1), 75–86.

https://doi.org/10.1016/j.ympev.2012.05.036

Dufour, C. M. S., Meynard, C., Watson, J., Rioux, C., Benhamou, S., Perez, J., du

Plessis, J. J., Avenant, N., Pillay, N., & Ganem, G. (2015). Space Use Variation in Co-

Occurring Sister Species: Response to Environmental Variation or Competition? PLOS

ONE, 10(2), e0117750. https://doi.org/10.1371/journal.pone.0117750

Dufour, C. M. S., Pillay, N., Avenant, N., Watson, J., Loire, E., & Ganem, G. (2018).

Habitat characteristics and species interference influence space use and nest-site

occupancy: implications for social variation in two sister species. Oikos, 128(4), 503–

516. https://doi.org/10.1111/oik.05357

Ebensperger, L. A., & Hurtado, M. J. (2005). On the Relationship between Herbaceous

Cover and Vigilance Activity of Degus (Octodon degus). Ethology, 111(6), 593–608.

https://doi.org/10.1111/j.1439-0310.2005.01084.x

Feder, M. E., & Block, B. A. (1991). On the Future of Animal Physiological Ecology.

Functional Ecology, 5(2), 136. https://doi.org/10.2307/2389251

Fuller, A., Mitchell, D., Maloney, S. K., & Hetem, R. S. (2016). Towards a mechanistic

understanding of the responses of large terrestrial mammals to heat and aridity

associated with climate change. Climate Change Responses, 3(1).

https://doi.org/10.1186/s40665-016-0024-1

Fuller, A., Mitchell, D., Maloney, S. K., Hetem, R. S., Fonsêca, V. F. C., Meyer, L. C.

R., van de Ven, T. M. F. N., & Snelling, E. P. (2021). How dryland mammals will

respond to climate change: the effects of body size, heat load and a lack of food and

water. Journal of Experimental Biology, 224(Suppl_1).

https://doi.org/10.1242/jeb.238113

Ganem, G., Dufour, C. M. S., Avenant, N. L., Caminade, P., Eiseb, S. J., Tougard, C.,

& Pillay, N. (2020). An update on the distribution and diversification of Rhabdomys sp.

(Muridae, Rodentia). Journal of Vertebrate Biology, 69(2), 1.

https://doi.org/10.25225/jvb.20013

Ganem, G., Meynard, C. N., Perigault, M., Lancaster, J., Edwards, S., Caminade, P.,

Watson, J., & Pillay, N. (2012). Environmental correlates and co-occurrence of three

mitochondrial lineages of striped mice (Rhabdomys) in the Free State Province (South

Africa). Acta Oecologica, 42, 30–40. https://doi.org/10.1016/j.actao.2012.01.003

Gordon, G., Brown, A. S., & Pulsford, T. (1988). A koala (Phascolarctos cinereus

Goldfuss) population crash during drought and heatwave conditions in south-western

Queensland. Australian Journal of Ecology, 13(4), 451–461.

https://doi.org/10.1111/j.1442-9993.1988.tb00993.x

Graham, T. W. (1991). Trace Element Deficiencies in Cattle. Veterinary Clinics of North

America: Food Animal Practice, 7(1), 153–215. https://doi.org/10.1016/s0749-

0720(15)30816-1

Gupta, U. C., Wu, K., & Liang, S. (2008). Micronutrients in Soils, Crops, and Livestock.

Earth Science Frontiers, 15(5), 110–125. https://doi.org/10.1016/s1872-

5791(09)60003-8

Hidalgo, M. C., Urea, E., & Sanz, A. (1999). Comparative study of digestive enzymes

in fish with different nutritional habits. Proteolytic and amylase activities. Aquaculture,

170(3–4), 267–283. https://doi.org/10.1016/s0044-8486(98)00413-x

Jury, M. R. (2021). Spreading of the semi-arid climate across South Africa. Journal of

Water and Climate Change, 12(8), 3734–3749. https://doi.org/10.2166/wcc.2021.187

Kasvosve, I., & Delanghe, J. (2002). Total Iron Binding Capacity and Transferrin

Concentration in the Assessment of Iron Status. Clinical Chemistry and Laboratory

Medicine, 40(10). https://doi.org/10.1515/cclm.2002.176

Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. Ecology

Letters, 7(12), 1225–1241. https://doi.org/10.1111/j.1461-0248.2004.00684.x

Kobbe, S., Ganzhorn, J. U., & Dausmann, K. H. (2010). Extreme individual flexibility of

heterothermy in free-ranging Malagasy mouse lemurs (Microcebus griseorufus).

Journal of Comparative Physiology B, 181(1), 165–173.

https://doi.org/10.1007/s00360-010-0507-5

Kruger, A. C. (2006). Observed trends in daily precipitation indices in South Africa:

1910–2004. International Journal of Climatology, 26(15), 2275–2285.

https://doi.org/10.1002/joc.1368

Le Couteur, D. G., Blyth, F. M., Creasey, H. M., Handelsman, D. J., Naganathan, V.,

Sambrook, P. N., Seibel, M. J., Waite, L. M., & Cumming, R. G. (2010). The Association

of Alanine Transaminase With Aging, Frailty, and Mortality. The Journals of

Gerontology Series A: Biological Sciences and Medical Sciences, 65A(7), 712–717.

https://doi.org/10.1093/gerona/glq082

Lian, X., Piao, S., Chen, A., Huntingford, C., Fu, B., Li, L. Z. X., Huang, J., Sheffield,

J., Berg, A. M., Keenan, T. F., McVicar, T. R., Wada, Y., Wang, X., Wang, T., Yang,

Y., & Roderick, M. L. (2021). Multifaceted characteristics of dryland aridity changes in

a warming world. Nature Reviews Earth & Environment, 2(4), 232–250.

https://doi.org/10.1038/s43017-021-00144-0

Lochmiller, R. L., Vestey, M. R., & Boren, J. C. (1993). Relationship between Protein

Nutritional Status and Immunocompetence in Northern Bobwhite Chicks. The Auk,

110(3), 503–510. https://doi.org/10.2307/4088414

López-Maury, L., Marguerat, S., & Bähler, J. (2008). Tuning gene expression to

changing environments: from rapid responses to evolutionary adaptation. Nature

Reviews Genetics, 9(8), 583–593. https://doi.org/10.1038/nrg2398

Masubelele, M. L., Hoffman, M. T., Bond, W. J., & Gambiza, J. (2014). A 50 year study

shows grass cover has increased in shrublands of semi-arid South Africa. Journal of

Arid Environments, 104, 43–51. https://doi.org/10.1016/j.jaridenv.2014.01.011

McClure, D. E. (1999). Clinical Pathology and Sample Collection in the Laboratory

Rodent. Veterinary Clinics of North America: Exotic Animal Practice, 2(3), 565–590.

https://doi.org/10.1016/s1094-9194(17)30111-1

Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic

plasticity: the problem and the evidence. Evolutionary Applications, 7(1), 1–14.

https://doi.org/10.1111/eva.12137

Meynard, C. N., Pillay, N., Perrigault, M., Caminade, P., & Ganem, G. (2012). Evidence

of environmental niche differentiation in the striped mouse (Rhabdomys sp.): inference

from its current distribution in southern Africa. Ecology and Evolution, 2(5), 1008–1023.

https://doi.org/10.1002/ece3.219

Muñoz-Garcia, A., Ben-Hamo, M., Pilosof, S., Williams, J. B., & Korine, C. (2022).

Habitat aridity as a determinant of the trade-off between water conservation and

evaporative heat loss in bats. Journal of Comparative Physiology B, 192(2), 325–333.

https://doi.org/10.1007/s00360-021-01425-2

Naumann, G., Alfieri, L., Wyser, K., Mentaschi, L., Betts, R. A., Carrao, H., Spinoni, J.,

Vogt, J., & Feyen, L. (2018). Global Changes in Drought Conditions Under Different

Levels of Warming. Geophysical Research Letters, 45(7), 3285–3296.

https://doi.org/10.1002/2017gl076521

Naya, D. E., Veloso, C., Sabat, P., & Bozinovic, F. (2011). Physiological flexibility and

climate change: The case of digestive function regulation in lizards. Comparative

Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 159(1),

100–104. https://doi.org/10.1016/j.cbpa.2011.02.005

Nicholson, S. E., & Selato, J. C. (2000). The influence of La Nina on African rainfall.

International Journal of Climatology, 20(14), 1761–1776. https://doi.org/10.1002/1097-

0088(20001130)20:14<1761::aid-joc580>3.0.co;2-w

Ostrowski, S., Williams, J. B., Mésochina, P., & Sauerwein, H. (2005). Physiological

acclimation of a desert antelope, Arabian oryx (Oryx leucoryx), to long-term food and

water restriction. Journal of Comparative Physiology B, 176(3), 191–201.

https://doi.org/10.1007/s00360-005-0040-0

Pajic, P., Pavlidis, P., Dean, K., Neznanova, L., Romano, R.-A., Garneau, D.,

Daugherity, E., Globig, A., Ruhl, S., & Gokcumen, O. (2019). Independent amylase

gene copy number bursts correlate with dietary preferences in mammals. eLife, 8.

https://doi.org/10.7554/elife.44628

Parmesan, C., Root, T. L., & Willig, M. R. (2000). Impacts of Extreme Weather and

Climate on Terrestrial Biota*. Bulletin of the American Meteorological Society, 81(3),

443–450. https://doi.org/10.1175/1520-0477(2000)081<0443:ioewac>2.3.co;2

Post, E., Brodie, J., Hebblewhite, M., Anders, A. D., Maier, J. A. K., & Wilmers, C. C.

(2009). Global Population Dynamics and Hot Spots of Response to Climate Change.

BioScience, 59(6), 489–497. https://doi.org/10.1525/bio.2009.59.6.7

Rimbach, R., Blanc, S., Zahariev, A., Gatta, M., Pillay, N., & Schradin, C. (2018).

Seasonal variation in energy expenditure in a rodent inhabiting a winter-rainfall desert.

Journal of Comparative Physiology B, 188(5), 877–888.

https://doi.org/10.1007/s00360-018-1168-z

Rimbach, R., Jäger, J., Pillay, N., & Schradin, C. (2018). Food Availability Is the Main

Driver of Seasonal Changes in Resting Metabolic Rate in African Striped Mice

(Rhabdomys pumilio). Physiological and Biochemical Zoology, 91(2), 826–833.

https://doi.org/10.1086/696828

Rintoul, J. L. P., & Brigham, R. M. (2014). The influence of reproductive condition and

concurrent environmental factors on torpor and foraging patterns in female big brown

bats (Eptesicus fuscus). Journal of Comparative Physiology B, 184(6), 777–787.

https://doi.org/10.1007/s00360-014-0837-9

Rocha, J. L., Godinho, R., Brito, J. C., & Nielsen, R. (2021). Life in Deserts: The Genetic Basis of Mammalian Desert Adaptation. Trends in Ecology & Evolution, 36(7), 637–650. https://doi.org/10.1016/j.tree.2021.03.007

Sabat, P., Gonzalez-Vejares, S., & Maldonado, K. (2009). Diet and habitat aridity affect osmoregulatory physiology: An intraspecific field study along environmental gradients in the Rufous-collared sparrow. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 152(3), 322–326. https://doi.org/10.1016/j.cbpa.2008.11.003

Saraç, F., & Saygılı, F. (2007). Causes of High Bone Alkaline Phosphatase. Biotechnology & Biotechnological Equipment, 21(2), 194–197. https://doi.org/10.1080/13102818.2007.10817444

Scantlebury, M., Bennett, N. C., Speakman, J. R., Pillay, N., & Schradin, C. (2006). Huddling in groups leads to daily energy savings in free‑living African Four‑Striped Grass Mice, Rhabdomys pumilio. Functional Ecology, 20(1), 166–173. https://doi.org/10.1111/j.1365-2435.2006.01074.x

Schoepf, I., Pillay, N., & Schradin, C. (2016). The pathophysiology of survival in harsh environments. Journal of Comparative Physiology B, 187(1), 183–201. https://doi.org/10.1007/s00360-016-1020-2

Schoepf, I., Pillay, N., & Schradin, C. (2017). Trade-offs between reproduction and health in free-ranging African striped mice. Journal of Comparative Physiology B, 187(4), 625–637. https://doi.org/10.1007/s00360-016-1054-5

Schulte-Hostedde, A. I., Zinner, B., Millar, J. S., & Hickling, G. J. (2005). RESTITUTION OF MASS–SIZE RESIDUALS: VALIDATING BODY CONDITION INDICES. Ecology, 86(1), 155–163. https://doi.org/10.1890/04-0232

Seebacher, F., & Franklin, C. E. (2012). Determining environmental causes of

biological effects: the need for a mechanistic physiological dimension in conservation

biology. Philosophical Transactions of the Royal Society B: Biological Sciences,

367(1596), 1607–1614. https://doi.org/10.1098/rstb.2012.0036

Seebacher, F., Davison, W., Lowe, C. J., & Franklin, C. E. (2005). A falsification of the

thermal specialization paradigm: compensation for elevated temperatures in Antarctic

fishes. Biology Letters, 1(2), 151–154. https://doi.org/10.1098/rsbl.2004.0280

Shapiro, R., & Heaney, R. P. (2003). Co-dependence of calcium and phosphorus for

growth and bone development under conditions of varying deficiency. Bone, 32(5),

532–540. https://doi.org/10.1016/s8756-3282(03)00060-7

Solmsen, N., Johannesen, J., & Schradin, C. (2011). Highly asymmetric fine-scale

genetic structure between sexes of African striped mice and indication for condition

dependent alternative male dispersal tactics. Molecular Ecology, 20(8), 1624–1634.

https://doi.org/10.1111/j.1365-294x.2011.05042.x

Somero, G. N. (2010). The physiology of climate change: how potentials for

acclimatization and genetic adaptation will determine 'winners' and 'losers.' Journal of

Experimental Biology, 213(6), 912–920. https://doi.org/10.1242/jeb.037473

Somero, G. N. (2012). The Physiology of Global Change: Linking Patterns to

Mechanisms. Annual Review of Marine Science, 4(1), 39–61.

https://doi.org/10.1146/annurev-marine-120710-100935

Speakman, J. R. (2007). The physiological costs of reproduction in small mammals.

Philosophical Transactions of the Royal Society B: Biological Sciences, 363(1490),

375–398. https://doi.org/10.1098/rstb.2007.2145

Stawski, C., & Rojas, A. D. (2016). Thermal physiology of a reproductive female

marsupial, Antechinus flavipes. Mammal Research, 61(4), 417–421.

https://doi.org/10.1007/s13364-016-0287-8

Sundar Ray, C., Singh, B., Jena, I., Behera, S., & Ray, S. (2017). Low Alkaline

Phosphatase (ALP) In Adult Population an Indicator of Zinc (Zn) and Magnesium (Mg)

Deficiency. Current Research in Nutrition and Food Science Journal, 5(3), 347–352.

https://doi.org/10.12944/crnfsj.5.3.20

Taylor, K. D., & Green, M. G. (1976). The influence of rainfall on diet and reproduction

in four African rodent species. Journal of Zoology, 180(3), 367–389.

https://doi.org/10.1111/j.1469-7998.1976.tb04683.x

Thaler, M. M., Gemes, D. L., & Bakken, A. F. (1972). Enzymatic Conversion of Heme

to Bilirubin in Normal and Starved Fetuses and Newborn Rats. Pediatric Research,

6(3), 197–201. https://doi.org/10.1203/00006450-197203000-00008

Thornthwaite, C. W. (1948). An Approach toward a Rational Classification of Climate.

Geographical Review, 38(1), 55. https://doi.org/10.2307/210739

Tieleman, B. I., Williams, J. B., Buschur, M. E., & Brown, C. R. (2003). PHENOTYPIC

VARIATION OF LARKS ALONG AN ARIDITY GRADIENT: ARE DESERT BIRDS

MORE FLEXIBLE? Ecology, 84(7), 1800–1815. https://doi.org/10.1890/0012-

9658(2003)084[1800:pvolaa]2.0.co;2

Tothova, C., Nagy, O., & Kovac, G. (2016). Serum proteins and their diagnostic utility

in veterinary medicine: a review. Veterinární Medicína, 61(9), 475–496.

https://doi.org/10.17221/19/2016-vetmed

Tshikae, B. P., Davis, A. L. V., & Scholtz, C. H. (2013). Does an aridity and trophic resource gradient drive patterns of dung beetle food selection across the Botswana Kalahari? *Ecological Entomology, 38(1),* 83–95. https://doi.org/10.1111/j.1365-2311.2012.01409.x

Vespasiani-Gentilucci, U., De Vincentis, A., Ferrucci, L., Bandinelli, S., Antonelli Incalzi, R., & Picardi, A. (2017). Low Alanine Aminotransferase Levels in the Elderly Population: Frailty, Disability, Sarcopenia, and Reduced Survival. *The Journals of Gerontology: Series A, 73(7),* 925–930. https://doi.org/10.1093/gerona/glx126

Walker, H. K., Hall, W. D., & Hurst, J. W. (1990). Clinical Methods: The History, Physical, and Laboratory Examinations. (1990). *Annals of Internal Medicine, 113(7),* 563. https://doi.org/10.7326/0003-4819-113-7-563_2

Webster, N. S., & Reusch, T. B. H. (2017). Microbial contributions to the persistence of coral reefs. *The ISME Journal, 11(10),* 2167–2174. https://doi.org/10.1038/ismej.2017.66

Willmer, P., Stone, G., & Johnston, I. (2009). Environmental physiology of animals. *John Wiley & Sons.*

Yousef, M. I., El Hendy, H. A., El-Demerdash, F. M., & Elagamy, E. I. (2002). Dietary zinc deficiency induced-changes in the activity of enzymes and the levels of free radicals, lipids and protein electrophoretic behavior in growing rats. *Toxicology, 175(1–3),* 223–234. https://doi.org/10.1016/s0300-483x(02)00049-5

Zhang, J., Zhang, N., Liu, S., Du, S., & Ma, G. (2022). Young Adults with Higher Salt Intake Have Inferior Hydration Status: A Cross-Sectional Study. *Nutrients, 14(2),* 287. https://doi.org/10.3390/nu14020287

Zhao, M.-M., Krebs, J., Cao, X., Cui, J., Chen, D.-N., Li, Y., Hua, L., Mann, J., & Yang,

J.-K. (2019). Helicobacter pylori infection as a risk factor for serum bilirubin change

and less favourable lipid profiles: a hospital-based health examination survey. BMC

Infectious Diseases, 19(1). https://doi.org/10.1186/s12879-019-3787-8

Zomer, R. J., Xu, J., & Trabucco, A. (2022). Version 3 of the Global Aridity

Index and Potential Evapotranspiration Database. Scientific Data, 9(1).

https://doi.org/10.1038/s41597-022-01493-1

1031