Title: *Negative impact of mild arid conditions in natural rodent populations*
revealed using markers of physiological condition in natura

Authors : Hamilcar Keilani[1*], Nico Avenant[2], Pierre Caminade[1], Neville Pillay[3],

Guila Ganem[1,3]

**1.	ISEM, Univ Montpellier, CNRS, IRD, Montpellier, Occitanie, France**

**2.	Department of Mammalogy, National Museum and Centre for**

Environmental Management, University of the Free State, Bloemfontein, Free

State, South Africa

**3.	School of Animal, Plant and Environmental Science, University of the**

Witwatersrand, Wits, Gauteng, South Africa

* Corresponding author; e-mail : hamilcar.keilani@umontpellier.fr,

hamilcarkeilani98@gmail.com

Open research statement

Data will be available from the Dryad repository (TBA)

Keywords

Aridity, blood metabolites, ecophysiology, liver, local adaptation, nutrition,

Rhabdomys dilectus dilectus, Rhabdomys bechuanae

Abstract

1.   Understanding how organisms respond to seasonal variations in their

environment can be a window to their potential adaptability, a classical

problem in evolutionary ecology. In the context of climate change, inducing

increased aridity and disruption of seasonality, it is crucial to study the extent

and limits of species responsiveness.

2.   Here, the physiological response to food and water shortage during

seasonally dry conditions were investigated. We studied populations of two

rodent species of the genus *Rhabdomys*, one arid and one mesic, in a semi-

arid zone where their range overlap in South Africa. We measured blood

concentrations of markers of kidney and liver function, as well as body

condition, at the onset and the end of the dry season.

3.   We found similar shifts in blood metabolite levels in the semi-arid

populations of the two species, indicating malnutrition consistent with the

observed degradation of habitat quality between the start and the end of the

dry season. Furthermore, regardless of the period, differences between the two

species in blood metabolite concentrations (ex: amylase, sodium, alkaline

phosphatase) were observed, suggesting contrasting diets and water

conservation abilities.

4.   Overall, we show that, as seasonal dry conditions worsen, organisms are

increasingly affected by reduced food availability, and local adaptation to arid

conditions may provide the arid species with an advantage to cope with semi-

arid conditions. Our study suggests that even mild arid conditions could have

a negative impact and questions resilience of animals to harsher arid

conditions.

Introduction

Organisms are constantly exposed to a wide range of environmental variations,

including short-term changes during their lifetime and longer-term changes across

generations (Lopez-Maury et al., 2008). In the coming decades, extreme and more

frequent droughts are predicted by most projections, resulting from changes in

precipitation patterns, increased temperature, and evaporation (Naumann et al.,

2018). These changes will induce great challenges to organisms. Under such

circumstances, divergent selection can lead to the evolution of local adaptations

where, in a given environment, resident phenotypes outperform nonresident ones

(Kawecki & Ebert, 2004).

As a result of climate change, organisms can be faced with warmer and drier

environmental conditions (Parmesan et al., 2000), likely to impact their capacity to

maintain homeostasis (i.e. the state of steady internal conditions allowing optimal

body functioning) (Davies, 2016; Fuller et al., 2016). In dry environments, food and

water can be scarce, generating strong selection on physiological attributes

maximizing energy and water availability for body function. Habitat generalists and

specialists' taxa may respond differently to environmental changes. Desert specialists

exhibit specific physiological and/or behavioral traits, resulting from genetic

adaptations to the environment, allowing them to continuously endure or evade

periods of resource restriction (Rocha et al., 2021). In contrast, generalist species

may seek to escape the effects of aridity through temporary avoidance (Abraham et

al., 2019) or plastic adjustments during unfavorable seasons (Kobbe et al., 2011).

Compared to mesic species, arid species have lower basal metabolic rates and

evaporative water loss (Muñoz-Garcia et al., 2022), adopt a more flexible diet

(Tshikae et al., 2013), and cope better with prolonged droughts (Boyers et al., 2021).

Local ecological adaptation can be described in several different and complementary

ways, such as through its genetic basis, transcriptomic, behavioral, or physiological

aspects. Adaptive evolution *sensu stricto* occurs when the genetic constitution of a

population changes because of natural selection (Merilä & Hendry, 2013). In contrast,

local adaptation *sensu lato*, also considers concepts such as phenotypic plasticity

(the ability of a genotype to produce distinct phenotypes when exposed to different

environments throughout its ontogeny; Pigliucci, 2005). It is usually assessed using

comparative population or species fitness estimates under novel environmental

conditions (Webster & Reusch, 2017). The response to local environmental

conditions is a major factor in the generation and maintenance of biodiversity,

impacting population dynamics, biogeographical ranges, and species interactions

(Blanquart et al., 2013; Post et al., 2009). Specifically, physiological limits can drive

and be driven by evolution, shape species distributions and niches, and define

species response capacities to future climate change, directly impacting risks of

extinction (Somero, 2012). For example, latitudinal niches are associated with

adaptive variation in thermal limits in marine invertebrates (Somero, 2010).

Studies of different physiological systems, such as the cardiovascular (Seebacher et

al., 2005, Berkel & Cacan, 2022) or digestive systems (Naya et al., 2011), in

response to different abiotic conditions, can improve our understanding of the causal

mechanisms of responses of organisms to climate variation (Seebacher & Franklin,

2012). Biotic factors, such as sex, age, diet, strain, or breeding status also influence

individual responses (McClure, 1999). With a focus on influence of air temperature,

water availability and energy resources on an organism's physiological state,

research in physiological ecology has addressed how variation in physiology evolves

and is maintained in relation to environmental conditions (Feder & Block, 1991).

Analysis of blood metabolites has successfully been used to investigate kidney and

liver physiology of wild populations (e.g. Al-Eissa et al., 2012). The kidney and the

liver both contribute to homeostasis in periods of food and water scarcity; indeed, the

kidney plays a crucial role in maintaining osmotic balance and the liver is involved in

dietary metabolism. Water loss can also mechanically induce high concentrations of

some blood markers and impact organism's capacity to evacuate products of

metabolic waste, such as urea, from the blood (Ostrowski et al., 2006). Hence, dry

conditions can put a strain on these organs and be detrimental to survival and

reproduction.

In the Succulent Karoo of South Africa, where the average daily precipitation during

the dry season is 0.26 mm *versus* 0.78 mm during the wet season, the probability of

survival of adult *R. pumilio,* another species of the same genus, were related to their

physiological response at the start of the dry season (Schoepf et al., 2017a). Schoepf

and collaborators found higher serum concentrations of albumin, glucose, potassium,

and lower concentrations of globulin, in animals that survived compared to those that

did not survive the dry season. At the peak of the dry season, individuals had lower

concentrations of glucose and phosphorus and higher concentrations in globulin and

urea nitrogen if they survived the dry season. These results strongly suggest that

such markers could be good indicators of individual fitness.

The two study species, *R.bechuanae* and *R.dilectus dilectus*, have different

environmental niches (du Toit et al. 2012, Meynard et al. 2012). Throughout most of

its distribution, *R.d.dilectus* occurs in mesic habitats with vegetation cover and nests

in dense grass, while *R.bechuanae* is found in semi-arid and arid habitats, thriving

predominantly in sparsely vegetated areas and nesting in bushes (Dufour et al.,

2015; Dufour et al., 2019). Moreover, differences in morphology and behavior

suggest adaptation to dry conditions in *R.bechuanae* (Ganem et al., 2020; Dufour et

al., 2019).

In our study, potential impact of increased aridity was addressed by investigating the

physiological consequences of seasonal variation in dry conditions, in two species of

the diurnal African striped mouse genus *Rhabdomys*. Here we address both

physiological responses to seasonal dry conditions and divergence in populations of

the two species in the same bioclimatic region.

Indeed, at the edge of their distributions in central South Africa, the two species

inhabit a semi-arid region within which they occur either as parapatric or sympatric

populations. This semi-arid zone, the expansion and contraction of which depends

on land use and precipitation (Lian et al., 2021), has experienced a recent trend

towards desiccation (Jury, 2021). Such drying conditions could generate additional

selective pressures to those already experienced by the semi-arid populations of the

two species. Short-term climatic patterns can also impact resource availability in this

region. Indeed, during La Niña years (like during this study), as part of the El Niño

Southern Oscillation phenomenon, there is a general association between regional

wetness and sea surface temperatures in the neighbouring Atlantic and Indian

Oceans, leading to wetter episodes throughout Southern Africa (Nicholson & Selato,

2000).

While many studies have harnessed physiology, genomics, and transcriptomics to

highlight potential adaptive interspecific or interpopulational contrasts, studies under

common natural environmental conditions are lacking (Rocha et al., 2021). These

approaches provide a mean to disentangle the roles of interpopulational or

interspecific variation from extrinsic environmental factors in shaping phenotypic

variation, taking advantage of the natural conditions experienced in the field,

including their complexity, with little human involvement. This study took place in a

natural semi-arid environment and asked how seasonal variation in dry conditions

influenced the physiological responses of *R.bechuanae* and *R.d.dilectus*. We

compared body condition, blood concentrations of markers of kidney and liver

functions and habitat characteristics of parapatric populations of the two species at

the start *versus* the end of the dry season. First, we expected that the habitat

available for the mice would be drier at the end compared to the onset of the dry

season, inducing a reduction of food and water, impairing liver and kidney function,

and impacting body condition. Second, assuming local adaptive plasticity, we

hypothesized that both species would be able to adjust their physiological responses

to seasonal changes in dry conditions. Third, we expected that *R.bechuanae*, having

evolved in arid environments, would perform better in semi-arid conditions and be

closer to the expected local optimal response to increased dry conditions than

R.d.dilectus (having evolved in mesic areas), as found for *Mus musculus* (Bittner et

al., 2021). Alternatively, if both species evolved specific adaptations to the semi-arid

environment, we expected no species differences (**Table 1**).

Materials and Methods

__Ethics statement__

Permits to sample and handle animals in the field were obtained from the DESTEA of

the Free State, the North West DEDECT and Northern Cape DENC (respectively,

n°202110000007994, NW 38956/04/2022, FAUNA 0389/2022). Animal handling was

performed under ethical clearance from the Languedoc-Roussillon ethical committee

for animal experimentation (n°2022041512256467 v5).

__Choice of sampling periods and sites__

We selected six study sites based on their geographical position and their aridity

index (AI). The AI calculation was based on Thornthwaite method (Thornthwaite,

1948; see **Supplementary Material** for full formula) using rainfall and temperature

data collected from 2010-2021 (South African Weather Service).

We selected parapatric populations of the two species in a semi-arid region of South

Africa ($0.2 < AI < 0.5$). In this region, a hot and wet season occurs from roughly

December to February, and a cool dry season from May to September. We sampled

six sites at the onset (May 2022) of the dry season: Barberspan Bird Sanctuary,

Benfontein Nature Reserve, Bloemhof Dam Nature Reserve, Gariep Dam Nature

Reserve, Kalkfontein Dam Nature Reserve and Wolwespruit Nature Reserve

(**Figures 1&3, Table 2**). We then resampled four of these sites at the end of the dry

season (September 2022). Two sites (Barberspan and Benfontein) were excluded

from the September sampling due to extreme weather conditions.

__Capture and field protocol__

We used small mammal PVC traps (LxHxW= 29.6x7.5x7.5 cm), baited with a mixture

of oats, peanut butter, salt and sunflower seeds, and added a piece of cotton wool.

They were placed approximately every 15 m along roughly 150 to 300 m transects.

Number and length of transects varied with habitat and terrain conditions. Overall,

trapping effort was 9688 trap nights (details in **Supplementary Tables 1&2**). Since

Rhabdomys is mostly diurnal, the traps were checked twice a day (8 AM and 3 PM).

On the field, all trapped striped mice were measured (body length and mass), their

sex and breeding status (breeding or non-breeding) assessed based on external

morphological features (individuals considered as breeders presented either: signs of

lactation, a perforated vagina, presence of a vaginal plug, or scrotal testis), and a 0.5

mm piece of tail was collected and kept in 98 % ethanol for species identification. All

striped mice were marked with a unique ear-tag before their release at the trapping

location unless they were kept for the physiological study (i.e. kept in their trap with

cotton wool and food).

Out of 903 trapped small mammals, 694 were *Rhabdomys*. For the physiological

study, we selected only adults, avoiding related individuals as much as possible, by

selecting mice trapped at least 100 m apart, except for breeding pairs (a male and a

female) that could be trapped in the same nest. This distance was based on data

obtained for *R. pumilio* indicating minimal relatedness (R=0 between males and

R=0.06 between females) at this distance (Solmsen et al. 2012). In each site,

trapping lasted 3 to 6 days.

Habitat

Vegetation composition

Earlier studies have shown inter-species differences in vegetation cover and

structure requirements (Dufour et al., 2015). Since habitat characteristics can vary

locally, we characterized the vegetation structure (i.e. grass *versus* woody

vegetation) at a microhabitat scale (around the trap), and at a mouse home-range

scale (Dufour et al., 2015). Around traps in which a *Rhabdomys* was trapped, the

percentage of the surface composed of dry grass, green grass, dry bushes, green

bushes, succulent plants, holes and uncovered surfaces was assessed within 2mx2m

(4m²) and the general vegetation within 10x10m (100m²) quadrats centered on the

trap position. For each 4m² quadrat we recorded in detail the percentage and type of

cover at the ground level within each of four 1x1m subunits (using a metal frame); the

results obtained for the four subunits were then averaged. For the 100m² quadrats

the assessment was made following a visual inspection at eye-level height; the

different cover percentages were estimated by the same observer. All together we

characterized 236 quadrats of each type (100m² and 4m²).

Normalised Differential Vegetation Index

NDVI, the satellite imagery-based index informative of ground vegetation greenness,

was retrieved from the Copernicus Open Access Data Hub (Copernicus Sentinel-2

data, 2023, calculated from 10-m resolution bands). While the two above-mentioned

quadrats were aimed at characterizing the structure/cover of the habitat, NDVI was

used as an index of habitat quality. To characterize each site and sampling session,

we used NDVI data available for the closest day to the beginning of a sampling

session, which was expected to represent the conditions experienced by the mice at

the time of capture**.**

Blood extraction and assessment of blood metabolites concentrations

In this study, a total of 273 adult mice were euthanized by means of cervical

dislocation on their day of capture, and a blood volume of around 100 µL was

collected in lithium-heparinized tubes by cardiac puncture. The breeding status was

confirmed for all individuals during dissection. The whole liver, the left kidney, and the

skull of every individual were also collected for a complementary study. Because the

two species could not be distinguished visually, the spleen was also harvested from

each mouse and subsequently analyzed for post-hoc species identification using

Cytochrome Oxidase I genotyping (as described in Ganem et al. 2020).

Levels of albumin, alkaline phosphatase, alanine aminotransferase, amylase, total

bilirubin, blood urea nitrogen, total protein, globulin, glucose, calcium, phosphorus,

potassium, and sodium were successfully measured in the whole blood of 257

individuals immediately after blood collection (Abaxis VETSCAN V2 technology with

the Comprehensive diagnostic profile cartridge), following the manufacturer's

instructions. The coefficient of variation (CV%) of measured levels of these markers

was determined from data obtained by analyzing a given blood sample twice.

Samples from six adult striped mice (5 *R.bechuanae*, 1 *R.d.dilectus*) were used for

these calculations. For these six samples, the calculated CV equaled 5.6 ± 1.9 %,

which is consistent with the intra-test CV% provided in the manufacturer's instruction

manual for other mammal species (**Supplementary Table 3**).

Data preparation and analysis

All statistical tests were carried out using the R software (version 4.2.1).

Habitat

<u>Vegetation composition</u>

Because the distribution of the variables describing the cover and vegetation type

did not meet the conditions necessary for further multivariate model based analyses,

we performed a Principal Component Analysis (package FactoMineR version 2.9)

including the 7 variables measured within the $4m^2$ quadrats and within the $100m^2$

quadrats aiming to transform and, as far as possible, reduce the number of variables.

In each case, over 80 % of the variance was explained by the first five Principal

Components (**Supplementary Tables 4 & 5**). The distribution of the coordinates of

all traps on the first 5 PCs complied to the statistical analysis constraints and were

used as response variables of a *PERMANOVA* (package *vegan*, version 2.6-4).

PERMANOVA is a multivariate statistical inference tool using permutational

algorithms (Anderson, 2001). This equivalent to *MANOVA* operates in a distribution-

free setting and is robust to non-normality of residuals as well as dispersion

heterogeneity, even in slightly unbalanced designs (Anderson, 2013). We tested

differences between sites nested within sessions (start and end of dry season), with

10 000 permutations, to assess whether the sites had similar enough habitat

characteristics to be considered as suitable replicates in other analyses.

<u>NDVI</u>

NDVI was averaged over a 60 m radius circular buffer centered at each trap and then

compared across sessions and sites using a GLM Analysis of Variance, with the site

nested within session. To meet statistical test prerequisites, the response variable

was log-transformed; the analysis was followed by Tukey multiple comparisons of

means post-hoc tests.

Age classes

Among trapped *Rhabdomys*, only adults were selected for this study based on their

size, general appearance (fur dullness, presence of scars), and/or breeding status.

Body length of trapped individuals varied between 5 and 12 cm. The body length of

adult individuals involved in the physiological study varied between 7.7 and 12 cm.

To distinguish between older and younger adults, mass was plotted against body

length (**Supplementary figure 6**) showing a curve defined by an allometric growth

equation with a steeper slope at the highest and lowest body length values. Since

bone growth can significantly alter the concentration of some physiological markers,

and growth rate varies with age/body length, we considered 4 length/age classes

among the adults (Class A: [7.7:9 cm[; Class B : [9:10 cm[; Class C : [10:11 cm[;

Class D [11 cm:]).

Body condition

We calculated a scaled mass index of body condition using the method described in

Peig and Green (2009; see **Supplementary Material** for full formula), to assess the

state of each individual's fat reserves (Schulte-Hostedde et al., 2005). To test

whether breeding status, age, habitat quality, increased dryness, and/or interspecific

differences influence body condition of studied individuals, linear mixed effect models

were computed with body condition as a response variable, site as a random factor

and session, species, sex, session*species*sex (including all two-way interactions),

breeding status (nested within sex), age class and NDVI as explanatory variables.

Assumptions of normality and homoscedasticity of residuals were checked using

diagnostic plots (*graphics* package v.3.6.2.). A Tukey multiple comparisons of means

post-hoc test was applied to determine which inter-level differences were driving the

significance of each factorial effect having more than two modalities.

Physiology

We first identified twelve outliers that we removed from further analyses. These

outliers were animals from which we collected smaller volume of blood samples than

recommended for the Vetscan or were heavily hemolyzed and flagged as such in the

Vetscan analysis output. In addition, as total protein values were used by the

Vetscan analyzer to infer the Globulin (GLOB) fraction of blood protein (with

GLOB=TP-ALB), we considered this marker as not informative and removed this

variable from our analyses. Finally, for 199 out of 257 samples, blood concentrations

of creatinine (CRE) were flagged as being below the instrument's sensitivity

threshold; we hence excluded CRE from subsequent analyses, as this imprecision

would have hindered analytical performance.

Due to covariance in variables such as blood physiological markers, multivariate

statistical techniques provide a unique insight into the main patterns and effects

driving variation in the data. As with habitat, multivariate normality was violated in our

physiological dataset, so permutational multivariate analysis of variance

(*PERMANOVA*) was performed. 10 000 permutations were used, with 236 blood

samples, 12 physiological parameters as response variables, site as a random factor,

and session, species, sex, breeding status (nested within sex), body condition, age

class, and NDVI as explanatory variables. A backwards stepwise model selection

was performed to retain the most parsimonious model.. We then sought to identify

which of the 12 physiological markers drove significantly the between-group

differences evidenced with the *PERMANOVA*. Because in the PERMANOVA

package a dedicated function was not available, twelve post-hoc univariate linear

mixed model (LMM) tests were performed (followed by backwards stepwise model

selection), initially using the parameters of the most parsimonious *PERMANOVA*

model. Assumptions of normality and homoscedasticity of residuals were checked

using diagnostic plots. In the case of total bilirubin, a permutation test had to be

performed instead of the *LMM*. Through a backwards stepwise model selection

process. For all tests, the significance level (α) was set at 0.05.

Results

Variation in habitat characteristics between species and sessions

Vegetation composition

We compared the vegetation characteristics surrounding each successful trap with a

PERMANOVA, using the first 5 Principal Component coordinates (**Supplementary**

Tables 4 & 5) of each quadrat as response variable, and site and session as

explanatory variables. We detected significant differences in overall habitat

characteristics between sessions (4m² and 100m²: p<0.001), and sites nested within

sessions (4m² and 100m²: p<0.001), indicating clear site differences and seasonal

effects on habitat structure (**Supplementary Figures 4 & 5**). Although a

permutational dispersion test indicated that the latter patterns could be attributable at

least in part to differences in dispersion between sessions or sites (see

Supplementary Material : Tables 6&7; Figures 1,2,3&4), some general trends

could be inferred from the data (**Supplementary Tables 7&8**). Indeed, quadrats from

R.d.dilectus sites (Barberspan, Bloemhof and Wolwespruit) tended to feature higher

proportions of grass (mean 4m²: 83.99 ± 1.12 %, mean 100m² : 71.42 ± 1.40 %) than

R.bechuanae sites (mean 4m²: 54.30 ± 2.57 %, mean 100m²: 53.12 ± 1.84 %).

Conversely, *R.bechuanae* sites (Benfontein, Gariep and Kalkfontein) tended to

feature higher proportions of bushes (mean 4m²: 31.07 ± 2.37 %, mean 100m²: :

27.35 ± 1.42%) than *R.d.dilectus* sites (mean 4m²: 7.33 ± 1.00 %, mean 100m² :

10.49 ± 1.03 %). Furthermore, differences between sessions could be observed;

quadrats surveyed at the end of the dry season tended to feature more dry grass

(mean 4m²: 60.82 ± 2.33 %, mean 100m²: 51.99 ± 1.87 %) than at its onset (mean

4m² : 50.75 ± 2.04 %, mean 100m²: 40.85 ± 1.39 %).

NDVI

The log-transformed NDVI of sites sampled at the two sessions (2 sites per species)

was also analyzed between sites and sessions using *ANOVA*. Site, session and the

interaction between these two variables (all $p<0.001$) contributed significantly to

explain the NDVI variation (**Table 3**). A post-hoc Tukey multiple comparisons of

means test showed that all sites had a significantly lower NDVI in September

compared to May (**Figure 4**), indicating a poorer state of the vegetation at the end of

the dry season. Furthermore, while these study locations were chosen in a region

with similar semi-arid bioclimatic features, Gariep Dam and Kalkfontein (where

R.bechuanae was targeted) seemed to stand out as having a lower NDVI than all

other sites sampled in both May and in September, with Wolwespruit and Bloemhof

(where *R.d.dilectus* was targeted) having the highest NDVI (**Figure 4**). Therefore, to

take into account the impact of variations in NDVI between the two species trapping

locations we included site as a random factor in analyses addressing body and

physiological conditions.

Variation of Body condition

A linear mixed model assessed the influence of session, species, breeding status,

age class and NDVI on body condition, including site as a random factor. The most

parsimonious model included sex ($p<0.001$), age class ($p<0.001$), breeding status

nested within sex ($p<0.01$) and the interaction between breeding status and session

($p<0.05$); session was not significant by itself $p=0.965$, while the effect of NDVI

verged on significance ($p=0.067$), and the interaction between session and sex

(p=0.683, (**Table 4**). Body condition was lower at the end of the dry season than at its

onset for non-breeding females compared to breeding females (post-hoc pairwise t-

test p<0.05, **Supplementary Table 10**).

Variation in physiological responses

A *PERMANOVA* was performed using the distance matrix of 12 physiological

markers as response variables and session, species, sex, breeding status, body

condition, age class, NDVI and site (random factor) as explanatory variables

(**Supplementary table 11**).

The most parsimonious model **(Table 5)** indicated an overall difference in

physiological parameter concentrations between *R.bechuanae* and *R.d.dilectus*

(p=0.002), as well as an influence of body condition (p=0.019), breeding status

(nested within sex) (p=0.028), that differed between the two species (interaction

effect, p=0.022), and the interaction between sex and species (p=0.020). Session

(p=0.065), and sex (p=0.084) verged on overall significance and were also included

in the model, indicating that physiological markers overall tend to shift between the

start and end of the dry season, and to vary between sexes. Importantly, both

species showed the same pattern of variation between sessions (no significant

species*session interaction effect, p=0.498, **Supplementary Table 11**).

Six physiological markers showed significant differences between the start and the

end of the dry season (**Figure 5**). Blood concentrations of markers of nutrition and

liver function (albumin, potassium, alanine aminotransferase, alkaline phosphatase,

and total bilirubin) were lower at the end compared to the onset of the dry season

(**Table 4**). Glucose also showed that trend, but not significantly (p=0.0577). In

contrast, globulin levels varied in the opposite direction (higher at the end compared

to the onset of the dry season).

The levels of four markers were significantly lower in *R.bechuanae* than in

R.d.dilectus : total bilirubin, alkaline phosphatase (ALP), sodium and potassium

(**Table 6, Figure 6**). However, amylase showed the opposite pattern between the

species.

Regardless of species, ~~ALP levels were negatively correlated with body condition,~~

~~while the levels of phosphorus exhibited the opposite pattern. Finally,~~ the levels of

glucose and ALP varied significantly with sex, tending to be lower in females than in

males (**Supplementary figure 7**).

Discussion

We asked how seasonal variation of dry conditions influenced the physiological

response of semi-arid parapatric populations of two closely related *Rhabdomys*

species that evolved under distinct environmental conditions (du Toit et al., 2012).

Consistent with our predictions, we found strong evidence for physiological

divergence between the species. We also observed a tendency for seasonal

variation of the physiological response in both species related to energetic resource

depletion at the end of the dry season, supported by the shifts in the levels of 6 blood

parameters between the onset and the end of the dry season.

Our study took place in a relatively wet year (La Niña). Vegetation growth and

senescence usually follow seasonal rainfall patterns; above-average rainfall widens

the temporal window for vegetation growth, improving habitat conditions at the start

of the typical dry season (Anyamba et al., 2002). Despite this, and consistent with our

predictions, milder dry conditions did alter habitat quality of our study sites (as

attested by shifts in NDVI and habitat structure).

Seasonal variation of physiological condition

Despite our study year being relatively wet, mild dry conditions impacted blood

marker concentrations in *R.bechuanae* and *R.d.dilectus*. These changes involved 6

out of 12 tested physiological markers, and included markers expected to vary in

periods of dietary restriction, such as albumin. Contrary to our prediction, we did not

find a significant interactive effect between session and species, suggesting that the

semi-arid populations of the two species were similarly affected by seasonal dry

conditions. However, a follow up study during an El Niño year may be necessary

before one can conclude.

Water and nutritional stress are expected to impact the body condition of individuals

(Fuller et al., 2021). ~~Interestingly, the body condition of sampled individuals was~~

~~higher at the end of the dry season.~~ As expected, body condition varied with breeding

status, with differences between breeding and non-breeding females heightened at

the end of the dry season. Additionally, several physiological markers varied

significantly between males and females, indicating their association with

physiological processes involved in reproduction. Reproduction in mammals has

direct physiological costs, associated with increased energy, protein and calcium

demands and indirectly through compensatory effects such as reduction in

thermogenesis, immune function and physical activity (Speakman, 2008; Rintoul &

Bringham, 2014; Stawski & Rojas, 2016; Schoepf et al., 2017b). It follows that sex-

specific differences in reproductive investments are associated with contrasts in body

weight, fat distribution and energy metabolism (Chen et al., 2012). In support, our

results indicate lower levels of blood glucose, and a lower body condition in females

compared with males. Moreover, albumin (ALB) and alkaline phosphatase (ALP)

were significantly elevated in males compared to females, which could also be

attributable to the pleiotropic effects of sex hormones (Havill et al., 2004; Sullivan et

al., 2007).

The two species responded similarly to increased dryness while maintaining body

condition, a proxy of fat reserves. Similarly, *R.pumilio* individuals tended to maintain

their body condition stable while reducing their energy expenditure and physical

activity during periods of limited food availability (Rimbach, Blanc et al., 2018;

Rimbach, Jäger et al., 2018). The semi-arid populations of *R.bechuanae* and

R.d.dilectus could be displaying a similar behavioral strategy.

~~Variation in ALP levels is linked to variation in soft tissue and/or bone growth (Walker~~

~~et al., 1990). As expected, ALP was also negatively correlated with body condition.~~

~~Individuals with the lowest body condition had lower phosphorus levels. Generally,~~

~~high phosphorus levels can be observed in dehydrated individuals in which~~

~~osmolality is high; indeed, low phosphorus levels related to survival in *R.pumilio* at~~

~~the peak of the dry season, where water balance was hardest to maintain (Schoepf et~~

~~al., 2017). However, blood levels of phosphorus also depend on its consumption and~~

~~hence availability in the food (Berndt & Kumar, 2009). Phosphorus reserve depletion~~

~~in smaller-sized adults might therefore be consistent with the role of phosphorus~~

~~intake as a promoter of bone and soft tissue growth (Shapiro & Heaney, 2003).~~

~~Furthermore, higher needs for phosphorus in smaller adults might be hindered by~~

~~reduced access to food. Indeed, foraging activity tends to vary between individuals~~

~~according to size and breeding status, possibly resulting in larger breeding individuals~~

~~gaining access to higher quality food (Ebensperger & Hurtado, 2005).~~

Blood protein levels are generally good indicators of physiological condition (Tothova

et al., 2016). They comprise two distinct and major components: albumin (ALB) and

globulins (GLOB). In our study, ALB was significantly lower at the end than at the

start of the dry season. ALB is synthesized in the liver and involved in the transport of

bilirubin and several hormones. Blood ALB levels are less prevalent in malnourished

individuals (Walker et al., 1990), supporting the hypothesis of reduction of quality

and/or availability of food at the end of the dry season. The Nubian ibex *Capra*

nubiana, a desert mammal, also displays low ALB during the dry season (Al-Eissa et

al., 2012), which could reflect poor nutritional conditions, kidney dysfunction leading

to poor water balance and high osmolality, or liver dysfunction causing low ALB

synthesis. We found that GLOB was significantly higher at the end than at the start of

the dry season. Higher GLOB levels were also reported for *R.pumilio* individuals that

did not survive the dry season (Schoepf et al., 2017). GLOB is a family of molecules

including immunity proteins and blood protein carriers (including transferrin). An

increase in GLOB levels usually indicates elevation of immunoglobulins and hence

could suggest heightened immune activity. Greater antibody production occurs in

response to a higher prevalence of respiratory infections during the dry season, e.g.

in the Nubian goat (Abdelatif et al., 2009), and/or reduced protein nutritional status in

young birds (Lochmiller et al., 1993). An increase in blood GLOB concentrations may

also reflect high levels of transferrin (Walker et al., 1990), a major GLOB that may

accumulate in the blood in cases of severe iron deficiency (Kasvosve & Delanghe,

2002). Hence, elevated blood GLOB levels at the end of the dry season may indicate

either or both depletion in iron resource (transferrin) and higher vulnerability to

infection (immunoglobulin), both resulting from or being aggravated by reduced

access to food. Lower levels of alkaline phosphatase (ALP), a liver enzyme, at the

end of the dry season in *Rhabdomys*, is also evidence of malnutrition and deficiency

in essential nutrients (proteins, magnesium, zinc) (Yousef et al., 2002; Saraç &

Saygili, 2007; Ray et al., 2017). Alanine aminotransferase (ALT), another marker of

liver function, also occurred at significantly lower concentrations at the end compared

to the start of the dry season, indicating a reduction in liver activity (Walker et al.,

1990) or malnutrition (Le Couteur et al., 2010). Indeed, as environmental resources

become scarce, essential nutrients, such as pyridoxine (or vitamin B6), that constitute

some ALT coenzymes, may be less available, leading to a reduction of ALT blood

levels (Vespasiani-Gentilucci et al., 2018).

Blood glucose, another major physiological marker of nutrition state, was also slightly

diminished at the end of the dry season, albeit not significantly. Fasting or low-energy

feeding results in lower blood glucose levels (Jensen et al., 2013).

Acute starvation also typically causes bilirubin (TBIL) to accumulate in the blood, as

the enzymatic process for conversion of heme into TBIL is stimulated (Thaler et al.,

1972). Yet, TBIL was lower at the end compared to the onset of the dry season,

rejecting the starvation hypothesis. Lower TBIL concentrations at the end of the dry

season in *Rhabdomys* could stem from low hemoglobin counts because the body

produces fewer red blood cells than usual; low hemoglobin can be induced by

several factors, among which dietary factors such as iron deficiencies (Clark, 2008).

Iron deficient anemia is also supported by higher GLOB concentrations observed at

the end of the dry season. Infection can likewise be a cause of both decreased TBIL

levels (Zhao et al., 2019) and higher GLOB concentrations at the end of the dry

season. Finally, malnutrition and/or lack of potassium intake are among the main

potential causes of the reduction of its levels in the blood (Their, 1986), which could

explain why they were the lowest at the end of the dry season, when the available

resources were lowest, in our study.

Overall, our results indicate that seasonal variation of some physiological markers

could reflect reduced primary productivity throughout the dry season, which is

coherent with our observation of habitat degradation (lower NDVI) at the end of the

dry season. Blood concentrations of various markers seem to indicate difficulties in

maintaining nutritional functions and the necessity of a higher immune output, caused

or exacerbated by food restriction. Indeed, individuals undergoing malnutrition during

dry conditions are expected to show low blood glucose concentrations, and

deficiencies in essential nutrients, leading to pathologies such as anemia (Gordon et

al., 1988). Indeed, animals foraging on soils potentially poor in nutrients such as iron,

zinc, and magnesium are more prone to malnutrition (Graham, 1991; Gupta et al.,

2008). Higher GLOB, lower TBIL and lower ALP blood concentrations in our study

might relate to such deficiencies.

We did not observe seasonal variation of sodium or blood urea nitrogen, blood

markers of kidney function or osmoregulation. Indeed, reduced environmental

moisture at the end of the dry season could have caused an overall increase in blood

osmolality, increasing blood concentrations of most biomarkers. Still, most markers

exhibited a significant reduction in blood concentrations. This suggests that,

compared to food availability, seasonal patterns in water availability may impose a

milder physiological cost to the striped mice throughout the dry season in this semi-

arid environment during a La Niña (wet) episode.

Interspecific differences

R.bechuanae is thought to have evolved in drier environments than *R.d.dilectus*.

Moreover, in the semi-arid region where they co-occur, we showed that *R.bechuanae*

populations occupy slightly drier habitats than *R.d.dilectus*. While body condition did

not differ significantly between species, some interspecific contrasts support our

hypothesis that *R.bechuanae* may have developed a better physiological capacity to

cope with dry conditions than *R.d.dilectus*.

Among the markers exhibiting a significant difference between *R.bechuanae* and

R.d.dilectus, all but amylase showed higher blood concentrations in *R.d.dilectus*.

Amylase is positively correlated with digestive activity and reflects differences in

feeding status or diet composition between the species (Hidalgo et al., 1999). Given

the role of alpha-amylase in digestion, lower blood amylase levels in *R.d.dilectus*

than *R.bechuanae* may indicate lower digestive activity due to reduced access to

food, or lower starch content in the diet. Pajic et al. (2019) found a correlation

between dietary starch content and the number of amylase gene copies in several

mammalian genomes, even in different habitats and with different diets. Since

amylase activity and amylase gene expression are directly correlated with amylase

gene copy number (Arendt et al., 2014), further studies could elucidate the potential

genetic basis of the different levels of amylase in our two study species.

Sodium (NA) and potassium (K) concentrations, which depend on both dietary and

water intake, were higher in *R.d.dilectus* than in *R.bechuanae*. These differences,

coupled with higher ALP and TBIL concentrations in *R.d.dilectus* compared to

R.bechuanae, could suggest a better nutrition state for *R.d.dilectus* than

R.bechuanae. However, higher levels of amylase in *R.bechuanae* cast doubt on this

interpretation. Instead, lower levels of blood NA, K, ALP, and TBIL, would be more

parsimoniously attributed to lower overall osmolality, perhaps due to better water

balance, in *R.bechuanae*. A better overall ability of *R.bechuanae* to conserve body

moisture could rely on physiological mechanisms such as more efficient excretion or

osmoregulation, or through feeding on a diet optimizing water intake. We hypothesize

that during the evolutionary history of *R.bechuanae*, selection pressures may have

favored specific strategies allowing an overall more efficient water regulation

compared to *R.d.dilectus*. These results, combined with the reduction in the levels of

K, ALP and TBIL at the end of the dry season suggest both species are exposed to

both water and nutrition stress.

We found relatively little evidence that the two species suffer differentially from lack of

food in their semi-arid environment during a dry season impacted by La Niña

phenomenon. Overall, energy levels, as indicated by blood glucose, and body

condition, did not vary between the species. Instead, it seems that the overall

differences between the two species can be accounted for by lower blood osmolality

in *R.bechuanae*, and on average higher blood osmolality in *R.d.dilectus,* as

evidenced by differences in NA and K concentrations, possibly due to overall poorer

water regulation. High average blood amylase levels in *R.bechuanae*, compared to

R.d.dilectus during the dry season, might indicate an interspecific difference in diet

composition or food intake, rather than protein-energy malnutrition. Little is known

about *R.bechuanae* and *R.d.dilectus* diets; the genus was characterised as a

generalist feeding on seeds and other plant material, and insects (Curtis & Perrin,

1979). Some studies showed variation in diet composition within *Rhabdomys* genus

associated with variation in local resources (Taylor & Green, 1976). The two species

could have adjusted their diet to the respective local availabilities of different food

resources within their home ranges. For example, during the dry season and in the

semi-arid zone, arthropod biomass and diversity is affected by seasonal patterns in

rainfall. Dalerum et al. (2017) found lower arthropod trapping success and diversity

during the dry season, in shrub habitats compared to any other biomes, in Benfontein

Game Reserve, one of our study sites. Thus, compared to seeds, the reliability of

arthropods as a food source during dry conditions could be lower in shrublands,

inhabited by *R.bechuanae* across its range and by both species in parapatry, than in

grasslands inhabited by *R.d.dilectus* in allopatry. Variation in seed composition could

also explain this interspecific difference in amylase concentrations. Indeed,

granivores prefer seeds that have higher protein content and lower secondary

metabolites but may be forced to take a wider range of seeds in dry areas than in

mesic ones (Wilmer et al., 2009). If the hypothesis of a relatively ancient adaptation

of *R.bechuanae* to dry habitats is true, it could be expected that it would also have

adjusted its diet to the specific conditions encountered in dry habitats. Within this

ecosystem, *R.bechuanae* could forage more efficiently compared with *R.d.dilectus*,

which evolved in a more mesic environment; its diet could also be richer in starch

(such as containing more seeds/grains than leaves or fibre). Higher starch intake

could help maintaining water balance, starch being the best substrate for metabolic

water production in a dry environment (Adolph, 1964). This is consistent with the

other interspecific differences revealed in this study, as diet can also significantly

affect osmoregulation (Sabat et al., 2009).

Perspectives and conclusions

Due to climate change, populations of many species experience changes in their habitats affecting their fitness and inducing shifts in their geographical distribution (Parmesan et al., 2006; Fuller et al., 2010). In some species, range shifts are expected to swell edge populations, as marginal habitats grow more favourable. For instance, edge populations of *Wilsonia citrina* hooded warblers have repeatedly expanded their ranges due to climate change (Melles et al., 2011). In other species, the quality of marginal habitats occupied by populations at the range edge may decline because of reduced food availability or nutrient richness, making these populations particularly vulnerable to climate change and inducing range contraction. For example, the koala *Phascolarctos cinereus* experiences declines in population sizes and habitat contraction due to droughts (Seabrook et al., 2011). As such, depending on their phenotypic plasticity, edge populations may hold the key to the survival of a species in a changing environment (Wu & Seebacher, 2022, Usui et al., 2023). *R.d.dilectus* edge populations may experience the same situation, as the example above, in the semi-arid zone, an hypothesis that could be tested addressing these populations dynamics, compared to *R.bechuanae* semi-arid populations.

In this study, we found that two closely related species with different ecological characteristics displayed similar seasonal negative shifts in blood concentrations of markers of nutrition and liver function in spite of a relatively mild seasonal drying of their habitats. We hence expect the physiological impact of harsher dry conditions, to affect the individuals' fitness even more dramatically. The fact that the habitat of *R.bechuanae* studied populations was the driest at both time points, combined with our findings concerning their overall better water balance and nutrition than

R.d.dilectus, point towards *R.bechuanae* coping better in the semi-arid environment

during the dry season. Though adaptation *per se* cannot be inferred, this comparative

study considering populations of the two species in the same biogeographic zone

adds to the growing evidence that these cryptic species of striped mice have

diverged phenotypically. Overall, these findings show the ecological relevance of

physiological approaches in investigating the effect of climatic variation on organisms

and question resilience of core populations of arid species and edge populations of

mesic species in the face of predicted enhanced aridification.

Acknowledgements

We thank R.Nokha who provided valuable and substantial field work contribution to

this project. Constructive and insightful comments were provided by C.Smadja and

C.Schradin helped improve a previous version of this paper. Technical support by

D.Greuet (INM), L.Paradis (Geomatics, ISEM) and GenSeq platform. We are grateful

to the rangers and personnel of Barberspan Bird Sanctuary, Benfontein Nature

Reserve, Bloemhof Dam Nature Reserve, Gariep Dam Nature Reserve, Kalkfontein

Dam Nature Reserve and Wolwespruit Nature Reserve who facilitated our work in

these locations. Funding: CNRS/NRF & OSU-OREME

Author contributions

665 G.G. conceived the idea and designed methodology with input from H.K; G.G., H.K.,

666 P.C., collected the data; H.K. analyzed the data with input from G.G.; H.K. wrote the

667 first draft of the manuscript with input from G.G.; N.P. and N.A. contributed critically

668 to the manuscript, secured funding for the project with G.G. and helped with field

669 logistics. All authors approved the final version of the manuscript.

Conflict of Interest statement

The authors declare they have no conflict of interest.

References

Abaxis, Inc. "VS2 Operator's Manual." (2009) Retrieved from

https://www.zoetis.es/_locale-assets/spc/vetscan-vs2-analizador.pdf

Abdelatif, Abdallah M., Mariam Y. Ibrahim, and Yahia Y. Hassan. "Seasonal variation

in erythrocytic and leukocytic indices and serum proteins of female Nubian goats."

Middle-Eastern Journal of Scientific Research, 4, no.3 (2009): 168-174. Retrieved

from https://idosi.org/mejsr/mejsr4(3)/9.pdf

Abraham, Joel O., Gareth P. Hempson, and A. Carla Staver. "Drought-response

Strategies of Savanna Herbivores." Ecology and Evolution 9, no. 12 (May 22, 2019):

7047–56. https://doi.org/10.1002/ece3.5270.

Adolph, Edward F. "Desert Animals: Physiological Problems of Heat and Water. Knut

Schmidt-Nielsen." Physiological Zoology 37, no. 3 (July 1964): 338–39.

https://doi.org/10.1086/physzool.37.3.30152404.

AL Eissa, M S, Alkahtani Saad, S A Al Farraj, A Alarifi Saud, B Al Dahmash, and Al

Yahya Hamad. "Seasonal Variation Effects on the Composition of Blood in Nubian

Ibex (Capra Nubiana) in Saudi Arabia." African Journal of Biotechnology 11, no. 5

(January 16, 2012): 1283–86. https://doi.org/10.5897/ajb11.2004.

Anyamba, Assaf, Compton J. Tucker, and Robert Mahoney. "From El Niño to La

Niña: Vegetation Response Patterns over East and Southern Africa during the 1997–

2000 Period." Journal of Climate 15, no. 21 (November 1, 2002): 3096–3103.

https://doi.org/10.1175/1520-0442(2002)015<3096:FENOTL>2.0.CO;2.

Arendt, Maja, Tove Fall, Kerstin Lindblad-Toh, and Erik Axelsson. "Amylase Activity

Is Associated with AMY2B Copy Numbers in Dog: Implications for Dog

Domestication, Diet and Diabetes." Animal Genetics 45, no. 5 (June 28, 2014): 716–

22. https://doi.org/10.1111/age.12179.

Berkel, Caglar, and Ercan Cacan. "The Differential Influence of Ambient Temperature

on Heart Rate in Male and Female Wild Pale-Throated Sloths (Bradypus

Tridactylus)." Research Square Platform LLC, August 12, 2022.

http://dx.doi.org/10.21203/rs.3.rs-1908943/v1.

~~Berndt, Theresa, and Rajiv Kumar. "Novel Mechanisms in the Regulation of~~

~~Phosphorus Homeostasis." Physiology 24, no. 1 (February 2009): 17–25.~~

~~https://doi.org/10.1152/physiol.00034.2008.~~

Bittner, Noëlle K. J., Katya L. Mack, and Michael W. Nachman. "Gene Expression

Plasticity and Desert Adaptation in House Mice*." Evolution 75, no. 6 (January 23,

2021): 1477–91. https://doi.org/10.1111/evo.14172.

Blanquart, François, Oliver Kaltz, Scott L. Nuismer, and Sylvain Gandon. "A Practical

Guide to Measuring Local Adaptation." Ecology Letters 16, no. 9 (July 15, 2013):

1195–1205. https://doi.org/10.1111/ele.12150.

Boyers, Melinda, Francesca Parrini, Norman Owen-Smith, Barend F. N. Erasmus,

and Robyn S. Hetem. "Contrasting Capabilities of Two Ungulate Species to Cope

with Extremes of Aridity." Scientific Reports 11, no. 1 (February 18, 2021).

https://doi.org/10.1038/s41598-021-83732-w.

Chen, Xuqi, Rebecca McClusky, Jenny Chen, Simon W. Beaven, Peter Tontonoz,

Arthur P. Arnold, and Karen Reue. "The Number of X Chromosomes Causes Sex

Differences in Adiposity in Mice." PLoS Genetics 8, no. 5 (May 10, 2012): e1002709.

https://doi.org/10.1371/journal.pgen.1002709.

Clark, Susan F. "Iron Deficiency Anemia." Nutrition in Clinical Practice 23, no. 2 (April

2008): 128–41. https://doi.org/10.1177/0884533608314536.

Curtis, B.A., and M.R. Perrin. "Food Preferences of the Vlei Rat (Otomys Irroratus)

and the Four-Striped Mouse (Rhabdomys Pumilio)." South African Journal of Zoology

14, no. 4 (January 1979): 224–29. https://doi.org/10.1080/02541858.1979.11447675.

Dalerum, Fredrik, J. Low de Vries, Christian W.W. Pirk, and Elissa Z. Cameron.

"Spatial and Temporal Dimensions to the Taxonomic Diversity of Arthropods in an

Arid Grassland Savannah." Journal of Arid Environments 144 (September 2017): 21–

30. https://doi.org/10.1016/j.jaridenv.2017.04.002.

Toit, Nina du, Bettine Jansen van Vuuren, Sonja Matthee, and Conrad A. Matthee.

"Biome Specificity of Distinct Genetic Lineages within the Four-Striped Mouse

Rhabdomys Pumilio (Rodentia: Muridae) from Southern Africa with Implications for

Taxonomy." Molecular Phylogenetics and Evolution 65, no. 1 (October 2012): 75–86.

https://doi.org/10.1016/j.ympev.2012.05.036.

Davies, Kelvin J.A. "Adaptive Homeostasis." Molecular Aspects of Medicine 49 (June

2016): 1–7. https://doi.org/10.1016/j.mam.2016.04.007.

Dufour, Claire M. S., Christine Meynard, Johan Watson, Camille Rioux, Simon

Benhamou, Julie Perez, Jurie J. du Plessis, Nico Avenant, Neville Pillay, and Guila

Ganem. "Space Use Variation in Co-Occurring Sister Species: Response to

Environmental Variation or Competition?" PLOS ONE 10, no. 2 (February 18, 2015):

e0117750. https://doi.org/10.1371/journal.pone.0117750.

Dufour, Claire M. S., Neville Pillay, Nico Avenant, Johan Watson, Etienne Loire, and

Guila Ganem. "Habitat Characteristics and Species Interference Influence Space Use

and Nest-site Occupancy: Implications for Social Variation in Two Sister Species."

Oikos 128, no. 4 (November 15, 2018): 503–16. https://doi.org/10.1111/oik.05357.

~~Ebensperger, Luis A., and María J. Hurtado. "On the Relationship between~~

~~Herbaceous Cover and Vigilance Activity of Degus (Octodon Degus)." Ethology 111,~~

~~no. 6 (May 31, 2005): 593–608. https://doi.org/10.1111/j.1439-0310.2005.01084.x.~~

Feder, M. E., and B. A. Block. "On the Future of Animal Physiological Ecology."

Functional Ecology 5, no. 2 (1991): 136. https://doi.org/10.2307/2389251.

Fuller, Andrea, Duncan Mitchell, Shane K. Maloney, and Robyn S. Hetem. "Towards

a Mechanistic Understanding of the Responses of Large Terrestrial Mammals to Heat

and Aridity Associated with Climate Change." Climate Change Responses 3, no. 1

(November 28, 2016). https://doi.org/10.1186/s40665-016-0024-1.

Fuller, Andrea, Duncan Mitchell, Shane K. Maloney, Robyn S. Hetem, Vinicius F. C.

Fonsêca, Leith C. R. Meyer, Tanja M. F. N. van de Ven, and Edward P. Snelling.

"How Dryland Mammals Will Respond to Climate Change: The Effects of Body Size,

Heat Load and a Lack of Food and Water." Journal of Experimental Biology 224, no.

Suppl_1 (February 15, 2021). https://doi.org/10.1242/jeb.238113.

Fuller, Andrea, Terence Dawson, Brian Helmuth, Robyn S. Hetem, Duncan Mitchell,

and Shane K. Maloney. "Physiological Mechanisms in Coping with Climate Change."

Physiological and Biochemical Zoology 83, no. 5 (September 2010): 713–20.

https://doi.org/10.1086/652242.

Ganem, Guila, Claire Marie Soleil Dufour, Nico L. Avenant, Pierre Caminade, Seth J.

Eiseb, Christelle Tougard, and Neville Pillay. "An Update on the Distribution and

Diversification of Rhabdomys Sp. (Muridae, Rodentia)." Journal of Vertebrate Biology

69, no. 2 (July 7, 2020): 1. https://doi.org/10.25225/jvb.20013.

Gordon, G., A. S. Brown, and T. Pulsford. "A Koala (Phascolarctos Cinereus

Goldfuss) Population Crash during Drought and Heatwave Conditions in

South‐western Queensland." Australian Journal of Ecology 13, no. 4 (December

1988): 451–61. https://doi.org/10.1111/j.1442-9993.1988.tb00993.x.

Graham, Thomas W. "Trace Element Deficiencies in Cattle." Veterinary Clinics of

North America: Food Animal Practice 7, no. 1 (March 1991): 153–215.

https://doi.org/10.1016/s0749-0720(15)30816-1.

Gupta, Umesh C., Kening Wu, and Siyuan Liang. "Micronutrients in Soils, Crops, and

Livestock." Earth Science Frontiers 15, no. 5 (September 2008): 110–25.

https://doi.org/10.1016/s1872-5791(09)60003-8.

Havill, L.M, M.C Mahaney, and J Rogers. "Genotype-by-Sex and Environment-by-

Sex Interactions Influence Variation in Serum Levels of Bone-Specific Alkaline

Phosphatase in Adult Baboons (Papio Hamadryas)." Bone 35, no. 1 (July 2004):

198–203. https://doi.org/10.1016/j.bone.2004.02.022.

Hidalgo, M.C, E Urea, and A Sanz. "Comparative Study of Digestive Enzymes in Fish

with Different Nutritional Habits. Proteolytic and Amylase Activities." Aquaculture 170,

no. 3–4 (January 1999): 267–83. https://doi.org/10.1016/s0044-8486(98)00413-x.

Havill, L.M, M.C Mahaney, and J Rogers. "Genotype-by-Sex and Environment-by-

Sex Interactions Influence Variation in Serum Levels of Bone-Specific Alkaline

Phosphatase in Adult Baboons (Papio Hamadryas)." Bone 35, no. 1 (July 2004):

198–203. https://doi.org/10.1016/j.bone.2004.02.022.

Jury, Mark R. "Spreading of the Semi-Arid Climate across South Africa." Journal of

Water and Climate Change 12, no. 8 (October 1, 2021): 3734–49.

https://doi.org/10.2166/wcc.2021.187.

Kasvosve, Ishmael, and Joris Delanghe. "Total Iron Binding Capacity and Transferrin

Concentration in the Assessment of Iron Status." Clinical Chemistry and Laboratory

Medicine 40, no. 10 (January 29, 2002). https://doi.org/10.1515/cclm.2002.176.

Kawecki, Tadeusz J., and Dieter Ebert. "Conceptual Issues in Local Adaptation."

Ecology Letters 7, no. 12 (November 12, 2004): 1225–41.

https://doi.org/10.1111/j.1461-0248.2004.00684.x.

Kobbe, Susanne, Jörg U. Ganzhorn, and Kathrin H. Dausmann. "Extreme Individual

Flexibility of Heterothermy in Free-Ranging Malagasy Mouse Lemurs (Microcebus

Griseorufus)." Journal of Comparative Physiology B 181, no. 1 (August 18, 2010):

165–73. https://doi.org/10.1007/s00360-010-0507-5.

Le Couteur, D. G., F. M. Blyth, H. M. Creasey, D. J. Handelsman, V. Naganathan, P.

N. Sambrook, M. J. Seibel, L. M. Waite, and R. G. Cumming. "The Association of

Alanine Transaminase With Aging, Frailty, and Mortality." The Journals of

Gerontology Series A: Biological Sciences and Medical Sciences 65A, no. 7 (May 24,

2010): 712–17. https://doi.org/10.1093/gerona/glq082.

Lian, Xu, Shilong Piao, Anping Chen, Chris Huntingford, Bojie Fu, Laurent Z. X. Li,

Jianping Huang, et al. "Multifaceted Characteristics of Dryland Aridity Changes in a

Warming World." Nature Reviews Earth & Environment 2, no. 4 (March 9, 2021):

232–50. https://doi.org/10.1038/s43017-021-00144-0.

Lochmiller, Robert L., Michelle R. Vestey, and Jon C. Boren. "Relationship between

Protein Nutritional Status and Immunocompetence in Northern Bobwhite Chicks."

The Auk 110, no. 3 (July 1993): 503–10. https://doi.org/10.2307/4088414.

López-Maury, Luis, Samuel Marguerat, and Jürg Bähler. "Tuning Gene Expression to

Changing Environments: From Rapid Responses to Evolutionary Adaptation." Nature

Reviews Genetics 9, no. 8 (August 2008): 583–93. https://doi.org/10.1038/nrg2398.

Melles, S. J., M.-J. Fortin, K. Lindsay, and D. Badzinski. "Expanding Northward:

Influence of Climate Change, Forest Connectivity, and Population Processes on a

Threatened Species' Range Shift." Global Change Biology 17, no. 1 (December 1,

2010): 17–31. https://doi.org/10.1111/j.1365-2486.2010.02214.x.

McClure, Diane E. "Clinical Pathology and Sample Collection in the Laboratory

Rodent." Veterinary Clinics of North America: Exotic Animal Practice 2, no. 3

(September 1999): 565–90. https://doi.org/10.1016/s1094-9194(17)30111-1.

Merilä, Juha, and Andrew P. Hendry. "Climate Change, Adaptation, and Phenotypic

Plasticity: The Problem and the Evidence." Evolutionary Applications 7, no. 1

(January 2014): 1–14. https://doi.org/10.1111/eva.12137.

Meynard, Christine N., Neville Pillay, Manon Perrigault, Pierre Caminade, and Guila

Ganem. "Evidence of Environmental Niche Differentiation in the Striped Mouse

(Rhabdomys Sp.): Inference from Its Current Distribution in Southern Africa." Ecology

and Evolution 2, no. 5 (April 12, 2012): 1008–23. https://doi.org/10.1002/ece3.219.

Muñoz-Garcia, Agustí, Miriam Ben-Hamo, Shai Pilosof, Joseph B. Williams, and

Carmi Korine. "Habitat Aridity as a Determinant of the Trade-off between Water

Conservation and Evaporative Heat Loss in Bats." Journal of Comparative

Physiology B 192, no. 2 (January 17, 2022): 325–33. https://doi.org/10.1007/s00360-

021-01425-2.

Naumann, G., L. Alfieri, K. Wyser, L. Mentaschi, R. A. Betts, H. Carrao, J. Spinoni, J.

Vogt, and L. Feyen. "Global Changes in Drought Conditions Under Different Levels of

Warming." Geophysical Research Letters 45, no. 7 (April 13, 2018): 3285–96.

https://doi.org/10.1002/2017gl076521.

Naya, Daniel E., Claudio Veloso, Pablo Sabat, and Francisco Bozinovic.

"Physiological Flexibility and Climate Change: The Case of Digestive Function

Regulation in Lizards." Comparative Biochemistry and Physiology Part A: Molecular

& Integrative Physiology 159, no. 1 (May 2011): 100–104.

https://doi.org/10.1016/j.cbpa.2011.02.005.

Nicholson, S.E., and J.C. Selato. "The Influence of La Nina on African Rainfall."

International Journal of Climatology 20, no. 14 (n.d.): 1761–76.

https://doi.org/10.1002/1097-0088(20001130)20:14<1761::AID-JOC580>3.0.CO;2-

W.

Ostrowski, Stéphane, Joseph B. Williams, Pascal Mésochina, and Helga Sauerwein.

"Physiological Acclimation of a Desert Antelope, Arabian Oryx (Oryx Leucoryx), to

Long-Term Food and Water Restriction." Journal of Comparative Physiology B 176,

no. 3 (November 9, 2005): 191–201. https://doi.org/10.1007/s00360-005-0040-0.

Pajic, Petar, Pavlos Pavlidis, Kirsten Dean, Lubov Neznanova, Rose-Anne Romano,

Danielle Garneau, Erin Daugherity, Anja Globig, Stefan Ruhl, and Omer Gokcumen.

"Independent Amylase Gene Copy Number Bursts Correlate with Dietary

Preferences in Mammals." eLife 8 (May 14, 2019).

https://doi.org/10.7554/elife.44628.

Parmesan, Camille. "Ecological and Evolutionary Responses to Recent Climate

Change." Annual Review of Ecology, Evolution, and Systematics 37, no. 1

(December 1, 2006): 637–69.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110100.

Parmesan, Camille, Terry L. Root, and Michael R. Willig. "Impacts of Extreme

Weather and Climate on Terrestrial Biota." Bulletin of the American Meteorological

Society 81, no. 3 (March 1, 2000): 443–50. https://doi.org/10.1175/1520-

0477(2000)081<0443:IOEWAC>2.3.CO;2.

Post, Eric, Jedediah Brodie, Mark Hebblewhite, Angela D. Anders, Julie A. K. Maier,

and Christopher C. Wilmers. "Global Population Dynamics and Hot Spots of

Response to Climate Change." BioScience 59, no. 6 (June 2009): 489–97.

https://doi.org/10.1525/bio.2009.59.6.7.

Rimbach, Rebecca, Stéphane Blanc, Alexandre Zahariev, Maria Gatta, Neville Pillay,

and Carsten Schradin. "Seasonal Variation in Energy Expenditure in a Rodent

Inhabiting a Winter-Rainfall Desert." Journal of Comparative Physiology B 188, no. 5

(June 8, 2018): 877–88. https://doi.org/10.1007/s00360-018-1168-z.

Rimbach, Rebecca, Jörg Jäger, Neville Pillay, and Carsten Schradin. "Food

Availability Is the Main Driver of Seasonal Changes in Resting Metabolic Rate in

African Striped Mice (Rhabdomys Pumilio)." Physiological and Biochemical Zoology

91, no. 2 (March 2018): 826–33. https://doi.org/10.1086/696828.

Rintoul, Jody L. P., and R. Mark Brigham. "The Influence of Reproductive Condition

and Concurrent Environmental Factors on Torpor and Foraging Patterns in Female

Big Brown Bats (Eptesicus Fuscus)." Journal of Comparative Physiology B 184, no. 6

(June 28, 2014): 777–87. https://doi.org/10.1007/s00360-014-0837-9.

Rocha, Joana L., Raquel Godinho, José C. Brito, and Rasmus Nielsen. "Life in

Deserts: The Genetic Basis of Mammalian Desert Adaptation." Trends in Ecology

& Evolution 36, no. 7 (July 2021): 637–50.

https://doi.org/10.1016/j.tree.2021.03.007.

Sabat, Pablo, Sandra Gonzalez-Vejares, and Karin Maldonado. "Diet and Habitat

Aridity Affect Osmoregulatory Physiology: An Intraspecific Field Study along

Environmental Gradients in the Rufous-Collared Sparrow." Comparative Biochemistry

and Physiology Part A: Molecular & Integrative Physiology 152, no. 3 (March

2009): 322–26. https://doi.org/10.1016/j.cbpa.2008.11.003.

Saraç, F., and F. Saygılı. "Causes of High Bone Alkaline Phosphatase."

Biotechnology & Biotechnological Equipment 21, no. 2 (January 2007): 194–97.

https://doi.org/10.1080/13102818.2007.10817444.

Seabrook, Leonie, Clive McAlpine, Greg Baxter, Jonathan Rhodes, Adrian Bradley,

and Daniel Lunney. "Drought-Driven Change in Wildlife Distribution and Numbers: A

Case Study of Koalas in South West Queensland." Wildlife Research 38, no. 6

(2011): 509. https://doi.org/10.1071/wr11064.

Schoepf, I., N. Pillay, and C. Schradin. "The Pathophysiology of Survival in Harsh

Environments." Journal of Comparative Physiology B 187, no. 1 (July 15, 2016): 183–

201. https://doi.org/10.1007/s00360-016-1020-2.

Schoepf, I., N. Pillay, and C. Schradin. "Trade-Offs between Reproduction and Health

in Free-Ranging African Striped Mice." Journal of Comparative Physiology B 187, no.

4 (February 4, 2017): 625–37. https://doi.org/10.1007/s00360-016-1054-5.

Schulte-Hostedde, Albrecht I., Bertram Zinner, John S. Millar, and Graham J.

Hickling. "RESTITUTION OF MASS–SIZE RESIDUALS: VALIDATING BODY

CONDITION INDICES." Ecology 86, no. 1 (January 2005): 155–63.

https://doi.org/10.1890/04-0232.

Seebacher, Frank, and Craig E. Franklin. "Determining Environmental Causes of

Biological Effects: The Need for a Mechanistic Physiological Dimension in

Conservation Biology." Philosophical Transactions of the Royal Society B: Biological

Sciences 367, no. 1596 (June 19, 2012): 1607–14.

https://doi.org/10.1098/rstb.2012.0036.

Seebacher, Frank, William Davison, Cara J Lowe, and Craig E Franklin. "A

Falsification of the Thermal Specialization Paradigm: Compensation for Elevated

Temperatures in Antarctic Fishes." Biology Letters 1, no. 2 (April 28, 2005): 151–54.

https://doi.org/10.1098/rsbl.2004.0280.

~~Shapiro, R, and R.P Heaney. "Co-Dependence of Calcium and Phosphorus for~~

~~Growth and Bone Development under Conditions of Varying Deficiency." Bone 32,~~

~~no. 5 (May 2003): 532–40. https://doi.org/10.1016/s8756-3282(03)00060-7.~~

Solmsen, N., J. Johannesen, and C. Schradin. "Highly Asymmetric Fine-Scale

Genetic Structure between Sexes of African Striped Mice and Indication for Condition

Dependent Alternative Male Dispersal Tactics." Molecular Ecology 20, no. 8 (March

3, 2011): 1624–34. https://doi.org/10.1111/j.1365-294x.2011.05042.x.

Somero, G. N. "The Physiology of Climate Change: How Potentials for

Acclimatization and Genetic Adaptation Will Determine 'Winners' and 'Losers.'"

Journal of Experimental Biology 213, no. 6 (March 15, 2010): 912–20.

https://doi.org/10.1242/jeb.037473.

Somero, George N. "The Physiology of Global Change: Linking Patterns to

Mechanisms." Annual Review of Marine Science 4, no. 1 (January 15, 2012): 39–61.

https://doi.org/10.1146/annurev-marine-120710-100935.

Speakman, John R. "The Physiological Costs of Reproduction in Small Mammals."

Philosophical Transactions of the Royal Society B: Biological Sciences 363, no. 1490

(August 8, 2007): 375–98. https://doi.org/10.1098/rstb.2007.2145.

Stawski, Clare, and A Daniella Rojas. "Thermal Physiology of a Reproductive Female

Marsupial, Antechinus Flavipes." Mammal Research 61, no. 4 (July 21, 2016): 417–

21. https://doi.org/10.1007/s13364-016-0287-8.

Sullivan, Jennifer C., Laura Semprun-Prieto, Erika I. Boesen, David M. Pollock, and

Jennifer S. Pollock. "Sex and Sex Hormones Influence the Development of

Albuminuria and Renal Macrophage Infiltration in Spontaneously Hypertensive Rats."

American Journal of Physiology-Regulatory, Integrative and Comparative Physiology

293, no. 4 (October 2007): R1573–79. https://doi.org/10.1152/ajpregu.00429.2007.

Sundar Ray, Chinmaya, Bratati Singh, Itishri Jena, Sudeshna Behera, and

Subhashree Ray. "Low Alkaline Phosphatase (ALP) In Adult Population an Indicator

of Zinc (Zn) and Magnesium (Mg) Deficiency." Current Research in Nutrition and

Food Science Journal 5, no. 3 (December 19, 2017): 347–52.

https://doi.org/10.12944/crnfsj.5.3.20.

Taylor, K. D., and M. G. Green. "The Influence of Rainfall on Diet and Reproduction

in Four African Rodent Species." Journal of Zoology 180, no. 3 (November 1976):

367–89. https://doi.org/10.1111/j.1469-7998.1976.tb04683.x.

Thaler, M Michael, David L Gemes, and Arne F Bakken. "Enzymatic Conversion of

Heme to Bilirubin in Normal and Starved Fetuses and Newborn Rats." Pediatric

Research 6, no. 3 (March 1972): 197–201. https://doi.org/10.1203/00006450-

197203000-00008.

Thornthwaite, C. W. "An Approach toward a Rational Classification of Climate."

Geographical Review 38, no. 1 (January 1948): 55. https://doi.org/10.2307/210739.

Tothova, C., O. Nagy, and G. Kovac. "Serum Proteins and Their Diagnostic Utility in

Veterinary Medicine: A Review." Veterinární Medicína 61, no. 9 (September 30,

2016): 475–96. https://doi.org/10.17221/19/2016-vetmed.

Tshikae, Balatlhane P., Adrian L. V. Davis, and Clarke H. Scholtz. "Does an Aridity

and Trophic Resource Gradient Drive Patterns of Dung Beetle Food Selection across

the Botswana Kalahari?" Ecological Entomology 38, no. 1 (January 22, 2013): 83–95.

https://doi.org/10.1111/j.1365-2311.2012.01409.x.

Usui, Takuji, David Lerner, Isaac Eckert, Amy L. Angert, Colin J. Garroway, Anna

Hargreaves, Lesley T. Lancaster, et al. "The Evolution of Plasticity at Geographic

Range Edges." Trends in Ecology & Evolution 38, no. 9 (September 2023):

831–42. https://doi.org/10.1016/j.tree.2023.04.004.

Vespasiani-Gentilucci, Umberto, Antonio De Vincentis, Luigi Ferrucci, Stefania

Bandinelli, Raffaele Antonelli Incalzi, and Antonio Picardi. "Low Alanine

Aminotransferase Levels in the Elderly Population: Frailty, Disability, Sarcopenia, and

Reduced Survival." The Journals of Gerontology: Series A 73, no. 7 (June 19, 2017):

925–30. https://doi.org/10.1093/gerona/glx126.

"Clinical Methods: The History, Physical, and Laboratory Examinations." Annals of

Internal Medicine 113, no. 7 (October 1, 1990): 563. https://doi.org/10.7326/0003-

4819-113-7-563_2.

Webster, Nicole S, and Thorsten B H Reusch. "Microbial Contributions to the

Persistence of Coral Reefs." The ISME Journal 11, no. 10 (May 16, 2017): 2167–74.

https://doi.org/10.1038/ismej.2017.66.

Willmer, Pat, Graham Stone, and Ian Johnston. Environmental Physiology of

Animals. John Wiley & Sons, 2009.

Wu, Nicholas C., and Frank Seebacher. "Physiology Can Predict Animal Activity,

Exploration, and Dispersal." Communications Biology 5, no. 1 (February 3, 2022).

https://doi.org/10.1038/s42003-022-03055-y.

Yousef, M.I., H.A. El Hendy, F.M. El-Demerdash, and E.I. Elagamy. "Dietary Zinc

Deficiency Induced-Changes in the Activity of Enzymes and the Levels of Free

Radicals, Lipids and Protein Electrophoretic Behavior in Growing Rats." Toxicology

175, no. 1–3 (June 2002): 223–34. https://doi.org/10.1016/s0300-483x(02)00049-5.

Zhang, Jianfen, Na Zhang, Shufang Liu, Songming Du, and Guansheng Ma. "Young

Adults with Higher Salt Intake Have Inferior Hydration Status: A Cross-Sectional

Study." Nutrients 14, no. 2 (January 11, 2022): 287.

https://doi.org/10.3390/nu14020287.

Zhao, Miao-Miao, Jeremy Krebs, Xi Cao, Jing Cui, Dong-Ning Chen, Yu Li, Lin Hua,

Jim Mann, and Jin-Kui Yang. "Helicobacter Pylori Infection as a Risk Factor for

Serum Bilirubin Change and Less Favourable Lipid Profiles: A Hospital-Based Health

Examination Survey." BMC Infectious Diseases 19, no. 1 (February 14, 2019).

https://doi.org/10.1186/s12879-019-3787-8.

Zomer, Robert J., Jianchu Xu, and Antonio Trabucco. "Version 3 of the Global Aridity

Index and Potential Evapotranspiration Database." Scientific Data 9, no. 1 (July 15,

2022). https://doi.org/10.1038/s41597-022-01493-1.

Tables

Table 1: Summary of main ecological characteristics of R.bechuanae and

R.d.dilectus and of our predictions on their physiological response during the

dry season

ECOLOGICAL CHARACTERISTICS	R.BECHUANAE	R.D.DILECTUS
ENVIRONMENTAL NICHE	Arid to semi-arid (Ganem et al., 2020)	Mesic to semi-arid (Ganem et al., 2020)
PREFERRED HABITAT	Bushy patches (Dufour et al., 2015)	Continuous cover / grassland (Dufour et al., 2015)
SOCIAL STRUCTURE	Group-living (Dufour et al., 2018)	Solitary / opportunistic group-living (Dufour et al., 2018)
SPECIES-SPECIFIC RESPONSE TO SEASONAL VARIATIONS OF DRY CONDITIONS: PREDICTIONS		
BODY CONDITION	↘↘ or ↘*	↘↘
NUTRITIONAL METABOLISM	↘↘ or ↘*	↘↘
LIVER ACTIVITY	↘↘ or ↘*	↘↘
WATER BALANCE	↘↘ or ↘*	↘↘
SENSITIVITY TO INFECTION	↗↗ or ↗*	↗↗
SPECIES DIFFERENCES: PREDICTIONS		
BODY CONDITION	+	-
LIVER ACTIVITY	+	-
WATER BALANCE	+	-
SENSITIVITY TO INFECTION	-	+

1000

1001 *Adaptation to harsher arid conditions could have enabled *R.bechuanae* to withstand growing aridity

1002 without suffering as many adverse effects as *R.d.dilectus* in the semi-arid zone. Alternatively, semi-

1003 arid populations of both species could both have developed local adaptation.

1004

1005 ***Table 2: Geographical position and Aridity Index of sampled sites for each***
1006 ***species***

SPECIES	SAMPLED SITES	GEOGRAPHICAL COORDINATES	ARIDITY INDEX[*]
R.bechuanae	Gariep Dam Nature Reserve	E25.55537; S30.57589	0.42 ± 0.07
	Kalkfontein Dam Nature Reserve	E25.28031; S29.52058	0.54 ± 0.07
	Benfontein Nature Reserve**	E24.81824; S28.82747	0.39 ± 0.07
R.d.dilectus	Bloemhof Dam Nature Reserve	E25.65144; S27.66063	0.35 ± 0.06
	Barberspan Bird Sanctuary**	E25.58354; S26.56538	0.39 ± 0.05
	Wolwespruit Nature Reserve	E26.26326; S27.41418	0.53 ± 0.10

1007 *Average Aridity index (see **Supplementary material** for formula) ± Standard Error

1008 of the Mean, based on climate data collected from 2010 to 2021 (South African

1009 Weather Service)

1010 ** Sampled only in May 2022 (dry season onset).

1011

1012 ***Table 3: Results of the Analysis of Variance (ANOVA) test addressing variation***
1013 ***in NDVI***

Variable	Df	R²	F	P
Session	1	0.445	442.757	**$< 2.2*10^{-16}$**
Site	3	0.348	115.385	**$< 2.2*10^{-16}$**
Session*Site	3	0.021	7.113	**$1.512*10^{-4}$**
Residual	229	0.185		

1014

1015

1016 ***Table 4: Results of the Analysis of Variance (ANOVA) test addressing variation***
1017 ***in body condition***

Variable	Df	R^2	F	P
Session	1	$1.271*10^{-5}$	0.0019	0.965
Sex	1	0.085	25.163	**$1.072*10^{-6}$**
NDVI	1	$8.007*10^{-4}$	0.239	0.626
Age class	3	0.091	8.990	**$1.201*10^{-5}$**
Breeding status	2	0.041	6.067	**0.003**
Session*Sex	1	$5.592*10^{-4}$	0.167	0.683
Session*Breeding status	2	0.0299	4.444	**0.012**
Residual	229	0.753		

1018

1019

1020 ***Table 5: Summary of the results of the best fitting Permutational multivariate***

1021 ***analysis of variance (PERMANOVA) model, addressing variation in***

1022 ***physiological response (site= random factor)***

1023

Variable	Df	R²	Pseudo-F	P
Session	1	0.011	2.882	0.067
Species	1	0.042	11.070	**0.003**
Sex	1	0.010	2.748	0.087
Breeding status	2	0.033	4.320	**0.011**
Species * Sex	1	0.016	4.232	**0.029**
Species * Breeding status	2	0.026	3.413	**0.037**
Residual	227	0.862		

1024

1025

Table 6: Summary of the results of univariate mixed models analyses of

variance addressing variation in responses of each of the 12 studied

physiological variables (best fitting models).

1029

Physiological marker	Variable(s) kept in Best Model and their significance	adjusted R²	F	df Model	df Residual	P (model)
Albumin	**Session*****, **Sex***	0.098	13.72	2	233	$2.338*10^{-6}$
Alkaline phosphatase	**Session*****, **Species***, **Sex*****, Breeding status+	0.260	17.43	5	229	< $1.281*10^{-14}$
Alanine aminotransferase	**Session***, Species, Sex, Breeding status, Species*Sex, Species*Breeding Status	0.060	2.884	8	227	0.004
Amylase	**Species*****, Sex, Breeding status, Species*Sex, Species*Breeding Status	0.185	8.603	7	228	$3.34*10^{-9}$
Calcium	Species	0.009	3.095	1	234	0.080
Globulin	**Session*****	0.291	97.44	1	234	< $2.2*10^{-16}$
Glucose	Session+, **Sex****	0.079	11.14	2	233	$2.397 *10^{-5}$
Potassium	**Session***, **Species***	0.068	9.569	2	233	$1.014*10^{-4}$
Sodium	Session, **Species*****, Sex	0.110	10.65	3	232	$1.39*10^{-6}$
Blood Urea Nitrogen	_Null Random effect model_	0.072	/	/	/	/
Phosphorus	Species, Sex, **Breeding status*** + random	0.248	/	/	/	($\square^2$ test vs. null model) $3.471*10^{-7}$

 Note: The full models, before backwards elimination, included session, species, body condition, sex,

 breeding status, sex*species and breeding status*species as variables, with breeding status nested

 within sex and site as a partially-crossed random factor nested within species.

 Bold: _significant effects (p < 0.05) following Bonferroni correction with:_

 _***: p < 0.001; **: 0.001 < p < 0.01 ; *: 0.01 < p < 0.05; + : 0.05<p<0.1_

Figure captions

Figure 1: Map of known occurrences of 6 main clades of Rhabdomys (translucent dots) and locations sampled in this study (full colour dots) in southern Africa. *Based on published data and unpublished data (origin details are available in doi: 0d1b3414-7e2a-11ea-a38d-00163e26bfb0). Base map: World Topographic Map Esri Standard, Aridity Index layer was computed from a 0.5° global grid, using data from the Version 3 of the Global Aridity Index and Potential Evapotranspiration Database (Zomer et al.,2022).*

Figure 2: Graphical summary of organization and composition of data used for analysis in this study. *Figures between brackets indicate number of modalities or types of categories included in the factor.*

Figure 3: Total monthly precipitation for all sampled sites as recorded in 2022.

Figure 4: Average Normalized Differential Vegetation Index calculated within a buffer circle of 60m radius around each successful trap per site and session. *NDVI was retrieved from the Copernicus Open Access Data Hub (Copernicus Sentinel-2 data [2023]). For each site and session, we retained NDVI data available for the closest day to the beginning of a sampling session, expected to represent the conditions experienced by the mice at the time of capture.*

Figure 5: Significant shifts in blood physiological marker concentrations between the onset and the end of the dry season (mean ± SE)

Figure 6: Significant interspecific differences involving four blood physiological marker concentrations (mean ± SE)

Figures

Based on published data and unpublished data (origin details are available in doi:

0d1b3414-7e2a-11ea-a38d-00163e26bfb0). Base map: World Topographic Map Esri

Standard, Aridity Index layer was computed from a 0.5° global grid, using data from

the Version 3 of the Global Aridity Index and Potential Evapotranspiration Database

(Zomer et al.,2022).

1067

1068

1069

1070

1071

1072

1073

1074

1075   ***<u>Figure 3: Total monthly precipitation for all sampled sites as recorded in 2022.</u>***

1076

1077

Figure 4: Average Normalized Differential Vegetation Index calculated within a

buffer circle of 60m radius around each successful trap per site and session.

1080   *NDVI was retrieved from the Copernicus Open Access Data Hub (Copernicus*

1081   *Sentinel-2 data [2023]). For each site and session, we retained NDVI data available*

1082   *for the closest day to the beginning of a sampling session, expected to represent the*

1083   *conditions experienced by the mice at the time of capture.*

1084

1085

1086

1087

1088

1089 ***Figure 5: Significant shifts in blood physiological marker concentrations***

1090 ***between the onset and the end of the dry season (mean ± SE)***

1091

1092

a : Alkaline phosphatase

b : Potassium

c : Sodium

d : Total bilirubin

1094   ***Figure 6: Sig______nt interspecific differences involving five_____od physiological***

1095   ***marker concentrations (mean ± SE)***

1096

1097