cGAS-STING is responsible for aging of telomerase deficient zebrafish

Naz Şerifoğlu[1], Giulia Allavena[1], Bruno Bastos-Lopes[1], Marta Marzullo[2,*], Pavlos Bousounis[3,4],

Eirini Trompouki[1] and Miguel Godinho Ferreira[1]

1- Institute for Research on Cancer and Aging of Nice (IRCAN), CNRS UMR7284, INSERM U1081,

Université Cote d'Azur, 06107 Nice, France.

2- Instituto Gulbenkian de Ciência, Oeiras, Portugal

3- Department of Cellular and Molecular Immunology, Max Planck Institute of Immunobiology
and Epigenetics, Freiburg, Germany

4- Faculty of Biology, University of Freiburg, Freiburg, Germany

*current address: Department of Biology and Biotechnologies, Sapienza University of Rome,

Rome, Italy.

30 Corresponding Author: Miguel-Godinho.FERREIRA@unice.fr

33   **Keywords:** Telomerase, cGAS-STING, inflammation, aging, zebrafish

Abstract (1088 characters/151 words)

Telomere shortening occurs in multiple tissues throughout aging. When telomeres become critically short, they trigger DNA damage responses and p53 stabilization, leading to apoptosis or replicative senescence. *In vitro*, cells with short telomeres activate the cGAS-STING innate immune pathway resulting in type I interferon inflammation and senescence. However, the consequences of these events to the organism are not yet understood. Here, we show that *sting* is responsible for premature aging of telomerase-deficient zebrafish. We generated *sting-/- tert-/-* double mutants and observed a thorough rescue of *tert-/-* phenotypes. At the cellular level, lack of cGAS-STING in *tert* mutants resulted in reduced senescence, increased cell proliferation, and low inflammation despite similar short telomeres. Critically, absence of *sting* function resulted in dampening of the DNA damage response and low p53 levels. At the organism level, *sting-/- tert-/-* zebrafish regained fertility, delayed cachexia, and cancer incidence, resulting in increased healthspan and lifespan of telomerase mutants.

Introduction (3612 characters /489 words)

Telomerase deficiency in humans results in the development of telomere biology disorders (TBDs) that include idiopathic pulmonary fibrosis, dyskeratosis congenita and aplastic anemia[1]. Common aspects of TBDs relate to accelerated telomere shortening, loss of tissue regeneration, premature aging phenotypes and shorter life span[2,3]. Zebrafish possess human-like telomere lengths that shorten to critical lengths during their lifetime. Like human TBDs, telomerase-deficient zebrafish (*tert-/-*) have accelerated telomere shortening, low cell proliferation, tissue damage, and reduced lifespan [4-6]. *tert-/-* zebrafish also develop chronic inflammation, increased infections, and accelerated incidence of cancer[7,8]. Like in humans[9], we have recently shown that not all organs age at the same rate[8]. The zebrafish intestine becomes dysfunctional earlier and triggers systemic aging. Reduced proliferation of intestine cells results in loss of tissue integrity, microbiota dysbiosis and systemic inflammation[8].

Type I interferon response and secretion of pro-inflammatory cytokines through activation of cGAS-STING is triggered by DNA damage, including telomeric damage, leading to the formation of micronuclei (MN)[10-13]. The cytosolic DNA sensor, cGAS (cGMP-AMP synthase), becomes activated upon binding to double-stranded DNA, encompassing both microbial and self-DNA[14]. Upon recognition of cytosolic DNA, cGAS initiates the production of the second messenger cGAMP that binds and activates the adaptor protein STING[15,16]. Subsequently, STING

recruits TBK1 (TANK-binding kinase 1) triggering the activation of IRF3 (IFN regulatory factor 3).

This activation cascade leads to the generation of type I interferons and inflammatory

cytokines[14]. cGAS-STING is involved in DNA repair, DNA damage responses (DDR) and cell

senescence (reviewed in[17]). These effects are achieved through the expression of interferon-

stimulated genes (ISGs) and the senescence-associated secretory phenotype (SASP)[12,16].

Secretion of these molecules modulates the proliferative capacity of surrounding cells in a

paracrine manner, propagating the senescence status (paracrine-SASP)[18].

Type I interferon response is increasingly linked to aging and neurodegenerative diseases

across different species. A recent significant study by the Ablasser lab using naturally aged mice

revealed that cGAS–STING signaling plays a pivotal role in the age-related type I interferon

response in neurodegeneration[19]. Transcriptional profiling revealed that cGAS–STING activation

initiated a gene expression program shared between neurodegenerative diseases and natural

aging. The authors also observed that there was an accumulation of mitochondrial DNA in the cell

cytoplasm of microglial cells, providing a potential mechanism through which the cGAS-STING

pathway may contribute to inflammation in the aging brain[19].

In vitro studies using human primary cells recently showed that cGAS-STING is activated

by short or dysfunctional telomeres (reviewed in[13]). However, it is currently unknown what are the

consequences of activation of cGAS-STING in response to short telomeres at the organism level.

Here, we show that telomere shortening triggers cGAS-STING pathway in skin, testis, kidney

marrow (the adult hematopoietic organ in fish) and intestine of zebrafish. This results in type I

interferon response, elevation in senescence levels and reduction in proliferative capacity of these

tissues. Absence of cGAS-STING restores cell proliferation, suppresses accelerated aging and

increases in both health and lifespan of the telomerase deficient zebrafish.

Results (14998 characters/ 2212 words)

Telomere shortening activates the cGAS STING and type I interferon *in vivo*

Our previous studies show that telomerase deficient zebrafish undergo accelerated

systemic inflammation[5–8,20,21]. Analysis of gene expression of proliferative (gut) and non-

proliferative tissues (muscle) derived from aged *tert-/-* (9-months old) and WT (36-months old)

zebrafish highlighted several genes related to type I interferon response[20]. To investigate if short

telomeres trigger type I inflammation and accelerated aging through the cGAS-STING pathway,

we combined *sting* zebrafish loss of function mutants (*sting^{sa35634/sa35634}*, hereby referred to as

sting-/-) with telomerase deficient zebrafish (*tert^{hu3430/hu3430}* or *tert-/-*) extensively characterized in

our previous studies[5–8,21]. Upon incross of double heterozygous fish, we first investigated if *tert-/-*

sting-/- double mutants had short telomeres comparable to their *tert-/-* siblings. We observed

that, by 9 months of age, *tert-/- sting-/-* had similar mean telomere length to *tert-/-* single mutants

in the skin and intestine (Figure 1A-D, Supplementary figure 1A-D). In testis, *tert-/- sting-/-*

zebrafish had slightly shorter telomeres than *tert-/-* (Figure 1B, Supplementary Figure 1B) and

slightly longer telomeres in the kidney marrow (Figure 1C, Supplementary Figure 1C). Overall, not

only telomere length was similar between *tert-/-* and *tert-/- sting-/-* but were significantly shorter

than WT and *sting-/-* siblings.

Previous studies using cell lines reported that short and dysfunctional telomeres lead to

the formation of MN[10–12]. To test whether we would observe an increase of MN upon telomere

shortening in zebrafish, we derived fibroblasts from the skin of 9-months old animals. We

observed a ten-fold increase in MN formation in *tert-/-* zebrafish when compared to the WT and

sting-/- siblings. Strikingly, *tert-/- sting-/-* had similar levels of MN as *tert-/-* siblings (Figure 1E).

Thus, telomere shortening results in MN accumulation *in vivo*.

After confirming the presence of shorter telomeres and MN in *tert-/-* and *tert-/- sting-/-*

zebrafish, we investigated if the cGAS-STING pathway was active *in vivo* by quantifying its

downstream targets. Therefore, using skin of 9-months-old zebrafish, we analyzed the

phosphorylation status of zebrafish Tbk1 and Irf3 and the transcription level of two members of

type I interferon response, *isg15* and *ifn-i* (Figure 1F-H). Comparing *tert-/-* to WT zebrafish we

observed ca. 3-fold increase in p-Irf3 and 2-fold increase in p-Tbk1 (Figure 1F). Consistently,

expression of *isg15* and *ifn-i* was increased by 10- and 2.5-fold, respectively (Figure 1G-H).

However, the phosphorylation profile of Tbk1 and Irf3 in *tert-/- sting-/-* mutants was similar to

their WT and *sting-/-* siblings (Figure 1F). *tert-/- sting-/-* mutants also lacked type I interferon
response, as observed by the reduced levels of *isg15* and *ifn-i* (Figure 1G-H).

We expanded our study to other proliferative tissues like testis, kidney marrow and
intestine since they all show signs of inflammation in aging zebrafish[5,8,21]. Comparing
transcription levels of *tert-/-* fish to WT fish, we observed 15-fold increase in expression for *isg15*
and *ifn-i* in testis; 20-fold increase in *isg15 and* 5 fold increase in kidney marrow; and, lastly, 7.5-
fold increase in *isg15* transcript and 2.5-fold increase in *ifn-i* in the intestine. Importantly, inactive
cGAS-STING pathway rescued the *isg15* and *ifn-i* levels of *tert-/- sting-/-* to those observed in WT
zebrafish in all tissues analyzed (Figure 1G, H). These results show that, despite the presence of
shorter telomeres and MN, the cGAS-STING pathway is inactive in *tert-/- sting-/-* mutants.

Type I interferon response can be initiated by mobilization of transposable elements (TEs).
During replicative senescence of human fibroblasts, L1 retrotransposable elements become
transcriptionally derepressed and activate a type-I interferon response[22]. TE transcription leads to
robust activation of RIG-I and MDA5 that recognize dsRNA and ssRNA and initiate a signaling
cascade, involving MAVS oligomerization, IRF3/IRF7 activation and the expression of type I
interferon response. We looked for the deregulation of TEs expression in zebrafish testis, kidney
marrow and intestine (Supplementary Fig. 2A-B). Multidimensional scaling (MDS) plots of TE
expression were non-overlapping for WT and *tert-/-* mutant in all tissues analyzed
(Supplementary Fig. 2A). Telomerase-deficient fish showed a clear upregulation of TEs in testis
and kidney marrow and, to a lesser extent, in the gut.

We then investigated whether the expressions of the RNA sensors *rig-I*, *mda5* and *mavs*
were increased in telomerase-deficient animals. Similar to TE expression, *rig-I* and *mda5* were
overexpressed in *tert-/-* kidney marrow and testis, respectively, compared to WT animals, but not
the intestine (Supplementary Fig. 2C-D). Recently, the Karlseder lab showed that both DNA
sensing (cGAS-STING) and RNA sensing (ZBP1-MAVS) innate immunity pathways orchestrate
telomere replicative crisis (M2) [23]. Their model proposes that short telomeres and MN trigger
cGAS-STING, whereas ZBP1 senses the telomeric lncRNA TERRA and activates MAVS. Since both
pathways result in the activation of type I interferon response, we tested if ZBP1-MAVS could also
contribute to IFN expression. As zebrafish lack a clear *zbp1* gene orthologue, we relied solely on
expression of *mavs* for this analysis. *mavs* mRNA was significantly increased in the testis, but not
in kidney marrow and intestine (Supplementary Figure 2E). Thus, we observed that telomere
shortening triggers multiple responses that include TE expression deregulation and RNA sensor
pathway activation which, in turn, reinforce the expression of type I interferon.

cGAS-STING is required for increased p53 levels in presence of DNA damage

We previously reported that telomere shortening leads to increased p53 levels as a consequence of activation of DDR in zebrafish [5,6,8,21]. To identify the cause for activation of cGAS-STING in prematurely aged *tert-/-* mutants, we analyzed DNA damage and activation of DDR in 9-month-old animals. First, we quantified the number of phosphorylated H2AX (γ-H2AX) stained cells by immunofluorescence in the tissues of interest. As in our previous work[8], γ-H2AX staining was dispersed through the nucleoplasm and aggregated in foci (Figure 2A insets). *tert-/-* zebrafish displayed 3-fold increase of γ-H2AX stained cells in the skin (Figure 2B), 4-fold in the testis (Figure 2C) and 5-fold increase both in the kidney marrow (Figure 2D) and the intestine (Figure 2E) when compared to WT and *sting-/-* mutants. Contrarily, *tert-/- sting-/-* mutants showed similar levels of γ-H2AX compared to the *tert-/-* sibling. (Figure 2A-D). Thus, consistent with shorter telomere length in *tert-/-* and *tert-/- sting-/-* mutants, we observed similar levels of the DNA damage marker γ-H2AX in these tissues.

Increased γ-H2AX levels in *tert-/-* zebrafish is accompanied by elevation in p53 protein levels[8,21]. *tert-/-* zebrafish exhibited an increase of 5-fold (Figure 2F,G) in the skin and testis (Figure 2F,H), 3-fold in the kidney marrow (Figure 2F,I) and 6-fold in the intestine (Figure 2F,J) when compared to WT and *sting-/-* mutants (Figure 2E-H). Surprisingly, p53 levels in the *tert-/- sting-/-* mutants were similar to the WT and *sting-/-* siblings (Figure 2F-J). Our results indicate that, despite similar levels of phosphorylated H2AX, cGAS-STING is required for elevated p53 levels in response to telomere shortening. This is in agreement with studies showing that p53 expression and stability are regulated by IFNs[24,25].

cGAS-STING is required for senescence and SASP caused by telomere shortening

Given that p53 was not elevated in aging *tert-/- sting-/-* zebrafish, despite the shorter telomere length, we decided to assess the remaining phenotypes linked with *tert-/-* premature aging. Previous studies in human cells showed that cGAS-STING is required for cell senescence[10–12]. Using 9-months-old zebrafish, we studied cell senescence using the SA-Beta-Galactosidase (SA-B-gal) assay in addition to expression of *cdkn2a/b* (p15/16) and *cdkn1a* (p21) by RT-qPCR. As previously observed [5,6,8,21], proliferative tissues of *tert-/-* mutants were strongly stained with SA-B-gal, but not those of WT or *sting-/-* siblings (Figure 3A). Consistently, we confirmed an elevated expression of *cdkn2a/b* and *cdkn1a* senescence markers in *tert-/-* mutants (Figure 3B-D). Senescence in *tert-/-* was accompanied by expression of SASP-related genes, namely *il1b, tgfb1b*

and *mmp15a*. As expected, none of these genes were elevated in WT or *sting-/-* siblings (Figure

3B-D).

With lower levels of p53, cell senescence was also reduced in *tert-/- sting-/-* tissues, as

observed by low SA-Beta-gal and expression of *cdkn2a/b* and *cdkn1a*. In absence of senescence,

expression of SASP factors were also reduced in *tert-/- sting-/-* mutants to the ones observed in

WT (Figure 3B-D). Thus, in agreement with previous *in vitro* studies, our results indicate that

cGAS-STING is required for senescence and SASP of proliferative tissues *in vivo*.

cGAS-STING controls cell proliferation and tissue integrity of *tert-/-* zebrafish

Replicative cell senescence is a barrier against cell proliferation in response to telomere

shortening. We thus investigated if absence of senescence in *tert-/- sting-/-* mutants would result

in increased cell proliferation. We examined proliferative tissues by immunofluorescence using

antibodies against PCNA, a marker for cell proliferation. As previously reported[5,6,8], cell

proliferation of *tert-/-* zebrafish is significantly reduced in proliferative tissues when compared to

WT and *sting-/-* siblings (Figure 4B-D). However, lower cell proliferation was recovered to WT

levels in *tert-/- sting-/-* zebrafish (Figure 4B-D). Thus, with lower cell senescence and p53 levels,

absence of cGAS-STING results in an increase in cell proliferation despite telomere shortening in

telomerase-deficient zebrafish.

Previously, we showed that telomere shortening impacts tissue integrity and results in

morphological defects of several tissues [5,6,8,21]. Inflammation of the intestine causes an increase in

thickness of the *lamina propria* in *tert-/-* zebrafish (mean of 17.5 micrometer, Figure 4F-G).

However, width of the *lamina propria* in *tert-/- sting-/-* zebrafish was similar to the WT and *sting-/-*

siblings (Figure 4G). Loss of gut tissue integrity activates the YAP-TAZ pathway in aging *tert-/-*

zebrafish[8]. In agreement, we observed an increase of *ctfg* and *cyr61* levels, targets of YAP-TAZ

pathway, (Figure 4L), in *tert-/-* zebrafish compared to WT and *sting-/-* siblings. However,

expressions of YAP-TAZ target genes were reduced to the WT levels in the *tert-/- sting-/-*

zebrafish (Figure 4L). Similarly, the remaining proliferative tissues also showed increased

expression of *ctgf* and *cyr61* in *tert-/-* zebrafish that were absent in *tert-/- sting-/-* siblings (skin:

2.5-fold and 5-fold, respectively, kidney marrow: ~10 fold, Figure 4M, Supplementary Figure 4B-

D). Thus, our results suggest that tissue integrity of the intestine and other tissues caused by

telomere shortening is rescued by an increase of cell proliferation upon inactivation of the cGAS-

STING pathway.

cGAS-STING causes premature aging of *tert-/-* zebrafish

Given the previous results at the cellular level, we investigated the consequences to the whole organism of absence cGAS-STING and Type I interferon in response to telomere shortening. We previously established male fertility and testicular atrophy as robust assays for aging zebrafish [5,6,26]. To measure testicular atrophy, we quantified the mature sperm area in HE stained sections of whole testis (Figure 4H). Percentage of mature sperm area of 9-month-old *tert-/-* zebrafish was reduced to ~5%, compared to ~50% in WT and *sting-/-* siblings (Figure 4I). However, the percentage of mature sperm area increased to ~40% in *tert-/- sting-/-* zebrafish (Figure 4I). Denoting the observed loss of tissue integrity in testis in *tert-/-* prematurely aged zebrafish, we saw increased expression of the YAP-TAZ pathway targets in *ctgf* and *cyr61* compared to WT and *sting-/-* siblings (Figure 4M). Consistent with our previous results, expression of *ctfg* and *cyr61* were reduced in *tert-/- sting-/-* to WT levels (Figure 4M).

We next asked if the rescue in the testis morphology of *tert-/- sting-/-* would restore fertility of aging *tert-/-* zebrafish. We previously reported that, from 6-month-old, *tert-/-* males become infertile paralleling the loss fertility observed in 18-month-old WT fish [5]. As expected, 9-month-old *tert-/-* males were unable to produce fertilized eggs when crossed with young WT females (Figure 5A). Strikingly, *tert-/- sting-/-* males were still fertile even if slightly lower than the WT and *sting-/-* siblings (Figure 5A). Our data shows that inhibition of cGAS-STING and type I interferon response is sufficient to restore fertility to aging *tert-/-* zebrafish.

As other vertebrates, incidence of cancer increases with age in zebrafish [5,26,27]. Similar to other age-associated phenotypes, the spontaneous cancer incidence is accelerated in younger ages in *tert-/-* zebrafish [5,26]. We quantified the rate of spontaneous tumor formation, mostly seminomas, in male zebrafish. We observed that *tert-/-* zebrafish developed tumors from the age of 11-month-old and by the age of 17 months (Figure 5D). Strikingly, *tert-/- sting-/-* mutants developed macroscopic tumors around the age of 12 months but were restricted to 5% of the population (Figure 5D). Importantly, spontaneous cancer incidence of *tert-/- sting-/-* zebrafish was not statistically different from the WT and *sting-/-* siblings.

Other phenotypes of aging, such as kyphosis (abnormal curvature of the spine), caused by increased weakness of the spinal bones, and cachexia (excessive muscle wasting), caused by muscle tissue atrophy, are present in younger *tert-/-* mutants and only appear later in WT zebrafish [5,6]. By the age of 15 months, 60% of *tert-/-* zebrafish showed aging phenotypes (Figure 5B) and weighed significantly less (Figure 5C) than WT and *sting-/-* siblings. However, the

incidence of aging phenotypes ameliorated in *tert-/- sting-/-* mutants (Figure 5B) and weight was
restored to WT levels (Figure 5C).

Finally, we compared the lifespan of *tert-/-* mutants to their *tert-/- sting-/-* siblings.
Whereas *tert-/-* mutants had a mean lifespan of 17 months-old, lifespan was extended to 24
months in *tert-/- sting-/-* mutants (Figure 5E). The mean lifespan of *tert-/- sting-/-* zebrafish was
not statistically different from WT and *sting-/-* siblings (Figure 5F). Overall, our results show that
by inhibiting cGAS-STING and, consequently, type I interferon, we observed an increase in
lifespan of a prematurely aging vertebrate model by 41%. More importantly, we recover most
age-associated phenotypes of aging *tert-/-* mutants, increasing their healthspan without the
increase in cancer incidence.

Discussion (7452 characters/1053 words)

Aging is accompanied by a wide range of physiological changes, such as chronic
inflammation[28]. Age-associated inflammation in absence of overt infection has been termed
inflammaging[29]. While the origins of inflammaging are mostly unclear, it is typically characterized
by high levels of pro-inflammatory cytokines, chemokines, acute phase proteins, and soluble
cytokine receptors in the serum [29,30]. Inflammaging was shown to contribute to the development
of age-associated diseases, such as neurodegenerative diseases, cardiovascular diseases and
cancer [28]. These are the main causes of morbidity and mortality in the elderly. Recent exciting
new data revealed that DNA damage and inflammation are connected by the cGAS-STING
pathway [10,12,31]. Moreover, chemical inhibition of STING suppresses aging-associated
inflammation and neurodegeneration[19]. Our study extends these observations by showing that
activation of cGAS-STING pathway caused by telomere shortening is responsible for premature
aging in zebrafish.

What triggers inflammaging upon telomere shortening? We documented several potential
triggers for type I interferon responses in our work. We observed an increase of MN in *tert-/-*
zebrafish. However, recent data has shown that, even though cGAS is recruited to MN, the
presence of chromatin might not lead to elevated cGAMP and STING activation[32]. Further
evidence from aging mice showed that mtDNA released from disrupted mitochondria was an
important source for cGAS-STING activation[19]. In our previous work[21], we showed that
telomerase mutants have dysfunctional mitochondria with disrupted membranes providing a
likely source for cytoplasmic mtDNA as an additional trigger. Moreover, mirroring what was
previously observed in human fibroblasts[22,23], we documented a tissue-specific elevation of TE

expression, accompanied by increased expression of the RNA sensing pathway Rig-I, Mda5 and

Mavs, that contribute and likely reinforce the activation type I interferon during aging. Finally,

expression of the telomeric lncRNA TERRA could also contribute to the activation of cGAS-STING

via the ZBP1/MAVS pathway, as reported for p53-deficient human cells undergoing crisis[23].

However, we were unable to detect a clear elevation of *mavs* expression in all tissues. A likely

explanation may relate to *tert-/-* zebrafish being p53 proficient and, therefore, their comparatively

longer telomeres may not express sufficient levels of TERRA to trigger its response.

Our work reveals that DDR and senescence triggered by short telomeres require an active

cGAS-STING pathway. Even though telomere length and γ-H2AX levels were similar between

tert-/- and *tert-/- sting-/-* zebrafish, p53 was only elevated in *tert-/-* mutants. Although a definite

explanation is currently unavailable, we suggest potential leads for this observation. First, IFN-b

signaling was shown to induce p53 transcription [25]. Therefore, cGAS-STING may be required to

promote p53 expression independently of canonical DDR. Second, the interferon-stimulated gene

ISG15, an ubiquitin-like protein, is involved in p53 degradation by the 20S proteasome. ISG15

primarily targets misfolded p53 and deletion of ISG15 results in suppression of p53 activity and

functions[24]. Thus, absence of cGAS-STING may lead to p53 destabilization. Third, cGAS-STING

may sensitize cells to DNA damage by lowering the threshold for DDR activation. In absence of

cGAS-STING, the ATM/ATR-p53 pathway may remain ineffective until genome instability is

triggered by telomere-end fusions during crisis. In this scenario, phosphorylation of γ-H2AX could

be achieved through parallel pathways, such as DNA-PKcs.

Absence of cGAS-STING in aging telomerase deficient zebrafish results in low p53 levels,

reduced senescence and increased cell proliferation, thus rescuing damage imposed to

proliferative tissues. This phenotype is also observed in *tp53-/- tert-/-* double mutant zebrafish[26].

Similar to late-generation telomerase deficient mice [33], lack of p53 leads to organismal rescue and

increased fertility allowing for extra generations with ever-shorter telomere mice[34]. As first

observed in tissue culture, upon telomere shortening, the first barrier to cell proliferation (M1) is

imposed by p53/Rb and occurs when telomeres are long enough to allow for further cell divisions

[35]. With loss of p53, the ensuing cell proliferation results in complete telomere deprotection,

genome instability and cell death during crisis (M2)[31]. In this context, loss of cGAS-STING

response is consistent with loss of p53 in aging *tert-/-* mutants. However, *tert-/- sting-/-* do not

completely phenocopy *tert-/- tp53-/-* mutants. Loss of cGAS-STING does not cause an increase in

spontaneous tumor incidence of either WT or *tert-/-* zebrafish. Zebrafish lacking p53, die

prematurely primarily from increased soft tissue tumors[36]. In contrast, *tert-/- sting-/-* zebrafish

lack elevated tumorigenesis characteristic of *tert-/- tp53-/-* mutants[26]. This may be attributed to
the downstream consequences of cGAS-STING and type I interferon response. Chronic
inflammation may be a key component of increased tumorigenesis in *tp53-/-* mutants. Lack of
inflammatory responses may protect *tert-/- sting-/-* zebrafish from early tumorigenesis in face of
increasing DNA damage.

We found that inhibiting cGAS-STING would restore tissue integrity and reduce
expression of YAP-TAZ target genes in aging *tert-/-* mutants[8]. The YAP-TAZ pathway was
recently shown to regulate cGAS-STING in stromal and contractile cells of aging mice[37].
Mechanotransduction by YAP-TAZ suppresses the activity of cGAS-STING, preventing
senescence *in vivo* and tissue degeneration of prematurely aging mouse models [37]. Our work now
shows that cGAS-STING is also required for YAP-TAZ activity upon telomere shortening. This is
likely to be the result of an indirect effect on tissue architecture. Restoring tissue integrity would
reduce the activation of YAP-TAZ mechanosignaling and modifications to the extracellular
matrix.

Loss of cGAS-STING and type I interferon improved the healthspan and the lifespan of
aging *tert-/-* mutants. Telomere shortening does not occur simultaneously in all tissues in humans
and zebrafish[5,38]. The gut of aging zebrafish presents early dysfunction and telomere shortening.
We have recently shown that expressing telomerase specifically in the gut of *tert-/-* mutants
prevents telomere shortening and gut dysfunction[8]. More importantly, maintaining telomere
length in the gut also reverses remote organ dysfunction and longevity of the entire organism.
Like *tert-/- sting-/-* zebrafish, gut-specific telomerase expression in *tert-/-* mutants reduces p53
levels and cell senescence in proliferative organs, namely testis and kidney marrow, despite short
telomeres and increased levels of γ-H2AX. These lead to increased cell proliferation and restored
tissue integrity. We propose that cGAS-STING and type I interferon responses initiated by
telomere shortening in specific organs of aging individuals result in systemic chronic
inflammation (inflammaging) deteriorating tissue integrity of remote organs by increasing DNA
damage and reducing cell proliferation. Thus, inhibition of STING and chronic inflammation in
organs primarily affected by telomere shortening, such as the gut and blood, would increase
healthspan and lifespan. This provides a new approach for treatments of telomere biology
disorders and to improve healthy aging.

Materials and Methods:

Ethics statement
The zebrafish work was conducted according to local and international institutional guidelines and was approved in Portugal by the Ethics Committee of the Instituto Gulbenkian de Ciência and approved by the competent Portuguese authority (Direcção Geral de Alimentação e Veterinária; approval no. 0421/000/000/2015) and in France by the Animal Care Committee of the Institute for Research on Cancer and Aging, Nice, the regional (CIEPAL Côte d'Azur no. 697) and national (French Ministry of Research no. 27673-2020092817202619) authorities.

Zebrafish lines and maintenance
Zebrafish were maintained in accordance with Institutional and National animal care protocols. To ensure telomere length comparisons and avoid the effects of haploinsufficiency of tert+/- heterozygous parental fish, we maintained double heterozygous stock lines (tert$^{AB/hu340}$ sting$^{AB/sa35631}$) as outcrosses to WT AB zebrafish. Experimental fish were obtained incrossing the stock fish. The overall characterization of these four genotypes was performed in F1 sibling animals at 9 months of age. Due to male sex bias in our crosses, that affected mostly tert-/- progeny, we were unable to obtain significant numbers of females for analysis and so all of our data except survival analysis are restricted to males.

Telomere restriction fragment (TRF) analysis by Southern blot
Isolated tissues were first lysed at 50°C overnight in lysis buffer (Fermentas #K0512) supplemented with 1mg/ml Proteinase K (Sigma Aldrich) and RNase A (1:100 dilution, Sigma Aldrich). Genomic DNA was then extracted by equilibrated phenol-chloroform (Sigma Aldrich) and chloroform-isoamyl alcohol extraction (Sigma Aldrich). The same amount of gDNA was digested with RSAI and HINFI enzymes (NEB) for 12 h at 37°C. After digestion, samples were loaded on a 0.6% agarose gel, in 0.5% TBE buffer, and run on an electrophoresis apparatus (Bio-Rad). The electrophoresis conditions were 110 V for 15 h. Gels were then processed for Southern blotting using a 1.6 kb telomere probe, (TTAGGG)n, labeled with [alpha-32P]-dCTP.

Fibroblast Derival
9-month-old fish zebrafish were sacrificed in 1g/L of MS-222 (Sigma Aldrich), and the skin was collected in PBS. After 3 washes in PBS + 2% antibiotics (Penicillin/Streptomycin, Gentamycin and Amphotericin b), the skin was dissociated for 5 minutes in Tryple (Gibco), cut into small pieces, and let to adhere O/N on coverslip coated by gelatin 2% in presence of few drops of FBS + 2% antibiotics. The day after, when the first fibroblasts were released from the skin, the well was filled with DMEM + 2% antibiotics. After 48h, fibroblasts were fixed and DAPI staining was performed.

Western Blot
Age and sex matched adult zebrafish were sacrificed in 1g/L of MS-222 (Sigma Aldrich) and collected tissues (skin, testis, kidney marrow and intestine) were immediately snap frozen liquid nitrogen. Tissues were homogenized in RIPA buffer (sodium chloride 150 mM; Triton-X-100 1%, sodium deoxycholate 0.5%, SDS 0.1%, Tris 50 mM, pH=8.0), including complete protease and phosphatase inhibitor cocktails (Roche diagnostics) with a motor pestle on ice. Homogenized tissues were incubated for 30 minutes on ice and centrifuged at 4°C, 13.000 rpm for 10 min. Supernatant was collected and stored at-80°C until use.
For each sample, 50 µg of protein was loaded per well, separated on 10% SDS-PAGE gels and transferred to Nitrocellulose Membrane (BioRad #1620097). The membranes were blocked in 5%

milk and then incubated with the primary antibody overnight at 4°C. Antibody complexes were visualized by enhanced chemiluminescence (ECL) after incubation with the appropriate HRP-conjugated secondary antibody. Antibodies concentrations: anti-p53 (1:1000, Anaspec, 55342), anti TBK1 (1:1000, CST, 3504), anti pTBK1 (1:1000, CST, 5483), anti IRF3 (1:1000, CST, 11904), anti pIRF3 (1:1000, CST, 29047).

Real-time quantitative PCR

Age and sex matched adult zebrafish were sacrificed in 1g/L of MS-222 (Sigma Aldrich) and collected tissues (skin, testis, kidney marrow and intestine) were immediately snap frozen liquid nitrogen. RNA extractions were performed in TRIzol (Invitrogen) by mashing tissues with a motorized pestle in a 1.5 mL eppendorf tube. After incubation at room temperature (RT) for 10 min TRIzol, chloroform extractions were performed. Quality of RNA samples was assessed through BioAnalyzer (Agilent 2100). Retro-transcription into cDNA was performed using QuantiTect Reverse Transcription kit (Qiagen).

Quantitative PCR (qPCR) was performed using FastStart Universal SYBR Green Master mix (Roche) and an StepOne+ Real time PCR Detection System (Applied Biosystems). qPCRs were carried out in duplicate for each cDNA sample. Relative mRNA expression was normalized against *rps11* mRNA expression using the 2^{-dCT} method.

	Gene ID:	Forward primer	Reverse primer
isg15	ZDB-GENE-021211-1	ACTCGGTGACGATGCAGC	TGGGCACGTTGAAGTACTGA
ifn-i	ZDB-GENE-030721-3	CAAGATACGCAAAGCCAGCA	GTGGCTTTTCACAACTCTCC
cdkn2a/b	ZDB-GENE-081104-306	GAGGATGAACTGACCACAGCA	CAAGAGCCAAAGGTGCGTTAC
cdkn1a	ZDB-GENE-070705-7	CAGCGGGTTTACAGTTTCAGC	TGAACGTAGGATCCGCTTGT
il1b	ZDB-GENE-040702-2	CGCTCCACATCTCGTACTCA	ATACGCGGTGCTGATAAACC
tgfb1b	ZDB-GENE-091028-1	ACCCCAGTTCAGCACACCATAG	TCGAAACTCGGCCTGGTAGA
mmp15a	ZDB-GENE-070817-4	GGGTCATGCTCTGGGGGTTGG	AGTGGTGACAGTCTCTGGAGATCCA
ctgf	ZDB-GENE-030131-102	ACTCCCCTCGTCAAAACACC	GGGACCGTATGTCTCCTCCT
cyr61	ZDB-GENE-040426-3	CCGTGTCCACATGTACATGGG	GGTGCATGAAAGAAGCTCGTC
mavs	ZDB-GENE-070112-1402	AGTAGAAGCCGCGAGAGGTA	GGCTTCGATCTCTTCCCGAT
rps11	ZDB-GENE-040426-2701	ACAGAAATGCCCCTTCACTG	GCCTCTTCTCAAAACGGTTG

Histology

Age and sex-matched adult zebrafish were sacrificed in 1g/L of MS-222 (Sigma Aldrich), fixed for 72 hours in 4% paraformaldehyde and decalcified in 0.5M EDTA for 48 hours at room temperature. Whole fish were then paraffin-embedded to perform five micrometer sagittal section slides. Slides were stained with hematoxylin (Sigma Aldrich) and eosin (Sigma Aldrich) for histopathological analysis. Microphotographs (N>=6 fish per genotype) were acquired in a Leica DM4000 B microscope coupled to a Leica DFC425 C camera (Leica Microsystems).

Immunofluorescence

Deparaffinized and rehydrated slides were microwaved 20 min at 550W in citrate buffer (10 mM
Sodium Citrate, pH 6.0) for antigen retrieval. Slides were washed twice in PBS for 5 minutes and
blocked for 1 hour at RT in 0.5% Triton, 1% DMSO, 5% normal goat serum in PBS (blocking
solution). Subsequently, slides were incubated overnight at 4°C with 1:50 dilution of primary
antibody in blocking solution. The following primary antibodies were used: mouse monoclonal
antibody against Proliferation Cell Nuclear Antigen (PCNA), sc56, (Santa Cruz), rabbit polyclonal
Histone H2A.XS139ph (phospho Ser139) GTX127342 (Genetex). Following two PBS washes,
overnight incubation at 4°C was performed 1:500 dilution of goat anti-rabbit secondary antibody
AlexaFluor 488 (Invitrogen) and goat anti-mouse secondary antibody AlexaFluor 488 (Invitrogen).
Finally, after DAPI staining (Sigma Aldrich), slides were mounted DAKO Fluorescence Mounting
Medium (Sigma Aldrich).
Immunofluorescence images were acquired on Delta Vision Elite (GE Healthcare) using an
OLYMPUS 60x/1.42 objective. For quantitative and comparative imaging, equivalent acquisition
parameters were used. The percentage of positive nuclei was determined by counting a total of
150-1000 cells per slide depending on the tissue (N>= 6 zebrafish per genotype).
445
Senescence Associated Beta Galactosidase Staining
Age and sex matched 9-month-old zebrafish were sacrificed in 1g/l of MS-222 (Sigma Aldrich,
MO, USA), fixed in 4% paraformaldehyde in PBS for 72 hour at 4°C, washed 3 times for 1 hour in
1x PBS pH 7.4 and 1 hour in 1x PBS pH 6.0 at 4°C. Beta Galactosidase staining was performed for
24 hours at 37°C in 5mM Potassium Ferrocyanide, 5 mM Potassium Ferricyanide, 2 mM MgCl2 and
1 mg/mL X-Gal, in 1x PBS pH 6.0. After staining, fish were washed 3 times for 5 minutes in 1x PBS
pH 7.0, and processed for decalcification and paraffin embedding. Paraffin blocks were sectioned
sagittally five micrometers in thickness and co-stained with nuclear fast red (Sigma Aldrich)
454
Fertility assays
In order to assess male fertility, 9-month-old males from the four different genotypes were
separately housed overnight in external breeding tanks with a single young (3-6 months old) WT
female. Breeding pairs were left to cross and lay eggs in the following morning and embryos were
collected approximately 4 hours post fertilization (hpf) and allowed to develop at 28°C.
Assessment of fertilized eggs and embryo viability was conducted between 4 and 6hpf. At least
12 independent crosses were conducted for each genotype to evaluate male fertility. Only
successful breeding trials, defined as events in which a clutch of eggs was laid by a female, were
scored.
464
Fixation for histology and tumor evaluation
Fish were screened weekly for the presence of macroscopic tumors [36]. Zebrafish were euthanized
with 1g/L of MS-222 (Sigma, MO, USA), followed by fixation in 10% neutral buffered formalin for
72 hr and decalcified in 0.5M EDTA for 48 hr at room temperature. Samples were then paraffin-
embedded to perform 5 mm sagittal section slides. Slides were stained with hematoxylin and
eosin and assessed for the presence of tumors.
471
RNA-seq analysis of TE families.
Raw FASTQ read files obtained from the SRA (PRJNA937311) were aligned to the reference
zebrafish genome (GRCz10/danRer10) using STAR v2.7.10a with the following parameters to
retain multi-mapping reads: --outMultimapperOrder Random --outSAMmultNmax 1 --
outFilterMismatchNmax 3 --winAnchorMultimapNmax 100 --outFilterMultimapNmax 100 --
alignSJDBoverhangMin 1. Expression of transposable element families was quantified using
TEtransctipts v2.2.393 with parameter –mode=multi to estimate transposable element

abundances from multimapped alignments using the pre-generated GENCODE danRer10 TE GTF
file from the Hammel lab FTP site (https://labshare.cshl.edu/shares/mhammelllab/www-
data/TEtranscripts/TE_GTF). Counts were normalized to counts per million (CPM) and differential
expression was assessed in R v3.1.4 (R Core Team, 2022) on the combined transposable element
/gene counts using the edgeR package v3.34.189 exact test.

Statistical Analysis
Graphs and statistical analyses were performed in GraphPad Prism8 software, using one-way
ANOVA test Tukey's post-correction or unpaired t-test. A critical value for significance of $p<0.05$
was used throughout the study. For survival analysis, Log-rank tests were performed using
GraphPad Prism8 to determine statistical differences of survival curves.

Data availability
All data generated or analyzed during this study are included in this published article and its
Extended information files.

Acknowledgments
We are grateful to our team members for fruitful discussions and advice. Especially, we thank Dr.
Hervé Técher for critically reading the manuscript. This work was supported by the Université
Côte d'Azur - Académie 4 (Installation Grant: Action 2 - 2019) and Institut National du Cancer
(INCa, PLBIO21-228). N.S. was supported by a PhD fellowship by La Ligue Contre le Cancer. E.T.
and P.B. thank the German Research Foundation, grant nos. 322977937/GRK2344. We are
grateful to the PEMAV fish facility, Imaging core facility (PICMI) and the Genomics facilities at the
IRCAN supported by FEDER, Région Provence Alpes-Côte d'Azur, Conseil Départemental 06,
ITMO Cancer Aviesan (plan cancer), Cancéropole Provence Alpes-Côte d'Azur, Gis Ibisa, CNRS
and Inserm. The funders had no role in study design, data collection and analysis, decision to
publish or preparation of the manuscript.

Author contributions
N.S. performed most experiments and carried out data analyses. G.A. performed data analysis
and revised the manuscript. B.L.-B. performed the skin fibroblast cell derivation and micronuclei
quantifications. M.M. proposed the idea for the study and performed the initial analysis of the
sting-/- tert-/- double mutants. RNA-seq analysis of transposon elements was performed by P.B.
and supervised by E.T. N.S. and M.G.F. designed the experiments and wrote the manuscript.
M.G.F. conceived the study, acquired funding and supervised the work.

Declaration of competing interests
The authors declare no competing interests.

References

1. Armanios, M. Syndromes of telomere shortening. *Annu Rev Genomics Hum Genet* **10**, 45–61 (2009).

2. Mitchell, J. R., Wood, E. & Collins, K. A telomerase component is defective in the human disease dyskeratosis congenita. *Nature* **402**, 551–5 (1999).

3. Opresko, P. L. & Shay, J. W. Telomere-associated aging disorders. *Ageing Res Rev* **33**, 52–66 (2017).

4. Anchelin, M. *et al.* Premature aging in telomerase-deficient zebrafish. *Dis Model Mech* (2013) doi:10.1242/dmm.011635.

5. Carneiro, M. C. *et al.* Short Telomeres in Key Tissues Initiate Local and Systemic Aging in Zebrafish. *PLoS Genet* **12**, e1005798 (2016).

6. Henriques, C. M., Carneiro, M. C., Tenente, I. M., Jacinto, A. & Ferreira, M. G. Telomerase is required for zebrafish lifespan. *PLoS Genet* **9**, e1003214 (2013).

7. Lex, K. *et al.* Telomere shortening produces an inflammatory environment that increases tumor incidence in zebrafish. *Proc Natl Acad Sci U S A* **117**, 15066–15074 (2020).

8. El Maï, M. *et al.* Gut-specific telomerase expression counteracts systemic aging in telomerase-deficient zebrafish. *Nat Aging* **3**, 567–584 (2023).

9. Oh, H. S.-H. *et al.* Organ aging signatures in the plasma proteome track health and disease. *Nature* **624**, 164–172 (2023).

10. Glück, S. *et al.* Innate immune sensing of cytosolic chromatin fragments through cGAS promotes senescence. *Nat Cell Biol* **19**, 1061–1070 (2017).

11. Yang, H., Wang, H., Ren, J., Chen, Q. & Chen, Z. J. cGAS is essential for cellular senescence. *Proc Natl Acad Sci U S A* **114**, E4612–E4620 (2017).

12. Dou, Z. *et al.* Cytoplasmic chromatin triggers inflammation in senescence and cancer. *Nature* **550**, 402–406 (2017).

13. Nassour, J., Przetocka, S. & Karlseder, J. Telomeres as hotspots for innate immunity and inflammation. *DNA Repair (Amst)* **133**, 103591 (2024).

14. Motwani, M., Pesiridis, S. & Fitzgerald, K. A. DNA sensing by the cGAS–STING pathway in health and disease. *Nat Rev Genet* **20**, 657–674 (2019).

15. Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z. J. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* **339**, 786–91 (2013).

16. Ishikawa, H. & Barber, G. N. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature* **455**, 674–8 (2008).

17. Zierhut, C. Potential cGAS-STING pathway functions in DNA damage responses, DNA replication and DNA repair. *DNA Repair (Amst)* **133**, 103608 (2024).

18. Glück, S. & Ablasser, A. Innate immunosensing of DNA in cellular senescence. *Curr Opin Immunol* **56**, 31–36 (2019).

19. Gulen, M. F. *et al.* cGAS-STING drives ageing-related inflammation and neurodegeneration. *Nature* **620**, 374–380 (2023).

20. Carneiro Madalena. The role of short telomeres as cause of natural aging in zebrafish. (Universidade Nova de Lisboa, Lisboa, Portugal, 2015).

21. El Maï, M., Marzullo, M., de Castro, I. P. & Ferreira, M. G. Opposing p53 and mTOR/AKT promote an in vivo switch from apoptosis to senescence upon telomere shortening in zebrafish. *Elife* **9**, (2020).

22. De Cecco, M. *et al.* L1 drives IFN in senescent cells and promotes age-associated inflammation. *Nature* **566**, 73–78 (2019).

23. Nassour, J. *et al.* Telomere-to-mitochondria signalling by ZBP1 mediates replicative crisis. *Nature* **614**, 767–773 (2023).

24. Huang, Y.-F., Wee, S., Gunaratne, J., Lane, D. P. & Bulavin, D. V. Isg15 controls p53 stability and functions. *Cell Cycle* **13**, 2200–10 (2014).

25. Takaoka, A. *et al.* Integration of interferon-alpha/beta signalling to p53 responses in tumour suppression and antiviral defence. *Nature* **424**, 516–23 (2003).

572   26.   Şerifoğlu, N., Lopes-Bastos, B. & Ferreira, M. G. Lack of telomerase reduces cancer
573   incidence and increases lifespan of zebrafish tp53M214K mutants. *Sci Rep* **14**, 5382 (2024).
574   27.   Spitsbergen, J. M., Buhler, D. R. & Peterson, T. S. Neoplasia and neoplasm-associated
575   lesions in laboratory colonies of zebrafish emphasizing key influences of diet and
576   aquaculture system design. *ILAR J* **53**, 114–25 (2012).
577   28.   López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging:
578   An expanding universe. *Cell* **186**, 243–278 (2023).
579   29.   Franceschi, C., Garagnani, P., Parini, P., Giuliani, C. & Santoro, A. Inflammaging: a new
580   immune-metabolic viewpoint for age-related diseases. *Nat Rev Endocrinol* **14**, 576–590
581   (2018).
582   30.   Li, X. *et al.* Inflammation and aging: signaling pathways and intervention therapies. *Signal*
583   *Transduct Target Ther* **8**, 239 (2023).
584   31.   Nassour, J. *et al.* Autophagic cell death restricts chromosomal instability during replicative
585   crisis. *Nature* **565**, 659–663 (2019).
586   32.   Sato, Y. & Hayashi, M. T. Micronucleus is not a potent inducer of the cGAS/STING
587   pathway. *Life Sci Alliance* **7**, (2024).
588   33.   Rudolph, K. L. *et al.* Longevity, Stress Response, and Cancer in Aging Telomerase-Deficient
589   Mice. *Cell* **96**, 701–712 (1999).
590   34.   Chin, L. *et al.* p53 deficiency rescues the adverse effects of telomere loss and cooperates
591   with telomere dysfunction to accelerate carcinogenesis. *Cell* **97**, 527–38 (1999).
592   35.   Saretzki, G., Sitte, N., Merkel, U., Wurm, R. E. & von Zglinicki, T. Telomere shortening
593   triggers a p53-dependent cell cycle arrest via accumulation of G-rich single stranded DNA
594   fragments. *Oncogene* **18**, 5148–58 (1999).
595   36.   Berghmans, S. *et al.* tp53 mutant zebrafish develop malignant peripheral nerve sheath
596   tumors. *Proc Natl Acad Sci U S A* **102**, 407–12 (2005).
597   37.   Sladitschek-Martens, H. L. *et al.* YAP/TAZ activity in stromal cells prevents ageing by
598   controlling cGAS-STING. *Nature* **607**, 790–798 (2022).
599   38.   Aubert, G., Baerlocher, G. M., Vulto, I., Poon, S. S. & Lansdorp, P. M. Collapse of telomere
600   homeostasis in hematopoietic cells caused by heterozygous mutations in telomerase genes.
601   *PLoS Genet* **8**, e1002696 (2012).
602
603

Figure Legends

Figure 1: Telomere shortening activates cGAS-STING pathway. a, quantification of mean telomere length measured by TRF analysis in the skin (n_{WT}= 5, $n_{tert-/-}$=5, $n_{sting-/-}$=5, $_{tert-/-\ sting-/-}$=4, WT *vs tert-/-* p= 0.017,tert-/- *vs sting-/-* p= 0.002, *sting-/- vs tert-/- sting-/-* p=0.008). **b,** quantification of mean telomere length measured by TRF analysis in the testis (n_{WT}= 4, $n_{tert-/-}$=6, $n_{sting-/-}$=5, $_{tert-/-\ sting-/-}$=6, WT *vs tert-/-* p=0.001, WT *vs tert-/- sting-/-* p<0.001, *tert-/- vs sting-/-* p= 0.016, *sting-/- vs tert-/- sting-/-* p=0.001). **c,** quantification of mean telomere length measured by TRF analysis in the kidney marrow (n_{WT}= 3, $n_{tert-/-}$=4, $n_{sting-/-}$=7, $_{tert-/-\ sting-/-}$=7, WT *vs tert-/-* p< 0.001, WT *vs tert-/- sting-/-* p= 0.001, *tert-/- vs sting-/-* p< 0.001, *sting-/- vs tert-/- sting-/-* p<0.001, *tert-/- vs tert-/- sting-/-* p=0.003). **d,** quantification of mean telomere length measured by TRF analysis in the intestine (n_{WT}= 6, $n_{tert-/-}$=6, $n_{sting-/-}$=6, $_{tert-/-\ sting-/-}$=6, WT vs *tert-/-* p= 0.001, WT vs *tert-/- sting-/-* p=0.015, *tert-/- vs sting-/-* p<0.001, *sting-/- vs tert-/- sting-/-* p=0.008, p= 0.623). **e,** representative immunofluorescence images and quantifications of MN formation in the fibroblasts derived from skin (n_{WT}= 1, $n_{tert-/-}$=1, $n_{sting-/-}$=1, $_{tert-/-\ sting-/-}$=1, WT vs *tert-/-* p= 0.040 , WT vs *tert-/- sting-/-* p=0.034, *tert-/-* vs *sting-/-* p= 0.047, *sting-/-* vs *tert-/- sting-/-* p=0.040). **f,** representative western blot images and quantification of downstream targets of cGAS-STING pathway (n_{WT}= 4-5, $n_{tert-/-}$=5-6, $n_{sting-/-}$=5-7, $_{tert-/-\ sting-/-}$=5, p-Irf3/Irf3: WT vs *tert-/-* p= 0.049, *tert-/-* vs *tert-/- sting-/-* p= 0.011; p-Tbk1/Tbk1: WT vs *tert-/-* p= 0.035, *tert-/-* vs *sting-/-* p= 0.002, *tert-/-* vs *tert-/- sting-/-* p= 0.042). **g,** RT-qPCR analysis of *isg15* gene expression in the skin, testis, kidney marrow and intestine (n_{WT}= 5-6, $n_{tert-/-}$=4-8, $n_{sting-/-}$=5-8, $_{tert-/-\ sting-/-}$=7-9, skin: WT vs *tert-/-* p= 0.024, *sting-/-* vs *tert-/-* p=0.011, N=5-7, *tert-/-* vs *tert-/- sting-/-* p=0.007; testis: WT vs *tert-/-* p=0.001, *sting-/-* vs *tert-/-* p=0.001, *tert-/-* vs *tert-/- sting-/-* p=0.002; kidney marrow: N=6-9, WT vs *tert-/-* p=0.045, *sting-/-* vs *tert-/-* p=0.022, *tert-/-* vs *tert-/- sting-/-* p= 0.019; intestine: WT vs *tert-/-* p<0.001, *sting-/-* vs *tert-/-* p<0.001, *tert-/-* vs *tert-/- sting-/-* p=0.001). **h,** RT-qPCR analysis of *ifn-i* gene expression in the skin, testis, kidney marrow and intestine (n_{WT}= 4-6, $n_{tert-/-}$=5-8, $n_{sting-/-}$=5-8, $_{tert-/-\ sting-/-}$=6-9, skin: *sting-/-* vs *tert-/-* p=0.004, *tert-/-* vs *tert-/- sting-/-* p=0.002; testis: WT vs *tert-/-* p=0.003, *sting-/-* vs *tert-/-* p=0.003, *tert-/-* vs *tert-/- sting-/-* p=0.003; kidney marrow: WT vs *tert-/-* p=0.008, *sting-/-* vs *tert-/-* p=0.004, *tert-/-* vs *tert-/- sting-/-* p= 0.011; intestine: WT vs *tert-/-* p=0.020, *sting-/-* vs *tert-/-* p=0.005, *tert-/-* vs *tert-/- sting-/-* p=0.014). Data are presented as the mean ± s.e.m.; *p<0.05; **p<0.01, ***p<0.001, using a one-way ANOVA and post hoc Tukey test.

Figure 2: cGAS-STING pathway inactivation attenuates DNA damage response. a, representative immunofluorescence images of DNA damage. **b,** quantification of DNA damage in skin (n_{WT}= 6, $n_{tert-/-}$=6, $n_{sting-/-}$=6, $_{tert-/-\ sting-/-}$=6; WT vs tert-/- p< 0.001, sting-/- vs tert-/- p< 0.001, *WT vs tert-/- sting-/- p< 0.001, sting-/- vs tert-/- sting-/- p< 0.001*). **c,** quantification of DNA damage in testis (n_{WT}= 6, $n_{tert-/-}$=6, $n_{sting-/-}$=6, $_{tert-/-\ sting-/-}$=6, N= 5-6, WT vs *tert-/-* p< 0.001, *sting-/-* vs tert-/- p< 0.001, WT vs *tert-/- sting-/-* p< 0.001, *sting-/-* vs tert-/- sting-/- p< 0.001), **d,** quantification of DNA damage in kidney marrow (n_{WT}= 6, $n_{tert-/-}$=6, $n_{sting-/-}$=6, $_{tert-/-\ sting-/-}$=6; WT vs *tert-/-* p< 0.001, *sting-/-* vs tert-/- p< 0.001, WT vs *tert-/- sting-/-* p< 0.001, *sting-/-* vs tert-/- sting-/- p<0.001). **e,** quantification of DNA damage in intestine (n_{WT}= 5, $n_{tert-/-}$=5, $n_{sting-/-}$=5, $_{tert-/-\ sting-/-}$=5; WT vs *tert-/-* p=0.006, sting-/- vs tert-/- p=0.008, WT vs tert-/- sting-/- p= 0.007, sting-/- vs tert-/- sting-/- p<0.001). **f,** representative western blot images of p53. **g,** quantification of p53 levels in the skin (n_{WT}= 4, $n_{tert-/-}$=4, $n_{sting-/-}$=4, $_{tert-/-\ sting-/-}$=4, WT vs *tert-/-* p< 0.001, *sting-/-* vs tert-/- p< 0.001, tert-/- vs *tert-/- sting-/-* p< 0.001). **h,** quantification of p53 levels in the testis (n_{WT}= 4, $n_{tert-/-}$=4, $n_{sting-/-}$=4, $_{tert-/-\ sting-/-}$=4, WT vs *tert-/-* p=0.050, *tert-/-* vs tert-/- sting-/- p=0.048). **i,** quantification of p53 levels in the kidney marrow (n_{WT}= 4, $n_{tert-/-}$=4, $n_{sting-/-}$=4, $_{tert-/-\ sting-/-}$=4, *sting-/-* vs *tert-/-* p= 0.017, *tert-/-* vs tert-/- sting-/- p= 0.022). **j,** quantification of p53 levels in the intestine (n_{WT}= 5, $n_{tert-/-}$=5, $n_{sting-/-}$=5, $_{tert-/-\ sting-/-}$=5, WT vs *tert-/-* p<0.001, *sting-/-* vs tert-/- p=0.006, *tert-/-* vs *tert-/- sting-/-* p=0.006). Data are presented as the mean ± s.e.m.; *p<0.05; **p<0.01, ***p<0.001, using a one-way ANOVA and post hoc Tukey test.

Figure 3: SASP induced by short telomeres are controlled by cGAS-STING pathway. a, representative images of senescence associated beta galactosidase staining in skin, testis, kidney marrow and intestine (n_{WT}= 3, $n_{tert-/-}$=3, $n_{sting-/-}$=3, $_{tert-/-\ sting-/-}$=3). **b,** RT-qPCR analysis of inflammatory markers and SASP factors in the skin (n_{WT}= 4-8, $n_{tert-/-}$=5-8, $n_{sting-/-}$=6-7, $_{tert-/-\ sting-/-}$=5-7; *cdkn2a/b:* WT vs *tert-/-* p=0.0001, *sting-/-* vs *tert-/-* p<0.0001, *tert-/-* vs tert-/- sting -/- p=0.004; *cdnk1a:* WT vs *tert-/-* p=0.025, *sting-/-* vs tert-/- p=0.014, *tert-/-* vs *tert-/- sting-/-* p=0.013; *il1b:* WT vs *tert-/-* p=0.023, *sting-/-* vs tert-/- p=0.031, *tert-/-* vs *tert-/- sting-/-* p=0.004,; *tgf1b: sting-/-* vs *tert-/-* p=0.014, *tert-/-* vs *tert-/- sting-/-* p=0.017; *mmp15a: sting-/-* vs *tert-/-*

p=0.011, tert-/- vs tert-/- sting-/- p=0.009). **c**, RT-qPCR analysis of inflammatory markers and SASP factors
in the testis (n_{WT}= 4-7, $n_{tert-/-}$=6-11, $n_{sting-/-}$=5-7, *tert-/- sting-/-*=5-8; *cdkn2a/b:* WT vs *tert-/- p=0.0234; cdnk1a:* WT vs
tert-/- p=0.013, tert-/- vs tert-/- sting-/- p=0.018; il1b: tert-/- vs tert-/- sting-/- p=0.030; tgf1b: WT vs *tert-/-*
p=0.029, sting-/- vs tert-/- p=0.020, tert-/- vs tert-/- sting-/- p=0.043; mmp15a: WT vs *tert-/- p=0.004, sting-/-*
vs *tert-/- p=0.016, tert-/- vs tert-/- sting-/- p=0.028*). **d**, RT-qPCR analysis of inflammatory markers and SASP
factors in the kidney marrow (n_{WT}= 5-9, $n_{tert-/-}$=5-11, $n_{sting-/-}$=6-11, *tert-/- sting-/-*=6-10; *cdkn2a/b:* WT vs *tert-/-*
p=0.035, sting-/- vs tert-/- p=0.009, tert-/- vs tert-/- sting-/- p<0.001; cdnk1a: WT vs *tert-/- p=0.002, sting-/- vs*
tert-/- p=0.001, tert-/- vs tert-/- sting-/- p=0.005; il1b: WT vs *tert-/- p=0.047, sting-/- vs tert-/- p=0.003, tert-/-*
vs *tert-/- sting-/- p=0.014; tgf1b: sting-/- vs tert-/- p=0.005; mmp15a:* WT vs *tert-/- p=0.009, sting-/- vs tert-/-*
p=0.011, tert-/- vs tert-/- sting-/- p=0.008). **e**, RT-qPCR analysis of inflammatory markers and SASP factors
in the intestine (n_{WT}= 5-7, $n_{tert-/-}$=5-8, $n_{sting-/-}$=5-6, *tert-/- sting-/-*=5-7; *cdkn2a/b:* WT vs *tert-/- p<0.001, sting-/- vs*
tert-/- p=0.001, tert-/- vs tert-/- sting-/- p=0.038; cdnk1a: WT vs *tert-/- p=0.007, sting-/- vs tert-/- p<0.001,*
tert-/- vs tert-/- sting-/- p=0.011; il1b: WT vs *tert-/- p=0.004, sting-/- vs tert-/- p=0.012, tert-/- vs tert-/- sting-/-*
p=0.009; tgf1b: WT vs *tert-/- p=0.034, sting-/- vs tert-/- p=0.003; mmp15a:* WT vs *tert-/- NS p=0.060, sting-/-*
vs *tert-/- p=0.021, tert-/- vs tert-/- sting-/- p=0.070*). All data are presented as the mean ± s.e.m.; *p<0.05;
**p<0.01, ***p<0.001, using a one-way ANOVA and post hoc Tukey test

Figure 4: cGAS-STING pathway control proliferation in telomeric dysfunction. **a**, representative
immunofluorescence images of proliferation and apoptosis. **b**, quantification of proliferation in the skin
(n_{WT}= 5, $n_{tert-/-}$=6, $n_{sting-/-}$=6, *tert-/- sting-/-*=6 WT vs *tert-/- p<0.001, sting-/- vs tert-/- p<0.001, tert-/- vs tert-/- sting-*
/- p=0.024). **c**, quantification of proliferation in the testis (n_{WT}= 6, $n_{tert-/-}$=6, $n_{sting-/-}$=6, *tert-/- sting-/-*=6, WT vs *tert-*
/- p<0.001, sting-/- vs tert-/- p<0.001, tert-/- vs tert-/- sting-/- p<0.001). **d**, quantification of proliferation in
the kidney marrow (n_{WT}= 8, $n_{tert-/-}$=8, $n_{sting-/-}$=6, *tert-/- sting-/-*=8, WT vs *tert-/- p=0.002, sting-/- vs tert-/- p=0.004*).
e, quantification of proliferation in the intestine (n_{WT}= 6, $n_{tert-/-}$=5, $n_{sting-/-}$=6, *tert-/- sting-/-*=8). **f**, representative
hematoxylin eosin staining of intestine, insets with yellow lines representative of lamina propria thickness
g, quantification of lamina propria width (n_{WT}= 4, $n_{tert-/-}$=4, $n_{sting-/-}$=4, *tert-/- sting-/-*=4, WT vs *tert-/- p=0.042,*
sting-/- vs tert-/- p=0.048). **h**, representative hematoxylin eosin staining of testis. **i**, quantification of mature
sperm area (n_{WT}= 4, $n_{tert-/-}$=4, $n_{sting-/-}$=4, *tert-/- sting-/-*=4, WT vs *tert-/- p<0.002, sting-/- vs tert-/- p<0.001, tert-/- vs*
tert-/- sting-/- p= 0.006). **l**, RT-qPCR analysis of YAP-TAZ pathway targets in the intestine (n_{WT}= 4, $n_{tert-/-}$=4,
$n_{sting-/-}$=5, *tert-/- sting-/-*=5, ctfg: WT vs *tert-/- p=0.016 sting-/- vs tert-/- p=0.005, tert-/- vs tert-/- sting-/- p=0.044;*
cyr61: WT vs *tert-/- p=0.030, sting-/- vs tert-/- p=0,023, tert-/- vs tert-/- sting-/- p=0.046*). **m**, RT-qPCR
analysis of YAP-TAZ pathway targets in the testis (n_{WT}= 7, $n_{tert-/-}$=8, $n_{sting-/-}$=6-7, *tert-/- sting-/-*=7-8, cftg: *tert-/- vs*
tert-/- sting-/- p=0.008, cyr61: tert-/- vs tert-/- sting-/- p=0.011). All data are presented as the mean ± s.e.m.;
*p<0.05; **p<0.01, ***p<0.001, using a one-way ANOVA and post hoc Tukey test.

Figure 5: cGAS-STING pathway rescues premature aging phenotypes. a, quantification of male fertile
capacity (n_{WT}= 15, $n_{tert-/-}$=10, $n_{sting-/-}$=14, *tert-/- sting-/-*=13), WT vs *tert-/- p<0.001, sting-/- vs tert-/- p<0.001,* WT vs
tert-/- sting-/- p=0.011, sting-/- vs tert-/- sting-/- p<0.001, tert-/- vs tert-/- sting-/- p=0.002). **b**, representative
images of adult zebrafish and quantification of aging phenotypes scored as kyphosis and cachexia (n_{WT}=
10, $n_{tert-/-}$=12, $n_{sting-/-}$=11, *tert-/- sting-/-*=12, WT vs *tert-/- p<0.001, sting-/- vs tert-/- p<0.001,* WT vs *tert-/- sting-/-*
p<0.001, sting-/- vs tert-/- sting-/- p=0.003, tert-/- vs tert-/- sting-/- p<0.001). **c**, quantification of weight in
adult zebrafish (n_{WT}= 12, $n_{tert-/-}$=13, $n_{sting-/-}$=11, *tert-/- sting-/-*=13, WT vs *tert-/- p<0.001, sting-/- vs tert-/- p=0.004,*
tert-/- vs tert-/- sting-/- p=0.012). **d**, quantification of seminoma (n_{WT}= 17, $n_{tert-/-}$=16, $n_{sting-/-}$=21, *tert-/- sting-/-*=18,
WT vs *tert-/- p=0.012, sting-/- vs tert-/- p=0.002, tert-/- vs tert-/- sting-/- p=0.027*). **e**, quantification of
survival n_{WT}= 24, $n_{tert-/-}$=38, $n_{sting-/-}$=32, *tert-/- sting-/-*=37, WT vs *tert-/- p=0.008, sting-/- vs tert-/- p<0.001, tert-/- vs*
tert-/- sting-/- p=0.010). Data are presented as the mean ± s.e.m. *p<0.05; **p<0.01, ***p<0.001, using a
one-way ANOVA and post hoc Tukey test. Seminoma occurrence and survival data were analyzed using
Log-rank tests, **p<0.01, ***p<0.001.

Figure 2

Figure 3

Figure 4

Figure 5