Title

Neuroprotective role of Hippo signaling by microtubule stability control in *C. elegans*

Short title: Hpo pathway maintains neuronal integrity

Authors

Hanee Lee[1], Junsu Kang[1†], Sang-Hee Lee[1‡], Dowoon Lee[1], Christine H. Chung[1], Junho Lee[1*]

Affiliations

[1] Department of Biological Sciences, Institute of Molecular Biology and Genetics,

Seoul National University, Seoul, 08826, South Korea

* Correspondences to

Junho Lee

Department of Biological Sciences, Seoul National University

Gwanak-ro 1, Gwanak-gu, Seoul, 08826, South Korea

elegans@snu.ac.kr

[†] Present address: Department of Cell and Regenerative Biology, University of Wisconsin
School of Medicine and Public Health, University of Wisconsin–Madison, Madison, WI
53705, USA

[‡] Present address: Korea Basic Science Institute, Ochang, Cheongju, Chungbuk, 28119,
South Korea

Abstract

The evolutionarily conserved Hippo (Hpo) pathway has been shown to impact early development and tumorigenesis by governing cell proliferation and apoptosis. However, its post-developmental roles are relatively unexplored. Here, we demonstrate its roles in post-mitotic cells by showing that defective Hpo signaling accelerates age-associated structural and functional decline of neurons in *C. elegans*. Loss of *wts-1*/LATS, the core kinase of the Hpo pathway, resulted in premature deformation of touch neurons and impaired touch responses in a *yap-1/*YAP-dependent manner, the downstream transcriptional co-activator of LATS. Decreased movement as well as microtubule destabilization by treatment with colchicine or disruption of microtubule stabilizing genes alleviated the neuronal deformation of *wts-1* mutants. Colchicine exerted neuroprotective effects even during normal aging. In addition, the deficiency of a microtubule-severing enzyme *spas-1* also led to precocious structural deformation. These results consistently suggest that hyper-stabilized microtubules in both *wts-1*-deficient neurons and normally aged neurons are detrimental to the maintenance of neuronal structural integrity. In summary, Hpo pathway governs the structural and functional maintenance of differentiated neurons by modulating microtubule stability, raising the possibility that the microtubule stability of fully developed neurons could be a promising target to delay neuronal aging. Our study provides potential therapeutic approaches to combat age- or disease-related neurodegeneration.

Keywords

Introduction

Neurons perceive extracellular signals and transmit the information to other cells, often over

long distances. For this to happen, neuronal cells extend their axons and dendrites from 1 μm

to more than 1 m and must maintain an intact morphology throughout their lifetime. The long,

elaborate neuronal structures are vulnerable to degeneration associated with organismal aging

or pathological conditions. Structural abnormalities and the consequent functional decline are

hallmarks of aged neurons or pathological neurons affected by neurodegenerative diseases

(Yankner et al. 2008; Bishop et al. 2010; Fjell and Walhovd 2010). Several studies have

shown that the aged brain exhibits structural abnormalities such as synaptic loss, neuronal

sprouting and restructuring, rather than neuronal cell death (reviewed in (Yankner et al.

2008)).

Microtubules form essential cytoskeletal structures for most eukaryotic cells. In neurons, they

serve as structural struts that allow the neuron to develop and maintain a specific shape

(Barnes and Polleux 2009; Marin et al. 2010). They also act as railroads for motor proteins

that deliver cellular cargo from cell bodies to distal axons (Hirokawa et al. 2010; Maday et al.

2014), as well as seeds for microtubule polymerization (Job et al. 2003). Microtubules of

varying lengths are observed in neurons, and they can be both stable and dynamic. Numerous

studies have suggested that both hyper- or un-stable microtubules are harmful for maintaining

neuronal morphology as the imbalance in microtubule dynamics contributes to neurological

diseases (reviewed in (Dubey et al. 2015)). Thus, microtubule regulators have been

implicated in a number of human diseases (Dubey et al. 2015; Matamoros and Baas 2016)

and signaling pathways including Notch pathway have been proposed as pharmaceutical

targets for cytoskeletal protection of postnatal neurons (Bonini et al. 2013). Microtubule

dynamics in normal neuronal aging, on the other hand, are poorly understood at the

organismal level.

The neurons of *Caenorhabditis elegans* provide a valuable model system because they share

fundamental characteristics with mammalian neurons in terms of their molecular basis and

functions. Major genetic pathways governing early neuronal development such as neuronal

cell migration or polarization were initially identified from numerous studies on *C. elegans*

(Hedgecock et al. 1990; Wadsworth et al. 1996; Culotti and Merz 1998; Ikegami et al. 2004).

Additionally, *C. elegans* neurons could be utilized to understand the molecular mechanism of

neuronal aging as they demonstrate age-associated structural and functional deterioration. In

particular, touch receptor neurons (TRNs), comprising 2 ALM, AVM, 2 PLM, and PVM,

show the most prominent decline in structure and function with aging (Pan et al. 2011; Tank

et al. 2011; Toth et al. 2012). In aged animals, TRNs have highly wavy or ectopically

branched processes, misshaped cell bodies with abnormal morphology or outgrown processes.

Moreover, *C. elegans* exhibits gradual decreases in touch responses as they age (Tank et al.

2011). DAF-2/IGF pathway, JNK/MAPK pathway and membrane activity of TRNs have

been identified to influence the speed of touch neuronal aging by modulating whole

organismal senescence or cell-autonomous functions (Pan et al. 2011; Tank et al. 2011; Toth

et al. 2012).

The Hippo (Hpo) kinase signaling pathway, which is evolutionarily conserved across animal

species, plays a pivotal role in the regulation of tissue size homeostasis (Harvey et al. 2013).

It regulates cell proliferation and apoptosis during early development and tumorigenesis.

When the Hpo pathway is active, LATS, the core kinase of the pathway, phosphorylates and

inhibits the nuclear localization of YAP, a transcriptional co-activator. In the nucleus, YAP

regulates target gene expression through interaction with the TEAD transcription factor

(Zhao et al. 2007; Zhao et al. 2008). The dysregulation of this pathway promotes

uncontrolled cell proliferation, leading to tumorigenesis and developmental problems (Huang

et al. 2005; Zhao et al. 2007; Harvey et al. 2013; Zheng and Pan 2019). In *C. elegans*, core

components and their genetic interaction of the Hpo pathway are conserved. Inhibition of

YAP-1/YAP by upstream WTS-1/LATS is needed for the maintenance of apicobasal

membrane polarity of the developing intestine (Kang et al. 2009; Lee et al. 2019). The

functions of the Hpo pathway in growth control are extensively established; however, its

post-developmental functions are relatively unknown.

Here, we establish a novel genetic animal model that reveals the postnatal roles of the Hpo

pathway in the differentiated neurons. Loss of *wts-1* leads to premature decline of TRNs both

in structure and function and downstream effectors, *yap-1* and *egl-44,* are required in this

phenomena. We employ genetic and chemical approaches to elucidate the cellular and

molecular mechanisms by which *wts-1*-deficient TRNs lead to a premature deformation. Our

results demonstrate that hyper-stabilized microtubules of the *wts-1* mutant or aged organism

are responsible for neuronal deformation. The fact that loss of *spas-1*, a putative microtubule-

severing enzyme, accelerates onset of age-associated neuronal deformation also supports the

contribution of hyper-stabilized microtubules in neuronal failures. Our study unveils the post-

developmental functions of the Hpo pathway in the maintenance of neuronal integrity and

highlights the importance of neuronal microtubule status in cellular aging and clinical

approaches.

Results

***wts-1* mutant showed impaired structure of touch receptor neurons**

A previous study demonstrated that the intestinal Hpo pathway is essential for the proper

development of the intestinal lumen and loss of *wts-1*, the core kinase of the pathway, leads

to early larval death (Lee et al. 2019). To define the non-intestinal roles of the Hpo pathway

in *C. elegans*, we used the mosaic *wts-1* mutant system of which larval lethality was

overcame with the intestine-specific expression of WTS-1. Since *wts-1* activity in all tissues

except the intestine is absent in this system, hereafter we will refer to it as the *wts-1* mutant

shortly.

We observed that the *wts-1* mutants showed extremely distinctive abnormalities in the

morphology of touch receptor neurons (TRNs). In the wild-type, ALM and its posterior

homolog PLM extended their neuronal processes along the anterior-posterior axis and

maintained structural integrity during adolescence (Fig. 1A, B). However, the *wts-1* mutant-

ALM and PLM exhibited ectopic swelling/waviness and branching (Fig. 1A, B), and other

various deformations, including extended or shortened distal processes or outgrowth of

processes from the cell body (somatic outgrowth) (Supplemental Fig. S1A, B). Atypical

structures of other TRNs, AVM and PVM, were also found in the mutants (Supplemental Fig.

S1C, D), indicating that *wts-1* is essential in mechanosensory neurons. As ALM and PLM

defects were much more severe, frequent and easier to observe, we focused on these neurons

for further study.

TRNs of *C. elegans* develop rapidly in their early life and govern escape behaviors in

response to aversive stimuli (Chalfie et al. 1985). ALM and PLM complete their development

during embryogenesis, whereas AVM and PVM develop until the first larval stage (L1)

(Chalfie et al. 1985). To determine whether *wts-1* is required for the development or

maintenance of TRNs, neuronal morphology of early larva was analyzed. The *wts-1* mutant

had intact, unaltered neuronal processes of ALM and PLM at the very early L1 stage and the

frequency of abnormal neurons gradually increased as the worms grew (Fig. 1C, D). At the

fourth larval (L4) stage, 98.89% of ALM and 84.44% of PLM, respectively, showed at least

one morphological abnormality in the mutant whereas they maintain intact structures in the

wild-type (Fig. 1D). Among the abnormalities, ectopic swelling and branching on the

processes were the most common (Supplemental Fig. S1E, F), followed by extension of the

distal process and somatic outgrowth of the cell body (Supplemental Fig. S1G, H). All these

structural deformations were gradually increased. Ectopic swelling or branching occurred

multiple places in a single process and the number of swelling/branching also increased as the

worms grew (Supplemental Fig. S1I).

To determine whether *wts-1* acts in the structural maintenance of neurons generally or TRNs

specifically, we analyzed the morphologies of other neurons in the mutants. Dopaminergic,

GABAergic, and cholinergic neurons were found to maintain structures comparable with

those of the wild-type at the same age (Supplemental Fig. S1J, K). It is noteworthy that ALN,

a cholinergic neuron that extends its process in association with ALM (White et al. 1986),

preserved their structural integrity in the mutants (Supplemental Fig. S1J). Thus, we

concluded that the loss of *wts-1* leads to wide range of structural deformities specifically in

TRNs and that this deformity is a matter of maintenance, rather than development of the

structures.

***wts-1* deficiency causes TRNs to deteriorate its structure and function**

TRNs govern the aversive movements of worms in response to gentle body touches (Chalfie

et al. 1985). To determine whether structurally deformed TRNs of *wts-1* affect neuronal

functions, we measured gentle touch responses of the mutant. At the L4 stage, control worms

responded to almost every gentle touch (9.40 responses/10 touches), whereas *wts-1* mutants

responded to 60% of stimuli (6.31/10) (Fig. 1E). On the first day of adulthood (1DA), control

worms showed slightly decreased touch responses compared with the L4 stage (8.25/10),

whereas *wts-1* mutants displayed more impaired responses (5.57/10) (Fig. 1E).

The structural and functional decline in the *wts-1* mutants is very similar with phenotypes of

normally aged animals. Morphological alteration and consequent functional decline are

typically observed in the aged nervous system across the animal kingdom (Yankner et al.

2008; Bishop et al. 2010; Fjell and Walhovd 2010). Aging of human brain in the absence of

disease is often accompanied by structural deformation, such as dendritic restructuring,

neuronal sprouting and synaptic losses, rather than severe neuronal degeneration or cell death

(Yankner et al. 2008). Consistently, the *C. elegans* nervous system exhibits age-associated

structural deformation. Among several neurons, TRNs exhibited the most robust decline in

the structure and function during aging (Pan et al. 2011; Tank et al. 2011; Toth et al. 2012).

Structural abnormalities of ALM or PLM appear from the 4[th] day of adulthood and occur in

almost every individual after the 15[th] day of adulthood (Toth et al. 2012). According to our

observations, the degree of morphological deformation of the L4 stage *wts-1* was similar to

that of the aged wild-type worms on the 15[th] day of adulthood. These results suggest that loss

of *wts-1* leads to structural and functional impairment of TRNs precociously.

yap-1 and _egl-44_ suppress premature deformation of _wts-1_ TRNs

To determine whether *yap-1* acts downstream of *wts-1* in this phenomenon, we assessed

genetic interaction of *yap-1* and *wts-1* mutant. In contrast to the *wts-1* mosaic mutant, the *wts-*

1; yap-1 mutant could survive without intestinal WTS-1 expression as previously reported

189 (Lee et al. 2019) and displayed intact neuronal processes without any swelling or branching

190 (Fig. 2A). Introduction of a mutation in *egl-44*, which is the worm homolog of TEAD and

191 functions downstream of *wts-1* along with *yap-1* in the developing intestine (Lee et al. 2019)*,*

192 was also sufficient to suppress the structural abnormalities of *wts-1* TRNs (Fig. 2A). To better

193 understand the basic features of structural dis-integrity in the *wts-1* mutant and mitigating

194 effects of the *yap-1* mutation, we examined TRNs from each genetic background using

195 electron microscopy. In wild-type TRNs, multiple microtubules with characteristic spherical

196 shapes were observed (Fig. 2C). The number of microtubules present in cross sections of

197 control ALM was comparable to the previous report (Chalfie and Thomson 1979) (n = 39, 48,

198 55, respectively (Supplemental Fig. S2B)). In contrast, in the *wts-1* mutant ALM, the number

199 of microtubules significantly decreased (n = 16, 4, 18, respectively (Supplemental Fig. S2C))

200 and the morphology was also irregular and non-spherical (Fig. 2C). Consistent with our

201 finding using a fluorescence marker, the abnormalities in the number and shape of

202 microtubules of TRNs were restored in the *wts-1; yap-1* mutant (n = 24, 30, 44, respectively

203 (Supplemental Fig. S2D)) (Fig. 2C). The loss of *yap-1* also restored the impaired touch

204 responses of the *wts-1* mutant. In comparison with the *wts-1* mutant, touch responses of the

205 *wts-1; yap-1* double mutant were significantly rescued (Fig. 2D, E).

206 Furthermore, to rule out the possibility that the ameliorating effect of *yap-1* mutation on

207 neuronal aging was an indirect result of the delayed organismal aging, we measured the

208 lifespan of each mutant. The *wts-1* mutant had a shorter lifespan than the wild-type,

209 suggesting that prematurely deformed neurons of the *wts-1* mutant were probably due to

210 accelerated organismal aging (Fig. 2F). However, given that the *wts-1; yap-1* mutant with

211 restored neurons had a slightly shorter lifespan than the *wts-1* mutant (Fig. 2F) and only

212 TRNs were affected in the mutant, premature deformation of the *wts-1* neurons appeared to

213   be a touch neuron-specific event, rather than being associated with whole body.

The Hpo pathway acts in a cell autonomous manner to maintain TRNs integrity

To ascertain whether this function of WTS-1 is cell-autonomous or not, we knocked down

WTS-1 selectively in TRNs by injecting tissue-specific RNAi constructs into wild-type

animals as previously described (Esposito et al. 2007). *S*ense and *a*nti*s*ense constructs

corresponding to a region containing kinase domain (sas1) or many exons (sas2) of the *wts-1*

gene were expressed under *mec-4* promoter which is specifically active in all six TRNs (Fig.

3A). For each construct, three transgenic lines were obtained and TRNs structure of these

lines were compared to that of the controls only harboring the transgenic marker, P*unc-*

122::RFP. While the control did not have any abnormalities of TRNs, every transgenic lines

showed significant disintegration of TRNs such as ectopic neuronal swellings or shortened

neuronal processes as seen in *wts-1* mutant at L4 stage (Fig. 3B–D). These neuronal

abnormalities induced by TRNs-specific *wts-1* RNAi was consistently observed in 1DA and

the penetrance of phenotype was slightly increased (Supplemental Fig. S3A–C). These results

documented that WTS-1 functions in TRNs to maintain intact neuronal structures.

Regarding the mechanism of action of the Hpo pathway in cells, YAP-1 likely acts in the

same cell with WTS-1. To clarify this, we rescued YAP-1 activity specifically in TRNs of

wts-1; yap-1 mutants. Touch neuronal expression of YAP-1 using a *mec-4* promoter was

sufficient to re-emerge ectopic neuronal swelling in the *wts-1; yap-1* mutants. Approximately

40% of transgenic worms exhibited ectopic swelling and branching on ALM or PLM. In

contrast, siblings containing no transgene preserved intact neuronal processes (Fig. 3E, F).

Overexpression of YAP is commonly observed in many human cancers and is associated

with poor prognosis of the diseases (Liu et al. 2013; Yuan et al. 2016; Guo et al. 2019).

Inducible overexpression of YAP is sufficient to increase liver size in mice through

transcriptional regulation of downstream genes that activate cell proliferation (Dong et al.

2007). To see if TRNs-specific overexpression of YAP-1 is sufficient to induce neuronal

abnormalities, we overexpressed P*mec-4*::YAP-1 in a wild-type and found that

overexpressing YAP-1 causes multiple neuronal defects in a dose-dependent manner (Fig.

3G–J). Lower expression of YAP-1 (1 ng/μl, final concentration in the injection mixture)

rarely affects neuronal structure of ALM or PLM (Fig. 3I, J). Transgenic worms injected

P*mec-4*::YAP-1 at 10 ng/μl final concentration slightly, but not significantly, elevated

neuronal defects of TRNs. However, higher expression of YAP-1 in TRNs (50 or 100 ng/μl)

resulted neuronal defects similar to TRNs-specific *wts-1* knockdown (Fig. 3G–J). It seems

that endogenous WTS-1 could inhibit overexpressed YAP-1 at lower concentration along

with endogenous YAP-1, however, fails to inhibit overexpressed YAP-1 at higher levels.

These results demonstrated that *wts-1* and *yap-1* act in a cell-autonomous manner to induce

structural disintegration of TRNs; thus, proper regulation of YAP-1 activity by WTS-1 is

important to maintain neuronal structural integrity.

Reduced movement alleviates the structural abnormalities of *wts-1*

Neuronal processes contain bundles of microtubules, which are essential structural

foundations that protect neurons from mechanical strain-induced damage (Tang-Schomer et

al. 2010; Krieg et al. 2017). We hypothesized that the absence of *wts-1* would affect the

microtubule stability in some way; thus, physical stresses generated by movement could

induce spontaneous degeneration of neurons. To test this hypothesis, we slowed the

movement of *wts-1* mutant via the knockdown of muscle machinery genes using RNAi

and observed neuronal morphology. As previously reported (MacLeod et al. 1977; Neumann

and Hilliard 2014), the knockdown of muscle machinery genes, including a myosin heavy

chain gene (*unc-54*), severely impaired organismal movement. Compared with the *wts-1*

mutants who were fed with an empty vector (L4440), worms fed with *unc-54* RNAi exhibited

uncoordinated movement and some of them were even immobilized (Fig. 4A). Impaired

movement resulting from *unc-54* knockdown were also appeared as reduced swimming of

animals (Fig. 4E). Consistent with our hypothesis, *unc-54* RNAi resulted in a significant

reduction in neuronal swelling/branching events in ALM as well as PLM (Fig. 4B–D). The

control ALM showed 7.67 lesions on average, whereas ALM treated with *unc-54* RNAi had

4.851 lesions at the L4 stage. For PLM, the events of swelling/branching occurred 6.59 times

on average in the controls, whereas 2.96 in the *unc-54* RNAi fed neurons at the same stage

(Fig. 4C). Additionally, *unc-54* RNAi significantly increased the proportion of intact PLM

without any swelling or branching (Fig. 4D). The knockdown of *unc-15* or *unc-95,* which

encodes paramyosin (a structural component of thick filaments) and muscle LIM domain

protein, respectively (Kagawa et al. 1989; Broday et al. 2004), showed movement defects as

well and exhibited ameliorative effects on neuronal structures (Fig. 4E, F). In contrast, *unc-*

89 and *deb-1* RNAi failed to induce distinguishable movement defects in our experiments

(Fig. 4E). As expected, *unc-89* RNAi did not mitigate neuronal swelling in the *wts-1* mutants

and *deb-1* RNAi only had limited effects on ALM, and no effect on PLM (Fig. 4F). These

observations indicate that the severity of movement defects is inversely correlated with the

severity of neuronal deformation and it strongly supports the hypothesis that *wts-1* neurons

are vulnerable to physical stresses generated by organismal movements.

Treatment with colchicine, a microtubule-destabilizing agent, reduced ectopic lesions in

wts-1 **neurons**

For further understanding of the status of microtubules in the *wts-1* mutant neurons, we

evaluated the effects of drugs regulating microtubule stability in the mutant neurons. We

transferred the *wts-1* mutants to the drug-containing plates at the L4 stage and scored the

number of lesions including ectopic swelling and branching along the processes of their L4

stage-offspring. Interestingly, colchicine, a microtubule-destabilizing agent (Vandecandelaere

et al. 1997), greatly reduced the morphological abnormalities of the *wts-1* mutant neurons

(Fig. 4G). The drug-untreated ALM showed 5.88 swelling/branching events on a process,

whereas the colchicine-treated ALM only had 2.95 swelling/branching. In the case of PLM,

colchicine treatment reduced neuronal lesions from 4.41 to 0.89 (Fig. 4H). Structurally intact

neurons were significantly increased by colchicine-treatment (Fig. 4I) and the beneficial

effect of colchicine on neuronal structures was consistently observed in TRNs of 1DA

offspring (Fig. 4J, K). Furthermore, treatment with colchicine significantly improved touch

responses of the mutants at L4 stage (Fig. 4L).

In contrast, treatment with paclitaxel, a microtubule-stabilizing agent (Schiff et al. 1979;

Vandecandelaere et al. 1997; Yvon et al. 1999), failed to mitigate neuronal abnormalities of

the *wts-1* mutants (Supplemental Fig. S4A–C). Because of the poor cuticle permeability of

paclitaxel, we introduced a mutation in the *bus-17* gene that encodes galactosyltransferase to

damage the cuticle of the *wts-1* mutant, as performed in previous studies (Bounoutas et al.

2009; Neumann and Hilliard 2014). The number of neuronal lesions was comparable between

the paclitaxel-treated *wts-1* neurons and untreated controls (Supplemental Fig. S4B). These

results show that treatment with colchicine, not paclitaxel, lessened the morphological and

functional alteration of *wts-1* neurons, suggesting that hyper-stabilized microtubules are

responsible for the structural and subsequent functional decline of the mutant neurons.

Hyper-stabilized microtubules are responsible for age-associated neurodegeneration

Next, we questioned whether colchicine could protect neurons from age-associated morphological alteration in wild-type worms. As colchicine treatment had a detrimental effect on the neurodevelopment of wild-type unlike in case of the *wts-1* mutant (Supplemental Fig. 5A), fully developed worms were transferred to colchicine-containing plates at the first day of adulthood (1DA), and neuronal morphologies were monitored in the same generation to assess age-associated morphological abnormalities. Among morphological alterations of TRNs, bead-like structures on the PLM did not accrue in the aged worms. Beaded PLM was most frequently observed in the young adults (2DA) and gradually decreased as the worms grew older (Supplemental Fig. 5I). This observation was consistent with the previous study (Toth et al. 2012); thus, we excluded beaded PLM from defective neurons in further analyses.

Notably, both ALM and PLM of colchicine-treated worms maintained intact structures even after 20 days of adulthood (20DA) (Fig. 5A–D). While untreated 20DA had approximately 73.1% and 57.5% defective ALM and PLM, respectively, colchicine-treated worms showed only 23.6% and 7.2% defective ALM and PLM, respectively (Fig. 5C, D). Colchicine treatment significantly reduced multiple structural abnormalities of TRNs, including somatic outgrowth of ALM and ectopic swelling or branching of PLM (Supplemental Fig. 5B–E). However, it failed to mitigate other morphological abnormalities such as beaded process of ALM and somatic outgrowth of PLM (Supplemental Fig. 5F–J). More importantly, colchicine treatment also improved touch responses of aged animals. Whereas there was no difference between touch responses of colchicine-treated and untreated 2DA animals, drug-treated 10DA animals showed significantly higher touch responses compared to the untreated

controls (Fig. 5E). However, touch responses of drug-treated 10DA were still lower than

those of drug-treated 2DA (Fig. 5E). It seems plausible, given that TRNs of colchicine-

treated 10DA were more structurally defective than that of colchicine-treated 2DA (Fig. 5C,

D). Considering that the same method of colchicine treatment did not affect worm lifespan

(Fig. 5F), the protective effects of colchicine on the structural and functional integrity of

TRNs are unlikely to be derived from the prolonged lifespan of worms.

Next, we examined that hyper-stabilization of neuronal microtubules by paclitaxel treatment

could mimic aged phenotype of TRNs. Age-synchronized *bus-17(br2)* mutants, in which

drug-permeability was enhanced as mentioned, were transferred to paclitaxel-containing

plates at L4 stage and their neuronal morphologies were scored at 5DA stage of the same

generation. Although Treatment with paclitaxel did not affect PLM structure, it significantly

increased neuronal defects of ALM (Supplemental Fig. 5K, L). Paclitaxel induced beaded

processes, somatic outgrowth and cell body abnormalities of ALM, as well as bipolar growth

which is reported to be induced by paclitaxel. Taken together, hyper-stabilized microtubules

could be responsible for age-related neuronal degeneration and neuronal status of the *wts-1*

mutant was similar to that of aged neurons.

Loss of *spas-1*, a putative microtubule-severing enzyme, accelerates structural decline of

touch receptor neurons

Spastin, an ATP-dependent microtubule-severing enzyme, is known to cause a human

neurodegenerative disease, hereditary spastic paraplegia (HSP) (Hazan et al. 1999; Svenson

et al. 2001). Hyper-stabilized microtubules by the loss of spastin activity have been

considered as a main contributor of disease pathology (Sherwood et al. 2004; Evans et al.

2005). SPAS-1, the worm homolog of spastin, possesses a microtubule-severing activity and

is expressed in the TRNs (Matsushita-Ishiodori et al. 2007; Brown and El Bejjani 2017). We

investigated whether hyper-stabilization of microtubules via genetic perturbation of *spas-1*

could cause premature structural defects of TRNs as observed in *wts-1* mutant. We found that

the *spas-1* mutants exhibit premature deformation of TRNs. From the 2nd day of adulthood

(2DA), the mutant ALM displayed an accelerated onset of neuronal deformity (Fig. 6A, C).

Although the frequency of defective PLM was not significantly increased in the mutants

except in *spas-1(tm683)* at 6DA (Fig. 6D), both the deletion mutants *tm683* and *ok1608*

exhibited more various and severe structural deformation of PLM than controls on the 2DA

and 6DA (Fig. 6B, Supplemental Fig. 6A). Ours and others' observation showing touch

neuronal expression of SPAS-1 suggest cell-autonomous function of SPAS-1 (Supplemental

Fig. 6B) (Brown and El Bejjani 2017). Consistent with this hypothesis, expression of SPAS-1

under *mec-4* promoter, as well as *spas-1* own promoter, restored premature deformation of

the *spas-1* mutant ALM (Fig. 6E, F). Human SPAST expression under the control of the *C.*

elegans spas-1 promoter displayed similar rescue effects, whereas the loss-of-function mutant

(K388R), which is associated with disease pathology (Fonknechten et al. 2000), failed to

restore the structural integrity of the mutant ALM (Fig. 6E). In the case of PLM, human

spastin expression or P*mec-4*-driven Ce_SPAS-1 expression slightly, but not significantly,

reduced the structural deformation of the mutants ($p = 0.0502$, $p = 0.1161$, respectively), and

other rescue constructs did not affect the PLM structures (Fig. 6E, F).

Next, we investigated that pharmacological destabilization of neuronal microtubules by

colchicine treatment could ameliorate structural defects of *spas-1* mutant. As done in *wts-1*

mutant background, F1 progenies of animals grown on the drug-contained plate were

observed at L4 stage. We found that colchicine treatment not only rescued structural defects

of *spas-1* mutant, but also increased touch responses of the mutant at 2DA (Fig. 6G, H).

In human spastin, K388R mutation completely abrogates the ATP binding in the ATPase

region and microtubule-severing activity (Evans et al. 2005). Taken together, microtubule-

severing activity of SPAST is likely conserved in *C. elegans* SPAS-1 and it is important to

protect neuronal structures from gradual deformation. Moreover, the fact that the loss of

microtubule-severing enzyme led to accelerated neuronal deformation also supports that

hyper-stabilized microtubules are responsible for the age-associated structural decline of

neurons.

Loss of microtubule-stabilizing genes *dlk-1* and *ptl-1* delayed premature deformation of

***wts-1* neurons**

We next studied the genetic interaction between *wts-1* and several genes that are known to

regulate microtubule stability, particularly those expected to act in TRNs. We selected six

potential microtubule-stabilizing genes (*atat-2*, *dlk-1*, *mec-17*, *mig-2*, *ptl-1* and *ptrn-1*) and

five microtubule-destabilizing genes (*kin-18*, *rho-1*, *unc-33, spas-1* and *elp-1*). A *wts-1; eri-1*

mutant were generated by genetic mating to enhance the efficiency of neuronal RNAi

(Kennedy et al. 2004) and we evaluated lesions in the TRNs of worms fed with the RNAi

vector of target genes. RNAi against microtubule-stabilizing genes, except for *atat-2* and

mec-17, could reduce the structural deformation of *wts-1; eri-1* mutant neurons

(Supplemental Fig. 7B). *atat-2* RNAi slightly alleviated ectopic lesions in ALM and PLM.

mec-17 RNAi failed to reduce PLM lesions but increased ALM lesions. In contrast, none of

the microtubule-destabilizing genes reduced ectopic lesions of the mutant neurons. Among

the candidate genes, *rho-1* RNAi resulted in early larval arrest, making it infeasible to test;

thus, RNAi experiments were performed for only four genes, which failed to reduce the

structural deformation of *wts-1* mutant neurons (Supplemental Fig. 7A).

To address the genetic interaction of microtubule-stabilizing genes and *wts-1* in detail, we

constructed double mutants with *wts-1* and *dlk-1*, *ptl-1*, and *ptrn-1*. Both *wts-1; dlk-1* and

wts-1; ptl-1 displayed improved TRNs structures. The total number of ectopic lesions was

significantly reduced and the proportion of intact neurons without any deformed structures

was also highly increased (Fig. 7A–C). In the case of *ptrn-1*, neuroprotective effects of the

gene knockdown were not reproduced in the double mutant (Supplemental Fig. 7C, D). The

fact that the loss of microtubule-stabilizing genes mitigates the structural deformation of the

wts-1 also supports the finding that the *wts-1* mutants have highly stabilized microtubules

which are responsible for the structural deformation.

ptl-1 encodes the sole *C. elegans* homolog of the microtubule-associated protein tau (MAPT)

(McDermott et al. 1996). In mammals, tau is predominantly localized to the neuronal axons

and promotes microtubule assembly and stability; moreover, mutations in the MAPT locus

are highly associated with several neurodegenerative diseases, such as FTD-17 (Goedert and

Spillantini 2000). In *C. elegans*, the loss of *ptl-1* itself results in premature structural

disintegration of TRNs (Chew et al. 2013). Given these observations with the *wts-1; ptl-1*

mutants and the revealed functions of MAPT, the structural decline seen in the *ptl-1* mutants

is probably due to the microtubule destabilization, unlike those observed in the *wts-1* mutants.

Consistent with this assumption, the loss of *yap-1* did not lessen structural deformation of *ptl-*

1, even increased the structural abnormalities of *ptl-1* (Fig. 7D, E). On the 5^{th} day of

adulthood, *ptl-1* mutant exhibited 41% defective ALM and 6% of defective PLM. In the *ptl-1;*

yap-1 mutant, cells with irregular shape, ectopic branching or somatic outgrowth were highly

increased; thus, 76% ALM and 25% PLM showed structural abnormalities (Fig. 7D, E). Loss

of *yap-1* could probably lead to microtubule destabilization and it has a redundant effect on

microtubule stability with microtubule destabilization due to *ptl-1* loss.

Discussion

In this study, we showed that the loss of *wts-1*, the core kinase of the Hpo pathway, results in

postnatal deformation of TRNs. Although *wts-1* mutants had intact neurons at the beginning,

they gradually exhibited structurally and functionally declined TRNs. The detailed

observation of morphological alteration in *wts-1* mutant suggests that the features of defective

TRNs in the *wts-1* mutant closely resemble those of aged TRNs (Toth et al. 2012). We also

defined that both *yap-1* and *egl-44* act as downstream of *wts-1* and that TRNs-specific rescue

of YAP-1 was sufficient to re-induce the neuronal deformation of *wts-1; yap-1* double mutant.

Neuronal defects induced by knockdown of *wts-1* or overexpression of YAP-1 selectively in

TRNs of wild-type animals also proved a cell-autonomous function of the Hpo pathway in

TRNs. Given that *daf-2*/IGF signaling modulates whole organismal aging, premature decline

of the *wts-1* mutants is more local and restricted in TRNs. These observations say that the

Hpo pathway of *C. elegans* has a neuroprotective effect in differentiated neurons in a cell-

autonomous manner, restricting YAP-1 from triggering premature neuronal decline.

In addition to its extensively studied roles in early development and tumorigenesis,

dysregulation of the Hpo pathway has been implicated in aging and pathologies of the

nervous system (Sahu and Mondal 2020; Gogia et al. 2021). Hyper-activation of the pathway

components such as MST1 and the consequent inhibition of YAP have been noted in several

neurodegenerative disease models (Matsumoto et al. 2014; Yamanishi et al. 2017; Mueller et

al. 2018; Tanaka et al. 2020). In these models, the inactivation of the pathway or the

activation of YAP ameliorates neuronal cell death; thus, the manipulation of the Hpo pathway

has been regarded to prevent pathologies associated with neuronal cell death. Our results

showing that the proper inhibition of YAP-1 by an upstream WTS-1 is required to maintain

neuronal integrity appear to contradict these previous observations. However, alterations that

occur in the aged brain or the *wts-1* mutants are structural alterations or failures in structural

maintenance, which are different from neuronal cell death that occurs in neurodegenerative

diseases. This could explain the conflicting function of YAP-1 in terms of neuroprotection.

Moreover, some examples present the detrimental effects of uncontrolled YAP.

Dysregulation of the pathway and ectopic activation of YAP have been observed in patients

with Alexander disease, which is a rare neurodegenerative disease that results in progressive

neuronal degeneration based on the loss of myelin (Wang et al. 2018). Further, the activation

of YAP also occurred in Müller cells during retinal degeneration (Hamon et al. 2017). The

present study shows that proper regulation of YAP is essential to maintain neuronal integrity

and adds important information to the research on anti-aging roles of the Hpo pathway after

development.

Genetic and pharmacological approaches suggest that defective, hyper-stabilized

microtubules in the *wts-1* mutant were responsible for premature deformation of neurons,

possibly in a 'wear-to-tear fashion' because of the mechanical strains generated by

organismal movements. Reduced movement or treatment with colchicine had significant

ameliorating effects on neuronal deformation of the *wts-1* mutant. Colchicine binds

irreversibly to tubulin dimers and prevents the addition of tubulin dimers to the fast growing

ends of microtubule (Ravelli et al. 2004). It has been widely used to cure acute gouty arthritis

and familial Mediterranean fever (FMF) (Dinarello et al. 1974; Zemer et al. 1974; Zemer et al.

1986). Our study demonstrated that colchicine reduced the severity of neuronal lesions and

even improved neuronal function in the *wts-1* mutant TRNs. Moreover, colchicine treatment

on fully developed animals largely reduced age-related failures in neuronal structures and

functions during normal aging. This supports the idea that neuronal deformation seen in the

wts-1 mutants is similar to that in aged organism and more importantly, hyper-stabilization of

neuronal microtubules would be a promising target to cure age-associated neuronal

deformation.

These observations are unexpected because microtubule-destabilizing agents mimic several

neuronal degenerative disease phenotypes, and microtubule stabilizing agents such as

paclitaxel and epothilone D exhibit neuroprotective effects in the pathological contexts

(Zhang et al. 2005; Shemesh and Spira 2011; Zhang et al. 2012). Some reports have shown

that colchicine has detrimental effects on cognitive function in animal disease models

specifically leading to cholinergic neuronal loss (Goldschmidt and Steward 1982; Veerendra

Kumar and Gupta 2002). In contrast, a study in older patients with FMF showed that long-

term colchicine treatment could protect against cognitive decline in patients (Leibovitz et al.

2006) and the neuroprotective functions of colchicine also have been reported (Pratt et al.

1994; Salama et al. 2012). In the case of hereditary spastic paraplegia (HSP) and paclitaxel-

induced peripheral neuropathy, hyper-stabilization of microtubule could be a cause of

neurodegenerative diseases (Cavaletti et al. 1995; Hazan et al. 1999; Trotta et al. 2004; Evans

et al. 2005; Lee and Swain 2006; Scripture et al. 2006; Gornstein and Schwarz 2014).

Mutated spastin is the most common cause of HSP, a hereditary, neurodegenerative disease

that affects the upper motor neurons (Hazan et al. 1999). Similar to the neuroprotective

effects of colchicine in the *wts-1* or aged neurons, vinblastine, a microtubule-destabilizing

agent, rescue axonal swellings in the HSP models (Fassier et al. 2013). Consistently, the loss

of *spas-1*, the worm homolog of spastin, led to premature deformation of TRNs, and

treatment with colchicine alleviated premature decline of *spas-1* mutant TRNs in both

structure and function. Additionally, the fact that impaired spastin activity in several animal

models results in sparser microtubule array at axons (Sherwood et al. 2004; Wood et al. 2006)

provides a possible link between hyper-stabilization and the sparseness of microtubules we

observed. These shared properties among *wts-1* mutant neurons, aged neurons and HSP

models strongly support that the hyper-stabilized microtubules are responsible for the

structural and functional failures of neurons in both senescent and pathological conditions. It

will be needed to determine that colchicine has more general neuroprotective effects on age-

or diseases-associated decline in other animal models and other microtubule destabilizing

agents show similar beneficial effects.

Mutations in several tubulin proteins, microtubule-binding proteins such as tau, and failures

in nerve attachment to the epidermis have been reported to induce similar deformities in

TRNs, as seen in the *wts-1* mutant. In several mutants of *mec-1*, which encodes an ECM

protein, TRNs fail to separate from body-wall muscles and they are prematurely degenerated

(Pan et al. 2011). In contrast to the *mec-1* mutant neurons, TRNs of the *wts-1* mutant

appeared to normally dissociate from the muscles and innervate the epidermal layer

(Supplemental Fig. S2A). Uncontrolled *yap-1* may affect the vesicular transport of membrane

proteins by altering plasma membrane polarity in the same way it acts in the intestine. Since

we were unable to identify transcriptional targets of YAP-1 associated with these phenomena,

the detailed mechanism of how the activated YAP-1 triggers microtubule hyper-stabilization

in TRNs cell-autonomously remains to be explored. The unique 15-protofilament

microtubules in TRNs (Chalfie and Thomson 1982), which may contribute to their

vulnerability to age-related deformation, could be influenced by transcriptional regulation

through YAP-1. Otherwise, YAP-1 may have broader effects in neuronal microtubules and

other characteristics of TRNs, such as their anatomical localization near the cuticle or their

long projections along the body axis, which could also affect age-related deformation. Given

that YAP is known to regulate transcription of secretory proteins, such as TGF-β ligand (Lee

et al. 2016), an extracellular factor possibly influences touch neuronal microtubules in a non-

cell-autonomous manner. Despite these limitations in understanding the underlying

mechanism, the phenotype of the *wts-1* mutant appears much earlier, and its phenotypic

penetrance is notably higher compared with any other known mutant. Moreover, to the best

of our knowledge, this is the one of the first reports to show the impact of the signaling

pathway on the specific neuronal aging. As we have shown in this study, defective TRNs of

the *wts-1* mutant could provide a neuronal model to investigate the potential therapeutic

targets for improving neuronal aging that occurs either prematurely or normally.

Materials and Methods

Worm maintenance and strains worms were maintained at 20°C as previously described

(Brenner 1974), unless noted otherwise. To visualized touch receptor neurons, we isolated

muIs35 [Pmec-7::GFP + lin-15(+)] V from CF1192 *egl-27(n170) II ; muIs35 [Pmec-7::GFP*

+ lin-15(+)] V by outcrossing 4 times with N2 wild-type. This strain was used as a wild-type

control and *muIs35* was transferred to each mutant background to track touch neurons in all

experiment, except that *spas-1* experiments. Since *spas-1* gene is localized at the same

chromosome with *muIs35*, CF702 *muIs32[Pmec-7::GFP + lin-15(+)] II* were used as control

and transferred in the each mutant background. Exact deletion site of *wts-1(ok753)* was

confirmed by PCR. It starts +4025 from ATG start codon and ends +4610. It also has a 'C'

insertion after deletion (581bp deletion and 1 insertion). Following strains were used. *wts-*

1(ok753) I; Ex [Popt-2:: WTS-1::GFP], wts-1(ok753) I; muIs35 V; Ex [Popt-2:: WTS-

1::GFP], wts-1(ok753) I; muIs35 V; yap-1(tm1416) X, wts-1(ok753) I; egl-44(ys39) II;

muIs35 V, wts-1(ok753) I; egl-44(ys41) II; muIs35 V, KP3948 *eri-1(mg366) IV; lin-15B(n744)*

X, FX683 *spas-1(tm683) V,* RB1411 *spas-1(ok1608) V,* KU12 *dlk-1(km12) I,* RB809 *ptl-*

1(ok621) III, ptrn-1(tm5597) X. CB7431 *bus-17(br2) X,* KP3948 *eri-1(mg366) IV; lin-*

15B(n744) X.

Molecular biology Cloning was performed with standard molecular biology techniques. All

expressions vector were based on pPD117.01 unless otherwise noted. Primer sequence

information is available upon request. To visualize dopaminergic, GABAergic and

cholinergic neurons of worms, *dat-1* promoter, *unc-47* promoter, and *cho-1* promoter

respectively, were fused to GFP using a standard fusion PCR method.

Fluorescence microscopy To monitor neuronal morphology, a fluorescence microscopy

(Axioplan2, Carl Zeiss, Inc) was used. All Fluorescence images were acquired using the

confocal microscope (ZEISS LSM700, Carl Zeiss, Inc.) and ZEN software (Carl Zeiss, Inc.).

RNAi For the *unc-54* RNA*i* construct, 2000 bps from 6th exon region was cloned into the

L4440 vector. For *unc-95, deb-1, mig-2, mec-17* and *rho-1*, we used constructs from Marc

Vidal libraries. The other genes were from the J. Ahringer libraries. RNAi feeding

experiments were done by standard methods. 4-6 L4 worms were transferred to plates with

bacteria expressing RNAi constructs, neuronal morphology was scored at L4 stage of F1

generations.

Fluorescence microscopy and phenotype scoring To monitor neuronal morphology, a fluorescence microscope (Axioplan2, Carl Zeiss, Inc) was used. All Fluorescence images were acquired using the confocal microscope (ZEISS LSM700, Carl Zeiss, Inc.) and ZEN software (Carl Zeiss, Inc.) To score mutant phenotype, 30 neuronal cells of stage-synchronized *wts-1(ok753)* mutants were observed for each stage and the experiments were repeated three times. To gain age-synchronized animals, 50 *wts-1* mutants were transferred to the new plate at the first day of adulthood and were removed leaving laid eggs 2 hours later. Since *wts-1(ok753)* develops slower than wild type control, L1 stage worms 16 hours after egg-laying, L2 worms 40 hrs later, L3 worms 54 hrs later and L4 worms 72 hrs later, and 1DA worms after 96hrs were observed. Morphological abnormalities not found in the L4 stage-wild type worms were considered as defects. To quantify phenotype severity of the mutant, total number of ectopic swellings and branching on a neuronal process was measured.

Gentle touch test Touch sensitivity was tested by stroking the animal with an eyebrow hair attached to a toothpick. To test anterior touch response, we stroked around the pharynx of the worm moving forward. In response of touch, the worm moving backward was scored as 'touch sensitive'. In case of posterior touch response, the worm moving backward was tested and tail of the worm was touched.

Drug treatment Colchicine (Sigma-Aldrich, C9754) was added as dry powder into hot agar at 0.1mM concentration. For *wts-1* mutant, 4~5 L4 worms were transferred to plates containing or not containing colchicine. 4 days later, neuronal morphology was scored in

their L4 progenies. To monitor colchicine effects on normal aging, we transferred about 300

synchronized 1day adult animals to each plates containing or not containing the drug and

transferred to new plates every 1~2day to remove F1 progenies. At 2, 4, 6, 8, 10, 15, 20 days

on their adulthoods, 20 touch neuronal cells were observed. In case of paclitaxel (Sigma-

Aldrich, T7402), DMSO was used as the solvent because paclitaxel is poorly soluble in water.

Final concentration of paclitaxel is 1μM. Plates only containing DMSO were used as the

control. The experiment was performed in the same way as the colchicine treatment into *wts-*

1.

Lifespan measurement Lifespan was measured the standard method with some modification.

For each strain or condition, 100 L4 worms were transferred to the plates (25 worms per one

plate, 4 plates) and living worms were transferred into new plates for every 1~3 days to

remove F1 progenies. Worms did not to respond to touch using a platinum wire were scored

to be dead. Animals that ruptured from vulva bursting, bagged, crawled off or burrowed into

the plates were excluded from the analysis. Every measurement was repeated three times and

yielded similar results. One representative experimental result was shown. Statistical analyses

were performed using OASIS2 (https://sbi.postech.ac.kr/oasis2/) (Han et al. 2016) and

significance were determined by Log-rank (Mantel-Cox) test.

Electron microscopy *C. elegans* animals were overlaid with 20% bovine serum albumin in

M9 buffer, and immediately high-pressure frozen using a Leica EM HPM100 apparatus

(Leica, Austria). Animals were transferred to the freeze substitution apparatus (Leica EM

AFS) under liquid nitrogen into a solution containing 2% osmium tetroxide and 2% water in

acetone. Samples were maintained at -90°C for 100 h, slowly warmed to -20°C (5°C per hour)

and maintained for 20 h, and slowly warmed to 0°C (6°C per hour). Three washes with cold

acetone were carried out at 0°C and samples were embedded in Embed-812 (EMS, USA).

After polymerization of the resin at 60 ˚C for 36 h, serial sections were cut with a diamond

knife on an ULTRACUT UC7 ultramicrotome (Leica, Austria) and mounted on formvar-

coated grids. Sections were stained with 4 % uranyl acetate for 10 min and lead citrate for 7

min. They were observed using a Tecnai G2 Spirit Twin transmission electron microscope

(FEI Company, USA).

References

Barnes AP, Polleux F. 2009. Establishment of axon-dendrite polarity in developing neurons.
Annu Rev Neurosci **32**: 347-381.
Bishop NA, Lu T, Yankner BA. 2010. Neural mechanisms of ageing and cognitive decline.
Nature **464**: 529-535.
Bonini SA, Ferrari-Toninelli G, Montinaro M, Memo M. 2013. Notch signalling in adult
neurons: a potential target for microtubule stabilization. *Ther Adv Neurol Disord* **6**:
375-385.
Bounoutas A, O'Hagan R, Chalfie M. 2009. The multipurpose 15-protofilament microtubules
in C. elegans have specific roles in mechanosensation. *Curr Biol* **19**: 1362-1367.
Brenner S. 1974. The genetics of Caenorhabditis elegans. *Genetics* **77**: 71-94.
Broday L, Kolotuev I, Didier C, Bhoumik A, Podbilewicz B, Ronai Z. 2004. The LIM
domain protein UNC-95 is required for the assembly of muscle attachment structures
and is regulated by the RING finger protein RNF-5 in C. elegans. *J Cell Biol* **165**:
857-867.
Brown G, El Bejjani R. 2017. SPAS-1 expression in neurons and vulva during L4 stage.
MicroPubl Biol **2017**.
Cavaletti G, Tredici G, Braga M, Tazzari S. 1995. Experimental peripheral neuropathy
induced in adult rats by repeated intraperitoneal administration of taxol. *Exp Neurol*
133: 64-72.
Chalfie M, Sulston JE, White JG, Southgate E, Thomson JN, Brenner S. 1985. The neural
circuit for touch sensitivity in Caenorhabditis elegans. *J Neurosci* **5**: 956-964.
Chalfie M, Thomson JN. 1979. Organization of neuronal microtubules in the nematode
Caenorhabditis elegans. *J Cell Biol* **82**: 278-289.
-. 1982. Structural and functional diversity in the neuronal microtubules of Caenorhabditis
elegans. *J Cell Biol* **93**: 15-23.
Chew YL, Fan X, Gotz J, Nicholas HR. 2013. PTL-1 regulates neuronal integrity and
lifespan in C. elegans. *J Cell Sci* **126**: 2079-2091.

Culotti JG, Merz DC. 1998. DCC and netrins. *Curr Opin Cell Biol* **10**: 609-613.

Dinarello CA, Wolff SM, Goldfinger SE, Dale DC, Alling DW. 1974. Colchicine therapy for
familial mediterranean fever. A double-blind trial. *N Engl J Med* **291**: 934-937.

Dong J, Feldmann G, Huang J, Wu S, Zhang N, Comerford SA, Gayyed MF, Anders RA,
Maitra A, Pan D. 2007. Elucidation of a universal size-control mechanism in
Drosophila and mammals. *Cell* **130**: 1120-1133.

Dubey J, Ratnakaran N, Koushika SP. 2015. Neurodegeneration and microtubule dynamics:
death by a thousand cuts. *Front Cell Neurosci* **9**: 343.

Esposito G, Di Schiavi E, Bergamasco C, Bazzicalupo P. 2007. Efficient and cell specific
knock-down of gene function in targeted C. elegans neurons. *Gene* **395**: 170-176.

Evans KJ, Gomes ER, Reisenweber SM, Gundersen GG, Lauring BP. 2005. Linking axonal
degeneration to microtubule remodeling by Spastin-mediated microtubule severing. *J*
Cell Biol **168**: 599-606.

Fassier C, Tarrade A, Peris L, Courageot S, Mailly P, Dalard C, Delga S, Roblot N, Lefevre
J, Job D et al. 2013. Microtubule-targeting drugs rescue axonal swellings in cortical
neurons from spastin knockout mice. *Dis Model Mech* **6**: 72-83.

Fjell AM, Walhovd KB. 2010. Structural brain changes in aging: courses, causes and
cognitive consequences. *Rev Neurosci* **21**: 187-221.

Fonknechten N, Mavel D, Byrne P, Davoine CS, Cruaud C, Bonsch D, Samson D, Coutinho
P, Hutchinson M, McMonagle P et al. 2000. Spectrum of SPG4 mutations in
autosomal dominant spastic paraplegia. *Hum Mol Genet* **9**: 637-644.

Goedert M, Spillantini MG. 2000. Tau mutations in frontotemporal dementia FTDP-17 and
their relevance for Alzheimer's disease. *Biochim Biophys Acta* **1502**: 110-121.

Gogia N, Chimata AV, Deshpande P, Singh A, Singh A. 2021. Hippo signaling: bridging the
gap between cancer and neurodegenerative disorders. *Neural Regen Res* **16**: 643-652.

Goldschmidt RB, Steward O. 1982. Neurotoxic effects of colchicine: differential
susceptibility of CNS neuronal populations. *Neuroscience* **7**: 695-714.

Gornstein E, Schwarz TL. 2014. The paradox of paclitaxel neurotoxicity: Mechanisms and
unanswered questions. *Neuropharmacology* **76 Pt A**: 175-183.

Guo LW, Chen YT, Luo J, Zheng JP, Shao GL. 2019. YAP1 overexpression is associated
with poor prognosis of breast cancer patients and induces breast cancer cell growth by
inhibiting PTEN. *Febs Open Bio* **9**: 437-445.

Hamon A, Masson C, Bitard J, Gieser L, Roger JE, Perron M. 2017. Retinal Degeneration
Triggers the Activation of YAP/TEAD in Reactive Muller Cells. *Invest Ophthalmol*
Vis Sci **58**: 1941-1953.

Han SK, Lee D, Lee H, Kim D, Son HG, Yang JS, Lee SV, Kim S. 2016. OASIS 2: online
application for survival analysis 2 with features for the analysis of maximal lifespan
and healthspan in aging research. *Oncotarget* **7**: 56147-56152.

Harvey KF, Zhang X, Thomas DM. 2013. The Hippo pathway and human cancer. *Nat Rev*
Cancer **13**: 246-257.

Hazan J, Fonknechten N, Mavel D, Paternotte C, Samson D, Artiguenave F, Davoine CS,
Cruaud C, Durr A, Wincker P et al. 1999. Spastin, a new AAA protein, is altered in
the most frequent form of autosomal dominant spastic paraplegia. *Nat Genet* **23**: 296-
303.

Hedgecock EM, Culotti JG, Hall DH. 1990. The unc-5, unc-6, and unc-40 genes guide
circumferential migrations of pioneer axons and mesodermal cells on the epidermis in
C. elegans. *Neuron* **4**: 61-85.

Hirokawa N, Niwa S, Tanaka Y. 2010. Molecular motors in neurons: transport mechanisms
and roles in brain function, development, and disease. *Neuron* **68**: 610-638.

Huang J, Wu S, Barrera J, Matthews K, Pan D. 2005. The Hippo signaling pathway
coordinately regulates cell proliferation and apoptosis by inactivating Yorkie, the
Drosophila Homolog of YAP. *Cell* **122**: 421-434.
Ikegami R, Zheng H, Ong SH, Culotti J. 2004. Integration of semaphorin-2A/MAB-20,
ephrin-4, and UNC-129 TGF-beta signaling pathways regulates sorting of distinct
sensory rays in C. elegans. *Dev Cell* **6**: 383-395.
Job D, Valiron O, Oakley B. 2003. Microtubule nucleation. *Curr Opin Cell Biol* **15**: 111-117.
Kagawa H, Gengyo K, McLachlan AD, Brenner S, Karn J. 1989. Paramyosin gene (unc-15)
of Caenorhabditis elegans. Molecular cloning, nucleotide sequence and models for
thick filament structure. *J Mol Biol* **207**: 311-333.
Kang J, Shin D, Yu JR, Lee J. 2009. Lats kinase is involved in the intestinal apical membrane
integrity in the nematode Caenorhabditis elegans. *Development* **136**: 2705-2715.
Kennedy S, Wang D, Ruvkun G. 2004. A conserved siRNA-degrading RNase negatively
regulates RNA interference in C. elegans. *Nature* **427**: 645-649.
Krieg M, Stuhmer J, Cueva JG, Fetter R, Spilker K, Cremers D, Shen K, Dunn AR, Goodman
MB. 2017. Genetic defects in beta-spectrin and tau sensitize C. elegans axons to
movement-induced damage via torque-tension coupling. *Elife* **6**.
Lee DH, Park JO, Kim TS, Kim SK, Kim TH, Kim MC, Park GS, Kim JH, Kuninaka S,
Olson EN et al. 2016. LATS-YAP/TAZ controls lineage specification by regulating
TGFbeta signaling and Hnf4alpha expression during liver development. *Nat Commun*
7: 11961.
Lee H, Kang J, Ahn S, Lee J. 2019. The Hippo Pathway Is Essential for Maintenance of
Apicobasal Polarity in the Growing Intestine of Caenorhabditis elegans. *Genetics* **213**:
501-515.
Lee JJ, Swain SM. 2006. Peripheral neuropathy induced by microtubule-stabilizing agents. *J*
Clin Oncol **24**: 1633-1642.
Leibovitz A, Lidar M, Baumoehl Y, Livneh A, Segal R. 2006. Colchicine therapy and the
cognitive status of elderly patients with familial Mediterranean fever. *Isr Med Assoc J*
8: 469-472.
Liu JY, Li YH, Lin HX, Liao YJ, Mai SJ, Liu ZW, Zhang ZL, Jiang LJ, Zhang JX, Kung HF
et al. 2013. Overexpression of YAP 1 contributes to progressive features and poor
prognosis of human urothelial carcinoma of the bladder. *BMC Cancer* **13**: 349.
MacLeod AR, Waterston RH, Fishpool RM, Brenner S. 1977. Identification of the structural
gene for a myosin heavy-chain in Caenorhabditis elegans. *J Mol Biol* **114**: 133-140.
Maday S, Twelvetrees AE, Moughamian AJ, Holzbaur EL. 2014. Axonal transport: cargo-
specific mechanisms of motility and regulation. *Neuron* **84**: 292-309.
Marin O, Valiente M, Ge X, Tsai LH. 2010. Guiding neuronal cell migrations. *Cold Spring*
Harb Perspect Biol **2**: a001834.
Matamoros AJ, Baas PW. 2016. Microtubules in health and degenerative disease of the
nervous system. *Brain Res Bull* **126**: 217-225.
Matsumoto H, Murakami Y, Kataoka K, Lin H, Connor KM, Miller JW, Zhou D, Avruch J,
Vavvas DG. 2014. Mammalian STE20-like kinase 2, not kinase 1, mediates
photoreceptor cell death during retinal detachment. *Cell Death Dis* **5**: e1269.
Matsushita-Ishiodori Y, Yamanaka K, Ogura T. 2007. The C. elegans homologue of the
spastic paraplegia protein, spastin, disassembles microtubules. *Biochem Biophys Res*
Commun **359**: 157-162.
McDermott JB, Aamodt S, Aamodt E. 1996. ptl-1, a Caenorhabditis elegans gene whose
products are homologous to the tau microtubule-associated proteins. *Biochemistry* **35**:
9415-9423.

Mueller KA, Glajch KE, Huizenga MN, Wilson RA, Granucci EJ, Dios AM, Tousley AR,
Iuliano M, Weisman E, LaQuaglia MJ et al. 2018. Hippo Signaling Pathway
Dysregulation in Human Huntington's Disease Brain and Neuronal Stem Cells. *Sci*
Rep **8**: 11355.
Neumann B, Hilliard MA. 2014. Loss of MEC-17 leads to microtubule instability and axonal
degeneration. *Cell Rep* **6**: 93-103.
Pan CL, Peng CY, Chen CH, McIntire S. 2011. Genetic analysis of age-dependent defects of
the Caenorhabditis elegans touch receptor neurons. *Proc Natl Acad Sci U S A* **108**:
9274-9279.
Pratt J, Roux M, Henneguelle E, Stutzmann JM, Laduron PM. 1994. Neuroprotective effects
of colchicine in the gerbil model of cerebral ischaemia. *Neurosci Lett* **169**: 114-118.
Ravelli RB, Gigant B, Curmi PA, Jourdain I, Lachkar S, Sobel A, Knossow M. 2004. Insight
into tubulin regulation from a complex with colchicine and a stathmin-like domain.
Nature **428**: 198-202.
Sahu MR, Mondal AC. 2020. The emerging role of Hippo signaling in neurodegeneration. *J*
Neurosci Res **98**: 796-814.
Salama M, Ellaithy A, Helmy B, El-Gamal M, Tantawy D, Mohamed M, Sheashaa H, Sobh
M, Arias-Carrion O. 2012. Colchicine protects dopaminergic neurons in a rat model
of Parkinson's disease. *CNS Neurol Disord Drug Targets* **11**: 836-843.
Schiff PB, Fant J, Horwitz SB. 1979. Promotion of microtubule assembly in vitro by taxol.
Nature **277**: 665-667.
Scripture CD, Figg WD, Sparreboom A. 2006. Peripheral neuropathy induced by paclitaxel:
recent insights and future perspectives. *Curr Neuropharmacol* **4**: 165-172.
Shemesh OA, Spira ME. 2011. Rescue of neurons from undergoing hallmark tau-induced
Alzheimer's disease cell pathologies by the antimitotic drug paclitaxel. *Neurobiol Dis*
43: 163-175.
Sherwood NT, Sun Q, Xue M, Zhang B, Zinn K. 2004. Drosophila spastin regulates synaptic
microtubule networks and is required for normal motor function. *PLoS Biol* **2**: e429.
Svenson IK, Ashley-Koch AE, Gaskell PC, Riney TJ, Cumming WJ, Kingston HM, Hogan
EL, Boustany RM, Vance JM, Nance MA et al. 2001. Identification and expression
analysis of spastin gene mutations in hereditary spastic paraplegia. *Am J Hum Genet*
68: 1077-1085.
Tanaka H, Homma H, Fujita K, Kondo K, Yamada S, Jin X, Waragai M, Ohtomo G, Iwata A,
Tagawa K et al. 2020. YAP-dependent necrosis occurs in early stages of Alzheimer's
disease and regulates mouse model pathology. *Nat Commun* **11**: 507.
Tang-Schomer MD, Patel AR, Baas PW, Smith DH. 2010. Mechanical breaking of
microtubules in axons during dynamic stretch injury underlies delayed elasticity,
microtubule disassembly, and axon degeneration. *FASEB J* **24**: 1401-1410.
Tank EM, Rodgers KE, Kenyon C. 2011. Spontaneous age-related neurite branching in
Caenorhabditis elegans. *J Neurosci* **31**: 9279-9288.
Toth ML, Melentijevic I, Shah L, Bhatia A, Lu K, Talwar A, Naji H, Ibanez-Ventoso C,
Ghose P, Jevince A et al. 2012. Neurite sprouting and synapse deterioration in the
aging Caenorhabditis elegans nervous system. *J Neurosci* **32**: 8778-8790.
Trotta N, Orso G, Rossetto MG, Daga A, Broadie K. 2004. The hereditary spastic paraplegia
gene, spastin, regulates microtubule stability to modulate synaptic structure and
function. *Curr Biol* **14**: 1135-1147.
Vandecandelaere A, Martin SR, Engelborghs Y. 1997. Response of microtubules to the
addition of colchicine and tubulin-colchicine: evaluation of models for the interaction
of drugs with microtubules. *Biochem J* **323** (**Pt 1**): 189-196.

Veerendra Kumar MH, Gupta YK. 2002. Intracerebroventricular administration of colchicine
produces cognitive impairment associated with oxidative stress in rats. *Pharmacol*
Biochem Behav **73**: 565-571.
Wadsworth WG, Bhatt H, Hedgecock EM. 1996. Neuroglia and pioneer neurons express
UNC-6 to provide global and local netrin cues for guiding migrations in C. elegans.
Neuron **16**: 35-46.
Wang L, Xia J, Li J, Hagemann TL, Jones JR, Fraenkel E, Weitz DA, Zhang SC, Messing A,
Feany MB. 2018. Tissue and cellular rigidity and mechanosensitive signaling
activation in Alexander disease. *Nat Commun* **9**: 1899.
White JG, Southgate E, Thomson JN, Brenner S. 1986. The structure of the nervous system
of the nematode Caenorhabditis elegans. *Philos Trans R Soc Lond B Biol Sci* **314**: 1-
340.
Wood JD, Landers JA, Bingley M, McDermott CJ, Thomas-McArthur V, Gleadall LJ, Shaw
PJ, Cunliffe VT. 2006. The microtubule-severing protein Spastin is essential for axon
outgrowth in the zebrafish embryo. *Hum Mol Genet* **15**: 2763-2771.
Yamanishi E, Hasegawa K, Fujita K, Ichinose S, Yagishita S, Murata M, Tagawa K, Akashi
T, Eishi Y, Okazawa H. 2017. A novel form of necrosis, TRIAD, occurs in human
Huntington's disease. *Acta Neuropathol Commun* **5**: 19.
Yankner BA, Lu T, Loerch P. 2008. The aging brain. *Annu Rev Pathol* **3**: 41-66.
Yuan Y, Li D, Li H, Wang L, Tian G, Dong Y. 2016. YAP overexpression promotes the
epithelial-mesenchymal transition and chemoresistance in pancreatic cancer cells. *Mol*
Med Rep **13**: 237-242.
Yvon AM, Wadsworth P, Jordan MA. 1999. Taxol suppresses dynamics of individual
microtubules in living human tumor cells. *Mol Biol Cell* **10**: 947-959.
Zemer D, Pras M, Sohar E, Modan M, Cabili S, Gafni J. 1986. Colchicine in the prevention
and treatment of the amyloidosis of familial Mediterranean fever. *N Engl J Med* **314**:
1001-1005.
Zemer D, Revach M, Pras M, Modan B, Schor S, Sohar E, Gafni J. 1974. A controlled trial of
colchicine in preventing attacks of familial mediterranean fever. *N Engl J Med* **291**:
932-934.
Zhang B, Carroll J, Trojanowski JQ, Yao Y, Iba M, Potuzak JS, Hogan AM, Xie SX,
Ballatore C, Smith AB, 3rd et al. 2012. The microtubule-stabilizing agent, epothilone
D, reduces axonal dysfunction, neurotoxicity, cognitive deficits, and Alzheimer-like
pathology in an interventional study with aged tau transgenic mice. *J Neurosci* **32**:
3601-3611.
Zhang B, Maiti A, Shively S, Lakhani F, McDonald-Jones G, Bruce J, Lee EB, Xie SX,
Joyce S, Li C et al. 2005. Microtubule-binding drugs offset tau sequestration by
stabilizing microtubules and reversing fast axonal transport deficits in a tauopathy
model. *Proc Natl Acad Sci U S A* **102**: 227-231.
Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, Xie J, Ikenoue T, Yu J, Li L et al. 2007.
Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact
inhibition and tissue growth control. *Genes Dev* **21**: 2747-2761.
Zhao B, Ye X, Yu J, Li L, Li W, Li S, Yu J, Lin JD, Wang CY, Chinnaiyan AM et al. 2008.
TEAD mediates YAP-dependent gene induction and growth control. *Genes Dev* **22**:
1962-1971.
Zheng Y, Pan D. 2019. The Hippo Signaling Pathway in Development and Disease. *Dev Cell*
50: 264-282.
844

Acknowledgments

We thank the Caenorhabditis Genetics Center and the National BioResource Project for
providing strains, Julie Ahringer for RNAi plasmids. We also thank Andrew Fire for the
worm expressing vectors. The plasmids containing human SPAST genes were gifts from
Jennifer Lippincott-Schwartz (Addgene plasmid #134461, #134463).

Funding

This work was supported by the Samsung Science and Technology Foundation under project
Number SSTF-BA1501-52 and National Research Foundation of Korea grant funded by the
Korean government (MEST) [2019R1A6A1A10073437]. H. Lee was supported by a
scholarship for basic researches, Seoul National University, Seoul, Korea.

Author contributions

Conceptualization: HL, JK, JL

Methodology: HL, JL

Investigation: HL, JK, SL, DL, CHC

Data curation: HL, JL

Supervision: JL

Writing—original draft: HL

Writing—review & editing: HL, JK, JL

Disclosure statement and competing interests

The authors state they have no competing interests or disclosures.

Data and materials availability

All data are available in the main text or the supplementary materials.

Figure 1. *wts-1* mutants show the premature structural and functional decline of touch receptor neurons (TRNs) (A, B) Representative images of TRNs ([A] ALM and [B] PLM) in the wild-type and *wts-1(ok753)* mutant at the L4 stage. In the mutant, morphologically disrupted ALM and PLM were observed to exhibit abnormalities including ectopic swelling/branching on the processes and posterior outgrowth (arrowheads). (C) Newly developed ALM and PLM in the wild-type and the *wts-1(ok753)* mutant at the early L1 stage. The intact neuronal process is indicated using arrowheads. Intestinal expression of the WTS-1 rescue construct (P*opt-2*::WTS-1::GFP) is indicated using an arrow. (A–C) TRNs were visualized by expressing GFP under the control of the *mec-7* promoter (*muIs35*[P*mec-7*::GFP]) and anterior is to the left unless otherwise noted. Scale bar = 20 μm. (D) Penetration of defective TRNs of the wild-type and the *wts-1(ok753)* at different developmental stages. Neurons displaying any morphological abnormalities such as swelling, branching, somatic outgrowth and extended distal process were scored as defective neurons. In one experiment, 30 cells were observed for each strain and each stage and the experiments were repeated 3 times. Statistical significance was determined by a two-way ANOVA followed by Bonferroni's post-test. (E) Quantified touch responses of the wild-type and the *wts-1(ok753)* mutant at L4 and 1st day of adulthood (1DA). N = 30. Statistical significance was determined by an unpaired t-test. ***, $p < 0.001$, **, $p < 0.01$, *, $p < 0.05$, n.s, not significant, compared to WT or control, unless otherwise marked on graph. All data are presented as means ± SEM, unless otherwise noted.

Figure 2. *yap-1* or *egl-44* suppresses premature neuronal decline in *wts-1* mutants. (A, B)
Representative images of (A) ALM and (B) PLM in wild-type, *wts-1(ok753)*, *wts-1(ok753);*
yap-1(tm1416) and *wts-1(ok753); egl-44(ys41)* at the L4 stage. Loss of *yap-1* or *egl-44*
completely restored structural integrity of the *wts-1* mutants. Scale bar = 20 μm. (C) Electron
microscope images of wild-type, *wts-1(ok753)* and *wts-1(ok753); yap-1(tm1416)* at 1DA.
TRNs are indicated using boxes. Scale bar = 500 nm. (D, E) Touch responses of wild-type,
wts-1(ok753) and *wts-1(ok753); yap-1(tm1416)* at (D) L4 stage and (G) 1DA stage. For each
strain and each stage, 90 animals were tested. Statistical significance was determined using a
one-way ANOVA, followed by the Tukey's multiple comparison test. (E) Survival curve of
wild-type, *wts-1(ok753)* and *wts-1(ok753); yap-1(tm1416)*. Worms were maintained at 20°C
and all lines used for the lifespan analysis have *muIs35*.

Figure 3. *wts-1-yap-1* act in a cell-autonomous manner to maintain touch neuronal integrity. (A) The targeting regions for touch neuronal specific RNAi of *wts-1*. Sense and *a*ntisense (sas) genomic fragments were cloned under the TRNs specific promoter, P*mec-4*. (B) Representative images and (C, D) quantified neuronal abnormalities of TRNs in the controls and *wts-1*-knockdowned animals. Both TRNs-specific-sas1- and sas2-based knock down of *wts-1* efficiently induces neuronal abnormalities as seen in *wts-1* mutant. (E–F) Touch neuronal rescue of YAP-1 is sufficient to re-induce neuronal dis-integrity in *wts-1(ok753); yap-1(tm1416)* mutant. (E) Representative images of TRNs of *wts-1(ok753); yap-1(tm1416)* mutant with the rescue construct or its sibling without transgenes. (F) Quantified results. 60 ALM and PLM of transgenic worms and their siblings were observed. (G, H) Touch neuronal overexpression of YAP-1 is sufficient to induce neuronal abnormalities in wild-type animals. Final concentration of P*mec-4*::GFP::YAP-1 in the injection mixture is 100 ng/μl. (I, J) Quantified neuronal defects of (I) ALM and (J) PLM induced by overexpression of P*mec-4*::GFP::YAP-1 at each concentration in the injection mixtures. (C, D, I, J) 90 ALM and PLM were observed in each of the three independent lines. Neuronal morphology was examined at L4 stage. Statistical significance was determined by a one-way ANOVA followed by the Dunnett's multiple comparison test. An unpaired t-test was used for (F). Red: Expression of injection marker, P*unc-122*::RFP, white arrowhead :neuronal swelling, yellow arrowhead: shortened process. Scale bar = 20 μm.

Figure 4. Abnormal microtubules are responsible for neuronal dis-integrity observed in the *wts-1* mutants. (A–D) Reduced movement resulting from *unc-54* knockdown diminishes ectopic swelling or branching in the *wts-1(ok753)* mutant. (A) *unc-54* knockdown leads to uncoordinated movement of worms (yellow arrowhead). (B, C) *unc-54* knockdown reduces the number of neuronal lesions on the ALM and PLM. (D) *unc-54* knockdown slightly, but significantly, increases the percentages of intact PLM and not ALM. Neurons without any structural defects, including swelling or branching, were considered intact. (E) Quantified motor deficits in animals knocking down various muscle machinery genes. RNAi against *unc-15, unc-54* or *unc-95* impaired swimming behavior, whereas *unc-89* and *deb-1* made no differences compared to control. (F) Quantified morphological abnormalities of TRNs in muscle machinery-knockdown animals. Knockdown of *unc-89* or *deb-1* did not alleviate structural decline of both ALM and PLM. (G–K) Treatment with colchicine reduces ectopic neuronal swelling and increases the percentages of intact neurons in the *wts-1(ok753)* mutants. F1 progenies grown on the drug-contained plates were scored at (H, I) L4 stage or (J, K) 1DA. (L) Colchicine treatment improved touch responses of *wts-1* mutant. The behavior test was done at L4 stage. Statistical significance was determined using an unpaired t-test (C, D, H, L) or a one-way ANOVA, followed by the Dunnett's multiple comparison test (E, F). The total number of cells or animals analyzed is indicated in each column. Asterisks indicate differences from L4440-fed or drug-untreated-control neurons. Ectopic neuronal swelling and branching were labeled with white arrowheads. Scale bar = 20μm.

Figure 5. Hyper-stabilized microtubules might be responsible for age-associated morphological deformation of touch receptor neurons. (A, B) Representative images of (A) ALM and (B) PLM of colchicine-treated or untreated wild-type worms. The age of worms is indicated in the image. (C, D) Quantified defects of (C) ALM or (D) PLM in colchicine-treated or untreated worms. Age-synchronized worms were transferred to drug-containing plates on the 1DA and phenotypes were scored on every 2^{nd}, 4^{th}, 6^{th}, 8^{th}, 10^{th}, 15^{th} and 20^{th} day of adulthood. At every time point and in each group, 20 neurons were scored per one experiment and the experiments were repeated 3 times. (E) Colchicine treatment alleviates impaired touch responses of aged animals. Touch responses were scored at 2DA and 10DA. Statistical significance was determined by a two-way ANOVA, followed by the Bonferroni's post-tests (C, D) or a one-way ANOVA with the Bonferroni's multiple comparison test (E). (F) Survival curves of colchicine-treated and untreated control animals. Scale bar = 20μm.

Figure 6. Loss of *spas-1*, a microtubule-severing enzyme, results in premature structural decline. (A, B) Representative images of defective TRNs of *spas-1* mutant on the 2DA. Both deletion mutants, *tm683* and *ok1608,* displayed age-associated morphological alterations of ALM or PLM including ectopic swelling and branching on the neuronal process (white arrowhead), somatic outgrowth and irregular shape of the cell body (yellow arrowhead) precociously. (C, D) Quantified results of structural defects of ALM or PLM in *spas-1* mutants. At every time point, 20 neurons were scored and the experiments were repeated 3 times. (E) Results of the rescue experiment of premature neuronal degeneration of *spas-1(tm683)* with SPAS-1, human SPAST(wt) and human SPAST(K388R). In all cases, the *C. elegans spas-1* promoter was used to induce the C-terminally mCherry tagged transgene. (F) Touch neuronal specific expression of SPAS-1 was sufficient to rescue neuronal defects of ALM. (E, F) Neuronal morphology were scored at 2DA. For each rescue construct, three independent transgenic lines were observed that yielded similar results and the results of one line are presented. (G, H) Quantified (G) structural defects of TRNs or (H) touch responses of colchicine-treated and untreated *spas-1(tm683)* mutant. Analyses were done at 2DA (N =30/one experiment, repeated 3times). Statistical significance was determined using a two-way ANOVA, followed by the Bonferroni's multiple comparison test (C, D), a one-way ANOVA with the Dunnett's multiple comparison test (E, F) or with the Turkey's multiple comparison test (G). Unpaired t-test was used in (H). Scale bar = 20 μm.

Figure 7. *wts-1-yap-1* affect neuronal integrity possibly by modulating microtubule stability (A–C) Loss of *dlk-1* or *ptl-1* significantly mitigates the structural deformation of *wts-1*-mutant neurons. (A) Representative images of ALM (upper panels) and PLM (lower panels) of *wts-1(ok753), dlk-1(km12), ptl-1(ok621), wts-1(ok753); dlk-1(km12)* and *wts-1(ok753); ptl-1(ok621)* at L4 stage. Ectopic lesions were labeled with arrowhead. (B) Average number of ectopic lesions per neuronal process of each strain. (C) Percentage of intact neurons of each mutant. Loss of *dlk-1* or *ptl-1* protects TRNs of the *wts-1* from premature deformation. (D, E) Loss of *yap-1* worsen the neuronal deformation as seen in the *ptl-1* mutant. (D) Unlike *ptl-1(ok621)* single mutant, *ptl-1(ok621); yap-1(tm1416)* double mutant exhibits severe deformation in ALM and PLM, such as irregular shape of cell body (yellow arrowhead) and ectopic branching (white arrowhead) of the neuronal process. (E) Percentage of undamaged touch neurons in *ptl-1(ok621)* and *ptl-1(ok621); yap-1(tm1416)*. (A–E) Neurons were analyzed at the L4 stage. The number of scored neurons is indicated in each column. Statistical significance was determined using a one-way ANOVA, followed by the Dunnett's multiple comparison test. Asterisks indicate differences from the *wts-1* or the *ptl-1* single mutant. Scale bar = 20 μm.

 Supplemental Figure legends

 Figure S1. Loss of *wts-1* leads to touch neuronal-specific structural decline. (A, B)
 Various morphological deformations observed in *wts-1* mutant (A) ALM, (B) PLM. In the
 mutants, both ALM and PLM showed ectopic swelling (a), branching (b) and somatic
 outgrowth (c). ALM also displays an extended distal process (d), posterior outgrowth (e), or
 abnormalities in the cell body (f). PLM exhibits extended (g) or shortened (h) process. TRNs
 were analyzed at the L4 or 1st day of adulthood (1DA) stage. Intestinal expression of the
 WTS-1 rescue construct is indicated using a yellow arrowhead. (C) Structural decline of
 AVM and PVM in the *wts-1* mutants. Somatic outgrowth or ectopic branching/swelling on
 the process is occasionally noted in both AVM and PVM. (D) Quantified structural defects of
 AVM and PVM in wild-type and *wts-1* mutants at different stages. Statistical significance
 was determined using a two-way ANOVA followed by Bonferroni's post-test. (E–H)
 Quantified structural defects of ALM and PLM observed in *wts-1* mutants at different stages.
 (D) Ectopic swelling or (E) branching on the neuronal process occurs the most frequently,
 followed by (F) somatic outgrowth and (G) extended distal process. Structural deformations
 of each category tend to increase as the worms grow. (I) The average number of ectopic
 lesions on a neuronal process of ALM or PLM of *wts-1* at different stages. The total number
 of ectopic swelling or branching was scored. (D–I) At each time point, 90 ALM or PLM were
 analyzed (N=30, repeated 3 times). (J) Representative images of dopaminergic, GABAergic
 and cholinergic neurons in *wts-1* mutants at the L4 stage. Each neuron was visualized using a
 fluorescent marker under the control of a specific promoter; *dat-1* promoter, *unc-47* promoter,
 cho-1 promoter, respectively. (K) Quantified structural defects of dopaminergic, GABAergic
 and cholinergic neurons in wild-type and *wts-1* mutants. N = 90. ***, $p < 0.001$, **, $p < 0.01$,
 *, $p < 0.05$, n.s, not significant, compared to WT or control, unless otherwise marked on
 graph. All data are presented as means ± SEM, unless otherwise noted. Scale bar = 20μm.

 Figure S2. *yap-1* suppresses structural deformities of *wts-1* mutant neurons. (A) Electron
 microscope images shows that TRNs (ALM) normally dissociate from the muscles (M) and
 innervate the epidermis in the *wts-1* mutants. TRNs are indicated using boxes. Scale bar = 1
 μm. (B–D) Microtubules of TRNs in (B) control wild-type, (C) *wts-1(ok753)* and (D) *wts-*
 1(ok753); yap-1(tm1416). Decreased number of microtubules in *wts-1* mutant ALM was
 restored in *wts-1(ok753); yap-1(tm1416)*. Microtubules are indicated using red numbers and
 the total number of microtubules present in each section is indicated in the upper left area of

each image. Imaging was performed at 1DA. Sections were cut from the head and serial

sections for 5~10 μm of one individual are presented. Scale bar = 500 nm.

Figure S3. Cell-autonomous function of *wts-1*. (A) Representative images of ALM (upper

panels) and PLM (lower panels) of control animals and *wts-1* TRNs-specific-knockdown

animals at 1DA. Both sas1 and sas2 constructs effectively induced structural defects in TRNs.

Ectopic neuronal swelling was labeled with white arrowheads and other structural defects,

including abnormal cell body, shortened process of PLM were labeled with yellow

arrowheads. (B, C) Quantified neuronal defects induced by (B) sas1- or (C) sas2-based

knockdown of *wts-1*. Neurons were observed at 1DA. N=30/one experiment, repeated 3 times.

Statistical significance was determined by a one-way ANOVA, followed by the Dunnett's

multiple comparison test. Scale bar = 20μm.

Figure S4. Paclitaxel treatment fails to mitigate *wts-1* neuronal defects. (A)

Representative images of ALM (upper panels) and PLM (lower panels) of *wts-1* mutants

treated with paclitaxel and untreated control worms. Unlike colchicine treatment, paclitaxel

treatment failed to lessen the touch neuronal deformation of the mutants. (B) Quantified

results of ectopic neuronal lesion scoring. In total, 60 cells were analyzed for each neuron. (C)

Quantified results of the percentage of intact ALM and PLM in the *wts-1(ok753); bus-*

17(br12) under the condition of paclitaxel-treated and untreated worms. Statistical

significance was determined using an unpaired t-test. Ectopic neuronal swelling was labeled

with arrowheads. Scale bar = 20μm.

Figure S5. Colchicine treatment affects touch receptor neurons morphology. (A)

Representative images of TRNs of 8-day-old adult wild-type worms fed with colchicine from

parental generations. Arrowheads indicate highly degenerated touch receptor neurons of F1

progenies. Almost all processes of ALM or PLM degraded and the cell bodies showed exhibit

irregular shape. (B–J) Quantified results of the effect of colchicine on age-related structural

deformations of touch neurons in each category. Colchicine (0.1 mM) was used on worms

from the 1DA and phenotypes were scored at every 2^{nd}, 4^{th}, 6^{th}, 8^{th}, 10^{th}, 15^{th} and 20^{th} day of

adulthood (N =20/one experiment, repeated 3 times). Among the structural deformations,

some features which are particularly associated with age, such as (B) somatic outgrowth and

(C) posterior outgrowth of ALM or (D) swelling and (E) branching of PLM have been

significantly reduced by colchicine-treatment. Colchicine-treatment has only limited effects

on (G) branching of ALM or no effect on (H–J) the other deformations. Statistical differences

were determined using a two-way ANOVA, followed by the Bonferroni's post-test. Asterisks indicate differences from untreated control ALM or PLM. (K, L) Paclitaxel treatment significantly increases neuronal defects of ALM. (K) Representative images of TRNs of control animals and paclitaxel-treated animals at 5DA. (L) Quantified neuronal defects in controls and drug-treated animals (N=30/one experiment, repeated 3times). Statistical significance was determined using an unpaired t-test. Scale bar = 20 μm.

Figure S6. *spas-1* also act cell autonomously to maintain neuronal morphology of TRNs. (A) Both *spas-1* deletion mutants displayed structural decline of TRNs precociously. Neurons were imaged at 6DA. Neuronal abnormalities were labeled with arrowheads. (B) SPAS-1 expressed in TRNs (Red: P*spas-1*:: *spas-1*:: mCherry, Green: P*mec-7*::GFP). Scale bar = 20μm.

Figure S7. Microtubule destabilizing genes, not stabilizing genes, mitigate morphological deformations of *wts-1* mutant neurons. (A, B) Knockdown of genes required for microtubule stabilization, except for *mec-17*, lessen the morphological abnormalities of *wts-1* mutant neurons. In contrast, the knockdown of genes acting as microtubule destabilizers did not affect the *wts-1* neuron. (C) ALM and PLM of the *wts-1(ok753); ptrn-1(tm5597)*. Unlike *dlk-1* or *ptl-1*, the *wts-1; ptrn-1* double mutant failed to recapitulate the knockdown effect of *ptrn-1*. Arrowheads labeled neuronal lesions. (D) Average number of the total ectopic lesions per neuronal process of the *wts-1; ptrn-1* mutant. (A–D) ALM or PLM at the L4 stage was scored. Statistical significances were determined by a one-way ANOVA, followed using the Dunnett's multiple comparison test. Asterisks indicate differences from L4440-fed ALM or PLM (A, B) or the *wts-1* single mutant neurons (D). Scale bar = 20μm.