Association between gene expression plasticity and regulatory network topology

Apolline J. R. Petit[1,*], Anne Genissel[2], and Arnaud Le Rouzic[1]

[1] *Université Paris-Saclay, CNRS, IRD, UMR EGCE, 91190, Gif-sur-Yvette, France*
[2] *Université Paris Saclay, INRAE, UR BIOGER, 91120, Palaiseau, France.*
[*] *Corresponding author: Université Paris-Saclay, CNRS, IRD, UMR EGCE, 91190, Gif-sur-Yvette, France.*
Email: apolline.petit@universite-paris-saclay.fr

Abstract

Over the past two decades, numerous studies have been conducted with the aim of identifying the the structures within regulatory networks responsible for gene expression patterns. Gene expression plasticity for instance has been associated with many network structures, but both theoretical results and empirical observations were often equivocal. Our objective was to decipher the regulatory causes of gene expression plasticity, with a particular focus on identifying the relevant structures in regulatory networks that might influence the sensitivity to environmental factors. We sought the common regulatory structures associated with gene expression plasticity between predictions from an evolutionary simulation model and the global regulatory network from *Escherichia coli*. Congruent results from empirical and theoretical approaches confirmed that selection promotes more regulation towards plastic genes and, as a consequence, plastic genes were more often regulated by feedforward loops than non-plastic genes. Selection tends to bias the distribution of regulatory loop motifs towards positive feedforward and diamond loops, but this enrichment in specific structures was the same in plastic and non-plastic genes. The inability to predict gene expression plasticity from the network regulatory structure opens new questions about the nature of the missing information in current systems biology databases and the evolutionary causes of gene expression plasticity.

Introduction

Understanding the mechanisms involved in the evolution of regulatory networks remains a major challenge in evolutionary and systems biology. It is widely acknowledged that biological gene networks are not random (Shen-Orr *et al.*, 2002; Burda *et al.*, 2011), but the interplay between natural selection, mutation bias, functional constraints, and genetic drift in the evolution of network topologies is not clearly established. Some regulatory structures, such as feedforward loops (FFLs) and feedback loops (FBLs) have attracted attention a lot because of their abundance in biological systems (Mangan and Alon, 2003; Alon, 2007). This enrichment could be the consequence of selection indirectly promoting motifs through their effect on gene expression patterns, including stability, plasticity, or oscillation.

Changes in gene expression is a molecular manifestation of phenotypic plasticity and it conditions the tempo and mode of the organism's response to environmental cues. It is now feasible to measure these changes at a large scale, by identifying differentially-expressed genes using RNA sequencing. At the gene level, plasticity is contingent on signal-sensitive regulation of transcription. Natural or artificial selection for plasticity might indirectly affect the underlying regulatory network structure. For instance, Vlková and Silander, 2022 demonstrated that *E. coli* promoters involved in the metabolism of specific substrate are often selected to maximise plasticity. Yet, both theoretical and empirical work falls short of consensus when linking gene expression patterns with specific network topologies. From the theoretical side, feedforward loops (Shen-Orr *et al.*, 2002; Basu *et al.*, 2004; Ma *et al.*, 2009; de Ronde *et al.*, 2012; Xiong *et al.*, 2019; Wang *et al.*, 2020), diamond loops (DMD) (de Ronde *et al.*, 2012; Xiong *et al.*, 2019), and feedback loops (Hornung and Barkai, 2008; Panovska-Griffiths *et al.*, 2013) have been reported to be involved in plastic (environmentally-sensitive) gene expression. Counterintuitively, other theoretical studies associate those same loop categories with robustness-related functions (FFL: Osella *et al.*, 2011; Kadelka *et al.*, 2013; Roy *et al.*, 2017, DMD: Khammash,

2021, FBL: Kwon and Cho, 2007; Wang *et al.*, 2020). From the empirical side, many studies report observation of these motifs in both plastic and robust systems. Two studies on synthetic bacteria (Basu *et al.*, 2004) and *Escherichia coli* (Sen *et al.*, 2014) brought experimental evidence of a link between feedforward loops and sensitivity to environmental signals. However these studies were restricted to a few genes, and the genome-wide picture remains unknown (Frei and Khammash, 2021; Weldemichael *et al.*, 2022). Overall, there is no clear evidence that specific network structures are involved in the regulation of plastic genes.

The structure of gene regulation can be described at several levels, i.e., the number of regulators, the number and the shape of regulatory loops (feedback, feedforward, or diamond) pointing to a gene, or the loop motif (the arrangement of activation and inhibition patterns within a loop). In regulatory networks, which of these levels — if any — is relevant for gene expression plasticity? Our ambition here is to tackle this question with a dual approach, coupling a theoretical model with a database analysis in order to derive consistent predictions and observations. For this purpose, we wish to (i) quantify the number of regulators and regulatory loops targeting plastic and non-plastic genes, and (ii) determine whether loop categories and loop motifs are relevant to predict plastic gene expression. Plastic genes were collected from 11 transcriptomic studies in which *E. coli* populations were submitted to several environmental conditions. The local regulatory structure of these genes was extracted from a transcriptional regulation database. Theoretical predictions were derived from evolutionary simulations, in which the regulatory interactions were free to evolve under a mutation-drift-selection process in populations. Gene expression plasticity was then measured as the response of individual genes to a environmental signal; both simulated and empirical datasets were analysed within the same statistical framework, allowing for direct comparison.

Results

The *E. coli* regulatory network was built from Ecocyc database (Keseler *et al.*, 2021) (see Fig. suppl. 1 for a graphic representation of the network). Regulations were encoded as qualitative factors ("activating" or "inhibiting"). Independently from the Ecocyc regulation data, a list of environmentally-dependent ("plastic") and independent ("non-plastic") genes was established based on 11 studies reporting differentially-expressed genes in response to various environmental changes. Out of the 4638 genes annotated in the reference annotation (RefSeq: GCF_000005845.2), 2144 genes were never reported as differentially expressed in these 11 studies, and were thus considered as "non-plastic". 1074 genes were differentially expressed in at least two studies, and therefore considered as "plastic" (Suppl. Table 1 and Fig. suppl. 2). The 1420 genes reported only once as differentially expressed were excluded from the analysis.

Computer simulations were performed by coupling a dynamic gene network model for the genotype-phenotype map, and an individual-based model for the population genetics (inspired from Wagner, 1994, 1996). Equilibrium gene expression was assimilated with a multivariate phenotypic trait, on which stabilizing selection was applied. The genotype was encoded as an interaction matrix across 36 genes: 1 sensor gene, which expression level informs the network about the environment, 10 transcription factors, able to affect the transcription of all other genes, 5 neutral genes, 20 selected genes (10 genes selected to be stable, which optimal expression was randomly determined at the beginning of each simulation and independent from the environment, and 10 genes selected to be plastic, which optimal expression depended on the environmental cue, fluctuating randomly every generation). Information about the environment was thus integrated into the gene regulatory network and could affect all gene expressions through cascading regulations. Populations of 20,000 clonal haploid individuals were submitted to 4,000 generations of selection, and the average regulatory network at the end of the simulations was used to reconstruct discrete gene networks, in which each gene-gene interaction could be "activating", "inhibiting", or "null". Genes were classified as "plastic" or "non-plastic" based on the amplitude of their response to the environmental cue (genes with intermediate response were discarded from the dataset). We obtained 33% of plastic genes and 12% of non-plastic genes.

Regulation of plastic *vs.* non-plastic genes

We first considered the number of regulatory interactions towards plastic and non-plastic genes. Plastic genes were influenced by more regulators than non-plastic genes, both in *E. coli* and in simulations

Figure 1: Regulation of plastic and non-plastic genes. A) Distribution of the number of regulations toward the genes. B) Number of feedforward, diamond, and feedback loops by which genes were regulated (only genes regulated by at least one loop are displayed). Simulation results have been downsampled to show the same number of data points in all panels.

(Fig. 1A). In *E. coli*, plastic genes were regulated by 3.2 genes on average (5.4 in simulations), while non-plastic genes were regulated by 1.3 genes on average (2.6 in simulations, to be compared to the strict minimum of 1 regulator necessary to deviate from the arbitrary constitutive expression). The distributions of *E. coli* plastic and non-plastic genes were found significantly different by a t-test (p-value lower than 0.001; statistical tests were conducted on *E. coli* data only, as they are meaningless for simulations, for which the sample size is potentially infinite). The number of feedforward, diamond, and feedback loops targeting plastic genes was also larger than non-plastic genes in both *E. coli* and in simulations, including in control simulations where mutations could accumulate freely with genetic drift (Fig. 1B). In absence of selection (drift simulations), genes that happened to evolve a plastic response were also characterized by a larger number of regulators, suggesting that plasticity tends to be passively induced by a larger number of regulators. In these drift simulations, there were more regulatory loops than in regular simulations for the same number of regulators, suggesting a selection against a too high number of loops.

Regulatory loops

In *E. coli*, 64% of plastic genes (Fig. 2A), and 30% of non-plastic genes (Fig. 2C) were regulated by at least one of the loop categories considered (*i.e.* feedforward, feedback, and diamonds, corresponding to the three loops that have been associated with plasticity in the literature). The most common observed pattern for *E. coli* genes was the regulation by both FFLs and DMDs, with an almost even distribution of both loop categories. Feedback loops (FBLs) were very rare in *E. coli* (68 genes regulated by FBLs out of 3214). Overall, the proportions of genes regulated by FFLs, DMDs, FBLs, and any combination thereof were strikingly similar between plastic and non-plastic genes, and none of these loop categories appeared to be preferentially associated with plastic gene expression.

In simulated gene networks, plastic genes were also more frequently regulated by feedforward, feedback, or diamond loops than non-plastic genes. Most genes were targeted by several categories of loops, and the proportion of FFLs *vs.* DMDs was even (Fig. 2B and D). Overall, there was no clear difference in the relative proportion of FFL and DMD involved in the regulation of plastic *vs.* non-plastic genes. Contrary to our findings for *E. coli*, FBLs were common in simulated networks for plastic genes, and were most often associated with FFL and DMD.

Loop motif frequencies

Loop categories remains a coarse-grained motif classification, as the precise regulatory function necessarily depends on the sign and the coherence of interactions. We have thus determined the motif frequencies within FFL and DMD loops (8 motifs for FFLs, Table 2, and 10 motifs for DMD, Table 3),

Figure 2: Genes classified by incoming regulation loops. A) Plastic genes in *E. coli* B) plastic genes in simulations, C) non-plastic genes in *E. coli*, D) non-plastic genes in simulations. FFL: Feedforward loops, FBL: Feedback loops, DMD: Diamond loops. "None" stands for genes which regulators were unknown, or which were regulated by other motifs (no loops, or large motifs involving more than four genes).

for both the *E. coli* dataset and for simulated networks. Feedback loops were too rare in *E. coli* networks (only 68), and thus not included in Fig. 3A and B.

In both *E. coli* and simulated gene networks, the most frequent feedforward motif was C1 (coherent, two genes activating), followed by I1 (incoherent, one gene activating and the other inhibiting) (Fig. 3A). Other motifs were found in similar proportions, with the exception of I3, frequent in simulations but almost absent in *E. coli*. In diamond loops, the most frequent motifs in both *E. coli* and simulations were PP (activation by both regulating genes) and PM (activation by one gene, inhibition by the other one). The motifs frequency was similar among all the studies used to identify plastic genes (Fig. suppl. 3). Motifs involving negative regulations (NN, NM, and NP) were more frequent in *E. coli* than in simulated networks; but overall, the motif distribution remained quite similar and the frequency of motifs found in *E. coli* and simulations was highly correlated (plastic genes: p-val = 4.3E − 05 ; r = 0.81; nonplastic genes: p-val = 1.7E − 04 ; r = 0.77) (Fig. 3B).

In contrast, the motif frequency distribution resulting from an evolutionary process occurring both in simulations and in *E. coli*, was substantially different from the randomly shuffled networks. In particular, the most frequent motifs (C1 and I3 for FFLs, PP and PM in DMDs) were enriched in evolved networks compared to the shuffled and drift controls. This enrichment was a consequence of selection in simulations, and it is reasonable to postulate that the same explanation holds for *E. coli*.

While there was a clear contrast between control and selected networks, differences in motif composition between plastic and non-plastic genes were minor for *E. coli*. Statistical tests in *E. coli* (Fisher's exact test) did not support any significant differences between plastic and non-plastic genes (p-val > 0.3 for every motif). The contrast between plastic and non-plastic genes was slightly larger in simulations, particularly in the frequency of the FFL motif C1, C2, I3, and the DMD motifs PM. These differences were not consistent with the non-significant trend in E. coli nor with drift simulations, and we deem it unlikely to reflect an inherent property of plastic regulation.

In a Principal Component Analysis on the relative proportion of each motifs of the three loop categories (Fig. 3C), PC1 (55% of the variance) was associated with the contrast between selected and random networks, while PC2 (19% of the variance) was discriminating *E. coli* vs. simulations. Plastic

Figure 3: Loop motifs regulating plastic and non-plastic genes. A) Loop motif frequencies within feedforward (top) and diamond (bottom) loop categories. Motif nomenclature matches Tables 2 and 3. Motif frequencies were normalized per gene (i.e., all genes have the same weight in the figure, irrespective of the number of regulatory motifs targeting them). Shuffled controls were obtained by permuting randomly all non-null interactions. Simulations runs in absence of selection are indicated as "drift". B) Correlation plot of DMD and FFL motifs frequency (featured in panel A) between simulation and *E. coli*. The two regression p-value were < 0.001. C) Principal Component Analysis on loop motif frequencies (FFL, DMD and FBL).

and non-plastic genes from the same conditions were found closer to each other than any other points, reinforcing that the contribution of plasticity to the distribution of motif frequency was minor.

Discussion

Our objective was to identify the gene regulatory structures involved in gene expression plasticity, from both evolutionary simulations and empirical regulation networks. We found that plastic genes indeed had specific regulatory characteristics: genes which expression depends on the environment had more regulators, and were the target of more regulatory loops. The analysis of the frequency of different kinds of regulatory loops and regulatory motifs showed that both real and simulated networks were not random, as a consequence of selection on gene expression levels. Yet, there were no plasticity-specific regulatory motifs, suggesting that environmental sensitivity is not encoded into the network topology.

Theoretical and empirical results strikingly similar Systems biology is rather convincing at describing general patterns in biological networks, based on measurement tools derived from graph theory (Mason and Verwoerd, 2007; Pavlopoulos *et al.*, 2011). For instance, it is often admitted that biological networks tend to be scale-free (Babu *et al.*, 2004), relatively scarce (Espinosa-Soto, 2018), and modular (Wagner *et al.*, 2007). Yet, there is a surprising lack of theory from which such patterns could be interpreted. Typically, the extent to which network evolution is driven by mutation bias, genetic drift, adaptation, or indirect selection remains widely unknown (Tsuda and Kawata, 2010; Halfon, 2017; Petit *et al.*, 2023). Here, we show how a simplified gene network evolutionary model, generally exploited to address genetic architecture-related questions, can predict empirical patterns with a striking accuracy: our simulations could retrieve, both qualitatively and quantitatively, the larger number of regulatory genes for plastic genes, motif enrichment and depletion in selected *vs.* randomized networks, the most abundant motifs, and the relationship between the number of regulation and number of motifs. This validates the capacity of such simple models (in which gene regulations are encoded in an interaction matrix (Wagner, 1994; Fierst, 2011)) to contribute to the theoretical corpus on gene network evolution, in complement to the more cumbersome physiologically-realistic models (Xiong *et al.*, 2019).

Nevertheless, we noticed a few differences between simulations and empirical patterns, that might be worthwhile to explore. The only obvious qualitative difference between both approaches was the ∼10-fold increase in the frequency of feedback loops in simulations compared to *E. coli* regulation network. Although feedback regulation exists in *E. coli* and is known to be involved into noise reduction or bistability (Smits *et al.*, 2006), feedback loops happen to be outnumbered by feedforward loops in the Ecocyc transcriptional dataset and in pathway databases in general (Karp, 2001; Kumar *et al.*, 2018). A possible explanation lies in the fact that most feedback mechanisms in prokaryotes rely on autoregulation (in *E. coli*, 50% of TFs self-regulate, Rosenfeld *et al.*, 2002), while the minimum size for a motif in our analysis was set to 2 genes. A more subtle discrepancy between theoretical and empirical networks is that simulations predicted a small but systematic difference between motif frequencies in plastic vs. non-plastic genes, which was not statistically supported in *E. coli*. Insufficient statistical power could partly explain this observation, but more generally, the classification of *E. coli* genes into plastic and non-plastic categories remained coarse-grained, and may not match the same biological properties: simulations consider the slope of the reaction norm to a unique environmental factor, while the *E. coli* analysis relies on the sensitivity to several stressors, which might not reflect exactly the same mechanisms. The observed convergence between theoretical and empirical observations in spite of these different measurements of plasticity strengthens our conclusions.

Plastic genes have more regulators We identified regulatory differences between plastic genes and non-plastic genes in both real and simulated data. Plastic genes were regulated by more genes, and were the target of more regulatory loops (feedforward loops, diamond loops, and feedback loops). These results reinforces previous observations, which associated the number of regulators with the complexity of the gene expression patterns (Promislow, 2005; Mochizuki, 2008). Nevertheless, the distributions of the number of regulators and the number of regulatory loops largely overlap between plastic and non-plastic genes, both in simulated and in empirical networks. As a consequence, the number of regulators remains a poor predictor of gene expression plasticity.

A direct benefit of using computer simulations to complement the analysis of empirical network lies in the possibility to go beyond association statistics, and to investigate cause-consequence relationships. We showed that (i) plasticity was statistically associated with a higher number of regulators, even in the drift simulations, (ii) selection for plasticity increased the number of regulators, (iii) selection decreased the number of regulatory loops while keeping a high number of regulators. We propose the following scenario for the changes in the regulatory networks associated with the evolution of plasticity: selection for plastic gene expression directly promotes the recruitment of cis-regulators, which indirectly increases the number of regulatory loops.

If some regulatory motifs were outstandingly more present in plastic genes than non-plastic genes, it would have been possible to confidently associate a plastic function to those motifs. However, no differences between the motifs regulating plastic genes and the motifs regulating non-plastic genes were found in *E. coli*. Yet loop motifs were non-randomly distributed, and our simulations confirmed that this pattern was due to selection on gene expression. The selection pressure on motifs thus appears to be identical for plastic and non-plastic genes. Given our findings and the conflicting literature on this topic, we find it likely that there is no regulation motifs with a dedicated "plasticity" function.

Motifs evolve due to indirect selection, i.e., as a consequence of selection on gene expression. The *E. coli* motif distribution we found was coherent with previous work (Mangan *et al.*, 2006; Cooper *et al.*, 2008), who also identified C1 and I1 as the most represented motifs. The high frequency of activating motifs like C1 for FF loops or PP for DMD loops could be a statistical consequence of the uneven balance between activation and inhibition within the network. C1 and PP are indeed very rare in drift simulations, featured by an equal number of activation and inhibition interactions. Yet, arguments related to activation/inhibition ratios are fragile, because it is difficult to discard that the uneven activation/inhibition balance could be a consequence, rather than a cause, of an evolutionary bias towards specific motifs. In any case, both C1 and PP were over-represented in *E. coli* compared to randomly-permuted networks, suggesting some specific preference for such motifs. An attractive hypothesis can be derived from the fact that C1 and I1 have been predicted to be efficient at buffering expression noise (Ghosh *et al.*, 2005; Osella *et al.*, 2011; Reeves, 2019; Chakravarty and Csikász-Nagy, 2021). Selection on noise filtering and gene expression robustness could thus enrich gene networks in some specific motifs like C1 and I1. This hypothesis could not be confirmed by our simulations, in which the gene network dynamics was deterministic, but alternative theoretical settings (Draghi and Whitlock, 2015; Schmutzer and Wagner, 2020) could be considered.

Conclusion   Our analysis indicated a higher prevalence of regulation toward plastic genes. Furthermore, the previously observed correlation between plasticity and the number of loop motifs was found to be a consequence of this association. Additionally, we revealed the limitations of regulatory network motifs in explaining environment-sensitive gene expression. We observed no topological feature associated with gene expression plasticity beyond the larger number of incoming regulations on average. Therefore, the plasticity of a given gene cannot be deduced solely from the properties of its local regulation network.

If not the network motif, which regulatory features could reveal whether the expression of a specific gene has been selected for robustness, stability, cyclic expression, or sensitivity to an environmental or physiological input? To what extent is it possible to predict gene expression profiles from pathway databases (Mercatelli *et al.*, 2020; Escorcia-Rodríguez *et al.*, 2023)? Three major hypotheses can be proposed: (i) the information about regulatory structures involved in plasticity is present in existing databases, but they are not caught by known descriptors of network structures and properties (loops, motifs, or graph metrics). For instance, it might depend on a more general regulatory context (Lipshtat *et al.*, 2008; Jiang and Hao, 2021), or on complex combinations of several intertwined network structures (Chakravarty and Csikász-Nagy, 2021). Alternatively, (ii) regulation databases remain the correct approach, but existing databases are not precise nor exhaustive enough. Typically, presence/absence data for regulatory interactions might be insufficient, and quantitative estimation of interaction strengths (weighted edges in the network graph (Freyre-González *et al.*, 2022)) could be necessary as suggested by others (Ingram *et al.*, 2006; Panovska-Griffiths *et al.*, 2013; Kim and Wysocka, 2023). In contrast, hypothesis (iii) presumes that regulation might be too complex to be abstracted in regulatory networks, which omit important epigenetic and cellular mechanisms, including the spatial organisation of the genome, methylation, phosphorylation, dimerization (Klumpe *et al.*, 2023), protein conformation, or "missing regulation" from intergenic sequences (Connally *et al.*, 2022).

265 The next challenge for the systems biology community may therefore not be to generate more data,
266 but to identify which data is relevant to understand the evolutionary properties of biological systems,
267 a task for which theoretical approaches might bring invaluable insights.

268 Material and Methods

269 Empirical data: *E. coli*

We used *E. coli* K-12 substr. MG1655 annotation (RefSeq: GCF_000005845.2). *E. coli* K-12 known
regulations have been obtained from Ecocyc (Keseler *et al.*, 2021, downloaded 29/01/2024). This
database contained 9534 regulatory interactions.

To conduct our analyses, we established a list of "plastic" genes, to be contrasted to "non-plastic"
genes. We selected 11 studies explicitly testing environmental conditions on gene expression in *E.*
coli, and proposing lists of differentially-expressed genes. The environmental factors were pH (Tucker
et al., 2002; Maurer *et al.*, 2005), growth medium (urine, gluconate, and lysogeny broth) (Feugeas *et al.*,
2016), aerosolization and liquid suspension (Ng *et al.*, 2018), exposure to apple juice (Bergholz *et al.*,
2009), magnesium stress, sodium stress and carbon source (Caglar *et al.*, 2017), temperature (White-
Ziegler *et al.*, 2007; Kim *et al.*, 2020), metabolic stress (low pH, low oxygen, low nutrients) (Bhatia
et al., 2022), oxydative stress (Wang *et al.*, 2009), and amino acid starvation (Durfee *et al.*, 2008).
Genes were considered as plastic when reported as differentially expressed in at least two independent
studies. Genes were considered as "non-plastic" if they were never reported as differentially expressed
in these studies.

284 Gene regulatory network model

We used a quantitative gene regulatory network model, inspired from A. Wagner's seminar work on
gene expression robustness and plasticity (Wagner, 1994, 1996). This setting, coupling a traditional
population genetics model with a gene network model as a genotype-to-phenotype map, was particu-
larly popular to explore the evolution of complex genetic architectures (Draghi and Wagner, 2008; Ng
and Kinjo, 2023; Petit *et al.*, 2023).

Gene network The gene network model was dynamic with discrete time-steps. Genotype consisted
in $n = 36$ genes encoding a gene network represented by a $n \times n$ gene regulatory matrix $\mathbf{W}$. Genes
were expressed quantitatively between $P = 0$ (no expression) and $P = 1$ (maximum expression).
Gene expression was computed dynamically: initial gene expressions were set to their basal level
(constitutive expression) $\kappa = 0.15$, representing the gene expression in absence of regulation, and
updated as a function of the concentration of the other genes in the network during 24 time steps:

$$\mathbf{P}_{t+1} = F(\mathbf{W}\mathbf{P}_t), \tag{1}$$

where $\mathbf{P}$ was a vector of n quantitative gene expressions scaled between 0 and 1 by a multivariate
sigmoid function $F(x_1, \ldots, x_n) = (f(x_1), \ldots, f(x_n))$, with $f(x) = [1 + (1/\kappa - 1)\exp(-x/\kappa(1-\kappa))]^{-1}$
as in (Rünneburger and Le Rouzic, 2016).

Individuals were characterized by the average expression of their n genes during the four last
time-steps $\bar{\mathbf{P}} = [\bar{P}_1, \ldots, \bar{P}_i, \ldots, \bar{P}_n]$, with $\bar{P}_i = \sum_{t=21}^{24} P_{it}$.

Reproduction, mutation, and selection Populations consisted in 20,000 haploid, clonally-
reproducing individuals, matching approximately *E. coli*'s life cycle. Simulations were carried on
for 4,000 non-overlapping generations.

The genotype $\mathbf{W}$ was affected by mutations of the regulation strength W_{ij}, at a rate of $\mu = 0.15$
per individual. Each gene had the same probability μ/n to be affected by a mutation. A mutated gene
i had a random regulation W_{ij} changed by a Gaussian random number of standard deviation 0.05.
This setting (many mutations of small effect) ensured that the distribution of alleles at all loci was
approximately Gaussian.

Successive generations were produced by drawing clones with a probability proportional to their
relative fitness. The individual fitness w was the product of two components: $w = w_e \times w_s$. The first
term w_e was determined by a stabilizing bell-shaped fitness function and decreased with the distance to

the gene expression optimum $\boldsymbol{\theta}$ (Lande, 1979): $w_e = \exp(-\sum_{i=1}^{n} s_i(\overline{P}_i - \theta_i)^2)$. Unless specified otherwise, the selection coefficient was $s_i = 0$ for "unselected" genes, and $s_i = 10$ for "selected" genes. The second term w_s corresponds to the selection for network stability, calculated as $w_s = \exp(-s' \sum_i^n V_i)$, where s' quantifies the selection against unstable gene expression, and $V_i = (1/4)\sum_{t=21}^{24}(P_{it} - \overline{P}_i)^2$ was the variance in the expression of gene i during the last 4 steps of the network dynamics. We set $s' = 46,000$, as in (Rünneburger and Le Rouzic, 2016; Petit *et al.*, 2023), which was a large penalty, reflecting the common idea that unstable networks are not viable (Siegal and Bergman, 2002).

Fluctuating selection and plasticity In our simulations, the environment was modelled as a numerical value E_g changing randomly every generation g, drawn from an uniform distribution $U(0.15, 0.85)$. The state of the environment was transmitted to the network via a sensor gene, as in (Draghi and Whitlock, 2012; Vadée-Le-Brun *et al.*, 2015). The model implementation follows Burban *et al.*, 2022, the expression of the sensor gene was constant $P_s = E_g$ during all the network dynamics. The sensor gene was free to evolve regulations toward any genes of the network, but it was not regulated, as its expression depends solely of the environment.

The other 35 network genes were divided in 4 categories. 10 genes were "Transcription factors" not subjected to direct selection; 10 "Plastic Target Genes" were directly selected to have their expression correlated to the environment; 10 "Non-plastic Target Genes" were directly selected to have a constant expression; and 5 unselected "control genes" were let free to evolve (Table 1).

Selection on plastic genes was modelled by changing the gene expression optimum every generation g as a function of the environment E_g, as $\theta_i = a_i E_g + b_i$. The intercept b_i and the slope a_i were randomly drawn for each gene i at the beginning of each simulation under the constraint that gene expression had to remain between 0 and 1 for any $E_g \in (0.15, 0.85)$, and stayed constant thorough generations. "Non-plastic" genes had their constant optimum drawn from a uniform distribution $U(0.15, 0.85)$.

The reaction norm is the function describing the trait response to a quantitative environmental variation. We measured the expression plasticity of a gene i as the slope β_i of the linear regression between a set of environments and the corresponding gene expression $\bar{P}_i$. Simulated genes were classified into the category "plastic genes" and "non-plastic genes" according to their reaction norm slope. Genes with $\beta_i \geq 0.5$ were considered as plastic and genes with $\beta_i \leq 0.025$ as non-plastic. Genes outside of these categories were not kept during the analysis.

	Sensor gene	Transcription factor (TF)	Plastic Target gene	Constant Target gene	Neutral gene
Selection s_i	0	0	10	10	0
Optimum θ_i			$a_i E_g + b_i$	b_i	
Evolution	Environment only	Free	Towards plasticity	Towards robustness	Drift
Regulating	All genes	All genes but sensor	TF	TF	None
Regulated by	None	All genes but Neutral	TF and Sensor	TF and Sensor	TF and Sensor

Table 1: Description of the five gene categories implemented in the gene regulatory network model.

Analysis

We used *E. coli* Ecocyc regulation data to construct a discrete 4638-gene network as a 4638×4638 square regulatory matrix **Ec**. A cell $\mathbf{Ec}_{ij} \in \{-1, 0, 1\}$ represents the regulation of gene j on i. Regulations that were both activating and inhibiting in the Ecocyc dataset were considered as null.

To study the regulatory motifs responsible for plasticity in the simulated networks with the same pipeline used for *E. coli*, we discretized the regulation matrix **W** into a matrix **Et**, based on the concept of "essentiality" (Burda *et al.*, 2011). We tested one by one every element of the **W** matrix by changing their value to 0. If it changed the reaction norm slope or the intercept by more than 0.01, then the regulation was considered as "essential": $\mathbf{Et}_{ij} = \text{sign}(\mathbf{W}_{ij})$, and $\mathbf{Et}_{ij} = 0$ otherwise.

	X → Y → Z	X → Y → Z	X → Y → Z	X → Y → Z
Coherent				
Nomenclature Abbreviation	Coherent 1 C1	Coherent 2 C2	Coherent 3 C3	Coherent 4 C4
Incoherent				
Nomenclature Abbreviation	Incoherent 1 I1	Incoherent 2 I2	Incoherent 3 I3	Incoherent 4 I4

Table 2: Three-genes feedforward loop nomenclature.

$X \to Y_n$ Sign	Inhibition	Activation	Mixed
Negative $Y_n \to Z$			
Nomenclature Abbreviation	Negative negative NN	Positive negative PN	Mixed negative MN
Positive $Y_n \to Z$			
Nomenclature Abbreviation	Negative positive NP	Positive positive PP	Mixed positive MP
Mixed $Y_n \to Z$			
Nomenclature	Negative mixed	Positive mixed	Mixed mixed 1 and 2
Abbreviation	NM	PM	MM1 ; MM2

Table 3: Diamond motifs nomenclature.

Motifs We investigated three loop categories: feedforward loops (corresponding to 8 unique motifs, Table 2), diamonds loops (10 unique motifs, Table 3), and feedbacks loops (of 4 different sizes). The nomenclature of the FFLs was based on (Mangan and Alon, 2003), the genes under consideration hold the position Z (regulated gene) in the motifs. FFLs were divided into two groups, coherent and incoherent. The coherence of a motif was defined by the product of all regulations of the motif: coherent if the product was positive, incoherent otherwise (Table 2). The nomenclature of diamonds was defined by the the signs of the $X \to Yn$ regulations (positive, negative, or mixed) for the first part of the name and the sign of $Yn \to Z$ for the second part (Table 3). Feedback loops were divided into subcategories by size, from 2 to 5 genes.

Control datasets In drifting populations, populations of 200 individuals evolved for 20000 generations with a mutation rate of $\mu = 0.4$. "Shuffled" controls were obtained by randomly shuffling the

values of non-null regulations, the number of regulations and loops were thus kept, but not the nature of loop motifs.

Availability and reproducibility

The simulation program was coded in C++, and is available at https://github.com/lerouzic/simevolv. The program was compiled and executed on the IFB-core cluster (https://www.france-bioinformatique.fr/en/ifb-core-cluster/). Simulation pipelines (in R and Bash) and analysis scripts (in R) are available at https://github/apetit/topoplast.

Aknowledgments

Simulations were performed on the Core Cluster of the Institut Français de Bioinformatique (IFB) (ANR-11-INBS-0013). AP was supported by the doctorate school SDSV (ED 577, Université Paris-Saclay). The authors aknowledge the support of the French Agence Nationale de la Recherche (ANR), under grant ANR-22-CE02-0026-01 (project EVOPLANET). The authors declare no conflict of interest.

References

Alon U. 2007. Network motifs: theory and experimental approaches. Nature Reviews Genetics. 8:450.

Babu MM, Luscombe NM, Aravind L, Gerstein M, Teichmann SA. 2004. Structure and evolution of transcriptional regulatory networks. Current opinion in structural biology. 14:283–291.

Basu S, Mehreja R, Thiberge S, Chen MT, Weiss R. 2004. Spatiotemporal control of gene expression with pulse-generating networks. Proceedings of the National Academy of Sciences. 101:6355–6360.

Bergholz TM, Vanaja SK, Whittam TS. 2009. Gene expression induced in *Escherichia coli* O157:H7 upon exposure to model apple juice. Applied and Environmental Microbiology. 75:3542–3553.

Bhatia RP, Kirit HA, Predeus AV, Bollback JP. 2022. Transcriptomic profiling of *Escherichia coli* K-12 in response to a compendium of stressors. Scientific Reports. 12:1–14.

Burban E, Tenaillon MI, Le Rouzic A. 2022. Gene network simulations provide testable predictions for the molecular domestication syndrome. Genetics. 220:iyab214.

Burda Z, Krzywicki A, Martin OC, Zagorski M. 2011. Motifs emerge from function in model gene regulatory networks. Proceedings of the National Academy of Sciences of the United States of America. 108:17263–17268.

Caglar MU, Houser JR, Barnhart CS, Boutz DR, Carroll SM, Dasgupta A, Lenoir WF, Smith BL, Sridhara V, Sydykova DK *et al*. 2017. The *E. coli* molecular phenotype under different growth conditions. Scientific Reports. 7:1–15.

Chakravarty S, Csikász-Nagy A. 2021. Systematic analysis of noise reduction properties of coupled and isolated feedforward loops. PLoS Computational Biology. 17:1–15.

Connally NJ, Nazeen S, Lee D, Shi H, Stamatoyannopoulos J, Chun S, Cotsapas C, Cassa CA, Sunyaev SR. 2022. The missing link between genetic association and regulatory function. eLife. 11:1–35.

Cooper MB, Loose M, Brookfield JF. 2008. Evolutionary modelling of feed forward loops in gene regulatory networks. BioSystems. 91:231–244.

de Ronde WH, Tostevin F, ten Wolde PR. 2012. Feed-forward loops and diamond motifs lead to tunable transmission of information in the frequency domain. Physical Review E. 86:021913.

Draghi J, Wagner GP. 2008. Evolution of evolvability in a developmental model. Evolution: International Journal of Organic Evolution. 62:301–315.

Draghi J, Whitlock M. 2015. Robustness to noise in gene expression evolves despite epistatic constraints in a model of gene networks. Evolution. 69:2345–2358.

Draghi JA, Whitlock MC. 2012. Phenotypic plasticity facilitates mutational variance, genetic variance, and evolvability along the major axis of environmental variation. Evolution: International Journal of Organic Evolution. 66:2891–2902.

Durfee T, Hansen AM, Zhi H, Blattner FR, Ding JJ. 2008. Transcription profiling of the stringent response in *Escherichia coli*. Journal of Bacteriology. 190:1084–1096.

Escorcia-Rodríguez JM, Gaytan-Nuñez E, Hernandez-Benitez EM, Zorro-Aranda A, Tello-Palencia MA, Freyre-González JA. 2023. Improving gene regulatory network inference and assessment: The importance of using network structure. Frontiers in Genetics. 14:1–14.

Espinosa-Soto C. 2018. On the role of sparseness in the evolution of modularity in gene regulatory networks. PLoS computational biology. 14:e1006172.

Feugeas JP, Tourret J, Launay A, Bouvet O, Hoede C, Denamur E, Tenaillon O. 2016. Links between Transcription, Environmental Adaptation and Gene Variability in *Escherichia coli*: Correlations between Gene Expression and Gene Variability Reflect Growth Efficiencies. Molecular Biology and Evolution. 33:2515–2529.

Fierst JL. 2011. A history of phenotypic plasticity accelerates adaptation to a new environment. Journal of evolutionary biology. 24:1992–2001.

Frei T, Khammash M. 2021. Adaptive circuits in synthetic biology. Current Opinion in Systems Biology. 28:100399.

Freyre-González JA, Escorcia-Rodríguez JM, Gutiérrez-Mondragón LF, Martí-Vértiz J, Torres-Franco CN, Zorro-Aranda A. 2022. System Principles Governing the Organization, Architecture, Dynamics, and Evolution of Gene Regulatory Networks. Frontiers in Bioengineering and Biotechnology. 10.

Ghosh B, Karmakar R, Bose I. 2005. Noise characteristics of feed forward loops. Physical Biology. 2:36–45.

Halfon MS. 2017. Perspectives on Gene Regulatory Network Evolution. Trends in Genetics. 33:436–447.

Hornung G, Barkai N. 2008. Noise Propagation and Signaling Sensitivity in Biological Networks: A Role for Positive Feedback. PLoS Computational Biology. 4:e8.

Ingram PJ, Stumpf MP, Stark J. 2006. Network motifs: structure does not determine function. BMC Genomics. 7:108.

Jiang Y, Hao N. 2021. Memorizing environmental signals through feedback and feedforward loops. Current Opinion in Cell Biology. 69:96–102.

Kadelka C, Murrugarra D, Laubenbacher R. 2013. Stabilizing gene regulatory networks through feed-forward loops. Chaos: An Interdisciplinary Journal of Nonlinear Science. 23:1–19.

Karp PD. 2001. Pathway databases: a case study in computational symbolic theories. Science. 293:2040–2044.

Keseler IM, Gama-Castro S, Mackie A, Billington R, Bonavides-Martínez C, Caspi R, Kothari A, Krummenacker M, Midford PE, Muñiz-Rascado L *et al*. 2021. The ecocyc database in 2021. Frontiers in Microbiology. 12.

Khammash MH. 2021. Perfect adaptation in biology. Cell Systems. 12:509–521.

Kim S, Kim Y, Suh DH, Lee CH, Yoo SM, Lee SY, Yoon SH. 2020. Heat-responsive and time-resolved transcriptome and metabolome analyses of *Escherichia coli* uncover thermo-tolerant mechanisms. Scientific Reports. 10:1–13.

Kim S, Wysocka J. 2023. Deciphering the multi-scale, quantitative cis-regulatory code. Molecular Cell. 83:373–392.

Klumpe HE, Garcia-Ojalvo J, Elowitz MB, Antebi YE. 2023. The computational capabilities of many-to-many protein interaction networks. Cell Systems. 14:430–446.

Kumar S, Mahajan S, Jain S. 2018. Feedbacks from the metabolic network to the genetic network reveal regulatory modules in E. coli and B. subtilis. PLOS ONE. 13:e0203311.

Kwon YK, Cho KH. 2007. Analysis of feedback loops and robustness in network evolution based on Boolean models. BMC Bioinformatics. 8:430.

Lande R. 1979. Quantitative genetic analysis of multivariate evolution, applied to brain: body size allometry. Evolution. 33:402–416.

Lipshtat A, Purushothaman SP, Iyengar R, Ma'ayan A. 2008. Functions of bifans in context of multiple regulatory motifs in signaling networks. Biophysical Journal. 94:2566–2579.

Ma W, Trusina A, El-Samad H, Lim WA, Tang C. 2009. Defining Network Topologies that Can Achieve Biochemical Adaptation. Cell. 138:760–773.

Mangan S, Alon U. 2003. Structure and function of the feed-forward loop network motif. Proceedings of the National Academy of Sciences of the United States of America. 100:11980–11985.

Mangan S, Itzkovitz S, Zaslaver A, Alon U. 2006. The incoherent feed-forward loop accelerates the response-time of the gal system of *Escherichia coli*. Journal of Molecular Biology. 356:1073–1081.

Mason O, Verwoerd M. 2007. Graph theory and networks in biology. IET systems biology. 1:89–119.

Maurer LM, Yohannes E, Bondurant SS, Radmacher M, Slonczewski JL. 2005. pH regulates genes for flagellar motility, catabolism, and oxidative stress in *Escherichia coli* K-12. Journal of Bacteriology. 187:304–319.

Mercatelli D, Scalambra L, Triboli L, Ray F, Giorgi FM. 2020. Gene regulatory network inference resources: A practical overview. Biochimica et Biophysica Acta - Gene Regulatory Mechanisms. 1863:194430.

Mochizuki A. 2008. Structure of regulatory networks and diversity of gene expression patterns. Journal of Theoretical Biology. 250:307–321.

Ng ETH, Kinjo AR. 2023. Plasticity-led evolution as an intrinsic property of developmental gene regulatory networks. Scientific Reports. 13:1–19.

Ng TW, Ip M, Chao CY, Tang JW, Lai KP, Fu SC, Leung WT, Lai KM. 2018. Differential gene expression in *Escherichia coli* during aerosolization from liquid suspension. Applied Microbiology and Biotechnology. 102:6257–6267.

Osella M, Bosia C, Corá D, Caselle M. 2011. The Role of Incoherent MicroRNA-Mediated Feedforward Loops in Noise Buffering. PLoS Computational Biology. 7:e1001101.

Panovska-Griffiths J, Page KM, Briscoe J. 2013. A gene regulatory motif that generates oscillatory or multiway switch outputs. Journal of the Royal Society Interface. 10.

Pavlopoulos GA, Secrier M, Moschopoulos CN, Soldatos TG, Kossida S, Aerts J, Schneider R, Bagos PG. 2011. Using graph theory to analyze biological networks. BioData mining. 4:1–27.

Petit AJR, Guez J, Le Rouzic A. 2023. Correlated stabilizing selection shapes the topology of gene regulatory networks. Genetics. 224. iyad065.

Promislow D. 2005. A regulatory network analysis of phenotypic plasticity in yeast. The American Naturalist. 165:515–523.

Reeves GT. 2019. The engineering principles of combining a transcriptional incoherent feedforward loop with negative feedback. Journal of Biological Engineering. 13:62.

Rosenfeld N, Elowitz MB, Alon U. 2002. Negative autoregulation speeds the response times of transcription networks. Journal of molecular biology. 323:785–793.

Roy S, Raj M, Ghosh P, Das SK. 2017. Role of motifs in topological robustness of gene regulatory networks. IEEE International Conference on Communications. 1.

Rünneburger E, Le Rouzic A. 2016. Why and how genetic canalization evolves in gene regulatory networks. BMC evolutionary biology. 16:239.

Schmutzer M, Wagner A. 2020. Gene expression noise can promote the fixation of beneficial mutations in fluctuating environments. PLoS computational biology. 16:e1007727.

Sen S, Kim J, Murray RM. 2014. Designing robustness to temperature in a feedforward loop circuit. In: . volume 2015-Febru. pp. 4629–4634. IEEE.

Shen-Orr SS, Milo R, Mangan S, Alon U. 2002. Network motifs in the transcriptional regulation network of *Escherichia coli*. Nature Genetics. 31:64–68.

Siegal ML, Bergman A. 2002. Waddington's canalization revisited: developmental stability and evolution. Proceedings of the National Academy of Sciences. 99:10528–10532.

Smits WK, Kuipers OP, Veening JW. 2006. Phenotypic variation in bacteria: the role of feedback regulation. Nature Reviews Microbiology. 4:259–271.

Tsuda ME, Kawata M. 2010. Evolution of gene regulatory networks by fluctuating selection and intrinsic constraints. PLoS computational biology. 6:e1000873.

Tucker DL, Tucker N, Conway T. 2002. Gene expression profiling of the pH response in *Escherichia coli*. Journal of Bacteriology. 184:6551–6558.

Vadée-Le-Brun Y, Rouzaud-Cornabas J, Beslon G. 2015. Epigenetic inheritance speeds up evolution of artificial organisms. In: . pp. 439–446. MIT Press.

Vlková M, Silander OK. 2022. Gene regulation in Escherichia coli is commonly selected for both high plasticity and low noise. Nature Ecology & Evolution. 6:1165–1179.

Wagner A. 1994. Evolution of gene networks by gene duplications: a mathematical model and its implications on genome organization. Proceedings of the National Academy of Sciences. 91:4387–4391.

Wagner A. 1996. Genetic redundancy caused by gene duplications and its evolution in networks of transcriptional regulators. Biological cybernetics. 74:557–567.

Wagner GP, Pavlicev M, Cheverud JM. 2007. The road to modularity. Nature Reviews Genetics. 8:921–931.

Wang J, Belta C, Isaacson SA. 2020. How Retroactivity Affects the Behavior of Incoherent Feedforward Loops. iScience. 23:101779.

Wang S, Deng K, Zaremba S, Deng X, Lin C, Wang Q, Tortorello ML, Zhang W. 2009. Transcriptomic response of *Escherichia coli* O157:H7 to oxidative stress. Applied and Environmental Microbiology. 75:6110–6123.

Weldemichael T, Asemoloye MD, Marchisio MA. 2022. Feedforward Loops: Evolutionary Conserved Network Motifs Redesigned for Synthetic Biology Applications. Applied Sciences. 12:8292.

White-Ziegler CA, Malhowski AJ, Young S. 2007. Human body temperature (37°C) increases the expression of iron, carbohydrate, and amino acid utilization genes in *Escherichia coli* K-12. Journal of Bacteriology. 189:5429–5440.

Xiong K, Lancaster AK, Siegal ML, Masel J. 2019. Feed-forward regulation adaptively evolves via dynamics rather than topology when there is intrinsic noise. Nature Communications. 10:2418.