The Nuclear Receptor NR4A3 Regulates Glucose Metabolism and Translation in Skeletal Muscle

Jonathon A.B. Smith[1], Brendan M. Gabriel[1], Anna Krook[1], Juleen R. Zierath[1,2], Nicolas J. Pillon[1*]

[1]Karolinska Institutet, Department of Physiology and Pharmacology, Stockholm, Sweden
[2]Karolinska Institutet, Department of Molecular Medicine and Surgery, Stockholm, Sweden

[*]For correspondence: Nicolas J. Pillon, nicolas.pillon@ki.se

Background: The transcription factor nuclear receptor subfamily 4 group A member 3 (NR4A3) is a pleiotropic regulator of skeletal muscle metabolism. NR4A3 expression is altered in response to changes in physical activity and associated with metabolic adaptations. However, the molecular mechanisms and consequences of altered NR4A3 expression remains largely unexplored. Here, we investigated the impact of NR4A3 on human skeletal muscle remodelling.

Methods: Published transcriptomic datasets from human bed rest and limb immobilisation studies were curated to meta-analyse the effect of physical inactivity on skeletal muscle NR4A3 levels. In primary human skeletal muscle myotubes, siRNA and lentivirus were used to silence and overexpress NR4A3, respectively. Basal and stimulated (insulin ± leucine) signal transduction was determined by immunoblot analysis. Effects on glucose, fatty acid, and protein metabolism were measured using radiolabelled substrate assays. Lactate production was assessed in culture supernatant by colourimetry. Cell morphology was analysed by immunocytochemistry and gene expression was quantified by RT-qPCR.

Results: Physical inactivity decreased skeletal muscle NR4A3 (-27%), concomitant with pathways related to mitochondrial function, cytoskeleton organization, chromatin regulation, protein synthesis and degradation. Silencing of nuclear-localised NR4A3 protein (-38%) in cultured myotubes decreased glucose oxidation (-18%) and increased lactate production (+23%) in vitro. This coincided with greater signalling downstream of AMPK and elevated rates of basal (+26%) and FCCP-stimulated (+55%) fatty acid oxidation. While NR4A3 silencing reduced protein synthesis (-25%), and mTORC1 signalling, overexpression of the canonical NR4A3 protein isoform (+290%) augmented translation and total cellular protein content. Moreover, partial restoration of NR4A3 levels rescued glucose oxidation in NR4A3-silenced myotubes and restored phosphorylation of mTORC1 substrates. NR4A3 depletion reduced myotube area (-48%) and further altered protein and mRNA expression of contractile elements in skeletal myotubes.

Conclusions: NR4A3 is central to the regulation of protein synthesis by directing mTORC1 signalling and ribosomal biogenesis in muscle. Downregulation of NR4A3 has adverse effects on metabolism, myotube size and contractile proteins, consistent with a physical inactivity/disuse phenotype. Thus, decrements in NR4A3 abundance may contribute to the deleterious health consequences resulting from sedentary lifestyles. Therapeutic targeting of NR4A3 could offer new avenues for combating metabolic dysfunction associated with skeletal muscle disuse atrophy.

Keywords Skeletal muscle, inactivity, metabolism, mitochondria, protein synthesis

Introduction

Engagement in physical activity initiates a cascade of integrative responses related to skeletal muscle contraction, such as increased energy expenditure, haemodynamics, and oxygen consumption (Smith, Murach et al., 2023). Depending on the modality, intensity, and volume of work performed, these homeostatic processes trigger a myriad of beneficial downstream events that serve to maintain or enhance aspects of human physiology, including cardiovascular function, skeletal muscle mass, and peripheral insulin sensitivity (Smith et al., 2023). Conversely, inactivity impairs skeletal muscle size and composition. These alterations have adverse effects on local tissue function and metabolism, which in turn impacts systemic health. Inactivity diminishes physical performance (Brook, Stokes et al., 2022, Manini, Clark et al., 2007), whole-body metabolic flexibility (Bergouignan, Antoun et al., 2013), and is associated with reduced quality of life (Yerrakalva, Hajna et al., 2023) and increased risk of non-communicable diseases like type 2 diabetes (Yuan, Li et al., 2023).

Changes in skeletal muscle mass are determined by rates of myofibrillar protein synthesis versus degradation. Physical inactivity favours signalling and transcriptional programmes that promote sustained periods of negative protein balance and muscle loss. Indeed, detectable muscle atrophy occurs within just four days of disuse (Brook et al., 2022, Shur, Simpson et al., 2024) and involves a complex molecular interplay. One of the major regulators of translation is the mammalian target of rapamycin complex 1 (mTORC1), which permits myofibrillar protein synthesis and muscle hypertrophy (You, McNally et al., 2019). However, mTORC1 acts in tandem with ribosomal content (among other processes) to regulate translation (West, Baehr et al., 2016). Detriments in both mTORC1 (Shad, Thompson et al., 2019) and ribosomal abundance (Figueiredo, D'Souza et al., 2021) are observed in skeletal muscle with inactivity. Alongside insulin- and amino acid-resistance (Breen, Stokes et al., 2013, Brook et al., 2022, Pavis, Abdelrahman et al., 2023, Shad et al., 2019, Shur et al., 2024), this downregulation of

translational machinery drives initial losses in skeletal muscle mass through dampened myofibrillar protein synthetic responses (Brook et al., 2022, Pavis et al., 2023). As an effector of the phosphatidylinositol-3-kinase (PI3K)-AKT pathway, mTORC1 is also central to proteostasis via temporal coordination of anabolic and catabolic processes (Kaiser, Milan et al., 2022). Nevertheless, the molecular transducers of this inactivity phenotype have not been fully elucidated.

Exercise training and physical inactivity are not mere opposing ends of a linear spectrum, but rather multifaceted processes with distinct underlying mechanisms. Understanding the dichotomy between exercise and inactivity necessitates recognising that they engage both common and distinct signalling and transcriptional pathways (Pillon, Gabriel et al., 2020). The NR4A family of stress-responsive orphan nuclear receptors are commonly induced by exercise. However, only *NR4A3* (also known as *NOR1*) shows opposite regulation to exercise and inactivity (Pillon et al., 2020). Notably, all members of the NR4A family have ties to skeletal muscle metabolism, development, or remodelling. Depletion of NR4A3 in primary human (Pillon et al., 2020) or mouse C2C12 (Paez, Ferrandi et al., 2023) skeletal muscle myotubes reduces mitochondrial oxidative capacity. Alternatively, transgenic overexpression of *Nr4a3* (Pearen, Eriksson et al., 2012) or *Nr4a1* (also known as *Nur77*) (Chao, Wroblewski et al., 2012) shifts skeletal muscle composition towards an oxidative phenotype, suggesting functional redundancy between these NR4A family members through shared NGFI-B (Wilson, Fahrner et al., 1991) or Nur (Maira, Martens et al., 1999) response elements. Furthermore, overexpression of *Nr4a3* (Pearen et al., 2012), *Nr4a1* (Chao et al., 2012), or *Nr4a2* (also known as *Nurr1*) (Amoasii, Sanchez-Ortiz et al., 2019) in skeletal muscle enhances physical performance, while deletion of *Nr4a1* impairs muscle size (Tontonoz, Cortez-Toledo et al., 2015). Silencing of *Nr4a3* in mouse C2C12 myotubes reduces mTORC1 signalling and global myosin heavy chain gene expression (Paez et al., 2023) but the molecular mechanisms underlying the metabolic, anabolic, and transcriptomic roles of NR4A3 remain largely unknown in human skeletal muscle.

Here, we investigated the impact of NR4A3 on metabolism and post-mitotic growth processes. We provide evidence that *NR4A3* is reduced during skeletal muscle disuse and that downregulation of *NR4A3* impairs glucose metabolism and protein synthesis, contributing towards atrophy of skeletal muscle cells.

Results and Discussion

Downregulation of *NR4A3* during inactivity correlates with transcripts involved in myogenic and metabolic pathways.

NR4A3 mRNA is induced by exercise and repressed during inactivity in human skeletal muscle (Pillon et al., 2020). We previously reported that gene expression profiles of primary human skeletal myotubes subjected to *NR4A3* silencing best reflects the human muscle transcriptome after periods of bedrest and limb immobilisation (Pillon et al., 2020). To build upon this observation and gain further insight into the regulation of *NR4A3*, we analysed eight published transcriptomic studies of the human skeletal muscle response to inactivity. The meta-analysis revealed that physical inactivity reduced *NR4A3* mRNA by 27%, but this decrease was not evident in all studies (**Figure 1A**). Grouping studies based on the duration of inactivity demonstrated that skeletal muscle *NR4A3* expression is transiently repressed during the first days of disuse before returning to baseline levels after 2 weeks (**Figure 1B**). Correlation of *NR4A3* with all other transcripts present in the eight studies allowed for the characterisation of pathways co-regulated with *NR4A3* during inactivity (**Figure 1C**). Gene set enrichment analysis showed that *NR4A3* was positively associated with mitochondrial function and negatively associated with pathways related to cytoskeleton organisation, chromatin regulation, protein synthesis, and degradation (**Figure 1C**). Furthermore, combined analysis of two datasets exploring the restoration of ambulatory behaviour after inactivity showed that reloading re-established *NR4A3* mRNA to pre-inactivity levels (**Figure 1D**), consistent with its role as a contraction-responsive transcription factor in skeletal muscle (Pillon et al., 2020).

Silencing *NR4A3* attenuates glucose oxidation by diverting glucose towards lactate and increases fatty acid oxidation.

To mimic the decreased levels observed in human skeletal muscle after inactivity, *NR4A3* was experimentally downregulated using RNA interference in differentiated primary human skeletal muscle cells. Silencing of *NR4A3* was associated with modest compensatory increases in other NR4A family members *NR4A1* (*NUR77*) and *NR4A2* (*NURR1*) (**Figure 2A**). NR4A3 protein was predominantly localised to the nucleus in myotubes and silencing efficiently reduced nuclear protein abundance (**Figure 2B**). This depletion of NR4A3 lowered basal and FCCP-stimulated (uncoupled) glucose oxidation (**Figure 2C**) independent of glucose transport. Rather, several phosphorylation events in the canonical insulin signalling cascade were potentiated by silencing of *NR4A3* (**Supplementary Figure 1**), including site thr308

on AKT and downstream inactivating phosphorylation of both AKT substrate of 160 kDa (AS160, also known as TBC1D4) and glycogen synthase kinase 3 (GSK3), which facilitate glucose transporter 4 (GLUT4) translocation and glycogen storage, respectively. Furthermore, basal and insulin-mediated glucose uptake and incorporation into glycogen were unaffected (**Supplementary Figures 1H, 1I**).

Upon *NR4A3* silencing, lactate production and release into culture medium was augmented (**Figure 2D**), indicating a shift in glucose fate that was also reported in mouse C2C12 myotubes (Pearen, Myers et al., 2008). Accordingly, mRNA expression of the alpha isoform of lactate dehydrogenase (*LDHA*) increased after *NR4A3* silencing, while expression of the beta isoform decreased (*LDHB*) (**Figure 2E**). Despite unaltered total LDH protein abundance (**Supplementary Figure 1K**), the modification of LDH isoform composition indicates preferential production of lactate from glycolysis and/or less conversion of lactate to pyruvate (Liang, Liu et al., 2016). Lactate release upon NR4A3 depletion could be further supported by the reduction of pyruvate kinase muscle 2 (*PKM2*) mRNA (**Figure 2E**), as inhibiting noncanonical PKM2 activity increased lactate release from C2C12 myoblasts (Blum, Gheller et al., 2021). For glucose to be oxidised in the mitochondria, pyruvate must be metabolised to acetyl coenzyme A (acetyl-CoA) via the pyruvate dehydrogenase (PDH) complex. *NR4A3* silencing upregulated pyruvate dehydrogenase kinase 4 (*PDK4*) gene expression (**Figure 2E**) consistent with attenuated glucose oxidation (Pin, Novinger et al., 2019) (**Figure 2C**). However, the inhibitory PDK4-target phosphorylation site on PDH E1 component subunit alpha (PDHA1^{ser293}) was unaltered (**Figure 2F**), indicating PDH activity was not perturbed.

We previously reported that mitochondrial respiration (in the presence of glucose, glutamine, and pyruvate) and protein subunits of the electron transport chain were diminished by *NR4A3* silencing in primary human skeletal myotubes (Pillon et al., 2020). This aligns with findings in C2C12 myotubes, where *Nr4a3* depletion decreased mitochondrial membrane potential in the absence of changes in mitochondrial DNA (Paez et al., 2023). Instead, downregulation of *Nr4a3* negatively affected the expression of respiratory complex genes, as well as transcripts for mitochondrial ribosomal proteins (Paez et al., 2023). Additionally, *Nr4a3* silencing reduced the mRNA abundance of mitofusin-2 (*Mfn2*) but increased dynamin 1 like expression (*Dnm1l*; also known as *Drp1*) (Paez et al., 2023). These data suggest that loss of NR4A3 attenuates mitochondrial fusion dynamics (instead favouring fission), mitochondrial reticulum connectivity, and

assembly of electron transport chain complexes in a manner separate from alterations in mitochondrial abundance. Our finding that myotube's response to the mitochondrial uncoupler FCCP was maintained after *NR4A3* RNA interference further supports not only retained mitochondrial content, but also the presence of functional mitochondria (**Figure 2C**). Thus, the attenuation of glucose oxidation in NR4A3-depleted myotubes appears to be a consequence of greater diversion of glucose towards lactate, which would reduce glucose-derived acetyl-CoA entry into the tricarboxylic acid (TCA) cycle.

Skeletal muscle cells are highly glycolytic in culture, preferentially oxidising glucose at rates 100-fold faster than long-chain lipids (Abdelmoez, Sardon Puig et al., 2020). Therefore, we explored the consequence of NR4A3-dependent reductions in glucose oxidation on lipid metabolism in primary human skeletal myotubes. NR4A3 depletion upregulated basal and FCCP-stimulated rates of fatty acid oxidation (**Figure 3A**) without altering the intracellular composition of select lipid species. Whereas the ratio of triacylglycerides, diacylglycerides, free fatty acids, and other lipids were modified by FCCP, *NR4A3* RNA interference was without effect (**Figure 3B**). Compatible with greater fatty acid oxidation, *NR4A3* silencing increased phosphorylation of the AMP-activated protein kinase α subunit (AMPKα^{thr172}) and its substrates acetyl-CoA carboxylase (ACC^{ser79}) and TBC domain family member 1 (TBC1D1^{ser237}) (**Figure 3C**). This signalling coalesced with activating phosphorylation of hormone-sensitive lipase (HSL^{ser660}), reduced LIPIN1 protein abundance (**Figure 3D**), and downregulation of total ACC protein (**Figure 3C**), as well as the mRNA of its cytosolic isoform (*ACCα*) (**Figure 3E**). Further profiling of transcripts involved in fatty acid metabolic pathways revealed that NR4A3 depletion altered the composition of AMPKα isoforms and fatty acid transport genes (**Figure 3E**). Compared to control, *NR4A3* silencing increased the expression of fatty acid transport protein 1 (*FATP1*) and reduced the expression of fatty acid binding protein 3 (*FABP3*) (**Figure 3E**). Together, the protein and transcriptional changes induced after RNA interference implicate NR4A3 in physiological mechanisms compensating for energy stress due to impaired glucose oxidation.

NR4A3 silencing in primary human myotubes downregulates translation, mTORC1 signalling, and pre-rRNA abundance.

Metabolic assays are commonly normalised to protein content to account for differences in seeding density. We observed that silencing of *NR4A3* consistently reduced the total protein and RNA abundance of cultured human myotubes (**Figures 4A, 4B**). Over 80% of cellular RNA is

ribosomal (rRNA) and rates of translation are proportional to total RNA content in skeletal muscle (Millward, Garlick et al., 1973). These observations, combined with the association of reduced *NR4A3* levels during bed rest and limb immobilisation inactivity (**Figure 1**), led us to explore the putative effects of *NR4A3* silencing on protein synthesis. Relative rates of translation can be measured *in vitro* by the surface sensing of translation (SUnSET) method, which detects puromycin incorporation into nascent peptide chains by immunoblot analysis (Goodman, Mabrey et al., 2011). Using this technique, we observed reduced puromycilation of proteins upon *NR4A3* silencing both at baseline and after insulin plus leucine stimulation (**Figure 4C**), suggesting impaired protein synthesis with NR4A3 downregulation.

The efficiency and total capacity for translation in skeletal muscle is determined by mTORC1 signalling and ribosomal content, respectively (Figueiredo et al., 2021, West et al., 2016). *In vitro*, AMPK impedes mTORC1 activity through phosphorylation of tuberous sclerosis complex 2 (TSC2) (Inoki, Zhu et al., 2003) and raptor (Gwinn, Shackelford et al., 2008). Consistent with energy-stress dependent upregulation of AMPK (**Figure 3C**), the ratio of phosphorylated-to-total protein of AMPK target sites on TSC2 (ser1387) and raptor (ser792) were increased by NR4A3 depletion (**Figures 4D-E**; **Supplementary Figure 2A**) indicative of AMPK interference of mTORC1. Furthermore, *NR4A3* silencing reduced the protein abundance of raptor, mTOR, and ribosomal protein S6 (RPS6) (**Figures 4D-E**; **Supplementary Figures 2B-D**). These collective AMPK-related and unrelated changes coalesced to inhibit mTOR^{ser2448} phosphorylation and activation of downstream mTORC1-substrates, including RPS6$^{ser235/236}$ and 4EBP1$^{thr37/46}$ phosphorylation, with negative consequences for skeletal muscle protein synthesis following NR4A3 depletion (**Figures 4D**, **4E**; **Supplementary Figures 2C-E**).

In agreement with diminished translation, gene expression analyses confirmed that *NR4A3* RNA interference decreased mRNA abundance of *RPS6*, as well as eukaryotic translation initiation factor 4E (*EIF4E*) and 45S pre-ribosomal RNA (5' external transcribed spacer; *5' ETS*) (**Figure 4F**). The 45S pre-rRNA polycistronic precursor is transcribed by RNA polymerase I (Pol I) and gives rise to the 18S, 5.8S, and 28S rRNAs (von Walden, 2019). These rRNAs are subsequently assembled with Pol II-dependent ribosomal proteins and Pol III-transcribed 5S rRNA into the functional 40S (small) and 60S (large) ribosomal subunits (von Walden, 2019). As such, Pol I activity is the rate-limiting step in ribosomal biogenesis. Due to the short half-life of the ETS region (Popov, Smirnov et al., 2013), measurement of the 45S pre-RNA 5' ETS is a reliable indicator of Pol I-mediated ribosomal transcription (von Walden, 2019). Thus, our results imply that lower levels of total RNA from *NR4A3*-silenced myotubes (**Figure 4B**) are a consequence of attenuated ribosomal biogenesis, resulting in reduced ribosomal mass, and that NR4A3 downregulation impedes both translational efficiency (i.e. mTORC1) and capacity (i.e. ribosomal abundance).

Despite an overall reduction in protein synthesis (**Figure 4C**), *NR4A3*-depleted myotubes still responded to insulin plus leucine stimulation, as suggested by the interaction between *NR4A3* silencing and treatment for mTOR^{ser2488}, RPS6$^{ser235/236}$, and the inhibitory AKT-target phosphorylation site on TSC2 (thr1462) (**Figure 4D**; **Supplementary Figures 2C**, **2D**, **2F**). AKT^{thr308} levels were greater upon insulin treatment (**Supplementary Figures 1A, 1E**) and tended to increase with insulin plus leucine stimulation (**Figure 4D-E**; **Supplementary Figure 2G**). Furthermore, mRNA of the large neutral amino acid transporter small subunit 1 (*LAT1*; also known as *SLC7A5*) was upregulated by *NR4A3* silencing (**Figure 4F**). LAT1 is the dominant antiporter for leucine uptake into cells (Gauthier-Coles, Vennitti et al., 2021). The rise in intracellular leucine is then detected through biochemical sensing mechanisms, including disassociation of sestrin-2 (SESN2) with GATOR2 (Saxton, Knockenhauer et al., 2016) and activation of leucyl-tRNA synthetase 1 (LARS1) (Han, Jeong et al., 2012), with subsequent activation of mTORC1. As such, increased LAT1 could feasibly support higher intramuscular leucine concentrations under *NR4A3*-silenced conditions, which could have offset any detrimental effects of increased *SESN2* or reduced *LARS1* mRNA (**Figure 4F**). Hence, canonical insulin signalling, combined with enhanced leucine transport, provides a mechanism by which anabolic sensitivity of NR4A3-depleted myotubes is retained via the AKT-mTORC1 pathway.

We next investigated the impact of NR4A3 downregulation on markers of inflammation and protein degradation. Although the contribution of protein breakdown towards human disuse muscle atrophy is unclear (Brook et al., 2022, Pavis et al., 2023, Shur et al., 2024), the ubiquitin-proteasomal (UPS) and autophagy-lysosomal systems are markedly induced during murine models of inactivity (Min, Smuder et al., 2011, Talbert, Smuder et al., 2013). NR4A3 depletion upregulated activating transcription factor 3 (*ATF3*) and DNA damage-inducible transcript 3 (*DDIT3*; also known as *CHOP*) mRNA, whilst reducing IkBα protein abundance (**Supplementary Figures 3A-C**), signifying possible inflammation-mediated

sarcoplasmic reticulum stress. However, other surrogates of inflammatory nuclear factor kappa B (NF-kB) signalling and the unfolded protein response, such as interleukin 6 (*IL-6*) and *ATF4*, were unaltered (**Supplementary Figure 3A**). Major proteolytic pathways use ubiquitination as a signal for select protein degradation (Pohl & Dikic, 2019). Indeed, overexpression of the muscle-specific E3 ubiquitin ligase muscle ring-finger 1 (MuRF1 or TRIM63) is sufficient to induce ubiquitination and muscle atrophy in mice (Baehr, Hughes et al., 2021). The *MuRF1* promoter is a direct target of forkhead box O3 (FOXO3a) (Bollinger, Witczak et al., 2014) and inhibitory phosphorylation of FOXO3a and FOXO1 were lower after *NR4A3* interference (**Supplementary Figures 3D-E**), indicative of greater FOXO transcriptional activity. Consistent with this, mRNA expression of *MuRF1* and *FOXO3* were also elevated (**Supplementary Figure 3A**). However, these findings were contrasted by downregulation of both MURF1 and FOXO3a protein levels, while FOXO1 and the alternative muscle-enriched E3 ligase MAFbx (also known as atrogin-1) were unperturbed (**Supplementary Figures 3D-G**). Furthermore, in contrast to the stark impairment of protein synthesis, *NR4A3* silencing did not alter global protein ubiquitination in primary skeletal myotubes (**Supplementary Figure 3H**). Rather, the proteases calpain-1 (including the activated, autolysed form) and caspase 3 (inactive zymogen form) were decreased upon NR4A3 depletion, as were alpha subunit proteins of the 20S core particle proteasome (Pan 20Sα) (**Supplementary Figures 3I-K**), whereas proteasome 20S subunit alpha 1 (*PSMA1*) and proteasome 20S subunit beta 2 (*PSMB2*) transcripts were induced and attenuated, respectively (**Supplementary Figure 3A**). Additionally, despite skeletal muscle-specific *NR4A3* transgenic mice displaying greater autophagic flux (Goode, Pearen et al., 2016), *NR4A3* interference had no impact on autophagy markers unc-51 like autophagy activating kinase 1 (ULK1), ubiquitin-binding protein P62 (also known as sequestrosome-1), or microtubule-associated proteins 1A/1B-light chain 3 (LC3-I and LC3-II) (**Supplementary Figures 3L-N**).

Altogether, silencing *NR4A3* in skeletal myotubes triggered AMPK-dependent and independent inhibition of mTORC1, reducing ribosomal biogenesis and protein translation. Pathways related to proteolysis were negligibly affected, suggesting NR4A3 downregulation primarily reduces skeletal muscle protein content via multi-level attenuation of protein synthesis.

Overexpression of the canonical *NR4A3* isoform increases protein synthesis in primary human myotubes.

The *NR4A3* gene produces four transcripts encoding three isoforms of the NR4A3 protein. The RNA interference method described in previous figures targeted an exon common to all isoforms, precluding the analysis of differential effects of select *NR4A3* variants. Thus, we explored lentiviral overexpression of the canonical *NR4A3* isoform (*NR4A3-203*; NM_006981.4) in primary human skeletal myotubes. Overexpression of the *NR4A3-203* (*NR-203*) transcript by >140-fold increased total NR4A3 protein levels by >4.5-fold without altering the abundance of other NR4A family members (**Figure 5A**) or affecting aspects of glucose and fatty acid metabolism (**Supplementary Figures 4A-D**). However, consistent with the effects of *NR4A3* silencing, *NR4A3-203* overexpression enhanced protein synthesis at baseline and after insulin plus leucine treatment (**Figure 5B**). This result was further supported by assessment of radiolabelled phenylalanine incorporation into protein (**Figure 5C**). Here, canonical *NR4A3* overexpression increased rates of translation under both unstimulated conditions and after foetal bovine serum (FBS) plus leucine treatment when combined with proteasomal blockade (lactacystin) or mTORC1-inhibition (rapamycin) (**Figure 5C**). That total cellular protein content of *NR4A3-203* overexpressing myotubes was greater in these same conditions implies the canonical NR4A3 isoform may upregulate overall protein turnover in a manner partly independent from mTORC1 (**Figure 5D**). Indeed, no differences in total RNA concentrations or mTORC1 signalling were observed after *NR4A3-203* overexpression (**Supplementary Figures 4E, 4F**), suggesting that *NR4A3* silencing and overexpression phenotypes manifest through different mechanisms. This hypothesis was further substantiated by the failure of *NR4A3-203* overexpression to modulate transcripts affected by silencing (**Supplementary Figure 3G**). Nevertheless, overexpression of the canonical *NR4A3* isoform in the presence of siRNA targeting all *NR4A3* variants (**Figure 6A-C**) partially restored glucose oxidation (**Figure 6D**) concomitant with recovery of signalling downstream mTORC1: S6K, RPS6 and 4EBP1 (**Figure 6E**, **Supplementary Figures 5A-C**). Despite these effects, AMPKα^{thr172} phosphorylation remained elevated after restoration of *NR4A3-203* (**Supplementary Figure 5D**), which may contribute to sustained augmentation of fatty acid oxidation rates (**Supplementary Figure 5E**). Overall, these results posit the *NR4A3-203* isoform as a major contributor towards the negative metabolic outcomes

associated with NR4A3-dependent downregulation of protein synthesis.

NR4A3 silencing alters primary human myotube size and structure.

NR4A3 is decreased during inactivity associated with muscle disuse (**Figure 1A, 1B**). We therefore examined how changes in translation and protein synthesis induced by *NR4A3* silencing impact on myotube size and contractile apparatus. Immunostaining for fast-type myosin heavy chain isoforms (MYH1/2; type-IIX/type-IIA) revealed a striking decrease in myotube size (**Figure 7A**). This difference was driven by a reduction in myotube area (**Figure 7B**) without any change in the number of nuclei (**Figure 7C**) or the ability of myoblasts to fuse into myotubes (**Figure 7D**).

NR4A3 depletion reduced the protein abundance of fast (MYH1/2) and slow (MYH7) myosin heavy chain isoforms (**Figure 7E**; **Supplemental Figure 6**) and altered the expression of transcripts encoding for proteins integral to skeletal muscle contraction and structure. This included downregulation of *MYH1*, titin (*TTN*), capping actin proteins of muscle Z-line (*CAPZ* family), and troponin isoforms *TTNT2* and *TTNT3* (**Figure 6F**). Altogether, the protein and mRNA levels of major structural and contractile proteins of skeletal muscle cells were disrupted by *NR4A3* silencing, associated with diminished myotube size.

Conclusions

NR4A3 is markedly induced following a single exercise bout, but not after habitual exercise training (Pillon et al., 2020). Similarly, *NR4A3* is transiently downregulated during reduced physical activity and quickly upregulated upon reloading, but unchanged after prolonged immobilisation (>2 weeks). The onset of muscle fibre atrophy occurs rapidly with disuse (Brook et al., 2022, Shur et al., 2024) and is most aggressive across the initial two weeks of unloading (Hardy, Inns et al., 2022), which coincides with the lowest levels of *NR4A3* mRNA. This expression pattern suggests a crucial role for NR4A3 in regulating skeletal muscle function during the acute transition phases associated with intense tissue remodelling. The first two weeks of unloading correspond to lower postprandial (Breen et al., 2013) and total daily (Shad et al., 2019) rates of myofibrillar protein synthesis. Here, we provide evidence that NR4A3 can directly influence translation in skeletal muscle. Furthermore, NR4A3 downregulation in myotubes recapitulated adverse effects observed in skeletal muscle tissue following inactivity (Eggelbusch, Charlton et al., 2024). This includes impaired glucose metabolism, reduced protein synthesis, altered expression of contractile apparatus, and diminished myotube size. Thus, our results shed light on the necessary involvement of NR4A3 in the control of skeletal muscle protein synthesis and metabolism during disuse atrophy.

Several lines of evidence from both *in vitro* and *in vivo* studies suggest that NR4A3 is associated with metabolic responses in skeletal muscle. Rats bred for fitness have skeletal muscle enrichment of NR4A3, associated with higher mitochondrial content and increased running capacity (Stephenson, Stepto et al., 2012). In transgenic mice, *Nr4a3* overexpression promotes skeletal muscle remodelling towards an oxidative phenotype (Pearen et al., 2012), suggesting that NR4A3 plays a causal role in metabolic reprogramming of skeletal muscle. Observations from *in vivo* studies corroborate with results from *in vitro* experiments in mouse C2C12 and primary human myotubes demonstrating that silencing of NR4A3 changes gene expression profiles towards anaerobic programmes (Pearen et al., 2008) and impairs mitochondrial function (Paez et al., 2023, Pearen et al., 2008, Pillon et al., 2020). Such effects may account for the lower adenosine triphosphate (ATP) concentrations in C2C12 myotubes after *Nr4a3* silencing (Pearen et al., 2008). Our data add mechanistic evidence that NR4A3 downregulation profoundly impacts substrate utilisation, shifting metabolism towards increased lactate production and fatty acid oxidation, while decreasing glucose oxidation. Additionally, we corroborate findings in C2C12 (Paez et al., 2023) that the mTORC1 complex is a major signalling node inhibited by NR4A3 depletion. Activation of AMPK led to the phosphorylation of TSC2 and raptor at sites that blunt mTORC1 transduction. However, we also observed decreased mTOR and RPS6 protein abundance, a phenomenon unlikely triggered exclusively by the activation of AMPK. Indeed, rates of translation and ribosomal biogenesis were attenuated in response to *NR4A3* RNA interference, indicating that reductions in NR4A3 impair cellular metabolism by obstructing cell-wide anabolic pathways. In agreement, rescuing protein synthesis via overexpression of the canonical NR4A3 isoform restored glucose oxidation in *NR4A3*-silenced myotubes.

Our study connects lower levels of NR4A3 with inactivity paradigms and indicates that attenuation of NR4A3 in human skeletal muscle has adverse effects on protein synthesis and metabolic responses. Muscle mass and strength are important predictors of mortality in intensive care (Weijs, Looijaard et al., 2014). Therefore, NR4A3 may be a key molecular transducer, downregulation of which contributes to the deleterious health consequences associated with sedentary lifestyles. Understanding and addressing NR4A3 regulation could have far-

reaching implications for global health by mitigating the adverse effects of inactivity, aging, and metabolic diseases.

References

Abdelmoez AM, Sardon Puig L, Smith JAB, Gabriel BM, Savikj M, Dollet L, Chibalin AV, Krook A, Zierath JR, Pillon NJ (2020) Comparative profiling of skeletal muscle models reveals heterogeneity of transcriptome and metabolism. *Am J Physiol Cell Physiol* 318: C615-C626

Amoasii L, Sanchez-Ortiz E, Fujikawa T, Elmquist JK, Bassel-Duby R, Olson EN (2019) NURR1 activation in skeletal muscle controls systemic energy homeostasis. *Proc Natl Acad Sci U S A* 116: 11299-11308

Baehr LM, Hughes DC, Lynch SA, Van Haver D, Maia TM, Marshall AG, Radoshevich L, Impens F, Waddell DS, Bodine SC (2021) Identification of the MuRF1 Skeletal Muscle Ubiquitylome Through Quantitative Proteomics. *Function* 2

Bergouignan A, Antoun E, Momken I, Schoeller DA, Gauquelin-Koch G, Simon C, Blanc S (2013) Effect of contrasted levels of habitual physical activity on metabolic flexibility. *J Appl Physiol (1985)* 114: 371-9

Blum JE, Gheller BJ, Benvie A, Field MS, Panizza E, Vacanti NM, Berry D, Thalacker-Mercer A (2021) Pyruvate Kinase M2 Supports Muscle Progenitor Cell Proliferation but Is Dispensable for Skeletal Muscle Regeneration after Injury. *J Nutr* 151: 3313-3328

Bollinger LM, Witczak CA, Houmard JA, Brault JJ (2014) SMAD3 augments FoxO3-induced MuRF-1 promoter activity in a DNA-binding-dependent manner. *Am J Physiol Cell Physiol* 307: C278-87

Breen L, Stokes KA, Churchward-Venne TA, Moore DR, Baker SK, Smith K, Atherton PJ, Phillips SM (2013) Two weeks of reduced activity decreases leg lean mass and induces "anabolic resistance" of myofibrillar protein synthesis in healthy elderly. *J Clin Endocrinol Metab* 98: 2604-12

Brook MS, Stokes T, Gorissen SHM, Bass JJ, McGlory C, Cegielski J, Wilkinson DJ, Phillips BE, Smith K, Phillips SM, Atherton PJ (2022) Declines in muscle protein synthesis account for short-term muscle disuse atrophy in humans in the absence of increased muscle protein breakdown. *J Cachexia Sarcopenia Muscle* 13: 2005-2016

Chao LC, Wroblewski K, Ilkayeva OR, Stevens RD, Bain J, Meyer GA, Schenk S, Martinez L, Vergnes L, Narkar VA, Drew BG, Hong C, Boyadjian R, Hevener AL, Evans RM, Reue K, Spencer MJ, Newgard CB, Tontonoz P (2012) Skeletal muscle Nur77 expression enhances oxidative metabolism and substrate utilization. *J Lipid Res* 53: 2610-9

Eggelbusch M, Charlton BT, Bosutti A, Ganse B, Giakoumaki I, Grootemaat AE, Hendrickse PW, Jaspers Y, Kemp S, Kerkhoff TJ, Noort W, van Weeghel M, van der Wel NN, Wesseling JR, Frings-Meuthen P, Rittweger J, Mulder ER, Jaspers RT, Degens H, Wust RCI (2024) The impact of bed rest on human skeletal muscle metabolism. *Cell Rep Med* 5: 101372

Figueiredo VC, D'Souza RF, Van Pelt DW, Lawrence MM, Zeng N, Markworth JF, Poppitt SD, Miller BF, Mitchell CJ, McCarthy JJ, Dupont-Versteegden EE, Cameron-Smith D (2021) Ribosome biogenesis and degradation regulate translational capacity during muscle disuse and reloading. *J Cachexia Sarcopenia Muscle* 12: 130-143

Gauthier-Coles G, Vennitti J, Zhang Z, Comb WC, Xing S, Javed K, Broer A, Broer S (2021) Quantitative modelling of amino acid transport and homeostasis in mammalian cells. *Nat Commun* 12: 5282

Goode JM, Pearen MA, Tuong ZK, Wang SC, Oh TG, Shao EX, Muscat GE (2016) The Nuclear Receptor, Nor-1, Induces the Physiological Responses Associated With Exercise. *Mol Endocrinol* 30: 660-76

Goodman CA, Mabrey DM, Frey JW, Miu MH, Schmidt EK, Pierre P, Hornberger TA (2011) Novel insights into the regulation of skeletal muscle protein synthesis as revealed by a new nonradioactive in vivo technique. *FASEB J* 25: 1028-39

Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, Turk BE, Shaw RJ (2008) AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol Cell* 30: 214-26

Han JM, Jeong SJ, Park MC, Kim G, Kwon NH, Kim HK, Ha SH, Ryu SH, Kim S (2012) Leucyl-tRNA synthetase is an intracellular leucine sensor for the mTORC1-signaling pathway. *Cell* 149: 410-24

Hardy EJO, Inns TB, Hatt J, Doleman B, Bass JJ, Atherton PJ, Lund JN, Phillips BE (2022) The time course of disuse muscle atrophy of the lower limb in health and disease. *J Cachexia Sarcopenia Muscle* 13: 2616-2629

Inoki K, Zhu T, Guan KL (2003) TSC2 mediates cellular energy response to control cell growth and survival. *Cell* 115: 577-90

Kaiser MS, Milan G, Ham DJ, Lin S, Oliveri F, Chojnowska K, Tintignac LA, Mittal N, Zimmerli CE, Glass DJ, Zavolan M, Ruegg MA (2022) Dual roles of mTORC1-dependent activation of the ubiquitin-proteasome system in muscle proteostasis. *Commun Biol* 5: 1141

Liang X, Liu L, Fu T, Zhou Q, Zhou D, Xiao L, Liu J, Kong Y, Xie H, Yi F, Lai L, Vega RB, Kelly DP, Smith SR, Gan Z (2016) Exercise Inducible Lactate Dehydrogenase B Regulates Mitochondrial Function in Skeletal Muscle. *J Biol Chem* 291: 25306-25318

Maira M, Martens C, Philips A, Drouin J (1999) Heterodimerization between members of the Nur subfamily of orphan nuclear receptors as a novel mechanism for gene activation. *Mol Cell Biol* 19: 7549-57

Manini TM, Clark BC, Nalls MA, Goodpaster BH, Ploutz-Snyder LL, Harris TB (2007) Reduced physical activity increases intermuscular adipose tissue in healthy young adults. *Am J Clin Nutr* 85: 377-84

Massart J, Zierath JR, Chibalin AV (2014) A simple and rapid method to characterize lipid fate in skeletal muscle. *BMC Res Notes* 7: 391

Millward DJ, Garlick PJ, James WP, Nnanyelugo DO, Ryatt JS (1973) Relationship between protein synthesis and RNA content in skeletal muscle. *Nature* 241: 204-5

Min K, Smuder AJ, Kwon OS, Kavazis AN, Szeto HH, Powers SK (2011) Mitochondrial-targeted antioxidants protect skeletal muscle against immobilization-induced muscle atrophy. *J Appl Physiol (1985)* 111: 1459-66

Paez HG, Ferrandi PJ, Pitzer CR, Mohamed JS, Alway SE (2023) Loss of NOR-1 represses muscle metabolism through mTORC1-mediated signaling and mitochondrial gene expression in C2C12 myotubes. *FASEB J* 37: e23050

Pavis GF, Abdelrahman DR, Murton AJ, Wall BT, Stephens FB, Dirks ML (2023) Short-term disuse does not affect postabsorptive or postprandial muscle protein fractional breakdown rates. *J Cachexia Sarcopenia Muscle* 14: 2064-2075

Pearen MA, Eriksson NA, Fitzsimmons RL, Goode JM, Martel N, Andrikopoulos S, Muscat GE (2012) The nuclear receptor, Nor-1, markedly increases type II oxidative muscle fibers and resistance to fatigue. *Mol Endocrinol* 26: 372-84

Pearen MA, Myers SA, Raichur S, Ryall JG, Lynch GS, Muscat GE (2008) The orphan nuclear receptor, NOR-1, a target of beta-adrenergic signaling, regulates gene expression that controls oxidative metabolism in skeletal muscle. *Endocrinology* 149: 2853-65

Pillon NJ, Gabriel BM, Dollet L, Smith JAB, Sardon Puig L, Botella J, Bishop DJ, Krook A, Zierath JR (2020) Transcriptomic profiling of skeletal muscle adaptations to exercise and inactivity. *Nat Commun* 11: 470

Pin F, Novinger LJ, Huot JR, Harris RA, Couch ME, O'Connell TM, Bonetto A (2019) PDK4 drives metabolic alterations and muscle atrophy in cancer cachexia. *FASEB J* 33: 7778-7790

Pohl C, Dikic I (2019) Cellular quality control by the ubiquitin-proteasome system and autophagy. *Science* 366: 818-822

Popov A, Smirnov E, Kovacik L, Raska O, Hagen G, Stixova L, Raska I (2013) Duration of the first steps of the human rRNA processing. *Nucleus* 4: 134-41

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 43: e47

Saxton RA, Knockenhauer KE, Wolfson RL, Chantranupong L, Pacold ME, Wang T, Schwartz TU, Sabatini DM (2016) Structural basis for leucine sensing by the Sestrin2-mTORC1 pathway. *Science* 351: 53-8

Shad BJ, Thompson JL, Holwerda AM, Stocks B, Elhassan YS, Philp A, LJC VANL, Wallis GA (2019) One Week of Step Reduction Lowers Myofibrillar Protein Synthesis Rates in Young Men. *Med Sci Sports Exerc* 51: 2125-2134

Shur NF, Simpson EJ, Crossland H, Constantin D, Cordon SM, Constantin-Teodosiu D, Stephens FB, Brook MS, Atherton PJ, Smith K, Wilkinson DJ, Mougin OE, Bradley C, Macdonald IA, Greenhaff PL (2024) Bed-rest and exercise remobilization: Concurrent

adaptations in muscle glucose and protein metabolism. *J Cachexia Sarcopenia Muscle* 15: 603-614

Smith JAB, Murach KA, Dyar KA, Zierath JR (2023) Exercise metabolism and adaptation in skeletal muscle. *Nat Rev Mol Cell Biol* 24: 607-632

Stephenson EJ, Stepto NK, Koch LG, Britton SL, Hawley JA (2012) Divergent skeletal muscle respiratory capacities in rats artificially selected for high and low running ability: a role for Nor1? *J Appl Physiol (1985)* 113: 1403-12

Talbert EE, Smuder AJ, Min K, Kwon OS, Szeto HH, Powers SK (2013) Immobilization-induced activation of key proteolytic systems in skeletal muscles is prevented by a mitochondria-targeted antioxidant. *J Appl Physiol (1985)* 115: 529-38

Tontonoz P, Cortez-Toledo O, Wroblewski K, Hong C, Lim L, Carranza R, Conneely O, Metzger D, Chao LC (2015) The orphan nuclear receptor Nur77 is a determinant of myofiber size and muscle mass in mice. *Mol Cell Biol* 35: 1125-38

von Walden F (2019) Ribosome biogenesis in skeletal muscle: coordination of transcription and translation. *J Appl Physiol (1985)* 127: 591-598

Weijs PJ, Looijaard WG, Dekker IM, Stapel SN, Girbes AR, Oudemans-van Straaten HM, Beishuizen A (2014) Low skeletal muscle area is a risk factor for mortality in mechanically ventilated critically ill patients. *Crit Care* 18: R12

West DW, Baehr LM, Marcotte GR, Chason CM, Tolento L, Gomes AV, Bodine SC, Baar K (2016) Acute resistance exercise activates rapamycin-sensitive and -insensitive mechanisms that control translational activity and capacity in skeletal muscle. *J Physiol* 594: 453-68

Wilson TE, Fahrner TJ, Johnston M, Milbrandt J (1991) Identification of the DNA binding site for NGFI-B by genetic selection in yeast. *Science* 252: 1296-300

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, Fu X, Liu S, Bo X, Yu G (2021) clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innovation (Camb)* 2: 100141

Yerrakalva D, Hajna S, Suhrcke M, Wijndaele K, Westgate K, Khaw KT, Wareham N, Brage S, Griffin S (2023) Associations between change in physical activity and sedentary time and health-related quality of life in older english adults: the EPIC-Norfolk cohort study. *Health Qual Life Outcomes* 21: 60

You JS, McNally RM, Jacobs BL, Privett RE, Gundermann DM, Lin KH, Steinert ND, Goodman CA, Hornberger TA (2019) The role of raptor in the mechanical load-induced regulation of mTOR signaling, protein synthesis, and skeletal muscle hypertrophy. *FASEB J* 33: 4021-4034

Yuan S, Li X, Liu Q, Wang Z, Jiang X, Burgess S, Larsson SC (2023) Physical Activity, Sedentary Behavior, and Type 2 Diabetes: Mendelian Randomization Analysis. *J Endocr Soc* 7: bvad090

Acknowledgments

This study was supported by Swedish Research Council (2015-00165, 2018-02389), Novo Nordisk Foundation (NNF22OC0077741), Swedish Diabetes Foundation (DIA2023-824), Swedish Research Council for Sport Science (P2023-0093), Knut and Alice Wallenberg (2021.0249), Strategic Research Program in Diabetes at Karolinska Institutet (2009-1068), Knut and Alice Wallenberg (2021.0249), European Foundation for the Study of Diabetes/Novo Nordisk Foundation Future Leaders Award (NNF21SA0072747), and Diabetes Wellness Sverige (PG21-6524)

The computations were enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS), partially funded by the Swedish Research Council through grant agreement no. 2022-06725. Figure 4E was created with BioRender.com.

Methods

Transcriptomic meta-analysis. Transcriptomic studies of inactivity in human trials were selected using the MetaMEx database (Pillon et al., 2020). Studies were annotated with the sex and age of participants, as well as the total duration of inactivity. Statistics were performed individually for each study and the mean, variance and n size were used to fit a random effects model to the data. The restricted maximum-likelihood (REML) method was applied using the R package metaphor. Obtained p-values were adjusted using the Benjamini & Hochberg method. To evaluate the effect of inactivity duration, all transcriptomic studies were merged by gene name and split in three groups: less than one week, one-two weeks or more than two weeks. A linear model was used to test the overall effect of inactivity duration, as well as pair-wise comparisons to the pre-inactivity group, using Empirical Bayes Statistics for Differential Expression with the limma package (Ritchie, Phipson et al., 2015). Gene set enrichment analysis was performed using ClusterProfiler (Wu, Hu et al., 2021) on genes ranked based on Spearman correlation with *NR4A3*. Gene ontology biological processes were considered significant at FDR<0.05. The effect of reloading was estimated with the same method as above after merging GSE21496 and GSE24215.

Primary human cell culture. Primary cells were isolated from *vastus lateralis* skeletal muscle biopsies derived from ten healthy volunteers (age: 36 ± 15 years; BMI: 24.0 ± 1.9 Kg.m^{-2}; biological sex: eight males, two females). The regional ethical review board in Stockholm approved protocols. Cells were cultured and differentiated as described previously (Abdelmoez et al., 2020). Successful myotube formation was monitored under the microscope and cells were used for terminal experiments ten days after the initiation of differentiation. The absence of mycoplasma contamination was routinely confirmed by PCR.

Silencing by RNA interference. On days six and eight of differentiation, myotubes were transfected with 10 nM of either Silencer Select siRNA Negative Control No. 2 (no. 4390847) or Silencer Select siRNA s15542 targeting *NR4A3* (Life Technologies, Foster City, CA). Transfections were performed for 5 h in OptiMEM reduced serum media with Lipofectamine RNAiMAX transfection reagent (Invitrogen, Carlsbad, CA). Terminal experiments were performed ≈48 h after the second transfection.

Plasmid transformation, expansion, and purification. cDNA of the canonical *NR4A3* transcript (NR4A3-203; NM_006981.4) was synthesised into a modified pLenti CMV Puro DEST (w118-1) vector (Addgene plasmid #17452) by GENEWIZ (Azenta Life Science, Leipzig, Germany). One Shot TOP10 Chemically Competent *E. coli* cells (Invitrogen, ThermoFisher Scientific) were used for bacterial transformation and expansion. Modified pLenti CMV Puro DEST (w118-1) empty vector (EV) and NR4A3-overexpressing (*NR-203*^{Oex}) plasmid DNA was then purified using endotoxin-free buffers (Cat. No. 19048) and columns (Cat. No. 10083) from Qiagen, before assessing DNA concentration and purity by spectrophotometry.

Lentivirus production. HEK293T cells were grown in DMEM media (Gibco #10569) supplemented with 10% foetal bovine serum, 1% penicillin-streptomycin, and 500 µg/mL geneticin (G418). Twenty-four hours prior to transfection, HEK293T cells were seeded into T-225 flasks coated with poly-L-lysine. Once ≈90% confluent, lentiviral harvest was conducted by transfecting each T-225 with a transfection reagent mix consisting of 69 µg of EV or *NR-203*^{Oex} plasmid DNA, second generation packaging vectors (69 µg of psPAX2 DNA and 45 µg of pMD2.G DNA), 549 µg polyethylenimine (PEI; 3 µg per µg of total plasmid DNA) and PBS at a final volume of 4.5 mL. The transfection mix was then added to 40.5 mL of pre-warmed growth medium (without geneticin) and added to HEK293T cells for 12-14 h. At this point, the transfection media was removed and replaced with fresh medium for 2 h before switching to new growth media supplemented with 5 mM sodium butyrate. Thirty hours later, viral media was collected and passed through a 0.45 µM low-protein binding filter (Millipore) and concentrated by low-speed centrifugation (3300 g for 30 min at 4°C) using 100 kDa molecular weight cut-off Centricon Plus-70 cartridges (UFC710008). Three volumes of concentrated virus medium were subsequently mixed with one volume of Lenti-X concentrator

(Takara Bio) and rotated at 4°C overnight before centrifugation (1500 g for 45 min at 4°C). Supernatant was discarded and the pellet resuspended in PBS, aliquoted, and stored at -80°C. Virus concentration was determined using Lenti-X GoStix Plus (Takara Bio) after one freeze-thaw cycle to be consistent with experimental conditions.

Transduction of primary skeletal myotubes. On day 7 of differentiation, myotubes were transduced with either EV or *NR-203*[Oex] lentivirus (concentration equivalent to 16.5 ng of p24.mL^{-1} per 9.6 cm^2 of cells) in post-fusion medium supplemented with 5 ug.mL^{-1} of polybrene. After 18-20 h, viral medium was removed, myotubes were washed once with PBS and then switched to normal post-fusion media (Abdelmoez et al., 2020). Terminal experiments were performed ≈48 h later.

RNA extraction and analysis. RNA was isolated from cultured muscle cells using TRIzol-cholorform extraction according to manufacturer's instruction (Invitrogen). RNA concentration and purity were determined by spectrophotometry. All equipment, software, and reagents for performing reverse transcription and RT-qPCR were from Thermo Fisher Scientific. The High-Capacity cDNA Reverse Transcription kit was used for cDNA synthesis according to manufacturer's instructions. RT-qPCR was performed on a StepOne Plus system using Taqman or Sybr Green technologies (probe sequences for all analysed genes are available upon request). Relative gene expression was calculated by the comparative ΔΔCt method using the geometric mean of at least two of the following reference genes: *18S*, *B2M*, *GUSB*, *HPRT1* or *TBP*. The most stable reference gene combination was determined within experimental conditions.

Glucose uptake. Myotubes were washed once with sterile PBS and incubated for 4 h in un-supplemented, serum-free low-glucose Dulbecco's Modified Eagle Medium (DMEM) (5.5 mM glucose, Gibco #21885) (i.e. serum starvation medium). Cells were then switched to fresh serum starvation medium in the absence or presence of insulin (120 nM) for 1 h. Myotubes were washed once with warm glucose- and serum-free DMEM (Gibco #11966) and glucose uptake was measured in the same medium by adding 1 uL.mL^{-1} of 2-[1,2-^{3}H]Deoxy-D-glucose (MT911, Moravek) and 10 μM unlabelled 2-Deoxy-D-glucose for 15 min (Abdelmoez et al., 2020). Cell monolayers were washed three times with ice-cold PBS and lysed in 1 mL 0.03% SDS. 0.5 mL of the cell lysate was counted in a liquid scintillation counter (TRI-CARB 4910TR, PerkinElmer) and protein content of the remaining lysate was measured for normalisation of results (BCA Protein Assay Kit; #23225, Thermo Fisher Scientific, Rockford, IL).

Glycogen synthesis. Incorporation of glucose into glycogen was performed using similar methods detailed elsewhere (Abdelmoez et al., 2020). Briefly, myotubes were nutrient-starved as in glucose uptake experiments and then treated with 0, 10, or 120 nM of insulin in fresh serum-starvation medium for 30 min before adding 2 μL.mL^{-1} D-[^{14}C(U)] glucose (NEC042B005MC; Perkin Elmer) for a further 90 min. Cells were washed three times with ice-cold PBS and lysed in 0.5 mL 0.03% SDS. Thereafter, [^{14}C]-labelled glycogen was extracted from 0.4 mL of the lysate and counted in a liquid scintillation counter (Abdelmoez et al., 2020). Cellular protein concentration was determined by bicinchoninic acid (BCA) assay from the remaining 0.1 mL of cell lysate and used to normalize results.

Glucose oxidation. Glucose oxidation was performed using methods as described (Abdelmoez et al., 2020). Myotubes were washed with PBS and incubated with D-[^{14}C(U)] glucose in serum starvation medium in the presence or absence of 2 μM carbonyl cyanide p-trifluoro methoxyphenylhydrazone (FCCP) for 4 h. Thereafter, [^{14}C]-CO$_2$ was released from the medium by adding of 150 μL 2 M HCl per mL of medium and captured in 300 μL of 2 M NaOH for 1 h. The captured [^{14}C]-CO$_2$ was counted using a liquid scintillation counter. Cells were washed three times with ice-cold PBS and homogenised in 0.4 mL of 0.5 M NaOH. After complete lysis, 0.1 mL of 2 M HCl was added to neutralise pH and protein was assessed using the BCA assay for normalisation of results.

Lactate assay. Lactate released into culture supernatant was measured in post-fusion medium (Abdelmoez et al., 2020) after 48 h of basal or (2 μM) FCCP-stimulated conditions using a colourimetric assay as described (Abdelmoez et al., 2020). Assay buffer contained Tris (50 mM pH 8), NAD (7.5 mM), N-methylphenazonium methyl sulfate (250 μM), p-iodonitrotetrazolium violet (500 μM), and lactate dehydrogenase (4 U.mL^{-1}). Medium was passed through 3 kDa filters by centrifugation (14000 g for 15 min at 4°C) and 50 μL of sample was mixed with 150 μL of assay buffer. After a 20-min incubation the absorbance was read at 490 nm. Results were normalised to total cellular protein content as measured by BCA assay after lysis in 0.03% SDS.

Fatty acid oxidation. [^{14}C(U)] palmitic acid (NEC534050UC; Perkin Elmer) oxidation was performed using the same experimental procedure as glucose oxidation, except that 1 μL.mL^{-1} of isotope was added to base glucose-free DMEM (Gibco #11966) supplemented with 25 μM of unlabelled BSA-conjugated palmitate.

Thin-layer chromatography (TLC). Myotubes were incubated for 6 h in the same medium as described for [^{14}C(U)] palmitic acid oxidation. Cells

were then washed three times with ice-cold PBS and hydrophobic lipids were extracted as detailed elsewhere (Massart, Zierath et al., 2014). The resulting lipid pellet was eluted in 50 µL of a methanol:chloroform solution (1:1) and loaded on a TLC plate (Silica Gel G 250 µm 20×20 cm, Analtech, DE, USA). Lipid species were separated for 30 min in a loading chamber filled with 100 mL of a hexane:diethylether:acetic acid mixture (80:20:3). The loaded TLC plate was then transferred to an exposure cassette (GE Healthcare) and exposed to an X-ray film for four weeks at -80°C before developing in an X-ray developer machine. Identification of the appropriate bands was determined against standards for 1,2-Dioctanyl [1-^{14}C] rac-glycerol (1,2-DAG), 1,3-Dioleoyl-rac-glycerol [oleoyl-1-^{14}C] (1,3-DAG) (American Radiolabeled Chemicals Inc.), [^{14}C(U)] palmitic acid (Perkin Elmer), and [^{14}C] triolein (TG). Densitometric quantification was performed using Image Lab software (Bio-Rad).

Phenylalanine incorporation into protein. Myotubes were washed once with PBS and incubated in low-glucose base DMEM for 4 h. After starvation, media was changed to low-glucose DMEM supplemented with an additional 25 µM unlabelled phenylalanine and 4 µL of [^{14}C]phenylalanine (NEC284E050UC; Perkin Elmer) per mL for 6 h, with or without 20% FBS plus 10 mM leucine, and in the presence or absence of 100 nM rapamycin or 10 µM lactacystin. Cells were then washed three times with ice-cold PBS and lysed in 0.5 mL 0.03% SDS. Protein concentration was measured from 0.1 mL of cell lysate by BCA protein assay. Protein from the remaining cell lysate was precipitated in 50% Trichloroacetic acid with 1% BSA overnight at −20°C, followed by centrifugation (12000 g for 15 min at 4°C). Supernatant was discarded and the protein pellet washed twice in acetone with centrifugation after each wash (12000 g for 15 min at 4°C). Acetone was discarded and the protein pellet dissolved in 0.5 M NaOH for 1 h in a heating block at 65°C with shaking. The dissolved pellet homogenate was transferred to scintillation vials and the amount of [^{14}C] phenylalanine incorporation determined by scintillation counting.

Subcellular fractionation. Separation of sarcoplasmic and nuclear fractions was performed on fresh cells, without freezing. Myotubes were washed once in ice-cold PBS and scraped in cell lysis buffer (10 mM HEPES pH 7.5, 10 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% NP-40) supplemented with 1 mM PMSF, 0.5 mM Na$_3$VO$_4$ and 1x Protease Inhibitor Cocktail Set 1 (PIC) (Calbiochem). Lysates were mixed by pipetting and rotated at 4°C for 20 min then subjected to centrifugation (12000 g for 10 min at 4°C). The sarcoplasmic fraction (supernatant) was collected and stored at -80°C. The remaining pellet was washed three times in ice-cold cell lysis buffer by inverting the Eppendorf multiple times per wash and then resuspended in nuclear extraction buffer (20 mM HEPES pH 7.5, 400 mM NaCl, 1 mM EDTA, 1 mM DTT) supplemented with 1 mM PMSF, 0.5 mM Na$_3$VO$_4$ and 1x PIC. Resuspension was performed in the same volume collected for the sarcoplasmic fraction to retain the sarcoplasmic to nuclear ratio of proteins. Samples were next rotated at 4°C for 30 min and debris pelleted by centrifugation (12000 g for 15 min at 4°C). The nuclear fraction supernatant was collected and stored at -80°C until lysate preparation for immunoblotting.

Surface sensing of translation (SUnSET) assay. Relative rates of in vitro protein synthesis were determined using the SUnSET method as described (Goodman et al., 2011). Briefly, myotubes were washed once in PBS and incubated in low-glucose base DMEM for 4 h. Media was then changed to fresh low-glucose vehicle DMEM or low-glucose DMEM supplemented with 120 nM insulin and 10 mM leucine, with or without 100 nM rapamycin or 178 µM cycloheximide (*NR-203*^{Oex} experiment only). After 1 h of treatment, puromycin (1 µM) was added to the medium and cells were incubated for a further 30 min before collection. Medium was aspirated and cells were washed once in ice-cold PBS. Samples were then harvested in protein lysis buffer (137 mM NaCl, 2.7 mM KCl, 1 mM MgCl$_2$, 1% Triton X-100, 10% glycerol, 20 mM Tris pH 7.8, 10 mM NaF, 1 mM EDTA, 1 mM PMSF, 0.5 mM Na$_3$VO$_4$ and 1x PIC), rotated for 30 min at 4°C and subjected to centrifugation (12,000 g for 15 min at 4°C). Supernatants were stored at -80°C until protein determination by BCA assay and subsequent sample preparation for immunoblotting.

Insulin stimulation for assessment of signalling. For interrogation of the canonical insulin signalling cascade, myotubes were incubated in serum starvation medium for 4 h. Cells were then stimulated with 0, 10, or 120 nM of insulin in fresh serum starvation medium for 20 min and lysed as mentioned for SUnSET assay samples.

Immunoblot analysis. Samples were prepared for SDS-PAGE with Laemmli buffer (60 mM Tris pH 6.8, 2% w/v SDS, 10% v/v glycerol, 0.01% w/v bromophenol blue, 1.25% v/v β-mercaptoethanol). Equal amounts of protein were loaded and separated on Criterion XT Bis-Tris Gels (Bio-Rad, Hercules, CA) and transferred to polyvinylidene fluoride membranes (Merck, Germany). Membranes were stained with Ponceau S to confirm transfer quality and even loading of samples. Membranes were then blocked with 5% non-fat milk in Tris-buffered saline supplemented with Tween-20 (TBST; 20 mM tris-HCl pH 7.6,

137 mM NaCl, 0.02% Tween-20) for 1 h at room temperature, before an overnight incubation with primary antibodies at 4°C with gentle rocking. Primary antibodies were diluted in TBS plus 0.1% w/v bovine serum albumin and 0.1% w/v NaN_3 or 5% non-fat milk in TBST (specific antibodies and dilutions are available upon request). Membranes were next washed with TBST and incubated with species-appropriate horseradish peroxidase-conjugated secondary antibody (1:25000 in TBST with 5% non-fat milk). Proteins were visualised by enhanced chemiluminescence (Amersham ECL Western Blotting Detection Reagent, UK) and quantified by densitometry (QuantityOne or Image Lab software, Bio-Rad).

Total protein and RNA. Unless stated otherwise, total protein concentrations reported in figures are from BCA protein assay measurements of basal conditions in metabolic assays. Compared to sample lysis for immunoblot analysis, homogenisation of cell monolayers for normalisation of metabolic assays was performed in larger volumes of buffer. Thus, these measurements more accurately reflect total protein concentration per well. In cases where total protein and RNA was determined for multiple experiments using primary cells from the same donor, values were log2 transformed, Z-scored and the average Z-score across assays was used for statistical analysis.

Immunocytochemistry. Myotubes were immuno-stained for fast myosin heavy chain isoforms (MYH1/2, 1:250; sc-53088, Santa Cruz Biotechnology) using methods as described (Abdelmoez et al., 2020), with the addition of nuclear counterstaining. After 90 min incubation with Alexa Fluor 594 goat anti-mouse (A-11005) secondary antibody (1:500) at room temperature

with gentle rocking, cells were washed twice with 0.025% Tween-20 in PBS and counterstained with 300 nM 4′,6-diamidino-2-phenylindole (DAPI) in PBS for 5 min at room temperature. Myotubes were then washed twice in PBS and kept in PBS protected from light at 4°C until imaging. MYH1/2 and nuclei images were taken using 20x magnification on a Zeiss Axio Vert.A1 inverted fluorescent microscope, equipped with ZEN software (Carl Zeiss Microscopy GmbH, Germany). Images were obtained from three random fields of view per well across 3 technical replicate wells, producing nine images per condition for each donor. MYH1/2 images were converted to 8-bit binary and quantification of MYH1/2 area (μm^2) per field was automated using ImageJ software (Fiji) to prevent measurement bias. Nuclei were counted manually using the ImageJ cell-counter plugin. Fused myonuclei were considered nuclei counted within the MYH1/2 area automatically detected by ImageJ software in the MYH1/2 analysis and fusion index calculated as the percentage of fused versus total nuclei per field of view.

Statistics. Analyses were performed using either R 4.1.0 (www.r-project.org) or GraphPad Prism 10.0.3 software (GraphPad Software Inc.). Exact p-values are specified for comparisons where probability met $p<0.1$. Data are presented as box-and-whisker plots with Tukey distribution and pairing by donor unless specified otherwise in figure legends. Normality was assessed using Shapiro-Wilk test before applying appropriate parametric or non-parametric tests. In 2-way ANOVA analyses, Tukey ladder transformation was used when residuals violated assumptions of normality. Statistical tests and sample sizes are described in figure legends. Bioinformatic analyses are described in their respective method sections.

Main Text Figures

Figure 1. Downregulation of *NR4A3* during inactivity correlates with transcripts involved in myogenic and metabolic pathways. A. Meta-analysis of transcriptomic inactivity studies in human skeletal muscle. **B.** Transcriptomic studies were pooled by the duration of inactivity protocols: less than one week, one-two weeks or more than two weeks. Datasets were normalized as described in methods. **C.** Gene set enrichment analysis with the Gene Ontology dataset was performed on genes ranked on Spearman correlation with *NR4A3* across all transcriptomic studies of inactivity. **D.** The two studies that explored effects of reloading after inactivity were merged and analysed as described in methods.

Figure 2. *NR4A3* silencing in primary human myotubes attenuates glucose oxidation by diverting glucose towards lactate production. Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** mRNA expression of NR4A family members *NR4A3*, *NR4A1* and *NR4A2* measured by RT-qPCR. n = 6, 2-way ANOVA (silencing x gene) with Šidák correction. **B.** Protein level of NR4A3 in cytosolic and nuclear fractions assessed by immunoblot analysis. Results are mean ± SEM, n = 2. **C.** Rates of radiolabelled glucose oxidation under basal and (2 µM) FCCP-stimulated conditions over 4 h. n = 6, 2-way ANOVA (silencing x FCCP) with Šidák correction. **D.** Lactate concentration in cell supernatant measured after 48 h of basal or (2 µM) FCCP-stimulated conditions. n = 7, 2-way ANOVA (silencing x FCCP) with Šidák correction. Overall statistical model effects are stated in figures. **E.** mRNA expression of genes involved in glucose metabolism measured by RT-qPCR. Results are box-and-whisker plots with Tukey distribution. n = 6, paired t-tests with FDR correction. **F.** Representative immunoblot and quantification of pyruvate dehydrogenase E1 Subunit Alpha 1 (PDHA1) phosphorylation at ser[293]. n = 5, 2-way ANOVA (silencing x insulin + leucine).

Figure 3. *NR4A3* silencing upregulates fatty acid oxidation and AMPK signalling. Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** Rates of radiolabelled [14]C palmitic acid oxidation under basal and (2 μM) FCCP-stimulated conditions over 4 h. n = 6, 2-way ANOVA (silencing x FCCP) with Šidák correction. **B.** Detection of select lipid species by thin layer chromatography (TLC) of lipid extracts from myotubes after 6 h of basal or (2 μM) FCCP-stimulated conditions. TG = triacylglycerides, FFA = free fatty acids, DAG = diacylglycerol, PA + other = phosphatidic acid plus all other non-migrating lipid species. Results are the mean ± SEM of each lipid species as a percentage of total detected lipid species within condition. n = 5, 3-way ANOVA (silencing x lipid species x FCCP). Overall statistical model effects are stated in figures. **C.** Immunoblot of AMPK-signalling proteins from a representative donor and quantification of phosphorylated-to-total protein ratios. n = 5, 2-way ANOVA (silencing x insulin + leucine) with Šidák correction. Only basal data are shown. **D.** Representative immunoblot and quantification of proteins involved in lipid storage, mobilisation, and transport. n = 5, 2-way ANOVA (silencing x insulin + leucine) with Šidák correction. Only basal data are shown and overall effects of silencing in the model are presented (si*N* effect, p<0.1). **E.** mRNA expression of genes involved in lipid metabolism measured by RT-qPCR. Results are box-and-whisker plots with Tukey distribution. n = 6, paired t-tests with FDR correction.

Figure 4. *NR4A3* silencing impairs mTORC1 signalling, ribosomal biogenesis, and translation. Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** Total protein per well and **B.** RNA per µL. Results are the average Z-score of log2-transformed values across experiments. n = 8 for protein and n = 9 for RNA, paired t-test. **C.** Representative immunoblot and quantification of cellular protein synthesis assessed by protein puromycilation (i.e. SUnSET method). n = 5, 2-way ANOVA (silencing x insulin + leucine) with Šidák correction. Overall statistical model effects are stated in the figure. **D.** Immunoblot of proteins involved in AMPK-mediated mTORC1 inhibition and AKT-mTORC1 pathways from a representative donor. **E.** Schematic representation of immunoblot analysis. n = 5, 2-way ANOVA (silencing x insulin + leucine) with Šidák correction. Quantification of each depicted signalling event can be found in **Supplementary Figure 2**. **F.** mRNA expression of genes involved in amino acid (AA) metabolism and protein synthesis measured by RT-qPCR. Results are box-and-whisker plots with Tukey distribution. n = 6, paired t-tests with FDR correction. 'Ribo-' = ribosomal.

Figure 5. Overexpression of the canonical *NR4A3-203* isoform increases protein synthesis. Primary skeletal muscle cells were transduced with an empty vector control plasmid (EV) or a plasmid containing variant *NR4A3-203* (*NR-203*^Oex). **A.** mRNA expression of NR4A family members *NR4A3*, *NR4A1* and *NR4A2* measured by RT-qPCR. n = 6, 2-way ANOVA (overexpression x gene) with Šidák correction. Also, protein level of NR4A3 from two indicative donors. **B.** Representative immunoblot and quantification of cellular protein synthesis assessed by protein puromycilation (i.e. SUnSET method). n = 5, 3-way ANOVA (overexpression x rapamycin x insulin+leucine) with FDR correction. Veh = vehicle, Rapa = rapamycin, and Cyclo = cycloheximide treatments, respectively. **C.** Rates of radiolabelled phenylalanine incorporation to protein under basal and (20%) FBS plus (10 mM) leucine-stimulated conditions over 6 h, with or without mTORC1- (rapamycin) or proteasomal- (lactacystin) inhibition. n = 6, 3-way ANOVA (overexpression x FBS+Leucine x compound) with FDR correction. **D.** Total protein per well from same experiment as C. Overall statistical model effects are stated in figures.

Figure 6. Overexpression of the canonical NR4A3 isoform partially restores glucose oxidation after global *NR4A3* silencing in myotubes, concomitant with rescued mTORC1 signalling. A. Schematic representation of the *NR4A3* 'rescue' experiment. **B.** *NR4A3* mRNA expression measured by RT-qPCR. n = 6, 1-way ANOVA with Tukey correction. **C.** Protein level of NR4A3 from one indicative donor. Con = siScr/EV, si*N* = si*NR4A3*/EV, and *N*^Oex = si*NR4A3*/*NR-203*^Oex conditions, respectively. +'ve = positive control from nuclear fraction lysate of *NR-203*^Oex. **D.** Rates of radiolabelled glucose oxidation under basal and (2 µM) FCCP-stimulated conditions over 4 h. n = 7, 2-way ANOVA (overexpression x FCCP) with Fisher's LSD post-test. Overall statistical model effects are stated in relevant figures. **E.** Immunoblot of proteins in the mTORC1 pathway from a representative donor. Quantification of each depicted signalling event can be found in **Supplementary Figure 5**.

Figure 7. *NR4A3* silencing reduces myotube size and alters the expression of muscle contractile and structural apparatus. Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** Immunocytochemistry of fast myosin heavy chain isoforms (MYH1/2, red) and nuclei (DAPI, blue) from a representative donor. Scale bar = 100 µm. Quantification of immunocytochemistry for **B.** MYH1/2 area, **C.** total nuclei, and **D.** fusion index per field of view. Results are violin plots with median and interquartile range. Circles represent measurements from different fields of view across three technical replicates per donor. n = 3, nested paired t-test. **E.** Representative immunoblot and quantification of proteins involved in contraction, structure, and myogenic transcription. Results for MYH1/2, MYH7, DES, and MYOG are the average Z-score of log2-transformed values across experiments (n = 9, paired t-test). Results for pan-Actin are n = 5, 2-way ANOVA (silencing x insulin + leucine) with only basal data shown. **F.** mRNA expression of genes involved in myogenesis, muscle structure and contraction measured by RT-qPCR. Results are box-and-whisker plots with Tukey distribution. n = 6, paired t-tests with FDR correction. 'SR' = sarcoplasmic reticulum.

Supplementary Figures

Supplementary Figure 1. *NR4A3*-silenced myotubes retain responsiveness to insulin stimulation. Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** Representative immunoblot and **B-G.** quantification of proteins and phosphorylation events in the canonical insulin signalling cascade. n = 6, 2-way ANOVA (silencing x insulin) with Šidák correction. Overall statistical model effects are stated in figures. **H.** Rates of radiolabelled 2-Deoxy-D-glucose uptake (n = 7) and **I.** glycogen synthesis (n = 6) under basal and insulin-stimulated conditions. 2-way ANOVA (silencing x insulin) with Šidák correction. **J.** Representative immunoblot and quantification of total glycogen phosphorylase, muscle associated (PYGM) and **K.** lactate dehydrogenase (LDH). n = 5, 2-way ANOVA (silencing x insulin + leucine). Only basal data are shown.

Supplementary Figure 2. *NR4A3* **silencing attenuates mTORC1 signalling.** Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** Immunoblot quantification of phosphorylated-to-total protein ratios for AMPK target sites on raptor (ser792) and TSC2 (ser1387). **B.** Immunoblot quantification of AKT-mTORC1 activating phosphorylation sites and total protein levels for Raptor, **C.** mTOR, **D.** RPS6, **E.** 4EBP1 (with images from a representative donor), **F.** TSC2, and **G.** AKT. All immunoblot analyses are n = 5, 2-way ANOVA (silencing x insulin + leucine) with Šidák correction; except 4EBP1, which is n = 6, 2-way ANOVA (silencing x insulin) with Šidák correction, and AKT(ser473) and AKT(thr308) phosphorylation, which are n = 5, paired t-tests. Overall statistical model effects are stated in figures. si*N* = main silencing effect, I+L = main insulin + leucine effect, si*N* x I+L = interaction effect.

Supplementary Figure 3. *NR4A3* **silencing minimally impacts pathways controlling protein degradation.** Primary skeletal muscle cells were exposed to a control scramble sequence (siScr) or a silencing RNA targeting *NR4A3* (si*NR4A3*). **A.** mRNA expression of genes related to the negative regulation of muscle mass, endoplasmic reticulum (ER) stress, oxidative stress, and inflammation measured by RT-qPCR. Results are box-and-whisker plots with Tukey distribution. n = 6, paired t-tests with FDR correction. **B.** Immunoblot of proteins involved in inflammation, ubiquitin-proteasomal, and autophagy-lysosomal degradation pathways from a representative donor. **C-G.** Immunoblot quantification of **C.** NF-κB inhibitor IκBα, basal and insulin-mediated inhibitory phosphorylation of transcription factors **D.** FOXO3a, **E.** FOXO1, and E3 ubiquitin ligases **F.** MURF1 and **G.** MAFbx. **H.** Representative immunoblot and quantification of cellular protein ubiquitination. **I-K.** Immunoblot quantification of **I.** Calpain-1 and **J.** Caspase 3 proteases, and **K.** alpha subunits of the 20S core particle proteasome (Pan 20Sα). **L-N.** Immunoblot quantification of autophagy markers **L.** ULK1 phosphorylation at ser317, and total **M.** P62 and **N.** LC3-I/LC3-II abundance. All immunoblot analyses are n = 5, 2-way ANOVA (silencing x insulin + leucine) with Šidák correction. Overall statistical model effects are stated in figures.

Supplementary Figure 4. Overexpression of the canonical *NR4A3* isoform increases translation without altering glucose and fatty acid metabolism, RNA abundance, or mTORC1 signalling. Primary skeletal muscle cells were transduced with an empty vector control plasmid (EV) or a plasmid containing variant *NR4A3-203* (*NR-203*[Oex]). **A.** Basal and stimulated rates of radiolabelled 2-Deoxy-D-glucose uptake, **B.** glycogen synthesis, **C.** glucose oxidation, and **D.** [14]C palmitic acid oxidation. n = 6, 2-way ANOVA (silencing x stimulation). Overall statistical model effects are stated in figures. **E.** Total RNA concentration (ng.µL[-1]). n = 6, paired t-test. **F.** Immunoblot of proteins in the mTORC1 and ERK signalling cascades from a representative donor. Samples are those presented in main **Figure 5B**. No difference in phosphorylation or total protein levels were detected with *NR-203*[Oex]. n = 5, 3-way ANOVA (overexpression x rapamycin x insulin + leucine). **G.** Volcano plot of all mRNA transcripts assessed after *NR-203*[Oex] measured by RT-qPCR. Genes were the same as analysed in *NR4A3*-silenced conditions. n = 6, paired t-tests with FDR correction.

Supplementary Figure 5. Reconstitution of the canonical *NR4A3* isoform after global NR4A3-depletion restores mTORC1 signalling but does not ameliorate AMPKα phosphorylation or fatty acid oxidation. Primary skeletal muscle cells were depleted of all *NR4A3* transcripts using siRNA and then transduced with an empty vector control plasmid or a plasmid containing variant *NR4A3-203* as described in main **Figure 6A**. **A-C.** Immunoblot quantification of total protein levels and phosphorylation events in the P70S6K-RPS6 and 4EBP1 branches of mTORC1 signalling. **D.** Representative immunoblot and quantification of total and phosphorylated AMPKα. All immunoblot analyses are n = 6, and either 1-way ANOVA with Tukey correction or Friedman test with Dunn's correction. **E.** Rates of radiolabelled palmitic acid oxidation under basal and (2 µM) FCCP-stimulated conditions over 4 h. n = 6, 2-way ANOVA (overexpression x FCCP) with Fisher's LSD post-test. Overall statistical model effects are stated in figures.

A

Supplementary Figure 6. The effect of *NR4A3* silencing on contractile proteins is specific to siRNA targeting *NR4A3* transcripts. Primary skeletal muscle cells were exposed to a control scramble sequence (Scr) or a silencing RNA targeting *NR4A3* (si*N*). **A.** Representative immunoblot of myosin heavy chain isoforms IIX (MYH1) and IIA (MYH2) from three independent donors. 'NT' = non-transfected condition.