Estradiol and Flutamide Effects on the Song System of Developing Male Zebra Finches

William Grisham[1] & Mary Ellen McCormick[2]

[1]Department of Psychology and Brain Research Institute, University of California, Los Angeles, CA 90095-1563

[2] Mary Ellen McCormick, Huntington Hospital, 100 West California Blvd Pasadena, CA 91105

address correspondence to:

William Grisham, Ph.D.
Department of Psychology
502 Portola Plaza
University of California, Los Angeles
Los Angeles, CA 90095-1563
Email: wgrisham@g.ucla.edu

Abstract

Estradiol (E$_2$) masculinizes the developing song system of female zebra finches (Taeniopygia castanotis) if it is administered in early life, but its effect is blocked with the co-administration of an antiandrogen (Flutamide). The effects of E$_2$ on the developing male song system are not uniform and reports of Flutamide administration in developing male zebra finches differ in their findings. Therefore. this study was conducted to further explore the effects of administering E$_2$ alone, Flutamide (Flut) alone, or the two in combination during early post-hatch development. Brains and testes were examined after day 100. The results showed definite demasculinizing effects of early E$_2$ on the song nucleus HVC (proper name)—its volume and neuron number were markedly reduced. Nonetheless, early E$_2$ hypermasculinized HVC neuronal size. Flut slightly hypermasculinized RA volume (Robust nucleus of the Arcopallium), which replicates one prior study but the absence of additional effects is at odds with others. Early E2 resulted in markedly reducing testes size, which is likely to be a consequence of hijacking endogenous endocrine feedback mechanisms. Arguments are put forward suggesting 1) early E2 action on HVC could be an anachronistic consequence of actions on the genotype of developing male versus females or 2) a disruption of endocrine mechanisms inducing inappropriate hormonal states during development. These possibilities are not mutually exclusive.

Keywords: songbird, androgen antagonist, estradiol, endocrine system, sex differences

Introduction

Song behavior in zebra finches (Taeniopygia castanotis) is sexually dimorphic; males sing and females do not. This sex difference in behavior is reflected by dramatic sex differences in its neural underpinnings. Males have larger song nuclei such as HVC, RA—robust nucleus of the arcopallium, and Area X (Grisham & Arnold, 1995), with more neurons in HVC and RA (Nordeen & Nordeen, 1988; Schlinger & Arnold, 1991), and larger neurons in HVC, lMAN—Lateral Magnocellular nucleus of the Anterior Nidopallium, and RA (Nottebohm & Arnold, 1976; Grisham & Arnold, 1995; Arnold & Saltiel, 1979; Nordeen et al., 1992).

The female's song system can be masculinized by administering estradiol (E_2) during development (Adkins-Regan et al., 1994; Gurney & Konishi, 1980; Gurney, 1981, 1982; Jacobs, Grisham & Arnold, 1995; Simpson & Vicario, 1991). The influence of early E_2 on song system development in males is less clear, however.

Two studies report that E_2 has no effect on the song system of developing male zebra finches either when administered at hatching (Gurney & Konishi, 1980) or *in ovo* (Wade et al., 1997). Nonetheless, Tang and Wade (2009) found early E_2 in males increased Area X volume when examined at day 25 posthatch but had no impact on HVC. Conversely, when males were treated for the first 25 days posthatch with estradiol benzoate (EB) and examined at day 25, the volume of Area X and HVC were demasculinized (Mathews & Arnold, 1991), but not when males were treated until day 20

then examined at day 60 when early EB treatment hypermasculinized male neuronal sizes in lMAN and HVC (Mathews & Arnold, 1991).

Mixed effects of androgenic manipulations in development also have been reported. Blocking androgen action with Flutamide (Flut) at hatching hypermasculinized RA volume and the number of its neurons (Schlinger & Arnold, 1991). But, life-long exposure to Flut demasculinized RA neuron number and slightly decreased HVC volume (Grisham et al., 2007). Also, when it was administered at day 20 to castrates, it demasculinized both lMAN and Area X but did not influence other song regions (Bottjer & Hewer, 1992). Blocking synthesis of an androgen, dihydrotestosterone, at hatching reduced the number and density of RA neurons (Grisham et al., 1997).

Administering Flut at hatching along with E$_2$ blocked the usual E$_2$ induced masculization of female zebra finch song system (Grisham et al., 2002). Given the mixed results of both estrogen and androgenic manipulations of males in early life, we decided to re-examine the effects of these treatments as well as their combination. We assessed the effects of both E$_2$ and Flut alone and in combination on hatchling males, particularly since we had found such dramatic effects on the combined treatment in hatchling females.

Methods

All aspects of this study were approved by the Animal Research Committee of the University of California, Los Angeles. Further, the ARRIVE guidelines 2.0 were followed.

Estradiol (E$_2$) and flutamide (Flut) were administered in a 2x2 design. Males were implanted within 1-3 days of hatching under the skin of the breast with E$_2$ pellets, Flut pellets, both, or neither. The E$_2$ implants were made by mixing finely ground E$_2$ with Medical grade Silastic glue in a ratio of 1:6 and extruding this mixture into polyethylene tubing (Clay Adams no. 7411, i.d. 0.58 mm, o.d. 0.965 mm). These implants were cut into 2 mm sections, weighed, and their average dose calculated to be 83 μg (batches of pellets ranged from 73 μg to 87 μg E$_2$). The Flut implants were made in a manner described by Schlinger and Arnold (1991). Briefly, Flut was mixed with Silastic glue and then extruded through a 1cc syringe without a needle. The resulting strips were left to cure overnight, quartered, weighed, and cut into lengths that would result in a dose of 200 μg. Each bird received two pellets (E$_2$ $_+$ blank, E$_2$ + Flut, Flut + blank, or two blanks) except for unimplanted controls.

All dependent measures were made blind to the birds' treatment group. At sacrifice (101-136 days of age), the birds were deeply anesthetized and then perfused with 0.75% saline followed by 10% formalin in saline. Birds were only included if the pellets could be found at sacrifice. (Birds in the blank+blank group were included regardless.) Even after having been implanted for 101 days or more, the E$_2$ pellets all appeared to contain E$_2$ crystals (they had white opacities). The Flut pellets were clear and appeared depleted. The testes were dissected and stored in 10% formalin until they were

weighed; at which time they were cleaned of extraneous tissue, dabbed dry, and weighed to the nearest 0.1 mg (if testes weighed less than .1 mg, they were assigned a 0.1 mg weight). Testes weights were taken from 26 adult males (n = 5 E$_2$ treated; n = 4 E$_2$+Flut treated; n=8 Flut treated; and 9 untreated/blanks) and both testes were averaged for each individual.

The brains were dissected, stored in 10% formalin, and frozen-sectioned at 40 μm in the coronal plane. Brain measures were taken from 28 individuals (n = 5 E$_2$-treated; n = 4 E$_2$+Flut treated; n = 8 Flut treated; n = 11 untreated/blanks), but due to histological problems HVC data was n = 26. Data and records can be accessed via UCLA Dataverse at https://doi.org/10.25346/S6/VNSKRP .

The brains were examined with a light microscope connected to a computer via a videocamera. Cross-sectional areas of song system nuclei and their individual neurons were measured using NIH Image (http://rsb.info.nih.gov/nih-image/).

The volumes of Area X, lMAN, HVC, and RA were calculated by means of the cylindrical method: tracing the cross-sectional areas on every third section magnified at 6.25X, adding these areas, and multiplying by the sampling interval (120 μm). The cross-sectional volumes of each song system nucleus were averaged across hemispheres for each animal unless one hemisphere was damaged.

The area of individual neurons was measured at 800X. Twenty-five neurons were sampled through the rostral-caudal extent of the nucleus in each hemisphere for a total of fifty neurons in each song control nucleus of each animal. Neurons were distinguished from glia by their dark staining, ample cytoplasm, and nuclei containing only one or two nucleoli. Glia were distinguished by light staining, little cytoplasm, often with several

nucleoli in each nucleus. The number of neurons in HVC was determined by counting the number of nucleoli in twenty-five frames (each 45,350 μm^3) that were sampled throughout its rostral-caudal extent in both hemispheres. Counting a small profile like nucleoli produces counts as reliable as using an optical dissector (Tramontin et al., 1998). The average density of neurons was calculated for each animal and multiplied by the mean volume of the given song system nucleus.

Dependent variables were analyzed via JASP https://jasp-stats.org/ and using a 2x2 ANOVA: estradiol treatment vs. none as one factor and Flut treatment vs. none as the other factor. Testes weights were analyzed by an ANOVA with 2x2 between (E$_2$ vs. none and flut vs. none) and one within variable (side).

Results

HVC volume was significantly demasculinized (reduced) by E$_2$ treatment, $F(1,22) = 8.38$, $p < 0.01$, $\eta^2 = 0.272$ (Figure 1A), as was the number of HVC neurons, $F(1,22) = 4.51$, $p < 0.05$, $\eta^2 = 0.168$ (Figure 1B). Despite E$_2$'s demasculinizing effect on the volume of HVC and on the number of its neurons, it hypermasculinized the size of HVC neurons, $F(1,24) = 6.00$, $p < 0.05$, $\eta^2 = 0.195$ (Figure 1C). There was a weak effect/strong trend for Flut hypermasculinizing RA volume, $F(1,24) = 4.133$, $p = 0.053$, $\eta^2 = 0.131$ (Figure 1D). No other brain measures were affected by either E$_2$ or Flut alone or in combination (all p values > 0.09).

Testes weight was markedly reduced by E$_2$ treatment, $F(1,69) = 74.63$, $p < 0.0001$, $\eta^2 = 0.475$ (Figure 2A). There was no effect of Flut alone or an E$_2$ x Flut

interaction on testes size, all $\underline{p}s > 0.50$ (Figure 2A). Both HVC volume and the number of HVC neurons significantly correlated with mean testicular weight, $r(22) = .576$, $\underline{p} < 0.01$ and $r(22) = .487$, $\underline{p} < 0.02$, respectively (Figure 2B & 2C).

Discussion

Early administration of E₂ to male hatchlings dramatically reduced HVC volume and the number of its neurons when examined in adulthood (Fig. 1A, 1B), which contrasts sharply with the masculinizing effects of early E₂ in females (Adkins-Regan et al., 1994; Grisham & Arnold, 1995; Gurney & Konishi, 1980; Gurney, 1981, 1982; Jacobs et al., 1995; Simpson & Vicario, 1991). Early E₂ administration to males could 1) differentially affect the male and female genome, possibly triggering temporally inappropriate genetic cascades, and/or 2) interfer with hormonal feedback mechanisms and gonadal function.

There is evidence of early E₂ differentially altering BDNF gene expression in male but not female HVC (Dittrich et al., 1999). This result suggests that early E₂ disrupts the temporal sequence of male gene expression, which could alter the adult phenotype. Early E₂ administration also has opposite effects on gene expression in females versus males; the number of HVC neurons expressing 17β-hydroxysteroid dehydrogenase mRNA decreased in males but increased in females when birds were examined at day 25 (Thompson et al., 2011). Early E₂ also alters genes in female Area X toward the masculine pattern whereas it induces more female-like patterns in male lMAN and RA (Choe et al., 2021), the latter of which is a target of HVC projections (Benezra et al., 2018). Thus, early E₂ could have its action on male HVC via its axonal projections to RA.

Temporal differences in early E$_2$ administration effects may explain some discrepancies in the literature. Notably, E$_2$ at hatching decreases the number of BrdU-labeled cells added to HVC in males at puberty whereas it increases the number of labeled cells in females (Tang and Wade, 2009). Choe et al. (2021) did not detect any effect of early E$_2$ on male HVC volume, but we did. Nonetheless, several methodological differences could explain this discrepancy: different dose regimens; different ways of defining HVC, and very different time points when the song systems were measured (day 30, vs. > 100 days of age). HVC neurons are added well after day 30 (Walton et al., 2012; Diez et al., 2021), which could also explain the discrepancy between the data of Choe et al. (2021) and ours. In agreement with our findings, however, Choe et al. (2021) also found early E$_2$ increased HVC cell size in males.

Hormonal mechanisms could have played a role on the brain and gonadal development in the present study. The dramatic decrease in testes size induced by early E$_2$ administration (Fig. 2A) suggests that there are estrogen-mediated endocrine feedback mechanisms in zebra finches. This finding is paralleled in other studies on zebra finches (Mathews & Arnold, 1991), and other bird species (Casto & Ball, 1996; Lorenz, 1954; Soma et al., 2000). So, our early E$_2$ exposure could have permanently altered hormonal feedback mechanisms resulting in small testicular size.

Notably, HVC volume and the number of its neurons were significantly correlated with testes size (Fig 2B, 2C). The testes in our early E$_2$-treated birds were more like those of babies rather than adults suggesting that they never matured. The best estimate of the onset of puberty in zebra finches is about Day 60-70 posthatch (Bölting & von Englehardt, 2017; Pröve, 1983) and we did not examine the birds until after Day 100.

HVC neurons are still being added up until Day 70 (Diez et al., 2021) and even into adulthood (Walton et al., 2012). Testosterone increases the recruitment and/or survival of HVC neurons in adult female canaries (Rasika et al., 1994), an effect that is dependent upon the androgenic rather than estrogenic metabolites of testosterone (Fusani et al., 2002). If the markedly reduced testes size in our birds resulted in reduced testicular secretions not only before but also after puberty, then demasculinizing effects could be a consequence.

The number of androgen target cells in HVC normally increases during adolescence in male zebra finches (Bottjer, 1987; Tang & Wade, 2010). Early E_2 treatment decreases androgen receptor levels in male but not female HVC (Thompson et al., 2011), so there may have been not only low androgen levels, but also fewer receptors upon which to act.

Our early antiandrogen (Flut) treatment showed a strong trend toward hypermasculinizing RA volume (Figure 1D) similar to Schlinger and Arnold (1991)—we used the same protocol as they did. We did not replicate the Flut effects of Bottjer and Hewer (1992) or Grisham et al. (2007) on the song system or on testes size (Fig. 2A), but the protocol of these other studies was quite different in timing of Flut administration. Our Flut pellets in this experiment were likely to be depleted before the birds were examined, which could explain the difference in outcomes.

In conclusion, the alteration of the male song system by early E_2 or Flut may be explained either by E_2 regulating genes in abberant fashions and/or by the altering steroid hormone profile of the developing males. The latter would probably result from the apparent lack of testes development, which would lead to a crucial lack of androgens

impacting HVC development. These two explanations are not mutually exclusive, and in fact may be intertwined. Consideration of timing and duration of treatments as well as impacts on endocrine systems in development could unconfound the discrepancies in the literature.

.

Figure Captions

Figure 1 A) Mean HVC volume and B) number of HVC neurons as a function of treatment group. Males receiving E$_2$ or E$_2$ +Flut were demasculized. C) Mean HVC neuron size—males in E$_2$-treated groups had significantly larger neurons. D) RA volume--males receiving Flut were hypermasculinized * indicates a strong trend $p < 0.06$. Error bars = SEM.

Figure 2 A) Mean testes weight as a function of treatment--E$_2$ treated birds had markedly reduced testes weights. Error bars = SEM. B) Scatterplot of HVC volume and C) HVC neuron numbers as a function of testes size.

Acknowledgements:

Thanks to Dr. Juli Wade and Dr. Arthur P. Arnold who provided valuable feedback on an earlier version of this manuscript. Further thanks to Dr. Art Arnold for allowing us to use

his facilities. Thanks to Natalie Schottler who drew the figures and made the final edits and Kay Yang-Stayner and Janet Lee who helped with the execution of the study. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

Adkins-Regan, E., Mansukhani, V., Seiwert, C., & Thompson, R. (1994). Sexual differentiation of brain and behavior in the zebra finch: Critical periods for effects of early estrogen treatment. *Journal of Neurobiology, 25*(7), 865-877. https://doi.org/10.1002/neu.480250710

Benezra, S. E., Narayanan, R. T., Egger, R., Oberlaender, M., & Long, M. A. (2018). Morphological characterization of HVC projection neurons in the zebra finch (Taeniopygia guttata). *Journal of Comparative Neurology, 526*(10), 1673-1689. https://doi.org/10.1002/cne.24437

Bölting, S., & von Engelhardt, N. (2017). Effects of the social environment during adolescence on the development of social behaviour, hormones and morphology in male zebra finches (Taeniopygia guttata). *Frontiers in Zoology, 14*, 5. https://doi.org/10.1186/s12983-017-0190-4

Bottjer, S. W. (1987). Ontogenetic changes in the pattern of androgen accumulation in song-control nuclei of male zebra finches. *Journal of Neurobiology, 18*(2), 125-139. https://doi.org/10.1002/neu.480180202

Bottjer, S. W., & Hewer, S. J. (1992). Castration and antisteroid treatment impair vocal learning in male zebra finches. *Journal of Neurobiology, 23*(4), 337-53. https://doi.org/10.1002/neu.480230402

Casto, J. M., & Ball, G. F. (1996). Early administration of 17beta-estradiol partially masculinizes song control regions and alpha2-adrenergic receptor distribution in European starlings (Sturnus vulgaris). *Hormones and Behavior, 30*(4), 387-406. https://doi.org/10.1006/hbeh.1996.0044

Choe, H. N., Tewari, J., Zhu, K. W., Davenport, M., Matsunami, H., & Jarvis, E. D. (2021). Estrogen and sex-dependent loss of the vocal learning system in female zebra finches. *Hormones and Behavior, 129*, 104911. https://doi.org/10.1016/j.yhbeh.2020.104911

Diez, A., An, H. Y., Carfagnini, N., Bottini, C., & MacDougall-Shackleton, S. A. (2021). Neurogenesis and the development of neural sex differences in vocal control regions of songbirds. *Journal of Comparative Neurology, 529*(11), 2970-2986. https://doi.org/10.1002/cne.25138

Dittrich, F., Feng, Y., Metzdorf, R., & Gahr, M. (1999). Estrogen-inducible, sex specific expression of brain-derived neurotrophic factor mRNA in a forebrain song control nucleus of the juvenile zebra finch. *Proceedings of the National Academy of Sciences (PNAS), 96*(14), 8241-8246. https://doi.org/10.1073/pnas.96.14.8241

Fusani, L., Metzdorf, R., Hutchison, J. B., & Gahr, M. (2002). Aromatase inhibition affects testosterone-induced masculinization of song and the neural song system in female canaries. *Journal of Neurobiology, 54*(2), 370-379. https://doi.org/10.1002/neu.10141

Grisham, W., & Arnold, A. P. (1995). A direct comparison of the masculinizing effects of testosterone, androstenedione, estrogen, and progesterone on the development of the zebra

finch song system. *Journal of Neurobiology, 26*(2), 163-170.
https://doi.org/10.1002/neu.480260202

Grisham, W., Lee, J., McCormick, M. E., Yang-Stayner, K., & Arnold, A. P. (2002). Antiandrogen blocks estrogen-induced masculinization of the song system in female zebra finches. *Journal of Neurobiology, 51*(1), 1-8. https://doi.org/10.1002/neu.10028

Grisham, W., Park, S. H., Hsia, J. K., Kim, C., Leung, M. C., Kim, L., & Arnold, A. P. (2007). Effects of long-term flutamide treatment during development in zebra finches. *Neuroscience Letters, 418*(1), 92-96. https://doi.org/10.1016/j.neulet.2007.03.002

Grisham, W., Tam, A., Greco, C. M., Schlinger, B. A., & Arnold, A. P. (1997). A putative 5α-reductase inhibitor demasculinizes portions of the zebra finch song system. *Brain Research, 750*(1-2):122-128. https://doi.org/10.1016/s0006-8993(96)01336-4

Gurney, M. E. (1981). Hormonal control of cell form and number in the zebra finch song system. *The Journal of Neuroscience, 1*(6), 658-673. https://doi.org/10.1523/JNEUROSCI.01-06-00658.1981

Gurney, M. E. (1982). Behavioral correlates of sexual differentiation in the zebra finch song system. *Brain Research, 231*(1), 153-172. https://doi.org/10.1016/0006-8993(82)90015-4

Gurney, M. E., & Konishi, M. (1980). Hormone-Induced Sexual Differentiation of Brain and Behavior in Zebra Finches. *Science, 208*(4450), 1380-1382. https://doi.org/10.1126/science.208.4450.1380

Jacobs, E. C., Grisham, W., & Arnold, A. P. (1995). Lack of a synergistic effect between estradiol and dihydrotestosterone in the masculinization of the zebra finch song system. *Journal of Neurobiology, 27*(4), 513-519. https://doi.org/10.1002/neu.480270406

Lorenz, F. W. (1954). Effects of Estrogens on Domestic Fowl and Applications in the Poultry Industry. In R. S. Harris, G. F. Marrian, K. V. & Thimann (Eds.), *Vitamins and Hormones, 12* (pp. 235-275). Academic Press. https://doi.org/10.1016/S0083-6729(08)61014-6

Mathews, G. A., & Arnold, A. P. (1991). Tamoxifen fails to block estradiol accumulation, yet is weakly accumulated by the juvenile zebra finch anterior hypothalamus: An autoradiographic study. *Journal of Neurobiology, 22*(9), 970-975. https://doi.org/10.1002/neu.480220908

Nordeen, E. J., Grace, A., Burek, M. J., & Nordeen, K. W. (1992). Sex-dependent loss of projection neurons involved in avian song learning. *Journal of Neurobiology, 23*(6), 671-679. https://doi.org/10.1002/neu.480230606

Nordeen, E. J., & Nordeen, K. W. (1988). Sex and regional differences in the incorporation of neurons born during song learning in zebra finches. *The Journal of Neuroscience, 8*(8), 2869-2864. https://doi.org/10.1523/JNEUROSCI.08-08-02869.1988

Nottebohm, F., & Arnold, A. P. (1976). Sexual Dimorphism in Vocal Control Areas of the Songbird Brain. *Science, 194*(4261), 211-213. https://doi.org/10.1126/science.959852

Pröve, E. (1983). Hormonal Correlates of Behavioural Development in Male Zebra Finches. In J. Balthazart, E. Pröve, & R. Gilles (Eds) *Hormones and Behaviour in Higher Vertebrates* (pp. 368-374). Springer. https://doi.org/10.1007/978-3-642-69216-1_26

Rasika, S., Nottebohm, F., & Alvarez-Buylla, A. (1994). Testosterone increases the recruitment and/or survival of new high vocal center neurons in adult female canaries. *Proceedings of the*

National Academy of Sciences (PNAS), 91(17), 7854-7858.
https://doi.org/10.1073/pnas.91.17.7854

Schlinger, B. A., & Arnold, A. P. (1991). Androgen effects on the development of the zebra finch song system. *Brain Research, 561*(1), 99-105. https://doi.org/10.1016/0006-8993(91)90754-J

Simpson, H. B., & Vicario, D. S. (1991). Early estrogen treatment of female zebra finches masculinizes the brain pathway for learned vocalizations. *Journal of Neurobiology, 22*(7), 777-793. https://doi.org/10.1002/neu.480220711

Soma, K. K., Sullivan, K. A., Tramontin, A. D., Saldanha, C. J., Schlinger, B. A., & Wingfield, J. C. (2000). Acute and chronic effects of an aromatase inhibitor on territorial aggression in breeding and nonbreeding male song sparrows. *Journal of Comparative Physiology A, 186*, 759-769. https://doi.org/10.1007/s003590000129

Tang, Y. P., & Wade, J. (2009). Effects of estradiol on incorporation of new cells in the developing zebra finch song system: Potential relationship to expression of ribosomal proteins L17 and L37. *Developmental Neurobiology, 69*(7):462-475. https://doi.org/10.1002/dneu.20721

Tang, Y. P., & Wade, J. (2010). Sex- and age-related differences in ribosomal proteins L17 and L37, as well as androgen receptor protein, in the song control system of zebra finches. *Neuroscience, 171*(4), 1131-1140. https://doi.org/10.1016/j.neuroscience.2010.10.014

Thompson, J. B., Dzubur, E., Wade, J., & Tomaszycki, M. (2011). The effects of estradiol on 17β-hydroxysteroid dehydrogenase type IV and androgen receptor expression in the developing zebra finch song system. *Brain Research, 1401*, 66-73. https://doi.org/10.1016/j.brainres.2011.05.031

Tramotin, A. D., Smith, G. T., Breuner, C. W., & Brenowitz, E. A. (1998). Seasonal plasticity and sexual dimorphism in the avian song control system: Stereological measurement of neuron density and number. *Journal of Comparative Neurology, 396*(2), 186-192. https://doi.org/10.1002/(SICI)1096-9861(19980629)396:2<186::AID-CNE4>3.0.CO;2-X

Wade, J., Gong, A., & Arnold, A. P. (1998). Effects of embryonic estrogen on differentiation of the gonads and secondary sexual characteristics of male zebra finches. *Journal of Experimental Zoology, 278*(6), 405-411. https://doi.org/10.1002/(SICI)1097-010X(19970815)278:6<405::AID-JEZ8>3.0.CO;2-S

Walton, C., Pariser, E., & Nottebohm, F. (2012). The Zebra Finch Paradox: Song Is Little Changed, But Number of Neurons Doubles. *The Journal of Neuroscience, 32*(3), 761-774. https://doi.org/10.1523/JNEUROSCI.3434-11.2012