Alteration of nociceptive Schwann cells in a mouse model of peripheral neuropathy in prediabetic condition

Guillaume Rastoldo[1], Ali K. Jaafar[1], Aurélie-Paulo-Ramos[1], Katy Thouvenot[1], Cynthia Planesse[1], Matthieu Bringart[1], Marie-Paule Gonthier, Gilles Lambert and Steeve Bourane[1]*

1- Université de La Réunion, INSERM, UMR 1188 Diabète athérothrombose Réunion Océan Indien (DéTROI), Saint-Pierre, La Réunion, France.

Abstract

Diabetic peripheral neuropathy (DPN) characterized by progressive and symmetrical sensory abnormalities is one of the earliest and main complications of diabetes. DPN is characterized by heterogeneous sensory symptoms such as chronic pain, tingling, burning or loss of sensation. Nociceptive Schwann cells (nSCs), a recently identified subtypes of dermal Schwann cells support terminal nerve fibers in mouse skin and contribute to mechanical sensation and neuropathic pain. While terminal nerve fibers density is basically evaluated in DPN models, there is currently few data about nSCs number and integrity during early stage of DPN. In the present study, we used a mouse model of prediabetes by using high-fat diet (HFD) fed mice to determine if there are quantitative differences in terminal nerve fiber density, nSCs number and cellular extensions between control and prediabetic neuropathic mice. In this study, we identified L1CAM as a reliable marker of nSCs currently characterized by the expression of S100β and Sox10. Interestingly, we observed a decrease in intraepidermal nerve fiber density (IENFD) associated to a significant reduction of nSCs in the glabrous foot skin of neuropathic mice. Overall, this study identifies L1CAM as a new marker of nSCs and indicates that these cells are impaired during prediabetic peripheral neuropathy.

Keywords: DPN; Nociceptive Schwann Cells; Type 2 diabetes; Mechanical hypoalgesia; Thermal hyperalgesia

Introduction

Diabetes is one of the most prevalent chronic diseases in the world. The epidemic threshold has been reached with over 537 million people affected worldwide. According to estimates by the International Diabetes Federation, this number is expected to reach 783 million patients by 2045. One of the first and main complications of diabetes is diabetic peripheral neuropathy (DPN), affecting more than 60% of diabetic patients (Feldman et al., 2017). Prediabetes represents the early stages of glucose dysregulation and precedes the development of type 2 diabetes. Prediabetes and diabetes are the most common causes of peripheral neuropathy in Europe and the United States (Feldman et al., 2017). DPN involves damage to the peripheral nervous system, i.e., the sensory and motor nerves as well as the autonomic nervous system that controls our organs. The most common symptoms of DPN are an impairment of sensitivity to pain and touch (Themistocleous et al., 2016). The early stages of the pathology is characterized by abnormal sensory symptoms such as pain, tingling, burning sensations, electrical discharges, pain evoked by non-painful mechanical stimuli (allodynia), increased sensitivity to noxious stimuli (hyperalgesia) (Themistocleous et al., 2016) significantly altering the patients' quality of life. These symptoms are difficult to treat with conventional analgesics.

Other sensory deficits such as numbness are also observed with progressive loss of sensation in the lower limbs often leading to tissue damage with the development of ulcers, increasing the risk of amputation (Callaghan et al., 2012a). Diabetes is the leading cause of lower extremity amputations, with a 15-fold increase in risk compared with non-diabetic populations (Margolis et al., 2011). Reduced sensation in the feet also leads to progressive loss of balance resulting in an increased risk of falls, fractures and death (Green et al., 2010; Malik et al., 2005).

Despite decades of research, there is currently no effective treatment for DPN due to an incomplete understanding of its mechanisms and key players. Much attention in DPN studies has been dedicated to the damage of small, non-myelinated nerve fibers (C-fibers) responsible for sensitivity to heat and pain. The degeneration and regeneration of these C-fibers are typically associated with diabetic patient phenotypes. Skin biopsy analysis in clinic serves as a late-stage confirmation of DPN presence (Lauria et al., 2022), lacking a diagnostic tool for predicting C-fiber degeneration related to pain.

While C-fibers was traditionally believed to terminate freely in the epidermis, recent work by Abdo et al. challenges this perception (Abdo et al., 2019). Their findings reveal a close connection between these C-fibers and specialized nociceptive Schwann cells (nSC) capable of independently generating pain signals. These cells are currently identified by the expression of S100β and Sox10 (Hu et al., 2022; Ojeda-Alonso et al., 2023; Rinwa et al., 2021). Conversely, eliminating these nSC alone seems sufficient to induce neuropathic pain in mice (Rinwa et al., 2021). Considering these recent discoveries, it is conceivable that neuropathies involving C-fiber degeneration, as seen in DPN, may result from early damage to nSCs. A recent report

supports this hypothesis and demonstrated a decreased in nSC density in Streptozotocin-
induced type 1 diabetes mice (Hu et al., 2022). However, no study has evaluated the integrity
of nSC in a type 2 diabetes model such as HFD or in prediabetes conditions.

To explore this hypothesis, our research delves into the impact of prediabetes on the integrity
of nSC during pathology development, an aspect that remains unexplored in the realm of DPN
investigations.

Methods
75

Research Design
77

78 All reported experiments were performed at the GIP-CYROI platform's animal facility,
79 conducted in accordance with the French and European Community Guidelines for the Use of
80 Animals in Research (86/609/EEC and 2010/63/EU), and were approved by the local Ethics
81 Committee (n°114) for animal experimentation and the French authorities (APAFIS#31763-
82 2018020912188806, approved on May 21, 2021).

Animals
84

85 Twenty C57BL/6J mice (male, 6 weeks old) were purchased from Janvier (Le Genest Saint Isle,
86 France) and housed with 5 mice per cage in a temperature-controlled environment with a 12–
87 12h light/dark photocycle. After 2 weeks of adaptation, mice were divided in two groups
88 (n=10/group) and placed on either a control diet (10% kcal %fat; SF13-081) or a high-fat diet
89 (45% kcal %fat; SF04-001) from Specialty Feeds during 16 weeks.

Transgenic mouse line and Tamoxifen injection
91

Mice expressing a tamoxifen-inducible Cre recombinase under the control of the PLP
promoter (PLP-CreER^{T2}; (Leone et al., 2003)) were crossed to Ai14 mice, which have a LoxP-
flanked stop cassette excisable by Cre. At 6 weeks old, the PLP-CreER^{T2}; Ai14 progeny were
injected intraperitoneally with tamoxifen (100 µg/g of mice) for 5 consecutive days. Tamoxifen
was prepared in a mixture of 9/10 corn oil and 1/10 ethanol, and stored at 4 °C in a light-proof
container. After 2 weeks, the mice were euthanized and the tissue collected for
immunostaining.

Genotyping
100

101 Genotyping of progeny from PLP-CreER[T2] and Ai14 mice were done with PCR using CreDN2
102 (GAT CTC CGG TAT TGA AAC TCC AGC), CreUP2 (GCT AAA CAT GCT TCA TCG TCG G), Ai14mut-
103 Fwd (CTG TTC CTG TAC GGC ATG G) and Ai14mut-Rev (GGC ATT AAA GCA GCG TAT CC).

Oral glucose tolerance test (OGTT)

106 An OGTT was performed after 16 weeks of diet in all mice. After 6 hours of fasting, one drop
107 of tail blood was collected and glycemia was analyzed using a standard glucometer (One Touch
108 Profile, Lifescan Inc., Milpitas, CA, USA, #6 strips). Oral gavage with glucose 30% (2g/kg body
109 weight) into conscious mice was performed and glycemic levels measured after 15, 30, 45, 60,
110 90 and 120 min post-administration.

Nerve conduction velocity

Mice were anesthetized by using isoflurane (IsoFlo®, Centravet, France) and placed on a
heating pad. Once they were deeply anesthetized, motor nerve conduction velocities (MNCV)
and sensory nerve conduction velocities (SNCV) were determined as previously described
(Obrosova et al., 2004; Oh et al., 2010; Schulz et al., 2014). Using the Keypoint 4, Dantec-
Medtronic electroneuromyography device, we performed both proximal and distal nerve
stimulations of the sciatic nerve, registering the transmitted potentials with sensing
electrodes at the gastrocnemius muscle. We used stainless steel subdermal needle electrodes
to deliver single square-wave supramaximal stimulation with 0.2 millisecond impulses. MNCV
was calculated by subtracting the distal from the proximal latency (measured in milliseconds)
from the stimulus artifact of the take-off of the evoked potential, and the difference was
divided into the distance between both stimulating electrodes (measured in millimeters using
a Vernier caliper). For SNCV, recording electrodes were placed on the dorsum of the foot and
stimulating electrodes on the ankle. For SNCV, the latency of onset (milliseconds) of the
sensory nerve action potential after supramaximal antidromic stimulation of the sural nerve
at the ankle was divided by the distance between the recording and stimulation electrodes
(measured in millimeters using a Vernier caliper).

Tissue Harvest

After 16 weeks of diet, mice received a subcutaneous injection of Buprenorphine (0.05 mg/kg,
Buprecare®, Centravet, France) and were anesthetized by using isoflurane (3%). The mice were
then euthanized by intracardiac puncture and blood samples were collected into Eppendorf
tubes and centrifuged at 2000 rpm for 10 min at 4°C. Serum was collected and stored at −80°C
until analysis. Intracardiac perfusion was performed with 20 mL of phosphate buffered saline
(PBS) and 20 mL of 4% paraformaldehyde (PFA) dissolved in PBS. The hind-paw plantar
surfaces of each mouse were collected and post-fixed overnight in 4% PFA solution. Biopsies

of foot skin were then rinsed and cryoprotected into 30% sucrose dissolved in PBS at 4°C. The
samples were then embedded in optimal cutting temperature (OCT, Tissue-Tek®, USA)
compound and stored at −80°C until sectioning.

Immunohistochemistry

Tissus sections (30 μm) were collected from the hind-paw plantar surface and washed
(3x10min) with PBS. Blocking and permeabilization were done by incubation (1h) in 5% donkey
anti-serum and 0.3% Triton X-100. For intraepidermal nerve fibers density, sections were
incubated with the primary antibody anti-PGP9.5 (1:500, #14730-1-AP, ProteinTech) for 2
days; first overnight at 4°C and the next day sections were incubated overnight at room
temperature. The same protocol was applied for L1 cell adhesion molecule (L1CAM, 1:500,
#20659-1-AP, ProteinTech) antibody but the sections were incubated only overnight at 4°C.
Sections were then rinsed in PBT (PBS + 0.1% triton) and incubated for 1 h at room
temperature with the secondary antibody Alexa Fluor 594 donkey anti-rabbit (1:500, Abcam;
ab150064), or Alexa Fluor 488 donkey anti-Rabbit (1:500, Invitrogen; A-21206).

Identification and quantification of nociceptive Schwann cells

Nociceptive Schwann cells were identified according to Hu et al. criteria (Hu et al., 2022).
Briefly, positive staining of S100β, Sox10 and DAPI identified nSCs. However, our investigation
revealed that the traditional markers for nociceptive Schwann cells (S100β and Sox10)
exhibited colocalization with L1CAM staining (**Supplementary Figure 1A**). Furthermore,
L1CAM staining colocalized with Td-tomato expression in a tamoxifen-induced PLP-*Cre*ER^{T2};
R26TdTomato double transgenic mouse (**Supplementary Figure 1B**). Cre-mediated
recombination driven by the proteolipid protein 1 (Plp1) promoter is employed specifically to
target Schwann cells (Doerflinger et al., 2003; Gomez-Sanchez et al., 2017; Grove et al., 2017).
These findings confirm that L1CAM is a reliable marker of nociceptive Schwann cells and can
independently be used without S100β and Sox10 to identify these cells.

To quantify nSCs, their localization within the foot skin is important. Commonly, nSCs connect
to PGP9.5+ nerve endings and are located within 25 μm depth in the subepidermal zone at
the junction between dermis and epidermis. For quantification, we exclude L1CAM positive
staining in glands or large peripheral fiber bundles found deeper in the dermis. The number
of nSCs was reported per mm length of foot skin.

To quantify nSCs cellular extensions, we measured cell fluorescence using ImageJ. We selected
the region of interest (ROI) at the subepidermal zone in each image of the foot skin and
measured the mean fluorescence value in that area using 'Mean gray value'.

Confocal imaging

Intraepidermal nerve fibers density and nociceptive Schwann cells number were evaluated under a laser scanning confocal microscope Eclipse (Nikon) with a 40x objective. Following immunostaining, 30 µm thick Z-stack images were taken and analysis was performed on orthogonal maximum intensity projection of the original confocal Z-stack images.

Metabolic markers

Plasma hemoglobin A_1C levels were determined using DCA® Vantage Analyzer (Siemens Healthineers, France) and hemoglobin A_1C reagent kit (#23-312018). Serum triglycerides (TG) and total cholesterol levels were determined using colorimetric assays (Diasys®, #157109910021 and #113009910021, Germany). Plasma insulin and leptin concentrations were measured using a mouse insulin and leptin ELISA kit (Mercodia®, #10-1247-01, Sweden and RayBiotech®, Peachtree Corners, GA, USA). All metabolic markers were measured after 16 weeks of diet.

Behavioral tests

Von Frey test

Mice were habituated to the testing apparatus for 1h/day during a week before behavioral testing. After this acclimation, animals were placed on an elevated wire grid and the plantar surface of the hindpaw was stimulated with calibrated von Frey monofilaments (0.008-6 g). The paw withdrawal threshold for the von Frey assay was determined by SUDO up-down method (Bonin et al., 2014). Testing was performed every 4 weeks after the start of the diet and up to 16 weeks.

Hargreaves test

To measure radiant heat pain by Hargreaves test (Hargreaves et al., 1988), animals were put in plastic boxes and the plantar paw surface was exposed to a beam of radiant heat according to the Hargreaves method. Paw withdrawal latency was then recorded and beam intensity was adjusted to result in a latency around 10 seconds in control animals. High speed of skin heating (6.5°C/sec) preferentially activates Aδ-fibers (Yeomans and Proudfit, 1996). Based on the study of Deuis and Vetter (Deuis and Vetter, 2016), we can assume that we used a medium to high speed of temperature heating (5°C/sec during 10 sec to obtain 50°C) which could indicate a predominant activation of Aδ-fibers on this behavioral test. Testing began after exploration and grooming behaviors ended. The heat stimulation was repeated 5 times at an interval of 10 min for each animal and the mean was calculated. A cutoff time of 30 seconds was set to prevent tissue damage. Testing was performed every 4 weeks after the start of the diet and up to 16 weeks.

Brush Test

To measure dynamic allodynia, mice were habituated to the testing apparatus (the same that
for von-Frey test) for 1h/day during a week before behavioral testing. After this acclimation,
a soft paintbrush (Lefranc & Bourgeois® - Brush no. 4 #175160) was used to gently stroke the
plantar surface of the hindpaw in the direction from heel to toe. The test was repeated five
times, with intervals of 5 min. For each test, no evoked movement was scored as 0, and
walking away or occasionally brief paw lifting (~1 sec or less) was scored as 1. For each mouse,
the cumulative scores of the five tests were used to form a percentage response (100%
response equivalent to 5 evoked movements during the 5 stimulations of the paw). Testing
was performed every 4 weeks after the start of the diet and up to 16 weeks.

Pinprick Test

To measure acute mechanical pain, mice were habituated to the testing apparatus (the same
that for von-Frey test) for 1h/day during a week before behavioral testing. After this
acclimation, for pinprick test, we touched the plantar surface of the hindpaw with a pin
(Bourane et al., 2015) and measured numbers of withdrawal response as for the brushing
tests. For each mouse, the cumulative scores of the five tests were used to form a percentage
response (100% response equivalent to 5 evoked movements during the 5 stimulations of the
paw). Testing was performed every 4 weeks after the start of the diet and up to 16 weeks.

Statistical analysis

We first analyzed the normality of all the data and showed that the results of serum
triglycerides and total cholesterol levels, insulinemia, motor NCV, sensory NCV, IENFD and
L1CAM-positive cells were normally distributed. Thus, unpaired Student's *t*-tests (two tailed)
were used for statistical comparisons. For behavioral data, a two-way ANOVA for repeated
measures followed by Bonferroni's post-hoc test was used to determine statistically significant
differences. A *P* value of less 0.05 was considered significant.

Results

1. Body weight gain, serum triglycerides and total cholesterol levels

In order to determine the effect of high-fat diet (HFD) on the development of peripheral
neuropathy, we fed mice with a control diet (10% kcal %fat) or a HFD (45% kcal %fat) for 16
weeks (n=10/group, **Figure 1A**). As expected, HFD-fed mice gained significant weight
compared to control diet-fed mice. The difference became significant from week 11 onward
(**Figure 1B**). Mice on the control diet gradually gained 10 g in body weight, whereas HFD-fed
mice gained 18 g over the course of the study. In line with weight gain, blood samples analysis

revealed that HFD-fed mice exhibited higher circulating levels of leptin compared to control
diet-fed mice (**Figure 1C**).Triglycerides levels were not significantly altered (**Figure 1D**).
However, HFD-fed mice showed significantly higher total plasma cholesterol level than mice
fed with the control diet (P < 0.01; **Figure 1E**).

Figure 1: Study design and HFD-induced dyslipidemia. (A) Study Design used to assess the effect of a
HFD (45% kcal %fat) vs. a control diet (10% kcal %fat) in two groups of mice (n=10/group). (**B**) Body
weight of control diet-fed mice and HFD-fed mice during the 16 weeks of diet. (**C**) Leptinemia of control
and HFD mice at 16 weeks of diet. (**D**) Plasma levels triglycerides of control and HFD mice at 16 weeks

of diet. (**E**) Total cholesterolemia of control and HFD mice at 16 weeks of diet. *$P < 0.05$, **$P < 0.01$,
***$P < 0.001$: control diet vs. HFD. Data are presented as means ± SEM.

2. Glucose homeostasis / Glucose regulation

Fasting glucose levels were assessed and exhibited a significant increase in HFD-fed mice
compared to mice fed with the control diet ($P < 0.001$; **Figure 2A**). However, HbA1c levels
were not significantly altered in mice fed with HFD compared to control diet-fed mice (**Figure
2B**). Plasma insulin level in HFD-fed mice was significantly increased compared to control mice
($P < 0.001$; **Figure 2C**) confirming impaired glucose tolerance and insulin resistance. We
performed an oral glucose tolerance test (OGTT) after 16 weeks of diet on both groups. We
observed that glucose tolerance was impaired in HFD-fed mice compared to control mice
(**Figure 2D**) with the AUC of blood glucose concentration significantly higher in HFD-fed mice
compared to control diet-fed mice ($P < 0.01$; **Figure 2E**).

Figure 2: HFD-induced impaired glucose tolerance. (A) Fasting blood glucose of control diet-fed mice
and HFD-fed mice at 16 weeks of diet. **(B)** Percentage of glycated hemoglobin (HbA1c) of control diet-
fed mice and HFD-fed mice at 16 weeks of diet. **(C)** Plasma insulin of control diet-fed mice and HFD-
fed mice at 16 weeks of diet. **(D)** Oral glucose tolerance test (OGTT) curves performed at 16 weeks of
diet in control and HFD-fed mice. **(E)** Area under the curve (AUC) during OGTT for control diet-fed mice
and HFD-fed mice. *P < 0.05, **P < 0.01, ***P < 0.001: control diet vs. HFD. Data are presented as
means ± SEM (n=10/group).

3. Nerve conduction studies

At week 16, nerve conduction velocity (NCV) was measured in the sciatic and sural nerves of
each animal. HFD-fed mice did not show any significant change in motor (**Figure 3A**) or sensory
(**Figure 3B**) NCV compared to control mice.

Figure 3: Nerve conduction studies. (A) Motor nerve conduction velocity (MNCV) performed at 16
weeks of diet in control and HFD-fed mice. **(B)** Sensory nerve conduction velocity (SNCV) performed at
16 weeks of diet in control and HFD-fed mice**.** Data are presented as means ± SEM (n=10/group).

4. Thermal and mechanical pain sensitivity

To evaluate peripheral neuropathy in mice we performed various behavioral sensory tests
every 4 weeks for 16 weeks of diet. HFD-fed mice exhibited an increase in paw withdrawal
threshold as early as 8 weeks (4 weeks HFD vs. 8 weeks HFD, P < 0.01, **Figure 4A**) and were
significantly different from their control diet-fed counterparts after 12 weeks of diet (12 weeks
control diet vs. 12 weeks HFD, P < 0.01, **Figure 4A**), indicating a decrease in mechanical
sensitivity. In parallel, we observed an increase in thermal sensitivity of HFD-fed mice
compared to control mice from week 8 (**Figure 4B**). The latency of hindpaw withdrawal in
response to noxious thermal stimulus was significantly decreased in HFD-fed mice compared
to control diet-fed mice at 12 and 16 weeks (P < 0.001). No significant change in haptic

mechanical sensitivity (**Figure 4C**) evaluated with the brush test or acute mechanical pain sensitivity (**Figure 4D**) assessed by the pinprick test was noticed over 16 weeks between both groups.

Figure 4: Mechanical and thermal sensitivity changes induced by HFD. (A) Quantification of static mechanical sensitivity was assessed every 4 weeks in control and HFD-fed mice using von Frey filaments. **(B)** Quantification of thermal sensitivity by Hargreaves test every 4 weeks in control and HFD-fed mice. **(C)** Quantification of haptic mechanical sensitivity by brush test every 4 weeks in control and HFD-fed mice. **(D)** Quantification of acute mechanical pain by pinprick test every 4 weeks in control- and HFD-fed mice. **$P < 0.01$, ***$P < 0.001$: control diet vs. HFD. #$P < 0.05$, ##$P < 0.01$: HFD 4weeks vs. HFD 8, 12 or 16 weeks. Data are presented as means ±SEM (n=10/group).

5. Epidermal innervation and Nociceptive Schwann cells

In order to fully characterized neuropathy in our model, we analyzed the intraepidermal nerve fiber density (IENFD) in the glabrous skin of control and HFD-fed mice. Our results show a significant reduction of foot skin innervation in HFD-fed compared with control mice after 16 weeks of diet ($P < 0.001$; **Figure 5A**, **5B**). Additionally, we investigated the integrity of

nociceptive Schwann cells that are present in the skin at the junction between the epidermis and the dermis by using immunohistochemistry against the L1 Cell Adhesion Molecule (L1CAM) that we found highly expressed in these cell types (**Supplementary Figure 1A, B**). To validate the specificity of L1CAM expression in nociceptive Schwann cells, we first verified its colocalization with Sox10 and S100β, two markers previously described to be expressed in these cells (Hu et al., 2022; Ojeda-Alonso et al., 2023; Rinwa et al., 2021) (**Supplementary Figure 1A**). To go even further, we confirmed the expression of L1CAM in Td-tomato positive nSCs in PLP-CreER^{T2};Ai14 double transgenic mouse foot skin (**Supplementary Figure 1B**), (Gomez-Sanchez et al., 2017; Leone et al., 2003). Altogether, these results indicate that L1CAM is a reliable marker of nSCs.

Using L1CAM to identified nSCs, we observed that, both mean fluorescence intensity (**Figure 5C,D**) and mean number of L1CAM-positive cells and cellular extensions (**Figure 5C,E**) were significantly decreased in HFD-fed mice compared with control mice ($P < 0.001$ and $P < 0.01$ respectively). Altogether, these results indicate a decrease in both sensory nerve endings and nociceptive Schwann cells in the foot skin of HFD mice.

Figure 5: HFD-induced small nerve fiber and nociceptive Schwann cells degeneration in the foot skin.
(A) PGP9.5 immunofluorescence with DAPI staining in foot skin section of control (n=5) and HFD-fed mice (n=5). **(B)** Quantitation of IENFD is presented as the number of fibers/mm of epidermis. **(C)** L1CAM immunofluorescence with DAPI staining in foot skin section of control (n=5) and HFD-fed mice (n=5).

Arrowhead indicate nerve fibers in the epidermis of the foot skin. Arrows indicate nociceptive Schwann
cells and their cellular extensions at the border of the epidermis and the dermis. **(D)** Mean fluorescence
intensity quantification of L1CAM immunostaining at the localization of nociceptive Schwann cells (at
the border of epidermis and dermis). **(E)** Quantification of L1CAM-positive cells and their cellular
extensions presented in number/mm of epidermis. $**P < 0.01$, $***P < 0.001$: control diet vs. HFD. Data
are presented as means ±SEM. Scale bar: 50 μm.

Discussion

In the present study, we used HFD (45% fat) fed mice during 16 weeks to generate a
mouse model of peripheral neuropathy in prediabetic condition and compared dyslipidemia,
glucose tolerance, and sensory deficits of the mice with the corresponding control diet (10%
fat). We demonstrated that HFD-fed mice develop a prediabetic phenotype with increased
weight gain, elevated leptinemia, hypercholesterolemia, impaired glucose tolerance, and
elevated fasting blood glucose and insulin resistance compared with control mice. Our
combined findings show that the HFD model recapitulate metabolic dysfunction observed in
prediabetes. The analysis of IENFD of the foot skin revealed an important reduction of terminal
fibers and nSCs in HFD-fed mice compared with mice fed with a control-diet. We also observed
behavioral sensory deficits in HFD-fed mice, with the development of mechanical
hyposensitivity and thermal hypersensitivity. Overall, our results indicate that HFD-fed mice
exhibit peripheral neuropathy in prediabetic conditions.

The literature on HFD models is extensive and each diet seems to have its specificities, which
leads to the development of different phenotypes in animals. Indeed, whether it is between
high fat diets of 45 or 60% of fat or even on the composition of the diet between those using
lard-based vs. plant-based or fish oil fats, the results obtained are not similar (For review see
(Eid and Feldman, 2021)). Rodent studies using a 60% fat HFD are potentially not relevant as
those using a 45% fat diet in term of human food consumption (Speakman, 2019). Therefore,
the choice and a proper characterization of the diet is important to know where the model
stands in terms of clinical significance. Mice fed on a high-fat diet progressively develop
metabolic disorders, including weight gain, insulin resistance, dyslipidemia and impaired
glucose tolerance. The HFD used in the present study recapitulates all of these deficits after
16 weeks of diet, with some discrepancies from the literature depending on the time spent on
diet. Indeed, when mice are fed with HFD during 34 weeks they developed a more pronounced
metabolic phenotype as demonstrated by increased plasma triglycerides (TG) and elevated
glycated hemoglobin (Vincent et al., 2009; Xu et al., 2019) which we did not find in the present
study. However, the lack of a difference in TG levels was also found in a similar study by Sajic
and colleagues after the same time spent on HFD (16 weeks) (Sajic et al., 2021). For HbA1c
levels, fasting blood glucose was only evaluated at the end of the experiments and, despite
we demonstrated hyperglycemia in HFD-fed mice, we do not know since when chronic

hyperglycemia started. Importantly, glycation of hemoglobin starts 6 weeks following
persistent hyperglycemia on rats (Nagisa et al., 2003) and then last until the lifespan of the
red blood cell, which is about 3 months in human, 60 days in rats and 40 days in mice (Van
Putten, 1958). It is then possible that in our study the chronic hyperglycemia of HFD mice did
not last long enough to be able to have an impact on the level of glycated hemoglobin
evaluated at 16 weeks. Considering the other metabolic parameters we measured in our
study, we can assume that our HFD-fed mice were pre-diabetic.

To characterize the progression of peripheral neuropathy, we followed the guidelines of the
diabetic neuropathy study group of the European Association for the Study of Diabetes
(Neurodiab). Namely, we assessed three key features present in human pathology
(nocifensive behavior, nerve conduction velocity, and IENFD) considering that the presence of
alterations in two of these three assessments are necessary to establish neuropathy in rodents
(Biessels et al., 2014). In this study, we demonstrated that HFD-fed mice exhibited mechanical
hypoalgesia and thermal hyperalgesia coupled to a severe loss of intraepidermal nerve fibers
but no impairment of motor or sensory nerve conduction velocity. According to the criteria of
Neurodiab, we can assert that our HFD-fed mice develop peripheral neuropathy in a context
of prediabetes.

In order to identify possible damage to small sensory fibers, we performed a quantification of
intraepidermal nerve fibers density considered as the gold standard for the evaluation of small
fiber neuropathies (Yu, 2021). Damage to small sensory fibers is one of the earliest
manifestations of DPN that can be observed in prediabetic conditions (Sumner et al., 2003).
We found a drastic reduction of 70% of epidermal innervation in the hind paw of the HFD-fed
mice in comparison to control-fed mice. This reduction of nerve fibers density is in accordance
with the literature using an equivalent HFD (Coppey et al., 2011; Sajic et al., 2021; Vincent et
al., 2009). Studies have shown that small nerve fibers penetrating the epidermis are mostly
nociceptive (Lauria et al., 2022) and these fibers undergo degeneration and regeneration in
patients with impaired glucose tolerance and at the early stages of type 2 diabetes (Green et
al., 2010; Ziegler et al., 2014). Once neuropathy progresses, degenerated fibers do not
regenerate. However, not all authors agree that the severity of IENFD loss is correlated with
sensory deficits and disease severity (Arimura et al., 2013; Nolano et al., 2020; Shun et al.,
2004; Smith et al., 2001). Until recently, it was thought that because unmyelinated C-fibers
lack the protection and nutrient supply that myelinated Schwann cells provide to A-fibers,
these unmyelinated C-fibers were more likely to be affected at the early stages of diabetic
neuropathy. However, recently a specialized cutaneous type of Schwann cells, termed
nociceptive Schwann cells , in close relationship with nociceptive C-fibers in the dermis has
been discovered (Abdo et al., 2019). These cells contribute to terminal axon support and play
a critical role in skin mechanosensitivity and neuropathic pain (De Logu et al., 2017; Ojeda-
Alonso et al., 2023; Raut et al., 2024; Rinwa et al., 2021). Therefore, nociceptive endings in the
epidermis are not "free" but form a mesh-like organ with a mutual dependence between C-
fibers and the nSCs. The absence of nSCs induces a retraction of epidermal nerve fibers and

development of mechanical, cold, and heat hyperalgesia (Rinwa et al., 2021). In the present study, we identified L1CAM as a reliable marker for nSCs, as an alternative to the conventional S100β / Sox10 staining methods highlighted by (Hu et al., 2022; Ojeda-Alonso et al., 2023; Rinwa et al., 2021). Interestingly, previous research used another cell adhesion molecule, NCAM to identify terminal Schwann cells in the skin of diabetic patients (Reinisch et al., 2008). As outlined by Hu et al., the precise localization of Schwann cells serves as a crucial criterion in distinguishing nSCs (Hu et al., 2022). These distinctive cells are arranged within a 25 μm depth in the subepidermal region, excluding glands and large fiber bundles. Based on these criteria and the expression of L1CAM, we demonstrated a significant decrease in the number of nSCs and their cellular extensions in HFD-fed mice compared to control mice. Interestingly, a similar results was obtained in a mouse model of type 1 diabetes which develop mechanical and thermal hyperalgesia (Hu et al., 2022).

Prediabetes represents the earliest stage of glucose dysregulation and hyperglycemia a primary factor that can induces apoptosis in Schwann cells (Liu et al., 2020). High glucose levels found in diabetic patients can lead to oxidative stress, inflammatory reactions, endoplasmic reticulum stress, autophagy, nitrification, and disruption of various signaling pathways such as PI3K/AKT, ERK, PERK/Nrf2, and Wnt/β-catenin, all of which are involved in Schwann cells apoptosis and the pathogenesis of DPN (Li et al., 2017; Liu et al., 2020; Wu et al., 2023; Zhang et al., 2021). Inhibiting apoptosis of Schwann cells under high-glucose conditions is considered a promising approach to treat DPN (Wang et al., 2023). While glycemic control has little impact of the development and progression of neuropathy in type 2 diabetic patients (Callaghan et al., 2012b), it is important to recognize that DPN is influenced by various metabolic parameters. In addition to elevated glucose levels, dyslipidemia is another factor commonly observed in diabetic patients and despite its prevalence, the impact of dyslipidemia on Schwann cells integrity has received relatively little attention into the literature (Jende et al., 2019; Rumora et al., 2022; Stino et al., 2020). Overall, the above-mentioned literature focused on Schwann cells in general, without any particular attention to the subpopulation of nociceptive Schwann cells despite its major role in skin mechanosensitivity and neuropathic pain. Therefore, the elucidation of mechanisms implicated in the alteration of nSCs in the skin is crucial for understanding and developing potential treatments for DPN.

Conclusion

In this study, we used HFD to induce a mouse model of peripheral neuropathy in prediabetic condition. We observed key metabolic dysfunctions associated with prediabetes, including increased weight gain, hypercholesterolemia, and impaired glucose tolerance. Behavioral analysis uncovered early-onset of peripheral neuropathy with mechanical hypoalgesia and thermal hyperalgesia. We also identified L1CAM as a new marker of nociceptive Schwann cells previously characterized by the expression of S100β and Sox10. Interestingly, we observed a significant decrease in number of nociceptive Schwann cells and their extensions in the foot

456   skin of neuropathic mice, highlighting the vulnerability of these cells in prediabetic conditions.

457   Overall, this research enhances our understanding of the early events in peripheral

458   neuropathy during prediabetic states, paving the way for future investigations into the

459   molecular mechanisms underlying nociceptive Schwann cells alterations and potential

460   strategies to thwart this complication of diabetes.

Bibliography

Abdo, H., Calvo-Enrique, L., Lopez, J.M., Song, J., Zhang, M.-D., Usoskin, D., El Manira, A., Adameyko, I., Hjerling-Leffler, J., Ernfors, P., 2019. Specialized cutaneous Schwann cells initiate pain sensation. Science 365, 695–699. https://doi.org/10.1126/science.aax6452

Arimura, A., Deguchi, T., Sugimoto, K., Uto, T., Nakamura, T., Arimura, Y., Arimura, K., Yagihashi, S., Nishio, Y., Takashima, H., 2013. Intraepidermal nerve fiber density and nerve conduction study parameters correlate with clinical staging of diabetic polyneuropathy. Diabetes Res. Clin. Pract. 99, 24–29. https://doi.org/10.1016/j.diabres.2012.09.026

Biessels, G.J., Bril, V., Calcutt, N.A., Cameron, N.E., Cotter, M.A., Dobrowsky, R., Feldman, E.L., Fernyhough, P., Jakobsen, J., Malik, R.A., Mizisin, A.P., Oates, P.J., Obrosova, I.G., Pop-Busui, R., Russell, J.W., Sima, A.A., Stevens, M.J., Schmidt, R.E., Tesfaye, S., Veves, A., Vinik, A.I., Wright, D.E., Yagihashi, S., Yorek, M.A., Ziegler, D., Zochodne, D.W., 2014. Phenotyping animal models of diabetic neuropathy: a consensus statement of the diabetic neuropathy study group of the EASD (Neurodiab). J. Peripher. Nerv. Syst. JPNS 19, 77–87. https://doi.org/10.1111/jns5.12072

Bonin, R.P., Bories, C., De Koninck, Y., 2014. A simplified up-down method (SUDO) for measuring mechanical nociception in rodents using von Frey filaments. Mol. Pain 10, 26. https://doi.org/10.1186/1744-8069-10-26

Bourane, S., Grossmann, K.S., Britz, O., Dalet, A., Del Barrio, M.G., Stam, F.J., Garcia-Campmany, L., Koch, S., Goulding, M., 2015. Identification of a spinal circuit for light touch and fine motor control. Cell 160, 503–515. https://doi.org/10.1016/j.cell.2015.01.011

Callaghan, B.C., Cheng, H.T., Stables, C.L., Smith, A.L., Feldman, E.L., 2012a. Diabetic neuropathy: clinical manifestations and current treatments. Lancet Neurol. 11, 521–534. https://doi.org/10.1016/S1474-4422(12)70065-0

Callaghan, B.C., Little, A.A., Feldman, E.L., Hughes, R.A.C., 2012b. Enhanced glucose control for preventing and treating diabetic neuropathy. Cochrane Database Syst. Rev. 6, CD007543. https://doi.org/10.1002/14651858.CD007543.pub2

Coppey, L., Davidson, E., Lu, B., Gerard, C., Yorek, M., 2011. Vasopeptidase inhibitor ilepatril (AVE7688) prevents obesity- and diabetes-induced neuropathy in C57Bl/6J mice. Neuropharmacology 60, 259–266. https://doi.org/10.1016/j.neuropharm.2010.09.008

De Logu, F., Nassini, R., Materazzi, S., Carvalho Gonçalves, M., Nosi, D., Rossi Degl'Innocenti, D., Marone, I.M., Ferreira, J., Li Puma, S., Benemei, S., Trevisan, G., Souza Monteiro de Araújo, D., Patacchini, R., Bunnett, N.W., Geppetti, P., 2017. Schwann cell TRPA1 mediates neuroinflammation that sustains macrophage-dependent neuropathic pain in mice. Nat. Commun. 8, 1887. https://doi.org/10.1038/s41467-017-01739-2

Deuis, J.R., Vetter, I., 2016. The thermal probe test: A novel behavioral assay to quantify thermal paw withdrawal thresholds in mice. Temp. Multidiscip. Biomed. J. 3, 199–207. https://doi.org/10.1080/23328940.2016.1157668

Doerflinger, N.H., Macklin, W.B., Popko, B., 2003. Inducible site-specific recombination in myelinating cells. genesis 35, 63–72. https://doi.org/10.1002/gene.10154

Eid, S.A., Feldman, E.L., 2021. Advances in diet-induced rodent models of metabolically acquired peripheral neuropathy. Dis. Model. Mech. 14, dmm049337. https://doi.org/10.1242/dmm.049337

Feldman, E.L., Nave, K.-A., Jensen, T.S., Bennett, D.L.H., 2017. New Horizons in Diabetic Neuropathy: Mechanisms, Bioenergetics, and Pain. Neuron 93, 1296–1313. https://doi.org/10.1016/j.neuron.2017.02.005

Gomez-Sanchez, J.A., Pilch, K.S., van der Lans, M., Fazal, S.V., Benito, C., Wagstaff, L.J., Mirsky, R., Jessen, K.R., 2017. After Nerve Injury, Lineage Tracing Shows That Myelin and Remak Schwann Cells Elongate Extensively and Branch to Form Repair Schwann Cells, Which Shorten Radically on Remyelination. J. Neurosci. Off. J. Soc. Neurosci. 37, 9086–9099. https://doi.org/10.1523/JNEUROSCI.1453-17.2017

517   Green, A.Q., Krishnan, S., Finucane, F.M., Rayman, G., 2010. Altered C-fiber function as an indicator
518   of early peripheral neuropathy in individuals with impaired glucose tolerance. Diabetes Care
519   33, 174–176. https://doi.org/10.2337/dc09-0101
520   Grove, M., Kim, H., Santerre, M., Krupka, A.J., Han, S.B., Zhai, J., Cho, J.Y., Park, R., Harris, M., Kim, S.,
521   Sawaya, B.E., Kang, S.H., Barbe, M.F., Cho, S.-H., Lemay, M.A., Son, Y.-J., 2017. YAP/TAZ
522   initiate and maintain Schwann cell myelination. eLife 6, e20982.
523   https://doi.org/10.7554/eLife.20982
524   Hargreaves, K., Dubner, R., Brown, F., Flores, C., Joris, J., 1988. A new and sensitive method for
525   measuring thermal nociception in cutaneous hyperalgesia. Pain 32, 77–88.
526   https://doi.org/10.1016/0304-3959(88)90026-7
527   Hu, X., Agarwal, N., Zhang, M.-D., Ernfors, P., Kuner, R., Nyengaard, J.R., Karlsson, P., 2022.
528   Identification and quantification of nociceptive Schwann cells in mice with and without
529   Streptozotocin-induced diabetes. J. Chem. Neuroanat. 123, 102118.
530   https://doi.org/10.1016/j.jchemneu.2022.102118
531   Jende, J.M.E., Groener, J.B., Rother, C., Kender, Z., Hahn, A., Hilgenfeld, T., Juerchott, A., Preisner, F.,
532   Heiland, S., Kopf, S., Pham, M., Nawroth, P., Bendszus, M., Kurz, F.T., 2019. Association of
533   Serum Cholesterol Levels With Peripheral Nerve Damage in Patients With Type 2 Diabetes.
534   JAMA Netw. Open 2, e194798. https://doi.org/10.1001/jamanetworkopen.2019.4798
535   Lauria, G., Faber, C.G., Cornblath, D.R., 2022. Skin biopsy and small fibre neuropathies: facts and
536   thoughts 30 years later. J. Neurol. Neurosurg. Psychiatry jnnp-2021-327742.
537   https://doi.org/10.1136/jnnp-2021-327742
538   Leone, D.P., Genoud, S., Atanasoski, S., Grausenburger, R., Berger, P., Metzger, D., Macklin, W.B.,
539   Chambon, P., Suter, U., 2003. Tamoxifen-inducible glia-specific Cre mice for somatic
540   mutagenesis in oligodendrocytes and Schwann cells. Mol. Cell. Neurosci. 22, 430–440.
541   https://doi.org/10.1016/s1044-7431(03)00029-0
542   Li, R., Wu, Y., Zou, S., Wang, X., Li, Y., Xu, K., Gong, F., Liu, Y., Wang, J., Liao, Y., Li, X., Xiao, J., 2017.
543   NGF Attenuates High Glucose-Induced ER Stress, Preventing Schwann Cell Apoptosis by
544   Activating the PI3K/Akt/GSK3β and ERK1/2 Pathways. Neurochem. Res. 42, 3005–3018.
545   https://doi.org/10.1007/s11064-017-2333-6
546   Liu, Y.-P., Shao, S.-J., Guo, H.-D., 2020. Schwann cells apoptosis is induced by high glucose in diabetic
547   peripheral neuropathy. Life Sci. 248, 117459. https://doi.org/10.1016/j.lfs.2020.117459
548   Malik, R.A., Tesfaye, S., Newrick, P.G., Walker, D., Rajbhandari, S.M., Siddique, I., Sharma, A.K.,
549   Boulton, A.J.M., King, R.H.M., Thomas, P.K., Ward, J.D., 2005. Sural nerve pathology in
550   diabetic patients with minimal but progressive neuropathy. Diabetologia 48, 578–585.
551   https://doi.org/10.1007/s00125-004-1663-5
552   Margolis, D.J., Malay, D.S., Hoffstad, O.J., Leonard, C.E., MaCurdy, T., de Nava, K.L., Tan, Y., Molina,
553   T., Siegel, K.L., 2011. Incidence of diabetic foot ulcer and lower extremity amputation among
554   Medicare beneficiaries, 2006 to 2008: Data Points #2, in: Data Points Publication Series.
555   Agency for Healthcare Research and Quality (US), Rockville (MD).
556   Nagisa, Y., Kato, K., Watanabe, K., Murakoshi, H., Odaka, H., Yoshikawa, K., Sugiyama, Y., 2003.
557   Changes in glycated haemoglobin levels in diabetic rats measured with an automatic affinity
558   HPLC. Clin. Exp. Pharmacol. Physiol. 30, 752–758. https://doi.org/10.1046/j.1440-
559   1681.2003.03902.x
560   Nolano, M., Tozza, S., Caporaso, G., Provitera, V., 2020. Contribution of Skin Biopsy in Peripheral
561   Neuropathies. Brain Sci. 10, 989. https://doi.org/10.3390/brainsci10120989
562   Obrosova, I.G., Li, F., Abatan, O.I., Forsell, M.A., Komjáti, K., Pacher, P., Szabó, C., Stevens, M.J., 2004.
563   Role of poly(ADP-ribose) polymerase activation in diabetic neuropathy. Diabetes 53, 711–
564   720. https://doi.org/10.2337/diabetes.53.3.711
565   Oh, S.S., Hayes, J.M., Sims-Robinson, C., Sullivan, K.A., Feldman, E.L., 2010. The effects of anesthesia
566   on measures of nerve conduction velocity in male C57Bl6/J mice. Neurosci. Lett. 483, 127–
567   131. https://doi.org/10.1016/j.neulet.2010.07.076

Ojeda-Alonso, J., Calvo-Enrique, L., Paricio-Montesinos, R., Kumar, R., Zhang, M.-D., Poulet, J.F.A., Ernfors, P., Lewin, G.R., 2023. Sensory Schwann cells set perceptual thresholds for touch and selectively regulate mechanical nociception. https://doi.org/10.1101/2022.02.04.477749

Raut, N.G.R., Maile, L.A., Oswalt, L.M., Mitxelena, I., Adlakha, A., Sprague, K.L., Rupert, A.R., Bokros, L., Hofmann, M.C., Patritti-Cram, J., Rizvi, T.A., Queme, L.F., Choi, K., Ratner, N., Jankowski, M.P., 2024. Schwann cells modulate nociception in neurofibromatosis 1. JCI Insight 9. https://doi.org/10.1172/jci.insight.171275

Reinisch, C.M., Traxler, H., Piringer, S., Tangl, S., Nader, A., Tschachler, E., 2008. Rarefaction of the Peripheral Nerve Network in Diabetic Patients Is Associated With a Pronounced Reduction of Terminal Schwann Cells. Diabetes Care 31, 1219–1221. https://doi.org/10.2337/dc07-1832

Rinwa, P., Calvo-Enrique, L., Zhang, M.-D., Nyengaard, J.R., Karlsson, P., Ernfors, P., 2021. Demise of nociceptive Schwann cells causes nerve retraction and pain hyperalgesia. Pain 162, 1816–1827. https://doi.org/10.1097/j.pain.0000000000002169

Rumora, A.E., Kim, B., Feldman, E.L., 2022. A Role for Fatty Acids in Peripheral Neuropathy Associated with Type 2 Diabetes and Prediabetes. Antioxid. Redox Signal. 37, 560–577. https://doi.org/10.1089/ars.2021.0155

Sajic, M., Rumora, A.E., Kanhai, A.A., Dentoni, G., Varatharajah, S., Casey, C., Brown, R.D.R., Peters, F., Hinder, L.M., Savelieff, M.G., Feldman, E.L., Smith, K.J., 2021. High Dietary Fat Consumption Impairs Axonal Mitochondrial Function In Vivo. J. Neurosci. 41, 4321–4334. https://doi.org/10.1523/JNEUROSCI.1852-20.2021

Schulz, A., Walther, C., Morrison, H., Bauer, R., 2014. In vivo electrophysiological measurements on mouse sciatic nerves. J. Vis. Exp. JoVE 51181. https://doi.org/10.3791/51181

Shun, C.-T., Chang, Y.-C., Wu, H.-P., Hsieh, S.-C., Lin, W.-M., Lin, Y.-H., Tai, T.-Y., Hsieh, S.-T., 2004. Skin denervation in type 2 diabetes: correlations with diabetic duration and functional impairments. Brain J. Neurol. 127, 1593–1605. https://doi.org/10.1093/brain/awh180

Smith, A.G., Ramachandran, P., Tripp, S., Singleton, J.R., 2001. Epidermal nerve innervation in impaired glucose tolerance and diabetes-associated neuropathy. Neurology 57, 1701–1704. https://doi.org/10.1212/wnl.57.9.1701

Speakman, J.R., 2019. Use of high-fat diets to study rodent obesity as a model of human obesity. Int. J. Obes. 2005 43, 1491–1492. https://doi.org/10.1038/s41366-019-0363-7

Stino, A.M., Rumora, A.E., Kim, B., Feldman, E.L., 2020. Evolving concepts on the role of dyslipidemia, bioenergetics, and inflammation in the pathogenesis and treatment of diabetic peripheral neuropathy. J. Peripher. Nerv. Syst. JPNS 25, 76–84. https://doi.org/10.1111/jns.12387

Sumner, C.J., Sheth, S., Griffin, J.W., Cornblath, D.R., Polydefkis, M., 2003. The spectrum of neuropathy in diabetes and impaired glucose tolerance. Neurology 60, 108–111. https://doi.org/10.1212/wnl.60.1.108

Themistocleous, A.C., Ramirez, J.D., Shillo, P.R., Lees, J.G., Selvarajah, D., Orengo, C., Tesfaye, S., Rice, A.S.C., Bennett, D.L.H., 2016. The Pain in Neuropathy Study (PiNS): a cross-sectional observational study determining the somatosensory phenotype of painful and painless diabetic neuropathy. Pain 157, 1132–1145. https://doi.org/10.1097/j.pain.0000000000000491

Van Putten, L.M., 1958. The life span of red cells in the rat and the mouse as determined by labeling with DFP32 in vivo. Blood 13, 789–794.

Vincent, A.M., Hayes, J.M., McLean, L.L., Vivekanandan-Giri, A., Pennathur, S., Feldman, E.L., 2009. Dyslipidemia-induced neuropathy in mice: the role of oxLDL/LOX-1. Diabetes 58, 2376–2385. https://doi.org/10.2337/db09-0047

Wang, X., Xu, G., Liu, H., Chen, Z., Huang, S., Yuan, J., Xie, C., Du, L., 2023. Inhibiting apoptosis of Schwann cell under the high-glucose condition: A promising approach to treat diabetic peripheral neuropathy using Chinese herbal medicine. Biomed. Pharmacother. 157, 114059. https://doi.org/10.1016/j.biopha.2022.114059

618   Wu, K., Deng, F., Mao, X., Zhou, D., Shen, W., 2023. Ferroptosis involves in Schwann cell death in
619   diabetic peripheral neuropathy. Open Med. 18, 20230809. https://doi.org/10.1515/med-
620   2023-0809

621   Xu, L., Lin, X., Guan, M., Zeng, Y., Liu, Y., 2019. Verapamil Attenuated Prediabetic Neuropathy in High-
622   Fat Diet-Fed Mice through Inhibiting TXNIP-Mediated Apoptosis and Inflammation. Oxid.
623   Med. Cell. Longev. 2019, 1896041. https://doi.org/10.1155/2019/1896041

624   Yeomans, D.C., Proudfit, H.K., 1996. Nociceptive responses to high and low rates of noxious
625   cutaneous heating are mediated by different nociceptors in the rat: electrophysiological
626   evidence. Pain 68, 141–150. https://doi.org/10.1016/S0304-3959(96)03177-6

627   Yu, Y., 2021. Gold Standard for Diagnosis of DPN. Front. Endocrinol. 12, 719356.
628   https://doi.org/10.3389/fendo.2021.719356

629   Zhang, X., Zhao, S., Yuan, Q., Zhu, L., Li, F., Wang, H., Kong, D., Hao, J., 2021. TXNIP, a novel key factor
630   to cause Schwann cell dysfunction in diabetic peripheral neuropathy, under the regulation of
631   PI3K/Akt pathway inhibition-induced DNMT1 and DNMT3a overexpression. Cell Death Dis.
632   12, 1–13. https://doi.org/10.1038/s41419-021-03930-2

633   Ziegler, D., Papanas, N., Zhivov, A., Allgeier, S., Winter, K., Ziegler, I., Brüggemann, J., Strom, A.,
634   Peschel, S., Köhler, B., Stachs, O., Guthoff, R.F., Roden, M., German Diabetes Study (GDS)
635   Group, 2014. Early detection of nerve fiber loss by corneal confocal microscopy and skin
636   biopsy in recently diagnosed type 2 diabetes. Diabetes 63, 2454–2463.
637   https://doi.org/10.2337/db13-1819

 Figure Legends

 Figure 1: Study design and HFD-induced dyslipidemia. (A) Study Design used to assess the
 effect of a HFD (45% kcal %fat) vs. a control diet (10% kcal %fat) in two groups of mice
 (n=10/group). (**B**) Body weight of control diet-fed mice and HFD-fed mice during the 16 weeks
 of diet. (**C**) Leptinemia of control and HFD mice at 16 weeks of diet. (**D**) Plasma levels
 triglycerides of control and HFD mice at 16 weeks of diet. (**E**) Total cholesterolemia of control
 and HFD mice at 16 weeks of diet. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$: control diet vs. HFD.
 Data are presented as means ± SEM.

 Figure 2: HFD-induced impaired glucose tolerance. (A) Fasting blood glucose of control diet-
 fed mice and HFD-fed mice at 16 weeks of diet. **(B)** Percentage of glycated hemoglobin
 (HbA1c) of control diet-fed mice and HFD-fed mice at 16 weeks of diet. **(C)** Plasma insulin of
 control diet-fed mice and HFD-fed mice at 16 weeks of diet. **(D)** Oral glucose tolerance test
 (OGTT) curves performed at 16 weeks of diet in control and HFD-fed mice. **(E)** Area under the
 curve (AUC) during OGTT for control diet-fed mice and HFD-fed mice. *$P < 0.05$, **$P < 0.01$,
 ***$P < 0.001$: control diet vs. HFD. Data are presented as means ± SEM (n=10/group).

 Figure 3: Nerve conduction studies. (A) Motor nerve conduction velocity (MNCV) performed
 at 16 weeks of diet in control and HFD-fed mice. **(B)** Sensory nerve conduction velocity (SNCV)
 performed at 16 weeks of diet in control and HFD-fed mice**.** Data are presented as means ±
 SEM (n=10/group).

 Figure 4: Mechanical and thermal sensitivity changes induced by HFD. (A) Quantification of
 static mechanical sensitivity was assessed every 4 weeks in control and HFD-fed mice using
 von Frey filaments. **(B)** Quantification of thermal sensitivity by Hargreaves test every 4 weeks
 in control and HFD-fed mice. **(C)** Quantification of haptic mechanical sensitivity by brush test
 every 4 weeks in control and HFD-fed mice. **(D)** Quantification of acute mechanical pain by
 pinprick test every 4 weeks in control- and HFD-fed mice. **$P < 0.01$, ***$P < 0.001$: control
 diet vs. HFD. #$P < 0.05$, ##$P < 0.01$: HFD 4weeks vs. HFD 8, 12 or 16 weeks. Data are presented
 as means ±SEM (n=10/group).

 Figure 5: HFD-induced small nerve fiber and nociceptive Schwann cells degeneration in the
 foot skin. (A) PGP9.5 immunofluorescence with DAPI staining in foot skin section of control
 (n=5) and HFD-fed mice (n=5). **(B)** Quantitation of IENFD is presented as the number of
 fibers/mm of epidermis. **(C)** L1CAM immunofluorescence with DAPI staining in foot skin

section of control (n=5) and HFD-fed mice (n=5). Arrowhead indicate nerve fibers in the
epidermis of the foot skin. Arrows indicate nociceptive Schwann cells and their cellular
extensions at the border of the epidermis and the dermis. **(D)** Mean fluorescence intensity
quantification of L1CAM immunostaining at the localization of nociceptive Schwann cells (at
the border of epidermis and dermis). **(E)** Quantification of L1CAM-positive cells and their
cellular extensions presented in number/mm of epidermis. $**P < 0.01$, $***P < 0.001$: control
diet vs. HFD. Data are presented as means ±SEM. Scale bar: 50 μm.

Supplementary Figure 1: Colocalization of classical nociceptive Schwann cell markers with
L1CAM. (A) L1CAM immunofluorescence colocalized with Sox10 and S100β classical markers
of nociceptive Schwann cells in the foot skin. **(B)** L1CAM immunofluorescence colocalized with
Td-tomato expressing nociceptive Schwann cells in a tamoxifen-induced PLP-*Cre*ER[T2];
R26TdTomato double transgenic mouse. White arrows head indicate nociceptive Schwann
cells and their cellular extensions at the border of the epidermis and dermis. Scale bar: 50 μm.

692