1

The gut contractile organoid: a novel model for studying the gut motility regulated by coordinating signals between interstitial cells of Cajal and smooth muscles

Rei Yagasaki[1,2], Ryo Nakamura[1,3], Yuuki Shikaya[1,4], Ryosuke Tadokoro[1,5], Ruolin Hao[6,7], Zhe Wang[6,7], Mototsugu Eiraku[6,7], Masafumi Inaba[1] and Yoshiko Takahashi[1,8]

1: Department of Zoology, Graduate School of Science, Kyoto University Sakyo-ku, Kyoto 606-8502, Japan

2: Present Address: WPI Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

3: Present Address: Ushimado Marine Institute, Okayama University, Setouchi, Okayama 701-4303, Japan

4: Present Address: Graduate School of Science, University of Hyogo, Kamigori-cho, Hyogo 678-1297, Japan

5: Present Address: Department of Bioscience, Okayama University of Science, Okayama, Japan

6: Institute for Life and Medical Sciences, Kyoto University, Sakyo-ku, Kyoto 606-8507, Japan

7: Department of Polymer Chemistry, Graduate School of Engineering, Kyoto University, Kyoto 615-8510, Japan

8: Author to whom all correspondence should be addressed.

email: yotayota@develop.zool.kyoto-u.ac.jp

Keywords:

gut contraction, interstitial cells of Cajal, smooth muscle cells, organoid, gap junction

Abstract

The gut undergoes peristaltic movements regulated by intricate cellular interactions. How these interactions emerge in the developing gut remains poorly explored due to a lack of model system. We here developed a novel contractile organoid that is derived from the muscle layer of chicken embryonic hindgut. The organoid contained smooth muscle cells (SMCs) and interstitial cells of Cajal (ICCs; pacemaker) with few enteric neurons, and underwent periodic contractions. The organoid was formed by self-organization with morphological arrangements of ICCs (internal) and SMCs (peripheral), allowing identification of these cells in live. GCaMP-Ca^{2+} imaging analyses revealed that Ca^{2+} transients between ICC-ICC, SMC-SMC or SMC-ICC were markedly coordinated. Pharmacological studies further suggested a role of gap junctions in ICC-to-SMC signaling, and also possible mechanical feedback from SMC's contraction to ICC's pace-making activities. In addition, two organoids with different rhythm became synchronized when mediated by SMCs, unveiling a novel contribution of SMCs to ICC's pace-making. The gut contractile organoid developed in this study offers a useful model to understand how the rhythm coordination between/among ICCs and SMCs is regulated and maintained during gut development.

Introduction

In the gut, ingested material is conveyed properly along the gut axis by the gut movements called peristalsis, which is recognized as wave-like propagation of a local constriction. The physiology of gut peristalsis has extensively been studied in adults, where the peristalsis plays pivotal roles in effective transportation and digestion/absorption of inter-luminal contents. And many cases of gut-related pathology are associated with peristaltic disfunctions. The gut peristaltic movements are achieved by intricate regulations of intercellular functions among multiple cell types. At the site of origin of peristaltic waves (OPW), a local constriction emerges along the circumferential axis, and this is soon followed by a progressive wave of the contraction along the gut axis. During these processes, the circumferential constriction demands multiple SMCs to achieve simultaneous contraction. However, due to the complex structure of the gut, how such synchronization/coordination in SMCs is regulated remains largely undetermined.

It is known in vertebrates that the embryonic gut undergoes peristaltic movements even without experience of food intake. The embryonic gut therefore serves as a powerful model to study the intrinsic mechanisms underlying the peristalsis contrasting with the adult gut where ingested content influences the gut motility increasing complexity in analyses. We have recently reported using chicken embryos that sites of OPW are randomly distributed along the gut axis at early stages, and they later become confined to specific sites, and that this confinement of OPWs enables rhythmic and patterned peristaltic movements (Shikaya et al., 2022). One of the long-standing and important questions is how the synchronized/coordinated contraction is achieved and maintained.

In a long history of gut motility studies, it has been known that enteric nervous system (ENS), smooth muscle cells (SMCs) and interstitial cells of Cajal (ICCs) play important roles in peristaltic movements (Barajas-Lopez et al., 1989; Camborova et al., 2003; Chevalier et al., 2020; Huizinga et al., 1995; Kito et al., 2005; Liu et al., 1998; Rumessen & Thuneberg, 1996; Sanders et al., 1991; Takaki, 2003; Takayama et al., 2002; Thomsen et al., 1998). It has widely been accepted that: 1) at early embryonic stages, gut movement/peristalsis does not require ENS activity, 2) ICCs serve as a pacemaker dictating their rhythm to SMCs, 3) the contraction is executed by SMC and not by ICCs, since thick fibers of myosin are found solely in SMCs. A series of elaborate electrophysiological studies showed that slow waves (a type of changes in membrane potential characteristic of gut movements) occur spontaneously in ICCs but not in SMCs, and that these slow waves lead to voltage-dependent Ca^{2+} influx evoking action potential, which is somehow transmitted to SMCs to execute gut contraction in register with ICC's

pace-making rhythm (Baker et al., 2021; Torihashi et al., 2002). The knowledge that ICCs

act as a pacemaker was supported by compelling evidence obtained by c-Kit-deficient

mouse mutants (W/W^v), in which ICC differentiation was severely affected leading to a

failure of gut peristalsis (Huizinga et al., 1995; Torihashi et al., 1999). However, it

remains largely unexplored how the intrinsic/spontaneous rhythm in a single ICC

becomes synchronized among multiple ICCs which constitute intricate networks in the

gut. It has also been under debate to what extend the gap junction contributes to cell-cell

communications between ICCs, SMCs or ICC-SMC. One reason is that it has been

difficult to stably maintain SMCs and ICCs in cell culture condition, and also to

distinguish ICCs from SMCs in the living gut (these two types of cells originate from the

same progenitor of splanchnopleural mesoderm during development). Thus, a novel

model system has been awaited to circumvent these obstacles. Recently, it was reported

that differentiation states of mouse hindgut-derived cells were successfully maintained

for a relatively long period of time in a serum-free culture medium (Wang et al., 2018).

In that study, many types of gut-derived cells including not only ENS, ICC, SMCs but

also glial cells and serosa were observed, and this large and complex cell mass was seen

to undergo rhythmic contractions *in vitro*.

It has widely been appreciated that organoids can serve as a powerful model and

tool to circumvent such obstacles of organ complexity. Relatively simple structures of

organoids allow analyses at higher resolution than *in vivo*, and also permit analyses of

cell behaviors in three-dimensional (3D) environment, which might be different from

behaviors *in vitro* confined to two dimensions. In the current study, we have developed a

novel organoid called "gut contractile organoid" by culturing chicken hindgut-derived

cells in a serum-free medium. The gut contractile organoid undergoes periodic

contractions, and it is essentially composed of ICCs and SMCs, the former residing

centrally whereas the latter peripherally, allowing distinction between the two cell types

in living organoids. These advantages enabled GCaMP-live imaging of Ca^{2+} dynamics,

and revealed coordinated oscillations of Ca^{2+} transients between ICC-ICC, SMC-SMC

and SMC-ICC. Pharmacological studies further suggested a role of gap junctions in an

ICC-to-SMC signaling, and also a possible mechanical feedback from SMC's

contractions to ICC's pace-making activities. In addition, by regarding an organoid as a

single oscillator unit and by placing these oscillators separately in a hydrogel mold, we

found that two oscillators with different rhythm become synchronized when mediated by

SMCs, supporting the notion of SMC's contribution to ICC's pace-making. The gut

contractile organoid developed in this study must be useful to unveil the intrinsic

mechanisms underlying the rhythm coordination and its maintenance between/among
ICCs and SMCs during peristaltic movements in the embryonic gut.
121
122
Results
Spheroids were formed from muscle layer-derived cells of the embryonic hindgut
To develop a culture condition that would facilitate analyses of gut motility, we
dissected the muscle layer (also called *tunica muscularis*) from the hindgut of chicken
embryos of embryonic day 15 (E15) by removing the serosa and intestinal epithelium
(mucosa) (Fig. S1). The isolated muscle layer was dissociated into single cells to prepare
5.0×10^5 cells per culture dish. We started analyses with a culture condition with FBS-free
medium and Matrigel as substrate as previously described for cultures in mice of
gastrointestinal cells including serosa (Wang et al., 2018). We tested three kinds of FBS-
free media: DMEM/Ham's F-12, Ham's F-12, and Neurobasal media (see Materials and
Methods). When cultured in DMEM/Ham's F-12 or Ham's F-12, dissociated cells formed
very small aggregates containing several cells at day 1, the morphology of which did not
change significantly until day 5 (Fig. 1A). In clear contrast, when cultured in the
Neurobasal medium, cells formed clusters that were interconnected by elongated cells
with neighboring clusters as early as day 1. These clusters grew as larger aggregates by
day 3, and became spherical by day 5. Such spheroids were not observed in the conditions
with Ham's F-12 or DMEM/Ham's F-12. We also tested different substrates, Poly-Lysine
or collagen for dish coating with the Neurobasal medium, but neither one yielded spheroid.
To know how the spherical aggregates were formed under the condition of Neurobasal
medium and Matrigel, we obtained time-lapse images at two-hour intervals. Originally
sparse clusters that were loosely connected by elongated cells merged each other forming
progressively larger clusters (Fig. 1B, Movie 1). In the following experiments, we focused
on the spheroids formed under the condition of Neurobasal medium and Matrigel coating.
146
The gut muscle layer-derived spheroids displayed periodic contractions
The spheroids underwent reiterative contractions at day 3, and these contractions
were observed at least until day 7 of culture (Fig. 2A for normalized intensities of 3
representative organoids for each stage, Movie 2). Time-lapse imaging of the
contractions combined with quantitative assessments by MATLAB (MathWorks, see
Materials and Methods) showed that interval periods between two successive peaks were
13.3 seconds (sec), 15.4 sec and 19.6 sec for cultures at day 3, day 5 and day 7,
respectively (median values) (Fig. 2B; contraction intervals).

The contracting spheroid was composed of ICCs and SMCs

To determine the cell types comprising the spheroids, we performed immunostaining with antibodies against the chicken c-Kit protein for ICCs (Yagasaki et al., 2022), and αSMA and desmin for SMCs. It is known that ICCs and SMCs are derived from common progenitors which are c-Kit$^+$/αSMA$^+$, and that after differentiation ICCs and SMCs are c-Kit$^+$/αSMA$^-$ and c-Kit$^-$/αSMA$^+$, respectively (Duband et al., 1993; Kluppel et al., 1998). In the clusters at day 3, c-Kit$^+$/αSMA$^-$ signals were detected in the internal region, and c-Kit$^+$/αSMA$^+$ signals at the periphery (Fig. 3A, Day 3), suggesting that internal cells were differentiated ICCs, whereas peripheral cells were ICC/SMC progenitors. At day 5 onward, c-Kit$^+$/αSMA$^-$ cells and c-Kit$^-$/αSMA$^+$ cells were segregated which were located internally and peripherally, respectively (Fig. 3A, Day 5), and this spatial segregation of the cells remained unchanged until day 7 (Fig. 3A, Day 7), the latest stage examined in this study. The peripherally lining cells were also positive for desmin staining, supporting that these cells were SMCs (Fig. 3B). Infection with RCAS-GapEGFP into a forming spheroid further visualized multipolar ICCs in the internal region (Fig. 3C), a morphology known to be characteristic of ICC-MY which would normally reside in the layer of myenteric plexus (Mei et al., 2009; Sanders et al., 2014). ICCs of a different type were also recognized, which were elongated, thin, and lining underneath the peripheral SMCs (Fig. 3A, Day 7). It is possible that these were ICC-IM, known to be embedded in and tightly associated with smooth muscles in the gut (Huizinga et al., 2011; Iino & Horiguchi, 2006). Unexpectedly, the spheroid contained few neural cells (ENS), if any, revealed by anti-Tuj1 antibody (Fig. S2A, B). Indeed, when tetrodotoxin was added to culture medium, contraction intervals of organoids were comparable to those of control (before the administration) (Fig. S2C) confirming little contribution by ENS. We counted cell numbers of each cell type per organoid, and found that contrasting with the intact gut, the proportion of SMCs was smaller than that of ICCs (Fig. S2B) possibly due to a cell type-specific loss during culture preparation.

Since ICCs and SMCs were spatially segregated in the sphenoid, we stained with anti N-cadherin antibody (E-cadherin is positive solely in the mucosa/endoderm) (Graham et al., 2017; Grosse et al., 2011). N-cadherin signals were seen in the internal ICCs, and not in the peripheral SMCs (Fig. 3D). Thus, it is likely that N-cadherin plays a role in the segregation.

In summary, the hindgut-derived spheroid displays three prominent characteristics: (1) a dominant occupation by ICCs and SMCs with neglectable

191 contribution by ENS, (2) self-organization ability of internal ICCs encapsuled by a thin
192 layer of SMCs, (3) recurrent and stable contractions. Based on these characteristics, we
193 designated this spheroid as "gut contractile organoid".
194

Contraction-associated intracellular Ca^{2+} transients were coordinated between
195
ICCs and SMCs in the gut contractile organoid
It has been reported that Ca^{2+} dynamics are important for pacemaker activity in
ICCs and contractions of SMCs. Ca^{2+} flows into ICCs via voltage-dependent Ca^{2+}
channels and propagates to SMCs, causing the gut muscle contractions (Baker et al.,
2021). We therefore investigated the Ca^{2+} dynamics in our gut contractile organoids.
Organoid-forming cells were infected with a RCAS-vector encoding GCaMP6s, a Ca^{2+}
indicator that emits EGFP fluorescence in response to Ca^{2+} influx, and mRuby3 as a
reporter (Fig. 4A). GCaMP6s-organoids were subjected to time-lapse imaging analyses
by confocal microscopy. As expected, the oscillatory rhythm of Ca^{2+} transients as a whole
organoid was highly concomitant with that of contractions (Fig. 4B, Movie 4).
Taking advantage of the spatial segregation between ICCs (internal) and SMCs
(peripheral) (Fig. 3A, B), we compared the Ca^{2+} oscillatory rhythm among ICCs and
SMCs (homotypically) and between ICC-SMC (heterotypically). We set up 3 regions of
interests (ROIs) in the SMC layer with one ROI corresponding to one cell, and captured
the Ca^{2+} transients. The three ROIs exhibited a synchronous pattern of Ca^{2+} oscillations
(Fig. 4C, SMCs). Similarly, Ca^{2+} oscillations in three ROIs in the central region (ICCs)
were synchronous (Fig. 4C, ICCs). These data highlight active communications taking
place intercellularly within SMCs and ICCs, respectively. We further compared Ca^{2+}
oscillations between ICC-SMC by setting up one ROI in each of ICC and SMC. Again,
the Ca^{2+} rhythm was synchronized between these heterotypic cell types (Fig. 4C,
ICC/SMC). With a deeper scrutinization, a peak of Ca^{2+} transient in ICC preceded that in
SMC with a time lag (also called latency) of 700 msec in average (Fig. 4C, ICC/SMC;
Fig. 4D, 117 peaks for 14 organoids), suggestive of a signal propagation from ICC to
SMC consistent with previous reports (Baker et al., 2021).
220

Gap junction plays a role in ICC-to-SMC signaling
221
Although a series of gut motility studies have proposed an importance of gap
junctions, rigid evidence has been limited due to a lack of experimental model. Our gut
contractile organoid should prove useful for clarifying the roles of gap junctions in the
synchronous motility, since intercellular synchronization was observed between/among
identifiable cells (ICCs and SMCs) in live as shown above. We performed

pharmacological assessments using gap junction inhibitors including carbenoxolone
(CBX) and 18β-glycyrrhetinic acid (18β-GA; Chevalier, 2018; Takeda et al., 2005), two
of the most widely used pharmacological inhibitors for gap junction studies.
Following inhibitor administration into day 7 culture medium, organoids were
allowed to rest for 30 min to exclude possible effects by the administration, for example,
turbulence of the medium. Contrary to our expectation, none of 100 μM concentrations
of CBX or 18β-GA showed detectable effects on the recurring contractions of an organoid
with frequency/interval comparable to the control (Fig. S3A and B, Movie 3). These
undetectable effects were not due to insufficient penetrance of the drugs into the organoid,
since the drug administration at day 3, when organoid-forming cells were still at 2D in
culture, also yielded undetectable effects assessed at day 7 (Fig. S3C). The inhibiting
activity of the drugs used here was verified using embryonic heart cultured cells (Fig.
S3D).
Such unchanged patterns were also observed for the synchronization/
coordination of Ca^{2+} transients either in ICCs or SMCs in experiments conducted in a
way similar to Fig. 4 (Fig. 5A, B, Movie 5). Same cells in an organoid were tracked for
their Ca^{2+} transients before and after the inhibitor administration. With a closer look,
however, while overall synchronization was retained between ICCs and SMCs, the
preceding peak of Ca^{2+} transient in ICC was abolished (Fig. 5C, D). Collectively, while
the contribution by gap junction to the periodic contraction and intercellular
synchronization in the Day 7 organoid is relatively limited, the ICC-to-SMC signals
require gap junction-mediated communications, at least partly.
249

Blebbistatin and Nifedipine ceased organoidal contractions and oscillatory
patterns of Ca^{2+} transients
Toward searching for factors that regulate the coordination between/among ICCs
and SMCs, we tested blebbistatin, a specific inhibitor of myosin II, which was expected
to cease organoidal contractions. Experimental procedures were similar to those for gap
junction inhibitors. We found that blebbistatin ceased periodic contractions of organoids
in a concentration-dependent manner: while intervals were shorter with smaller amplitude
at 5 μM, contractions were ceased completely at 10 μM (Fig. 6A, B, Movie 6). The 10
μM-treated specimens resumed contractions following medium-wash out, showing that
the organoids were alive (Fig. 6B).
We examined Ca^{2+} transients in these contraction-inhibited organoids. Markedly,
periodic Ca^{2+} transients were extinguished not only in SMCs but also in ICCs, yielding
no/little synchronous Ca^{2+} patterns among and between ICCs and SMCs (Fig. 6C-F).

Although a possible direct inhibition of non-muscle myosin II in ICCs cannot be excluded,

these findings raised an interesting possibility that the contractility feeds back to ICCs to

generate/maintain their periodic rhythm.

This notion was further corroborated by similar experiments using Nifedipine, a

blocker of L-type Ca^{2+} channel known to function in gut SMCs (Chevalier et al., 2024;

Der et al., 2000). Organoidal contractions were completely ceased by 1 μM Nifedipine,

and resumed after washout, at least partly (Fig. 7A, B). In Nifedipine-treated organoids,

Ca^{2+} transients (GCaMP) in ICCs were erased, which resumed following washout (Fig.

7C).

Inter-organoidal coordination was mediated by SMCs

During analyses with our novel organoids, we noticed that they easily fuse to each

other, suggesting that organoids grow by progressive fusion (Fig. 8A, Movie 7). This also

raised the possibility that the synchronous Ca^{2+} transients among organoid-constituting

cells (Fig. 4C) might be a consequence of phasic coordination upon the fusion of multiple

organoids that had shown different oscillatory phases. To test this possibility, we

transferred two organoids in a petri dish, and allowed them to fuse. Before the fusion,

Ca^{2+} oscillatory rhythm was indeed out of phase in the two organoids (Fig. 8B, Movie 8).

Markedly, upon fusion by 24 hours, their rhythm became in phase/synchronous (Fig. 8B,

Movie 8). Intriguingly, they underwent a "pause" of oscillation during fusion (see

Discussion).

Since we noticed that cellular protrusions were often observed around the time of

organoidal fusion, we reasoned that these cellular processes would mediate the fusion and

its subsequent synchronization of Ca^{2+} transients (Fig. 8D). To test this, we developed a

3-well hydrogel with narrow channels connecting the wells (Fig. 9A). Organoids were

placed separately in each well, which prevented the fusion of organoidal bodies, but

allowed extension of cellular processes through the narrow channels and contact with

each other (Fig. 9B). After 72 hours of placement of organoids, cellular processes as well

as several cell bodies were present within the channel, and Ca^{2+} transients became

synchronized among the three organoids (Fig. 9C, Movie 9). However, with careful

examination, we also noticed that organoid-derived cells crawled out from the wells to

cover the top surface of the hydrogel, connecting the three organoids (Fig. 9D). Thus,

another possibility was raised that these crawled out cells would mediate the inter-

organoidal synchronization.

To test this possibility, we prepared a similar 3-well hydrogel, but in this case the

three wells were disconnected (no channels) so that organoid-derived cellular processes

were not able to connect each other (Fig. 9E). By 3 days of culture, the top surface of the
hydrogel was indeed covered by cells in a similar way to Fig. 9D. These cells were
positive for αSMA but negative for c-Kit (Fig. 9F), showing that they were SMCs that
were somehow detached and crawled out from the peripheral layer of their "host"
organoids. Importantly, coincided with the top-coverage by the SMCs, the three
organoids in the disconnected wells displayed synchronized Ca^{2+} transients, highlighting
the role of SMCs in mediating coordination between organoids (Fig. 9G, Movie 10). Gap
junction inhibitor yielded no/little effects on the Ca^{2+} transient coordination (Fig. 9H,
Movie 11).
308
Discussion
We have developed a novel gut contractile organoid, which displays several
unique characteristics: 1) it undergoes recurrent contractions, 2) differentiation states of
ICCs (c-Kit$^+$/αSMA$^-$) and SMCs (c-Kit$^-$/αSMA$^+$) are maintained at least until day 7 in
the organoid, 3) the organoid is composed essentially of two types of cells, ICCs and
SMCs, with few ENS cells, if any, 4) ICCs (internal) and SMCs (peripheral) can be
distinguished for their localization in a living organoid, allowing 5) GCaMP-visualization
of Ca^{2+} transients and assessments of cell interactions between and among ICCs and
SMCs. These characteristics circumvent, at least partly, obstacles that have hampered
analyses in the research of gut peristalsis, such as unstable differentiation state of ICCs
and SMCs in cultures, and difficulties in identifying these cells in living preparations. In
studies of gut movements, how ICCs generate/maintain their periodic rhythm and how
ICCs and SMCs interact with each other have been long-standing questions, and our
organoids offer powerful advantages to address these fundamental questions and to
understand the cellular mechanisms underlying the gut contractions/peristaltic motility at
least in the embryonic gut.
Contrasting with many cases in organoid studies that aim at a *maximum*
recapitulation of the intact organ, our gut contractile organoid is composed of a (probably)
minimum number of cell types that suffice the generation and/or maintenance of rhythmic
contractions, allowing high-resolution analyses at the cellular level. For example, adding
ENS components to our organoid would allow the clarification of the role of ENS in the
gut contraction. While the majority of internal cells in the organoid are ICCs (c-
Kit$^+$/αSMA$^-$), the possibility cannot be excluded that platelet-derived growth factor
receptor-α positive (PDFGRα$^+$/c-Kit$^-$/αSMA$^-$) cells, another type of interstitial cells
(fibroblast-like cell) known to mediate neural activity to SMCs in the mouse gut (Sanders

et al., 2024; Sanders et al., 2016), are included in our organoid. Available antibody against
the chicken PDFGRα protein is awaited.
336

Coordinated Ca^{2+} transients in ICC/SMC populations in the gut contractile organoid

Measurement and quantification analyses with ICCs and SMCs that are
identifiable in the living organoid revealed exquisite coordination of Ca^{2+}
transients/oscillation homotypically in both ICC-ICC and SMC-SMC combinations, and
heterotypically between SMC-ICC (Fig. 4C). Notably, a peak of Ca^{2+} transient in ICCs
precedes that of SMCs with a time lag (also called latency) of 700 msec, implying a
signaling from ICC to SMC. These observations are consistent with previous studies
using ICC- and SMC-specific transgenic mice that expressed GCaMP and RCaMP,
respectively (Baker et al., 2021). In that study, the authors assessed Ca^{2+} transients in
submucosal ICCs (ICC-SM) and compared them with those in their adjacent circular
muscles, and showed that the rise of GCaMP signal (ICC) preceded that of RCaMP
(SMC) with a latency of 56 +/- 14 msec. With these observations, they concluded that
ICC-SM sends signals to its adjacent SMC. In our study, time-lapse imaging was mostly
performed with 700 msec intervals. Further studies with shorter intervals are awaited to
know whether the latency time would be shorter than 700 msec in average in our
organoids.
354

Contribution of gap junction to ICC-to-SMC signaling

Effects by the block of gap junction by CBX or 18β-GA were relatively limited
in our organoid assays: organoidal contraction rhythm and synchronous patterns of Ca^{2+}
transients remained unchanged between SMC-SMC and ICC-ICC. In contrast, signaling
from ICCs to SMCs was affected, in which the preceding peak of Ca^{2+} transient in ICCs
was abolished. This effect was seen by CBX but not by 18β-GA (Fig. 5C, S3E). Further
studies are required to clarify which connexin(s) plays a role in the generation of latency.
The contribution of gap junction to the ICC-to-SMC signaling was previously reported in
mouse gut acting from intramuscular ICCs (ICC-IM) to their adjacent circular smooth
muscles. However, interpretation of the role of gap junction in ICC-SMC interactions, in
general, has been under big debate. Some studies reported that CBX or 18β-GA failed to
inhibit these interactions or peristaltic motilities (Komuro et al., 1996; Rohr et al., 1998;
Schultz et al., 2003), or that electron microscopy did not detect structures of gap junction
in longitudinal muscles (Cousins et al., 2003; Daniel & Wang, 1999; Gabella & Blundell,
1981). Contribution of gap junctions to the gut motility appears to be highly variable in

different regions of gastrointestinal tract, for example, stomach versus colon (Iino et al.,
2007; Yang et al., 2012) . And a detection of gap junction structures or mRNA/protein of
connexins does not necessarily mean that the gap junction is functional. Dominant
negative form of gap junction might be useful. Currently, ICC- or SMC-specific gene
manipulations in our organoid are not available, and further studies are needed. One
possibility is that while gap junctions are important for the onset of rhythm coordination,
once the rhythm is established, other mechanisms might be employed, for example,
mechanical feedback from muscles to ICCs (see below). Similar notion has recently been
shown for coordinated Ca^{2+} signaling in cardiac muscle cells (Fukui et al., 2021).
379

Possible feedback from SMC's contractility to ICC's oscillatory rhythm
Blebbistatin extinguished the contraction of the organoids, which was
concomitant with abrogation of Ca^{2+} transients in ICCs (Fig. 6). Since it has been reported
that thick myosin fibers necessary for the contraction are found only in SMCs but not in
ICCs (Gherghiceanu & Popescu, 2005; Rumessen & Thuneberg, 1991, 1996; Sun et al.,
2006), it is likely that ICCs do not have contractile ability. Our observations therefore
raise the possibility that ICC's pace-making activity requires mechanical feedback from
contracting SMCs. This notion is also supported by additional finding that Nifedipine, an
L-type Ca^{2+} channel blocker known to be expressed in SMCs (Chevalier et al., 2024; Der
et al., 2000), erases both organoidal contractions and Ca^{2+} transients in ICCs (Fig. 7). The
possibility of the SMC-to-ICC signaling is further corroborated by other findings
obtained in this study showing that SMCs mediate inter-organoidal rhythm coordination
(Fig. 9) as more discussed below.
It is of note that the reciprocal interactions between pace-making cells and their
effectors have been reported in studies of neural circuit establishment. Spontaneous
activities emerging in motor neurons during peristaltic locomotion of larvae in
Drosophila are regulated by feedback from their governing muscle's contractions, so that
motor neurons get organized to display coordinated and stable oscillatory activities (Zeng
et al., 2021). Thus, it is tempting to speculate that during early gut peristalsis, ICCs that
have initiated their spontaneous activities receive feedback from their governing SMCs
to generate more stable coordination of pace-making activity among ICCs. Indeed, we
have previously reported that in the very early embryonic gut, origins of peristaltic wave
(OPWs) are randomly distributed along the gut axis, but these unstable OPWs later
become confined to specific sites displaying more stable and coordinated pace-making
patterns (Shikaya et al., 2022).
405

Interactions between ICCs and SMCs in the gut contractile organoid

The organoid developed in this study was derived from the muscle layer (also called *tunica muscularis*) of chicken E15 hindgut devoid of mucosa and serosa, in which myenteric ICCs (ICC-MY), intramuscular ICCs (ICC-IM), and submucosal ICCs (ICC-SM) are localized in a way similar to those in mice, shown by staining with antibody against the chicken c-Kit protein (Yagasaki et al., 2022). In the current study, c-Kit antibody-staining showed two types of cells in morphology in the gut contractile organoid: one is multipolar and N-cad-positive ICCs located centrally, and the other is ICCs that are thin in shape, N-cad negative, and lining beneath the most external layer of SMCs. Based on the knowledge obtained in studies with mammalian species that ICC-MY are multipolar whereas ICC-IM are bipolar and tightly associated with adjacent SMCs (Huizinga et al., 2011; Iino & Horiguchi, 2006), it is conceivable in our organoids that the central cells are ICC-MY, and the peripherally lining ones are ICC-IM. In addition, it has been reported in mice that signaling from ICC-IM to SMCs is gap junction-dependent. Collectively, our observations that gap junction-dependent signal found between ICC (central ICC) and SMC in the organoid (Fig. 5C, D) could be interpreted as follows: ICC-MY (central) signals to ICC-IM (second-most peripheral), which in turn acts on the external SMCs mediated partly by gap junction. Such sequences of signaling (ICC-MY to ICC-IM to SMCs) have also been proposed in the intact gut in mammals, although rigid evidence has not been known. At present, direct comparison of Ca^{2+} transients between the ICC-IM-like cells and SMCs in our organoid was technically very difficult since these two cells were both thin and tightly associated each other.

The gut contractile organoid provides useful model and tool for studying phase coordination of oscillatory rhythm

To understand the gut peristaltic movements, which reiterate at specific sites along the gut axis (Shikaya et al., 2022), deciphering the mechanisms underlying the coordination/synchronization of oscillators among multiple cells is critical. Exploiting the finding obtained in the current study that multiple organoids easily fuse each other in vitro, we have found that two organoids with different oscillatory rhythm eventually coordinate their phases upon the fusion (Fig. 8). This suggests the ability of ICCs to adjust their rhythm to their neighbors. It is tempting to speculate that during "pausing time" of oscillation upon organoidal fusion (Fig. 8B), they might communicate each other to adjust to a unified rhythm.

During identification of the rhythm-adjustment mediators using the 3-well hydrogel (without connecting channels), SMCs that were unexpectedly crawled out from

the organoids and covered the top surface of the hydrogel were able to mediate the rhythm coordination among organoids (Fig. 9). Whether cellular thin protrusions connected with each other observed in the channel-connected 3-well hydrogel mediate the coordination remains undetermined. Nevertheless, our findings have revealed a novel role of SMCs in mediating rhythm coordination, and as discussed above, these support the notion of the SMC-to-ICC signaling, which is unprecedented. It is unlikely that gap junction plays a major role in such signaling since gap junction inhibitors yielded no detectable effects in SMC-mediated organoidal phase synchronization (Fig. 9H).

In summary, the gut contractile organoid developed in this study serves as a powerful model to study the establishment and maintenance of oscillatory rhythm (pace-making) and their coordination in the multicellular systems.

Materials and Methods
Chicken embryos

Fertilized chicken eggs were obtained from the Yamagishi poultry farms (Wakayama, Japan) and Takeuchi Farm (Nara, Japan). Embryos were staged according to embryonic days. All animal experiments were conducted with the ethical approval of Kyoto University (#202110).

Culture preparation of hindgut-derived cells

A hindgut was dissected from E15 chicken embryos (Fig. S1A), and cut into small pieces. After treating with 25 U dispase (Fujifilm Wako, 383-02281) /phosphate buffer saline (PBS: 0.14 M NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4-$12H_2O$, 1.8 mM KH_2PO_4) at 38.5 °C for 40 min, serosa and intestinal epithelium were removed using forceps. The muscle layer was minced into smaller pieces and treated with 0.2 mg/ml collagenase (Fujifilm Wako, 038-22361) and 0.25 % trypsin/PBS at 37.0 °C for 30 min. The reaction was stopped with 1 % FBS/PBS followed by centrifugation at 800 rpm for 3 min. The pellet was resuspended and washed in PBS followed by centrifugation. They were suspended in culture medium, and 5.0×10^5 cells were plated on 14-mm diameter glass-bottom dishes (Matsunami, D11130H) which had been treated with undiluted Matrigel (Corning, 354248) at 38.5°C for 20 min. Poly-lysine and collagen-coated dishes were purchased (Matsunami, D11131H, D11134H). D-MEM /Ham's F-12 (Wako, 048-29785), Ham's F-12 (Wako, 087-08335) and Neurobasal medium (Gibco, 21103-049) with $1\times$ B-27 supplement (Gibco, 17504044) and 0.5 mM L-glutamine were tested. After seeding, time-lapse images were obtained with CM20 (Evident) under 5% CO_2 and

478   38.5°C with 2-hour intervals.

Assessment of contraction in the spheroid/cluster and organoids

Time-lapse images were obtained using confocal microscopy (Nikon, A1R) under
5% CO_2 and 38.5°C. Region of interest (ROI) was set using the Time Measurement
function in Nikon NIS Elements, and changes were detected in the mode "stDev Intensity"
and exported as individual CSV files. Bright-field changes were normalized to values
between 0 and 1. For drug experiments, the values in the control group were used for
normalization.

Using MATLAB (MathWorks), the changes were plotted, and peaks with a
minimum prominence of 0.25 and a maximum width of 10 seconds were detected as
contractions of spheroid/cluster or organoid using the findpeaks function. To determine
the contraction frequency, we analyzed intervals between contractions and displayed as
box plots.

Immunocytochemistry

An organoid was fixed in acetic acid/ethanol (1:5) for 10 min at room
temperature (RT), and washed in PBS for 10 min at RT. The specimens were
permeabilized in 0.1% Tween-20 in PBS for 10 min at RT, followed by washing in PBS
for 10 min at RT. After blocking with 1 % blocking reagent for 1 hour at RT, specimens
were incubated overnight at 4 ℃ with dilution of 1:300 anti-c-Kit (Yagasaki et al., 2022),
1:300 Tuj-1 (RSD, MAB1195), 1:400 anti-αSMA antibody (Sigma-Aldrich, A5228),
anti-Desmin antibody (Novus Biologicals, NBP1-97707) and/or 1:200 anti-N-cadherin
antibodies (TAKARA, M110) in 1% blocking regent (Roche, 1096176)/PBS. Following
three times washing in PBS for 10 min each at RT, specimens were incubated for 1.5 hr
at RT with 1:500 anti-rabbit IgG(H+L)-Alexa 488-conjugated antibody (Donkey;
Invitrogen, A21206), anti-mouse IgG_{2a}-Alexa 568-conjugated antibody (Goat; Invitrogen,
A21134), anti-rat IgG (H+L)-Alexa 488-conjugated antibody (Goat; Invitrogen, A11006)
and 1:2000 DAPI. After washing three times in PBS for 10 min at RT, fluorescent images
were obtained using the Nikon A1R confocal microscope.

Plasmids

pAAV-hSynapsin1-GCaMP6s-P2A-mRuby3 was purchased from Addgene
(112007). Full-length cDNA of GCaMP6s was PCR-amplified:
forward 5'- GCGTACCACTGTGGCATCGATGCCACCATGGGTTCTCA -3',
reverse 5'- GCCCGTACATCGCATCGATTTACTTGTACAGCTCGT -3'.

The retroviral vector RCAS-EGFP was digested with Cla I to remove EGFP, into which
a DNA fragment was inserted by In-Fusion HD Cloning Kit (TAKARA) to produce
RCAS-GCaMP6s-P2A-mRuby3. RCAS-GapEGFP is as previously described (Murai et
al., 2015)
518

Preparation of retroviral vector particles
RCAS-GCaMP6s-P2A-mRuby3 was transfected into the chicken fibroblast line
cells DF1 using Lipofectamin 2000 (Invitrogen). Transfected cells were cultured in 10
cm culture dish until confluent. The supernatant of transfected DF1 was collected for viral
precipitation, from which retroviral particles were prepared using Retro-Concentin^{TM}
Virus Precipitation Solution (SBI, RV100A-1).
525

Intracellular Ca^{2+} imaging in the gut contractile organoid
On day 2 or 3 of cell culture, a 10 mg/ml polybrene solution (final concentration:
4 µg/ml; Nacalai, 12996-81) and 20 µl of Opti-MEM containing the aforementioned viral
particles were added to the culture medium to transfect GCaMP6s into organoid-forming
cells. The culture medium was replaced with fresh medium on day 5, and Ca^{2+} imaging
was performed on day 7. Time-lapse images were acquired using a confocal microscope
(Nikon, A1R) at intervals of either 700 or 450 msec. Fluorescence intensity from each
region of interest (ROI) was exported into an Excel file using the time-measurement
function of the NIS-Elements software (Nikon). Fluorescence intensity traces were
plotted using the Excel data, with the start of measurement set as time zero. For peak
latency analysis, calcium transients in ICCs and SMCs were identified using the
"findpeaks" function in MATLAB (MathWorks). Temporal differences between peak
signals in ICC and SMC were calculated to show the latency.
539

Drug administration
Carbenoxolone (CBX; nacalai tesque, 32775-51) /H_2O, 18beta-Glycyrrhetinic
acid (18β-GA; abcam, ab142579)/DMSO, (-)-Blebbistatin (FUJIFILM Wako, 021-
17041) and Nifedipine (FUJIFILM Wako, 141-05783) were prepared. Time-lapse images
were acquired for a single organoid before and after the drug administration. Following
the drug adding, organoids were allowed to rest for 30 min to avoid possible effect of
turbulence of the medium, and time-lapse images were taken for 5 min.
547

Three-well hydrogel fabrication
The photoinitiator P2CK was synthesized as previously reported (Li et al., 2013).

The target product was verified using proton NMR and Bruker's TopSpin software.
Gelatin-Norbornene was synthesized based on previous report (Van Hoorick et al., 2018).
The final reaction mixture was transferred to a dialysis tube (5-6 kDa; Spectra Por, cat.
no. 132680T) for dialysis against pure water at 40°C for 3 days. After dialysis, the
solution's pH was carefully adjusted to 8.0.

A piece of 3-well hydrogel was fabricated on a glass bottom dish (Matsunami)
from 100 μL of a solution containing 20 wt % Gelatin-Norbornene, 2 mM of the
photoinitiator P2CK, and 20 mM of crosslinker (DTT) (TCI, #D1071) using a two-photon
microscope with controllable laser power in 3D space according to the voxel file input
(Olympus, in house customized). The voxel files specifying the 3D shape of the hydrogel
were designed with Fusion 360 software. (Autodisk). Organoids were transferred to the
hydrogel using a glass capillary.

563   Culture of embryonic heart-derived cells

A heart was dissected from E10 chicken embryo, and cut into small pieces. The
tissue fragments were treated with 0.125% trypsin in PBS at 37.0 °C for 30 min. The
reaction was terminated by adding 10 % FBS in D-MEM/Ham's F-12, followed by
centrifugation at 800 rpm for 3 min. Pellet was resuspended in D-MEM/Ham's F-12
supplemented with 10% FBS, and the cells were plated in poly-L-lysine-coated glass-
bottom dishes (Matsunami).

On day 1, time-lapse images were acquired using confocal microscopy under 5%
CO_2 at 38.5 °C, with 700 msec intervals. Cell contractions were detected using the "stDev
Intensity" mode in the Time Measurement function of Nikon NIS-Elements.

575   <u>Acknowledgement</u>

We thank Dr. Scott Gilbert for careful reading of the manuscript and discussion.
We also thank National BioResource Project (Chicken-Quail, Nagoya University) for
their technical help. This work was supported by JSPS KAKENHI Grant Numbers;
23H04933, 20H03259, 20K20520, 20K21425 for Y. T., and 21K06198, 23H04702 for
M. I., and FY 2022 Kusunoki 125 of Kyoto University 125[th] Anniversary Fund for M. I.,
and Ginpu Funds (Kyoto University) and incu-be fund (Leave a Nest Co., Ltd.) for R. Y..
R. Y. is an ex-fellow of JSPS.

585   <u>References</u>

586   Baker, S. A., Leigh, W. A., Del Valle, G., De Yturriaga, I. F., Ward, S. M., Cobine, C. A., Drumm,
587   B. T., & Sanders, K. M. (2021). Ca(2+) signaling driving pacemaker activity in
588   submucosal interstitial cells of Cajal in the murine colon. *Elife*, *10*.
589   https://doi.org/10.7554/eLife.64099

590   Barajas-Lopez, C., Berezin, I., Daniel, E. E., & Huizinga, J. D. (1989). Pacemaker activity recorded
591   in interstitial cells of Cajal of the gastrointestinal tract. *Am J Physiol*, *257*(4 Pt 1), C830-
592   835. https://doi.org/10.1152/ajpcell.1989.257.4.C830

593   Camborova, P., Hubka, P., Sulkova, I., & Hulin, I. (2003). The pacemaker activity of interstitial
594   cells of Cajal and gastric electrical activity. *Physiol Res*, *52*(3), 275-284.
595   https://www.ncbi.nlm.nih.gov/pubmed/12790758

596   Chevalier, N. R. (2018). The first digestive movements in the embryo are mediated by
597   mechanosensitive smooth muscle calcium waves. *Philos Trans R Soc Lond B Biol Sci*,
598   *373*(1759). https://doi.org/10.1098/rstb.2017.0322

599   Chevalier, N. R., Ammouche, Y., Gomis, A., Teyssaire, C., de Santa Barbara, P., & Faure, S. (2020).
600   Shifting into high gear: how interstitial cells of Cajal change the motility pattern of the
601   developing intestine. *Am J Physiol Gastrointest Liver Physiol*, *319*(4), G519-g528.
602   https://doi.org/10.1152/ajpgi.00112.2020

603   Chevalier, N. R., Dacher, N., Jacques, C., Langlois, L., Guedj, C., & Faklaris, O. (2019).
604   Embryogenesis of the peristaltic reflex. *J Physiol*, *597*(10), 2785-2801.
605   https://doi.org/10.1113/jp277746

606   Cousins, H. M., Edwards, F. R., Hickey, H., Hill, C. E., & Hirst, G. D. (2003). Electrical coupling
607   between the myenteric interstitial cells of Cajal and adjacent muscle layers in the guinea-
608   pig gastric antrum. *J Physiol*, *550*(Pt 3), 829-844.
609   https://doi.org/10.1113/jphysiol.2003.042176

610   Daniel, E. E., & Wang, Y. F. (1999). Gap junctions in intestinal smooth muscle and interstitial
611   cells of Cajal. *Microsc Res Tech*, *47*(5), 309-320. https://doi.org/10.1002/(SICI)1097-
612   0029(19991201)47:5<309::AID-JEMT2>3.0.CO;2-K

613   Duband, J. L., Gimona, M., Scatena, M., Sartore, S., & Small, J. V. (1993). Calponin and SM 22
614   as differentiation markers of smooth muscle: spatiotemporal distribution during avian
615   embryonic development. *Differentiation*, *55*(1), 1-11. https://doi.org/10.1111/j.1432-
616   0436.1993.tb00027.x

617   Fukui, H., Chow, R. W., Xie, J., Foo, Y. Y., Yap, C. H., Minc, N., Mochizuki, N., & Vermot, J.
618   (2021). Bioelectric signaling and the control of cardiac cell identity in response to
619   mechanical forces. *Science*, *374*(6565), 351-354.
620   https://doi.org/10.1126/science.abc6229

621   Gabella, G., & Blundell, D. (1981). Gap junctions of the muscles of the small and large intestine.

Cell Tissue Res, 219(3), 469-488. https://doi.org/10.1007/BF00209987

Gherghiceanu, M., & Popescu, L. M. (2005). Interstitial Cajal-like cells (ICLC) in human resting mammary gland stroma. Transmission electron microscope (TEM) identification. *J Cell Mol Med, 9*(4), 893-910. https://doi.org/10.1111/j.1582-4934.2005.tb00387.x

Graham, H. K., Maina, I., Goldstein, A. M., & Nagy, N. (2017). Intestinal smooth muscle is required for patterning the enteric nervous system. *J Anat, 230*(4), 567-574. https://doi.org/10.1111/joa.12583

Grosse, A. S., Pressprich, M. F., Curley, L. B., Hamilton, K. L., Margolis, B., Hildebrand, J. D., & Gumucio, D. L. (2011). Cell dynamics in fetal intestinal epithelium: implications for intestinal growth and morphogenesis. *Development, 138*(20), 4423-4432. https://doi.org/10.1242/dev.065789

Huizinga, J. D., Martz, S., Gil, V., Wang, X. Y., Jimenez, M., & Parsons, S. (2011). Two independent networks of interstitial cells of cajal work cooperatively with the enteric nervous system to create colonic motor patterns. *Front Neurosci, 5*, 93. https://doi.org/10.3389/fnins.2011.00093

Huizinga, J. D., Thuneberg, L., Kluppel, M., Malysz, J., Mikkelsen, H. B., & Bernstein, A. (1995). W/kit gene required for interstitial cells of Cajal and for intestinal pacemaker activity. *Nature, 373*(6512), 347-349. https://doi.org/10.1038/373347a0

Iino, S., & Horiguchi, K. (2006). Interstitial cells of cajal are involved in neurotransmission in the gastrointestinal tract. *Acta Histochem Cytochem, 39*(6), 145-153. https://doi.org/10.1267/ahc.06023

Iino, S., Horiguchi, S., Horiguchi, K., & Nojyo, Y. (2007). Interstitial cells of Cajal in the gastrointestinal musculature of W mutant mice. *Arch Histol Cytol, 70*(3), 163-173. https://doi.org/10.1679/aohc.70.163

Kito, Y., Ward, S. M., & Sanders, K. M. (2005). Pacemaker potentials generated by interstitial cells of Cajal in the murine intestine. *Am J Physiol Cell Physiol, 288*(3), C710-720. https://doi.org/10.1152/ajpcell.00361.2004

Kluppel, M., Huizinga, J. D., Malysz, J., & Bernstein, A. (1998). Developmental origin and Kit-dependent development of the interstitial cells of cajal in the mammalian small intestine. *Dev Dyn, 211*(1), 60-71. https://doi.org/10.1002/(SICI)1097-0177(199801)211:1<60::AID-AJA6>3.0.CO;2-5

Komuro, T., Tokui, K., & Zhou, D. S. (1996). Identification of the interstitial cells of Cajal. *Histol Histopathol, 11*(3), 769-786. https://www.ncbi.nlm.nih.gov/pubmed/8839765

Li, Z. Q., Torgersen, J., Ajami, A., Mühleder, S., Qin, X. H., Husinsky, W., Holnthoner, W., Ovsianikov, A., Stampfl, J., & Liska, R. (2013). Initiation efficiency and cytotoxicity of novel water-soluble two-photon photoinitiators for direct 3D microfabrication of

hydrogels. *Rsc Advances*, *3*(36), 15939-15946. https://doi.org/10.1039/c3ra42918k

Liu, L. W., Thuneberg, L., & Huizinga, J. D. (1998). Development of pacemaker activity and

interstitial cells of Cajal in the neonatal mouse small intestine. *Dev Dyn*, *213*(3), 271-282.

https://doi.org/10.1002/(SICI)1097-0177(199811)213:3<271::AID-AJA4>3.0.CO;2-R

Mei, F., Han, J., Huang, Y., Jiang, Z. Y., Xiong, C. J., & Zhou, D. S. (2009). Plasticity of interstitial

cells of cajal: a study in the small intestine of adult Guinea pigs. *Anat Rec (Hoboken)*,

292(7), 985-993. https://doi.org/10.1002/ar.20928

Murai, H., Tadokoro, R., Sakai, K., & Takahashi, Y. (2015). In ovo gene manipulation of

melanocytes and their adjacent keratinocytes during skin pigmentation of chicken

embryos. *Dev Growth Differ*, *57*(3), 232-241. https://doi.org/10.1111/dgd.12201

Rohr, S., Kucera, J. P., & Kleber, A. G. (1998). Slow conduction in cardiac tissue, I: effects of a

reduction of excitability versus a reduction of electrical coupling on microconduction. *Circ*

Res, *83*(8), 781-794. https://doi.org/10.1161/01.res.83.8.781

Rumessen, J. J., & Thuneberg, L. (1991). Interstitial cells of Cajal in human small intestine.

Ultrastructural identification and organization between the main smooth muscle layers.

Gastroenterology, *100*(5 Pt 1), 1417-1431.

https://www.ncbi.nlm.nih.gov/pubmed/2013387

Rumessen, J. J., & Thuneberg, L. (1996). Pacemaker cells in the gastrointestinal tract: interstitial

cells of Cajal. *Scand J Gastroenterol Suppl*, *216*, 82-94.

https://doi.org/10.3109/00365529609094564

Sanders, K. M., Drumm, B. T., Cobine, C. A., & Baker, S. A. (2024). Ca(2+) dynamics in

interstitial cells: foundational mechanisms for the motor patterns in the gastrointestinal

tract. *Physiol Rev*, *104*(1), 329-398. https://doi.org/10.1152/physrev.00036.2022

Sanders, K. M., Kito, Y., Hwang, S. J., & Ward, S. M. (2016). Regulation of Gastrointestinal

Smooth Muscle Function by Interstitial Cells. *Physiology (Bethesda)*, *31*(5), 316-326.

https://doi.org/10.1152/physiol.00006.2016

Sanders, K. M., Publicover, N. G., & Ward, S. M. (1991). Involvement of interstitial cells of Cajal

in pacemaker activity of canine colon. *J Smooth Muscle Res*, *27*(1), 1-11.

https://doi.org/10.1540/jsmr.27.1

Sanders, K. M., Ward, S. M., & Koh, S. D. (2014). Interstitial cells: regulators of smooth muscle

function. *Physiol Rev*, *94*(3), 859-907. https://doi.org/10.1152/physrev.00037.2013

Schultz, T., Daniel, V., & Daniel, E. E. (2003). Does ICC pacing require functional gap junctions

between ICC and smooth muscle in mouse intestine? *Neurogastroenterol Motil*, *15*(2),

129-138. https://doi.org/10.1046/j.1365-2982.2003.00401.x

Shikaya, Y., Takase, Y., Tadokoro, R., Nakamura, R., Inaba, M., & Takahashi, Y. (2022).

Distribution Map of Peristaltic Waves in the Chicken Embryonic Gut Reveals Importance

of Enteric Nervous System and Inter-Region Cross Talks Along the Gut Axis. *Front Cell*
Dev Biol, *10*, 827079. https://doi.org/10.3389/fcell.2022.827079

Sun, X., Yu, B., Xu, L., Dong, W., & Luo, H. (2006). Interstitial cells of Cajal in the murine
gallbladder. *Scand J Gastroenterol*, *41*(10), 1218-1226.
https://doi.org/10.1080/00365520600708800

Takaki, M. (2003). Gut pacemaker cells: the interstitial cells of Cajal (ICC). *J Smooth Muscle Res*,
39(5), 137-161. https://doi.org/10.1540/jsmr.39.137

Takayama, I., Horiguchi, K., Daigo, Y., Mine, T., Fujino, M. A., & Ohno, S. (2002). The interstitial
cells of Cajal and a gastroenteric pacemaker system. *Arch Histol Cytol*, *65*(1), 1-26.
https://doi.org/10.1679/aohc.65.1

Takeda, Y., Ward, S. M., Sanders, K. M., & Koh, S. D. (2005). Effects of the gap junction blocker
glycyrrhetinic acid on gastrointestinal smooth muscle cells. *Am J Physiol Gastrointest*
Liver Physiol, *288*(4), G832-841. https://doi.org/10.1152/ajpgi.00389.2004

Thomsen, L., Robinson, T. L., Lee, J. C., Farraway, L. A., Hughes, M. J., Andrews, D. W., &
Huizinga, J. D. (1998). Interstitial cells of Cajal generate a rhythmic pacemaker current.
Nat Med, *4*(7), 848-851. https://doi.org/10.1038/nm0798-848

Torihashi, S., Fujimoto, T., Trost, C., & Nakayama, S. (2002). Calcium oscillation linked to
pacemaking of interstitial cells of Cajal: requirement of calcium influx and localization of
TRP4 in caveolae. *J Biol Chem*, *277*(21), 19191-19197.
https://doi.org/10.1074/jbc.M201728200

Torihashi, S., Nishi, K., Tokutomi, Y., Nishi, T., Ward, S., & Sanders, K. M. (1999). Blockade of
kit signaling induces transdifferentiation of interstitial cells of cajal to a smooth muscle
phenotype. *Gastroenterology*, *117*(1), 140-148. https://doi.org/10.1016/s0016-
5085(99)70560-3

Van Hoorick, J., Gruber, P., Markovic, M., Rollot, M., Graulus, G. J., Vagenende, M., Tromayer,
M., Van Erps, J., Thienpont, H., Martins, J. C., Baudis, S., Ovsianikov, A., Dubruel, P., &
Van Vlierberghe, S. (2018). Highly Reactive Thiol-Norbornene Photo-Click Hydrogels:
Toward Improved Processability. *Macromol Rapid Commun*, *39*(14), e1800181.
https://doi.org/10.1002/marc.201800181

Wang, Q., Wang, K., Solorzano-Vargas, R. S., Lin, P. Y., Walthers, C. M., Thomas, A. L., Martín,
M. G., & Dunn, J. C. Y. (2018). Bioengineered intestinal muscularis complexes with long-
term spontaneous and periodic contractions. *PLoS One*, *13*(5), e0195315.
https://doi.org/10.1371/journal.pone.0195315

Yagasaki, R., Shikaya, Y., Kawachi, T., Inaba, M., Takase, Y., & Takahashi, Y. (2022). Newly raised
anti-c-Kit antibody visualizes morphology of interstitial cells of Cajal in the developing
gut of chicken embryos. *Dev Growth Differ*, *64*(8), 446-454.

https://doi.org/10.1111/dgd.12808

Yang, P., Yu, Z., Gandahi, J. A., Bian, X., Wu, L., Liu, Y., Zhang, L., Zhang, Q., & Chen, Q. (2012).

The identification of c-Kit-positive cells in the intestine of chicken. *Poult Sci, 91*(9), 2264-

2269. https://doi.org/10.3382/ps.2011-02076

Zeng, X., Komanome, Y., Kawasaki, T., Inada, K., Jonaitis, J., Pulver, S. R., Kazama, H., & Nose,

A. (2021). An electrically coupled pioneer circuit enables motor development via

proprioceptive feedback in Drosophila embryos. *Curr Biol, 31*(23), 5327-5340 e5325.

https://doi.org/10.1016/j.cub.2021.10.005

    Figure legends

    Figure 1. Culture of muscle layer-derived cells prepared from embryonic hindgut.

    (A) Culture of muscle layer-derived cells prepared from embryonic hindgut with FBS

    free-media and substrates. **(B)** Long-term time-lapse imaging after seeding on Matrigel

    with Neurobasal media. The images show the ability of these cells to self-assemble at 0,

    20, 40, 60, 80 and 100 hours taken from Movie 1. Scale bars: 100 μm (A, B).

Figure 2. Spheroids formed in Neurobasal medium and Matrigel exhibited reiterated contractions.

(A) Clusters/spheroids at days 3, 5, and 7 exhibited reiterative contractions. Graphs show normalized contraction intensities visualized using the Time Measurement function. Arrowheads indicate contraction peaks defined by a peak prominence > 0.25 and a peak width ≤ 10 seconds. **(B)** Contraction intervals in clusters/spheroids from day 3 to day 7. Each dot represents a single contraction interval. Median values: day 3, 13.3; day 5, 15.4; day 7, 19.6. Sample sizes: day 3, $n = 35$, peak count = 646; day 5, $n = 48$, peak count = 748; day 7, $n = 21$, peak count = 250. Scale bar: 50 μm (A).

Figure 3. Clusters/spheroids are composed of internally located ICCs and peripherally located SMCs.

(A) Co-staining with anti-c-Kit and anti-αSMA antibodies. White arrowheads indicate co-expression of c-Kit and αSMA at day 3. Yellow arrowheads indicate cells expressing αSMA but not c-Kit at day 7. A schematic diagram illustrates the spatial arrangement of

cells within a day 7 spheroid (green: ICCs; magenta: SMCs). **(B)** Staining with anti-

Desmin antibody, a marker for mature smooth muscle cells. **(C)** Cell morphology within

the spheroid at day 6 visualized by RCAS-gapEGFP expression. **(D)** Staining of day 5

spheroids with anti-N-cadherin antibody. A white arrowhead indicates N-cadherin–

positive cells, while a yellow arrowhead indicates N-cadherin–negative cells in the outer

region of the spheroid. Scale bars: 30 µm (A–D), 10 µm (inset a in C).

Figure 4. Ca²⁺ imaging of the gut contractile organoid revealed intercellular synchronization.

(**A**) RCAS-GCaMP6s-P2A-mRuby3 plasmid. (**B**) Ca^{2+} imaging of gut contractile organoid during relaxation and contraction. Ca^{2+} dynamics (green) and normalized values of contraction (grey) of gut contractile organoid. (**C**) Simultaneous measurement of intercellular Ca^{2+} dynamics between ICC-ICC, SMC-SMC or ICC-SMC. Three or two ROIs in Ca^{2+} signal-positive cells were set in a single organoid. Graphs show Ca^{2+} dynamics in the ROIs. Magnified view shows that a peak of Ca^{2+} signal in ICC (green) preceded that in SMC (magenta). (**D**) Peak latency in ICC/SMC. Median value = 0.7; Sample size: $n = 14$, peak count = 117. Scale bar: 50 μm (B).

Figure 5. Gap junction inhibitor exerted limited effects on the synchronization of Ca²⁺ dynamics.

Ca²⁺ synchronization among two or three ROIs in GCaMP6s-expressing organoids was evaluated before and after treatment with 100 µM CBX. The synchronization was unaffected between **(A)** SMC-SMC and **(B)** ICC-ICC, but was partially affected between **(C)** ICC–SMC. Magnified views **(a, b)** highlight that the preceding Ca²⁺ peak in ICC (green) observed before CBX treatment (a) was abolished after treatment (b). **(D)** Peak latency between ICC and SMC. Median values: Before, 1.167; CBX 20 µM, 0.701; CBX

100 µM, 0.0. Statistical significance was assessed using Welch's t-test. ***$p < 0.001$, n.s.,

$p \geq 0.05$ ($p = 0.27$ for Before vs. CBX 20 µM). Sample sizes: Before: $n = 10$, peak

count $= 50$; CBX 20 µM: $n = 5$, peak count $= 34$; 100 µM: $n = 5$, peak count $= 51$, Scale

bar: 50 µm (A).

Figure 6. The organoidal contraction is important for Ca^{2+} dynamics in ICCs.

GCaMP6s-expressing organoids were cultured with Blebbistatin. **(A)** Organoidal

contractions were extinguished at 10 μM. **(B)** Contraction intervals before and after

Blebbistatin, and upon washout. Median values: Before, 33.46; Blebbistatin 5 μM,

12.86; 10 μM, N/A; Washout, 23.95. Statistical significance was assessed using

Welch's t-test. ****p < 0.001*, **p < 0.05*. Sample size: *n* = 3. **(C-E)** Comparison of Ca^{2+}

dynamics in SMC-SMC, ICC-ICC, and ICC-SMC. Three or two ROIs were assessed

before and after administrations of 10 μM Blebbistatin. **(F)** Ca^{2+} transients in a single

ICC at 0 μM (before), 5 μM, and 10 μM. Scale bar: 50 μm (B).

Figure 7. Nifedipine ceased organoidal contractions and Ca²⁺ activities in ICCs.

Similar experiment to Fig. 6. **(A)** Organoidal contractions were extinguished at 1 µM. **(B)**

Contraction intervals before and after administration, and upon washout. Median values:

Before, 25.7; Nifedipine 1 µM, N/A; Washout, 22.2. Statistical significance was assessed

using Welch's t-test. ******$p < 0.01$. Sample size: $n = 4$. Peak counts: Before, 101; Nifedipine

1 µM, N/A; Washout, 46. **(C)** Ca²⁺ transients in a single ICC before and after

administration of 1 µM nifedipine, and following washout. Scale bar: 50 µm (B).

Figure 8. Ca²⁺ transients in multiple organoids undergo synchronization upon organoidal fusion.

(A) Time-lapse imaging of organoidal fusion. **(B)** When two organoids that originally displayed independent Ca²⁺ rhythm fused to each other, their rhythm became synchronized after fusion (24 h). **(C)** Contraction intervals before and after fusion. Median values: before fusion, 13.65; after fusion, 21.04. Statistical significance was assessed using Welch's t-test. ***$p < 0.001$. Sample size: $n = 3$ pairs. Peak counts: before, 138; after, 90. **(D)** Cellular protrusions between two neighboring organoids. White arrowheads show three protrusions from the left organoid. Scale bars: 100 μm (A, B, D), 20 μm (inset (a) in (D)).

Figure 9. Smooth muscle cells mediate Ca²⁺ synchronization between organoids.
(A) Diagram of a 3-well hydrogel in which one organoid was placed per well. The three
wells were connected with narrow channels, and this gel mold does not allow organoidal
bodies to fuse to each other, but allow them to extend/migrate protrusions/cells through
the channel **(B)**. **(C)** After 3 days, the three organoids displayed synchronization of Ca²⁺
dynamics. **(D)** Some organoid-derived cells crawled out from the wells, and covered the
top surface of the hydrogel, resulting in bridging the three unfused organoids. **(E)**

Diagram of a 3-well hydrogel *without* channels. **(F)** The surface-covering cells were
identified as SMCs (αSMA-positive, c-Kit-negative). **(G)** In the hydrogel without
channels but with surface-covered SMCs, Ca^{2+} dynamics in the three organoids were
synchronized. **(H)** The Ca^{2+} synchronization shown in (G) was not altered by treatment
with 18β-GA. Blue planes indicate focal planes. Scale bars: 50 μm (B); 100 μm (C, D, F,
G, H).
844

Supplementary Figure S1

(A) Chicken embryonic hindgut at E15. It was dissected from the bottom of the cecum to
the front of the cloaca (white lines). (B) Three layers of E15 hindgut: serosa, muscle layer,
intestinal epithelium. Remak's ganglion was also removed. Scale bars: 1 mm in A, B

Supplementary Figure S2

(A) Co-staining of Day 7 organoids with anti-c-Kit- and Tuj1 antibodies. White arrowhead shows a Tuj1$^+$ cell. **(B)** Representation of Tuj1-positive, αSMA-single positive, c-Kit-single positive cells in an organoid at day 7. Median values: Tuj1$^+$, 0.92; αSMA$^+$, 40; c-Kit$^+$, 53. Sample size: Tuj1$^+$, $n = 12$; αSMA$^+$, c-Kit$^+$, $n = 3$ **(C)** Contraction intervals before and after administration of TTX. Median values: Before, 23.4; TTX, 25.35. Sample size: $n = 9$. Peak counts: Before, 166; TTX, 148. Statistical significance was assessed using Welch's t-test. *n.s., $p \geq 0.05$* ($p = 0.11$). Scale bar: 30 μm (A).

 Supplementary Figure S3

Effects of gap junction inhibitors on organoidal contractions and ICC-SMC latency. **(A)**

Organoids at day 7 exhibited rhythmic contractions before and after administration of

100 μM CBX. Graphs show normalized contraction intensities visualized using the Time

Measurement function. Arrowheads indicate contraction peaks defined by a peak

prominence > 0.25 and a peak width ≤ 10 seconds. **(B)** Contraction intervals before and

after treatment with CBX (100 μM), H_2O (vehicle control for CBX), 18β-GA (100 μM),

and DMSO (vehicle control for 18β-GA). Median values: CBX, 14.0 (before), 11.9

(after); H_2O, 14.7 (before), 15.4 (after); 18β-GA, 14.7 (before), 7.7 (after); DMSO, 19.95

(before), 17.5 (after). Sample sizes and peak counts: CBX, $n = 9$ (before), $n = 9$ (after),

peak count $= 150$ (before), 196 (after); H_2O, $n = 10$ (before), $n = 10$ (after), peak

count $= 173$ (before), 171 (after); 18β-GA, $n = 7$ (before), $n = 8$ (after), peak count $= 129$;

DMSO, $n = 7$ (before), $n = 6$ (after), peak count $= 193$ (before), 92 (after). **(C)**

Contraction intervals at day 7 in organoids cultured with CBX, H_2O, 18β-GA, and DMSO

from day 3 to day 7. Median values: H_2O, 22.75; CBX, 32.5; DMSO, 18.85; 18β-GA,

27.95. Sample sizes and peak counts: H_2O, $n = 9$, peak count $= 136$; CBX, $n = 11$, peak

count $= 135$; DMSO, $n = 9$, peak count $= 209$; 18β-GA, $n = 18$, peak count $= 268$. **(D)**

Peak latency of Ca^{2+} transients between ICC and SMC before and after administration of

18β-GA (20 μM, 100 μM). Statistical significance was assessed using Welch's t-test. n.s.,

$p = 0.27$ for Before vs. 18β-GA 20 μM; $p = 0.25$ for Before vs. 18β-GA 100 μM; $p = 0.46$

for 18β-GA 20 μM vs. 100 μM. Sample sizes and peak counts: Before, $n = 5$, peak

count $= 62$; 18β-GA 20 μM, $n = 3$, peak count $= 52$; 18β-GA 100 μM, $n = 2$, peak

count $= 57$. **(E)** Contractions of cultured embryonic heart cells. Graphs show normalized

contraction intensities using the Time Measurement function. The box plot shows the

number of contractions per 5 minutes. Median values: CBX, 30.75 (before), 15(CBX

100μM), 55.75 (washout); 18β-GA: 45.5 (before), 8.5 (18β-GA 100μM), 51 (washout).

Statistical significance was assessed using Welch's t-test. **$p < 0.01$, *$p < 0.05$; n.s., p

≥ 0.05 ($p = 0.13$ for Before vs. washout of CBX; $p = 0.26$ for Before vs. washout of 18β-

GA). Sample sizes: CBX, $n = 4$; 18β-GA, $n = 4$. Scale bar: 50 μm (A).

Movie 1

Long-term time-lapse imaging after seeding. Time lapse was taken every 2 hours for a total of 100 hours. This video corresponds to Fig. 1B.

Movie 2

Periodic contractions of the cluster/spheroid on day 3, 5 and 7. Time-lapse images were taken with 700 msec intervals for 5 min. This video corresponds to Fig. 2A.

Movie 3

Periodic contractions of a day 7 organoid with CBX. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. S3.

Movie 4

Ca^{2+} dynamics in day 7 gut contractile organoid are concomitant with its contractions. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 4.

Movie 5

Ca^{2+} dynamics in a gut contractile organoid with CBX. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 5.

Movie 6

Ca^{2+} dynamics in day 7 gut contractile organoid with Blebbistatin. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 6.

Movie 7

Live imaging during fusion of multiple organoids. Time-lapse images were obtained every 10 minutes for 4 hours. This video corresponds to Fig. 8A.

Movie 8

Ca^{2+} transients in two gut contractile organoids before and after fusion. Time-lapse images were obtained with 450 msec intervals for 5 min. This video corresponds to Fig. 8B.

Movie 9

Ca^{2+} transients in three gut contractile organoids in 3-well hydrogel with channels.
Time-lapse images were obtained with 700 msec intervals for 5 min. This video
corresponds to Fig. 9C.

Movie 10

Ca^{2+} transients in three gut contractile organoids in 3-well hydrogel *without*
channels. Time-lapse images were obtained with 450 msec intervals for 5 min. This
video corresponds to Fig. 9G.

Movie 11

Similar assay to movie 10 with 18β-GA administration. Time-lapse images were
obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 9H.