1

The gut contractile organoid: a novel model for studying the gut motility regulated by coordinating signals between interstitial cells of Cajal and smooth muscles

Rei Yagasaki[1,2], Ryo Nakamura[1,3], Yuuki Shikaya[1,4], Ryosuke Tadokoro[1,5], Ruolin Hao[6,7], Zhe Wang[6,7], Mototsugu Eiraku[6,7], Masafumi Inaba[1] and Yoshiko Takahashi[1,8]

1: Department of Zoology, Graduate School of Science, Kyoto University Sakyo-ku, Kyoto 606-8502, Japan

2: Present Address: WPI Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

3: Present Address: Ushimado Marine Institute, Okayama University, Setouchi, Okayama 701-4303, Japan

4: Present Address: Sorbonne Université, CNRS, Institut de la Mer de Villefranche, UMR7009 Laboratoire de Biologie du Développement de Villefranche-sur-Mer (LBDV), Villefranche-sur-Mer, France

5: Present Address: Department of Bioscience, Okayama University of Science, Okayama, Japan

6: Institute for Life and Medical Sciences, Kyoto University, Sakyo-ku, Kyoto 606-8507, Japan

7: Department of Polymer Chemistry, Graduate School of Engineering, Kyoto University, Kyoto 615-8510, Japan

8: Author to whom all correspondence should be addressed.

email: yotayota@develop.zool.kyoto-u.ac.jp

Keywords:
gut contraction, interstitial cells of Cajal, smooth muscle cells, organoid, gap junction

<u>Abstract</u>

The gut undergoes peristaltic movements regulated by intricate cellular
interactions. However, they have poorly been explored due to a lack of model system.
We here developed a novel contractile organoid that is derived from the muscle layer of
chicken embryonic hindgut. The organoid contained smooth muscle cells (SMCs) and
interstitial cells of Cajal (ICCs; pacemaker) with few enteric neurons, and underwent
periodic contractions. The organoid formed by self-organization with morphological
arrangements of ICCs (internal) and SMCs (peripheral), allowing identification of these
cells in live. GCaMP-Ca^{2+} imaging analyses revealed that Ca^{2+} transients between ICC-
ICC, SMC-SMC or SMC-ICC were markedly coordinated. Pharmacological studies
further showed that gap junctions play a role in ICC-to-SMC signaling, and also possible
feedback from SMC's contraction to ICC's pace-making activities. In addition, two
organoids with different rhythm became synchronized when mediated by SMCs,
unveiling a novel contribution of SMCs to ICC's pace-making. The gut contractile
organoid developed in this study offers a useful model to understand the mechanisms
underlying the rhythm coordination between/among ICCs and SMCs during gut
peristaltic movements.

47
48

Introduction

In the gut, ingested material is conveyed properly along the gut axis by the gut movements called peristalsis, which is recognized as wave-like propagation of a local constriction. The physiology of gut peristalsis has extensively been studied in adults, where the peristalsis plays pivotal roles in effective transportation and digestion/absorption of inter-luminal contents. And many cases of gut-related pathology are associated with peristaltic disfunctions. The gut peristaltic movements are achieved by intricate regulations of intercellular functions among multiple cell types. At the site of origin of peristaltic waves (OPW), a local constriction emerges along the circumferential axis, and this is soon followed by a progressive wave of the contraction along the gut axis. During these processes, the circumferential constriction demands multiple SMCs to achieve simultaneous contraction. However, due to the complex structure of the gut, how such synchronization/coordination in SMCs is regulated remains largely undetermined.

It is known in vertebrates that the embryonic gut undergoes peristaltic movements even without experience of food intake. The embryonic gut therefore serves as a powerful model to study the intrinsic mechanisms underlying the peristalsis contrasting with the adult gut where ingested content influences the gut motility increasing complexity in analyses. We have recently reported using chicken embryos that sites of OPW are randomly distributed along the gut axis at early stages, and they later become confined to specific sites, and this confinement of OPWs enables rhythmic and patterned peristaltic movements (Shikaya et al., 2022). One of the long-standing and important questions is how the synchronized/coordinated contraction is achieved.

In a long history of gut motility studies, it has been known that enteric nervous system (ENS), smooth muscle cells (SMCs) and interstitial cells of Cajal (ICCs) play important roles in peristaltic movements (Barajas-Lopez et al., 1989; Camborova et al., 2003; Chevalier et al., 2020; Huizinga et al., 1995; Kito et al., 2005; Liu et al., 1998; Rumessen & Thuneberg, 1996; Sanders et al., 1991; Takaki, 2003; Takayama et al., 2002; Thomsen et al., 1998). It has widely been accepted that: 1) at early embryonic stages, gut movement/peristalsis does not require ENS activity, 2) ICCs serve as a pacemaker dictating their rhythm to SMCs, 3) the contraction is executed by SMC and not by ICCs, since thick fibers of myosin are found solely in SMCs. A series of elaborate electrophysiological studies showed that slow waves (a type of changes in membrane potential characteristic of gut movements) occur intrinsically in ICCs but not in SMCs, and that these slow waves lead to voltage-dependent Ca^{2+} influx evoking action potential, which is somehow transmitted to SMCs to execute gut contraction in register with ICC's pace-making rhythm (Baker et al., 2021; Torihashi et al., 2002). The knowledge that ICCs

act as a pacemaker was supported by compelling evidence obtained by c-Kit-deficient
mouse mutants (W/Wᵛ), in which ICC differentiation was severely affected leading to a
failure of gut peristalsis (Huizinga et al., 1995; Torihashi et al., 1999). However, it
remains largely unexplored how the intrinsic/spontaneous rhythm in a single ICC
becomes synchronized among multiple ICCs which constitute intricate networks in the
gut. It has also been under debate to what extend the gap junction contributes to cell-cell
communications between ICCs, SMCs or ICC-SMC. One reason is that it has been
difficult to stably maintain SMCs and ICCs in cell culture condition, and also to
distinguish ICCs from SMCs in the living gut (these two types of cells originate from the
same progenitors of splanchnopleural mesoderm during development). Thus, a novel
model system has been awaited to circumvent these obstacles. Recently, it was reported
that differentiation states of mouse hindgut-derived cells were successfully maintained
for a relatively long period of time in a serum-free culture medium (Wang et al., 2018).
In that study, many types of gut-derived cells including not only ENS, ICC, SMCs but
also glial cells and serosa were observed, and this large and complex cell mass was seen
to undergo rhythmic contractions *in vitro*.

It has widely been appreciated that organoids can serve as a powerful models
and tools to circumvent such obstacles of organ complexity. Their relatively simple
structures allow analyses at higher resolution than *in vivo*, and also permit analyses of cell
behaviors in three-dimensional (3D) environment, which might be different from
behaviors *in vitro* behaviors confined to two dimensions. In the current study, we have
developed a novel organoid called "gut contractile organoid" by culturing chicken
hindgut-derived cells in a serum-free medium. The gut contractile organoid undergoes
periodic contractions, and it is essentially composed of ICCs and SMCs, the former
residing centrally whereas the latter peripherally, allowing distinction between the two
cell types in living organoids. These advantages enabled GCaMP-live imaging of Ca^{2+}
dynamics and revealed coordinated oscillations of Ca^{2+} transients between ICC-ICC,
SMC-SMC and SMC-ICC. Pharmacological studies further suggested that gap junctions
play a role in an ICC-to-SMC signaling, and also a possible feedback from SMC's
contractions to ICC's pace-making activities. In addition, by regarding a single organoid
as an oscillator unit and by placing oscillators separately in a hydrogel mold, we found
that two oscillators with different rhythm become synchronized when mediated by SMCs,
supporting the notion of SMC' contribution to ICC's pace-making. The gut contractile
organoid developed in this study must be useful to unveil the intrinsic mechanisms
underlying the rhythm coordination between/among ICCs and SMCs during peristaltic
movements in the embryonic gut.

Results

Spheroids were formed from muscle layer-derived cells of the embryonic hindgut

To develop a culture condition that would facilitate analyses of gut motility, we
dissected the muscle layer (also called *tunica muscularis*) from the hindgut of E15
embryos by removing the serosa and intestinal epithelium (mucosa) (Fig. S1). The
isolated muscle layer was dissociated into single cells to prepare 5.0×10^5 cells per culture
dish. We started analyses with a culture condition with FBS-free medium and Matrigel
as substrate as previously described for cultures in mice of gastrointestinal cells including
serosa (Wang et al., 2018). We tested three kinds of FBS-free media: DMEM/Ham's F-
12, Ham's F-12, and Neurobasal media (see Materials and Methods). When cultured in
DMEM/Ham's F-12 or Ham's F-12, dissociated cells formed very small aggregates
containing several cells at day 1, the morphology of which did not change significantly
until day 5 (Fig. 1A). In clear contrast, when cultured in the Neurobasal medium, cells
formed clusters that were interconnected by elongated cells with neighboring clusters as
early as day 1. These clusters grew as larger aggregates by day 3, and became spherical
by day 5. Such spheroids were not observed in the conditions with Ham's F-12 or
DMEM/Ham's F-12. We also tested different substrates, Poly-Lysine or collagen for dish
coating with the Neurobasal medium, but neither one yielded spheroid. To know how the
spherical aggregates were formed under the condition of Neurobasal medium and
Matrigel, we took time-lapse images at two-hour intervals. Originally sparse clusters that
were loosely connected by extended cells merged each other forming progressively larger
clusters (Fig. 1B, Movie 1). In the following experiments, we focused on the spheroids
formed under the condition of Neurobasal medium and Matrigel coating.

The gut muscle layer-derived spheroids displayed periodic contractions

The spheroids underwent reiterative contractions at day 3, and these contractions
were observed at least until day 7 of culture (Fig. 2A, Movie 2). Time-lapse imaging of
the contractions combined with quantitative assessments by ImageJ (see Materials and
Methods) showed that the average frequencies of contractions in each cluster/spheroid
at day 3, day 5, and day 7, were 3.32, 3.04, and 2.97 contractions/min, respectively (Fig.
2B), which largely corresponded to the rhythm of the E16 chicken embryonic intact gut
(Chevalier et al., 2019).

The contracting spheroid was composed of ICCs and SMCs

To determine the cell types comprising the spheroids, we performed
immunostaining with antibodies for the chicken c-Kit protein (for ICCs; Yagasaki et al.,
2022) and αSMA (for SMCs). It is known that ICCs and SMCs are derived from common
progenitors which are c-Kit$^+$/αSMA$^+$, and that after differentiation ICCs and SMCs are c-
Kit$^+$/αSMA$^-$ and c-Kit$^-$/αSMA$^+$, respectively (Duband et al., 1993; Kluppel et al., 1998).
In the clusters at day 3, c-Kit$^+$/αSMA$^-$ signals were detected in the internal region, and c-
Kit$^+$/αSMA$^+$ signals at the periphery (Fig. 3A, Day 3), suggesting that internal cells were
differentiated ICCs, whereas peripheral cells were ICC/SMC progenitors. At day 5
onward, c-Kit$^+$/αSMA$^-$ cells and c-Kit$^-$/αSMA$^+$ cells were segregated which were located
internally and peripherally, respectively (Fig. 3A, Day 5), and this spatial segregation of
the cells remained unchanged until day 7 (Fig. 3A, Day 7), the latest stage examined in
this study. Infection of RCAS-GapEGFP into a forming spheroid further visualized
multipolar ICCs in the internal region (Fig. 3B), known to be characteristic of ICC-MY
which would normally reside in the layer of myenteric plexus (Mei et al., 2009; Sanders
et al., 2014). ICCs of a different type were also recognized, which were elongated, thin,
and lining underneath the peripheral SMCs (Fig. 3A, Day 7). It is possible that they were
ICC-IM, known to be embedded in and tightly associated with smooth muscles in the gut
(Huizinga et al., 2011; Iino & Horiguchi, 2006). Unexpectedly, the spheroid contained
few neural cells (ENS), if any, revealed by anti-Tuj1 antibody (Fig. S2).
Since ICCs and SMCs were spatially segregated in the sphenoid, we stained with
anti N-cadherin antibody (E-cadherin is positive solely in the mucosa/endoderm)
(Graham et al., 2017; Grosse et al., 2011). N-cadherin signals were seen in the internal
ICCs, and not in the peripheral SMCs (Fig. 3C). Thus, it is likely that N-cadherin plays a
role in the segregation.
In summary, the hindgut-derived spheroid displays three prominent
characteristics: (1) a dominant occupation by ICCs and SMCs with neglectable
contribution by ENS, (2) self-organization ability of internal ICCs encapsuled by a thin
layer of SMCs, (3) contractions with periodic rhythm. Based on these characteristics, we
designated this spheroid as "gut contractile organoid".
186

187   Contraction-associated intracellular Ca^{2+} transients were coordinated between
188   ICCs and SMCs in the gut contractile organoid
It has been reported that Ca^{2+} dynamics are important for pacemaker activity in
ICCs and contractions of SMCs. Ca^{2+} flows into ICCs via voltage-dependent Ca^{2+}
channels and propagates to SMCs, causing the gut muscle contractions (Baker et al.,
2021). We therefore investigated the Ca^{2+} dynamics in our gut contractile organoids.

Organoid-forming cells were infected with a RCAS-vector encoding GCaMP6s, a Ca^{2+}
indicator that emits EGFP fluorescence in response to Ca^{2+} influx, and mRuby3 as a
reporter (Fig. 4A). GCaMP6s-organoids were subjected to time-lapse imaging analyses
by confocal microscopy. As expected, the oscillations rhythm of Ca^{2+} transients as a
whole organoid was concomitant with that of contractions (Fig. 4B, Movie 4).
Taking advantage of the spatial segregation between ICCs (internal) and SMCs
(peripheral) (Fig. 3A), we compared the Ca^{2+} oscillation rhythm among ICCs and SMCs
(homotypically) and between ICC-SMC (heterotypically). We set up 3 regions of interests
(ROIs) in the central region, with one ROI corresponding to one cell, and captured the
Ca^{2+} transients. The three ROIs exhibited a synchronous pattern of Ca^{2+} oscillations (Fig.
4C, ICCs). Similarly, Ca^{2+} oscillations in three ROIs in the SMC layer were synchronous
(Fig. 4C, SMCs). These data highlight active communications taking place intercellularly
within ICCs and SMCs, respectively. We further compared Ca^{2+} oscillations between
ICC-SMC. Setting up with one ROI in an ICC, and one in a SMC. Again, the Ca^{2+} rhythm
was synchronized between these heterotypic cell types (Fig. 4C, ICC/SMC). With a closer
look, a rise of the Ca^{2+} transient in the ICC preceded that in SMC with a time lag (also
called latency) of 690 msec in average (Fig. 4C, ICC/SMC), suggestive of a propagation
of Ca^{2+} signals from ICC to SMC consistent with previous reports (Baker et al., 2021).
211

Gap junction plays a role in ICC-to-SMC signaling

Although a series of gut motility studies have proposed an importance of gap
junctions, rigid evidence has been limited due to a lack of experimental model. Our gut
contractile organoid should prove useful for clarifying the roles of gap junctions in the
synchronous motility, since intercellular synchronization was observed between/among
identifiable cells (ICCs and SMCs) as shown above. We performed pharmacological
assessments using gap junction inhibitors including carbenoxolone (CBX) and 18β-
glycyrrhetinic acid (18β-GA; Chevalier, 2018; Takeda et al., 2005), two of the most
widely used pharmacological inhibitors used to study gap junctions.
Following inhibitor administration into day 7 culture medium, organoids were
allowed to rest for 30 min to exclude possible effects by the administration, for example,
turbulence of the medium. With CBX or 18β-GA in the culture medium, the periodic
contraction of an organoid was retained with almost the same frequency as the control
(Fig. S3, Movie 3). We then examined whether the intercellular
synchronization/coordination was affected by the inhibitors. Similarly to Fig. 4, three
ROIs were set either in ICCs or SMCs territories for assessment of Ca^{2+} transients. The
same cells in a single organoid were tracked for their Ca^{2+} transients before and after the

inhibitor administration. Contrary to our expectation, none of 10 μM, 20 μM and 100 μM
concentrations of these inhibitors showed detectable inhibition of the synchronous
patterns of Ca^{2+} transients (Fig. 5A, B show 100 μM of CBX, Movie 5).

In contrast, when an ICC and SMC were compared, the preceding rise of Ca^{2+}
transient in ICC was abolished, although overall synchronization was retained between
these the cells (Fig. 5C). Collectively, while the contribution by gap junction to the
periodic contraction and intercellular synchronization in the organoid is relatively limited,
the ICC-to-SMC signals require gap junction-mediated communications, at least partly.
237

238   Blebbistatin abolished organoidal contractions and oscillatory patterns of Ca^{2+}
239   transients

Toward searching for factors that regulate the coordination between/among ICCs
and SMCs, we tested blebbistatin, a specific inhibitor of myosin II, which was expected
to cease organoidal contractions. Experimental procedures were similar to those for gap
junction inhibitors. We found that blebbistatin ceased periodic contractions of organoids
in a concentration-dependent manner, with 10 μM ceasing it completely (Fig. 6A, Movie
6). We examined Ca^{2+} transients in these contraction-inhibited organoids. Markedly,
periodic Ca^{2+} transients not only in SMCs, but also in ICCs were extinguished, resulting
in no/little synchronous Ca^{2+} patterns among and between ICCs and SMCs (Fig. 6B-E).
Although the possibility is not excluded that blebbistain directly inhibited non-muscle
myosin II in ICCs, these findings imply that the contractility feeds back to ICCs to
generate their periodic rhythm.
251

252   Inter-organoidal coordination was mediated by SMCs

During analyses with our novel organoids, we found that they easily fuse to each
other, suggesting that organoids grow by progressive fusion (Fig. 7A, Movie 7). This also
raised the possibility that the synchronous Ca^{2+} transients among organoid-constituting
cells (Fig. 4C) might be a consequence of phasic coordination upon the fusion of multiple
organoids that had shown different oscillatory phases. To test this possibility, we
transferred two organoids in a petri dish, and allowed them to fuse. Before the fusion,
Ca^{2+} oscillatory rhythm was indeed out of phase in the two organoids (Fig. 7B, Movie 8).
Markedly, upon fusion by 24 hours, their rhythm became in phase/synchronous (Fig. 7B,
Movie 8).

Since we noticed that cellular protrusions were often observed around the time of
organoidal fusion, we reasoned that these cellular processes would mediate the fusion and
its subsequent synchronization of Ca^{2+} transients (Fig. 7C). To test this, we developed a

3-well hydrogel with narrow channels connecting the wells (Fig. 8A). Organoids were
placed separately in each well, which prevented the fusion of organoidal bodies, but
allowed extension of cellular processes through the narrow channels and contact with
each other (Fig. 8B). After 72 hours of placement of organoids, cellular processes as well
as several cell bodies were present within the channel, and Ca^{2+} transients became
synchronized among the three organoids (Fig. 8C, Movie 9). However, with careful
examination, we also noticed that organoid-derived cells crawled out from the wells to
cover the top surface of the hydrogel, connecting the three organoids (Fig. 8D). Thus,
another possibility was raised that these crawled out cells would mediate the inter-
organoidal synchronization.
To test this possibility, we prepared a 3-well hydrogel in which the three wells
were disconnected (no channels) so that organoid-derived cellular processes were not able
to connect each other (Fig. 8E). By 3 days of culture, the top surface of the hydrogel was
indeed covered by cells in a similar way to Fig. 8D. These cells were positive for αSMA
but negative for c-Kit (Fig. 8F), indicating that they were SMCs that were somehow
detached and crawled out from the peripheral layer of the organoids. Importantly,
coincided with the top-coverage by the SMCs, the three organoids in the disconnected
wells displayed synchronized Ca^{2+} transients, highlighting the role of SMCs in mediating
coordination between organoids (Fig. 8G, Movie 10). Gap junction inhibitor yielded
no/little effects on the Ca^{2+} transient coordination (Fig. 8H, Movie 11).
285
286

Discussion
We have developed a novel gut contractile organoid, which displays several
unique characteristics: 1) it undergoes periodic contractions, 2) differentiation states of
ICCs (c-Kit$^+$/αSMA$^-$) and SMCs (c-Kit$^-$/αSMA$^+$) are maintained at least until day 7 in
the organoid, 3) the organoid is composed essentially of two types of cells, ICCs and
SMCs, with few ENS cells, if any, 4) ICCs (internal) and SMCs (peripheral) can be
distinguished for their localization in a living organoid, allowing 5) GCaMP-visualization
of Ca^{2+} transients and assessments of cell interactions between and among ICCs and
SMCs. These characteristics circumvent, at least partly, obstacles that have hampered
analyses in the research of gut peristalsis, such as unstable differentiation state of ICCs
and SMCs in cultures, and difficulties in identifying these cells in living preparations. In
studies of gut movements, how ICCs generate/maintain their periodic rhythm and how
ICCs and SMCs interact with each other have been long-standing questions, and our

organoids offer powerful advantages to address these and to understand the cellular
mechanisms underlying the gut contractions/peristaltic motility.

Contrasting with many cases in organoid studies that have aimed at a *maximum*
recapitulation of the intact organ, our gut contractile organoid is composed of a (probably)
minimum number of cell types that suffice the generation and/or maintenance of rhythmic
contractions, allowing high-resolution analyses at the cellular level. For example, adding
ENS components to our organoid would allow the clarification of the role of ENS in the
gut contraction. While the majority of internal cells in the organoid are ICCs (c-
Kit$^+$/αSMA$^-$), the possibility cannot be excluded that platelet-derived growth factor
receptor-α positive (PDFGRα^+/c-Kit$^-$/αSMA$^-$) cells, another type of interstitial cells
(fibroblast-like cell) known to mediate neural activity to SMCs in the mouse gut (Sanders
et al., 2024; Sanders et al., 2016), are included in our organoid. Available antibody against
the chicken PDFGRα protein is awaited.

314   Coordinated Ca^{2+} transients in ICC/SMC populations in the gut contractile
315   organoid
Measurement and quantification analyses with ICCs and SMCs that are
identifiable in the living organoid revealed exquisite coordination of Ca^{2+}
transients/oscillation homotypically between ICC-ICC and SMC-SMC and
heterotypically between SMC-ICC (Fig. 4C). Notably, a rise (upstroke) of Ca^{2+} transient
in ICCs precedes that of SMCs with a time lag (also called latency) of 690 msec,
suggesting a directed signaling from ICC to SMC. These observations are consistent with
previous studies using ICC- and SMC-specific transgenic mice that expressed GCaMP
and RCaMP, respectively (Baker et al., 2021). In that study, the authors assessed Ca^{2+}
transients in submucosal ICCs (ICC-SM) and compared them with those in their adjacent
circular muscles, and showed that the rise of GCaMP signal preceded that of RCaMP with
a latency of 56 +/- 14 msec. With these observations, they concluded that ICC-SM signals
to its adjacent SMCs. In our study, time-lapse imaging was mostly performed with 700
msec intervals. Further studies with shorter intervals are awaited to know the latency time
would be shorter than 690 msec in average.

331   Contribution of gap junction to ICC-to-SMC signaling
Effects by the block of gap junction by CBX or 18β-GA were relatively limited
in our organoid assays: synchronous patterns of Ca^{2+} transients remained unchanged
between SMC-SMC and ICC-ICC. In contrast, signaling from ICCs to SMCs was
specifically affected by these inhibitors, in which the preceding rise of Ca^{2+} transient in

ICCs was abolished. The contribution of gap junction to the ICC-to-SMC signaling was
previously reported in mouse gut, which was from intramuscular ICCs (ICC-IM) to their
adjacent circular smooth muscles. However, interpretation of the role of gap junction in
ICC-SMC interactions, in general, has been under big debate. Some studies reported that
CBX or 18β-GA failed to inhibit these interactions or peristaltic motilities (Komuro et al.,
1996; Rohr et al., 1998; Schultz et al., 2003), or that electron microscopy did not detect
structures of gap junction in longitudinal muscles (Cousins et al., 2003; Daniel & Wang,
1999; Gabella & Blundell, 1981). Contribution of gap junctions to the gut motility appears
to be highly variable in different regions of gastrointestinal tract, for example, stomach
versus colon (Iino et al., 2007; Yang et al., 2012) . And a detection of gap junction
structures or mRNA/protein of Connexins (Cx; a component of gap junction) does not
necessarily mean that the gap junction is functional. Dominant negative form of gap
junction might be useful. Currently, ICC- or SMC-specific gene manipulations in our
organoid are not available, and further studies are awaited. It is intriguing that the
coordination between SMC-SMC and ICC-ICC was unaffected by the inhibitors in our
organoids. The possibility cannot be excluded that gap junctions that are not inhibited by
CBX or 18β-GA might operate. However, since these two inhibitors have popularly been
used to block a wide range of gap junctions, alternative possibility arises that cell
communications particularly between SMC-SMC are mediated by other mechanisms than
gap junctions, including mechanical signals as recently shown for coordinated Ca^{2+}
signaling in cardiac muscle cells (Fukui et al., 2021).
357

Possible feedback from SMC's contractility to ICC's oscillatory rhythm

Blebbistatin prevented the contraction of the organoids, which was concomitant
with abrogation of Ca^{2+} transients in ICCs. Since it has been reported that thick myosin
fibers necessary for the contraction are found only in SMCs but not in ICCs
(Gherghiceanu & Popescu, 2005; Rumessen & Thuneberg, 1991, 1996; Sun et al., 2006),
it is likely that ICCs do not have contractile ability. Our observations therefore promote
the possibilities that ICC's pace-making activity requires its own non-muscle myosin II,
and/or mechanical feedback by contracting SMCs. Currently, techniques that distinguish
between these possibilities are unavailable. Nevertheless, if it is the latter case, it would
indicate a signaling from SMC to ICC, the direction of which is unprecedented, since
previous literatures had shown that it was solely an ICC-to-SMC signaling process that
dictated the rhythm of SMC's contraction. We presume that the SMC's contribution to
ICC's pace-making is highly possible since this notion is further supported by other

findings obtained in this study showing that SMCs mediate inter-organoidal phase
coordination (Fig. 8) as more discussed below.
It is of note that the reciprocal interactions between pace-making cells and their
effectors have been reported in studies of neural circuit establishment. Spontaneous
activities emerging in motor neurons during embryogenesis are regulated by feedback
from their governing muscle's contractions so that motor neurons get organized to display
coordinated and stable oscillatory activities (Zeng et al., 2021). Thus, it is tempting to
speculate that during early gut peristalsis, ICCs might receive feedback from their
governing SMCs to generate more stable coordination of pace-making activity among
ICCs. Indeed, we have previously reported that in the very early embryonic gut, origins
of peristaltic wave (OPW) are randomly distributed along the gut axis, but they later
become confined to specific sites showing that ICCs undergo more stable and coordinated
pace-making (Shikaya et al., 2022).
384
Interactions between ICCs and SMCs in the gut contractile organoid
The organoid developed in this study was derived from the muscle layer (also
called *tunica muscularis*) of chicken E15 hindgut devoid of mucosa and serosa, in which
myenteric ICCs (ICC-MY), intramuscular ICCs (ICC-IM), and submucosal ICCs (ICC-
SM) are localized in a way similar to those in mice, shown by staining with antibody
against the chicken c-Kit protein (Yagasaki et al., 2022). In the current study, c-Kit
antibody-staining showed two types of cells in morphology in the gut contractile
organoid: one is multipolar and N-cad-positive ICCs located centrally, and the other is
ICCs that are thin in shape, N-cad negative, and lining beneath the most external layer of
SMCs. Based on the knowledge obtained in studies with mammalian species that ICC-
MY are multipolar whereas ICC-IM are bipolar and tightly associated with adjacent
SMCs (Huizinga et al., 2011; Iino & Horiguchi, 2006), it is conceivable in our organoids
that the central cells are ICC-MY, and the peripherally lining ones are ICC-IM. In
addition, it has been reported in mice that signaling from ICC-IM to SMCs is gap
junction-dependent. Collectively, our observations that gap junction-dependent signal
found between ICCs (central ICC) and SMCs in the organoid (Fig. 5C) could be
interpreted as follows: ICC-MY (central) signals to ICC-IM (second-most peripheral),
which in turn signals to the external SMCs mediated partly by gap junction. Such
sequences of signaling (ICC-MY to ICC-IM to SMCs) has also been proposed in the
intact gut in mammals, although rigid evidence has not been known. Direct comparison
of Ca^{2+} transients between the ICC-IM-like cells and SMCs in our organoid was

technically very difficult since these two cells were both thin and tightly associated each
other.
408
The gut contractile organoid provides a useful model and tool for studying phasic
coordination of oscillatory rhythm
To understand the gut peristaltic movements, which reiterate at specific sites
along the gut axis (Shikaya et al., 2022), deciphering the mechanisms underlying the
coordination/synchronization of oscillators among multiple cells is critical. Exploiting the
finding obtained in the current study that multiple organoids easily fuse each other in vitro,
we have found that two organoids with different oscillatory rhythm eventually coordinate
their phases upon the fusion (Fig. 7). This suggests the ability of ICCs to adjust their
rhythm to their neighbors. During identification of the rhythm-adjustment mediators
using the 3-well hydrogel, contrasting with our original expectation, SMCs that were
crawled out from the organoids and covered the top surface of the hydrogel were able to
mediate the rhythm coordination among organoids (Fig. 8). Whether cellular thin
protrusions connected with each other in narrow channels mediate the coordination
remains undetermined. Nevertheless, our findings have revealed a novel role of SMCs in
mediating rhythm coordination, and as discussed above, this supports the notion of a
SMC-to-ICC signaling. It is unlikely that gap junction plays a major role in such signaling
since gap junction inhibitor yielded no detectable effects in the assays using the 3-well
hydrogel.
In summary, the gut contractile organoid developed in this study serves as a
powerful model to study the establishment and maintenance of oscillatory rhythm (pace-
making) and their coordination in the multicellular systems.
430
431
Materials and Methods
Chicken embryos
Fertilized chicken eggs were obtained from the Yamagishi poultry farms
(Wakayama, Japan) and Takeuchi Farm (Nara, Japan). Embryos were staged according
to embryonic days. All animal experiments were conducted with the ethical approval of
Kyoto University (#202110).
438
Culture preparation of hindgut-derived cells
A hindgut was dissected from E15 chicken embryos (Fig. S1A), and cut into
small pieces. After treating with 25 U dispase (Fujifilm Wako, 383-02281) /phosphate

buffer saline (PBS: 0.14 M NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4-$12H_2O$, 1.8 mM

KH_2PO_4) at 38.5 °C for 40 min, serosa and intestinal epithelium were removed using

forceps. The muscle layer was minced into smaller pieces and treated with 0.2 mg/ml

collagenase (Fujifilm Wako, 038-22361) and 0.25 % trypsin/PBS at 37.0 °C for 30 min.

The reaction was stopped with 1 % FBS/PBS followed by centrifugation at 800 rpm for

3 min. The pellet was resuspended and washed in PBS followed by centrifugation. The

pellet was resuspended in culture medium, and 5.0×10^5 cells were plated in a glass bottom

dish (Matsunami, D11130H) treated with Matrigel (Corning, 354248) prepared in

advance at 38.5 °C for 20 min before use. Poly-lysine and collagen-coated dishes were

purchased (Matsunami, D11131H, D11134H). D-MEM /Ham's F-12 (Wako, 048-29785),

Ham's F-12 (Wako, 087-08335) and Neurobasal medium (Gibco, 21103-049) with $1\times$ B-

27 supplement (Gibco, 17504044) and 0.5 mM L-glutamine were tested. After seeding,

time-lapse images were obtained with CM20 (Evident) under 5% CO_2 and 38.5°C with

2-hour intervals.

Assessment of contraction in the spheroid/cluster

The time-lapse images were obtained by confocal microscopy (Nikon, A1R)

under 5 % CO_2 and 38.5 °C. The bright field images were converted to the sequence of

Tiff images. Using ImageJ (National Institutes of Health, USA), these sequences were

divided into foreground and background automatically (Auto Threshold, mode "default")

and graphed by Plot Z-axis profile.

Immunocytochemistry

An organoid was fixed in acetic acid/ethanol (1:5) for 10 min at room

temperature (RT), and washed in PBS for 10 min at RT. The specimens were

permeabilized in 0.1% Tween-20 in PBS for 10 min at RT, followed by washing in PBS

for 10 min at RT. After blocking with 1 % blocking reagent for 1 hour at RT, specimens

were incubated overnight at 4 ℃ with dilution of 1:300 anti-c-Kit (Yagasaki et al., 2022),

1:300 Tuj-1 (RSD, MAB1195), 1:400 anti-αSMA antibody (Sigma-Aldrich, A5228),

and/or 1:200 anti-N-cadherin antibodies (TAKARA, M110) in 1% blocking regent

(Roche, 1096176)/PBS. Following three times washing in PBS for 10 min each at RT,

specimens were incubated for 1.5 hr at RT with 1:500 anti-rabbit IgG(H+L)-Alexa 488-

conjugated antibody (Donkey; Invitrogen, A21206), anti-mouse IgG_{2a}-Alexa 568-

conjugated antibody (Goat; Invitrogen, A21134), anti-rat IgG (H+L)-Alexa 488-

conjugated antibody (Goat; Invitrogen, A11006) and 1:2000 DAPI. After washing three

times in PBS for 10 min at RT, fluorescent images were obtained using the Nikon A1R

confocal microscope.

Plasmids

pAAV-hSynapsin1-GCaMP6s-P2A-mRuby3 was purchased from Addgene

(112007). Full-length cDNA of GCaMP6s was PCR-amplified:

forward 5'- GCGTACCACTGTGGCATCGATGCCACCATGGGTTCTCA -3',

reverse 5'- GCCCGTACATCGCATCGATTTACTTGTACAGCTCGT -3'.

The retroviral vector RCAS-EGFP was digested with Cla I to remove EGFP, into which

a DNA fragment was inserted by In-Fusion HD Cloning Kit (TAKARA) to produce

RCAS-GCaMP6s-P2A-mRuby3. RCAS-GapEGFP is as previously described (Murai et

al., 2015)

Preparation of retroviral vector particles

RCAS-GCaMP6s-P2A-mRuby3 was transfected into the chicken fibroblast line

cells DF1 using Lipofectamin 2000 (Invitrogen). Transfected cells were cultured in 10

cm culture dish until confluent. The supernatant of transfected DF1 was collected for viral

precipitation, from which retroviral particles were prepared using Retro-Concentin™

Virus Precipitation Solution (SBI, RV100A-1).

Intracellular Ca^{2+} imaging in the gut contractile organoid

At day 2 or day 3 of cell culture, 10 mg/ml polybrene solution (final

concentration 4 µg/ml, Nacalai, 12996-81) and 20 µl opti-mem with above mentioned

virus particles were added to the culture medium to transfect GCaMP6s into organoid-

forming cells. The medium was replaced with fresh medium at day 5 and Ca^{2+} imaging

was performed at day 7. Time-lapse images were obtained using the confocal microscope

(Nikon, A1R) with 700 or 450 msec intervals, and exported intensity of each region of

interest (ROI) into an excel file by time measurement (Nikon, NIS-Elements). From the

excel file, the graphs were plotted with the fluorescence intensity by setting the beginning

of the measurement to zero.

Drug administration

Carbenoxolone (CBX; nacalai tesque, 32775-51) /H_2O, 18beta-Glycyrrhetinic

acid (18β-GA; abcam, ab142579)/DMSO and (-)-Blebbistatin (FUJIFILM Wako, 021-

17041) were prepared. Time-lapse images were acquired for a single organoid before and

after the drug administration. Following the drug adding, organoids were allowed to rest

for 30 min to avoid possible effect of turbulence of the medium, and time-lapse images

were taken for 5 min.
515

Three-well hydrogel fabrication

The photoinitiator P2CK was synthesized as previously reported (Li et al., 2013).
The target product was verified using proton NMR and Bruker's TopSpin software.
Gelatin-Norbornene was synthesized based on previous report (Van Hoorick et al., 2018).
The final reaction mixture was transferred to a dialysis tube (5-6 kDa; Spectra Por, cat.
no. 132680T) for dialysis against pure water at 40°C for 3 days. After dialysis, the
solution's pH was carefully adjusted to 8.0.
A piece of 3-well hydrogel was fabricated on a glass bottom dish (Matsunami)
from 100 μL of a solution containing 20 wt % Gelatin-Norbornene, 2 mM of the
photoinitiator P2CK, and 20 mM of crosslinker (DTT) (TCI, #D1071) using a two-photon
microscope with controllable laser power in 3D space according to the voxel file input
(Olympus, in house customized). The voxel files specifying the 3D shape of the hydrogel
were designed with Fusion 360 software. (Autodisk).
529
530

Acknowledgement

We thank Dr. Scott Gilbert for careful reading of the manuscript and discussion.
We also thank National BioResource Project (Chicken-Quail, Nagoya University) for
their technical help. This work was supported by JSPS KAKENHI Grant Numbers;
23H04933, 20H03259, 20K20520, 20K21425 for Y. T., and 21K06198, 23H04702 for
M. I., and FY 2022 Kusunoki 125 of Kyoto University 125[th] Anniversary Fund for M. I.,
and Ginpu Funds (Kyoto University) and incu-be fund (Leave a Nest Co., Ltd.) for R. Y..
R. Y. is an ex-fellow of JSPS.

539
540

References

Baker, S. A., Leigh, W. A., Del Valle, G., De Yturriaga, I. F., Ward, S. M., Cobine, C. A., Drumm,
B. T., & Sanders, K. M. (2021). Ca(2+) signaling driving pacemaker activity in
submucosal interstitial cells of Cajal in the murine colon. *Elife, 10.*
https://doi.org/10.7554/eLife.64099
Barajas-Lopez, C., Berezin, I., Daniel, E. E., & Huizinga, J. D. (1989). Pacemaker activity recorded
in interstitial cells of Cajal of the gastrointestinal tract. *Am J Physiol, 257*(4 Pt 1), C830-
835. https://doi.org/10.1152/ajpcell.1989.257.4.C830
Camborova, P., Hubka, P., Sulkova, I., & Hulin, I. (2003). The pacemaker activity of interstitial

cells of Cajal and gastric electrical activity. *Physiol Res*, *52*(3), 275-284.
https://www.ncbi.nlm.nih.gov/pubmed/12790758

Chevalier, N. R. (2018). The first digestive movements in the embryo are mediated by
mechanosensitive smooth muscle calcium waves. *Philos Trans R Soc Lond B Biol Sci*,
373(1759). https://doi.org/10.1098/rstb.2017.0322

Chevalier, N. R., Ammouche, Y., Gomis, A., Teyssaire, C., de Santa Barbara, P., & Faure, S. (2020).
Shifting into high gear: how interstitial cells of Cajal change the motility pattern of the
developing intestine. *Am J Physiol Gastrointest Liver Physiol*, *319*(4), G519-g528.
https://doi.org/10.1152/ajpgi.00112.2020

Chevalier, N. R., Dacher, N., Jacques, C., Langlois, L., Guedj, C., & Faklaris, O. (2019).
Embryogenesis of the peristaltic reflex. *J Physiol*, *597*(10), 2785-2801.
https://doi.org/10.1113/jp277746

Cousins, H. M., Edwards, F. R., Hickey, H., Hill, C. E., & Hirst, G. D. (2003). Electrical coupling
between the myenteric interstitial cells of Cajal and adjacent muscle layers in the guinea-
pig gastric antrum. *J Physiol*, *550*(Pt 3), 829-844.
https://doi.org/10.1113/jphysiol.2003.042176

Daniel, E. E., & Wang, Y. F. (1999). Gap junctions in intestinal smooth muscle and interstitial
cells of Cajal. *Microsc Res Tech*, *47*(5), 309-320. https://doi.org/10.1002/(SICI)1097-
0029(19991201)47:5<309::AID-JEMT2>3.0.CO;2-K

Duband, J. L., Gimona, M., Scatena, M., Sartore, S., & Small, J. V. (1993). Calponin and SM 22
as differentiation markers of smooth muscle: spatiotemporal distribution during avian
embryonic development. *Differentiation*, *55*(1), 1-11. https://doi.org/10.1111/j.1432-
0436.1993.tb00027.x

Fukui, H., Chow, R. W., Xie, J., Foo, Y. Y., Yap, C. H., Minc, N., Mochizuki, N., & Vermot, J.
(2021). Bioelectric signaling and the control of cardiac cell identity in response to
mechanical forces. *Science*, *374*(6565), 351-354.
https://doi.org/10.1126/science.abc6229

Gabella, G., & Blundell, D. (1981). Gap junctions of the muscles of the small and large intestine.
Cell Tissue Res, *219*(3), 469-488. https://doi.org/10.1007/BF00209987

Gherghiceanu, M., & Popescu, L. M. (2005). Interstitial Cajal-like cells (ICLC) in human resting
mammary gland stroma. Transmission electron microscope (TEM) identification. *J Cell*
Mol Med, *9*(4), 893-910. https://doi.org/10.1111/j.1582-4934.2005.tb00387.x

Graham, H. K., Maina, I., Goldstein, A. M., & Nagy, N. (2017). Intestinal smooth muscle is
required for patterning the enteric nervous system. *J Anat*, *230*(4), 567-574.
https://doi.org/10.1111/joa.12583

Grosse, A. S., Pressprich, M. F., Curley, L. B., Hamilton, K. L., Margolis, B., Hildebrand, J. D., &

Gumucio, D. L. (2011). Cell dynamics in fetal intestinal epithelium: implications for
intestinal growth and morphogenesis. *Development, 138*(20), 4423-4432.
https://doi.org/10.1242/dev.065789

Huizinga, J. D., Martz, S., Gil, V., Wang, X. Y., Jimenez, M., & Parsons, S. (2011). Two
independent networks of interstitial cells of cajal work cooperatively with the enteric
nervous system to create colonic motor patterns. *Front Neurosci, 5*, 93.
https://doi.org/10.3389/fnins.2011.00093

Huizinga, J. D., Thuneberg, L., Kluppel, M., Malysz, J., Mikkelsen, H. B., & Bernstein, A. (1995).
W/kit gene required for interstitial cells of Cajal and for intestinal pacemaker activity.
Nature, 373(6512), 347-349. https://doi.org/10.1038/373347a0

Iino, S., & Horiguchi, K. (2006). Interstitial cells of cajal are involved in neurotransmission in the
gastrointestinal tract. *Acta Histochem Cytochem, 39*(6), 145-153.
https://doi.org/10.1267/ahc.06023

Iino, S., Horiguchi, S., Horiguchi, K., & Nojyo, Y. (2007). Interstitial cells of Cajal in the
gastrointestinal musculature of W mutant mice. *Arch Histol Cytol, 70*(3), 163-173.
https://doi.org/10.1679/aohc.70.163

Kito, Y., Ward, S. M., & Sanders, K. M. (2005). Pacemaker potentials generated by interstitial cells
of Cajal in the murine intestine. *Am J Physiol Cell Physiol, 288*(3), C710-720.
https://doi.org/10.1152/ajpcell.00361.2004

Kluppel, M., Huizinga, J. D., Malysz, J., & Bernstein, A. (1998). Developmental origin and Kit-
dependent development of the interstitial cells of cajal in the mammalian small intestine.
Dev Dyn, 211(1), 60-71. https://doi.org/10.1002/(SICI)1097-
0177(199801)211:1<60::AID-AJA6>3.0.CO;2-5

Komuro, T., Tokui, K., & Zhou, D. S. (1996). Identification of the interstitial cells of Cajal. *Histol
Histopathol, 11*(3), 769-786. https://www.ncbi.nlm.nih.gov/pubmed/8839765

Li, Z. Q., Torgersen, J., Ajami, A., Mühleder, S., Qin, X. H., Husinsky, W., Holnthoner, W.,
Ovsianikov, A., Stampfl, J., & Liska, R. (2013). Initiation efficiency and cytotoxicity of
novel water-soluble two-photon photoinitiators for direct 3D microfabrication of
hydrogels. *Rsc Advances, 3*(36), 15939-15946. https://doi.org/10.1039/c3ra42918k

Liu, L. W., Thuneberg, L., & Huizinga, J. D. (1998). Development of pacemaker activity and
interstitial cells of Cajal in the neonatal mouse small intestine. *Dev Dyn, 213*(3), 271-282.
https://doi.org/10.1002/(SICI)1097-0177(199811)213:3<271::AID-AJA4>3.0.CO;2-R

Mei, F., Han, J., Huang, Y., Jiang, Z. Y., Xiong, C. J., & Zhou, D. S. (2009). Plasticity of interstitial
cells of cajal: a study in the small intestine of adult Guinea pigs. *Anat Rec (Hoboken),
292*(7), 985-993. https://doi.org/10.1002/ar.20928

Murai, H., Tadokoro, R., Sakai, K., & Takahashi, Y. (2015). In ovo gene manipulation of

melanocytes and their adjacent keratinocytes during skin pigmentation of chicken embryos. *Dev Growth Differ*, *57*(3), 232-241. https://doi.org/10.1111/dgd.12201

Rohr, S., Kucera, J. P., & Kleber, A. G. (1998). Slow conduction in cardiac tissue, I: effects of a reduction of excitability versus a reduction of electrical coupling on microconduction. *Circ Res*, *83*(8), 781-794. https://doi.org/10.1161/01.res.83.8.781

Rumessen, J. J., & Thuneberg, L. (1991). Interstitial cells of Cajal in human small intestine. Ultrastructural identification and organization between the main smooth muscle layers. *Gastroenterology*, *100*(5 Pt 1), 1417-1431. https://www.ncbi.nlm.nih.gov/pubmed/2013387

Rumessen, J. J., & Thuneberg, L. (1996). Pacemaker cells in the gastrointestinal tract: interstitial cells of Cajal. *Scand J Gastroenterol Suppl*, *216*, 82-94. https://doi.org/10.3109/00365529609094564

Sanders, K. M., Drumm, B. T., Cobine, C. A., & Baker, S. A. (2024). Ca(2+) dynamics in interstitial cells: foundational mechanisms for the motor patterns in the gastrointestinal tract. *Physiol Rev*, *104*(1), 329-398. https://doi.org/10.1152/physrev.00036.2022

Sanders, K. M., Kito, Y., Hwang, S. J., & Ward, S. M. (2016). Regulation of Gastrointestinal Smooth Muscle Function by Interstitial Cells. *Physiology (Bethesda)*, *31*(5), 316-326. https://doi.org/10.1152/physiol.00006.2016

Sanders, K. M., Publicover, N. G., & Ward, S. M. (1991). Involvement of interstitial cells of Cajal in pacemaker activity of canine colon. *J Smooth Muscle Res*, *27*(1), 1-11. https://doi.org/10.1540/jsmr.27.1

Sanders, K. M., Ward, S. M., & Koh, S. D. (2014). Interstitial cells: regulators of smooth muscle function. *Physiol Rev*, *94*(3), 859-907. https://doi.org/10.1152/physrev.00037.2013

Schultz, T., Daniel, V., & Daniel, E. E. (2003). Does ICC pacing require functional gap junctions between ICC and smooth muscle in mouse intestine? *Neurogastroenterol Motil*, *15*(2), 129-138. https://doi.org/10.1046/j.1365-2982.2003.00401.x

Shikaya, Y., Takase, Y., Tadokoro, R., Nakamura, R., Inaba, M., & Takahashi, Y. (2022). Distribution Map of Peristaltic Waves in the Chicken Embryonic Gut Reveals Importance of Enteric Nervous System and Inter-Region Cross Talks Along the Gut Axis. *Front Cell Dev Biol*, *10*, 827079. https://doi.org/10.3389/fcell.2022.827079

Sun, X., Yu, B., Xu, L., Dong, W., & Luo, H. (2006). Interstitial cells of Cajal in the murine gallbladder. *Scand J Gastroenterol*, *41*(10), 1218-1226. https://doi.org/10.1080/00365520600708800

Takaki, M. (2003). Gut pacemaker cells: the interstitial cells of Cajal (ICC). *J Smooth Muscle Res*, *39*(5), 137-161. https://doi.org/10.1540/jsmr.39.137

Takayama, I., Horiguchi, K., Daigo, Y., Mine, T., Fujino, M. A., & Ohno, S. (2002). The interstitial

cells of Cajal and a gastroenteric pacemaker system. *Arch Histol Cytol, 65*(1), 1-26. https://doi.org/10.1679/aohc.65.1

Takeda, Y., Ward, S. M., Sanders, K. M., & Koh, S. D. (2005). Effects of the gap junction blocker glycyrrhetinic acid on gastrointestinal smooth muscle cells. *Am J Physiol Gastrointest Liver Physiol, 288*(4), G832-841. https://doi.org/10.1152/ajpgi.00389.2004

Thomsen, L., Robinson, T. L., Lee, J. C., Farraway, L. A., Hughes, M. J., Andrews, D. W., & Huizinga, J. D. (1998). Interstitial cells of Cajal generate a rhythmic pacemaker current. *Nat Med, 4*(7), 848-851. https://doi.org/10.1038/nm0798-848

Torihashi, S., Fujimoto, T., Trost, C., & Nakayama, S. (2002). Calcium oscillation linked to pacemaking of interstitial cells of Cajal: requirement of calcium influx and localization of TRP4 in caveolae. *J Biol Chem, 277*(21), 19191-19197. https://doi.org/10.1074/jbc.M201728200

Torihashi, S., Nishi, K., Tokutomi, Y., Nishi, T., Ward, S., & Sanders, K. M. (1999). Blockade of kit signaling induces transdifferentiation of interstitial cells of cajal to a smooth muscle phenotype. *Gastroenterology, 117*(1), 140-148. https://doi.org/10.1016/s0016-5085(99)70560-3

Van Hoorick, J., Gruber, P., Markovic, M., Rollot, M., Graulus, G. J., Vagenende, M., Tromayer, M., Van Erps, J., Thienpont, H., Martins, J. C., Baudis, S., Ovsianikov, A., Dubruel, P., & Van Vlierberghe, S. (2018). Highly Reactive Thiol-Norbornene Photo-Click Hydrogels: Toward Improved Processability. *Macromol Rapid Commun, 39*(14), e1800181. https://doi.org/10.1002/marc.201800181

Wang, Q., Wang, K., Solorzano-Vargas, R. S., Lin, P. Y., Walthers, C. M., Thomas, A. L., Martín, M. G., & Dunn, J. C. Y. (2018). Bioengineered intestinal muscularis complexes with long-term spontaneous and periodic contractions. *PLoS One, 13*(5), e0195315. https://doi.org/10.1371/journal.pone.0195315

Yagasaki, R., Shikaya, Y., Kawachi, T., Inaba, M., Takase, Y., & Takahashi, Y. (2022). Newly raised anti-c-Kit antibody visualizes morphology of interstitial cells of Cajal in the developing gut of chicken embryos. *Dev Growth Differ, 64*(8), 446-454. https://doi.org/10.1111/dgd.12808

Yang, P., Yu, Z., Gandahi, J. A., Bian, X., Wu, L., Liu, Y., Zhang, L., Zhang, Q., & Chen, Q. (2012). The identification of c-Kit-positive cells in the intestine of chicken. *Poult Sci, 91*(9), 2264-2269. https://doi.org/10.3382/ps.2011-02076

Zeng, X., Komanome, Y., Kawasaki, T., Inada, K., Jonaitis, J., Pulver, S. R., Kazama, H., & Nose, A. (2021). An electrically coupled pioneer circuit enables motor development via proprioceptive feedback in Drosophila embryos. *Curr Biol, 31*(23), 5327-5340 e5325. https://doi.org/10.1016/j.cub.2021.10.005

694 <u>**Figures**</u>

A

B

696 **Figure 1 Culture of muscle layer-derived cells prepared from embryonic hindgut.**

697 (A) Culture of muscle layer-derived cells prepared from embryonic hindgut with FBS

698 free-media and substrates. (B) Long-term time-lapse imaging after seeding on Matrigel

699 with Neurobasal media. The images show the ability of these cells to self-assemble at 0,

700 20, 40, 60, 80 and 100 hours taken from Movie 1. Scale bars: 100 μm in A, B.

Figure 2 Forming spheroids exhibited reiterated contractions.

(A) Clusters/spheroids on day 3, 5 and 7 exhibited periodic contractions. Graphs show
contractions quantified by differences in the intensity of brightfield in sequential images.
(B) Frequency of contractions of the cluster/spheroid on day 3-7 (N=day 3: 46, day 5: 35,
day 7: 60). The number on each graph shows the average of contraction frequencies. Scale
bars: 50 μm in A.

Figure 3 The cluster/spheroid is composed of ICCs internally and SMCs peripherally. (A) Co-staining with anti-c-Kit- and αSMA antibodies. White arrowheads show co-expression of c-Kit and αSMA at Day 3. Yellow arrowheads indicate a cell expressing αSMA but not c-Kit at day 7. A diagram shows an image of cell arrangement

713   (green: ICC, magenta: SMC) in the spheroid at day 7. (B) Cell morphology in the spheroid

714   at day 6 revealed by RCAS-gapEGFP expression. (C) Staining of spheroids at day 5 with

715   anti-N-cadherin antibody. A white arrowhead shows expression of N-cadherin. A yellow

716   arrowhead shows a cell which does not express N-cadherin in the outer region of the

717   spheroid. Scale bars: 30 μm in A-C, 10 μm in a.

Figure 4 Ca²⁺ imaging of the gut contractile organoid revealed intercellular synchronization.

(A) RCAS-GCaMP6s-P2A-mRuby3 plasmid. (B) Ca^{2+} imaging of gut contractile organoid during relaxation and contraction. Ca^{2+} dynamics (green) and contraction (grey) of gut contractile organoid. (C) Simultaneous measurement of intercellular Ca^{2+} dynamics between ICC-ICC, SMC-SMC or ICC-SMC. Three or two ROIs with Ca^{2+} signal-positive cells were set in a single organoid. Graphs show Ca^{2+} dynamics in the

727   ROIs. Magnified view shows that a rise of Ca^{2+} signal in ICC (green) preceded that in

728   SMC (magenta). Scale bar: 50 μm in B.

Figure 5 Gap junction inhibitor exerted limited effects on the synchronization of Ca²⁺ dynamics.

GCaMP6s-expressing organoids were cultured with CBX, and three or two ROIs were assessed for their Ca²⁺ synchronization. A single organoid was subjected to the assessment before (0 μM; control) and after administrations of 100 μM CBX. The

736   synchronization was unaffected between (A) SMC-SMC and (B) ICC-ICC, but that in

737   (C) ICC-SMC was partially affected. Magnified view shows that the preceding rise of

738   Ca^{2+} in ICC (green) before CBX was abrogated after CBX. Scale bar: 50 µm in A.

Figure 6 The organoidal contraction is important for Ca²⁺ dynamics in ICCs.
GCaMP6s-expressing organoids were cultured with Blebbistatin. (A) Ca²⁺ oscillation was extinguished in the entire organoid by administration of Blebbistatin. (B-D) Ca²⁺ dynamics in (B) SMC-SMC, (C) ICC-ICC and (D) ICC-SMC. Three or two ROIs were assessed for their Ca²⁺ synchronization. A single organoid was subjected to the

746 assessment before (0 µM; control) and after administrations of 10 µM Blebbistatin. (E)

747 Ca^{2+} transients in a single ICC in 0 µM, 5 µM, and 10 µM. Scale bar: 50 µm in B.

Figure 7 Ca^{2+} dynamics in multiple gut contractile organoids undergo synchronization upon their fusion.

(A) Time-lapse imaging of organoidal fusion. (B) When two organoids that originally showed independent rhythm of Ca^{2+} fused to each other, their rhythm became synchronized after fusion (24 h). (C) Protrusions between two neighboring organoids. White arrowheads show three protrusions from the left organoid. Scale bars: 100 μm in A-C, 20 μm in a.

Figure 8 Smooth muscle cells mediate the Ca²⁺ synchronization between organoids.
(A) Diagram of 3-well hydrogel in which an organoid placed in each well. This gel mold does not allow organoidal bodies to fuse to each other, but allow them to extend/migrate protrusions/cells through a narrow channel connecting the wells (B). (C) After 3 days, three organoids displayed synchronization of Ca²⁺ dynamics. (D) Some organoid-derived cells crawled out from the wells and covered the top surface of the hydrogel, resulting in

bridging the three unfused organoids. (E) Diagram of 3-well hydrogel *without* channels (F) Surface-covering cells were SMCs (αSMA positive and c-Kit-negative). (G) In the hydrogel without channels but with surface-covered SMCs, Ca^{2+} dynamics in the 3 organoids were synchronized. (H) The Ca^{2+} synchronization shown in (G) was not altered by 18β-GA. Scale bars: 50 μm in B, 100 μm in C-H.

773 **Supplementary Figure S1**

774   (A) Chicken embryonic hindgut at E15. The hindgut was dissected from the

bottom of the cecum to the front of the cloaca (white lines). (B) Three layers of E15

hindgut: serosa, muscle layer, intestinal epithelium. Remak's ganglion was also removed.

Scale bars: 1 mm in A, B

Supplementary Figure S2

(A) Co-staining of Day 7 organoids with anti-c-Kit- and Tuj1 antibodies. White

arrowhead shows a Tuj1$^+$ cell. (B) Representation of Tuj1$^+$ cells in an organoid at day 7.

Scale bar: 30 μm in A.

Supplementary Figure S3

(A) Day 7 organoid exhibited periodic contractions both before (0 μM; control) and after adding CBX. Graphs show contractions quantified by the difference in intensity of brightfield in sequential images. (B) Relative changes to the control in contraction frequency in day 7 organoid after CBX. The number is an average. Scale bar: 50 μm in A.

Movie 1

Long-term time-lapse imaging after seeding. Time lapse was taken every 2 hours for a total of 100 hours. This video corresponds to Fig. 1B.

Movie 2

Periodic contractions of the cluster/spheroid on day 3, 5 and 7. Time-lapse images were taken with 700 msec intervals for 5 min. This video corresponds to Fig. 2A.

Movie 3

Periodic contractions of a day 7 organoid with CBX. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. S3.

Movie 4

Ca^{2+} dynamics in day 7 gut contractile organoid are concomitant with its contractions. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 4.

Movie 5

Ca^{2+} dynamics in a gut contractile organoid with CBX. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 5.

Movie 6

Ca^{2+} dynamics in day 7 gut contractile organoid with Blebbistatin. Time-lapse images were obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 6.

Movie 7

Live imaging during fusion of multiple organoids. Time-lapse images were obtained every 10 minutes for 4 hours. This video corresponds to Fig. 7A.

Movie 8

Ca^{2+} transients in two gut contractile organoids before and after fusion. Time-lapse images were obtained with 450 msec intervals for 5 min. This video corresponds to Fig. 7B.

Movie 9

Ca^{2+} transients in three gut contractile organoids in 3-well hydrogel with channels.
Time-lapse images were obtained with 700 msec intervals for 5 min. This video
corresponds to Fig. 8C.

Movie 10

Ca^{2+} transients in three gut contractile organoids in 3-well hydrogel *without*
channels. Time-lapse images were obtained with 450 msec intervals for 5 min.   This
video corresponds to Fig. 8G.

Movie 11

Similar assay to movie 10 with 18β-GA administration. Time-lapse images were
obtained with 700 msec intervals for 5 min. This video corresponds to Fig. 8H.