Metatranscriptomic Compendium of 55900 RNA Viruses Deciphers Human Health Implication and Ecological roles of RNA Virome in Global Wastewater Treatment Plants

Ling Yuan[1,2], Lin-xing Chen[3], Feng Ju[1,2,4,5]

[1] Environmental Microbiome and Biotechnology Laboratory (EMBLab), Research Center for Industries of the Future, Westlake University. Hangzhou 310030, China

[2] Zhejiang Provincial Key Laboratory of Intelligent Low-Carbon Biosynthesis, School of Engineering, Westlake University, Hangzhou 310030, China

[3] The Department of Environmental Science and Engineering, University of Science and Technology of China, Hefei, 230026, China

[4] Westlake Laboratory of Life Sciences and Biomedicine, School of Life Sciences, Westlake University, Hangzhou 310024, China

[5] Center for Future Foods, Muyuan Laboratory, Zhengzhou 450016, Henan Province, China.

* Correspondence to

Dr. Feng Ju

Associate Professor

Westlake University

Email: jufeng@westlake.edu.cn

Address: 600 Dunyu Road, Xihu District, Hangzhou 310030, China

Tel.: 571-87963205 (lab), 571-87380995 (office); Fax: 0571-85271986

Abstract

RNA viruses are widely recognized for their roles in causing human diseases, yet their diversity and functional roles in engineered systems remain largely unexplored. Wastewater treatment plants (WWTPs) are eco-friendly biotechnological systems where the roles of RNA viruses in process engineering and sanitation remain unclear. This study analyzed the largest RNA sequencing data (> 3.8 Tb) from global WWTPs to examine the diversity, host associations, and auxiliary metabolic functions of RNA viruses. We identified 11,414 RNA virus operational taxonomic units (vOTUs), expanding the known diversity of RNA viruses in engineered systems by 65%. The RNA viral community in WWTPs was dominated by prokaryotic viruses, including established RNA phage lineages (*Leviviricetes* and *Cystoviridae*) and potential novel clades within *Picobirnaviridae*, *Partitiviridae*, base-*Cystoviridae*, and base-*Howeltoviricetes*. Notably, the novel phage clade within base-*Howeltoviricetes* was discovered to prevail in WWTPs and diverse ecosystems globally, with a single vOTU in this clade associated with the pathogenic *Campylobacterota Aliarcobacter cryaerophilus*, suggesting its broad niche breadths and potential for RNA phage therapy applications. Additionally, metatranscriptomics-based identification of human RNA vOTUs (e.g., *Astrovirus*, *Rotavirus*, and *Norovirus*), and high-risk RNA viruses (e.g., SARS-CoV-2) identified by short-read alignments emphasized the utility of wastewater-based surveillance for public health monitoring. The discovery of auxiliary metabolic genes encoded by RNA viruses further revealed their involvement in critical host metabolic pathways such as translation and cellular respiration. These findings unveil the previously hidden diversity, health implications, and biochemical impacts of RNA viruses within WWTPs, underscoring their multifaceted roles in

47 these critical engineered systems.

48 Key Words: RNA virus, virus-host relationship, auxiliary metabolic gene, wastewater-

49 based surveillance, human health, metatranscriptomics

Introduction

RNA viruses are abundant and influential biological entities across natural ecosystems,

infecting both eukaryotes and prokaryotes. They play critical roles in human health[1-3],

host metabolism manipulation[4-6], and maintaining global biodiversity[7-9]. Recent

advancements in RNA virus research, driven by the detection of RNA-dependent

RNA polymerase (RdRp) in metatranscriptomic datasets, have dramatically expanded

our understanding of their diversity[6-8, 10-14]. Metatranscriptomics has become a

powerful tool for exploring RNA viruses across diverse ecosystems, revealing not

only viruses associated with human, livestock, and crop diseases [12, 15], but also

previously unculturable viruses. Recent large-scale surveys of RNA viruses in marine[6,

10, 11] and soil ecosystems[13, 16] have demonstrated that RNA viruses are integral

components of biodiversity in aquatic and terrestrial ecosystems.

Wastewater treatment plants (WWTPs) are among the world's most significant

biotechnological systems, designed to safeguard human, animal and environmental

health. These systems purify wastewater by leveraging enhanced microbial

metabolisms to remove organic pollutants and nutrients. Consequently, WWTPs

function as both physical and ecological connectors between human populations and

natural ecosystems, particularly aquatic environments[17]. Viruses play a pivotal role in

WWTPs, influencing the structure and dynamics of microbial communities[18] and

serving as valuable indicators of population health[19]. While considerable progress has

been made in understanding the diversity of DNA viruses in WWTPs[20, 21], insights

into RNA viruses are comparatively nascent[12, 22]. The RNA virosphere in wastewater

comprises represents a highly diverse assemblage, including prokaryotic RNA viruses,

human RNA viruses of feces and urine origins, and RNA viruses associated with

animals, plants, protozoa, and fungi[23].

Prokaryotic RNA viruses, particularly RNA phages, play a crucial role in

wastewater treatment plant (WWTP) ecosystems by infecting and lysing bacterial

hosts. This activity regulates bacterial populations and shapes the microbial

community structure, which is vital for the microbial processes driving WWTP

functionality. Understanding these dynamics requires the identification and

characterization of specific RNA phage clades, including novel lineages.

Simultaneously, human RNA viruses present in wastewater have emerged as powerful

tools for public health surveillance. Wastewater-based epidemiology (WBE), as

demonstrated during the COVID-19 pandemic, has provided valuable insights into

population-wide health trends and enabled early detection of outbreaks[19, 24]. Despite

these advancements, many questions remain unanswered, such as the comprehensive

spectrum of human RNA viruses present in WWTPs. Furthermore, while auxiliary

metabolic genes (AMGs) are widely found in DNA viruses[20, 25-28], it remains unclear

whether and how RNA viruses can affect host metabolism and biochemical cycling

through AMGs. Addressing these gaps on the diversity, host interactions, biochemical

roles, and fate of RNA viruses in WWTPs could uncover their novel ecological

functions and potentially improve the efficiency of WWTP processes.

To bridge these knowledge gaps, we conducted a comprehensive study to

investigate RNA viruses and their ecological roles in WWTPs worldwide. We

compiled the WWTP RNA Viruses Database (WRVirDB) by analyzing the largest

metatranscriptomic dataset of WWTPs to date (>3.8 Tb). This effort identified

abundant RNA phages, including both known and newly discovered clades, and

explored the presence of RNA virus-encoded AMGs, revealing their potential

ecological significance. Additionally, we mapped the global distribution of human

RNA viruses in wastewater and identified novel viral species, highlighting the utility

of wastewater-based surveillance (WBS) for environmental epidemiology and public

health management. This research aims to enhance our understanding of RNA viral

diversity, optimize wastewater treatment processes, and advance strategies for

monitoring public health through wastewater microbiome-based analysis.

Results

Metatranscriptomic profiling and cross-compartment distribution patterns of RNA viruses in the global WWTPs

To explore the diversity of RNA viruses in global WWTPs, we analyzed > 3.8 Tb of

RNA sequencing data from 557 publicly available metatranscriptomes spanning five

WWTP compartments (Fig. 1a), including influent (IN), denitrification sludge (DS),

activated sludge (AS), effluent (EF), and anaerobic digestion sludge (AD). A total of

55,900 RNA viral contigs were identified by iteratively hidden Markov model

(HMM)-based searching of RdRp genes and subsequent assessment of RNA virus

authenticity (See Methods and Extended Data Fig. 1). These RNA viral contigs were

further clustered into 11,414 RNA virus operational taxonomic units (vOTUs) at an

average nucleotide identity (ANI) of 90% over 80% alignment fraction of the shorter

sequence (AF) to empirically evaluate RNA viruses at the "species" rank

(Supplementary Table 2). Of these RNA vOTUs, 9,276 encoded "complete" (or near-

complete) RdRp domains. Although the recovery of RNA viral contigs was gradually

to be saturated after eight iterative searches, the overall diversity of unique RNA

vOTUs in the WWTPs showed no indication of reaching saturation (Fig. 1b),

suggesting that additional samples are required to assemble a more comprehensive

catalog of RNA viruses in WWTP systems.

Further analysis of α-diversity patterns of RNA vOTUs revealed a highly variable

richness of RNA viruses in the WWTP samples (Fig. 1c). Non-Metric

Multidimensional Scaling (NMDS) analysis based on vOTUs composition

demonstrated a clear separation of RNA viral communities by sample type (ANOSIM

R^2 = 0.821, P = 0.001; Fig. 1d). To estimate the absolute concentration of RNA

vOTUs, we leveraged 47 samples from Switzerland spiked with known copies of

mRNA internal standards[29]. The absolute concentration of RNA vOTUs estimated

based on the recovery of internal standards from metatranscriptomic reads reached ~

10^{12} copies/gram-biomass in sludge samples and ~ 10^{10} copies/gram-biomass in

influent and effluent wastewater samples (Fig. 1e). Notably, the effluent samples from

Switzerland displayed higher variations in terms of α-diversity and combined absolute

concentration of RNA vOTUs compared to the related influent samples

(Supplementary Fig. 1 and Fig. 1e), indicating that wastewater treatment processes

may exert uneven filtering or removal effects on the RNA virome, leading to more

dispersed RNA viral community structure in the effluent (Supplementary Text 1).

Knowledge expansion of phylogenetic diversity and taxonomic novelty of RNA

viruses in the global WWTPs

RdRp-based megataxon clustering and phylogeny analysis supported the five

established RNA virus phyla approved by the International Committee on Taxonomy

of Viruses (ICTV) (Supplementary Table 4, Fig. 2a and Supplementary Fig. 2). Most

of the RNA vOTUs that encoded "complete" RdRp domains (8,755 / 9,276) can be

posited into already established phyla. Moreover, 5,945 RNA vOTUs could be

classified into known viral families. The RNA vOTUs were mostly taxonomically

assigned to *Fiersviridae* (1,772) and *Steitzviridae* (580), both of which were known

viral families infecting prokaryotes, followed by *Botourmiaviridae* (488), which was

known to infect plants or fungi (Extended Data Fig. 2).

To assess our expansion of RNA virus diversity in engineered systems, the RNA

viruses identified in this study were clustered with the publicly available RNA viral

sequences from the NCBI Virus database and recent RNA virome studies[10-13, 16].

Clustering was conducted at both the genome level (GRvOTU set) and the RdRp gene

level (RCR90 set). Our findings substantially increased the diversity of unique RNA

viral clusters in engineered systems, from 9,709 previously known RNA viral clusters

to a total of 16,057 RNA viral clusters (Fig. 2b). At the viral family level, clustering

statistics based on genome and RdRp gene data were highly consistent

(Supplementary Table 5). Although WWTP RNA viruses showed a limited impact on

expanding the overall diversity for RNA viral families, they significantly enriched the

diversity of bacteriophages within engineered systems (Fig. 2a and Supplementary

Table 5). *Leviviricetes* and *Cystoviridae* are the only two groups known to infect

bacteria so far. The WWTP RNA viruses discovered in this study contributed 32.4%

to 73.0% of novel diversity to RNA viral families within these two clades in

engineered systems (Extended Data Fig. 3).

The RNA viruses identified in this study also supported the establishment of

potential new clades. A substantial number of RNA viruses encoding "complete"

RdRp could not be classified into established viral families (n = 3,331). These family-

unassigned viruses were grouped at broader taxonomic levels, including phylum-level

unclassified groups (111), phylum (1,762), class (300), or order (1,158) level. For

example, many WWTP RdRps (n = 1350) and reference RdRps could only be

classified into *Kitrinoviricota* phylum without further assigned taxonomic information

(Supplementary Fig. 2). These *Kitrinoviricota* RdRps were distributed among three

RdRp megataxa (2, 7 and 16, Supplementary Fig. 2 and Table 4). Furthermore, one

RdRp megataxon predominantly composed of WWTP-derived RdRps, emerged as a

sister lineage to *Ellioviricetes* in the phylogenetic tree, potentially representing a new

class within the *Negarnaviricota* phylum (Extended Data Fig. 4). Together, these

results depict the most complete landscape of phylogenetic and taxonomical diversity

of RNA viruses in the global WWTPs and underscored the necessity for establishing

new taxonomic frameworks for RNA viruses.

Tracking host associations of RNA prokaryotic viruses in the global WWTPs

Through host inference based on the established Orthornaviran taxonomy, prokaryotic

RNA viruses from known phage clades accounted for nearly one-third (3,324, 29.1%)

of the total RNA vOTUs richness (*Leviviricetes* n = 3,245 and *Cystoviridae* n = 79)

(Fig. 3a). The relative abundance patterns of RNA vOTUs with different host ranges

varied across sample types and countries but suggested the dominance of

bacteriophages in nearly all sludge samples. (Extended Data Fig. 5 and

Supplementary Text 2).

The identification of Shine-Dalgarno (SD) motifs was also used to infer

prokaryotic RNA viral clades. Among the RNA vOTUs classified to *Leviviricetes*,

83.3% of the open reading frames (ORFs) were associated with the SD motif.

Similarly, 84.9% of ORFs in *Cystoviridae* were linked to the SD motif. Furthermore,

several additional clades were discovered to be probable RNA phages due to their

enrichment of SD motifs. These included i) *Picobirnaviridae* (n = 87), ii) a subset of

Partitiviridae (n = 120), iii) a family-unclassified clade basal to *Cystoviridae* (n = 13)

(Extended Data Fig. 6), consistent with a recent study[12], and iv) a clade basal to

Howeltoviricetes (n = 194, Fig. 3b-c), which was reported here for the first time.

The base-*Howeltoviricetes* phage clade was phylogenetically classified under the

family *Mitoviridae* within the *Lenarviricota* phylum (Fig. 3b). Over 85% of the

predicted ORFs in this clade were associated with the SD motif (Fig. 3b-c). This

generalist phage clade was identified in 390 out of 557 WWTP metatranscriptomes.

Additionally, the GRvOTU clustering set facilitated the assessment of the distribution

of the base-*Howeltoviricetes* phage clade across public datasets. Based on the

phylogenetic tree of this clade (Fig. 3c), all viral sequence members belonging to this

clade were extracted from the GRvOTU set. The distribution of this clade was then

determined based on the types and geographical locations of the originating samples

of the viral sequence members. Base-*Howeltoviricetes* phage clade was found not

only present in WWTPs but was also widely distributed across diverse ecosystems

globally, including aquatic environments, soil, and samples associated with animals

and plants (Fig. 3d).

The CRISPR spacer matching linked one vOTU within the base-*Howeltoviricetes*

phage clade to its probable specific host. Several viral contigs from vOTU FD-

DS_117671 matched CRISPR spacers identified in bacterial genomes from the NCBI

RefSeq database and in contigs from WWTP metagenomic assemblies (Fig. 4a).

These matched sequences were taxonomically assigned to the phylum

Campylobacterota, with the most specific classification identifying *Aliarcobacter*

cryaerophilus, a globally emerging foodborne and zoonotic pathogen (Fig. 4a). Virus-

host abundance quantification throughout the WWTPs revealed significantly higher

relative abundance of both the vOTU representative and its host in wastewater

samples compared to sludge samples (Mann-Whitney U test, all $P < 10^{-6}$, Fig. 4b). In

time-series samples (n = 47, sampled approximately weekly), the abundance of the

vOTU representative exhibited a significant positive correlation with the abundance

of *Aliarcobacter cryaerophilus* (Spearman r = 0.76, p < 10□□), further supporting the

predicted virus-host relationship.

In most WWTP samples, *Leviviricetes* was the most abundant group of RNA

bacteriophages (Extended Data Fig. 5). However, base-*Howeltoviricetes* phages

exhibited substantial abundance in several samples and, in certain cases, became the

dominant phage group, contributing to over 50% abundance of the RNA viral

community. Examples included TU-EF_29550 in three BE WWTP samples and AH-

EF_62580 in SF50-SF52 samples (Fig. 4c). Genome extension of these high-

abundance base-*Howeltoviricetes* phages (See Methods) revealed that they likely

encode only RdRp (Fig. 4c). Functional annotation did not identify capsid, maturation

proteins, or other common RNA viral genes in the genomes of base-*Howeltoviricetes*

phages. This observation consisted with the gene composition of their closest

phylogenetic relatives, the *Mitoviridae*, a capsid-less viral family that usually encodes

only RdRp. Together, these findings provide first insights into the broad distribution,

host associations, and genomic features of base-*Howeltoviricetes* phages, a novel

RNA bacteriophage clade.

Prevalence, concentration, and fate of human RNA virus in the global WWTPs

Based on GRvOTU clustering, 21 RNA vOTUs were identified as putative human

RNA viruses, including 8 *Mamastrovirus*, 7 *Norovirus* and 3 *Rotavirus* (Fig. 5a).

These putative human RNA viruses were all known to potentially cause gastroenteritis

in humans. Absolute abundance estimations in the 47 WWTP samples from

Switzerland indicated that human RNA vOTUs were significantly reduced during the

treatment process, from influent to effluent (Fig. 5b). *Mamastrovirus* and *Rotavirus*

vOTUs were consistently detected in nearly all influent and sludge samples in

Switzerland, with an absolute abundance peak of $2.01 \times 10^{1\square}$ copies/liter (AD-

IN_141867 in TG-AS) for *Mamastrovirus* and $8.42 \times 10^{\square}$ copies/liter (FR-

INF_306656 in AH-DS) for *Rotavirus*. While the absolute abundance of

Mamastrovirus vOTUs did not differ significantly between influent and sludge,

Rotavirus vOTUs exhibited significantly higher concentrations in sludge (Mann-

Whitney U test, $P < 0.05$). In contrast, these two viral species exhibited lower

prevalence in 277 influent samples from the United States, with *Mamastrovirus*

detected in 34 samples and *Rotavirus* in only 5 samples. Norwalk virus was detected

in approximately one-third of influent samples in both Switzerland and the United

States.

RdRp-based phylogeny was also employed to identify potential human RNA

viruses. In addition to the human RNA vOTUs identified based on genome homology

match, three novel potential human RNA viruses were discovered through

phylogenetic analysis (Extended Data Fig. 7). IL13_9334 and KB7_72351 appeared

to be potential novel species within the *Husavirus* genus. Meanwhile, the RdRp of

TU-EFF_201332 was positioned between two *Nodaviridae* proteins (Extended Data

Fig. 7), which were recovered from human plasma samples and have been identified

as potential zoonotic viruses[30].

To detect human RNA viruses more sensitively, we searched clean RNA reads

from metatranscriptomes against the NCBI Virus Database with several evaluation

steps (See Methods). A total of 50 human RNA viral species were identified in the

global WWTP metatranscriptomes (Extended Data Fig. 8). SARS-COVID-2 was

detected in 151 (54.5%) influent samples collected between July 2020 and August

2021 from USA. Five other *Coronaviridae* viral species were detected, including

Alphacoronavirus 1, *Betacoronavirus*, Human coronavirus HKU15, 229E, and NL63

(Extended Data Fig. 8). Influenza and Hepatitis viruses were also detected in some

samples (Extended Data Fig. 8). Reads-based diversity analysis separated the

community structure of human RNA viruses in wastewater influent samples from

Switzerland and USA (Supplementary Fig. 3), revealing the geographical difference

of prevalent viruses and possibly health status between the corresponding populations.

Ecological roles of RNA viruses in the WWTPs revealed by auxiliary metabolism

To explore the auxiliary metabolism of RNA viruses, a total of 160 RNA viral contig-

encoding genes were identified as auxiliary metabolic genes (AMGs) through a series

of bioinformatics procedures including gene annotation, authenticity evaluation, and

3D protein structure annotation (Supplementary Table 6). The most prevalent

functional types of AMGs were peptidases or proteases (n = 28), followed by ATP-

binding proteins (n = 24). Both prokaryotic (n = 9) and eukaryotic (n = 7) ribosomal

proteins (RPs) were identified in RNA viral genomes. For instance, AH-IN_3822,

which encoded a 60S RP, was classified within *Botourmiaviridae*, a family known to

infect fungi or plants (Fig. 6). Phylogenetic analysis of this 60S RP suggested that it

was likely acquired from fungal *Mucoromycota* during infection (Supplementary Fig.

4). WT-DS_133589, which encoded a 40S ribosomal protein, was associated with the

new megataxon (Fig. 6 and Supplementary Fig. 5). BE-IN_10020 and AD-IN_243634,

both from the bacteriophage family *Fiersviridae*, encoded 50S and 30S RPs,

respectively (Fig. 6). Phylogenetic analyses indicated that BE-IN_10020 may acquire

50S RP from *Faecalibacterium* within the phylum *Bacillota*, while AD-IN_243634

may acquire 30S RP from *Chitinibacteraceae* within the phylum *Pseudomonadota*

(Supplementary Fig. 6-7). Five AMGs related to cellular respiration were identified in

the RNA viral genomes, including two glyceraldehyde-3-phosphate dehydrogenase,

one citrate synthase, one cytochrome C551, and one ribose-phosphate

pyrophosphokinase. One glyceraldehyde-3-phosphate dehydrogenase, involved in

glycolysis, was encoded by IL33_2153 from bacteriophage family *Atkinsviridae*,

while the citrate synthase gene, involved in the TCA cycle, was encoded by MN-

EF_134878 from bacteriophage family *Steitzviridae*, implying a prokaryotic origin for

these viral genes (Fig. 6). Additionally, AMGs involved in various other metabolic

pathways were also identified, including those related to nucleotide metabolism (n =

9), RNA modification (n = 7), amino acid metabolism (n = 7), and the metabolism of

cofactors and vitamins (n = 6).

Compared with DNA viruses, interpreting the expression activity of RNA viral

genes in metatranscriptomes is inherently challenging because RNA viral genomes

and transcripts were interwoven. Here, the abundance comparison between AMG and

RdRp within the viral genome in their originating sample was used to indicate the

potential expression of AMG. An AMG was considered potentially active if its

relative abundance (Tmean) exceeded that of the RdRp gene by more than fivefold

(Tmean(AMG) – Tmean(RdRp) > 5×). A total of 19 AMGs were inferred to be

potentially expressed in their respective originating samples (Supplementary Table 7).

The expressed AMGs included three RPs, two ribonucleases, one elongation factor,

and one RNA-binding protein, whose potential expression implied that their hosting RNA viruses may utilize AMGs to enhance transcription, translation, and RNA processing (Supplementary Table 7). Notably, nine unique M15 family peptidases were identified in WWTP RNA viral genomes, seven of which exhibited potential expression activity (Supplementary Table 7). The viruses encoding M15 family peptidases were all derived from the base-*Cystoviridae* clade, a novel bacteriophage group identified through SD motif analysis. In summary, the identified AMGs and their expression patterns point to the dynamic role of RNA viruses in manipulating host cellular machinery and imply multiple virus-host interactions.

Discussion

This study represents a significant advancement in understanding RNA viruses within wastewater treatment plants (WWTPs) by constructing the most extensive catalog of RNA viruses in these systems to date. The establishment of the WWTP RNA Viruses Database (WRVirDB) expands the phylogenetic and functional diversity of RNA viruses known in engineered systems, offering profound implications for public health monitoring and disease surveillance through wastewater-based epidemiology (WBE).

Advancements in RNA Virus Diversity and Phylogeny Our metatranscriptomic compendium has refined the global RNA-dependent RNA polymerase (RdRp) phylogenetic tree (Supplementary Text 3), providing critical insights into previously unidentified RNA viral clades, particularly RNA bacteriophages. Historically, studies on RNA viruses in either engineered or animal and human systems have focused on human pathogens[31-33]. However, our findings reveal that WWTPs harbor a diverse and abundant population of prokaryotic RNA viruses, which likely influence the microbial communities and processes within these systems (Fig. 2a and Extended Data Fig. 5).

The high diversity of putative prokaryotic RNA viruses is plausible, as bacteria are the dominant wastewater microbes and key drivers to the success of WWTPs [29, 34]. Notably, alongside well-characterized RNA phage clades such as *Leviviricetes* and *Cystoviridae*, we identified novel clades, including those within *base-Cystoviridae*, *Picobirnaviridae*, *Partitiviridae*, and *base-Howeltoviricetes* (Extended Data Fig. 6 and Fig. 3). This highlights the underappreciated ecological significance and underestimated diversity of RNA phages in WWTPs.

The high abundance of RNA phages in engineered systems and soil environments (Supplementary Fig. 8) is likely driven by the composition of host organisms, underscoring WWTPs as pivotal ecosystems for uncovering RNA phage diversity. These discoveries suggest that RNA phages may play critical roles in regulating microbial communities and maintaining system stability, positioning WWTPs as key environments for advancing our understanding of microbial ecology that can enlighten us on enhanced wastewater microbiome engineering and wastewater-based surveillance as further discussed.

Functional Contributions of RNA Viruses   Beyond expanding our understanding of RNA virus diversity, this study reveals their functional importance, particularly through the discovery of auxiliary metabolic genes (AMGs). Virus-encoded AMGs, widely studied in dsDNA viruses for their roles in biogeochemistry [25, 35, 36], are less understood in RNA viruses. Our findings demonstrate that RNA viruses encode a wide array of AMGs of both prokaryotic and eukaryotic origin (Supplementary Table 6). These genes, likely acquired through horizontal gene transfer during infection, indicate not only the biochemical roles of RNA viruses in influencing host metabolic pathways, energy flow, and nutrient cycling within WWTPs, but also potential virus-

host associations. These novel insights emphasize the significant, albeit

underexplored, ecological functions of RNA viruses in engineered environments like

the eco-friendly WWTPs.

Implications for Wastewater-Based Surveillance   This study highlights the

potential of metatranscriptomics-based methods for high-throughput detection of the

pool of animal and prokaryotic RNA viruses, generally supporting wastewater-based

surveillance (WBS) an emerging tool for tracking infectious diseases [19, 37]. Unlike

traditional RT-PCR-based methods, which are limited to specific pathogens,

metatranscriptomics enables the simultaneous detection of a broad spectrum of RNA

viruses, including novel species which expand the human RNA virus database (Fig. 5

and Extended Data Fig. 7-8). Our detection of SARS-CoV-2 and other human RNA

viruses responsible for gastroenteritis, influenza, and hepatitis at either the read or

viral operational taxonomic unit (vOTU) levels demonstrates the application potential

of this broad-spectrum omics approach. Additionally, spatial-temporal analysis of

human RNA viruses in WWTPs provides critical insights into their distribution,

dynamics, and persistence throughout the wastewater treatment process.

Particularly, absolute quantification of human viral loads in WWTPs improves the

accuracy of abundance trends compared with relative abundance profiling and

enhances the reliability of health assessments [29]. For example, fourteen human RNA

vOTUs, including *Mamastrovirus*, *Norovirus*, *Rotavirus*, *Husavirus*, and *Sapovirus*,

were detected in WWTP samples from Switzerland using quantitative

metatranscriptomcis with spiked mRNA internal standards (Fig. 5b), with the median

concentrations of the summed vOTUs in influent, sludge, and effluent samples

reached $1.43 \times 10^\square$, $1.76 \times 10^\square$, and $3.48 \times 10^\square$ copies/L, respectively. These

findings demonstrate the value of metatranscriptomics-based WBS not only in monitoring public health but also in providing early warnings for infectious diseases, supporting local and global health initiatives.

Research Significance and Concluding Remark   This comprehensive exploration of RNA viruses in WWTPs achieves several major breakthroughs. First, it disentangles the high diversity and novelty of prokaryotic RNA viruses in these systems. Second, it emphasizes the biochemical contributions of RNA virus-encoded AMGs, which may influence microbial and ecosystem-level processes. Third, it elucidates the distribution patterns of human RNA viruses, offering insights into their implications for public health management. However, the global diversity of RNA viruses in WWTPs warrants further exploration. Limitations include reliance on RdRp as a universal marker gene, potentially missing non-RdRp coding fragments of RNA viruses, and the constraints of current sequencing and bioinformatic methodologies. Despite these challenges, the genomic and functional data generated provides a valuable resource for future research. In conclusion, this study underscores the ecological and public health significance of RNA viruses in WWTPs. By highlighting their roles as both ecological drivers and health indicators, it positions RNA viruses as critical components of global wastewater and other engineered ecosystems and offers a foundation for advancing microbial ecology, public health, and disease surveillance methodologies.

Methods

Metatranscriptome collection

To build the first catalogue of RNA viruses of global WWTPs, a total of 557 available

metatranscriptomes of WWTP samples were downloaded from the public databases

including National Center for Biotechnology Information Sequence Read Archive

(NCBI SRA), Metagenomic for Rapid Annotations using Subsystems Technology

(MG-RAST) and National Genomics Data Center (NGDC) to identify RNA viruses.

These metatranscriptomic samples included 289 influent, 80 denitrification sludge, 61

activated sludge, 26 effluent, and 101 anaerobic digester sludge samples from 9

countries (Supplementary Table 1).   The 47 samples from Switzerland were

sequenced at a similar depth (8.26 ± 1.90 G per sample) and were spiked with mRNA

internal standards in the previous study[29], which enabled the absolute concentration

estimation of RNA viruses in these samples. The detailed information for these public

metatranscriptomes were listed in Supplementary Table 1.

Quality control, metatranscriptomic assembly and gene prediction

The raw sequencing reads were trimmed by Trimmomatic (v0.39)[38] with default

parameters. MEGAHIT[39] has been regarded as a successful assembly tool for

metatranscriptomes with efficient memory and computational time[10]. The trimmed

reads of each metatranscriptome were assembled into contigs using MEGAHIT

(v1.1.3)[39] with default parameters. Only contigs > 1 kbp were used for the downstream

analysis. ORFs were predicted from assembled contigs using Prodigal (v2.6.3)[40] with

option "-p meta".

Identification and evaluation of RNA viruses

The methodology of RNA virus identification referred to a previous study which

systemically searched RNA viruses in the global ocean metatranscriptomes[10]. In detail,

RNA viruses were identified by iterative HMM searches against virus RdRp HMM

profiles (Extended Data Fig. 1). The initial 65 RdRp HMM profiles were downloaded

from (doi:10.5281/zenodo.5731488)[10]. For every iteration, all the protein sequences

predicted from the assembled contigs, along with the public virus RdRp protein

sequences from NCBI Genbank release 253 (n = 3,694,787), and recent RNA virus

studies from ocean (n = 44,828[10] and n = 4,593[11]) and global metatranscriptomes from

diverse ecosystems (n = 329,202)[12] were searched against the RdRp HMM profiles by

hmmsearch (v3.3.2)[41] with the option "-A", which supported to trim the portion

aligned to the RdRp HMM profile. The hit protein sequences with a bit-score ≥ 30

and with ≥ 70% aligned portions to the best RdRp HMM profiles were retained. The

aligned portions of hit protein sequences were clustered into updated RdRp HMM

profiles by vFam pipeline (version February 2014)[42] with default parameters. The

RdRp profile HMMs were generated and updated for eight iterations to meet the near

saturated number of identified viral RdRps. A total of 56086 putative virus RdRps and

455 RdRp HMM profiles were identified after eight search-and-updated iterations.

The contigs encoded putative RdRps were regarded as putative RNA viral contigs.

To evaluate the authenticity of the putative RNA viruses, false-positive RNA

viruses were filtered by 1) a competitive hmmsearch approach[10] and 2) mapping

against corresponding metagenomic datasets. First, 56086 putative RdRps were

searched against the database merged by PfamA (v33.0)[43] HMM profiles and 455

RdRp HMM profiles constructed in this study. If the putative RdRp showed a best

match to a non-RdRp HMM profile, it was considered as a false-positive virus RdRp.

All of the 56086 putative RdRps passed the competitive hmmsearch test, and they all

showed a best match to the RdRp HMM profile. Then, based on the hypothesis that

RNA viruses (under *Orthornavirae* kingdom) would not be present in the DNA

fraction, the occurrence of putative RNA viral contigs in the metagenomes were

examined and used to evaluate false-positive RNA viruses. We downloaded additional

134 metagenomes originated from the samples same as part of the WWTP

metatranscriptomes. The raw DNA sequences were mapped to the putative RNA viral

contigs by Bowtie2 (v2.3.5.1)[44] in "--very-sensitive" mode. If >20% of the positions

in the putative RNA viral contig were mapped by DNA reads in any metagenome, this

RNA viral contig was considered as a false-positive RNA virus. In total, 165 putative

RNA viral contigs were identified as false-positive RNA viruses and were discarded

in this step.

To examine the completeness of virus RdRp domains, RNA viral contigs were

translated into protein sequences by transeq (EMBOSS:6.6.0.0)[45] with all six frames

and the standard translational code. The translated protein sequences were searched

against the 455 RdRp HMM profiles once more, and the sequences with a bit-score $\geq$

30 and with $\geq$ 90% aligned portions to the best RdRp HMM profiles were considered

as RNA viruses containing "complete" (or "near-complete") RdRp domains. Together,

55900 RNA viral contigs were identified from 557 WWTP metatranscriptomes,

including 44399 RNA viral contigs contained "complete" RdRp domains.

To check the alternative translation code usage of RNA vOTUs, the ORFs of

RNA vOTUs were predicted under all available codes with Prodigal (v2.6.3)[40]. The

codon producing the longest average length of ORFs was chosen for the

corresponding RNA vOTU.

RNA virus clustering

Clustering of WWTP RNA virus operational taxonomic units

The RNA viral contigs were clustered into RNA vOTUs of 90% ANI across 80%

of AF (parameters referred to reference[10], the scripts used for RNA vOTU clustering

were downloaded from https://bitbucket.org/berkeleylab/checkv/src/master/scripts/).

The clustered RNA vOTUs were pragmatically at approximately species level[10]. In

total, 55900 RNA viral contigs were clustered into 11414 RNA vOTUs, including

9276 RNA vOTU representatives contained "complete" RdRp domains.

Clustering of WWTP and publicly known RNA viruses

To evaluate the expansion of RNA viral diversity in this study, RNA viruses

identified from the WWTP metatranscriptomes were clustered alongside publicly

known RNA viruses at both the genome and RdRp levels. Public RNA viral genome

and RdRp sequences were retrieved from NCBI Virus database (accessed on August

13, 2024) and recent studies of RNA viruses in oceanic[10, 11], soil[13, 16], and global

metatranscriptomes from diverse ecosystems[12]. Based on the global RdRp

phylogenetic tree, no RNA vOTUs from WWTPs were assigned to the *Coronaviridae*

family. Due to the clustering efficiency considerations, reference sequences of SARS-

COVID-2 were excluded from the clustering analysis. Genome-level clustering was

performed using the same parameters as for WWTP vOTU clustering (90% ANI and

80% AF). RdRp-level clustering was conducted using MMseqs2[46] with the parameters

'--min-seq-id 0.9 -c 0.333 -e 0.1 --cov-mode 1 --cluster-mode 2'[12]. The resulting RNA

viral genome clusters and RdRp clusters were termed GRvOTU and RCR90,

respectively, and were classified into three categories, 'novel', 'shared', and

'reference', based on the data sources of their member sequences. All public RNA

viral genomes were also subjected to genome-level clustering (90% ANI and 80% AF)

to establish the background reference for known RNA viral diversity.

RdRp-based taxonomy classification

The taxonomic megataxon of RNA vOTUs were determined by a network-based

clustering method[10]. The public RdRp reference protein sequences that matched to

RdRp HMM profiles after eight hmmsearch iterations were collected and pre-

clustered at 50% amino-acid identity by Uclust (v8.0.1623)[47] with options "usearch –

cluster_fast -id 0.5 -sort length", and resulted in 30,504 public RdRp centroids. Then,

30,504 public RdRp centroids along with 9,276 WWTP RNA vOTUs containing

"complete" RdRp domains were merged and used for clustering megataxon of RNA

viruses. Pairwise comparisons of these merged RdRp sequences were conducted by

BLASTp (v2.11.0) with options "-gapopen 9 -gapextend 1 -word_size 3 -threshold

10". E-values were extracted and negative-log10-transformed by "mcxload" in MCL

(v14-137)[48] with options "--stream-mirror --stream-neg-log10 -stream-tf 'ceil(200)' ".

Transformed e-values were used for MCL[48] network clustering with option "-I 0.9" to

establish RdRp megataxon (approximately at class to order level). In total, 22 RdRp

megataxon were clustered and labeled by the public reference RdRp centroids with

known taxonomic classification within the megataxon. The accuracy of the label for

each megataxon was calculated as the number of public RdRp centroids affiliated

with the taxonomy divided by the total number of known public centroids (those with

labels at phylum level at least) in the megataxon. One megataxon (megataxon 22, n =

68) were mainly consisted of RdRps identified in this study (n = 56) with only two

known reference RdRps. This megataxon was regarded as novel megataxon and was

not labeled (Supplementary Table 4).

RdRp-based phylogenetic analysis

Establishment of the global viral RdRp tree

The public RdRp centroids, WWTP viral RdRps containing "complete" RdRp

domains, along with the group II introns and non-LTR retrotransposons were aligned

and used to generate the approximate global tree for viral RdRps. In detail, the protein

sequences of group II introns and non-LTR retrotransposons were downloaded from

Genbank Release 253. The RdRps within megataxon 18 were affiliated with viral

families of *Birnaviridae* and *Permutotetraviridae*, which encoded conserve motifs

with the order permutation "C-A-B" instead of the canonical permutation "A-B-C".

Such RdRp permutations can make RdRp sequences unalignable with those of the

canonical configuration[11]. Therefore, the RdRp sequences of megataxon 18 (n = 388)

were excluded from the analysis of global RdRp tree construction and analyzed

individually (Supplementary Fig. 6). In total, public RdRp centroid sequences (n =

30,217), group II introns (n = 950) and non-LTR retrotransposons (n = 83), and

WWTP RNA vOTUs containing "complete" RdRp domains (except for RdRps of

megataxon 18, n = 9,175), were aligned by MUSCLE (v5.1)[49] under "-super5" mode.

The aligned sequences were trimmed by Trimal (v1.4.rev15)[50] with sites having more

than 20% gaps removed. The alignment of trimmed sequences was used to construct

an approximate maximum likelihood tree by the FastTree (v2.1.11)[51] with options "-

wag -gamma". The constructed tree was illustrated in iTOL (v6)[52]. The tree was

rerooted between RdRps and RTs. The family-level taxonomic classification for

RdRps was assigned based on the placement of the sequences in the tree, requiring the

RdRps fell within a clade to be assigned to a specific family. The family-level

taxonomy classification should be consistent in both methods of megataxon clustering

and phylogenetic analysis.

Phylogenetic analysis for established RNA bacteriophage families

To construct phylogenetic trees for known RNA bacteriophage families, RdRp

representatives from RCR90 clusters occurred in the engineering systems were

selected for phylogenetic analysis. Sequence alignment was performing by MUSCLE

(v5.1)[49] and phylogenetic analysis was conducted with IQTREE (v2.1.4-beta)[53] using

parameters '-bb 1000'. The resulting trees were illustrated in iTOL (v6)[52].

Host prediction of RNA vOTUs

Virus-host relationships were predicted using the following approaches. (i) taxonomy-

based prior known host range, (ii) CRISPR spacer match, (iii) SD motif prediction, (iv)

species-level clustering, (v) RdRp phylogeny, and (vi) Reads-based alignment. The

second and third approaches were designed to identify RNA phages, while the fourth,

fifth, and sixth approaches aimed to identify human RNA viruses.

For taxonomy-based host range, the taxonomic classification for RNA vOTUs

(See RdRp-based taxonomy classification and RdRp-based phylogenetic analysis)

were used to retrieve known host range of each family determined by ICTV proposals.

Type-III and Type-□ CRISPR systems can target RNA and appear possibilities to

defend against RNA bacteriophages[54]. To identify potential RNA phages by CRISPR

spacer match, all RNA viral contigs were compared to prokaryotic CRISPR spacer

sequences to predict virus-prokaryote associations. CRISPR spacers were predicted

for all > 10 kb reference bacterial and archaeal genomes downloaded from NCBI

Refseq database (https://ftp.ncbi.nlm.nih.gov/genomes/refseq/) with CRT (2.0rev0)[55].

We also predicted CRISPR spacers from metagenomic contigs assembled from 134

corresponding WWTP metagenomes. The DNA contigs were assembled by WWTP

metagenomes using metaSPAdes (v3.13.0)[56] with default parameters. The predicted

CRISPR spacers were removed if (i) they included low complexity (consisting of 4–

6bp repeat motifs) or ambiguous bases, (ii) they were ≤ 20bp [25]. To link RNA viral

contigs to predicted CRISPR spacers, only BLASTn-short hits with 0 or 1 mismatch

over the whole spacer were considered. The existence, type, and location of CRISPR

systems were identified in potential host genomes using GenPept files for reference

hosts from the NCBI RefSeq database and gene annotations for WWTP metagenomic

assemblies. The ORFs from host DNA contigs were predicted by Prodigal (v2.6.3)[40]

and annotated by HHblits (v3.3.0)[57] with options "-n 1 -e 0.001" using

UniRef30_2023_02 database. Only hits with probability >90% were retained. The

predicted virus-host pairs were considered if the matched spacer array was next to an

RT-encoding Type III or Type VI CRISPR system in the host genome.

To identify potential RNA phages by SD motif prediction, the information of

"rbs_motif" for ORFs of RNA vOTUs with the genetic code identified as above (See

Identification and evaluation of RNA viruses) was extracted from Prodigal output

files. The canonical SD motifs were identified based on the reference list from the

previous study[12]. To identify potential novel prokaryotic RNA viral clades, the viral

contigs from two public datasets (the Global Ocean Virome[10] and The RNA Viruses in

Metatranscriptomes Discovery Project[12]) were recruited and also checked for the

presence of SD motif as described above. The proportion of SD motif-associated

ORFs for each leaf in the global RdRp tree was calculated as the number of SD-

carrying ORFs divided by the total number of ORFs with a true start ("start_type"

different from "Edge" in Prodigal output) within the WWTP RNA vOTU (or the

public centroid). The phylogenetic trees of potential prokaryotic RNA viral clades

indicated by SD motifs were pruned from the global tree.

To identify potential human RNA viruses by species-level clustering, the

GRvOTU clustering results (See RNA virus clustering) was used to identify human

RNA viruses. If a viral cluster from GRvOTU dataset included at least one known

human RNA virus, the WWTP RNA viral contig within this cluster was assigned to

the same species.

To identify potential human RNA viruses by RdRp phylogeny, the public human

viral RdRps and WWTP RNA vOTUs containing "complete" RdRp domains were

aligned and used to generate a phylogenetic tree. Public human viral RdRp domains

retrieved from NCBI genbank release 253 (See "Identification and evaluation of RNA

viruses") were clustered at 95% amino-acid identity to reduce sequence redundancy

by Uclust before phylogenetic analysis. Subsequent phylogenetic analysis was

performed in the same way as the global RdRp tree construction. The potential human

RNA RdRp was identified based on the placement of the sequence in the tree,

requiring the RdRp fell within a clade to be assigned to a known clade of human RNA

viruses.

To identify potential human RNA viruses by reads-based alignment, all the clean

RNA reads from 557 WWTP metatranscriptomes were BLASTn against complete

nucleotides of human RNA viruses in NCBI Virus database. The reads matched to

reference human RNA viral genomes with $\geq$ 90% identity along $\geq$ 80% reads length

were considered as potential fragments of the related human RNA viral species. The

matched RNA reads were then checked against the NCBI nt database using BLASTn.

Reads that matched reference cellular organisms (prokaryotes and eukaryotes) with

identity$\geq$ 90% along $\geq$ 80% read length were discarded. The relative abundance of

detected human RNA viral genomes was normalized to RPKM (Reads Per Kilobase

of transcript per Million mapped reads) based on reference genome length and the

size of metatranscriptome. The relative abundance of human RNA viral species was

calculated by summing the RPKM value of the viral organisms belonging to the viral

species. Only human RNA viral species with a total read count > 10 in the samples

were retained for analyses.

A mock metatranscriptomic dataset, composed of Illumina sequencing data from

bacterial transcriptomes and human RNA viral genomes, was generated to further

assess the feasibility of reads-based detection of human RNA viruses[58]. Eight Illumina

transcriptomic datasets from pure-culture bacteria isolated from WWTP samples were

downloaded from NCBI SRA Database (Supplementary Table 9). The mock Illumina

sequencing data for eight human RNA viruses were generated by Mason (v2.4.0)[59]

with different sequencing depth (from 20 to 40000 reads). The merged mock

metatranscriptome has 172.5 million reads in total with 72.3k simulated human RNA

viral reads (Supplementary Table 9). The mock metatranscriptome was also subjected

to the same BLASTn alignment steps against the NCBI Virus and NCBI nt databases.

The alignment results were used to evaluate the accuracy of reads-based detection of

human RNA viruses. In the mock community, 99.99% of reads belonging to the

human RNA viral genome were successfully recovered, and only 28 sequences from

bacterial transcriptomes were incorrectly recruited (Supplementary Table 9),

suggesting the reads-based examination for human RNA viruses from global WWTP

metatranscriptomes in this study is confident and accurate.

Functional gene annotation of RNA viral contigs

The functional annotation of predicted ORFs with the genetic code identified as above

(See Identification and evaluation of RNA viruses) was determined by one iteration of

HHblits (v3.3.0)[57] with options "-n 1 -e 0.001" using UniRef30_2023_02 database

(https://wwwuser.gwdg.de/~compbiol/uniclust/2023_02/UniRef30_2023_02_hhsuite.t

ar.gz). Only hits with >90% probability score were used for functional annotation. The

predicted ORFs were also searched against the NCBI nr database by BLASTp with a

bitscore > 50 were retained.

Identification and phylogenetic analysis of RNA virus-encoded AMGs

The RNA virus-encoded genes were regarded as AMGs if they met the following

requirements: (i) both HHblits best hit in the UniRef database and BLASTp best hit in

the NCBI nr database were originated from cellular organism. (ii) No DNA reads

from the 134 corresponding WWTP metagenomes could be perfectly mapped to the

AMG regions of the RNA viral contigs. (iii) The 3D protein structure annotation was

consistent with the homology-based annotation. The 3D protein structures of probable

AMG sequences were predicted with ESMFold[60], annotated with Foldseek

(9.427df8a)[61] (probability threshold set to 0.9), and visualized in PyMOL (v3.0.5)[62].

To investigate the probable evolutionary origin of RNA virus-encoded RPs, sequences

similar to four types of RPs were recruited from the NCBI nr database. The viral and

reference RP sequences were aligned by MUSCLE (v5.1)[49] and analyzed by IQTREE

(v2.1.4-beta)[53]. Phylogenetic trees were visualized in iTOL (v6)[52].

Quantitative profiling and diversity analysis of RNA vOTUs

Quantification of RNA vOTUs

To calculate relative abundance of RNA vOTUs in each sample, clean reads from

each metatranscriptome were first trimmed off polyA and polyp stretches using bbduk

(v37.62) (https://jgi.doe.gov/data-andtools/bbtools/) with options

"literal=AAAAAAAAAA,TTTTTTTTTT, hdist=2 k=10 minlength=30, ktrim=r

restrictright=20; ktrim=l restrictleft=20" for three times. WWTP RNA viral contigs

were also trimmed off polyA and polyT stretches. PolyA/T-trimmed clean reads in

each metatranscriptome were individually mapped against polyA/T-trimmed RNA

viral contigs using Bowtie2 (v2.3.5.1)[44] with options "--local -D 20 -R 3 -N 0 -L 16 -i

S,1,0.50 --very-sensitive --non-deterministic -I 0 -X 2000"[10]. The relative

abundance of RNA vOTUs in each sample was calculated by CoverM (v0.6.1,

https://github.com/wwood/CoverM) with options "--min-read-percent-identity 90 --

min-read-aligned-percent 75 -m trimmed_mean --min-covered-fraction 30", which

recruited reads that mapped with ≥ 90% identity ≥ 75% of the read length and

restricted RNA viral contigs should be covered ≥ 30% length horizontally by mapped

reads. The calculated Tmean abundances for RNA viral contigs were normalized by

the number of mapped reads as filtered by CoverM. The final relative abundance of

RNA vOTUs were calculated by summing the normalized abundances of contigs

belonging to the RNA vOTUs.

Quantification of AMGs

To determine the relative abundance of RNA viral ORFs in its originated sample,

the polyA/T-trimmed RNA clean reads from each metatranscriptome were mapped to

RNA viral ORFs individually using the '--very-sensitive' mode with Bowtie2

(v2.3.5.1)[44]. The relative abundance of each ORF was then calculated as the trimmed

mean abundance, determined using CoverM (v0.6.1) with the options "--min-read-

percent-identity 100 --min-read-aligned-percent 70 -m trimmed_mean --min-covered-

fraction 70".

Due to the intermingling of RNA viral genomes and transcriptional products in

metatranscriptomes, determining whether RNA viral genes are actively expressed is

challenging. However, the potential expression activity of AMG can be inferred by

comparing the relative abundance with RdRp within its respective viral genome. An

AMG was considered potentially active in its originated sample if: Tmean (AMG) –

Tmean (RdRp) > 5×. It was important to note that these bioinformatics steps identify

AMGs more likely to be actively expressed, though sequencing inaccuracies,

sampling size fractions and gene expression patterns (e.g., actively expressed AMGs

with lower expression levels than RdRp cannot be identified) might affect the

prediction.

Absolute quantification of RNA vOTUs in mRNA internal standards spiked samples

To estimate absolute abundance of RNA vOTUs in 47 metatranscriptomes from

Switzerland, the statistical information of spiked mRNA internal standards as referred

to reference[29] was used in this study. The absolute abundance of RNA vOTU in each

sample was measured in both "copies/g-VSS" and "copies/L" rank using the equation

1 and 2, respectively.

$$\text{Absolute abundance (copies/g-VSS)} = \frac{N_{spiked\,RIS}}{m_{biomass}} \times \frac{N_{mapped\,reads}/L_{vOTU}}{N_{RIS\,reads}/L_{RIS}} \quad (1)$$

$$\text{Absolute abundance (copies/L)} = \frac{N_{spiked\,RIS}}{V_{sample}} \times \frac{N_{mapped\,reads}/L_{vOTU}}{N_{RIS\,reads}/L_{RIS}} \quad (2)$$

$N_{spiked\,RIS}$ was the copy number of spiked mRNA internal standards in the sample,

$m_{biomass}$ was the mass of collected volatile suspended solids of the sample, $N_{mapped\,reads}$

was the number of reads mapped to the RNA vOTU in the metatranscriptome, L_{vOTU}

was the length of the RNA vOTU, $N_{RIS\ reads}$ was the number of reads mapped to the

mRNA internal standards in the metatranscriptome, L_{RIS} was the length of the mRNA

internal standards.

Similarly, the above formula was also applied to estimate the absolute

concentration of detected human RNA viruses in 47 samples from Switzerland at

reads level, using the length of matched reference genome to replace the length of

vOTU in the formula.

Diversity analysis of RNA vOTUs

The α-diversity of RNA vOTUs in the samples was measured with the Shannon's

index calculated by function "diversity" in "vegan" package[63] in R. The β-diversity of

RNA vOTUs in the samples was measured using the Bray-Curtis dissimilarity method

calculated by function "vegdist" in "vegan" package[63] in R, using the log-transformed

relative abundance table of RNA vOTUs in the samples as input. Then, the Non-

Metric Multidimensional Scaling (NMDS) analysis was performed using function

"metaMDS" in R package "vegan"[63]. The analysis of Similarity (ANOSIM) was

performed using function "anosim" in R package "vegan". The difference of Shannon

index H and absolute concentration of RNA vOTUs between pairwise compartments

from Switzerland was determined by the Mann–Whitney U test using the function

"wilcox.test" with option "paired = FALSE" in R.

Host taxonomy, host-virus abundance correlation and genome contents of base-

Howeltoviricetes bacteriophage clade

Several viral contigs from vOTU FD-DS_117671, belonging to the base-*Howeltoviricetes* phage clade, matched CRISPR spacer identified in both bacterial genomes from the NCBI RefSeq database and WWTP metagenomic contigs. Two potential host genomes from the NCBI RefSeq database were both classified as *Aliarcobacter Cryaerophilus*. To determine the taxonomic classification of two potential host contigs (TG-IN_1259 and ZR-IN_6153) from WWTP metagenomic assemblies, metagenome-assembled genomes (MAGs) were reconstructed from the respective samples (TG-IN and ZR-IN) using metaBAT2 (v2.12.1)[64]. MAG completeness and contamination were evaluated with CheckM (v1.0.12)[65], and taxonomy was assigned using GTDB-tk (v2.1.1)[66]. ZR-IN_6153 was successfully binned into a MAG with 69.35% completeness and 5.69% contamination, and classified as *Aliarcobacter sp.*. Since TG-IN_1259 could not be binned into any MAG, its taxonomy was determined using CAT (v5.2.3)[67].

The viral-host abundance of vOTU FD-DS_117671 was examined across two sample series: i) 47 samples from Switzerland, representing virus-host abundance changes throughout the wastewater treatment process. ii) 47 time-series samples from Luxembourg, capturing the temporal dynamics of virus-host abundance. From the Luxembourg dataset, only the first 47 of 68 samples were included in the correlation analysis because, according to the original metadata, these samples were collected at weekly intervals, ensuring consistent temporal resolution, and sequenced in the same batch, reducing technical variability. The FD-DS_117671 vOTU comprised 53 contigs. As FD-DS_117671 originated from the Swiss sample series, it was designated as the representative sequence for this vOTU in the Swiss dataset. In the Luxembourg sample series, SF52_112815 was selected as the representative sequence for this vOTU, as it was the longest contig recovered from the Luxembourg dataset. To

calculate the relative abundance of representative sequence in the related dataset,

polyA/T-trimmed clean reads in the corresponding metatranscriptomes were mapped

against representative sequence using Bowtie2 (v2.3.5.1)[44] under '--very-sensitive'

mode. The relative abundance of the vOTU representative sequence in each sample

was calculated as the trimmed mean abundance, determined using CoverM (v0.6.1,

https://github.com/wwood/CoverM) with the options "--min-read-percent-identity 90

--min-read-aligned-percent 75 -m trimmed_mean --min-covered-fraction 30" divided

by the size of the metatranscriptome (in Gb). The relative abundance of potential

bacterial host (*Aliarcobacter Cryaerophilus*) in the two datasets were determined by

mOTUs3[68] (v3.0.3). Then, the correlation between host and viral abundance was

calculated by the Spearman correlation coefficient.

In the BE samples from Bern in Switzerland and the SF50-SF52 samples from

Luxembourg, base-*Howeltoviricetes* phage clade showed remarkably high abundance

(abundance proportion > 50%). Therefore, near-complete genomes of the highly

abundant RNA phages may be reconstructed from these samples. For example, the

vOTU TU-EF_29550 from base-*Howeltoviricetes* phage clade accounted for 57.1% of

the total RNA vOTU abundance in the BE-AS sample. However, no RNA viral contig

members from vOTU TU-EF_29550 were assembled in BE-AS sample. To check this

discrepancy, all contigs from the BE-AS sample were aligned to TU-EF_29550 using

BLASTn. A total of 83 contigs were identified >95% identity over >95% coverage

with TU-EF_29550. These contigs were all less than 1 kb so they were not included in

the RNA virus identification step. The fragmented assembly was likely caused by the

microdiversity of the RNA virus. Among these 83 contigs, the longest contig (BE-

AS_5392, 863bp) was selected as the starting sequence for genome extension. BE-

AS_5392 shared 97.7% sequence identity with TU-EF_29550 over the full length.

Based on genome extension method described in the reference[69], BE-AS_5392 was

extended to a final length of 2,313 bp. Similarly, in the sample SF51 from

Luxembourg, the vOTU AH-EF_62580 from base-*Howeltoviricetes* phage clade

accounted for 76.6% of the total RNA vOTU abundance, and the related contig

SF51_6014 was extended to a final length of 2,031bp in the SF51 sample.

Data availability

788 The sequences of all RNA vOTUs and RdRps identified in this study and the detail

789 information of RNA vOTUs were available at https://github.com/emblab-

790 westlake/WRVirDB. The accession number in a publicly available database will be

791 provided during manuscript submission and upon request.

References

1.   Wu, D., Wu, T., Liu, Q. & Yang, Z. The SARS-CoV-2 outbreak: What we know. *International*
Journal of Infectious Diseases **94**, 44-48 (2020).

2.   Nimgaonkar, I., Ding, Q., Schwartz, R.E. & Ploss, A. Hepatitis E virus: advances and challenges.
Nature Reviews Gastroenterology & Hepatology **15**, 96-110 (2017).

3.   Hall, A.J. et al. Norovirus Disease in the United States. *Emerging Infectious Diseases* **19**, 1198-
1205 (2013).

4.   Moreno-Altamirano, M.M.B., Kolstoe, S.E. & Sánchez-García, F.J. Virus Control of Cell
Metabolism for Replication and Evasion of Host Immune Responses. *Frontiers in Cellular and*
Infection Microbiology **9** (2019).

5.   Gualdoni, G.A. et al. Rhinovirus induces an anabolic reprogramming in host cell metabolism
essential for viral replication. *Proceedings of the National Academy of Sciences* **115** (2018).

6.   Dominguez-Huerta G, Zayed A A, Wainaina J M & al., e. Diversity and ecological footprint of
Global Ocean RNA viruses. *Science* **376**, 1202-1208 (2022).

7.   Shi, M. et al. The evolutionary history of vertebrate RNA viruses. *Nature* **556**, 197-202 (2018).

8.   Shi, M. et al. Redefining the invertebrate RNA virosphere. *Nature* **540**, 539-543 (2016).

9.   Kamitani, M., Nagano, A.J., Honjo, M.N., Kudoh, H. & Kümmerli, R. RNA-Seq reveals virus–
virus and virus–plant interactions in nature. *FEMS Microbiology Ecology* **92** (2016).

10.   Zayed A A, Wainaina J M, al., e. & Sullivan, M.B. Cryptic and abundant marine viruses at the
evolutionary origins of Earth's RNA virome. *Science* **376**, 156-162 (2022).

11.   Wolf, Y.I. et al. Doubling of the known set of RNA viruses by metagenomic analysis of an
aquatic virome. *Nat Microbiol* **5**, 1262-1270 (2020).

12.   Neri, U. et al. Expansion of the global RNA virome reveals diverse clades of bacteriophages.
Cell **185**, 4023-4037 e4018 (2022).

13.   Chen, Y.-M. et al. RNA viromes from terrestrial sites across China expand environmental viral
diversity. *Nature Microbiology* **7**, 1312-1323 (2022).

14.   Callanan, J. et al. Expansion of known ssRNA phage genomes: From tens to over a thousand.
Science Advance **6**, eaay5981 (2020).

15.   Zhang, Y.-Z., Chen, Y.-M., Wang, W., Qin, X.-C. & Holmes, E.C. Expanding the RNA Virosphere
by Unbiased Metagenomics. *Annual Review of Virology* **6**, 119-139 (2019).

16.   Starr, E.P., Nuccio, E.E., Pett-Ridge, J., Banfield, J.F. & Firestone, M.K. Metatranscriptomic
reconstruction reveals RNA viruses with the potential to shape carbon cycling in soil. *Proc*
Natl Acad Sci U S A **116**, 25900-25908 (2019).

17.   Ju, F., Xia, Y., Guo, F., Wang, Z. & Zhang, T. Taxonomic relatedness shapes bacterial assembly
in activated sludge of globally distributed wastewater treatment plants. *Environ Microbiol* **16**,
2421-2432 (2014).

18.   Chen, Y., Wang, Y., Paez-Espino, D., Polz, M.F. & Zhang, T. Prokaryotic viruses impact
functional microorganisms in nutrient removal and carbon cycle in wastewater treatment
plants. *Nat Commun* **12**, 5398 (2021).

19.   Karthikeyan, S. et al. Wastewater sequencing reveals early cryptic SARS-CoV-2 variant
transmission. *Nature* **609**, 101-108 (2022).

20.   Yuan, L. & Ju, F. Potential Auxiliary Metabolic Capabilities and Activities Reveal Biochemical
Impacts of Viruses in Municipal Wastewater Treatment Plants. *Environmental Science &*
Technology **57**, 5485-5498 (2023).

21.   Shi, L.D. et al. A mixed blessing of viruses in wastewater treatment plants. *Water Res* **215**,
118237 (2022).

22.   Rothman, J.A. et al. RNA Viromics of Southern California Wastewater and Detection of SARS-
839   CoV-2 Single-Nucleotide Variants. *Applied and Environmental Microbiology* **87** (2021).

840   23.   Guajardo-Leiva, S., Chnaiderman, J., Gaggero, A. & Díez, B. Metagenomic Insights into the
841   Sewage RNA Virosphere of a Large City. *Viruses* **12** (2020).

842   24.   Yousif, M. et al. SARS-CoV-2 genomic surveillance in wastewater as a model for monitoring
843   evolution of endemic viruses. *Nature Communications* **14** (2023).

844   25.   Roux, S. et al. Ecogenomics and potential biogeochemical impacts of globally abundant ocean
845   viruses. *Nature* **537**, 689-693 (2016).

846   26.   Chen, L.X. et al. Large freshwater phages with the potential to augment aerobic methane
847   oxidation. *Nat Microbiol* **5**, 1504-1515 (2020).

848   27.   Yuan, L. et al. Seasonal succession, host associations, and biochemical roles of aquatic viruses
849   in a eutrophic lake plagued by cyanobacterial blooms. *Environment International* **193** (2024).

850   28.   Sun, M., Yuan, S., Xia, R., Ye, M. & Balcázar, J.L. Underexplored viral auxiliary metabolic genes

851   in soil: Diversity and eco-evolutionary significance. *Environmental Microbiology* **25**, 800-810

852   (2023).

853   29.   Ju, F. et al. Wastewater treatment plant resistomes are shaped by bacterial composition,
854   genetic exchange, and upregulated expression in the effluent microbiomes. *ISME J* **13**, 346-
855   360 (2019).

856   30.   Orf, G.S. et al. Metagenomic Detection of Divergent Insect- and Bat-Associated Viruses in
857   Plasma from Two African Individuals Enrolled in Blood-Borne Surveillance. *Viruses* **15** (2023).

858   31.   Stobnicka-Kupiec, A., Gołofit-Szymczak, M., Cyprowski, M. & Górny, R.L. Detection and
859   identification of potentially infectious gastrointestinal and respiratory viruses at workplaces
860   of wastewater treatment plants with viability qPCR/RT-qPCR. *Scientific Reports* **12** (2022).

861   32.   Gonzalez, R. et al. COVID-19 surveillance in Southeastern Virginia using wastewater-based
862   epidemiology. *Water Research* **186** (2020).

863   33.   Woolhouse, M.E.J. & Brierley, L. Epidemiological characteristics of human-infective RNA
864   viruses. *Scientific Data* **5** (2018).

865   34.   Ju, F. & Zhang, T. Bacterial assembly and temporal dynamics in activated sludge of a full-scale
866   municipal wastewater treatment plant. *ISME J* **9**, 683-695 (2015).

867   35.   Gazitua, M.C. et al. Potential virus-mediated nitrogen cycling in oxygen-depleted oceanic
868   waters. *ISME J* **15**, 981-998 (2021).

869   36.   Rosenwasser, S., Ziv, C., Creveld, S.G.V. & Vardi, A. Virocell Metabolism: Metabolic Innovations
870   During Host-Virus Interactions in the Ocean. *Trends Microbiol* **24**, 821-832 (2016).

871   37.   Levy, J.I., Andersen, K.G., Knight, R. & Karthikeyan, S. Wastewater surveillance for public
872   health. *Science* **379**, 26-27 (2023).

873   38.   Bolger, A.M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
874   data. *Bioinformatics* **30**, 2114-2120 (2014).

875   39.   Li, D., Liu, C.M., Luo, R., Sadakane, K. & Lam, T.W. MEGAHIT: an ultra-fast single-node solution
876   for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* **31**,
877   1674-1676 (2015).

878   40.   Hyatt, D. et al. Prodigal: prokaryotic gene recognition and translation initiation site
879   identification. *BMC Bioinformatics* **11**, 119 (2010).

880   41.   Finn, R.D., Clements, J. & Eddy, S.R. HMMER web server: interactive sequence similarity
881   searching. *Nucleic Acids Res* **39**, W29-37 (2011).

882   42.   Tse, H., Skewes-Cox, P., Sharpton, T.J., Pollard, K.S. & DeRisi, J.L. Profile Hidden Markov
883   Models for the Detection of Viruses within Metagenomic Sequence Data. *PLoS ONE* **9** (2014).

884   43.   Finn, R.D. et al. The Pfam protein families database: towards a more sustainable future.
885   *Nucleic Acids Res* **44**, D279-285 (2016).

886   44.   Langmead, B. & Salzberg, S.L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
887   357-359 (2012).

888   45.   Rice, P., Longden, I. & Bleasby, A. EMBOSS: the European molecular biology open software
889   suite[<emboss.pdf>. *Trends in genetics* **16**, 276-277 (2000).

890   46.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the
891   analysis of massive data sets. *Nature Biotechnology* **35**, 1026-1028 (2017).

892   47.   Edgar, R.C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**,
893   2460-2461 (2010).

894   48.   Enright, A.J., VAN, D.S. & Ouzounis, C.A. An efficient algorithm for large-scale detection of
895   protein families. *Nucleic Acids Res* **30**, 1575-1584 (2002).

896   49.   Edgar, R.C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

897   *Nucleic Acids Res* **32**, 1792-1797 (2004).

898   50.   Capella-Gutiérrez, S., Silla-Martínez, J.M. & Gabaldón, T. trimAl: a tool for automated
899   alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973
900   (2009).

901   51.   Price, M.N., Dehal, P.S. & Arkin, A.P. FastTree 2--approximately maximum-likelihood trees for
902   large alignments. *PLoS One* **5**, e9490 (2010).

903   52.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display
904   and annotation. *Bioinformatics* **23**, 127-128 (2007).

905   53.   Nguyen, L.-T., Schmidt, H.A., von Haeseler, A. & Minh, B.Q. IQ-TREE: A Fast and Effective
906   Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*
907   *Evolution* **32**, 268-274 (2015).

908   54.   Makarova, K.S. et al. Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and
909   derived variants. *Nature Reviews Microbiology* **18**, 67-83 (2019).

910   55.   Bland, C. et al. CRISPR recognition tool (CRT): a tool for automatic detection of clustered
911   regularly interspaced palindromic repeats. *BMC Bioinformatics* **8**, 209 (2007).

912   56.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P.A. metaSPAdes: a new versatile
913   metagenomic assembler. *Genome Res* **27**, 824-834 (2017).

914   57.   Steinegger, M. et al. HH-suite3 for fast remote homology detection and deep protein
915   annotation. *BMC Bioinformatics* **20** (2019).

916   58.   Liu, L. et al. High-quality bacterial genomes of a partial-nitritation/anammox system by an
917   iterative hybrid assembly method. *Microbiome* **8** (2020).

918   59.   Berlin, F. Mason — A Read Simulator for Second Generation Sequencing
919   Data.<mason201009.pdf>. *Technical Report* (2010).

920   60.   Lin, Z. et al. Evolutionary-scale prediction of atomic-level protein structure with a language
921   model. *Science* **379**, 1123-1130 (2023).

922   61.   van Kempen, M. et al. Foldseek: fast and accurate protein structure search. *Biorxiv* **2022.02.**
923   **07.479398** (2022).

924   62.   Schrodinger, LLC (2015).

925   63.   Dixon, P. VEGAN, a package of R functions for community ecology. *Journal of Vegetation*
926   *Science* **14**, 927-930 (2003).

64. Kang, D.D. et al. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome
reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

65. Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P. & Tyson, G.W. CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res **25**, 1043-1055 (2015).

66. Chaumeil, P.A., Mussig, A.J., Hugenholtz, P. & Parks, D.H. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).

67. Cambuy, D.D., Coutinho, F.H. & Dutilh, B.E. Contig annotation tool CAT robustly classifies
assembled metagenomic contigs and long sequences. *bioRxiv* (2016).

68. Milanese, A. et al. Microbial abundance, activity and population genomic profiling with
mOTUs2. *Nat Commun* **10**, 1014 (2019).

69. Chen, L.-X., Anantharaman, K., Shaiber, A., Eren, A.M. & Banfield, J.F. Accurate and complete
genomes from metagenomes. *Genome Research* **30**, 315-333 (2020).

Acknowledgements

This work was supported by the Zhejiang Provincial Natural Science Foundation of China under (Grant No. LR22D010001) and the HRHI program 202309010 of Westlake Laboratory of Life Sciences and Biomedicine at Westlake University. The authors thank Mr. Guoqing Zhang and Ms. Xinyu Huang at Westlake University for their helpful suggestions and technical support. We thank the Westlake University HPC Center for computation support.

Author contributions

F.J. conceived the project, obtained funding, and supervised the study. F.J. and L.Y. wrote the manuscript. L.Y. performed bioinformatics analysis, statistical analysis, and

951   data visualization. L.C refined viral genomics analysis and data visualization.   F.J.

952   finalized and all authors approved the final version of the manuscript.

953   **Competing interests**

954   The authors declare no competing interests

Fig. 1 Exploring RNA viruses in the global WWTP metatranscriptomes. a. Distribution of sample types and locations of 557 publicly available WWTP metatranscriptomes and the brief bioinformatic workflow for RNA vOTUs identification. b. The number of RNA viral contigs detected across eight hmmsearch iterations (upper panel) and the accumulation curve of unique RNA vOTUs identified in the samples (below panel). c. The α-diversity of RNA vOTUs measured using Shannon's index across five sample types in all WWTP samples. d. NMDS analysis of abundance distribution of RNA vOTUs in all WWTP samples. e. The absolute abundance of all identified RNA vOTUs expressed as copies per gram (left) and copies per liter (right), is shown for four sample types in the 47 WWTP samples from Switzerland. Statistical significance was determined using the Mann-Whitney U test, with significance levels represented as ∗∗∗∗ for adjusted $P < 0.00001$. IN: influent, DS: denitrification sludge, AS: activated sludge, EF: effluent, AD: anaerobic digester sludge.

Fig. 2 The phylogenetic and taxonomical expansion of RNA viral diversity. a. The global RdRp-based phylogenetic tree. The tree consists of reference RdRp centroids and vOTUs encoded "complete" RdRp domains, classified into known RNA viral families. Each family clade is accompanied by two pie charts, representing the distribution of its clusters within the GRvOTU set that occurred in engineering systems (left pie) and all ecosystems (right pie). In each pie chart, the segments indicate the proportions of novel (members only identified in the WWTPs), reference (members only identified in the reference dataset), and shared (members present in both datasets) clusters. The numbers adjacent to the pie charts show the count of 'novel' clusters within the GRvOTU set for the family. Host range icons next to family names represent their known host ranges. Icons highlighted in red indicate a subset of *Partitiviridae* and *Picobirnaviridae* identified as bacteriophages, challenging previously assumed host ranges (see Section: Tracking Host Associations of RNA Prokaryotic Viruses in Global WWTPs). b. Expansion of RNA viral diversity in engineering systems, showing the number of known RNA viral clusters (left) and the distribution of 'novel', 'shared', and 'reference' clusters (right) within the GRvOTU set.

Fig. 3 Expanding RNA bacteriophage diversity from global WWTPs. a. The number of WWTP vOTUs from known and novel RNA bacteriophage clades. b. Phylogenetic placement of the Base-*Howeltoviricetes* phage clade within the *Lenarviricota* phylum. c. Phylogenetic tree of the Base-*Howeltoviricetes* phage clade pruned from the global RdRp tree. Blue outer bars represent the proportion of ORFs associated with the SD motif in the corresponding reference RdRp centroids or vOTUs, while gray bars indicate cases where no ORFs with true start codons were identified, precluding SD motif assessment. The vOTU marked with a star is linked to a probable host based on CRISPR spacer matching. d. Global distribution of the Base-*Howeltoviricetes* phage clade. A viral cluster from the GRvOTU set is considered part of the Base-*Howeltoviricetes* phage clade if it contains sequence members that belong to the clade's phylogenetic tree. Sequence members from these clusters are traced back to their corresponding original public samples, and the number of Base-*Howeltoviricetes* phage sequences assembled in these samples is counted. The distribution of this phage clade is then visualized according to sample types and geographic locations.

Fig. 4 Host association and gene composition of Base-*Howeltoviricetes* phages. a. CRISPR spacer matching links vOTU FD-DS_117671 within the Base-*Howeltoviricetes* phage clade to *Campylobacterota*. Arrows indicate the blastn-short alignments between host spacers and FD-DS_117671, with numbers indicating the count of matching RNA viral contigs within the vOTU. NZ_CP026656.1 and NZ_SNQM01000020.1 are genomes from the NCBI RefSeq database, while TG-IN_1259 and ZR-IN_6153 are WWTP metagenomic contigs. ZR-IN_6153 is binned into a MAG belonging to *Aliarcobacter* sp. Coding genes point right for the (+) sense strand and to the left for the (−) sense strand. b. Relative abundances of FD-DS_117671 and *Aliarcobacter cryaerophilus* during wastewater treatment processes (47 samples from Switzerland, upper panel) and in time-series samples (47 sludge samples from Luxembourg, lower panel). Viral abundance is quantified as the Tmean calculated by CoverM normalized by sample size, and host abundance is estimated using motus3. c. Genome extension for Base-*Howeltoviricetes* phages. Two examples of genome extension are shown. TU-EF_29550 dominates the RNA viral community in BE WWTP samples, with genome extension starting from BE-AS_5392. Similarly, AH-EF_62580 dominates SF50-52 samples, with its genome extension starting from SF51_6014.

Fig. 5 Distribution and absolute concentration of human RNA vOTUs in the WWTPs. a. Heatmap depicting the abundance distribution of 21 human RNA vOTUs identified by species-level clustering. Each row represents a human RNA vOTU and each column represents a sample. Upper annotation columns illustrate the sample location and type. b. Bubble plot showing the estimated absolute concentration of human RNA vOTUs (in copies per liter) in the 47 samples from Switzerland which spiked with mRNA internal standards.

Fig. 6 The AMGs encoded by RNA viral genomes revealed the involvement of RNA viruses in various metabolic processes. AMGs are highlighted in red. Coding genes point right for the (+) sense strand and to the left for the (−) sense strand. Below the gene flow, the 3D structures of the AMGs predicted by ESMfold and annotated by Foldseek are displayed. Known hosts are indicated alongside the affiliated taxonomy.

Extended Data Fig. 1 The bioinformatic pipeline for studying RNA viruses in the WWTPs in this study. a. A total of 557 metatranscriptomes were downloaded from public databases and used for RNA virus exploration. Among the collected metatranscriptomes, 47 samples from Switzerland were spiked with mRNA internal standards enabling the estimation of the absolute concentration of RNA viruses in these samples. The metatranscriptomes were first quality controlled and then assembled into contigs. b. After gene prediction, putative viral RdRps were identified by an iterative hmmsearch approach. The authenticity of identified RdRps was assessed by a comparative hmmsearch of putative viral RdRps against RdRp and non-RdRp profiles as well as checking the occurrence of putative viral RdRp-encoding sequences in the metagenomes. c. The RNA vOTUs were clustered at 90% average nucleotide identity (ANI) over 80% alignment fraction of the shorter sequence (AF). Then, to systemically investigate the diversity and ecological roles of RNA viruses in the WWTPs, bioinformatic analyses including taxonomic and phylogenetic classification, RNA bacteriophage identification, human RNA virus identification, and AMG annotation and expression detection were conducted for WWTP RNA viruses.

Extended Data Fig. 2 Taxonomy-based richness of RNA viruses in the WWTPs. The richness percentage of RNA vOTUs at the family level. Among 9276 RNA vOTUs encoding "complete" RdRp domains, 5945 RNA vOTUs were classified into established viral families. Host range icons next to family names represent their known host ranges.

Extended Data Fig. 3. Phylogenetic trees of established RNA bacteriophage families. Each tree includes RdRps from RCR90 set occurred in at least one engineering sample. Leaf colors indicate the type of the corresponding RCR90 RdRp cluster as either "novel," "shared," or "reference."

Extended Data Fig. 4. RdRp phylogenetic tree of the newly identified megataxon. This megataxon primarily comprises RdRps from the WWTPs and unclassified reference RdRp centroids. The outer color strip indicates the origin of the RdRp sequences (either from this study or from reference datasets). Among the reference sequences, only ND_053632_1 is annotated with a taxonomic label (*Bunyavirales*), whereas the remaining reference sequences lack taxonomic classification.

Extended Data Fig. 5. The abundance proportion of RNA viruses in the WWTPs. The abundance proportion of RNA vOTUs in the samples from Switzerland (a) and all samples (b) grouped by their host ranges suggested by affiliated taxonomy. The abundance proportion of RNA bacteriophage clades in all samples (c). The host range of each RNA viral lineage was determined based on: 1) ICTV proposals and 2) previously unknown bacteriophage clades identified in this study (*Picobirnaviridae*, part of *Partitiviridae*, Base-*Cystoviridae*, and part of Base-*Howeltoviricetes*).

Extended Data Fig. 6. RNA bacteriophage clades in phylum *Pisuviricota* identified through Shine-Dalgarno motifs. a. Phylogenetic placement of RNA bacteriophage clades within the *Pisuviricota* phylum. All RNA viral families in the tree are assigned to the *Durnavirales* order, except for (base-)*Cystoviridae*, which is currently assigned to *Duplornaviricota* by ICTV but was placed under *Pisuviricota* in both this study and a recent study (Neri et al.). b–d. Phylogenetic trees of (base-)*Cystoviridae* (b), *Picobirnaviridae*, and a subset of *Partitiviridae* (d) were pruned from the global RdRp tree. Blue outer bars represent the proportion of ORFs associated with the SD motif in the corresponding reference RdRp centroids or vOTUs, while gray bars indicate cases where no ORFs with true start codons were identified, precluding SD motif assessment. The RdRps of *Cystoviridae* and base-*Cystoviridae* are phylogenetically intermingled. Thus, their phylogeny is displayed together in Figure b, with an outer color strip indicating their corresponding taxonomy.

Extended Data Fig. 7 Potential human RNA viruses identified through RdRp phylogeny. RdRps labeled in red represent sequences from WWTP RNA vOTUs, while the remaining sequences are reference RdRp domain sequences obtained from the NCBI Genbank Database. Human RNA viruses identified through both species-level clustering (See Methods) and RdRp phylogeny are marked with a blue star. Human RNA viruses identified exclusively through RdRp phylogeny are marked with a red star and are considered novel human RNA viruses, at least at the species level.

Extended Data Fig. 8 Detection of potential human RNA viruses through short-read alignments. The heatmap displays the RPKM values of human RNA viruses identified through read-based alignment. The alignment was performed using blastn to align clean RNA reads from the metatranscriptomes against the NCBI Virus database, with a 90% identity threshold and 80% read length coverage as cutoffs. A constructed mock community validated that the read-based approach for detecting human RNA viruses in global WWTP metatranscriptomes is reliable (See Methods).

Supplementary Fig. 1. α- and β-diversity of RNA vOTUs throughout the wastewater treatment process. a. α-diversity of RNA vOTUs measured in Shannon's index across 4 sample types in 47 WWTP samples from Switzerland. b. NMDS analysis of the abundance distribution of RNA vOTUs in 47 WWTP samples from Switzerland. c. ANOSIM analysis showing the dissimilarity rank of RNA vOTU compositions between and within different WWTP compartments in Switzerland

.

Supplementary Fig. 2 Complete global RdRp phylogenetic tree. The phylogenetic tree includes of all reference RdRp centroids and RNA vOTUs encoding "complete" RdRp domains. The color of each clade in the tree represented the lowest taxonomic level assigned to that clade. The number next to each clade indicates that the majority of its members belong to the corresponding megataxon. Specific labels for each megataxon are provided in Supplementary Table 4.

Supplementary Fig. 3. NMDS analysis of the short reads-based distribution of human RNA virus in influent samples from Switzerland and the USA.

Supplementary Fig. 4. Phylogenetic tree of RNA virus-encoded 60S ribosomal protein and reference sequences from the NCBI nr database.

Supplementary Fig. 5. Phylogenetic tree of RNA virus-encoded 40S ribosomal protein and reference sequences from the NCBI nr database.

Supplementary Fig. 6. Phylogenetic tree of RNA virus-encoded 50S ribosomal protein and reference sequences from the NCBI nr database.

Supplementary Fig. 7. Phylogenetic tree of RNA virus-encoded 50S ribosomal protein and reference sequences from the NCBI nr database.

Supplementary Fig. 8. Abundance proportions of the top ten most abundant RNA viral families in this study and other ecosystems surveyed by Neri et al.