Title: The best of two worlds: using stacked generalization for integrating expert range maps in

species distribution models

Authors: Julian Oeser[1], Damaris Zurell[2], Frieder Mayer[3], Emrah Çoraman[3,4], Nia Toshkova[5],

Stanimira Deleva[5], Ioseb Natradze[6], Petr Benda[7,8], Astghik Ghazaryan[9], Sercan Irmak[4,10], Nijat

Alisafa Hasanov[11], Gulnar Gahraman Guliyeva[11], Mariya Gritsina[12], and Tobias Kuemmerle[1,13]

Affiliations:

[1] Geography Department, Humboldt-Universität zu Berlin, Unter den Linden 6, 10099 Berlin,

Germany

[2] Ecology and Macroecology, Inst. for Biochemistry and Biology, University of Potsdam, 14469

Potsdam, Germany

[3] Museum für Naturkunde, Leibniz-Institut für Evolutions- und Biodiversitätsforschung, Berlin

10115, Germany

[4] Eurasia Institute of Earth Sciences, Department of Ecology and Evolution, Istanbul Technical

University, Maslak, Istanbul 34469, Türkiye

[5] National Museum of Natural History, Bulgarian Academy of Sciences, Sofia 1000, Bulgaria

[6] Institute of Zoology of Ilia State University, Tbilisi 0162, Georgia

[7] Department of Zoology, Natural History Museum Prague, 115 79 Praha 1, Czech Republic

[8] Department of Zoology, Faculty of Science, Charles University in Prague, 128 44 Praha 2,

Czech Republic

[9] Department of Zoology, Yerevan State University, Yerevan 0025, Armenia

[10] Science and Technology Research and Application Center, Balikesir University 10145,

Balıkesir, Türkiye

[11] Institute of Zoology, Ministry of Science and Education, 1004 Baku, Azerbaijan

[12] Institute of Zoology, Academy of Science of Uzbekistan, 100053 Tashkent, Uzbekistan

[13] Integrative Research Institute on Transformation in Human Environment Systems, Humboldt-

Universität zu Berlin, Berlin, Germany

Acknowledgements:

We gratefully acknowledge Alexander Buknikashvili, Christian Dietz, Amit Dolev, Heliana

Dundarova, Eran Levin, and Panagiotis Georgiakakis for providing bat occurrence data. Giorgi

Sheklashvili, Andrei Kandaurov, Irina Rakhmatulina, Kursantbek Altybaev, Pirimkul

Mamatkalykov, Abdurashit Nizamievf, Alexey Dudashvili Svetlana V. Baskakova, Georgiy V.

Shakula, Kudaibergen Amirekul, Artemis Kafkaletou Diez, Ioanna Salvarina, and Eleni

Papadatou helped with the collection of bat occurrence data. In addition, we are thankful to

Christian Dietz for help with the taxonomic classification of bat occurrence records.

Abstract

Species distribution models (SDMs) are powerful tools for assessing suitable habitat across large

areas and at fine spatial resolution. Yet, the usefulness of SDMs for mapping species' realized

distributions is often limited, since data biases or missing information on dispersal barriers or

biotic interactions hinder them from accurately delineating species' range limits. One way to

overcome this limitation is to integrate SDMs with expert range maps, which provide coarse-

scale information on the extent of species' ranges that is complementary to information offered

by SDMs. Here, we propose a new approach for integrating expert range maps in SDMs based on

an ensemble method called stacked generalization. Specifically, our approach relies on training a

meta-learner regression model using predictions from one or more SDM algorithms alongside the

distance of training points to expert-defined ranges as predictor variables. We demonstrate our

approach with an occurrence dataset for 49 bat species covering four biodiversity hotspots in the

Eastern Mediterranean, Western Asia, and Central Asia. Our approach offers a flexible method to

integrate expert range maps with any combination of SDM modeling algorithms, thus facilitating

the use of algorithm ensembles. In addition, it provides a novel, data-driven way to account for

uncertainty in expert-defined ranges not requiring prior knowledge about their accuracy, which is

often lacking. Our approach holds considerable promise for better understanding species

distributions, and thus for biogeographical research and conservation planning. In addition, our

work highlights the overlooked potential of stacked generalization as an ensemble method in

species distribution modeling.

1 Introduction

While global biodiversity is declining rapidly (Pimm et al. 2014), our knowledge about species'

distributions often remains limited (Diniz-Filho, De Marco Jr, and Hawkins 2010). This lack of

detailed information for many regions and taxa, referred to as the Wallacean shortfall (Hortal et

al. 2015), translates not only into knowledge gaps in biogeography and ecology, but also into real

barriers for conservation planning to ensure that limited conservation funding is spent most

effectively (Hochkirch et al. 2021). Species distribution models (SDMs) have become a central

tool for addressing the Wallacean shortfall, allowing to characterize species' niches by combining

occurrence records with environmental predictors for predicting species' distributions (Elith and

Leathwick 2009; Guisan and Thuiller 2005). Yet, although SDMs can accurately assess the

environmental suitability of habitats (i.e., map *potential distributions*), they typically lack

information on other factors limiting species' ranges, such as barriers to dispersal or biotic

interactions (i.e., competitive exclusion). This, in turn, means that the usefulness of SDMs for

mapping *realized distributions* of species can be limited, as their inability to identify range limits

often results in an overprediction of species' ranges, particularly when distributions are modelled

across large geographic extents (Calabrese et al. 2014; Merow, Wilson, and Jetz 2017; Soberón

2007). While methods for capturing dispersal and biotic interactions within SDMs have been

proposed (Ovaskainen et al. 2016; Zurell 2017), their applicability is often limited due to a lack

of adequate datasets or missing knowledge about underlying ecological processes.

A widely-applicable solution to improve SDMs' capability for assessing realized

distributions lies in their combination with external information on species' range limits

(Domisch, Wilson, and Jetz 2016; Fletcher Jr. et al. 2019; Merow et al. 2017). Most commonly,

range information is available in the form of expert-based range maps, which offer estimates of

species' range extents derived from occurrence information as well as expert knowledge about

geographical, biotic, or environmental range limits. The most important database of range maps

(particularly for terrestrial animals) is offered by the International Union for the Conservation of

Nature (IUCN), which provides expert-defined ranges for more than 150,000 species (IUCN

2022). Although widely available, expert range maps are frequently criticized for being coarse in

resolution (meaning that species will often be absent from many areas within the expert-defined

range), incomplete in terms of species coverage or outdated (Higino et al. 2023; Ramesh et al.

2017). Despite these shortcomings, expert range maps often provide the best-available (or only)

information on range limits for many species. More importantly, they provide information that is

highly complementary to data generated by SDMs (Merow et al. 2017). While range maps

characterize a species' extent of occurrence (i.e., its range limits), SDMs offer fine-scale

representations of suitable habitats, making approaches combining both datasets promising for

improving distribution assessments (Domisch et al. 2016; Ellis-Soto et al. 2021; Merow et al.

2017).

Several approaches have sought to combine these relative strengths of expert range maps

and SDMs, such as using range maps directly as predictors in SDMs (Domisch et al. 2016) or

adding spatial offset terms to models that are fit via point process models or related approaches

(e.g., Maxent; Merow et al., 2017). The latter approach is particularly promising as it allows to

account for uncertainty in expert range maps by incorporating user-defined decay curves that

reflect *a priori* expectations about the accuracy of expert range boundaries. Applying this

approach, however, can be challenging for two reasons. First, defining spatial offsets and decay

curves can be difficult if prior information on the accuracy of range maps is missing, potentially

leading to bias introduced by decisions on the strength and decay of the offset term. Second,

while the use of algorithm ensembles has become a key approach in species distribution

modeling (Araújo et al. 2019; Araújo and New 2007), several widely-used and well-performing

machine learning algorithms (e.g., random forests or support vector machines) do not feature

offset terms.

Here, we suggest stacked generalization (Wolpert 1992) as an alternative approach for

integrating external range information enabling flexible combinations of multiple SDM

algorithms. Designed as an ensemble method for combining multiple modeling algorithms,

stacked generalization uses the predictions of models built at one level as the input for a meta-

learner built at a second level (Naimi and Balzer 2018). Although being widely applied in

machine learning (Sesmero, Ledezma, and Sanchis 2015), and despite the general proliferation of

algorithm ensembles in SDM studies (Buisson et al. 2010; Hao et al. 2019), stacked

generalizations have rarely been used with SDMs (but see Bonannella et al., 2022; El Alaoui &

Idri, 2023). Here, we demonstrate the use of stacked generalization as an approach for integrating

expert range information with one or more SDM algorithms. Using available occurrence datasets

for characterizing expert map accuracy, our approach offers an alternative, data-driven method to

integrate expert range maps in SDMs while accounting for their uncertainty.

In the following, we first introduce our approach and highlight issues important to

consider in its application. Then, we assess our approach by applying it to a presence-only

occurrence dataset for 49 bat species collected across a large geographic extent covering four

biodiversity hotspots in the Eastern Mediterranean, Western and Central Asia. Specifically, we

compare the predictive performance as well as resulting distribution maps of (1) single-algorithm

SDMs, (2) ensembles of SDM algorithms built with stacked generalization, and (3) stacked

generalizations including expert range maps.

2. Stacked generalization for integrating expert range information in SDMs

Stacked generalization is an ensemble method for combining multiple models, often built with different algorithms, using their individual predictions as training data in a meta-learner (Naimi and Balzer 2018; Wolpert 1992). Here, we apply this approach to integrate one or more SDM algorithms with expert range maps. By using the expert map as an additional model containing complementary information to SDMs (i.e., coarse-scale estimate of range limits), our approach is taking advantage of stacked generalizations working best if heterogeneous input models are combined (Sesmero et al. 2015).

Multiple potential approaches exist to combine SDMs with expert range maps via stacked generalization. One approach is creating a predictor variable for the meta-learner by assigning a fixed value ratio to training points lying inside vs. outside the expert-defined ranges, thereby allowing to control how much weight is given to the expert map (Merow et al. 2017). However, this approach assumes that the probability of occurrence is the same at any distance outside the expert range, although a continuously decreasing probability with increasing distance from the expert range should be expected (Merow et al. 2017). Therefore, we instead use the spatial distance of the training points to the expert range boundaries as a predictor in the meta-learner (Figure 1). This predictor, hereafter referred to as *distance term*, describes the (relative) probability of observing the modeled species within a given distance of the expert-defined range, thereby characterizing the uncertainty of the expert map. This approach is conceptually similar to including spatial offsets with decay curves in point process models (Merow et al. 2017), and results in predicted habitat suitability values smoothly decreasing outside the expert range. However, in contrast to user-defined offsets, the distance term of the meta-learner is derived from the occurrence records used to train SDMs. While using the same datasets for fitting SDMs and assessing the uncertainty of the expert range maps might introduce bias if the collection of

occurrence records is influenced by knowledge about expert ranges (Merow et al. 2017), such a

data-driven approach will be particularly useful when accurate and representatively sampled

occurrence records are available or if prior knowledge about the accuracy of expert range maps is

lacking.

While the approach by Merow et al. (2017) allows to control the shape of the decay curve

by choosing several curve parameters, in stacked generalizations, the analyst can influence the

shape of the fitted distance term through the choice of the meta-learner algorithm or the

functional form of the distance term. As a baseline approach we here use logistic regression as a

meta-learner, which is widely used in stacked generalizations and results in distance terms

following a logistic function similar to the smooth decay curves proposed by Merow et al.

(2017). Conceptually, adding the distance term to a logistic regression meta-learner can be seen

as adding a constant 'offset' to all areas inside the expert-defined range (i.e., areas with distance

= 0). This offset is described by the intercept of the logistic regression and expresses the

(relative) probability of observing the species inside the expert range given suitability predictions

of 0 from all SDM algorithms. Predictions by the meta-learner will decrease with increasing

distance from the expert range according to the distance term (Figure 1).

Figure 1: *Schematic overview of stacked generalization for combining SDM algorithms with*

expert range maps. Predictions of multiple SDM algorithms are used together with the distance

of occurrence data to the expert range as predictor variables in a logistic regression meta-

learner, which then is used to predict the species' distribution. Maps show examples for one bat

species in our dataset (Nyctalus noctula). Map panel for expert range shows IUCN range in grey

with presence records colored according to their distance to the IUCN range. Shown maps are in

Albers equal area projection.

By relating individual species' occurrences to expert ranges, our approach accommodates

species-to-species variability in the uncertainty of expert ranges. However, due to a lack of

presence records or highly accurate expert range maps, in some cases only few or no presence

records might lie outside expert ranges, which will cause (quasi-)complete separation in the meta-

learner. We propose two potential solutions to this issue. First, if species-specific distance terms

should be used, bias-reduced logistic regression can be applied for fitting meta-learners (Firth

1993). This commonly recommended strategy for dealing with (quasi-)complete separation in

logistic regressions ensures finite parameter estimates and results in responses (i.e., distance

terms in our case) that are less steep compared to standard maximum likelihood estimation

(Heinze and Schemper 2002). Second, when occurrence data from multiple related taxa are

available, species-specific distance terms of meta-learners might be replaced with 'target-group'

distance terms, which can be calculated by fitting a meta-learner based on training points from

multiple or all available species. In this case, the distance term characterizes the uncertainty

(probability of occurrences lying outside expert ranges) across all included taxa and does not vary

between species, similar to applying the same decay curve across species when integrating range

maps as spatial offsets in point process models (Merow et al. 2017).

3. Method application

3.1 Study area and bat occurrence data

Our study area covers 6.5 million km² and intersects four global biodiversity hotspots (following

Myers et al., 2000): The eastern part of the Mediterranean hotspot, the Caucasus hotspot, the

Irano-Anatolian hotspot, as well as partially covering the Mountains of Centrals Asia hotspot. To

delineate our study area, we fully included all countries in which the sampling of our bat

occurrence records was primarily conducted (Afghanistan, Albania, Armenia, Azerbaijan,

Bulgaria, Georgia, Greece, Iran, Israel, Montenegro, Syria, Turkey). The borders of our study

area were defined based on ecoregion boundaries (Olson et al. 2001). Our study area represents

the contact zone between the Western and Eastern Palearctic species pools, where information on

the distribution of bats remains limited.

Figure 2: Extent of the study area, shown as black polygon and intersecting global biodiversity

hotspots shown as colored polygons. Map is in Albers equal area projection.

We collected and harmonized bat occurrence datasets from various sources, including national

databases, field records, and literature data (see Appendix S1 for an overview of all data sources).

In total, we gathered 37,714 occurrence records from 61 taxa. To ensure the quality of records

used for model training, we removed all instances in which a species-level identification was

impossible or problematic (e.g., uncertain identification within complexes of morphologically

highly similar species). In addition, we removed records collected before 1970 to avoid a

temporal mismatch between occurrence records and predictor variables (Milanesi, Della Rocca,

and Robinson 2020).

Where appropriate, we reclassified records to account for recent genetic analyses that

have led to a subdivision of species complexes into multiple cryptic species. This reclassification

was done based on available information on the distribution of cryptic species (see Appendix S2

for details on taxonomic revisions within species complexes as well as an overview of species).

To remove spatial duplicates and reduce sampling bias, we thinned occurrence records (Boria et

al. 2014). As thinning records may reduce model performance for rare species (Steen et al. 2021),

we classified species according to the percentile values of sample prevalence (i.e., number of

raster cells with presence records) into three classes (low, intermediate, and high prevalence). We

then thinned records with minimum distances of 1km, 5km, and 10km for species with low,

intermediate, and high prevalence, respectively.

As expert information on species range limits, we compiled IUCN range maps for all

species. This led to four species being excluded from modelling since no range map was

available. Finally, we selected species with a minimum of 30 remaining records to ensure robust

training data sets for building SDMs, resulting in 9,650 presence records from 49 species.

3.2 Species distribution modeling

We used presence-background SDMs (Elith and Leathwick 2009) to characterize the distributions

of bats in our study area. For modeling, we compiled a set of 40 candidate predictor variables,

indicating four key dimensions of habitat suitability for bats: climate, land cover and vegetation

productivity, topography and geology, and human pressure and modification (Table 1). While

target resolution of our SDMs was 1km², we derived all predictor variables at three spatial scales

(1km², 5km² and 10km²), resulting in 120 candidate variables. Including coarser scales derived

through moving window averaging allows better characterizing habitat conditions at the scale of

bat home ranges (e.g., available forest cover within the surrounding area of a bat roost).

Table 1: Overview of environmental predictor variables used in species distribution models.

Category	Predictor	Available time steps	Data source
Climate	19 bioclimatic variables	1981-2010 (average)	CHELSA climate data (Karger et al. 2017)
Land cover and vegetation productivity	Six land-cover proportions (agriculture, forest, shrubs, herbaceous vegetation, bare and sparse vegetation, water)	1992-2020 (annual)	ESA CCI land cover
	Nine Landsat-based spectral-temporal metrics (cumulative, minimum and seasonality metrics for Tasseled Cap greenness, brightness, and wetness indices)	1990, 1995, 2000, 2005, 2010, 2015	Landsat satellite imagery (Oeser et al. 2020)
Topography and geology	Terrain ruggedness index	-	(Amatulli et al. 2018)
	Presence of karstifiable rocks	-	World Karst Aquifer Map (Chen et al. 2017)
Human pressure and modification	Human modification index	1990, 2000, 2010, 2015, 2017	(Theobald et al. 2020)
	Accessibility (travel time to cities)	2015	(Weiss et al. 2018)
	Nighttime-lights	1992-2018	(Zhao et al. 2022)
	Forest landscape integrity index	2019	(Grantham et al. 2020)

We sampled background points using a target group bias grid, created from kernel density

estimation based on all presence records in our dataset. Using the density of bat occurrence

records as sampling weights for background points allows characterizing sampling effort and

helps to mitigate the influence of sampling bias in presence-background SDMs (Barber et al.

2022; Inman et al. 2021; Syfert, Smith, and Coomes 2013). For each species, we sampled

background points equal to ten times the number of available presence records.

We used three SDM algorithms: Maxent (R-package *dismo*; Hijmans et al., 2020),

random forests (R-package *randomForest*; Cutler & Wiener, 2022), and boosted generalized

additive models (GAMs, R-package *mboost*; Hothorn et al., 2022). Following recommendations

by Valavi et al. (2021), we used down-sampled random forests, in which subsamples of the

background points are used within each individual tree in order to correct for class imbalances. In

a first modeling step, we performed variable selection by fitting univariate models (with default

parameters) for all 120 candidate variables and evaluating their predictive performance using the

area under the receiver operating characteristic curve (AUC) and Pearson correlation between the

predicted and observed presence (COR) in a five-fold cross validation (Valavi, Guillera-Arroita,

et al. 2021). For selecting the best-performing model, we combined AUC and COR into a single

performance score by rescaling their values across all tested models to a 0-1 scale and calculating

the mean of rescaled AUC and COR values. Based on this combined performance score, for each

species, we selected the set of variables offering the best predictive performance while having

correlation coefficients $|r|<0.7$ (Dormann et al. 2013). Using the selected variables in a second

five-fold cross validation, we tuned algorithm parameters for all species (selecting regularization

multipliers for Maxent, *mtry* and *maxnodes* parameters in random forest, and the number of

boosting iterations in boosted GAMs; see Appendix S4 for details).

3.3 Stacked generalizations

We implemented stacked generalizations in two ways. First, we created pure algorithm

ensembles (hereafter *SDM ensembles*) solely relying on the predictions of the three SDM

algorithms as predictors in the meta-learner. Second, we created *expert-informed ensembles*

additionally including information from IUCN range maps. Additionally, we compared two

approaches for adding distance terms to the meta-learner. First, we used species-specific distance

terms, using the distance of species-level training points to the species' IUCN range as a

predictor in the meta-learner. Second, we calculated a target-group distance term, which we

derived by fitting a logistic regression to the distances of all bat occurrence records in our dataset

(i.e., all 49 species). To deal with (quasi-)complete separation in species-specific distance terms,

we used bias-reduced logistic regression implemented in the R-package *brglm2* (Kosmidis et al.

2023) for fitting meta-learners.

A critical consideration when using stacked generalizations is the risk of overfitting the

meta-learner. A widely adopted strategy for this purpose, referred to as *Super Learner* (van der

Laan, Polley, and Hubbard 2007; Naimi and Balzer 2018), uses out-of-sample predictions (i.e.,

from cross validation) for training the meta-learner. To assess the effect of overfitting on stacked

generalizations, we compared meta-learners trained on out-of-sample vs. in-sample predictions

(i.e., with and without the *Super Learner* strategy).

We compared the predictive performance of all three tested modeling approaches using

five-fold cross validation: (1) single-algorithm SDMs (i.e., Maxent, random forest, and boosted

GAMs), (2) SDM ensembles, (3) expert-informed ensembles. To create distribution maps for all

species, we predicted all models for the most recent time step (target year for prediction: 2020).

To compare mapped distribution patterns between SDM ensembles and expert-informed

ensembles, we calculated two metrics: First, species-wise niche breadth using Levins' B2 metric,

describing the uniformity of predicted suitability in geographic space (Warren et al. 2021), and

second range overlaps calculated using Schoener's D metric, describing the similarity of

predicted suitability between species pairs (Warren, Glor, and Turelli 2008). We hypothesized

that expert-informed ensembles should result in overall lower niche breadths and lower range

overlaps, since integrating information on species' range limits should correct for the

overprediction of species ranges by SDMs due to missing information on the effect of dispersal

limitations and biotic interactions (Merow et al. 2017).

4 Results

The accuracy of IUCN range maps varied considerably across bat species. On average, 73% of

presence records fell inside expert-defined ranges (inter-quartile range: 22%), with records lying

at an average distance of 50 km of expert-defined range boundaries (inter-quartile range: 30 km).

These differences in the accuracy of expert range maps translated into considerable variation in

species-specific distance terms and thus clear differences in how predicted suitability values

declined outside expert ranges when using expert-informed ensembles. In the case of accurate

expert ranges, suitability sharply declined outside expert ranges, leading to the exclusion of (often

large) areas identified as environmentally suitable by SDMs but lying outside species' ranges

(e.g., *Myotis myotis* in Figure 3). Conversely, when occurrence records indicated that expert

range maps were inaccurate, SDMs clearly dominated the predictions of expert-informed

ensembles, allowing to identify areas outside IUCN ranges as likely occupied by species (e.g.,

Taphozous nudiventris in Figure 3). When using target-group instead of species-specific distance

terms, suitability values declined at similar rates outside expert ranges across species (Appendix

S4).

Figure 3: *Comparison of distribution maps (A+B) and decline in predicted occurrence*

probabilities outside expert ranges (C+D) for two example species with high (Myotis myotis) and

low expert map accuracy (Taphozous nudiventris). Distribution maps based on IUCN ranges

(including available presence records), SDM ensembles, and expert-informed ensembles are

shown. Expert-informed ensembles correspond to models built with species-specific distance

terms. Plots of decline in predicted occurrence probabilities outside expert ranges (C+D) are

based on loess smooth to the data. Maps are in Albers equal area projection.

Considering predictive performance, stacked generalization ensembles outperformed single-

algorithm SDMs. However, training on out-of-sample predictions was necessary to achieve

optimal predictive performance (i.e., using *Super Learner* approach; Figure 4). Specifically,

expert-informed ensembles built with species-specific distance terms achieved the highest

predictive performance according to both AUC and COR values, followed by SDM ensembles

and expert-informed ensembles with target-group distance terms (Figure 4; since COR values

showed no qualitative difference to AUC, we only show AUC values here).

Figure 4: *Predictive performance of modeling approaches for 49 bat species in Eastern*

Mediterranean, Western Asia and Central Asia according to AUC values.

Performance improvements of expert-informed ensembles compared to SDM ensembles

generally diminished with the average distance of presence records to expert ranges (i.e.,

increasing performance gains with higher expert map accuracy; Figure 5A). Performance

improvements tended also to be higher for species with fewer available occurrence records as

well as for species with smaller range extents (Figure 5B+C), but these relationships were

considerably weaker than for expert map accuracy.

Figure 5: *Improvement in predictive performance of expert-informed ensembles compared to*

SDM ensembles in relationship to expert map accuracy (mean distance of presence records to

expert range, including points inside the range with distance = 0). Data for expert-informed

ensembles with species-specific distance terms are shown, with linear trend plotted on top.

Considering mapped distribution patterns, expert-informed ensembles resulted in lower niche

breadths (i.e., less uniform distribution of predicted suitability in geographic space) for 83% of

species compared to SDM ensembles. On average, species-wise niche breadths obtained from

expert-informed ensembles were 21% lower compared to niche breadths derived from SDM

ensembles (Figure 6A). Range overlaps between species pairs (i.e., similarity of predicted

suitability) derived from expert-informed ensembles were lower than overlaps predicted by SDM

ensembles in 90% of the cases. On average, overlaps were 30% lower in expert-informed

ensembles compared to those obtained from SDM ensembles(Figure 6B).

Figure 6: *Distribution of (A) niche breadths and (B) range overlaps of bat species according to*

SDM ensembles vs. expert-informed ensembles.

5. Discussion

Addressing the Wallacean shortfall is critical to biogeographical research and conservation

planning, yet accurately mapping species' realized distributions through species distribution

modeling presents a significant challenge. Here, we developed a new approach for integrating

expert information on range limits in species distribution models by making use of stacked

generalization, an ensemble method widely applied in machine learning but still underexplored in

the context of SDMs. Testing our approach with a dataset covering 49 bat species demonstrated

its flexibility and promise for improving species distribution mapping, allowing to combine the

key strength of SDMs (characterizing environmentally suitable habitats) with that of expert range

maps (characterizing range limits) without requiring prior knowledge about expert range maps or

having to rely on specific modeling algorithms. In a broader context, we add to the growing

toolbox of integrated SDM approaches, providing an important step towards more accurate

assessments of species' distributions.

The application of our approach showed that it effectively enables the exclusion of areas

lying outside species' realized range limits, while preserving fine-scale predictions of habitat

suitability, which offer a key strength of SDM approaches (Mainali et al. 2020). At the same

time, when enough presence records are recorded outside expert-defined ranges, expert range

maps exert minimal influence on mapped distributions, demonstrating the flexibility of our

approach towards varying levels of expert map accuracy. We did not have an independent

validation dataset on species' absence available, precluding us from performing a more detailed

assessment of how our approach affects the accuracy of mapped distributions. Yet, we found

improved predictive performance of expert-informed ensembles compared to pure SDM

ensembles within our presence-background dataset, with performance improvements depending

on the accuracy of expert range maps.

Our approach offers an alternative to applying user-defined spatial offsets in point-process

models as proposed by Merow et al. (2017). Choosing between stacked generalization and spatial

offsets as ways to integrate expert range maps boils down to selecting different styles of

modeling approaches: relying either on prior knowledge when using spatial offsets or on available occurrence datasets when using stacked generalizations for characterizing expert map accuracy. The appropriateness of using stacked generalizations thus hinges on whether available occurrence records can accurately capture expert map accuracy. As highlighted by Merow et al. (2017), using occurrence records for characterizing expert map accuracy requires that their collection is independent of expert range maps (i.e., expert range maps not affecting sampling intensity or species identification), otherwise they will give a biased view on expert map accuracy. However, in many cases occurrence records provide a more comprehensive and up-to-date picture of species distributions compared to expert range maps. Moreover, occurrence records will often be the best available (or only) type of data allowing to evaluate expert range maps, as other *a priori* information on their accuracy is difficult to obtain. Our application of stacked generalizations for 49 bat species highlighted the advantage of allowing for species-to-species variation in the uncertainty of expert-defined ranges, with models achieving the highest predictive performance when using species-specific distance terms. Thus, given that expert map accuracies are expected to vary strongly across species, stacked generalization provides a simple yet effective data-driven approach without the need for manually adjusting prior expectations when assessing many species at once. If occurrence datasets for individual species are deemed too incomplete or biased for characterizing expert map accuracy, target-group distance can be used. Both these options are conceptually very similar to manually defining spatial offsets in point process models based on available evidence on expert map accuracies (Merow et al. 2017), yet eliminate the need for subjective decisions potentially biasing results. In sum, our approach provides an easily and widely applicable data-driven alternative for integrating expert range information in SDMs, proving particularly useful when accurate and comprehensive occurrence datasets are available.

An additional key advantage of our approach lies in its flexibility to combine expert range

maps with any combination of modeling algorithms, thereby facilitating the use of algorithm

ensembles. In contrast to the use of spatial offsets in point process models, stacked

generalizations can be easily combined with machine learning algorithms that do not include

offset terms. This enables the use of algorithms such as random forest, often found among the

best-performing in comparisons of SDM algorithms (Valavi, Guillera-Arroita, et al. 2021), also

featuring the highest discriminative accuracy (i.e., AUC values) in our dataset. With SDM

ensembles performing better than any individual modeling algorithm in our dataset, our results

also point towards the potential of stacked generalizations as a method for combining modeling

algorithms more generally. In line with findings on the importance of avoiding overfitting in

stacked generalizations (van der Laan et al. 2007; Naimi and Balzer 2018), we only achieved

improved performance when using out-of-sample predictions for training the meta-learner

("Super Learner" approach). It has been shown that in large samples, the Super Learner approach

performs at least as well as the best-performing individual algorithm (van der Laan and Dudoit

2003; van der Laan et al. 2007). Yet, despite its potential, stacked generalization has remained

neglected in the context of species distribution modeling (El Alaoui and Idri 2023), with studies

typically relying on unweighted or weighted model averaging for combining algorithms and

stacked generalization not being considered in systematic assessments of SDM ensemble

methods (Hao et al. 2020). We therefore recommend stacked generalization as a versatile

approach for combining SDM algorithms, which should be included in future comparisons of

SDM ensemble methods.

In most cases, the integration of expert range maps resulted in considerably less uniform

occurrence predictions and decreased range overlap between species, likely reflecting more

realistic predictions of bat distributions in our study area. Both SDMs and expert range maps tend

to overpredict occurrence of species since they are missing information on factors limiting

species' ranges (dispersal and competition in the case of SDMs, habitat suitability in the case of

expert ranges). Integrating both data sources can therefore improve estimates of individual

species' distributions as well as species richness (Ellis-Soto et al. 2021). Additionally, by

disentangling environmental constraints from other limiting factors, the combination of SDMs

and expert ranges can help to better understand the influence of non-environmental factors

affecting range limits (i.e., biotic interactions and dispersal). For example, contrasting potential

range overlaps derived from SDMs with realized range overlaps derived from expert-informed

models can provide a window into the potential role of interspecific competition in shaping

species' ranges (Novella-Fernandez et al. 2021). In sum, our approach has broad applicability in

ecological research and conservation planning, for example for updating species' conservation

status, assessing conservation priorities through more accurate species richness mapping, and by

providing new ecological insights into factors determining species' range limits.

Our approach adds to the growing toolbox of integrated species distribution modeling

approaches by providing a flexible and easily applicable approach for integrating SDMs with

readily available information on species' range limits. As SDMs have become one of the most

widely used tools in ecological and biogeographical research, an increasing recognition of their

shortcomings has developed (A. Lee-Yaw et al. 2022; Franklin 2010). Recently, integrated

modeling approaches have been proposed that try to enhance SDMs by combining them with

additional sources of information (Fletcher Jr. et al. 2019; Miller et al. 2019). Integrated SDM

approaches already offer key innovations for improving the mapping of species' realized

distributions (Jung 2023; Miller et al. 2019). The broader adoption of these methods, combined

with a rapid growth in the availability of biodiversity data will be critical for filling knowledge

gaps about the distribution of species and overcome the Wallacean shortfall.

References

A. Lee-Yaw, Julie, Jenny L. McCune, Samuel Pironon, and Seema N. Sheth. 2022. "Species
Distribution Models Rarely Predict the Biology of Real Populations." *Ecography*
2022(6):e05877. doi: 10.1111/ecog.05877.

Amatulli, Giuseppe, Sami Domisch, Mao-Ning Tuanmu, Benoit Parmentier, Ajay Ranipeta,
Jeremy Malczyk, and Walter Jetz. 2018. "A Suite of Global, Cross-Scale Topographic
Variables for Environmental and Biodiversity Modeling." *Scientific Data* 5(1):180040.
doi: 10.1038/sdata.2018.40.

Araújo, Miguel B., Robert P. Anderson, A. Márcia Barbosa, Colin M. Beale, Carsten F.
Dormann, Regan Early, Raquel A. Garcia, Antoine Guisan, Luigi Maiorano, Babak
Naimi, Robert B. O'Hara, Niklaus E. Zimmermann, and Carsten Rahbek. 2019.
"Standards for Distribution Models in Biodiversity Assessments." *Science Advances*
5(1):eaat4858. doi: 10.1126/sciadv.aat4858.

Araújo, Miguel B., and Mark New. 2007. "Ensemble Forecasting of Species Distributions."
Trends in Ecology & Evolution 22(1):42–47. doi: 10.1016/j.tree.2006.09.010.

Barber, Robert A., Stuart G. Ball, Roger K. A. Morris, and Francis Gilbert. 2022. "Target-Group
Backgrounds Prove Effective at Correcting Sampling Bias in Maxent Models." *Diversity
and Distributions* 28(1):128–41. doi: 10.1111/ddi.13442.

Bonannella, Carmelo, Tomislav Hengl, Johannes Heisig, Leandro Parente, Marvin N. Wright,
Martin Herold, and Sytze de Bruin. 2022. "Forest Tree Species Distribution for Europe
2000–2020: Mapping Potential and Realized Distributions Using Spatiotemporal Machine
Learning." *PeerJ* 10:e13728. doi: 10.7717/peerj.13728.

Boria, Robert A., Link E. Olson, Steven M. Goodman, and Robert P. Anderson. 2014. "Spatial
Filtering to Reduce Sampling Bias Can Improve the Performance of Ecological Niche
Models." *Ecological Modelling* 275:73–77. doi: 10.1016/j.ecolmodel.2013.12.012.

Buisson, Laëtitia, Wilfried Thuiller, Nicolas Casajus, Sovan Lek, and Gaël Grenouillet. 2010.
"Uncertainty in Ensemble Forecasting of Species Distribution." *Global Change Biology*
16(4):1145–57. doi: 10.1111/j.1365-2486.2009.02000.x.

Calabrese, Justin M., Grégoire Certain, Casper Kraan, and Carsten F. Dormann. 2014. "Stacking
Species Distribution Models and Adjusting Bias by Linking Them to Macroecological
Models." *Global Ecology and Biogeography* 23(1):99–112. doi: 10.1111/geb.12102.

Chen, Zhao, Augusto S. Auler, Michel Bakalowicz, David Drew, Franziska Griger, Jens
Hartmann, Guanghui Jiang, Nils Moosdorf, Andrea Richts, Zoran Stevanovic, George
Veni, and Nico Goldscheider. 2017. "The World Karst Aquifer Mapping Project:
Concept, Mapping Procedure and Map of Europe." *Hydrogeology Journal* 25(3):771–85.
doi: 10.1007/s10040-016-1519-3.

Cutler, Fortran original by Leo Breiman and Adele, and R. port by Andy Liaw and Matthew
Wiener. 2022. *randomForest: Breiman and Cutler's Random Forests for Classification*
and Regression [computer program]. Version 4.7-1.1. https://cran.r-
project.org/web/packages/randomForest/index.html.

Diniz-Filho, Jose Alexandre Felizola, Paulo De Marco Jr, and Bradford A. Hawkins. 2010.
"Defying the Curse of Ignorance: Perspectives in Insect Macroecology and Conservation
Biogeography." *Insect Conservation and Diversity* 3(3):172–79. doi: 10.1111/j.1752-
4598.2010.00091.x.

Domisch, Sami, Adam M. Wilson, and Walter Jetz. 2016. "Model-Based Integration of Observed
and Expert-Based Information for Assessing the Geographic and Environmental
Distribution of Freshwater Species." *Ecography* 39(11):1078–88. doi:
10.1111/ecog.01925.

Dormann, Carsten F., Jane Elith, Sven Bacher, Carsten Buchmann, Gudrun Carl, Gabriel Carré,
Jaime R. García Marquéz, Bernd Gruber, Bruno Lafourcade, Pedro J. Leitão, Tamara
Münkemüller, Colin McClean, Patrick E. Osborne, Björn Reineking, Boris Schröder,
Andrew K. Skidmore, Damaris Zurell, and Sven Lautenbach. 2013. "Collinearity: A
Review of Methods to Deal with It and a Simulation Study Evaluating Their
Performance." *Ecography* 36(1):27–46. doi: 10.1111/j.1600-0587.2012.07348.x.

El Alaoui, Omar, and Ali Idri. 2023. "Predicting the Potential Distribution of Wheatear Birds
Using Stacked Generalization-Based Ensembles." *Ecological Informatics* 75:102084. doi:
10.1016/j.ecoinf.2023.102084.

Elith, Jane, and John R. Leathwick. 2009. "Species Distribution Models: Ecological Explanation
and Prediction Across Space and Time." *Annual Review of Ecology, Evolution, and*
Systematics 40(1):677–97. doi: 10.1146/annurev.ecolsys.110308.120159.

Ellis-Soto, Diego, Cory Merow, Giuseppe Amatulli, Juan L. Parra, and Walter Jetz. 2021.
"Continental-Scale 1 Km Hummingbird Diversity Derived from Fusing Point Records
with Lateral and Elevational Expert Information." *Ecography* 44(4):640–52. doi:
10.1111/ecog.05119.

Firth, David. 1993. "Bias Reduction of Maximum Likelihood Estimates." *Biometrika* 80(1):27–
38. doi: 10.2307/2336755.

Fletcher Jr., Robert J., Trevor J. Hefley, Ellen P. Robertson, Benjamin Zuckerberg, Robert A.
McCleery, and Robert M. Dorazio. 2019. "A Practical Guide for Combining Data to
Model Species Distributions." *Ecology* 100(6):e02710. doi: 10.1002/ecy.2710.

Franklin, Janet. 2010. "Moving beyond Static Species Distribution Models in Support of
Conservation Biogeography." *Diversity and Distributions* 16(3):321–30. doi:
10.1111/j.1472-4642.2010.00641.x.

Grantham, H. S., A. Duncan, T. D. Evans, K. R. Jones, H. L. Beyer, R. Schuster, J. Walston, J. C.
Ray, J. G. Robinson, M. Callow, T. Clements, H. M. Costa, A. DeGemmis, P. R. Elsen, J.

Ervin, P. Franco, E. Goldman, S. Goetz, A. Hansen, E. Hofsvang, P. Jantz, S. Jupiter, A. Kang, P. Langhammer, W. F. Laurance, S. Lieberman, M. Linkie, Y. Malhi, S. Maxwell, M. Mendez, R. Mittermeier, N. J. Murray, H. Possingham, J. Radachowsky, S. Saatchi, C. Samper, J. Silverman, A. Shapiro, B. Strassburg, T. Stevens, E. Stokes, R. Taylor, T. Tear, R. Tizard, O. Venter, P. Visconti, S. Wang, and J. E. M. Watson. 2020. "Anthropogenic Modification of Forests Means Only 40% of Remaining Forests Have High Ecosystem Integrity." *Nature Communications* 11(1):5978. doi: 10.1038/s41467-020-19493-3.

Guisan, Antoine, and Wilfried Thuiller. 2005. "Predicting Species Distribution: Offering More than Simple Habitat Models." *Ecology Letters* 8(9):993–1009. doi: 10.1111/j.1461-0248.2005.00792.x.

Hao, Tianxiao, Jane Elith, Gurutzeta Guillera-Arroita, and José J. Lahoz-Monfort. 2019. "A Review of Evidence about Use and Performance of Species Distribution Modelling Ensembles like BIOMOD." *Diversity and Distributions* 25(5):839–52. doi: 10.1111/ddi.12892.

Hao, Tianxiao, Jane Elith, José J. Lahoz-Monfort, and Gurutzeta Guillera-Arroita. 2020. "Testing Whether Ensemble Modelling Is Advantageous for Maximising Predictive Performance of Species Distribution Models." *Ecography* 43(4):549–58. doi: https://doi.org/10.1111/ecog.04890.

Heinze, Georg, and Michael Schemper. 2002. "A Solution to the Problem of Separation in Logistic Regression." *Statistics in Medicine* 21(16):2409–19. doi: 10.1002/sim.1047.

Higino, Gracielle T., Francis Banville, Gabriel Dansereau, Norma Rocio Forero Muñoz, Fredric Windsor, and Timothée Poisot. 2023. "Mismatch between IUCN Range Maps and Species Interactions Data Illustrated Using the Serengeti Food Web." *PeerJ* 11:e14620. doi: 10.7717/peerj.14620.

Hijmans, Robert J., Steven Phillips, and John Leathwick and Jane Elith. 2020. *dismo: Species Distribution Modeling. R package.* [computer program]. Version 1.3-3. https://CRAN.R-project.org/package=dismo.

Hochkirch, Axel, Michael J. Samways, Justin Gerlach, Monika Böhm, Paul Williams, Pedro Cardoso, Neil Cumberlidge, P. J. Stephenson, Mary B. Seddon, Viola Clausnitzer, Paulo A. V. Borges, Gregory M. Mueller, Paul Pearce-Kelly, Domitilla C. Raimondo, Anja Danielczak, and Klaas-Douwe B. Dijkstra. 2021. "A Strategy for the next Decade to Address Data Deficiency in Neglected Biodiversity." *Conservation Biology* 35(2):502–9. doi: 10.1111/cobi.13589.

Hortal, Joaquín, Francesco de Bello, José Alexandre F. Diniz-Filho, Thomas M. Lewinsohn, Jorge M. Lobo, and Richard J. Ladle. 2015. "Seven Shortfalls That Beset Large-Scale Knowledge of Biodiversity." *Annual Review of Ecology, Evolution, and Systematics* 46(1):523–49. doi: 10.1146/annurev-ecolsys-112414-054400.

Hothorn, Torsten, Peter Buehlmann, Thomas Kneib, Matthias Schmid, Benjamin Hofner, Fabian Otto-Sobotka, Fabian Scheipl, and Andreas Mayr. 2022. *mboost: Model-Based Boosting* [computer program]. Version 2.9-7. https://cran.r-project.org/web/packages/mboost/index.html.

Inman, Richard, Janet Franklin, Todd Esque, and Kenneth Nussear. 2021. "Comparing Sample Bias Correction Methods for Species Distribution Modeling Using Virtual Species." *Ecosphere* 12(3):e03422. doi: 10.1002/ecs2.3422.

IUCN. 2022. "The IUCN Red List of Threatened Species." *IUCN Red List of Threatened Species*. Retrieved May 2, 2023 (https://www.iucnredlist.org/en).

Jung, Martin. 2023. "An Integrated Species Distribution Modelling Framework for Heterogeneous Biodiversity Data." *Ecological Informatics* 102127. doi: 10.1016/j.ecoinf.2023.102127.

Karger, Dirk Nikolaus, Olaf Conrad, Jürgen Böhner, Tobias Kawohl, Holger Kreft, Rodrigo Wilber Soria-Auza, Niklaus E. Zimmermann, H. Peter Linder, and Michael Kessler. 2017. "Climatologies at High Resolution for the Earth's Land Surface Areas." *Scientific Data* 4(1):170122. doi: 10.1038/sdata.2017.122.

Kosmidis, Ioannis, Euloge Clovis Kenne Pagui, Kjell Konis, and Nicola Sartori. 2023. *brglm2: Bias Reduction in Generalized Linear Models* [computer program]. Version 0.9. https://cran.r-project.org/web/packages/brglm2/index.html.

van der Laan, Mark, and Sandrine Dudoit. 2003. "Unified Cross-Validation Methodology For Selection Among Estimators and a General Cross-Validated Adaptive Epsilon-Net Estimator: Finite Sample Oracle Inequalities and Examples." *U.C. Berkeley Division of Biostatistics Working Paper Series*.

van der Laan, Mark, Eric C. Polley, and Alan E. Hubbard. 2007. "Super Learner." *Statistical Applications in Genetics and Molecular Biology* 6(1). doi: 10.2202/1544-6115.1309.

Mainali, Kumar, Trevor Hefley, Leslie Ries, and William F. Fagan. 2020. "Matching Expert Range Maps with Species Distribution Model Predictions." *Conservation Biology* 34(5):1292–1304. doi: 10.1111/cobi.13492.

Merow, Cory, Adam M. Wilson, and Walter Jetz. 2017. "Integrating Occurrence Data and Expert Maps for Improved Species Range Predictions." *Global Ecology and Biogeography* 26(2):243–58. doi: 10.1111/geb.12539.

Milanesi, Pietro, Francesca Della Rocca, and Robert A. Robinson. 2020. "Integrating Dynamic Environmental Predictors and Species Occurrences: Toward True Dynamic Species Distribution Models." *Ecology and Evolution* 10(2):1087–92. doi: 10.1002/ece3.5938.

Miller, David A. W., Krishna Pacifici, Jamie S. Sanderlin, and Brian J. Reich. 2019. "The Recent Past and Promising Future for Data Integration Methods to Estimate Species'

Distributions." *Methods in Ecology and Evolution* 10(1):22–37. doi: 10.1111/2041-210X.13110.

Myers, Norman, Russell A. Mittermeier, Cristina G. Mittermeier, Gustavo A. B. da Fonseca, and Jennifer Kent. 2000. "Biodiversity Hotspots for Conservation Priorities." *Nature* 403(6772):853–58. doi: 10.1038/35002501.

Naimi, Ashley I., and Laura B. Balzer. 2018. "Stacked Generalization: An Introduction to Super Learning." *European Journal of Epidemiology* 33(5):459–64. doi: 10.1007/s10654-018-0390-z.

Novella-Fernandez, Roberto, Javier Juste, Carlos Ibáñez, Hugo Rebelo, Danilo Russo, Antton Alberdi, Andreas Kiefer, Laura Graham, Hynek Paul, Charles Patrick Doncaster, and Orly Razgour. 2021. "Broad-Scale Patterns of Geographic Avoidance between Species Emerge in the Absence of Fine-Scale Mechanisms of Coexistence." *Diversity and Distributions* 27(9):1606–18. doi: 10.1111/ddi.13375.

Oeser, Julian, Marco Heurich, Cornelius Senf, Dirk Pflugmacher, Elisa Belotti, and Tobias Kuemmerle. 2020. "Habitat Metrics Based on Multi-Temporal Landsat Imagery for Mapping Large Mammal Habitat." *Remote Sensing in Ecology and Conservation* 6(1):52–69. doi: 10.1002/rse2.122.

Olson, David M., Eric Dinerstein, Eric D. Wikramanayake, Neil D. Burgess, George V. N. Powell, Emma C. Underwood, Jennifer A. D'amico, Illanga Itoua, Holly E. Strand, John C. Morrison, Colby J. Loucks, Thomas F. Allnutt, Taylor H. Ricketts, Yumiko Kura, John F. Lamoreux, Wesley W. Wettengel, Prashant Hedao, and Kenneth R. Kassem. 2001. "Terrestrial Ecoregions of the World: A New Map of Life on Earth: A New Global Map of Terrestrial Ecoregions Provides an Innovative Tool for Conserving Biodiversity." *BioScience* 51(11):933–38. doi: 10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2.

Ovaskainen, Otso, David B. Roy, Richard Fox, and Barbara J. Anderson. 2016. "Uncovering Hidden Spatial Structure in Species Communities with Spatially Explicit Joint Species Distribution Models." *Methods in Ecology and Evolution* 7(4):428–36. doi: 10.1111/2041-210X.12502.

Pimm, S. L., C. N. Jenkins, R. Abell, T. M. Brooks, J. L. Gittleman, L. N. Joppa, P. H. Raven, C. M. Roberts, and J. O. Sexton. 2014. "The Biodiversity of Species and Their Rates of Extinction, Distribution, and Protection." *Science* 344(6187):1246752. doi: 10.1126/science.1246752.

Ramesh, Vijay, Trisha Gopalakrishna, Sahas Barve, and Don J. Melnick. 2017. "IUCN Greatly Underestimates Threat Levels of Endemic Birds in the Western Ghats." *Biological Conservation* 210:205–21. doi: 10.1016/j.biocon.2017.03.019.

Sesmero, M. Paz, Agapito I. Ledezma, and Araceli Sanchis. 2015. "Generating Ensembles of Heterogeneous Classifiers Using Stacked Generalization." *WIREs Data Mining and Knowledge Discovery* 5(1):21–34. doi: 10.1002/widm.1143.

Soberón, Jorge. 2007. "Grinnellian and Eltonian Niches and Geographic Distributions of
Species." *Ecology Letters* 10(12):1115–23. doi: 10.1111/j.1461-0248.2007.01107.x.

Steen, Valerie A., Morgan W. Tingley, Peter W. C. Paton, and Chris S. Elphick. 2021. "Spatial
Thinning and Class Balancing: Key Choices Lead to Variation in the Performance of
Species Distribution Models with Citizen Science Data." *Methods in Ecology and*
Evolution 12(2):216–26. doi: 10.1111/2041-210X.13525.

Syfert, Mindy M., Matthew J. Smith, and David A. Coomes. 2013. "The Effects of Sampling
Bias and Model Complexity on the Predictive Performance of MaxEnt Species
Distribution Models." *PLOS ONE* 8(2):e55158. doi: 10.1371/journal.pone.0055158.

Theobald, David M., Christina Kennedy, Bin Chen, James Oakleaf, Sharon Baruch-Mordo, and
Joe Kiesecker. 2020. "Earth Transformed: Detailed Mapping of Global Human
Modification from 1990 to 2017." *Earth System Science Data* 12(3):1953–72. doi:
10.5194/essd-12-1953-2020.

Valavi, Roozbeh, Jane Elith, José J. Lahoz-Monfort, and Gurutzeta Guillera-Arroita. 2021.
"Modelling Species Presence-Only Data with Random Forests." *Ecography* 44(12):1731–
42. doi: 10.1111/ecog.05615.

Valavi, Roozbeh, Gurutzeta Guillera-Arroita, José J. Lahoz-Monfort, and Jane Elith. 2021.
"Predictive Performance of Presence-Only Species Distribution Models: A Benchmark
Study with Reproducible Code." *Ecological Monographs* n/a(n/a):e01486. doi:
10.1002/ecm.1486.

Warren, Dan L., Richard E. Glor, and Michael Turelli. 2008. "Environmental Niche Equivalency
Versus Conservatism: Quantitative Approaches to Niche Evolution." *Evolution*
62(11):2868–83. doi: 10.1111/j.1558-5646.2008.00482.x.

Warren, Dan L., Nicholas J. Matzke, Marcel Cardillo, John B. Baumgartner, Linda J. Beaumont,
Michael Turelli, Richard E. Glor, Nicholas A. Huron, Marianna Simões, Teresa L.
Iglesias, Julien C. Piquet, and Russell Dinnage. 2021. "ENMTools 1.0: An R Package for
Comparative Ecological Biogeography." *Ecography* 44(4):504–11. doi:
10.1111/ecog.05485.

Weiss, D. J., A. Nelson, H. S. Gibson, W. Temperley, S. Peedell, A. Lieber, M. Hancher, E.
Poyart, S. Belchior, N. Fullman, B. Mappin, U. Dalrymple, J. Rozier, T. C. D. Lucas, R.
E. Howes, L. S. Tusting, S. Y. Kang, E. Cameron, D. Bisanzio, K. E. Battle, S. Bhatt, and
P. W. Gething. 2018. "A Global Map of Travel Time to Cities to Assess Inequalities in
Accessibility in 2015." *Nature* 553(7688):333–36. doi: 10.1038/nature25181.

Wolpert, David H. 1992. "Stacked Generalization." *Neural Networks* 5(2):241–59. doi:
10.1016/S0893-6080(05)80023-1.

Zhao, Chenchen, Xin Cao, Xuehong Chen, and Xihong Cui. 2022. "A Consistent and Corrected
Nighttime Light Dataset (CCNL 1992–2013) from DMSP-OLS Data." *Scientific Data*
9(1):424. doi: 10.1038/s41597-022-01540-x.

Zurell, Damaris. 2017. "Integrating Demography, Dispersal and Interspecific Interactions into
Bird Distribution Models." *Journal of Avian Biology* 48(12):1505–16. doi:
10.1111/jav.01225.