The *Mycobacterium tuberculosis* complex pangenome is small and driven by

sub-lineage-specific regions of difference

Mahboobeh Behruznia[1,2], Maximillian Marin[3], Maha Farhat[3,4], Jonathan C Thomas[1], Maria

Rosa Domingo-Sananes[1], Conor J Meehan[1,5*]

[1] Department of Biosciences, Nottingham Trent University, Nottingham, UK

[2] Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham,
Birmingham, UK

[3] Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115, USA

[4] Pulmonary and Critical Care Medicine, Massachusetts General Hospital, Boston, MA 02114, USA

[5] Unit of Mycobacteriology, Institute of Tropical Medicine, Antwerp, Belgium

Corresponding author: conor.meehan@ntu.ac.uk

Authors ORCHID IDs: 0000-0002-3302-7272 (MB), 0000-0002-9108-3328 (MM), 0000-0002-

3871-5760 (MF), 0000-0002-1599-9123 (JCT), 0000-0002-3339-8671 (MRDS), 0000-0003-

0724-8343 (CJM)

Abstract

The *Mycobacterium tuberculosis* complex (MTBC) is a group of bacteria causing tuberculosis

(TB) in humans and animals. Understanding MTBC genetic diversity is crucial for insights into

its adaptation and traits related to survival, virulence, and antibiotic resistance. While it is

known that within MTBC diversity is characterised by large lineage-specific deletions

(regions of difference [RD]), a comprehensive pangenomic analysis incorporating both

coding and non-coding regions remains unexplored. We utilised a curated dataset

representing various MTBC genomes, including under-represented lineages to quantify the

true diversity of the MTBC pangenome. The MTBC was found to have a small, closed

pangenome with distinct genomic features and RDs both between lineages (as previously

known) and between sub-lineages. The accessory genome was identified to be a product of

genome reduction, showing both lineage-specific and independent deletions. This variation

has implications for traits like virulence, drug resistance, and metabolism. The study provides

a comprehensive understanding of the MTBC pangenome, highlighting the importance of

genome reduction in its evolution and showing that within-lineage genome content diversity

is present. The findings underline the significance of genomic variations in determining the

pathogenic traits of different MTBC lineages.

Introduction

The *Mycobacterium tuberculosis* complex (MTBC) is a group of closely related bacteria that cause tuberculosis (TB) in humans and other animals. Worldwide, TB is the 13th leading cause of death and is anticipated to once again become the top infectious killer following global efforts to combat the COVID-19 pandemic (WHO, 2023). The MTBC comprises both human-associated lineages (L1 to L9) and animal-associated lineages (La1, La2, La3, *M. microti* and *M. pinnipedii*) (Coscolla et al., 2021, Zwyer et al., 2021). The human-adapted MTBC exhibits a strong phylogeographical population structure, with some lineages occurring globally and others showing a strong geographical restriction (Gagneux, 2018). For example, L2 and L4 are widespread globally, with L2 dominating in Southeast Asia and L4 in Europe (Napier et al., 2020). L1 and L3 occur mainly in regions around the Indian Ocean (Menardo et al., 2021). L5 and L6 are highly restricted to West Africa (de Jong et al., 2010) and L7, L8 and L9 are exclusively found in East Africa (Blouin et al., 2012, Ngabonziza et al., 2020, Coscolla et al., 2021). Animal-associated lineages refer to lineages of the MTBC that primarily infect and cause disease in animals but can also infect humans in close contact with infected animals (Brites et al., 2018). Little is known about the geographic distribution and host range of animal-associated lineages (Zwyer et al., 2021).

Various factors such as human migration, local adaptation to host population, and host immune response are thought to have contributed to the current phylogeographical distribution of human-adapted MTBC members (Darwin and Stanley, 2022, Hershberg et al., 2008). Additionally, different lineages of MTBC are known to differ in virulence, metabolism and transmission potential, which can affect the spread of certain lineages and sub-lineages (Coscolla, 2017, Gagneux et al., 2006). Understanding the genetic diversity within MTBC is

essential for gaining insights into how these bacteria adapt to different host populations and

environments, and identify traits that may be important for survival, virulence, and antibiotic

resistance.

MTBC is recognised for its clonal population structure and little genetic diversity between

different strains, caused by the absence of horizontal gene transfer, plasmids, and

recombination (Godfroid et al., 2018, Chiner-Oms et al., 2022). Single nucleotide

polymorphisms (SNPs) and large sequence polymorphisms in the so-called regions of

difference (RDs) represent the source of MTBC genetic diversity (Gagneux, 2018). RDs are

known to differentiate different MTBC lineages from each other (Gagneux, 2018), but within

lineage diversity is thought to be mostly driven by SNPs (Napier et al., 2020). While this has

been shown to not be the case in some lineages (Sanoussi et al., 2021), these larger genomic

structural differences have not been fully explored. Deletions in some RDs can surpass 10 kb

in length and encompass groups of genes with diverse functions, which can be linked to

factors like virulence and metabolic diversity (Bespiatykh et al., 2021, Bottai et al., 2020).

Additionally, the *IS6110* element has a notable impact on the genetic diversity of the MTBC.

Transposition of *IS6110* into various coding or regulatory regions, and genomic deletions

through homologous recombination between proximal copies, can also translate into

significant genotypic variation at strain level (Alonso et al., 2013, Antoine et al., 2021).

The pangenome represents the set of all genes present in a species. It comprises both the

core genome, which consists of genes shared by all members of the species, and the

accessory genome, which includes genes present in some but not all members (Brockhurst

et al., 2019). Previous research has focused on pangenomic investigations of particular MTBC

lineages (Baena et al., 2023, Ceres et al., 2022, Sanoussi et al., 2021). A recent study

investigated the MTBC pangenome focusing on protein-coding sequences (Silva-Pereira et

al., 2024). However, there is a gap in comprehensive pangenomic analyses encompassing

both coding and non-coding regions. Despite the name, pangenome analyses tend to not

focus on full genome comparisons, but rather on protein-coding sequences. This focus

neglects non-coding RNAs and pseudogenes as well as essential elements in intergenic

regions, such as promoters, terminators, and regulatory binding sites, despite their

demonstrated role in selection and phenotypic impact (Tonkin-Hill et al., 2023). Moreover,

the inference drawn from pangenomic analyses is often influenced by the use of short-read

datasets. It has been shown that using short-read datasets can artificially inflate the size of

the pangenome estimates, leading to incorrect assumptions about evolutionary scenarios

(Marin, 2024). Here, we used a diverse collection of closed MTBC genomes including human

and animal-associated lineages to develop our understanding of MTBC pangenome and

evolution incorporating both coding and non-coding regions. The objectives of this study

are: (i) to establish the true size of MTBC pangenome using a highly curated dataset, (ii) to

assess specific biological functions that are prevalent in the core and accessory genome and

(iii) to assess genetic variation between lineages and sub-lineages that may affect virulence,

drug resistance, metabolism and evolutionary characteristics. We find that the MTBC has a

small, closed pangenome and genome reduction is the driving evolutionary force that

shapes the genomic evolution of the species. We identified distinct genomic features among

lineages and sub-lineages, including sub-lineage specific RDs, that can account for variations

in virulence, metabolism, and antibiotic resistance.

Methods

Summary of sequencing data and quality control

Previous studies have shown that closed genomes should be used for best estimation of

pangenome size (Marin et al., 2024). From a combination of previously published research

and publicly available data, 322 non-duplicated complete and closed (single contig) genome

assemblies were selected for having (i) long-read sequencing (Nanopore or PacBio) or (ii)

hybrid sequencing with Illumina across the MTBC lineages. Information related to the

geographical location, source of isolation, and assembly method can be found in

Supplementary File 1. The quality of all assemblies was assessed using BUSCO v5.4.7 (Manni

et al., 2021) and only assemblies with ≥ 95% completeness were used for further analysis.

The assemblies were set to start on *dnaA* gene using the fixstart command of Circlator v1.5.5

(Hunt et al., 2015). Complete circular assemblies were annotated using PGAP v6.4 with

default options (Tatusova et al., 2016). This approach is recommended by previous analyses

(Marin et al., 2024) as PGAP tends to annotate pseudogenes in cases of pseudogenisation

and frameshift, rather than annotating them as new coding sequences. Visualisations of the

PGAP annotation were made using the DNA Features Viewer Python library (Zulkower and

Rosser, 2020). Lineage classification was performed using TB-profiler v4.4.2 (Phelan et al.,

2019).

Nanopore sequencing and assembly

We sequenced lineages and sub-lineages that either had no sequencing data available in

public databases or were under-represented in the collection. These isolates belonged to L5

(n=2, L5.1 and L5.2), L4.6 (n=3, L4.6.1, L4.6.1.1 and L4.6.1.2), L4.7 (n=3), L9 (n=1), La2 (*M.*

caprae, n =1) and *M. pinnipedii* (n=1). A list of genomes sequenced in this study and their

assembly information can be found in Table S1. Assembled genomes were deposited at the

NCBI under the Bioproject PRJNA1085078.

DNA extraction suitable for PacBio SMRT sequencing was performed using the Genomic-tip

100/G (Qiagen Inc) as described previously (Ngabonziza et al., 2020). DNA was enriched in

samples with low DNA concentration (1-60 ng/μl) using AMPureXP Beads (SPRIselect,

Beckman Coulter) in a 1:1 ratio. Samples were incubated for 10 minutes at room

temperature followed by pelleting the beads on a magnetic stand. The supernatant was

discarded, and DNA was eluted from the bead using 10-20 μl of nuclease-free water

(ThermoFisher Scientific). DNA concentration was quantified using a Qubit fluorometer and

1X dsDNA High Sensitivity (HS) assay kit (ThermoFisher Scientific). Nanopore sequencing was

performed using the native barcoding kit 24 V14 (SQK-NBD114.24) according to the

manufacturer's instruction on the MinION Mk1C platform with R10.4.1 flow cells. Sequence

adaptors were trimmed using Porechop v0.2.4 with "--end_threshold 95" and "--

middle_threshold 85" options (https://github.com/rrwick/Porechop.git). Sequences were

assembled using flye v2.9.2 with default options (Kolmogorov et al., 2019) and iteratively

polished using Racon v1.5.0 (Vaser et al., 2017) and the "consensus" subcommand of

Medaka v1.8.0 (https://github.com/nanoporetech/medaka.git). Complete assemblies

underwent Circlator, BUSCO and PGAP steps as above.

Pangenome analysis

Panaroo was used to characterise the MTBC protein-coding pangenome (Tonkin-Hill et al.,

2020) using the "--merge_paralogs", "--clean-mode strict" and "--core_threshold 0.95"

options. Ceres et al. (2022) showed that genes impacted by pseudogenisation could emerge

as accessory genes. To find split genes due to pseudogenisation, the reference pangenome

created by Panaroo was compared to the H37Rv genome using BLASTn with a similarity e-

value cutoff of 1×10^{-10} (Zhang et al., 2000). The BLASTn result was then filtered to only

include queries with ≥90% identity and ≥75% of the H37Rv gene length covered using in-

house Python scripts (https://github.com/conmeehan/pathophy). A core genome alignment

was constructed using the built-in MAFFT aligner included in Panaroo (Katoh et al., 2002).

The core phylogeny was created using RAxML-NG v1.2.0 with a GTR substitution model

(Kozlov et al., 2019). Tree visualisations were created using the Treeio package v1.20.0

(Wang et al., 2020).

Pangenome graph construction

To look at total pangenome content (both coding and non-coding), multiple whole genomes

were aligned into a graph using Pangraph v0.7.2 (Noll et al., 2023). The DNA sequences from

gene blocks present in at least one sub-lineage but completely absent in others were

extracted. These sequences were subjected to BLAST against the H37Rv genome, along with

RvD1 and TbD1 genes to identify their corresponding Rv loci within the H37Rv genome and

determine sub-lineage-specific regions of difference.

Functional genome analysis

The reference pangenome created by Panaroo was translated using transeq v5.0.0 (Rice et

al., 2000) to obtain the protein sequences. COG functional categories were assigned using

eggNOG Mapper v5.0.2 using default options (Huerta-Cepas et al., 2017). Genes that did not

return a match within the eggNOG database were included in the not found category.

Functional domain annotation was performed with InterProScan v5.63-95 (Jones et al.,

2014). For the gene association analysis, the accessory genome (genes present in less than

95% of genomes) was used as input for Coinfinder v1.2.1 (Whelan et al., 2020). STRING

database (https://string-db.org/) was used to infer interaction networks of genes. Predicted

functional partners with a high confidence interaction score (≥0.7) by STRING v12.0 were

considered in this study.

Statistical analysis

Statistical analysis was conducted using R studio v2023.12.0+369 (R Core Team, 2023).

Principal component analysis (PCA) based on gene presence and absence data from Panaroo

and Pangraph derived accessory genome contents was performed using the prcomp package

in R. The fviz_contrib function in Factoextra (v1.0.7) was used to identify the contribution of

important variables from the PCA results (Kassambara and Mundt, 2020).

Results

Overview of the genomic dataset

We combined a total of 335 whole-genome sequences covering all known MTBC lineages. This

included 324 genomes published previously, as well as 11 genomes from sub-lineages with few or no

representatives in our collection to better cover the known phylogeny of MTBC (Supplementary Table

S1). Except Antarctica, our collection includes genomes from every continent, and different host

species including humans, cattle, voles and BCG vaccine strains (Figure 1-A). Africa represents the

highest diversity of MTBC genomes among other continents, which is consistent with the fact that

the continent is known to have the most diverse MTBC strains (Figure 1-B) (Gagneux, 2018). Most L1

genomes have been collected from East Africa, South and Southeast Asia, where this lineage is

geographically common. L3 genomes were mostly isolated from India, while L2 and L4 genomes have

been collected from different continents, representing the global distribution of these lineages. L5

and L6 genomes were exclusively found in West Africa, where they contribute significantly to the

overall burden of TB in this region (Coscolla et al., 2021).

Pangenome characteristics and functional diversity

Panaroo estimated a protein-coding pangenome consisting of 4,252 genes, including 3,639

core genes (present in more than 99% of genomes) and 613 accessory genes including 258

soft core, 218 shell and 137 cloud genes. After mapping all the genes in the pangenome to

H37Rv, we identified 127 genes that were misclassified as accessory genes, including 25 out

of 258 soft core, 24 out of 218 shell and 78 out of 137 cloud genes (Supplementary Table

S2). This approach increased the core genome size from 3,639 to 3,746 and reduced the

accessory genome from 613 to 506. This shift occurred because 107 genes out of 127 genes

detected as false accessories were re-classified as part of the core genome. To examine

genomic diversity among lineages, we investigated variations in the size of the accessory

genome among different lineages. This analysis revealed significant differences in accessory

genome size among different lineages (Kruskal-Wallis test, p= 2.2e-16). Post-hoc Dunn tests

with Bonferroni correction identified significant pairwise differences between lineages

(Supplementary Table S3). L6, La1 and *M. microti* have significantly smaller accessory

genomes compared to others (Figure 2, Supplementary Table S3). L2 has a significantly

smaller accessory genome in comparison with L1, L3, and L4.

Functional annotation of core and accessory genes showed that genes of unknown function were the

largest functional group in both (26.8 % and 12.0% respectively). The core genome of MTBC encodes

a high proportion of genes involved in lipid transport and metabolism (category I) and cellular

transport and secretion systems (category N), indicating the importance of these biological functions

in defining the species. Functional breakdown of the accessory genome also revealed variations in

genes involved in DNA replication, repair and mobile elements (category L) and lipid transport and

metabolism (category I) across the species. A breakdown of the functional composition of the core

and accessory genome is shown in Figure 3.

Population structure analysis

The population structure was assessed using SNPs in the core genome and PCA plots were

constructed based on presence-absence patterns in the accessory genome determined with Panaroo

and Pangraph (Figure 4). In the PCAs, genomes group together based on their lineage, indicating

variations in accessory genome are driven mainly by lineage structure (Figure 4-B and 4-C). The first

principal component (PC1) highlights the contrast between L2 from other members of MTBC. In both

PCA plots, L2 formed a distinct group, defined primarily by RD105 and RD207 deletions. RD105

serves as a classic marker for all L2 genomes and RD207 deletion is found in all L2.2 sub-lineages; the

single L2.1 genome is separated from the L2.2 cluster due to the absence of RD207. The second

principal component (PC2) demonstrates the difference between modern human-associated lineages

(L2, L3 and L4) from other lineages. Important variables driving this separation are the TbD1 gene

deletion in modern lineages, and gene deletions associated with RD11 (prophage phiRv2) in L6, L7,

La1 and La3 genomes and some L1 sub-lineages. L1 genomes exhibit considerable genetic diversity

primarily due to RD3 (related to prophage phiRv1) and RD11. The latter was identified in all L1.1 sub-

lineages and certain L1.2 sub-lineages (L1.2.1, L1.2.2, and L1.2.2.2). The positioning of the single L8

genome, marked with a positive sign at the centre of PCAs, implies that L8, serving as the ancestor of

the MTBC, exhibits similarities in the accessory genome with other lineages.

The accessory genome of MTBC

Among the 506 filtered accessory genes from Panaroo output, 107 were identified as being specific

to certain lineages and sub-lineages (Figure 5-A, Supplementary Table S4). Similarly, we identified 82

lineage and sub-lineage-specific genomic regions ranging from 270 bp to 9.8kb, encompassing a total

of 116 genes and 4 non-coding RNA using the Pangraph approach (Figure 5-B, Supplementary Table

S4 and Supplementary File S1). Our analysis revealed both previously known RDs and new deletions

in the accessory genome, suggesting that it originates from genome deletions. Lineage and sub-

lineage-specific genomic regions were consistently identified using both methods, including TbD1,

RD9, RD10, RD105, RD303, RD312, RD316, RD702, RD711, RD713, RD750, ND1, RDoryx, and N-

RD25bov/cap (Figure 6).

Panaroo failed to identify partial gene deletions caused by RD9, RD10, RD13, RD105, RD207,

and RD702. For example, RD702 is present in all L6 and L9 genomes and covers a 641 bp

region (Figure 7). RD702 affects less than a quarter of the *bglS* gene (470 bp from 2,076 bp)

and *mymT* gene (126 bp from 162 bp gene) in the 5' to 3' direction as well as a regulatory

RNA (ncRv0186c) in the 3' to 5' direction. Panaroo did not identify the 470 bp deletion

downstream of the *bglS* and this gene was classified as being in the core genome. These

results could be explained by the fact that Panaroo uses a relaxed alignment threshold along

with neighbourhood gene information to cluster gene families (Tonkin-Hill et al., 2020). This

can lead to the grouping of partially deleted genes with those that are complete. The

Pangraph alignment approach identified partial gene deletion and non-coding regions of the

DNA that were impacted by genomic deletion. However, the graph approach failed to

identify four structural variants, including a gene deletion corresponding to RD8 (Rv3618) in

L6 and animal-associated genomes. Despite these limitations, both methods yielded

consistent results indicating that the accessory genome of MTBC is small and is acquired

vertically from a common ancestor within the lineage. Gene association analysis using

coinfinder confirmed that most accessory genes (as determined by Panaroo) appear in pairs

or groups in larger multi-gene RDs (as determined by Pangraph) (Supplementary Figure S1).

Thus, the small accessory genome is contained within an even smaller number of RDs.

Coincident gene blocks that were associated with lineages were RD8, RD9, RD10, and

RD178, whereas those that were distributed across the phylogeny belonged to RD3 and

RD11.

Discussion

MTBC accessory genome is small and is acquired vertically

The MTBC pangenome size has been reported to vary across studies. In a recent investigation conducted by Marin (2024), the accessory genome ranged from 314 to 2951 genes, as a function of the assembly quality, software and specific parameters used. Genome sequencing using short-read technology has some limitations that can result in incomplete assemblies, particularly in regions with repetitive sequences that are difficult to sequence (Meehan et al., 2019). About 10% of the coding sequences of the MTBC genome are composed of repetitive polymorphic sequences known as the PE (Proline-Glutamic acid) and PPE (Proline-Proline-Glutamic acid) families, implicated in host-pathogen interactions and virulence (Cole et al., 1998). Incomplete assemblies can impact the accurate detection of genes and may lead to artificial inflation of the estimated pangenome size (Ceres et al., 2022, Marin et al., 2024). We used a curated dataset to address these issues and estimate the true size of the MTBC pangenome. We showed that MTBC has a compact accessory genome that is acquired vertically. This finding is consistent with recent research, which found a small accessory genome with a clonal genetic structure in MTBC (Baena et al., 2023, Ceres et al., 2022, Silva-Pereira et al., 2024). Using a BLASTn analysis approach, we identified core genes that were designated as accessory and further reduced the accessory genome from 613 to 506. For example, indel mutations observed in the *lppA* gene across 128 genomes resulted in the misclassification of this gene as an accessory (labelled as group_1700). Notably, over 50% of the cloud genes (genes present in less than 15% of strains) were found to be core genes. These genes mapped to the reference genes using the BLAST approach and appeared to be fragmented genes caused by misassemblies.

Attributes of the accessory genome

Attributes of the accessory genome

We employed a pangenomic approach to analyse the genetic diversity within the MTBC,

aiming to understand its evolution and functional potential. In addition to inferring a gene-

based pangenome with Panaroo, we also analysed variation in non-coding regions by

inferring a pangenome graph with Pangraph. Examining non-coding regions is particularly

important for species with limited genetic diversity, as they play a vital role in controlling

gene expression, stress response and host adaptation (Arnvig et al., 2014). Our study

demonstrated that, although both methods produced reliable outcomes, the whole-genome

alignment approach could capture higher genetic variations, especially those involving large

structural variants (>200 bp). Moreover, this method identified non-coding regions of the

genome that were affected by genomic deletions. Results from both methodologies revealed

that the accessory genome of MTBC is a product of gene deletions, that can be classified

into lineage-specific deletions and lineage-independent deletions, with the former category

being the more common. Coinfinder results also showed that coincident genes in the

accessory genome either belonged to particular lineages or were dispersed throughout the

phylogeny. Region of difference, which are large deletions often encompassing multiple

genes, can act as distinctive markers not only for lineages, as demonstrated previously, but

also for sub-lineages. These warrant further investigation to understand the biological

consequences of such deletions on traits like virulence, drug resistance, and metabolism,

along with SNP only analyses. On the other hand, we observed several genomic deletions

that appear to be lineage-independent and may indicate gene instability (Brosch et al.,

2002). This includes RD3 and RD11 corresponding to the two prophages phiRv1 and phiRv2.

One drawback of our study was that many of the lineages were poorly represented,

especially in the L5, L6, L9 and all animal-associated lineages. The production of more closed

genomes from the geographic regions where these are abundant (e.g. East and West Africa)

will provide further information about the RD distributions within the MTBC and better

understanding of the driving evolutionary forces.

Differential traits between lineages and sub-lineages

The size of the accessory genome content varied significantly among different lineages. L2

genomes exhibited a high degree of clonality and a smaller accessory genome. L2 strains are

characterised by a large number of *IS6110* copies, possibly due to the RD207, which affects

the CRISPR-Cas type III-A system, responsible for adaptive immunity against mobile genetic

elements (Alonso et al., 2013). While RD207 encompasses seven genes (Rv2814c-Rv2820c)

(Bespiatykh et al., 2021), our analysis revealed that only two genes, Rv2819c and Rv2820c,

are specifically absent in L2 and can serve as a genetic marker for this lineage. Other RDs

affecting all L2 genomes are RD3 (9.2 kb), and RD105 (3.4 kb). The latter is exclusively

present in all L2 genomes and is caused by the insertion of *IS6110* (Alonso et al., 2013).

Previous research proposed a potential role of RD105 in conferring drug resistance in L2.2

strains (Qin et al., 2019). RD207 (7.4 kb), linked to the insertion of the *IS6110* element, is

also present in all L2 genomes except for L2.1. L3 genomes exhibited significantly less

genomic deletion in comparison with L2, and RDs affecting all L3 genomes were RD316 (1.2

kb) and RD750 (765 bp). The genes affected by these RDs are involved in lipid metabolism

(category I) and secondary metabolite biosynthesis and transport respectively (category Q),

which may contribute to metabolic variations in L3 strains. L4 exhibits substantial genetic

diversity primarily due to variations in the presence and absence of specific RDs at the sub-

lineage level. For example, RD145 (4.7 kb) and RD182 (6.5 kb) were detected in L4.1.2.1

genomes isolated from Africa, Asia, Europe and South America. However, none of the

affected genes appears advantageous or essential for MTBC growth and survival and this

sub-lineage is successful worldwide (https://tbdr.lshtm.ac.uk/sra/browse). Other L4 sub-

lineage-specific RDs detected in this study with no known advantage are RD115 in L4.3.4,

RD727 in L4.6.1, RD219 in L4.8, RvD1 in L4.8 and L4.9. RD4 in La1, L5.3 Del in L5.3, and

RD750 in L3, RD147c in L1.

Nonetheless, some genomic deletions can affect the growth advantage gene group in

various lineages. Disruption of these genes has been shown to provide a fitness advantage

for the *in vitro* growth of H37Rv by analysis of saturated Himar 1 transposon libraries

(DeJesus et al., 2017). For example, RD174 specific to L4.3.4 intersects with RD743 specific

to L5, and both deletions affect two genes (Rv1995 and Rv1996) including *cmtR*

transcriptional regulator. The latter regulates bacterial stress response to metal deficiency

and is important during persistence in phagocyte (Flores-Valdez et al., 2020). While the

absence of these genes may provide a fitness advantage for *in vitro* growth, the impact of

this deletion is likely to be different in *in vivo* conditions. The low prevalence of L4.3.4 strains

in the TB Profiler database may suggest either the lower fitness of this sub-lineage, or its

presence in under-sampled geographic areas. RDs impacting the growth advantage gene

group were prevalent in L5 genomes (L5.3 Del, RD743 and RD711). For instance, RD711

affects the Rv1335 gene, which is annotated as a sulphur carrier protein involved in cysteine

biosynthesis by PGAP. This may imply that the genome reduction observed in L5 has evolved

to depend on host factors for essential nutrients, consequently making these strains difficult

to grow in a culture medium (Sanoussi et al., 2017).

L6 and animal-associated lineages accumulated a higher number of deletions including RD7,

RD8, RD9 and RD10. Comparative analysis of the accessory genome size in L6, La1, and *M.*

microti revealed significant differences compared to other lineages. The identified genomic

regions in L6, L9, and animal-associated lineages impact genes associated with virulence,

potentially explaining their reduced virulence in human hosts. RD7 contains the mammalian

cell entry (*mce3*) operon implicated in the invasion of host cells, while RD9 includes genes

related to biliverdin reduction, potentially aiding in protection against oxidative stress inside

macrophages (Szklarczyk et al., 2023). RD10 impacts genes related to lipid storage crucial for

MTBC transition to a dormant state (Deb et al., 2009). RD1BCG and RD1mic affect genes

encoding the ESX-1 secretion system, crucial for virulence and phagosomal escape, with

RD1BCG being deleted during BCG vaccine development (Szklarczyk et al., 2023). Overall,

the findings of this study suggest that, despite having a compact accessory genome, it can

account for certain biological distinctions among MTBC. Previous research has highlighted

the significance of random genetic drift in shaping the evolutionary dynamics of the species

(Hershberg et al., 2008). Gene loss, driven by genetic drift, is likely to be a key contributor to

the observed genetic diversity within the MTBC.

A need for more high-quality genomes

While this dataset represents high-quality MTBC genome sequences that cover most of its

known diversity, it suffers from sampling bias as some lineages are over-represented

whereas others have a few representatives. The over-representation of L2 and L4 in the

dataset is primarily attributed to their association with Bioprojects analysing TB outbreaks or

originating from countries with extensive sampling, highlighting a bias in sequencing efforts

towards more prevalent or historically studied lineages. Many sub-Saharan African countries

do not have a representative sample in our dataset, and this may explain the under-

representation of L5 and L6 in the dataset. These lineages contribute significantly to the

overall burden of TB in this region (Coscolla et al., 2021). Additionally, we found that certain

RDs that were previously assumed to be lineage-specific were identified in other lineages

(Supplementary table S4). The observed variation in RD distribution emphasizes the need for

an expanded genomic dataset to provide a more comprehensive understanding of their

prevalence, specificity, and implications in the evolution of the MTBC. Future sequencing

efforts should target high-burden TB countries, ensuring a comprehensive representation of

the population structure to enhance our understanding of MTBC diversity and evolution.

Conclusion

In conclusion, this study reveals several significant findings. Firstly, the MTBC has a small,

closed pangenome, predominantly driven by genome reduction. This observation underlines

the importance of genomic deletions in the evolution of the MTBC. The study also highlights

distinct genomic features across MTBC, including sublineage-specific RDs, which may play a

crucial role in variations in virulence, metabolism, and antibiotic resistance. The analysis

further reveals that the accessory genome of MTBC is a product of gene deletions, which can

be classified into lineage-specific and independent deletions. Finally, the study underscores

the impact of non-coding regions and large structural variations on MTBC's genetic diversity,

offering a new perspective on MTBC pangenomics.

Acknowledgements and funding

We would like to thank all the staff at the BCCM/ITM collection for the growth of strains for

sequencing in this study. We thank Alasdair Hubbard for assistance with Nanopore

sequencing.

C.J.M. and M.B. are supported by the Academy of Medical Sciences (AMS), the Wellcome

Trust, the Government Department of Business, Energy and Industrial Strategy (BEIS), the

British Heart Foundation and Diabetes UK and the Global Challenges Research Fund (GCRF)

via a Springboard grant [SBF006\1090].

Author contributions

M.B. contributed to conceptualization, methodology, formal analysis, investigation, data

curation, writing -original draft, writing – reviewing & editing, and visualization.

M.M. contributed to methodology, formal analysis, resources, data curation and writing –

reviewing & editing.

M.F. contributed to methodology, data curation, writing – reviewing & editing and

supervision.

J.C.T contributed to methodology, resources, and writing – reviewing & editing.

M.R.D.S. contributed to methodology, validation, formal analysis, investigation, writing –

reviewing & editing and visualisation.

C.J.M. contributed to conceptualization, methodology, formal analysis, investigation, writing

-original draft, writing – reviewing & editing, supervision, project administration and funding

acquisition.

References

ALONSO, H., SAMPER, S., MARTÍN, C. & OTAL, I. 2013. Mapping IS6110 in high-copy number
Mycobacterium tuberculosis strains shows specific insertion points in the Beijing
genotype. *BMC Genomics,* 14**,** 422.
ANTOINE, R., GAUDIN, C. & HARTKOORN, R. C. 2021. Intragenic Distribution of IS6110 in
Clinical Mycobacterium tuberculosis Strains: Bioinformatic Evidence for Gene
Disruption Leading to Underdiagnosed Antibiotic Resistance. *Microbiol Spectr,* 9**,**
e0001921.
ARNVIG, K. B., CORTES, T. & YOUNG, D. B. 2014. Noncoding RNA in Mycobacteria.
Microbiology Spectrum, 2**,** 10.1128/microbiolspec.mgm2-0029-2013.
BAENA, A., CABARCAS, F., OCAMPO, J. C., BARRERA, L. F. & ALZATE, J. F. 2023. Large genomic
deletions delineate Mycobacterium tuberculosis L4 sublineages in South American
countries. *PLOS ONE,* 18, e0285417.
BESPIATYKH, D., BESPYATYKH, J., MOKROUSOV, I. & SHITIKOV, E. 2021. A Comprehensive
Map of Mycobacterium tuberculosis Complex Regions of Difference. *mSphere,* 6**,**
e0053521.
BLOUIN, Y., HAUCK, Y., SOLER, C., FABRE, M., VONG, R., DEHAN, C., CAZAJOUS, G.,
MASSOURE, P.-L., KRAEMER, P., JENKINS, A., GARNOTEL, E., POURCEL, C. &
VERGNAUD, G. 2012. Significance of the Identification in the Horn of Africa of an
Exceptionally Deep Branching Mycobacterium tuberculosis Clade. *PLOS ONE,* 7**,**
e52841.
BOTTAI, D., FRIGUI, W., SAYES, F., DI LUCA, M., SPADONI, D., PAWLIK, A., ZOPPO, M.,
ORGEUR, M., KHANNA, V., HARDY, D., MANGENOT, S., BARBE, V., MEDIGUE, C., MA,
L., BOUCHIER, C., TAVANTI, A., LARROUY-MAUMUS, G. & BROSCH, R. 2020. TbD1
deletion as a driver of the evolutionary success of modern epidemic *Mycobacterium*
tuberculosis lineages. *Nature Communications,* 11**,** 684.
BRITES, D., LOISEAU, C., MENARDO, F., BORRELL, S., BONIOTTI, M. B., WARREN, R.,
DIPPENAAR, A., PARSONS, S. D. C., BEISEL, C., BEHR, M. A., FYFE, J. A., COSCOLLA, M.
& GAGNEUX, S. 2018. A New Phylogenetic Framework for the Animal-Adapted
Mycobacterium tuberculosis Complex. *Front Microbiol,* 9**,** 2820.
BROCKHURST, M. A., HARRISON, E., HALL, J. P. J., RICHARDS, T., MCNALLY, A. & MACLEAN, C.
2019. The Ecology and Evolution of Pangenomes. *Curr Biol,* 29**,** R1094-r1103.
BROSCH, R., GORDON, S. V., MARMIESSE, M., BRODIN, P., BUCHRIESER, C., EIGLMEIER, K.,
GARNIER, T., GUTIERREZ, C., HEWINSON, G., KREMER, K., PARSONS, L. M., PYM, A. S.,
SAMPER, S., VAN SOOLINGEN, D. & COLE, S. T. 2002. A new evolutionary scenario for
the *Mycobacterium tuberculosis* complex. *Proceedings of the National Academy of*
Sciences, 99**,** 3684-3689.
CERES, K. M., STANHOPE, M. J. & GRÖHN, Y. T. 2022. A critical evaluation of Mycobacterium
bovis pangenomics, with reference to its utility in outbreak investigation. *Microb*
Genom, 8.
CHINER-OMS, Á., LÓPEZ, M. G., MORENO-MOLINA, M., FURIÓ, V. & COMAS, I. 2022. Gene
evolutionary trajectories in *Mycobacterium tuberculosis* reveal temporal signs of
selection. *Proceedings of the National Academy of Sciences,* 119**,** e2113600119.
COLE, S. T., BROSCH, R., PARKHILL, J., GARNIER, T., CHURCHER, C., HARRIS, D., GORDON, S. V.,
EIGLMEIER, K., GAS, S., BARRY, C. E., TEKAIA, F., BADCOCK, K., BASHAM, D., BROWN,

D., CHILLINGWORTH, T., CONNOR, R., DAVIES, R., DEVLIN, K., FELTWELL, T., GENTLES,
S., HAMLIN, N., HOLROYD, S., HORNSBY, T., JAGELS, K., KROGH, A., MCLEAN, J.,
MOULE, S., MURPHY, L., OLIVER, K., OSBORNE, J., QUAIL, M. A., RAJANDREAM, M. A.,
ROGERS, J., RUTTER, S., SEEGER, K., SKELTON, J., SQUARES, R., SQUARES, S.,
SULSTON, J. E., TAYLOR, K., WHITEHEAD, S. & BARRELL, B. G. 1998. Deciphering the
biology of Mycobacterium tuberculosis from the complete genome sequence.
Nature, 393, **537-544.**

COSCOLLA, M. 2017. Biological and Epidemiological Consequences of MTBC Diversity. *Adv*
Exp Med Biol, 1019**,** 95-116.

COSCOLLA, M., GAGNEUX, S., MENARDO, F., LOISEAU, C., RUIZ-RODRIGUEZ, P., BORRELL, S.,
OTCHERE, I. D., ASANTE-POKU, A., ASARE, P., SÁNCHEZ-BUSÓ, L., GEHRE, F.,
SANOUSSI, C. N., ANTONIO, M., AFFOLABI, D., FYFE, J., BECKERT, P., NIEMANN, S.,
ALABI, A. S., GROBUSCH, M. P., KOBBE, R., PARKHILL, J., BEISEL, C., FENNER, L.,
BÖTTGER, E. C., MEEHAN, C. J., HARRIS, S. R., DE JONG, B. C., YEBOAH-MANU, D. &
BRITES, D. 2021. Phylogenomics of Mycobacterium africanum reveals a new lineage
and a complex evolutionary history. *Microb Genom,* 7.

DARWIN, K. H. & STANLEY, S. A. 2022. The aldehyde hypothesis: metabolic intermediates as
antimicrobial effectors. *Open Biol,* 12**,** 220010.

DE JONG, B. C., ANTONIO, M. & GAGNEUX, S. 2010. Mycobacterium africanum—Review of
an Important Cause of Human Tuberculosis in West Africa. *PLOS Neglected Tropical*
Diseases, 4**,** e744.

DEB, C., LEE, C.-M., DUBEY, V. S., DANIEL, J., ABOMOELAK, B., SIRAKOVA, T. D., PAWAR, S.,
ROGERS, L. & KOLATTUKUDY, P. E. 2009. A Novel In Vitro Multiple-Stress Dormancy
Model for Mycobacterium tuberculosis Generates a Lipid-Loaded, Drug-Tolerant,
Dormant Pathogen. *PLOS ONE,* 4**,** e6077.

DEJESUS, M. A., GERRICK, E. R., XU, W., PARK, S. W., LONG, J. E., BOUTTE, C. C., RUBIN, E. J.,
SCHNAPPINGER, D., EHRT, S., FORTUNE, S. M., SASSETTI, C. M. & IOERGER, T. R. 2017.
Comprehensive Essentiality Analysis of the Mycobacterium tuberculosis Genome via
Saturating Transposon Mutagenesis. *mBio,* 8.

FLORES-VALDEZ, M. A., ACEVES-SÁNCHEZ, M. J., PETERSON, E. J. R., BALIGA, N., BRAVO-
MADRIGAL, J., DE LA CRUZ-VILLEGAS, M., ARES, M. A., BORN, S., VOSKUIL, M., PÉREZ-
PADILLA, N. A., BURCIAGA-FLORES, M., CAMACHO-VILLEGAS, T. A. & ESPINOZA-
JORGE, M. G. 2020. Transcriptional portrait of M. bovis BCG during biofilm
production shows genes differentially expressed during intercellular aggregation and
substrate attachment. *Sci Rep,* 10**,** 12578.

GAGNEUX, S. 2018. Ecology and evolution of Mycobacterium tuberculosis. *Nat Rev*
Microbiol, 16**,** 202-213.

GAGNEUX, S., DERIEMER, K., VAN, T., KATO-MAEDA, M., DE JONG, B. C., NARAYANAN, S.,
NICOL, M., NIEMANN, S., KREMER, K., GUTIERREZ, M. C., HILTY, M., HOPEWELL, P. C.
& SMALL, P. M. 2006. Variable host-pathogen compatibility in *Mycobacterium*
tuberculosis. *Proceedings of the National Academy of Sciences,* 103**,** 2869-2873.

GODFROID, M., DAGAN, T. & KUPCZOK, A. 2018. Recombination Signal in Mycobacterium
tuberculosis Stems from Reference-guided Assemblies and Alignment Artefacts.
Genome Biol Evol, 10**,** 1920-1926.

HERSHBERG, R., LIPATOV, M., SMALL, P. M., SHEFFER, H., NIEMANN, S., HOMOLKA, S.,
ROACH, J. C., KREMER, K., PETROV, D. A., FELDMAN, M. W. & GAGNEUX, S. 2008. High

Functional Diversity in Mycobacterium tuberculosis Driven by Genetic Drift and Human Demography. *PLOS Biology,* 6**,** e311.

HUERTA-CEPAS, J., FORSLUND, K., COELHO, L. P., SZKLARCZYK, D., JENSEN, L. J., VON MERING, C. & BORK, P. 2017. Fast Genome-Wide Functional Annotation through Orthology Assignment by eggNOG-Mapper. *Mol Biol Evol,* 34**,** 2115-2122.

HUNT, M., SILVA, N. D., OTTO, T. D., PARKHILL, J., KEANE, J. A. & HARRIS, S. R. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biology,* 16**,** 294.

JONES, P., BINNS, D., CHANG, H. Y., FRASER, M., LI, W., MCANULLA, C., MCWILLIAM, H., MASLEN, J., MITCHELL, A., NUKA, G., PESSEAT, S., QUINN, A. F., SANGRADOR-VEGAS, A., SCHEREMETJEW, M., YONG, S. Y., LOPEZ, R. & HUNTER, S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics,* 30**,** 1236-40.

KASSAMBARA, A. & MUNDT, F. Extract and Visualize the Results of Multivariate Data Analyses [R package factoextra version 1.0.7]. 2020.

KATOH, K., MISAWA, K., KUMA, K. & MIYATA, T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res,* 30**,** 3059-66.

KOLMOGOROV, M., YUAN, J., LIN, Y. & PEVZNER, P. A. 2019. Assembly of long, error-prone reads using repeat graphs. *Nature Biotechnology,* 37**,** 540-546.

KOZLOV, A. M., DARRIBA, D., FLOURI, T., MOREL, B. & STAMATAKIS, A. 2019. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics,* 35**,** 4453-4455.

MANNI, M., BERKELEY, M. R., SEPPEY, M., SIMÃO, F. A. & ZDOBNOV, E. M. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol Biol Evol,* 38**,** 4647-4654.

MARIN, M. 2024. Analysis of the limited *Mycobacterium tuberculosis* pangenome reveals potential pitfalls of pan-genome analysis approaches

MEEHAN, C. J., GOIG, G. A., KOHL, T. A., VERBOVEN, L., DIPPENAAR, A., EZEWUDO, M., FARHAT, M. R., GUTHRIE, J. L., LAUKENS, K., MIOTTO, P., OFORI-ANYINAM, B., DREYER, V., SUPPLY, P., SURESH, A., UTPATEL, C., VAN SOOLINGEN, D., ZHOU, Y., ASHTON, P. M., BRITES, D., CABIBBE, A. M., DE JONG, B. C., DE VOS, M., MENARDO, F., GAGNEUX, S., GAO, Q., HEUPINK, T. H., LIU, Q., LOISEAU, C., RIGOUTS, L., RODWELL, T. C., TAGLIANI, E., WALKER, T. M., WARREN, R. M., ZHAO, Y., ZIGNOL, M., SCHITO, M., GARDY, J., CIRILLO, D. M., NIEMANN, S., COMAS, I. & VAN RIE, A. 2019. Whole genome sequencing of Mycobacterium tuberculosis: current standards and open issues. *Nature Reviews Microbiology,* 17**,** 533-545.

MENARDO, F., RUTAIHWA, L. K., ZWYER, M., BORRELL, S., COMAS, I., CONCEIÇÃO, E. C., COSCOLLA, M., COX, H., JOLOBA, M., DOU, H. Y., FELDMANN, J., FENNER, L., FYFE, J., GAO, Q., GARCÍA DE VIEDMA, D., GARCIA-BASTEIRO, A. L., GYGLI, S. M., HELLA, J., HIZA, H., JUGHELI, L., KAMWELA, L., KATO-MAEDA, M., LIU, Q., LEY, S. D., LOISEAU, C., MAHASIRIMONGKOL, S., MALLA, B., PALITTAPONGARNPIM, P., RAKOTOSAMIMANANA, N., RASOLOFO, V., REINHARD, M., REITHER, K., SASAMALO, M., SILVA DUARTE, R., SOLA, C., SUFFYS, P., BATISTA LIMA, K. V., YEBOAH-MANU, D., BEISEL, C., BRITES, D. & GAGNEUX, S. 2021. Local adaptation in populations of Mycobacterium tuberculosis endemic to the Indian Ocean Rim. *F1000Res,* 10**,** 60.

NAPIER, G., CAMPINO, S., MERID, Y., ABEBE, M., WOLDEAMANUEL, Y., ASEFFA, A., HIBBERD, M. L., PHELAN, J. & CLARK, T. G. 2020. Robust barcoding and identification of Mycobacterium tuberculosis lineages for epidemiological and clinical studies. *Genome Medicine,* 12**,** 114.

NGABONZIZA, J. C. S., LOISEAU, C., MARCEAU, M., JOUET, A., MENARDO, F., TZFADIA, O., ANTOINE, R., NIYIGENA, E. B., MULDERS, W., FISSETTE, K., DIELS, M., GAUDIN, C., DUTHOY, S., SSENGOOBA, W., ANDRÉ, E., KASWA, M. K., HABIMANA, Y. M., BRITES, D., AFFOLABI, D., MAZARATI, J. B., DE JONG, B. C., RIGOUTS, L., GAGNEUX, S., MEEHAN, C. J. & SUPPLY, P. 2020. A sister lineage of the Mycobacterium tuberculosis complex discovered in the African Great Lakes region. *Nature Communications,* 11**,** 2917.

NOLL, N., MOLARI, M., SHAW, L. P. & NEHER, R. A. 2023. PanGraph: scalable bacterial pan-genome graph construction. *Microbial Genomics,* 9.

PHELAN, J. E., O'SULLIVAN, D. M., MACHADO, D., RAMOS, J., OPPONG, Y. E. A., CAMPINO, S., O'GRADY, J., MCNERNEY, R., HIBBERD, M. L., VIVEIROS, M., HUGGETT, J. F. & CLARK, T. G. 2019. Integrating informatics tools and portable sequencing technology for rapid detection of resistance to anti-tuberculous drugs. *Genome Medicine,* 11**,** 41.

QIN, L., WANG, J., LU, J., YANG, H., ZHENG, R., LIU, Z., HUANG, X., FENG, Y., HU, Z. & GE, B. 2019. A deletion in the RD105 region confers resistance to multiple drugs in Mycobacterium tuberculosis. *BMC Biol,* 17**,** 7.

R CORE TEAM 2023. R: A language and environment for statistical computing. R Foundation for Statistical Computing. *R Core Team.* Vienna, Austria.

RICE, P., LONGDEN, I. & BLEASBY, A. 2000. EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet,* 16**,** 276-7.

SANOUSSI, C. N., APOS, DIRA, COSCOLLA, M., OFORI-ANYINAM, B., OTCHERE, I. D., ANTONIO, M., NIEMANN, S., PARKHILL, J., HARRIS, S., YEBOAH-MANU, D., GAGNEUX, S., RIGOUTS, L., AFFOLABI, D., DE JONG, B. C. & MEEHAN, C. J. 2021. Mycobacterium tuberculosis complex lineage 5 exhibits high levels of within-lineage genomic diversity and differing gene content compared to the type strain H37Rv. *Microbial Genomics,* 7.

SANOUSSI, C. N. D., AFFOLABI, D., RIGOUTS, L., ANAGONOU, S. & DE JONG, B. 2017. Genotypic characterization directly applied to sputum improves the detection of Mycobacterium africanum West African 1, under-represented in positive cultures. *PLOS Neglected Tropical Diseases,* 11**,** e0005900.

SILVA-PEREIRA, T. T., SOLER-CAMARGO, N. C. & GUIMARÃES, A. M. S. 2024. Diversification of gene content in the Mycobacterium tuberculosis complex is determined by phylogenetic and ecological signatures. *Microbiology Spectrum,* 0**,** e02289-23.

SZKLARCZYK, D., KIRSCH, R., KOUTROULI, M., NASTOU, K., MEHRYARY, F., HACHILIF, R., GABLE, A. L., FANG, T., DONCHEVA, N. T., PYYSALO, S., BORK, P., JENSEN, L. J. & VON MERING, C. 2023. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Res,* 51**,** D638-d646.

TATUSOVA, T., DICUCCIO, M., BADRETDIN, A., CHETVERNIN, V., NAWROCKI, E. P., ZASLAVSKY, L., LOMSADZE, A., PRUITT, K. D., BORODOVSKY, M. & OSTELL, J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res,* 44**,** 6614-24.

TONKIN-HILL, G., GLADSTONE, R. A., PÖNTINEN, A. K., ARREDONDO-ALONSO, S., BENTLEY, S.
D. & CORANDER, J. 2023. Robust analysis of prokaryotic pangenome gene gain and
loss rates with Panstripe. *Genome Res,* 33**,** 129-140.
TONKIN-HILL, G., MACALASDAIR, N., RUIS, C., WEIMANN, A., HORESH, G., LEES, J. A.,
GLADSTONE, R. A., LO, S., BEAUDOIN, C., FLOTO, R. A., FROST, S. D. W., CORANDER, J.,
BENTLEY, S. D. & PARKHILL, J. 2020. Producing polished prokaryotic pangenomes with
the Panaroo pipeline. *Genome Biology,* 21**,** 180.
VASER, R., SOVIĆ, I., NAGARAJAN, N. & ŠIKIĆ, M. 2017. Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Res,* 27**,** 737-746.
WANG, L. G., LAM, T. T., XU, S., DAI, Z., ZHOU, L., FENG, T., GUO, P., DUNN, C. W., JONES, B.
R., BRADLEY, T., ZHU, H., GUAN, Y., JIANG, Y. & YU, G. 2020. Treeio: An R Package for
Phylogenetic Tree Input and Output with Richly Annotated and Associated Data. *Mol
Biol Evol,* 37**,** 599-603.
WHELAN, F. J., RUSILOWICZ, M. & MCINERNEY, J. O. 2020. Coinfinder: detecting significant
associations and dissociations in pangenomes. *Microb Genom,* 6.
WHO. 2023. *Global tuberculosis report 2023* [Online]. World Health Organization. Available:
https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-
tuberculosis-report-2023 [Accessed].
ZHANG, Z., SCHWARTZ, S., WAGNER, L. & MILLER, W. 2000. A greedy algorithm for aligning
DNA sequences. *J Comput Biol,* 7**,** 203-14.
ZULKOWER, V. & ROSSER, S. 2020. DNA Features Viewer: a sequence annotation formatting
and plotting library for Python. *Bioinformatics,* 36**,** 4350-4352.
ZWYER, M., ÇAVUSOGLU, C., GHIELMETTI, G., PACCIARINI, M. L., SCALTRITI, E., VAN
SOOLINGEN, D., DÖTSCH, A., REINHARD, M., GAGNEUX, S. & BRITES, D. 2021. A new
nomenclature for the livestock-associated Mycobacterium tuberculosis complex
based on phylogenomics. *Open Res Eur,* 1**,** 100.

Figures

Figure 1: The geography of the dataset. (A) sample collection distribution by country; (B) the

number of genomes of each lineage included from each continent.

Figure 2: Boxplot showing the distribution of accessory genes within each lineage. Lineages with a

small number of genomes were excluded from the analysis. L6, La1 and *M. microti* have significantly

smaller accessory genomes compared to other lineages.

Figure 3: Analysis of the functional components in the core (A) and accessory (B) genome using EggNOG mapper and InterProScan.

648   **Figure 4: (A) phylogenetic tree based on MTBC core genome. PCA based on the accessory genome**

649   **data from Panaroo (B) and pangraph (C).**

Figure 5: MTBC Accessory genome identified by Panaroo (A) and Pangraph (B).

Figure 6: Sub-lineage specific regions of differences (RDs) identified using pangenome-based and genome alignment-based approaches. Sub-lineages are shown on the Y-axis coloured as per the legend and clustered based on the RD presence/absence patterns. RDs are listed on the X-axis, grouped by their pattern of presence/absence.

659　**Figure 7: Visual presentation of deletion caused by RD702.** The truncated *bglS* gene was classified as

660　a core gene with Panaroo but has specific deletions within, which can be problematic for

661　pangenomic analyses.

662