The evolutionary history of the ancient weevil family Belidae (Coleoptera: Curculionoidea) reveals the marks of Gondwana breakup and major floristic turnovers, including the rise of angiosperms

Xuankun Li[1,2,3*], Adriana E. Marvaldi[4,5*], Rolf G. Oberprieler[6], Dave Clarke[2,3], Brian D. Farrell[7], Andrea Sequeira[8], M. Silvia Ferrer[5], Charles O'Brien[9†], Shayla Salzman[10], Seunggwan Shin[11], William Tang[12], Duane D. McKenna[2,3]

[1] Department of Entomology, College of Plant Protection, China Agricultural University, Beijing 100193, People's Republic of China ORCID: 0000-0002-0622-2064 (XL)

[2] Department of Biological Sciences and [3] Center for Biodiversity Research, University of Memphis, Memphis, TN 38152, USA ORCID: 0000-0002-7823-8727 (DDM)

[4] CONICET, División Entomología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, 1900 La Plata, Buenos Aires, Argentina ORCID: 0000-0001-6029-4803 (AEM)[5] CONICET, Instituto Argentino de Investigaciones de Zonas Áridas, Entomología, CC 507, 5500 Mendoza, Argentina (AEM, MSF)

[6] CSIRO, Australian National Insect Collection, GPO Box 1700, Canberra, Australian Capital Territory 2601, Australia ORCID: 0000-0002-1837-580X (RGO)

[7] Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA. ORCID: 0000-0002-6843-0539 (BDF)

[8] Department of Biological Sciences, Wellesley College, Wellesley MA, 02481, USA ORCID: 0000-0001-6771-2991 (AS)

[9] 2313 West Calle Balaustre, Green Valley, Arizona 85622, USA (CO)

[10] Department of Entomology, University of Georgia, Athens, GA 30602 USA (ShSa)

[11] School of Biological Sciences, Seoul National University, Seoul 08826, Republic of Korea. ORCID: 0000-0002-4258-8661 (SeSh)

[12] Florida State Collection of Arthropods, Florida Department of Agriculture – DPI, P.O. Box 147100, Gainesville, FL 32614, USA. ORCID: 0000-0001-9009-3113 (WT)

* Correspondence authors: xuankun.li@cau.edu.cn (XL); marvaldi@fcnym.unlp.edu.ar (AEM).

†Deceased

Abstract

The rise of angiosperms to ecological dominance and the breakup of Gondwana during the Mesozoic marked major transitions in the evolutionary history of insect-plant interactions. To elucidate how contemporary trophic interactions were influenced by host plant shifts and palaeogeographical events, we integrated molecular data with information from the fossil record to construct a timetree for ancient phytophagous weevils of the beetle family Belidae. Our analyses indicate that crown-group Belidae originated approximately 138 Ma ago in Gondwana, associated with Pinopsida (conifer) host plants, with larvae likely developing in dead/decaying branches. Belids tracked their host plants as major plate movements occurred during Gondwana's breakup, surviving on distant, disjunct landmasses. Some belids shifted to Angiospermae and Cycadopsida when and where conifers declined, evolving new trophic interactions, including brood-pollination mutualisms with cycads and associations with achlorophyllous parasitic angiosperms. Extant radiations of belids in the genera *Rhinotia* (Australian region) and *Proterhinus* (Hawaiian Islands) have relatively recent temporal origins.

Keywords: angiosperms, beetles, biogeography, gymnosperms, host shifts, phytophagy, Pinopsida

Introduction

The evolutionary interplay between plant-feeding (phytophagous) insects and vascular plants has fundamentally shaped terrestrial biodiversity for over 400 million years (e.g., Ehrlich & Raven 1964; Farrell et al. 1992; Wilf 2008; Misof et al. 2014; Swain et al. 2022). The fossil record has provided insights into the historical context and evolutionary processes that have shaped modern insect-plant interactions (Labandeira & Currano 2013). However, (1) the degree to which contemporary trophic interactions reflect those in the past, e.g., representing relatively ancient primary associations versus more recent secondary associations, and (2) the relative roles of vicariance (including major palaeogeographical events) and dispersal in the evolution of host shifts, remain unclear.

Gymnosperms (cycads, conifers and relatives), the dominant plant group during the Mesozoic Era, thrived from the Permian to the Cretaceous period, dominating most terrestrial ecosystems and forming intimate associations with many groups of phytophagous insects (Anderson et al., 2007). The transition to the Cretaceous marked the rise of angiosperms (flowering plants), which gradually replaced gymnosperms in many ecological settings to become the dominant plant group (Anderson et al., 2007). This transition was not only a botanical shift but also marks a critical period in the evolutionary history of insect-plant interactions. Angiosperms introduced novel ecological niches and resources, leading to the diversification of phytophagous insects and a reconfiguration of insect-plant associations (e.g., Farrell 1998; Wilf 2008; McKenna et al. 2009, 2015; Swain et al. 2022). By the end of the Mesozoic, the

continents had rifted into nearly their present forms, though their positions would continue to change.

Gondwana split into South America, Africa, Australia, Antarctica and the Indian subcontinent, while

Laurasia became North America and Eurasia.

On account of their extraordinary taxonomic diversity and varied trophic interactions with

contemporary gymnosperms and flowering plants, beetles (order Coleoptera; > 400000 described extant

species) have been widely used as models to study the evolution of diversity at the insect-plant interface

(e.g., Farrell 1998; McKenna et al. 2009, 2019). The present paper addresses a significant gap in our

understanding of phylogeography and the evolution of host plant associations in the family Belidae (belid

weevils), an ancient group of phytophagous beetles whose extant species exhibit specialized trophic

associations with gymnosperms and flowering plants. By reconstructing the evolutionary history of belid

weevils and examining the dynamics of their host plant associations and geographical distributions over

time, this paper seeks to yield new insights into the evolution of modern insect-plant interactions.

Belidae, with approximately 360 described extant species in 40 genera and two subfamilies

(Marvaldi & Ferrer, 2014; O'Brien & Tang, 2015), comprise an early branch of Curculionoidea

(Oberprieler et al., 2007; McKenna et al., 2009, 2018; McKenna, 2011) (Fig. 1). Belidae are indicated to

have originated during the Jurassic in association with gymnosperms and subsequently colonized

angiosperms, as the latter diversified and rose to ecological dominance during the Late Cretaceous and

Paleogene (e.g., Farrell 1998; McKenna et al., 2009; Shin et al., 2018). They exhibit their highest generic

diversity in the Southern Hemisphere and collectively are associated with a variety of plants, including

conifers, cycads and a few families of angiosperms, including Arecaceae, Balanophoraceae, Hydnoraceae,

Celastraceae, Myrtaceae and Vitaceae (Marvaldi & Ferrer, 2014). The adults appear to feed mainly on

stem tissues but sometimes also on pollen, some being important pollinators, and the larvae develop in the

bark and woody tissues of decaying branches or twigs, gymnosperm strobili, flower buds or fruits. Like

Australian cycad-pollinating weevils (Curculionidae: Molytinae) (Hsiao et al., 2023), Neotropical Belidae

in the tribe Allocorynina develop as brood pollination mutualists in the pollen cones (or 'male strobili') of

cycads (e.g., Tang 1987; Marvaldi & Ferrer, 2014; Salzman et al., 2020), a biotic interaction that has

multiple independent origins in weevils (Curculionidae: Curculioninae) (e.g., Haran et al., 2023).

The historical biogeography and evolution of host associations of Belidae have long drawn the

attention of scientists (Kuschel, 1959; Zimmerman, 1994; Anderson, 2005; Marvaldi et al., 2006).

However, our understanding of belid evolution remains limited due to the lack of a dated phylogeny.

Since the landmark study of weevil family-level phylogeny by Kuschel (1995), a series of studies based

on morphological characters of adults and larvae have inferred the phylogenetic relationships of Belidae

and revised their tribal-level classification (Kuschel & Leschen, 2003; Anderson, 2005; Marvaldi, 2005;

Marvaldi et al., 2006). Phylogenetic hypotheses based on analyses of morphological data have been

proposed for all genera of both Belinae (Kuschel & Leschen, 2003) and Oxycoryninae (Marvaldi et al.,

2006; Anderson & Marvaldi, 2013). However, beyond including small numbers of exemplar taxa in

higher-level studies of weevils (e.g., McKenna et al., 2009; Shin et al., 2018), the phylogeny and

evolution of Belidae have not been explored using molecular data.

We conducted molecular phylogenetic analyses and divergence dating of belid weevils to

investigate their relationships and evolution. We also reconstructed ancestral states of host plant

associations and undertook a biogeographical analysis to explore geographical patterns of diversification

and the evolution of host plant organ and taxon associations. We integrated phylogenomic and Sanger

data for Belidae, sampling all seven tribes and 60 % of the extant genera[1]. We performed ancestral-state

reconstruction using the resulting chronogram with stochastic character mapping and event-based

likelihood ancestral-area estimation. We sought to answer two main questions about the distribution and

host plant associations of crown group Belidae: (1) is the development of larvae of Agnesiotidini and

Pachyurini (Belinae) and Oxycraspedina (Oxycoryninae) in conifer hosts an ancient, primary association

(Farrell 1998) or the result of more recent, secondary colonization, and (2) how did the interplay between

biogeographical process and host plant shifts influence and shape the trophic associations of belids?

Results

The 33t423g dataset comprised 97334 nucleotides, with 37154 parsimony-informative sites and 31.8 %

missing data. The 33t424g and 46t424g datasets contained 107199 nucleotides each, with 45046 and

45157 parsimony-informative sites and 33.3 % and 51.6 % missing data, respectively.

Backbone topologies generated from the combined dataset (46t424g) were consistent with those

from the reduced datasets (Figs 1, S1). Flanking regions were included in subsequent analyses because

this increased the resolution and backbone nodal support values for the reduced datasets. The family

Belidae, the subfamilies Belinae and Oxycoryninae, the tribes of Belinae (Agnesiotidini, Belini and

Pachyurini) and most subtribes of Oxycoryninae (Oxycraspedina, Metrioxenina, Aglycyderina and

Oxycorynina) were robustly supported as monophyletic (UFBoot ≥ 95 AND SH-aLRT ≥ 80) in the trees

resulting from the three partitioning schemes (Fig. 1). Tribal-level relationships were robustly recovered

in Belinae, but generic relationships were only weakly supported even excluding the taxa without AHE

data (33t423g and 33t424g) (Fig. S1). The tribe Oxycorynini, in its current concept (Marvaldi et al., 2006),

was not recovered as a monophylum, Oxycraspedina instead forming the sister-group of Aglycyderini +

Metrioxenini, and *Archicorynus* was found to be the sister-group of Allocorynina rather than of all other

---

[1] Combining Sanger sequences data with genomic data for phylogenetic inference has been demonstrated as a feasible approach to resolving deep-level relationships while adding taxa to the phylogeny for tracing the evolutionary history of characters (e.g. Zhang et al., 2016 Song et al., 2020; Li et al., 2022a, 2022b).

Oxycoryninae (Anderson & Marvaldi, 2013). The phylogenetic positions of Metrioxenini and

Archicorynini were unstable across different partitioning schemes (Fig. 1). This is most likely due to

insufficient data, as only Sanger data were generated for representatives of these two tribes. The

phylogeny estimate generated with the partition-by-locus scheme was selected as optimal because it

closely matches hypotheses based on the analysis of morphological characters (Marvaldi et al., 2006).

Therefore, the ML tree generated by the 46t424t dataset and partitioned by locus was used for

downstream analyses. Details of node supports are summarized in Fig. S2 and Tab. S2.

BEAST analyses with different tree priors/clock schemes yielded similar results, with the crown

age of Belidae ranging from 160.0 to 125.6 Ma (Tab. 1). The preferred BEAST analysis, applying a birth-

death tree prior with 13 unlinked molecular clocks, was identified using marginal-likelihood estimation

(Tab. 1). Divergence-time estimation results revealed an origin of stem Belidae in the Middle Jurassic at

167.8 Ma (95 % highest posterior density = 185.9–160.0 Ma) and of crown-group Belidae in the early

Lower Cretaceous at 138.5 Ma (154.9–125.6 Ma) (Fig. 2). The highest likelihood among six

biogeographical models tested was DIVALIKE with the unconstrained analysis (M0) (Tab. S3). Belidae

were reconstructed as having a Gondwanan origin, with the two extant subfamilies Belinae and

Oxycoryninae originating in the Australian and Neotropical regions, respectively (Fig. 2).

The most likely ancestral host plants of Belidae were reconstructed as Pinopsida (0.53) or

Angiospermae (0.43) (Fig. S2, Tab. S2). In Belinae, Pinopsida were the most likely ancestral host plants

(0.79), with a shift to Angiospermae occurring in Belini (0.65) and two independent colonizations of

Polypodiopsida (if indeed these are larval hosts and the two genera are not as closely related as indicated

by morphology; see *Discussion* below). The ancestral host plants of Oxycoryninae were recovered as

Angiospermae (0.76) or Pinopsida (0.21). Extant Oxycraspedina are today associated with Pinopsida and

Allocorynina with Cycadopsida (Fig. 2). Based on the ASR analysis, branches of the host plants were

used for larval development by ancestral Belidae with high possibility (0.87), with a single shift to

flowers and fruits (of Angiospermae) in Oxycorynina and at least three shifts to strobili, two to strobili of

Pinopsida in Pachyurini and Oxycraspedina and one to (pollen) cones of Cycadopsida in Allocorynina

(Fig. 2).

Discussion

Phylogenetic relationships

This study provides the first target-enrichment-based phylogeny estimate of the weevil family Belidae.

The monophyly of both subfamilies, Belinae and Oxycoryninae, was recovered, and Aglycyderini were

found deeply nested in the latter, supporting the now widely accepted two-subfamily system of belid

weevil classification (e.g., Marvaldi 2005). In Belinae, all three subtribes (Agnesiotidini, Belini and

Pachyurini) were recovered as monophyletic, supporting the classification of Kuschel & Leschen (2003).

Generic relationships in Belini were generally poorly resolved, especially when taxa with Sanger data

only were included (Figs 1, S1). The placements of *Trichophthalmus* and *Homalocerus* are likely to be

artifacts because they result from sparse and non-overlapping data (four Sanger and 13 AHE loci,

respectively). In contrast, morphological data suggest that these two genera are closely related (Kuschel,

1959; Vanin, 1976; Kuschel & Leschen, 2003) and together constitute the sister group of all Australian

Belini (Fig. 195 of Kuschel & Leschen, 2003). In Oxycoryninae, the tribe Oxycorynini in its current

concept was found to be polyphyletic, with Oxycraspedina being more closely related to Aglycyderini +

Metrioxenini (Fig. 1). A close relationship between Aglycyderini and Metrioxenini was also found in the

phylogeny estimate derived from morphological characters (Marvaldi et al. 2006). All other tribes and the

subtribes were recovered as monophyletic groups, in agreement with the results of phylogenies

reconstructed from morphological characters by previous authors (Marvaldi et al., 2006; Anderson &

Marvaldi, 2013). However, the placement of *Archicorynus* as the sister group of Allocorynina (albeit with

weak support, Fig. 1) differs from that indicated by morphological characters, which resolves the genus as

the sister group of all other Oxycoryninae (Anderson & Marvaldi, 2013).

Ancestral host plants and larval feeding habits

The larvae of ancestral Belidae were reconstructed as having developed endophytically in branches (not

strobili) of their host plants, which is consistent with some previous hypotheses (Marvaldi et al. 2002;

Marvaldi 2005). Even though our reconstruction provides support for their ancestral hosts most likely

being the Pinopsida, which is in accordance with our clade age estimates, angiosperm hosts cannot be

completely ruled out, considering that crown angiosperms are now thought to have originated earlier than

Belidae, in the interval between the late Permian and the latest Jurassic (256–149 Ma) (Barba-Montoya et

al., 2018). However, angiosperm families known to be belid hosts, such as Arecaceae, Hydnoraceae,

Balanophoraceae and Fabaceae, did not diversify until the Cretaceous (Barba-Montoya et al., 2018; Li et

al., 2019). During the early evolutionary history of Belidae, conifer lineages were dominant (Crisp &

Cook, 2011; Anderson et al. 2007; Wang & Ran, 2014), consistent with Pinopsida as the most likely

ancestral host plants of Belidae.

The ancestral host plants of Belinae were recovered as being Pinopsida. Based on our phylogeny

estimate, the host plant shift to Angiospermae may have occurred during the branching of Belini in the

Middle to Late Paleogene. Possible host shifts to ferns in the Paleogene in the Neotropical genera

Trichophthalmus and *Homalocerus* must be viewed with great reservation, as all host records of these

genera are only based on the collection of adults, without any evidence of adult feeding, oviposition or

larval development (Kuschel, 1959; Vanin, 1976). As mentioned above, *Homalocerus* and

Trichophthalmus might form the earliest branching lineage of Belini. Therefore, the more plausible

scenario for host plant shifts is a single shift to ferns from conifers during the Early to Middle Paleogene,

perhaps tracking the radiation of ferns in the late Cretaceous (Schneider et al., 2004; Schuettpelz & Pryer,

2009).

In Oxycoryninae, the South American genus *Oxycraspedus* (the only extant genus of

Oxycraspedina) retains the reconstructed ancestral host association with Pinopsida. Its current hosts,

Araucariaceae, were widely distributed in Mesozoic times, and when the oxycorynines diverged by the

Lower Cretaceous according to our dated phylogeny estimate, *Araucaria* still constituted an abundant and

widely distributed host for beetles (Sequeira et al, 2000). It is plausible, then, that the ancestors of the

remaining oxycorynine tribes/subtribes were originally associated with these conifers before they

declined by the Eocene (Kershaw & Wagstaff, 2001). The other oxycorynine lineages evidently adapted

to this decrease in conifer availability by host-shifting to various distantly related taxa of Angiospermae

and Cycadopsida.

As is true for weevils generally, the association of Metrioxenini with palms (Arecaceae) may date

to the Upper Cretaceous, given our dating and the fossil record of the family (Marvaldi et al, 2002;

Matsunaga & Smith, 2021). However, extant Metrioxenina may only be associated with the genus *Arenga*

(Marvaldi et al., 2006) — though possible associations with other members of the tribe Caryoteae and the

subfamily Coryphoideae have not been explored. The Aglycyderini evidently shifted onto angiosperms as

well, but their pattern of host associations is complex and indeterminate as the larvae of most species

develop in dead bark and twig tissues of a variety of plants. The larvae of Oxycorynina develop in the

flowers and fruits of achlorophyllous root-parasitic plants belonging to two distantly related angiosperm

families (Balanophoraceae and Hydnoraceae) (Ferrer et al. 2011). The sequence of their host shifts cannot

be reconstructed unequivocally, the possibilities include a parallel colonization of these taxa or a shift to

Balanophoraceae first and then to Hydnoraceae or vice versa. In any case, our results suggest that these

host shifts occurred in the Paleogene at *c.* 47.7 Ma (68.5–27.5 Ma), consistent with the estimated origins

of Balanophoraceae and Hydnoraceae (crown age of Hydnoraceae at 54.7 Ma (75–37 Ma) (Naumann et

al., 2013)).

All extant species of Allocorynina are known to develop in pollen cones of cycads of the genera

Dioon and *Zamia* (Marvaldi et al., 2006; O'Brien &Tang, 2015). Our analysis recovered an initial shift to

Dioon (hosts of *Parallocorynus* and *Protocorynus*) followed by a subsequent one from *Dioon* to *Zamia*

(hosts of *Notorhopalotria* and *Rhopalotria*) (Fig. 2). The shift to *Dioon* is indicated to have occurred

between the late Cretaceous and the early Paleogene, at 58.4 Ma (67.1–49.9 Ma), which is older than the

crown age of *Dioon* (24.6–7.5 Ma) but younger than the stem age (207.9–107.0 Ma) (Condamine et al.,

2015). Australian cycad weevils have also been found to have colonized cycads before their main

radiation (Hsiao et al., 2023). The cycad tree of life is known for long branches subtending generic-level

radiations (Nagalingum et al., 2011; Salas-Leiva et al., 2013; Condamine et al., 2015; Liu et al., 2022).

The switch from *Dioon* to *Zamia* is estimated to have occurred in the middle Neogene, at 11.0 Ma (13.5–

8.6 Ma), which is in line with the crown group age of *Zamia* (9.5 Ma, 22.1–9.0 Ma) (Calonje et al., 2019).

235   There appears to be a pronounced conservatism in the type of tissue consumed by belid larvae,

always involving parenchymatous parts of branches or reproductive structures (not pollen or seeds). The

type of plant organ used for larval development shows few shifts from the ancestral state (branches): in

Belinae to ovulate and/or pollen cones (or 'female and/or male strobili') of Pinopsida (arguably a shift;

possibly just opportunistic development alongside that in twigs and branches) in some Agnesiotidini and

Pachyurini, in Oxycoryninae to ovulate cones of Pinopsida in *Oxycraspedus*, fleshy flowers/fruits of

parasitic angiosperms in Oxycorynina and pollen cones of cycads in Allocorynina (there does not appear

to be any feeding on ovulate cones by Allocorynina in nature although they clearly visit them and are

presumably capable of feeding on them at least in the short term; Simon et al., 2023). As is generally true

for plant-feeding beetles (Farrell & Sequeira, 2004), host taxon associations are evidently more labile than

the use of tissue/organ for larval development. This is particularly noticeable even within some belid

genera, whose larvae have been found developing in similar tissues (under bark and in branches) of

different plant families, both between congeneric species (e.g., *Sphinctobelus*) and in single species (e.g.,

Isacantha, *Rhicnobelus*); however, there are some cases known in Belidae where the use of plant organs

is apparently plastic and opportunistic (e.g. *Apagobelus brevirostris* (Lea) reared from both stems and

cones of *Araucaria*; Zimmerman (1994)).

Palaeogeographical events and host plant shifts

A Gondwanan origin was recovered for Belidae at *c.* 138 Ma, when South America and Australia were

connected via Antarctica. The two subfamilies—Belinae and Oxycoryninae—originated and diverged

during the separation of East and West Gondwana (McIntyre et al., 2017) and both have lineages that still

develop on their ancestral host plant. In the East Gondwana clade, Belinae, the divergence time between

Agnesiotis and *Dicordylus* was estimated at 44.7 Ma (55.9–34.5 Ma), during the Paleocene–Eocene

separation of South America and Australia (Lawver et al., 1992; Briggs, 1995; McIntyre et al., 2017).

Thus, the evolutionary history of *Dicordylus* might best be explained by persisting in association with the

ancestral host plant despite vicariance. The history of the other two South American genera, *Homalocerus*

and *Trichophthalmus*, could potentially also be elucidated through the vicariance process (though we

cannot rule out the possibility of dispersal to South America). This process may have offered an

opportunity for their common ancestor to encounter the new niches of South American ferns, thereby

facilitating their shift to new host plants. Such palaeogeographical events shaping host plant usage have

also been documented in other phytophagous arthropods (e.g., Calatayud et al., 2016).

Due to the lack of Afrocorynina in our sample and the weakly supported placements of

Archicorynini and Metrioxenina, the evolutionary history of Oxycoryninae could not be reconstructed

with high confidence. Nonetheless, a plausible scenario is that the common ancestor of Aglycyderini +

Metrioxenini shifted to angiosperms in the Lower to Middle Cretaceous at 99.9 Ma (113.9–88.9 Ma)

(McIntyre et al., 2017), diverged in the Ethiopian region by the Middle to Upper Cretaceous, and

subsequently dispersed to the Palearctic and Oriental regions, the tropical Pacific islands and as far south

as New Zealand (Aglycyderina). Additional taxon sampling is needed to illuminate whether this host shift

promoted the dispersal of Aglycyderini + Metrioxenini.

Adaptive radiations in Rhinotia *and* Proterhinus

In Belidae, notable taxonomic diversity is observed in *Rhinotia* and *Proterhinus*, with 87 and 168

described species, respectively (Marvaldi & Ferrer, 2014; Brown, 2019). In contrast, other genera of

Belidae comprise fewer than 15 known species (Marvaldi & Ferrer, 2014). Host plants of *Rhinotia* are

mostly *Acacia* (Fabaceae), the most species-rich plant genus in Australia, with more than 1000 endemic

species (Maslin, 2004; Lewis, 2005). *Acacia* is particularly dominant in arid and semi-arid areas of

Australia (Byrne et al., 2008). The crown age of *Rhinotia* is here estimated at 11.7 Ma (14.1–9.2 Ma),

which postdates the origin of crown *Acacia* (23.9–21 Ma) (Miller et al., 2013). The common ancestor of

Rhinotia might have colonized *Acacia* in the early to middle Miocene when Australia was warm and wet

(Martin, 2006), and co-diversified during the aridification of Australia from 10–6 Ma (Byrne et al., 2008).

The origin might also have been 20 Ma earlier and involved other now extinct taxa, with *Rhinotia* being a

surviving lineage, or it might have colonized and radiated during the aridification. A similar situation has

been postulated for the thrips subfamily Phlaeothripinae, which colonized *Acacia* and diversified into

more than 250 species in 35 genera (McLeish et al., 2007, 2013).

With 159 species described from the Hawaiian Islands, *Proterhinus* is another example of the

spectacular radiation of Hawaiian insects, along with, e.g., the nearly 1000 species of Drosophilidae

(Diptera) (O'Grady et al., 2011), more than 400 species of *Hyposmocoma* (Lepidoptera: Cosmopterigidae)

(Haines et al., 2014), over 110 species of *Plagithmysus* (Coleoptera: Cerambycidae) (Gressitt, 1975) and

more than 190 species of *Nesophrosyne* (Hemiptera: Cicadellidae) (Bennett & O'Grady, 2013). Little is

known about the host plants of *Proterhinus*, but what is known suggests that host plant ranges (e.g., in *P.*

deceptor Perkins, *P. obscurus* Sharp and *P. vestitus* Sharp) are remarkably broader than in other species

of Belidae (Legalov, 2009). Such host "jumps" are typical for endemic Hawaiian phytophagous insects,

e.g., *Carposina* (Lepidoptera: Carposinidae) (Medeiros et al., 2016) and *Nesosydne* (Hemiptera:

Delphacidae) (Goodman et al., 2012), but some lineages have high host plant specificity e.g., *Philodoria*

(Lepidoptera, Gracillariidae) (Johns et al., 2018) and *Nesophrosyne* (Hemiptera: Cicadellidae) (Bennett &

O'Grady, 2013). Due to limited taxon sampling, we could not estimate the crown age of *Proterhinus* nor

when and from where it may have arrived in Hawaii.

Materials and methods

Taxon sampling

Forty-six taxa were included in the present study, including eight outgroups representing the

chrysomeloid families Chrysomelidae and Orsodacnidae and the weevil families Anthribidae,

Cimberididae and Nemonychidae. Ingroup taxon sampling spanned 38 species of Belidae in 24 genera,

representing the seven tribes from both subfamilies and 60 % of extant belid genera. Genomic data from

33 taxa, 23 newly generated for this study, were used in phylogeny reconstruction. Sanger DNA sequence

data (CO1, 16S, 18S and 28S) from six outgroup and 34 ingroup species were also used, including newly

generated data for 31 species, and 13 species were represented by Sanger data only (Table S1).

DNA extraction, library preparation and Illumina DNA sequencing

Total genomic DNA was extracted from the legs, thoracic muscle or the whole body, depending on the

size of the specimen, using the G-Biosciences OmniPrep kit (G-Biosciences, Catalog #786-136, St. Louis,

MO, U.S.A.), following the manufacturer's protocol, except that samples were incubated for 15 hours

instead of 15 minutes. Final DNA extractions were eluted with 60 uL of nuclease-free water and treated

with RNaseA. The remaining body parts were preserved in 95 % ethanol as vouchers. Genomic DNA QC

statistics were generated for each extracted specimen using a Qubit fluorometer, and DNA quality

(fragmentation/degradation and contamination with RNA) was further assessed via gel electrophoresis.

The extracted DNA was fragmented by sonication with a Q800R2 Sonicator (Illumina TruSeq),

using 50 uL of the DNA extractions in 0.2 mL strip tubes, targeting a modal fragment size of 350 base

pairs. Genomic DNA libraries were constructed using the NEBNext Ultra II DNA Library Prep Kit (NEB

#E7645L) with NEBNext Multiplex Oligos for Illumina (Dual Index Primers Sets 1 and 2) (NEB

#E7600S and #E7780S), with two-sided size selection around a mode of 480 base pairs. Target

enrichment through hybridization followed the myBaits Hybridization Capture for Targeted NGS

(Version 5), with 65 °C chosen for the hybridization temperature. We used the published AHE Coleoptera

Probe set (Haddad et al. 2018; Shin et al. 2018) and targeted 599 nuclear loci.

Enriched libraries were amplified using KAPA HiFi HotStart ReadyMix. PCR cycling consisted

of an initial denaturing step at 98 °C for 2 min, followed by eight cycles of denaturing at 98 °C for 20 s,

annealing at 60 °C for 30 s, elongation at 72 °C for 45 s and a final elongation step at 72 °C for 5 min.

The 192 enriched and multiplexed libraries were sequenced using 150-bp paired-end reads on an Illumina

HiSeq Lane at Novogene Corporation Inc. (Sacramento, CA, USA).

DNA isolation, PCR amplification and Sanger DNA sequencing

DNA extraction and PCR amplification of Sanger data were performed at IADIZA-CONICET (Mendoza,

Argentina) and Wellesley College (MA, USA). Total genomic DNA was extracted from adult voucher

specimens using an adapted "salting-out" protocol (Sunnuks & Hales 1996) or the DNeasy Blood and

Tissue Kit (QIAGEN, MD, USA). Tissue was processed from one to two legs or part of the thorax. The

extracted DNA was stored at -20 °C. Four molecular markers (two nuclear and two mitochondrial) were

used in this study: 18S rDNA (entire), 28S rDNA (regions D2, D3), 16S rDNA (regions IV, V) and COI

("barcode" or 5' region). The primers used for amplification and sequencing of the four Sanger loci and

PCR conditions are as described by Marvaldi et al. (2018). The PCR products were purified and bi-

directionally sequenced with the Sanger method, using the Sequencing Service of "Unidad de Genómica

de INTA-Castelar" (Buenos Aires, Argentina) and, in Wellesley, using an ABI PRISM 3100 Genetic

Analyzer (Applied Biosystems, Foster City, CA, USA). Electropherograms were edited and contig-

assembled using ProSeq v.2.91 (Filatov 2002) and sometimes Sequencher v.5 (GeneCodes Corp.). All

sequences were deposited in GenBank under accession numbers provided in Table S1.

Sequence assembly and orthology prediction

The dataset preparation procedure for AHE-targeted loci used is as outlined by Breinholt et al. (2018). A

reference set was prepared using genomic coding sequences (CDS) from nine coleopteran genomes:

Anoplophora glabripennis (Motschulsky) (Cerambycidae, GCA_000390285.2), *Aethina tumida* Murray

(Nitidulidae, GCA_001937115.1), *Callosobruchus maculatus* (Fabricius) (Chrysomelidae,

GCA_900659725.1), *Dendroctonus ponderosae* (Hopkins) (Curculionidae, GCA_020466585.1),

Diabrotica virgifera LeConte (Chrysomelidae, GCA_003013835.2), *Gonioctena quinquepunctata*

(Fabricius) (Chrysomelidae, GCA_018342105.1), *Leptinotarsa decemlineata* Say (Chrysomelidae,

GCA_000500325.2), *Sitophilus oryzae* (Linnaeus) (Curculionidae, GCA_002938485.2) and *Tribolium*

castaneum (Herbst) (Tenebrionidae, GCA_000002335.3). Raw reads were assembled using an iterative

baited assembly (IBA) after filtering with Trim Galore! v.0.4.0 (bioinformatics.babraham.ac.uk).

Orthology was determined using the *T. castaneum* genome as a reference, and single-hit and genome

mapping location criteria were used with NCBI Blastn (Camacho et al., 2009). Cross-contamination

checks were conducted with USEARCH (Edgar, 2010), and sequences with > 99 % identity across

different subfamilies were identified and removed. Cleaned sequences were aligned in MAFFT v.7.245

(Katoh & Standley, 2013), and isoform consensuses were generated using FASconCAT-G 1.02 (Kück &
Longo, 2014).
Following the method outlined by Teasdale et al. (2016) and Li et al. (2022a), we used a blast-
based method to extract Sanger genes from genomic sequences. In short, raw sequence data were
assembled using SOAPdenovo v.2 (Li et al., 2015), the four Sanger genes were identified using an all-by-
all tBlastx search, reads were mapped on potential orthologous sequences using BBmap v.35.85
(Bushnell, 2014), and a final consensus sequence was generated after variants were called using GATK
v.4.1.1.0 (McKenna et al., 2010). Extracted sequences were compared with Sanger sequencing results,
and sequences with higher quality and longer reads were maintained for subsequent phylogenetic analysis.
The sequences of the ribosomal markers (nuclear 18S and 28S and mitochondrial 16S) were aligned using
information on the secondary structure of the arthropod rRNA genes to identify homologous positions as
well as regions of ambiguous alignment to be excluded from analyses (Gillespie et al., 2006; Marvaldi et
al. 2009, 2018).
378
Dataset preparation
AHE probe sets comprise highly conserved coding probe regions (i.e., exons) and more variable,
generally, non-coding flanking regions (e.g., introns or intergenic regions) located on flanks of the probe
region (Lemmon et al., 2012; Haddad et al., 2018; Shin et al. 2018). AliView v1.18 (Larsson, 2014) was
used to manually check each nucleotide alignment to separate "flanks" from "probe regions" and ensure
that the probe region was in the correct open reading frame (ORF). We used a long-branch detection
protocol to investigate the possibility of external contamination, paralogous sequences and significant
sequencing/assembly errors (longbranchpruner.pl available on Osiris, http://galaxy-
dev.cnsi.ucsb.edu/osiris/). We produced maximum-likelihood (ML) gene trees from nucleotide (NT)-
probe-region multiple-sequence alignments (MSAs) in IQ-TREE v.2.0.6 (Nguyen et al., 2015),
conducting a full model test for each gene. We pruned tip sequences that exceeded eight standard
deviations from the mean tip length of the gene tree from NT MSAs. Loci with < 40 % taxon coverage
were excluded.
Five hundred eighty loci were assembled across 35 taxa, and 419 loci were selected for
phylogenetic inference. Four Sanger genes were included, and flanking regions were concatenated and
treated as a single locus. Cleaned MSAs were concatenated using Phyx v.1.1 (Brown et al., 2017) to
generate the dataset with 46 taxa and 424 loci for phylogenetic inference (46t424g). The 13 taxa
represented only by Sanger data were excluded to evaluate the impact of missing data and flanking
regions, generating two datasets, 33t424g with 33 taxa and 424 loci and 33t423g with 33 taxa and 423
loci (flanking regions excluded).

Phylogenetic analyses and tests of node support

We conducted ML phylogenetic analyses in IQ-TREE v2.1.3 (Nguyen et al., 2015). For the 33t423g and 33t424g datasets, three partitioning schemes were used: (1) unpartitioned, (2) partitioned by locus, and (3) partitioned by locus and codon position. All schemes were model-tested using ModelFinder (Kalyaanamoorthy et al. 2017) as implemented in IQ-TREE. The best partitioning scheme for the 33t423g dataset was found after merging possible partitions (using the '-MFP+MERGE' command) and determining the best scheme under the Bayesian information criterion (BIC). For the 33t424g and 46t424g datasets, the best schemes of the 33t423g dataset were used with the GTR+ASC model added for the flanking regions. An initial 1000 parsimony trees were generated in IQ-TREE with the command '-ninit 1000', and 100 trees with the fewest steps were used to initialize the candidate set (-ntop 100), considering all possible nearest-neighbor interchanges (-allnni). These 100 trees were maintained in the candidate set during the ML tree search (-nbest 100), and unsuccessful runs were terminated after 1000 iterations (-nstop 1000). Perturbation strength was set to 0.2 (-pers 0.2), as recommended for datasets with many short sequences. We used nearest-neighbor interchange (NNI) branch swapping to improve the tree search and limit overestimating branch supports due to severe model violations ('-bnni' command). Node supports were computed with 1000 UFBoot replicates ('-B' command) (Minh et al. 2013; Hoang et al. 2018) and SH-aLRT ('-alrt' command) (Guindon et al. 2010).

Both concatenation and gene coalescence approaches were used for tree estimation on dataset 46t424g. For concatenated analyses, partitioning schemes and parameters used were the same as those for the 33t424g and 33t423g datasets. Nodes were classified as "robustly supported" when they were recovered with support values of UFBoot ≥ 95 AND SH-aLRT ≥ 80, as "moderately supported" when UFBoot ≥ 95 OR SH-aLRT ≥ 80 and as "weakly supported" when UFBoot < 95 AND SH-aLRT < 80 (Minh et al. 2013; Hoang et al. 2018).

The best evolutionary model for coalescence analyses was found using ModelFinder Plus ('-MFP' command) for each gene, followed by likelihood tree searches using a partitioning scheme generated in IQ-TREE. We applied nearest-neighbor interchange (NNI) branch swapping to improve the tree search and limit overestimating branch supports due to severe model violations ('-bnni' command). Nodal support was computed with 1000 UFBoot replicates ('-B' command) (Minh et al. 2013; Hoang et al. 2018). Multi-species-coalescent (MSC) analyses based on these single trees were calculated using ASTRAL-III v5.6.2 (Zhang et al., 2018). Nodes were classified as "robustly supported" when recovered with support values ≥ 0.95.

Four-cluster likelihood mapping (FcLM; Strimmer & von Haeseler, 1997) was performed in IQ-TREE using the 33t423g and 33t424g datasets and ModelFinder-determined partitions (partition by locus

433   and codon position) to quantify support for the sister relationship between Allocorynina and

434   Oxycraspedina. The four schemes used to define the four-taxon clusters are Allocorynina, Aglycyderini,

435   Oxycraspedina and outgroups.

436   Quartet sampling (Pease et al., 2018) provided an alternate examination of branch support.

437   Quartet sampling of internal node scores included a set of three scores: quartet concordance (QC: values

438   near 1 being more concordant and near -1 more discordant), quartet differential (QD: the more equal the

439   frequencies of discordant topologies, the closer to 1; 0 indicating that only one other discordant topology

440   was found) and quartet informativeness (QI: 1 for all replicates informative, 0 for no replicates

441   informative) (Pease et al., 2018).

442   Concordance and disagreement among genes on the selected ML tree generated by the 46t424g

dataset were estimated using the gene concordance factor (gCF) implemented in IQ-TREE (Minh et al.,

2020), as concatenated analyses can return well-supported trees even when the level of gene incongruence

is high (e.g., Jeffroy et al., 2006; Kumar et al., 2012). Nodes were classified as "robustly supported" when

gCF was higher than gene discordance factor gDF1 and gDF2, "weakly supported" when gCF equaled

gDF1 or gDF2 or both and "not supported" when gCF was lower than gDF1 or gDF2 (more loci

supporting an alternative topology).

Divergence time estimation

Divergence times were estimated in a Bayesian framework using BEAST v1.10.4 (Suchard et al. 2018) on

the high-performance computing clusters at the University of Memphis and the China Agricultural

University. SortaDate (Smith et al. 2018) was used to reduce the nucleotide alignment to a

computationally tractable matrix (50 loci) using combined results of clock-likeness (get_var_length.py)

and least topological conflict with the species tree (get_bp_genetrees.py). Four Sanger genes were

included in the BEAST analyses to ensure that all taxa were present in the reduced matrix, regardless of

the SortaDate result. Three different initial partitioning strategies were used: 1) unpartitioned; 2) four

partitions (three codon positions for protein-coding genes and one partition for non-protein coding genes);

3) 144 partitions by locus and nucleotide codon position. The best-fit model and partitioning schemes

were selected by PartitionFinder2 (Lanfear et al. 2017) using the greedy algorithm and the BIC scores

across all models included in BEAST (models = beast). The reduced concatenated data matrix was

imported into BEAUTi (Drummond et al., 2012). Substitution and clock models were unlinked among

partitions, and tree models were linked. An uncorrelated relaxed-molecular-clock model (Drummond et al.

2006) and a lognormal prior were applied, two tree priors were tested for each partitioning scheme, Yule

(pure birth) and birth–death, and a fixed cladogram based on the topology generated by the concatenated

ML analysis was used.

Four internal nodes and the root were constrained for calibration based on the fossil record of

Belidae. Several supposed belid fossils have been described, but only a few could be confidently placed

based on evident synapomorphies. After a careful examination following Parham et al. (2012), five fossils

were selected for calibration of four internal nodes: 1) *Sinoeuglypheus daohugouensis* Yu et al., 2019

(together with other undescribed Daohugou specimens) for the stem of Cimberididae, with a minimum

age of 160 Ma based on the age of the boundary between the Oxfordian and Callovian (ICC 2023); 2)

Talbragarus averyi Oberprieler & Oberprieler, 2012 for the stem of Rhinorhynchinae, with a minimum

age of 147.3 Ma based on analysis of zircons (Bean 2006); 3) *Preclarusbelus vanini* Santos et al., 2007

and *Cratonemonyx martinsnetoi* Legalov, 2014 for the stem of Oxycoryninae, with a minimum age of 113

Ma based on the Crato Formation as Upper Aptian following Santos et al. (2011); 4) *Pleurambus*

strongylus Poinar & Legalov, 2014 for the stem of Allocorynini, with a minimum age of 15 Ma

(Iturralde-Vinent & MacPhee 1996, 2019). The lower margin of the estimated timing of origin of

Phytophaga (195 Ma) (McKenna et al. 2019) was used for the maximum age of the stem of Cimberididae

and Rhinorhynchinae.

Three independent analyses of each clock scheme and tree-prior combination were run to check

for convergence. We evaluated the convergence and mixing of MCMC chains in Tracer version 1.6

(Rambaut et al. 2018) to ensure that the effective sample sizes (ESS) exceeded 200. The resulting tree

files were combined and resampled with a frequency of 100000 in LogCombiner (BEAST package) and a

burn-in of 30 %. Subsampled trees were summarized as a maximum-clade-credibility tree using

TreeAnnotator (Rambaut and Drummond 2015), with median heights as node heights. Path sampling and

stepping-stone sampling (Xie et al. 2011; Baele et al. 2012, 2013) were performed as part of all BEAST

analyses to identify the best tree prior and clock scheme combination.

Ancestral host plant reconstructions

Ancestral reconstruction analyses were performed for two host plant characters on the dated phylogeny: (i)

host plant higher taxa (four states): Angiospermae, Cycadopsida, Pinopsida and Polypodiopsida; (ii) host

plant organ (three states): branches, strobili and flower and fruit (Table S1). Stochastic character mapping

was conducted with the "make.simmap" command in the R package phytools (Revell, 2012) with 1000

simulations. Character state transitions were assumed to have equal rates ("ER" option).

Ancestral area estimation

We recognize five bioregions that best account for the distribution of the sampled species in Belidae:

Neotropical region (N), Australian region (A), tropical Pacific islands (I), Oriental region (O) and

Palearctic + Ethiopian regions (P) (Table S1). We performed an event-based likelihood-ancestral-area

estimation using BioGeoBEARS (Matzke, 2014). Three models were used: (1) DEC (Dispersal

Extinction Cladogenesis (Ree & Smith, 2008)), (2) DIVALIKE (a likelihood-based implementation of

dispersal vicariance analysis, originally parsimony-based (Ronquist, 1997)), (3) BAYAREALIKE (a

likelihood implementation of BayArea, originally Bayesian (Landis et al., 2013)). All models were also

evaluated under a constrained analysis (M1), in which we considered palaeogeographical events that

occurred in the past 160 Ma over four-time slices (160–125 Ma, 125–35 Ma, 35–28 Ma, 28 Ma to present;

sliced by three events: separation of East and West Gondwana, separation of South America and Australia

(McIntyre et al., 2017), origin of Hawaii islands (Price & Clague, 2007)) and geographical distance

variation for a total of six scenarios. The maximum allowed ancestral area was restricted to two. Time

slices for geographical events used were those of McIntyre et al. (2017). The Akaike Information

Criterion (AIC (Burnham & Anderson, 1998)) and the corrected Akaike Information Criterion (AICc

(Burnham & Anderson, 2002)) were calculated to select the best-fitting model.

Acknowledgments

This study was funded by grants from the United States National Science Foundation DEB:1355169 and

2110053 to DDM. XL was supported by the 2115 Talent Development Program of China Agricultural

University. AEM acknowledges support by CONICET through grant PIP 3108. The authors acknowledge

the China Agricultural University and the University of Memphis for providing high-performance

computing platforms and support that contributed to the research results reported in this study.

References

Anderson, R.S. (2005). **New Oxycoryninae from Central and South America: phylogenetic and**

biogeographical implications (Coleoptera: Belidae) *Systematic Entomology* **30**:644–652

https://doi:10.1111/j.1365-3113.2005.00304.x

Anderson, J.M., Anderson H.M., Cleal C.J (2007) *Brief history of the gymnosperms: classification,*

biodiversity, phytogeography and ecology South African National Biodiversity Institute. South

Africa, Pretoria, 2007 pp.x + 280 pp

Anderson, R.S., Marvaldi, A.E (2013). **Finding unexpected beetles in odd places:** *Archicorynus*

kuscheli **Anderson and Marvaldi, a new genus and species representing a new tribe,**

Archicorynini, of Oxycoryninae (Coleoptera: Belidae) from Nicaragua *The Coleopterists*

Bulletin 61–71

Baele G., Lemey P., Bedford T., Rambaut A., Suchard M.A., Alekseyenko, A.V (2012) **Improving the**

accuracy of demographic and molecular clock model comparison while accommodating

phylogenetic uncertainty *Molecular Biology and Evolution* **29**:2157–2167

https://doi.org/10.1093/molbev/mss084

Baele G., Li W.L.S., Drummond A.J., Suchard M.A., Lemey P (2013) **Accurate model selection of**

relaxed molecular clocks in Bayesian phylogenetics *Molecular Biology and Evolution* **30**:239–

243 https://doi.org/10.1093/molbev/mss243

Barba-Montoya, J., Dos Reis, M., Schneider, H., Donoghue, P. C., Yang, Z (2018) **Constraining**

uncertainty in the timescale of angiosperm evolution and the veracity of a Cretaceous

Terrestrial Revolution *New Phytologist* **218**:819–834

Bean, L.B. (2006) **The leptolepid fish *Cavenderichthys talbragarensis* (Woodward, 1895) from the**

Talbragar Fish Bed (Late Jurassic) near Gulgong, New South Wales *Records-Western*

Australian Museum **23**:43–76

Bennett, G.M., O'Grady, P.M (2013) **Historical biogeography and ecological opportunity in the**

adaptive radiation of native Hawaiian leafhoppers (Cicadellidae: Nesophrosyne) *Journal of*

Biogeography **40**:1512–1523

Breinholt, J.W., Earl, C., Lemmon, A.R., Lemmon, E.M., Xiao, L., Kawahara, A.Y (2018) **Resolving**

relationships among the megadiverse butterflies and moths with a novel pipeline for

anchored phylogenomics *Systematic Biology* 67:78–93 https://doi:10.1093/sysbio/syx048

Briggs, J.C. (1995) ***Global biogeography*** Elsevier Science, Amsterdam. XVII + 452 pp

Brown, J.W., Walker, J.F., Smith, S.A (2017) **Phyx: Phylogenetic tools for unix** *Bioinformatics*

33:1886–1888 https://doi:10.1093/bioinformatics/ btx063

Brown, S.D. (2019) **A new species of *Proterhinus* Sharp, 1878 (Coleoptera: Curculionoidea: Belidae)**

from Miti'aro, Cook Islands, South Pacific *Zootaxa* **4664**:481–496

Byrne, M., Yeates, D.K., Joseph, L., Kearney, M., Bowler, J., Williams, M.A.J., Cooper, S., Donnellan,

S.C., Keogh, J.S., Leys, R., Melville, J., Murphy, D.J., Porch, N., Wyrwoll, K.-H (2008) **Birth of a**

biome: insights into the assembly and maintenance of the Australian arid zone biota *Molecular*

Ecology **17**:4398–4417

Burnham, K.P., Anderson, D.R (1998) **Practical use of the information-theoretic approach** In *Model*

selection and inference (eds KP Burnham, DR Anderson), pp. 75–117. New York, NY: Springer

Burnham, K.P., Anderson, D.R (2002) ***Model selection and multimodel inference: a practical***

information-theoretic approach, 2nd edn New York, NY: Springer. XXVI + 488 pp

Bushnell, B. (2014) **BBMap: A Fast, Accurate, Splice-aware Aligner (No. LBNL-7065E)** Lawrence

Berkeley National Lab. (LBNL), Berkeley, CA

Calatayud, J., Hórreo, J.L., Madrigal-Gonzalez, J., Migeon, A., Rodriguez, M.A., Magalhaes, S., Hortal, J
(2016) **Geography and major host evolutionary transitions shape the resource use of plant**
parasites *Proceedings of the National Academy of Sciences* **113**:9840–9845

Calonje, M., Meerow, A.W., Griffith, M.P., Salas-Leiva, D., Vovides, A.P., Coiro, M., Francisco-Ortega,
J (2019) **A time-calibrated species tree phylogeny of the New World cycad genus *Zamia* L.**
(Zamiaceae, Cycadales) *International Journal of Plant Sciences* **180**:286–314

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L (2009)
BLAST+: architecture and applications *BMC Bioinformatics* **10**:421 https://doi:10.1186/1471-
2105-10-421

Condamine, F. L., Nagalingum, N. S., Marshall, C. R., Morlon, H (2015) **Origin and diversification of**
living cycads: a cautionary tale on the impact of the branching process prior in Bayesian
molecular dating *BMC Evolutionary Biology* **15**:1–18

Crisp, M. D., Cook, L.G (2011) **Cenozoic extinctions account for the low diversity of extant**
gymnosperms compared with angiosperms *New Phytologist* **192**:997–1009

Drummond, A.J., Ho, S.Y.W., Phillips, M.J., Rambaut, A (2006) **Relaxed phylogenetics and dating**
with confidence *PLoS Biology* **4**:699–710 https://doi.org/10.1371/journal.pbio.0040088

Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A (2012) **Bayesian phylogenetics with BEAUti and**
the BEAST 1.7 *Molecular Biology and Evolution* **29**:1969–1973
https://doi.org/10.1093/molbev/mss075

Edgar, R.C. (2010) **Search and clustering orders of magnitude faster than BLAST** *Bioinformatics*
26:2460–2461 https://doi:10.1093/bioinformatics/btq461

Ehrlich, P.R., Raven, P.H (1964) **Butterflies and plants: a study in coevolution** *Evolution* 586–608.

Farrell, B.D. (1998) **"Inordinate Fondness" explained: why are there So many beetles?** *Science*
281:555–559

Farrell, B.D., Mitter, C., Futuyma, D.J (1992) **Diversification at the insect-plant Interface** *Bioscience*
42:34–42

Farrell, B.D., Sequeira, A.S (2004) **Evolutionary rates in the adaptive radiation of beetles on plants**
Evolution **58**:1984–2001

Ferrer, M.S., Marvaldi, A.E., Sato, H.A., Gonzalez, A.M (2011) **Biological notes on two species of**
***Oxycorynus* (Coleoptera: Belidae) associated with parasitic plants of the genus**
***Lophophytum* (Balanophoraceae), and new distribution records in Argentina**. *Revista de la*
Sociedad Entomológica Argentina **70**:351–355

Filatov, D.A. (2002) **ProSeq: a software for preparation and evolutionary analysis of DNA sequence**
data sets *Molecular Ecology Notes* **2**:621–624

Gillespie, J.J., Johnston, J.S., Cannone, J.J., Gutell, R.R (2006) **Characteristics of the nuclear (18S, 5.8**
601   **S, 28S and 5S) and mitochondrial (12S and 16S) rRNA genes of *Apis mellifera* (Insecta:**
602   **Hymenoptera): structure, organization, and retrotransposable elements** *Insect Molecular*
603   *Biology* **15**:657–686

604   Goodman, K.R., Welter, S.C., Roderick, G.K (2012) **Genetic divergence is decoupled from ecological**
605   **diversification in the Hawaiian *Nesosydne* planthoppers** *Evolution* **66**:2798– 2814

606   Gressitt, J.L. (1975) **Evolution of the endemic Hawaiian cerambycid-beetles** *International Biological*
607   *Program, Technical Report No. 71. U. S.* 1–46

608   Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O (2010) **New algorithms**
609   **and methods to estimate maximum-likelihood phylogenies: assessing the performance of**
610   **PhyML 3.0.** *Systematic Biology* **59**:307–321 https://doi:10.1093/sysbio/ syq010

611   Haddad, S., Shin, S., Lemmon, A.R., Lemmon, E.M., Svacha, P., Farrell, B., Ślipiński, A.D., Windsor, D.,
612   McKenna, D.D (2018) **Anchored hybrid enrichment provides new insights into the**
613   **phylogeny and evolution of longhorned beetles (Cerambycidae)** *Systematic Entomology*
614   **43**:68–89 https://doi: 10.1111/syen.12257

615   Haines, W.P., Schmitz, P., Rubinoff, D (2014) **Ancient diversification of *Hyposmocoma* moths in**
616   **Hawaii** *Nature Communications* **5**:p.3502

617   Haran, J., Li, X., Allio, R., Shin, S., Benoit, L., Oberprieler, R.G., Farrell, B.D., Brown, S.D.J., Leschen,
618   R.A., Kergoat, G.J., McKenna, D.D (2023) **Phylogenomics illuminates the phylogeny of flower**
619   **weevils (Curculioninae) and reveals ten independent origins of brood-site pollination**
620   **mutualism in true weevils** *Proceedings of the Royal Society B*, **290**:p.20230889

621   Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S. (2018) **UFBoot2: improving the**
622   **ultrafast bootstrap approximation** *Molecular Biology and Evolution* **35**:518–522
623   https://doi:10.1093/molbev/msx281

624   Hsiao, Y., Oberprieler, R.G., Zwick, A., Zhou, Y.L. and Ślipiński, A. (2023) **Museomics unveil**
625   **systematics, diversity and evolution of Australian cycad-pollinating weevils** *Proceedings of*
626   *the Royal Society B* **290**:p.20231385

627   Iturralde-Vinent, M.A., MacPhee, R.D.E (1996) **Age and paleogeographical origin of Dominican**
628   **amber** *Science* **273**:1850–1852

629   Iturralde-Vinent, M.A., MacPhee, R.D.E (2019) **Remarks on the age of Dominican amber**
630   *Palaeoentomology* **2**:236–240

631   Jeffroy, O., Brinkmann, H., Delsuc, F., Philippe, H (2006) **Phylogenomics: the beginning of**
632   **incongruence?** *Trends in Genetics* **22**:225–231 https://doi:10.1016/j.tig.2006.02.003

Johns, C.A., Toussaint, E.F., Breinholt, J.W. and Kawahara, A.Y (2018) **Origin and macroevolution of**

micro-moths on sunken Hawaiian Islands *Proceedings of the Royal Society B* **285**:p.20181047

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K., von Haeseler, A., Jermiin, L.S (2017) **ModelFinder: fast**

model selection for accurate phylogenetic estimates *Nature Methods* **14**:587–589

https://doi.org/10.1038/nmeth.4285

Katoh, K., Standley, D.M (2013) **MAFFT multiple sequence alignment software version 7:**

improvements in performance and usability *Molecular Biology and Evolution* **30**:772–780

https://doi:10.1093/molbev/mst010

Kershaw, P., Wagstaff, B (2001) **The southern conifer family Araucariaceae: history, status, and**

value for paleoenvironmental reconstruction *Annual Review of Ecology and Systematics* **32**:397–

414

Kumar, S., Filipski, A.J., Battistuzzi, F.U., Pond, S.L.K., Tamura, K (2012) **Statistics and truth in**

phylogenomics *Molecular Biology and Evolution* **29**:457–472 https://doi:10.1093/molbev/msr202

Kuschel, G. (1959) **Nemonychidae, Belidae y Oxycorynidae de la fauna chilena, con algunas**

consideraciónes biogeográficas (Coleoptera Curculionoidea, aporte 28) *Investigaciones*

zoológicas chilenas **5**:229–271

Kuschel, G. (1995) **A phylogenetic classification of Curculionoidea to families and subfamilies**

Memoirs of the Entomological Society of Washington **14**:5–33

Kuschel, G., Leschen, R.A.B. (2003) *Appendix 1. Phylogenetic relationships of the genera of Belinae.*

In 'Nemonychidae, Belidae, Brentidae (Insecta: Coleoptera: Curculionoidea)'. (Author: G.

Kuschel.) pp. 48–55. Fauna of New Zealand, Volume 45 (Manaaki Whenua Press: Lincoln, New

Zealand.)

Kück, P., Longo, G.C (2014) **FASconCAT-G: extensive functions for multiple sequence alignment**

preparations concerning phylogenetic studies *Frontiers in Zoology* **11**:81

https://doi.org/10.1186/s12983-014-0081-x

Labandeira, C.C., Currano, E.D (2013) **The fossil record of plant-insect dynamics** *Annual Review of*

Earth and Planetary Sciences **41**:287–311

Landis, M.J., Matzke, N.J., Moore, B.R., Huelsenbeck, J.P (2013) **Bayesian analysis of biogeography**

when the number of areas is large *Systematic Biology* **62**:789–804

https://doi:10.1093/sysbio/syt040

Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T., Calcott, B. (2017) **PartitionFinder 2: new**

methods for selecting partitioned models of evolution for molecular and morphological

phylogenetic analyses *Molecular Biology and Evolution* **34**, 772–773

Larsson A. (2014) **AliView: A fast and lightweight alignment viewer and editor for large datasets**

Bioinformatics **30**:3276–3278 https://doi:10.1093/ bioinformatics/btu531

Lawver, L.A., Gahagan, L.M., Coffin, M.F (1992) **The development of paleoseaways around**
Antarctica *Anarctic Research Series vol. 56*:7–30

Legalov, A.A. (2009) **Contribution to the knowledge of the Mesozoic Curculionoidea (Coleoptera)**
Amurian Zoological Journal **1**:283–295

Legalov, A.A. (2014) Taxonomic names, in the Mesozoic stage of evolution of the family Nemonychidae
(Coleoptera, Curculionoidea). *Paleontological Journal* **48**:851–944

Lemmon, A.R., Emme, S.A., Lemmon, E.M (2012) **Anchored hybrid enrichment for massively high-**
throughput phylogenomics *Systematic Biology* **61**:727–744 https://doi:10.1093/sysbio/sys049

Lewis, G.P. (2005) *Tribe Acacieae.* pp. 187–192 In Lewis, G.P., Schrire, B., MacKinder, B. & Lock, M.
(eds.), Legumes of the world. Kew: Royal Botanic Gardens.

Li, D., Liu, C.M., Luo, R., Sadakane, K. and Lam, T.W (2015) **MEGAHIT: an ultra-fast single-node**
solution for large and complex metagenomics assembly via succinct de Bruijn graph
Bioinformatics **31**:1674–1676

Li, H.T., Yi, T.S., Gao, L.M., Ma, P.F., Zhang, T., Yang, J.B., Gitzendanner, M.A., Fritsch, P.W., Cai, J.,
Luo, Y., Wang, H (2019) **Origin of angiosperms and the puzzle of the Jurassic gap** *Nature*
Plants **5**:461–470

Li, X., Hamilton, C.A., St Laurent, R., Ballesteros-Mejia, L., Markee, A., Haxaire, J., Rougerie, R.,
Kitching, I.J., Kawahara, A.Y (2022b) **A diversification relay race from Caribbean-**
Mesoamerica to the Andes: historical biogeography of *Xylophanes* **hawkmoths** *Proceedings of*
the Royal Society B **289**:20212435 https://doi:10.1098/rspb.2021.2435

Li, X., St Laurent, R., Earl, C., Doorenweerd, C., van Nieukerken, E.J., Davis, D.R., Johns, C.A.,
Kawakita, A., Kobayashi, S., Zwick, A., Lopez-Vaamonde, C (2022a) **Phylogeny of gracillariid**
leaf-mining moths: evolution of larval behaviour inferred from phylogenomic and Sanger data
Cladistics **38**:277–300

Liu, Y., Wang, S., Li, L., Yang, T., Dong, S., Wei, T., Wu, S., Liu, Y., Gong, Y., Feng, X., Ma, J (2022)
The Cycas genome and the early evolution of seed plants *Nature Plants* **8**:389–401

Martin, H.A. (2006) **Cenozoic climatic changes and the development of the arid zone vegetation of**
Australia *Journal of Arid Environments* **66**:533–563

Marvaldi, A.E. (2005) **Larval morphology and biology of oxycorynine weevils and the higher**
phylogeny of Belidae (Coleoptera, Curculionoidea) *Zoologica Scripta* **34**:37–48

Marvaldi, A.E., Del Río, M.G., Pereyra, V.A., Rocamundi, N., Lanteri, A.A. (2018) **A combined**
molecular and morphological approach to explore the higher phylogeny of entimine weevils
(Coleoptera: Curculionidae), with special reference to South American taxa *Diversity* **10**:95
https://doi.org/10.3390/d10030095

Marvaldi, A.E., Duckett, C.N., Kjer, K.M., Gillespie, J.J (2009) **Structural alignment of 18S and 28S**
rDNA sequences provides insights into phylogeny of Phytophaga (Coleoptera: Curculionoidea
and Chrysomeloidea) *Zoologica Scripta* **38**:63–77

Marvaldi, A.E., Ferrer, M.S (2014) *Belidae Schoenherr, 1826*. Handbook of Zoology: Arthropoda.
Insecta.

Marvaldi, A.E., Oberprieler, R.G., Lyal, C.H.C., Bradbury, T., Anderson, R.S (2006) Phylogeny of the
Oxycoryninae *sensu lato* (Coleoptera: Belidae) and evolution of host-plant associations.
Invertebrate Systematics 20:447–476

Marvaldi, A.E., Sequeira, A.S., O'Brien, C.W., Farrell, B.D (2002) **Molecular and morphological**
phylogenetics of weevils (Coleoptera, Curculionoidea): do niche shifts accompany
diversification? *Systematic Biology* **51**:761–785

Matsunaga, K.K., Smith, S.Y (2021) **Fossil palm reading: using fruits to reveal the deep roots of palm**
diversity *American Journal of Botany* 108:472–494

Maslin, B.R. (2004) *Classification and phylogeny of Acacia. In Evolution of ecological and*
behavioural diversity: Australian Acacia thrips as model organisms. Edited by Crespi BJ, Morris
DC, Mound LA. Canberra: Australian Biological Resources Study and Australian National Insect
Collection, CSIRO 97–112

Matzke, N.J. (2014) **Model selection in historical biogeography reveals that founder-event speciation**
is a crucial process in island clades *Systematic Biology* **63**:951–970
https://doi:10.1093/sysbio/syu056

McIntyre, S.R., Lineweaver, C.H., Groves, C.P., Chopra, A (2017) **Global biogeography since Pangaea**
Proceedings of the Royal Society B: Biological Sciences **284**:20170716

McKenna, D.D. (2011) **Temporal lags and overlap in the diversification of weevils and flowering**
plants: Recent advances and prospects for additional resolution *American Entomologist* 57:54–
55

McKenna, D.D., Clarke, D.J., Anderson, R., Astrin, J.J., Brown, S., Chamorro, L., Davis, S.R., De
Medeiros, B., Del Rio, M.G., Haran, J., Kuschel, G (2018) **Morphological and molecular**
perspectives on the phylogeny, evolution, and classification of weevils (Coleoptera:
Curculionoidea): Proceedings from the 2016 International Weevil Meeting *Diversity* 10:64
https://doi:10.3390/d10030064

McKenna, D.D., Farrell, B.D., Caterino, M.S., Farnum, C.W., Hawks, D.C., Maddison, D.R., Seago, A.E.,
Short, A.E., Newton, A.F. and Thayer, M.K (2015) **Phylogeny and evolution of S taphyliniformia**
and S carabaeiformia: forest litter as a stepping stone for diversification of nonphytophagous
beetles *Systematic Entomology* **40**:35–60.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M. and DePristo, M.A., 2010. **The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data** *Genome Research* **20**:1297–1303

McKenna, D.D., Sequeira, A.S., Marvaldi, A.E., Farrell, B.D (2009) **Temporal lags and overlap in the diversification of weevils and flowering plants** *Proceedings of the National Academy of Sciences* **106**:7083–7088

McKenna, D.D., Shin, S., Ahrens, D., Balke, M., Beza-Beza, C., Clarke, D.J., Donath, A., Escalona, H.E., Friedrich, F., Letsch, H., Liu, S (2019) **The evolution and genomic basis of beetle diversity** *Proceedings of the National Academy of Sciences* **116**:24729–24737

McLeish, M.J., Chapman, T.W., Schwarz, M.P (2007) **Host-driven diversification of gall-inducing *Acacia* thrips and the aridification of Australia** *BMC Biology* **5**:1–13

McLeish, M.J., Miller, J.T., Mound, L.A (2013) **Delayed colonisation of *Acacia* by thrips and the timing of host-conservatism and behavioural specialisation** *BMC Evolutionary Biology* **13**:1–15

Medeiros, M.J., Bianchi, G.L., Taschetta, L.R., Oboyski, P.T (2016) **A review of Polynesian *Carposina* Herrich-Schäffer (Lepidoptera: Carposinidae), with descriptions of four new species** *Zoological Journal of the Linnean Society* **177**:135–146

Miller, J.T., Murphy, D.J., Ho, S.Y., Cantrill, D.J., Seigler, D (2013) **Comparative dating of *Acacia*: combining fossils and multiple phylogenies to infer ages of clades with poor fossil records** *Australian Journal of Botany* **61**:436–445

Minh, B.Q., Hahn, M.W., Lanfear, R (2020) **New Methods to Calculate Concordance Factors for Phylogenomic Datasets** *Molecular Biology and Evolution* **37**:2727–2733 https://doi:10.1093/molbev/msaa106

Minh, B.Q., Nguyen, M.A.T., von Haeseler, A (2013) **Ultrafast approximation for phylogenetic bootstrap** *Molecular Biology and Evolution* **30**:1188–1195 https://doi:10.1093/ molbev/mst024

Misof, B., Liu, S., Meusemann, K., Peters, R.S., Donath, A., Mayer, C., Frandsen, P.B., Ware, J., Flouri, T., Beutel, R.G., Niehuis, O (2014) **Phylogenomics resolves the timing and pattern of insect evolution** *Science* **346**:763–767

Nagalingum, N.S., Marshall, C.R., Quental, T.B., Rai, H.S., Little, D.P., Mathews, S (2011) **Recent synchronous radiation of a living fossil** *Science* 334:796–799

Naumann, J., Salomo, K., Der, J.P., Wafula, E.K., Bolin, J.F., Maass, E., Frenzke, L., Samain, M.S., Neinhuis, C., DePamphilis, C.W., Wanke, S (2013) **Single-copy nuclear genes place haustorial Hydnoraceae within Piperales and reveal a Cretaceous origin of multiple parasitic angiosperm lineages** *PLoS One* 8:e79204

Nguyen, L.T., Schmidt, H.A., von Haeseler, A., Minh, B.Q (2015) **IQ-TREE: a fast and effective**
stochastic algorithm for estimating maximum-likelihood phylogenies *Molecular Biology and*
Evolution **32**:268–274 https://doi:10.1093/molbev/msu300

Oberprieler, R., Marvaldi, A., Anderson, R.S (2007) **Weevils, weevils, weevils everywhere. In:**
Linnaeus tercentenary: progress in invertebrate taxonomy *Zootaxa* **1668**:491–520

Oberprieler, R.G., Oberprieler, S.K (2012) *Talbragarus averyi* **gen.** *et* **sp. n., the first Jurassic weevil**
from the southern hemisphere (Coleoptera: Curculionoidea: Nemonychidae) *Zootaxa*
3478:256–266

O'Brien, C. W., Tang, W (2015) **Revision of the New World cycad weevils of the subtribe**
Allocorynina, with description of two new genera and three new subgenera (Coleoptera:
Belidae: Oxycoryninae) *Zootaxa* **3970**:1–87

O'Grady, P.M., Lapoint, R.T., Bonacum, J., Lasola, J., Owen, E., Wu, Y., DeSalle, R (2011)
Phylogenetic and ecological relationships of the Hawaiian *Drosophila* **inferred by**
mitochondrial DNA analysis *Molecular Phylogenetics and Evolution* **58**:244–256

Parham, J.F., Donoghue, P.C., Bell, C.J., Calway, T.D., Head, J.J., Holroyd, P.A., Inoue, J.G., Irmis, R.B.,
Joyce, W.G., Ksepka, D.T., Patané, J.S (2012) **Best practices for justifying fossil calibrations**
Systematic Biology **61**:346–59

Pease, J.B., Brown, J.W., Walker, J.F., Hinchliff, C.E., Smith, S.A. (2018) **Quartet Sampling**
distinguishes lack of support from conflicting support in the green plant tree of life
American Journal of Botany **105**:385–403 https://doi:10.1002/ajb2.1016

Poinar, G.O., Legalov, A.A (2014) *Pleurambus strongylus* **n. gen., n. sp. (Coleoptera: Belidae) in**
Dominican amber *Historical Biology* **26**:670–674

Price, J.P., Clague, D.A (2002) **How old is the Hawaiian biota? Geology and phylogeny suggest**
recent divergence *Proceedings of the Royal Society of London. Series B: Biological Sciences*
269:2429–2435 https://doi:10.1098/rspb.2002.2175

Rambaut, A., Drummond, A.J (2015) **TreeAnnotator v1.8.2** *MCMC Output Analysis.*

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A (2018) **Posterior summarization in**
Bayesian phylogenetics using Tracer 1.7 *Systematic Biology* **67**:901–904

Ree, R.H., Smith, S.A (2008) **Maximum likelihood inference of geographic range evolution by**
dispersal, local extinction, and cladogenesis *Systematic Biology* **57**:4–14
https://doi:10.1080/10635150701883881

Revell, L.J. (2012) **Phytools: an R package for phylogenetic comparative biology (and other things)**
Methods in Ecology and Evolution **3**:217–223

Ronquist, F. (1997) **Dispersal-vicariance analysis: a new approach to the quantification of historical**

biogeography *Systematic Biology* **46**:195–203 https://doi:10.1093/sysbio/46.1.195

Salas-Leiva, D.E., Meerow, A.W., Calonje, M., Griffith, M.P., Francisco-Ortega, J., Nakamura, K.,

Stevenson, D.W., Lewis, C.E., Namoff, S (2013) **Phylogeny of the cycads based on multiple**

single-copy nuclear genes: congruence of concatenated parsimony, likelihood and species tree

inference methods *Annals of Botany* 112:1263–1278

Salzman, S., Crook, D., Crall, J.D., Hopkins, R., Pierce, N.E (2020) **An ancient push-pull pollination**

mechanism in cycads *Science Advances* 6:p.eaay6169

Santos, M.F.A., Mermudes, J.R.M., Fonseca, V.M.M. (2007) **Descrição de um novo gênero e espécie de**

Belinae (Belidae, Curculionoidea, Coleoptera) da Formação Santana (Membro Crato,

Cretáceo Inferior) da Bacia do Araripe, Nordeste do Brasil *Paleontologia: Cenários de Vida*

441–447

Santos, M.F.D.A., Mermudes, J.R.M., Fonseca, V.M.M.D. (2011) **A specimen of Curculioninae**

(Curculionidae, Coleoptera) from the Lower Cretaceous, Araripe Basin, north-eastern Brazil

Palaeontology **54**:807–814

Sequeira, A.S., Normark, B.B. and Farrell, B.D (2000) **Evolutionary assembly of the conifer fauna:**

distinguishing ancient from recent associations in bark beetles *Proceedings of the Royal Society*

of London. Series B: Biological Sciences **267**:2359–2366

Schneider, H., Schuettpelz, E., Pryer, K. M., Cranfill, R., Magallón, S., Lupia, R (2004) **Ferns**

diversified in the shadow of angiosperms *Nature* **428**:553–557

Schuettpelz, E., Pryer, K.M (2009) **Evidence for a Cenozoic radiation of ferns in an angiosperm-**

dominated canopy *Proceedings of the National Academy of Sciences* **106**:11200–11205

Shin, S., Clarke, D.J., Lemmon, A.R., Moriarty Lemmon, E., Aitken, A.L., Haddad, S., Farrell, B.D.,

Marvaldi, A.E., Oberprieler, R.G., McKenna, D.D (2018) **Phylogenomic data yield new and**

robust insights into the phylogeny and evolution of weevils *Molecular Biology and Evolution*

35:823–836

Simon, A., Salzman, S., Specht, C.D., Raguso, R.A (2023) **Behavior and feeding of two beetle**

pollinators of *Zamia integrifolia* **(Cycadales):** *Rhopalotria slossoni* **(Coleoptera: Belidae) and**

Pharaxanotha floridana **(Coleoptera: Erotylidae)** *Florida Entomologist* **106**:199–202

Smith, S.A., Brown, J.W., Walker, J.F (2018) **So many genes, so little time: a practical approach to**

divergence-time estimation in the genomic era *PLoS ONE*, **13**:e0197433.

https://doi:10.1101/114975

Song, H., Béthoux, O., Shin, S., Donath, A., Letsch, H., Liu, S., McKenna, D.D., Meng, G., Misof, B.,

Podsiadlowski, L., Zhou, X (2020) **Phylogenomic analysis sheds light on the evolutionary**

pathways towards acoustic communication in Orthoptera *Nature Communications* **11**:1–16

Strimmer, K., von Haeseler, A (1997) **Likelihood-mapping: a simple method to visualize phylogenetic**
content of a sequence alignment *Proceedings of the National Academy of Sciences USA* **94**:6815–
6819 https://doi:10.1073/pnas.94.13.6815

Suchard, M.A., Lemey, P., Baele, G., Ayres, D.L., Drummond, A.J., Rambaut, A. (2018) **Bayesian**
phylogenetic and phylodynamic data integration using BEAST 1.10 *Virus Evolution* **4**:vey016
https://doi:10.1093/ve/vey016

Sunnucks, P., Hales, D.F (1996) **Numerous transposed sequences of mitochondrial cytochrome**
oxidase I-II in aphids of the genus *Sitobion* (Hemiptera: Aphididae) *Molecular Biology and*
Evolution **13**:510–524

Swain, A., Maccracken, S.A., Fagan, W.F., Labandeira, C.C (2022) **Understanding the ecology of host**
plant–insect herbivore interactions in the fossil record through bipartite networks
Paleobiology **48**:239–260. https://doi:10.1017/pab.2021.20

Tang, W. (1987) **Insect pollination in the cycad *Zamia pumila* (Zamiaceae)** *American Journal of*
Botany **74**:90–99 https://doi.org/10.1002/j.1537-2197.1987.tb08582.x

Teasdale, L.C., Köhler, F., Murray, K.D., O'Hara, T. and Moussalli, A (2016) **Identification and**
qualification of 500 nuclear, single-copy, orthologous genes for the Eupulmonata (Gastropoda)
using transcriptome sequencing and exon capture *Molecular Ecology Resources* **16**:1107–1123

Vanin, S.A. (1976) **Taxonomic revision of the South American Belidae (Coleoptera)** *Arquivos de*
Zoologia **28**:1–75

Wang, X.Q., Ran, J.H (2014) **Evolution and biogeography of gymnosperms** *Molecular Phylogenetics*
and Evolution **75**:24–40

Wilf, P. (2008) **Insect-damaged fossil leaves record food web response to ancient climate change and**
extinction *New Phytologist* **178**:486–502

Xie, W., Lewis, P.O., Fan, Y., Kuo, L., Chen, M.H (2011) **Improving marginal likelihood estimation**
for Bayesian phylogenetic model selection *Systematic Biology* **60**:150–160
https://doi.org/10.1093/sysbio/syq085

Yu, Y.L., Davis, S.R., Shih, C.K (2019) **The earliest fossil record of Belidae and its implications for**
the early evolution of Curculionoidea (Coleoptera) *Journal of Systematic Palaeontology*
17:2105–2117

Zhang C, Rabiee M, Sayyari E, Mirarab S. (2018) **ASTRAL-III: Polynomial time species tree**
reconstruction from partially resolved gene trees *BMC Bioinformatics* **19**:15–30
https://doi.org/10.1186/s12859-018-2129-y

Zhang, J., Gordon, E.R., Forthman, M., Hwang, W.S., Walden, K., Swanson, D.R., Johnson, K.P., Meier,
R., Weirauch, C (2016) **Evolution of the assassin's arms: insights from a phylogeny of combined**

transcriptomic and ribosomal DNA data (Heteroptera: Reduvioidea) *Scientific Reports* **6**:22177

https://doi.org/10.1038/sre p22177

Zimmerman, E.C. (1994) *Australian Weevils (Coleoptera: Curculionoidea). Volume I. Orthoceri.*

Anthribidae to Attelabidae. The Primitive Weevils. CSIRO Australia, Melbourne, xxxii + 741 pp

 Figure legends

Figure 1. Phylogeny estimate of Belidae. Topology obtained from analysis of the 46t424g matrix via

maximum likelihood (ML), partitioned by gene. Squares along branches are ML SH-aLRT and UFBoot

values for 1–3: 1. partition by gene, 2. unpartitioned, and 3. partition by gene and codon positions. Astral

support for 4, Quartet Concordance (QC) for 5, and gene concordance factor (gCF) for 6. Branches

without squares indicate the highest support in all analyses. Taxa displayed on the right: a. *Cyrotyphus*

vestitus (Agnesiotidini) (photo credit: Rolf Oberprieler), b. *Pachyura australis* (Pachyurini) (photo credit:

Simon Grove), c. *Isacanthodes ganglionicus* (Belini) (photo credit: Rolf Oberprieler), d. *Homalocerus* sp.

(Belini) (photo credit: Jeff Gruber), e. *Rhinotia* sp. (Belini) (photo credit: Rolf Oberprieler), f.

Oxycorynus missionis (Oxycorynini, Oxycorynina) (photo credit: Adriana Marvaldi), g. *Rhopalotria*

slossoni (Oxycorynini, Allocorynina) (photo credit: Shayla Salzman).

Figure 2. Belid timetree. Chronogram generated using BEAST and a birth-death tree prior with 13

unlinked molecular clocks; pale green diamonds on nodes indicate calibration points. Scale is in Ma.

Horizontal bars depict the 95% HPD density of each estimate, cyan bars generated by birth-death tree

prior, grey bars generated by Yule tree prior. Ancestral area estimation under the likelihood-based

implementation of dispersal vicariance analysis and unconstrained dispersal rates (DIVALIKE, M0)

performed with BioGeoBEARS. The distribution of each species is mapped to the right of the

chronogram. A single most probable ancestral area is mapped at each node. Ancestral state reconstruction

of belid larval host plant group and host organ usage under the ER model.

Table 1. Marginal-likelihood estimate (MLE) scores for various BEAST analyses performed for this

study, and estimated ages (in Ma) for Belidae crown nodes for each tree prior/clock scheme in BEAST.

Notes: SS, stepping-stone sampling marginal-likelihood estimation; PS, path-sampling marginal-

likelihood estimation; median post-burn-in divergence times in millions of years (95% credibility

interval).

analysis	tree model	clock model	MLE SS	MLE PS	crown Belidae age (Ma)
A1	birth-death	1 ULRC	-142024.6921	-142025.2950	143.8706 [127.5175–159.9967]
A2	Yule	1 ULRC	-142024.2190	-142024.9031	144.1718 [128.1144–159.9990]
A3	birth-death	4 ULRC	-135931.3312	-135932.0701	140.7353 [125.4955–157.7755]
A4	Yule	4 ULRC	-135931.0581	-135931.8383	140.8507 [125.7566–157.9643]
A5	**birth-death**	**13 ULRC**	**-134225.1645**	**-134225.9867**	**138.4569 [125.5653–154.8667]**
A6	Yule	13 ULRC	-134226.8380	-134227.8971	138.3267 [125.5619–

154.5464]

901
902

Figure S1. (a) Belidae phylogenetic tree inferred from the 33t424g matrix via maximum likelihood (ML),

partitioned by gene and codon position. Squares along branches are ML SH-aLRT and UFBoot values.

Supporting values are based on the 33t424g matrix for 1–3 and the 33t423g matrix for 4–6. Partitioned by

gene and codon position for 1 and 4, partitioned by gene for 2 and 5, and unpartitioned for 3 and 6. (b)

Four-cluster likelihood mapping (FcLM) result based on the 33t424g matrix. (c) FcLM result based on

the 33t423g matrix.

Figure S2. Topology generated by maximum likelihood (ML) analysis of the 46t424g matrix partitioned

by gene, with node numbers.

Table S1. Taxa sampled in the present study, with the number of loci retained after all cleaning steps,

host plant group and organ coding schemes, and distribution coding schemes.

Table S2. Summary of node support generated by concatenated ML analysis, coalescence approach, gene

concordance factors (gCF), and Quartet Sampling; the posterior possibility of characters generated by

ancestral state reconstructions.

Table S3. Results of the BioGeoBEARS analyses.

Orsodacne cerasi
Diabrotica undecimpunctata
Bunyaeus montheithi
Rhynchitomacerinus kuscheli
Urodontus mesemoides
Dendropemon albopictus
Araecerus sp.1
Cimberis pilosa

Dicordylus binotatus
Agnesiotis pilosula — **Agnesiotidini**

Rhicnobelus metallicus
Apagobelus brevirostris
Basiliobelus flavovittatus
Basiliobelus lepidus — **Pachyurini**

Trichophthalmus miltomerus
Isacantha dermestiventris
Isacanthodes sp.1
Araiobelus sp.1
Homalocerus lyciformis
Rhinotiodes spinipennis
Rhinotia haemoptera
Rhinotia cf. *suturalis*
Rhinotia lineata
Rhinotia phoenicoptera — **Belini**

Oxycraspedus cornutus
Oxycraspedus cribicollis
Oxycraspedus minutus — **Oxycorynini** / Oxycraspedina

Metrioxena sp.2
Metrioxena sp.1 — **Metrioxenini**

Aglycyderes setifer
Aralius sp.1
Proterhinus sp.1 — **Aglycyderini**

Oxycorynus missionis
Hydnorobius parvulus
Hydnorobius hydnorae
Hydnorobius helleri — **Oxycorynini** / Oxycorynina

Archicorynus kuscheli — **Archicorynini**

Protocorynus bontai
Parallocorynus bicolor
Parallocorynus salasae
Notorhopalotria montgomeryensis
Notorhopalotria panamensis
Rhopalotria calonjei
Rhopalotria mollis
Rhopalotria dimidiata
Rhopalotria slossoni — **Oxycorynini** / Allocorynina

Belinae (Agnesiotidini, Pachyurini, Belini)

Oxycoryninae

0.2

Legend

Unpartitioned

PGM	1 2 3	PCM
Astral	4 5 6	gCF
	QC	

■ SH-aLRT ≥ 80 AND UFBoot ≥ 95
▨ SH-aLRT ≥ 80 OR UFBoot ≥ 95
□ SH-aLRT < 80 AND UFBoot < 95

■ Astral support ≥ 0.9
▨ 0.9 > Astral support ≥ 0.7
□ 0.7 > Astral support > 0

■ QC > 0.2
■ 0.2 ≥ QC > 0
■ 0 ≥ QC ≥ -0.05
■ QC < -0.05

■ gCF > gDF1 AND gCF > gDF2
□ gCF = gDF1 OR gCF = gDF2
■ gCF < gDF1 OR gCF < gDF2

◩ Clade not recovered

a
b
c
d
e
f
g
h