Hybrid Exb/Mot stators require substitutions distant from the chimeric pore to power flagellar rotation

Authors

Pietro Ridone[1], Matthew A. B. Baker[1].

[1]School of Biotechnology and Biomolecular Sciences, UNSW Sydney, Australia.

Abstract

Powered by ion transport across the cell membrane, conserved ion-powered rotary motors (IRMs) drive bacterial motility by generating torque on the rotor of the bacterial flagellar motor. Homologous heteroheptameric IRMs have been structurally characterized in ion channels such as Tol/Ton/Exb/Gld, and most recently in phage defense systems such as Zor. Functional stator complexes synthesized from chimeras of PomB/MotB (PotB) have been used to study flagellar rotation at low ion-motive force achieved via reduced external sodium concentration. The function of such chimeras is highly sensitive to the location of the fusion site, and these hybrid proteins have thus far been arbitrarily designed. To date, no chimeras have been constructed using interchange of components from Tol/Ton/Exb/Gld and other ion powered motors with more distant homology.

Here we synthesised chimeras of MotAB, PomAPotB and ExbBD to assess their capacity for cross-compatibility. We generated motile strains powered by stator complexes with B-subunit chimeras. This motility was further optimised by directed evolution. Whole genome sequencing of these strains revealed that motility-enhancing residue changes occurred in the A-subunit and at the peptidoglycan binding domain of the B-unit, which could improve motility. Overall, our work highlights the complexity of stator architecture and identifies the challenges associated with rational design of chimeric IRMs.

Importance

Ion-powered rotary motors (IRMs) underpin the rotation of one of nature's oldest wheels, the flagellar motor. Recent structures have shown that this complex drives ion conduction and even phage defence and thus appears to be a fundamental molecular module with diverse biological utility where electrical energy can be coupled to rotational force to execute work.

Here, we attempted to rationally design chimeric IRMs to explore the cross-compatibility of these ancient motors. We succeeded in making one working chimera of a flagellar motor and a non-flagellar transport system protein. This had only a short hybrid stretch in the ion-conducting channel, and function was subsequently significantly improved through additional substitutions at sites distant from this hybrid pore region. Our goal was to test the cross-compatibility of these homologous systems and highlight challenges arising when engineering new rotary motors from this fundamental nanomotor.

Introduction

Many bacteria are propelled by the rotation of flagella to navigate to better environments. The rotation of these flagella is powered by ion-powered rotary ion-pumps known as stators[1–3]. Recent structural characterization of the stator elements that are ion conductors revealed that these are heteroheptamers ($MotA_5MotB_2$[1, 2]), and this has now been observed in sodium powered stators ($PomA_5PomB_2$[3]), in ion pumps ($GldM_5GldN_2$[4], $ExbB_5ExbD_2$[5–7]) and most recently in phage defense systems ($ZorA_5ZorB_2$[8]). These ion-powered rotary machines (IRMs)[9] have diverse functions but high structural conservation implying a common origin. Thus, IRMs represent a key molecular complex with an origin at least as ancient as flagellar motility[10, 11]. As such there is significant interest in understanding what the structure and function of a potential common ancestor might be.

The flagellar motor itself is highly modular (Fig.1). This organization is routinely described as the filament, hook and basal body[12], an oversimplification which does not even include the IRMs as a separate module. We consider submodules of the motor to be identifiable domains that have a specific function: in the case of the stator-complex, this is commonly defined for the central B-unit as the transmembrane (TM) domain, the linker, and the peptidoglycan-binding domain (PGB, in gram negative species such as *E. coli*). As more structural information becomes available, it is also possible to clearly identify pore-facing, rotor-interacting, and lipid-facing domains in the pentameric A-unit[13–15].

Chimeric work in the stator complex is not new: the first Mot/Pom chimeras were designed in the 1990s with the target of generating a sodium-powered chimera that was functional in *E. coli* (16, 17). Mutagenesis and protein engineering work exploring the role of the plug, linker and peptidoglycan-binding domains in light of new structures continues[18–23] as well as efforts to reconstruct an ancestral stator to explore origins and divergence[14, 24]. However, to our knowledge no attempt has been made to combine homologous A_5B_2 pumps coupled to non-flagellar function with existing stator components.

In this work, we seek to explore the possibility of engineering functional stator chimeras. We explore compatibility of submodules of IRMs and test these for their capacity to restore motility in stator-deleted bacterial strains. We show the capacity of the overall flagellar complex to rescue motility through epistatic substitutions that affect residues far from chimeric interfaces, revealing the complex integration of submodules across the complex. In doing so, we explore the limits of function as well as the potential for customizing new functions and determining essential protein motifs which differentiate structure and function across the IRM classes.

Results

We designed chimeras of *motA* and *exbB* (the A-subunit) and *motB exbD* (the B-subunit) by performing amino acid alignments between homologs and by manual inspection of their 3D structures to select the residues connecting the chimeric portions of our constructs. We selected 4 A-subunit and 11 B-subunit constructs to clone into a plasmid vector (pBAD33) for *in vivo* testing in a stator-deleted strain (*E. coli* ΔmotAB) (SI Fig. 1A, 1B).

Each construct was maintained in culture on a soft agar swim plate for up to 2 weeks to allow for the isolation and propagation of up-motile populations. Out of all 14 tested constructs we isolated one *potB / exbD* variant ($PotD_{3.3}$) able to restore motility in presence of *pomA* (Fig. 2). The plasmid carried by this motile population was extracted and both stator genes were sequenced to check for additional mutations contributing to the phenotype. A missense mutation (*CAA* to *CCA*) causing the Q104P substitution was found in the plasmid-encoded chimeric construct $potD_{3.3}$. (SI Fig. 2). This stator B-subunit construct ($potD_{3.3}$ *Q104P*) was renamed "PotDα".

To verify that Q104P was the substitution that rescued motility, we deleted *pomA* from the mutant
plasmid and tested the function of PotDα in combination with various A-subunits.
PotDα was tested in presence of *motA* and all chimeric A-subunit constructs to check whether together
they could form a functional stator (Fig. 3A, 3B). Interestingly, none of our strains expressing *potDα* / A-
subunit combinations produced swim rings within 24 hr (SI Fig. 3A) whereas cells containing control
combinations (*motAmotB*, *motApotB* and *pomApotB*) all formed swim rings on plates and produced
observable swimming cells (Fig. 3C) powered by proton (*motA*) or sodium (*pomA*). Thus, none of our A-
subunit chimeras functioned in presence of any B subunit partner, whether chimeric or wild-type. Notably,
bacteria co-expressing the *mobB.v1 potDα* combination showed a slower growth phenotype when
compared to *motAmotB* control (SI Fig.3B), suggesting that the stator construct might leak ions although
being unable to generate torque.

Because the Q104P substitution alone was unable to rescue motility, we screened for improved swimmers
by directed evolution to isolate additional mutations in stator genes that might have contributed to
phenotypic rescue. We subjected our motile PotDα isolate and all non-motile, B-subunit chimeras to
repeated passaging on fresh swim plates at 3-4 days intervals. Initially, swim rings of the *pomApotDα*
strain spread noticeably slower than the *pomApotB* strain but after six passages motility had improved
considerably. Visible swim rings of *pomApotDα* formed within a 24-hr period (SI Fig. 4A) compared to the
multiple days required for the original flare (SI Fig. 2). The *potDα* variant was motile not only in the
presence of both *motA* and *pomA* but also when only *pomA* was present (tested in Δ*motB E. coli* RP3087,
motB on genome, *pomA* on plasmid; or *E. coli* Δ*motAB,* only *pomA* from plasmid). Plasmid sequencing of
this variant revealed an additional mutation in the plasmid-encoded *pomA* gene (causing the substitution
A86V). All other B-subunit constructs, which did not support motility, showed no rescue of motility after
6 passages.

We used genome-based directed evolution of motility(25) to attempt to further improve motility in our
chimeric variant. We replaced the native *motAB* locus of *E. coli* RP437 with the *pomApotDα* construct
cloned from the plasmid (SI Fig.4A, 4B). As observed in directed evolution experiments of strains
expressing the chimeras from a plasmid vector, flares developed over multiple days, and motility
improved after several passages (SI Fig. 4B.) A control genomic edit using the original *potD*$_{3.3}$ construct (*E.*
coli "PotD", SI Fig.5A, 5B), without additional mutations, did not adapt to rescue motility (SI Fig. 5C).
Whole-genome sequencing of members of our PotDα evolved lineage revealed a new missense mutation
in the chromosomal *pomA* gene (substitution H116Y) and additional mutations in the *flgM* and *icd* genes
(SI Fig. 4B, SI Table 2).

To repeat the effect of these stator mutations, we introduced them into the plasmid-encoded *pomApotDα*
constructs via site-directed mutagenesis and measured their effect on motility in our stator-deleted strain
(*E. coli* Δ*motAB*). Growth profiles of all our evolved and engineered strains showed no effect on growth
when *potDα* was expressed from plasmids, while growth was inhibited when the construct was expressed
from the edited locus on the chromosome (SI Fig. 6A). At the same time, we tested the effect on swim
rings in the presence and absence of sodium to determine the ion-selectivity of the chimeric stators (Fig.
4A, SI Fig. 6B). Our strains expressing chimeric stators appeared to require Na$^+$ for effective motility and
did not produce swim rings on K$^+$ swim plates. Motility was inhibited in presence of the Na$^+$-powered
stator blocker Phenamil (SI Fig. 6B). Our plasmid-based validation showed that the A86V/H116Y residue
changes in PomA improved the motility phenotype when co-expressed with *potDα* (Fig. 4B). Although the
Q104P change in PotD$_{3.3}$ was located at the N-terminal end of the PG-binding domain, both *pomA*
mutations affected residues on the cytoplasmic side of the stator near the surface that interacts with the
rotor (FliG, YcgR). Thus, mutations that improve motility can affect residues located far from each other
in the folded protein structure, indicating that they improve different aspects of stator function.

Discussion

We engineered 11 chimeras of the B subunit of the hybrid PomAPotB stator with the ExbD component of the ExbBD stator. These hybrids had varying extents of ExbD, ranging from 5.3% to 16.95 %. Of these, the PotD$_{3.3}$ construct with the smallest contribution from ExbD was able to generate torque and rescue motility following subsequent adaptation with directed evolution. None of our A-subunit chimeras were functional or rescued by directed evolution. However, some mutations in *pomA* improved motility when expressed with *potD$_{3.3}$*. These mutations affected regions of PomA that are far from the ion pore but proximal to the stator-rotor interface.

Fusion-sites are critical in the B-unit

It is challenging to determine why some chimeras are functional where others are not, especially considering that some B-unit chimeras differed only very slightly. Chimeras that differed by only 4 residues (PotD$_{3.0}$), in non-conserved regions adjacent to the pore were unable to rescue motility. This is indicative of the general challenge of rational design and the need to select fusion sites carefully and attempt multiple variants simultaneously to improve throughput. Single amino acid replacements within the TM domain of *Ec*MotB are likely to disrupt motility even when they are conservative[26].

Preservation of the pore proximal A-B interface does not guarantee function.

To date, no functional A-unit chimeras have been made, either from Mot/Pom complexes or across the ion-powered rotary motors more widely. Here, we matched A-B interfaces across the TM region of *Ec*MotB (residues I28-W45) and TM3/4 in *Ec*MotA (residues N139-N250) (SI Fig. 7A, 7B). Despite this, none were functional in driving torque-generation and motility. One combination of chimeric variants aimed at reconstituting a complete ExbBD-like pore (*mobB.v1 potDα*) caused growth-delaying effects when expressed, suggesting that it may be able to conduct ions without coupling this current to torque generation (SI Fig. 7C, 7D). Construct MobB v2, which contains the C-terminal domain of ExbB (SI Fig. 1), did not show any growth defects when it was expressed, suggesting that the C-terminal residues of MotA (residues 247- 295 of *Ec* MotA) are required for ion conduction[3, 27].

The pore interface of PomA is compatible with both PotB and PotDα, but not with MotB, PotD 4.3 (SI Fig. 1) or a PotB variant expressing the TM segment of MotB (PotB-H$^+$, SI Fig. 8), suggesting that the PomB portion of the chimera is crucial to allow PotDα to function in a sodium-dependent fashion. The MotAB stator thus appears to be less likely to allow functional chimeras with ExbD compared with the Pom stator. When testing the function of our chimeric variants we did not investigate whether the inability of some constructs to rescue motility was due to poor expression via Western blot or RT-PCR.

Despite this, our results indicate that regions outside the pore are important in improving engineered pores. However, when the lipid and rotor interfaces were preserved by using a single A-subunit with known functional domains in these locations (*mobB.v1*, *mobB.v2* and *potA*, Fig. 2A), these chimeras were still non motile.

Allosteric substitutions further improve motility in partially functional stators.

Motility was rescued in *potD$_{3.3}$* via an amino acid substitution at a residue (Q104) located in a different protein domain than the chimeric TM pore domain. The substitution introduced a proline on the first alpha helix of the large PGB domain. This substitution might contribute to the stability of the binding to

the peptidoglycan layer and the mechanosensitivity of the chimeric stator. A previously reported
substitution (R109P) found in a linker-deleted PotB variant rescued the motility of the host[28],
emphasizing that the PGB-linker connection is probably structured.

Subsequent mutations affecting sites far from the hybrid region improved motility and caused residue
substitutions near a known FliG interaction site of the A-subunit (PomA) at A86 and H116. Regions far
from the pore not only affect function but also repair a hybrid pore to enable function. Substitutions
affecting sites far from the pore have been shown to affect phenamil binding and resistance[29],
putatively allowing a defective stator to remain engaged with the rotor for longer. It is possible that
stabilizing the PG-binding of the stator complex would enable motility driven by a partially functional
stator. Such allosteric interactions undermine attempts to rationally design proteins through edits at only
a single functional site. Methods to examine allosteric networks across proteins can help these attempts
(30–32). Mutations in class III flagellar gene repressor FlgM (anti-FliA, σ^{28} factor) have been shown to
impact motility[33], but whether the FlgM S7Y substitution affected the efficiency of FlgM secretion
remains to be determined. The Ser7 residue in FlgM is distant from the interaction interface between FliA
and FlgM (SI Fig. 9) and might not affect the ability of FlgM to bind to FliA.

Limitations of testing chimeric constructs in gram-negative E. coli .

We sought to design chimeras that preserved ion conduction, torque generation and mechanosensitivity
and approached this via engineering conservative chimeras centered around the pore. The stator complex
has multiple functions, executed by specific domains of the complex that are either pore-facing (TM3-4
residues N139-N250 in *Ec*MotA), rotor-facing (residues L81-M138) or PG-binding (residues N102-L288 in
*Ec*MotB)[23]. Not only must the stator complex sense force to activate and transduce chemical potential
into torque, but it must also interact with the rotor to couple torque to filament rotation. The rotor itself
is a large molecular complex with multiple functions that include the conformational rearrangements
necessary to change the direction of rotation as well as the conformational changes necessary to generate
torque to drive rotation.

All chimeras in our work were only tested in an *E. coli* chassis. Motors in different species have different
protein and membrane composition, different sizes, as well as different stator dynamics(34–36). Thus, it
is possible our stators are functional but require rotor and membrane environments that are better
matched to their chimeric pores. Single-molecule *in vitro* bilayer measurements combined with
electrophysiology(37, 38) could provide a platform to assay the role that lipid composition and membrane
environment have in affecting the diffusion and ligand-binding of the IRM. This could elucidate which
submodules of the motor retain partial functionality, whereas in our *in vivo* motility assays we are only
able to test overall function as presence or absence of motility.

Directed and experimental evolution improves function rapidly.

As in our previous work, during iterative swim plating we observed mutants that upregulate filament
expression(25, 39). Our plasmid-based and genome-directed evolution generated convergent results.
Substitutions affecting residues far from the pore rescued motility within days. These screens allow us to
explore epistatic interactions that rescue motility, such as non-pore residues improving function in
environments depleted of ions for conduction. However, a simple swim plate does confound growth,
motility and chemotaxis, so it does not select only for mutations that alter the capacity for the stator
complex to generate torque[25]. Nevertheless, we do see substitutions that repair edits and improve

performance on swim plates. These effects can be correlated with other measurements, such as tethered
cell assays(40–42).

More specific screens that directly target motility(43) or cell energetics could be applied to select for
mutations that affect the function of specific submodules of the stator complex. Likewise, *in vitro* methods
using purified membrane protein complexes could facilitate determination of the *in situ* structure of the
stator within the membrane(44), or used to measure current flow and other biophysical parameters(5).
Fluorescently labelled stators could be visualized *in vitro* to detect conformational changes at the single-
molecule level(45, 46). Improved preparation of combinatoric libraries using newer approaches in
synthetic biology could improve throughput(47–50). We utilized medium throughput biological methods
that are still able to select one-in-a-million up-motile variants(29). Future high-throughput screens will
require automated colony passaging by liquid-handling or colony-picking robots.

Implications for the evolution and common ancestry of IRMs.

Our experimental results might reflect a closer evolutionary relationship between the Exb and Pom
systems than Exb and Mot. The Exb system is nevertheless largely incompatible with the BFM stator
systems given how many of our chimeras failed to confer motility. An in-depth phylogenetic and structural
assessment of the IRM protein family will be required to confirm this hypothesis.

Conclusions

Our work shows that even with as few as 12 amino acid substitutions, cut-site location and chimeric design
is challenging and requires high-throughput approaches to succeed. Recent structural characterization of
structural homologues of the stator complex, including Zorya, has shown the diversity of biological
applications where IRMs can transduce energy from ion gradients to mechanical work. Knowing the
structure of the common ancestor of these IRMs would shed light on the origin of the stator complex, as
well as other complexes, and this may be a key innovation of function in early life.

Construction of chimeras exchanging components between extant homologues does not necessarily
inform the structure and function of this common ancestor, but it does allow us to examine what are the
core elements of these complexes, as well as understand which modules are interchangeable and which
are not. This can inform the order of events under which the stator complex arose and diversified and
inform parsimony approaches to loss and gain of function in subsequent adaptation. Further studies with
new methods in statistical and structural phylogenetics are required to target ancestors of the stator
complex for characterization both structurally and biophysically.

Materials and Methods

E. coli strains, plasmids, and culture media

Synthetic stator chimeras were tested in stator-deleted derivative strains of *E. coli* RP437, either *E. coli*
Δ*motB* (RP3087(51)) or *E. coli* Δ*motAB* (SI Fig. 5). For genomic edits, *E. coli* RP437 was used as the parent
strain(52) to with the native *motAB* locus edited using No-SCAR methods, as previously (40). The
pSHU1234 (pBAD33 backbone, Cm+) plasmid encoding *pomA* and *potB* (29) was used as the vector to
clone and express all chimeric constructs.

Chimeric constructs were designed *in silico* and synthesized by Integrated DNA Technologies (IDT) as gene
fragments (gBlocks). Chimeras were cloned in plasmid pSHU1234 using restriction digestion with NdeI and

PstI (New England Biolabs). Plasmids carrying chimeric constructs were used as templates to generate the
double-stranded donor DNA used to replace the *motA* and *motB* gene on the RP437 chromosome with
pomA and *potD* variants, respectively (see Scar-less editing).

Point mutations in plasmids were generated using the QuikChange technique. Bicistronic plasmids
expressing both A and B stator subunits were converted to single-gene plasmids expressing only the B-
subunit via inverse deletion PCR. Single-construct plasmids expressing A-subunits were constructed via
restriction digestion (NdeI and PstI, New England Biolabs) and ligation using pBAD24 (Amp+) as a plasmid
backbone. The PotB TM replacement with the MotB TM domain (PotB-H$^+$ construct) was generated by
inverse deletion PCR followed by self-ligation of the linearized PCR product. Cloned constructs were
confirmed by Sanger sequencing (Ramaciotti Centre for Genomics, Kensington). Gene expression from all
plasmids used in this work was induced using 1 mM L-(+)-Arabinose (A3256, Sigma-Aldrich) in liquid and
solid media.

Liquid cell culturing was done using LB broth (85 mM NaCl or 85 mM KCl, 0.5% yeast extract (70161, Sigma-
Aldrich), and 1% Bacto tryptone). Cells were cultured on agar plates composed of LB broth and 1% Bacto
agar (BD Biosciences, USA). Swim plate cultures were performed on the same LB broth-based substrates
adjusted for agar content (0.25% Bacto agar). Inhibition of Na+-dependent motility was performed using
100 µM phenamil methanesulfonate (P203, Sigma-Aldrich).

261 Scar-less editing of *E. coli* with Cas9-assisted recombineering

This procedure was adapted from the no-SCAR method(53). The target strain to be edited (E. coli RP437)
was sequentially transformed first with the pKD-sgRNA-3'*motA* (Sm+) plasmid, encoding a single guide
RNA (sgRNA) sequence directed at the 3' end of *motA*, and then with the pCas9cr4 (Cm+) plasmid to yield
a parent strain harboring both plasmids. The overlap extension PCR technique(54) was used to assemble
linear double-stranded DNA (dsDNA) molecules using two (knock-out) or three (knock-in) starting dsDNA
fragments. The resulting donor DNA was electroporated in the plasmid-bearing host, and the successfully
edited clones were selected via colony PCR and Sanger sequencing of the *motAB* locus. A list of
oligonucleotide primers and synthetic gene fragments (gBlocks) used in this work is provided in
Supplementary Table 1.

271 Free-swimming assay preparation and analysis

272 Motile cells were grown overnight in tryptone buffer (TB: tryptone, 85 mM NaCl) at 30°C to OD600 of
273 ~0.5, diluted 100x in TB and loaded for imaging in a tunnel slide. A custom LabVIEW software was used as
274 previously(24, 29, 40) reported to measure the speed of swimming cells. Time-lapse videos were collected
275 using a camera (Chameleon3 CM3, Point Grey Research) recording 20-s-long videos at 20 frames/s.
276 Visualization of the data was performed using GraphPad Prism 8.

277 Sequencing and genomics

WGS of *E. coli* strains was performed using a MiSeq 2 × 150–base pair (bp) chip on an Illumina sequencing
platform. Sequencing was carried out at the Ramaciotti Centre for Genomics, Kensington and delivered
as demultiplexed fastQ files (quality control: >80% bases higher than Q30 at 2 × 150 bp). The SNP calling
and analysis were performed using Snippy(55–57). The short reads from sequencing were aligned to the
MG1655 reference *E. coli* genome (GenBank: U00096.2) and an *E. coli* RP437 (GenBank: CP083410.1) as
previously (25). A summary of all single nucleotide polymorphisms (SNPs) detected in this study is
provided in Supplementary Table 2.

Structural modelling

Sequence alignments were aligned using MAFFT. Models of stator monomers and heptameric complexes was performed using AlphaFold2 on Galaxy.au, using the Australian Alphafold Service (AAS). The software was run in multimer mode using FASTA files as input(58, 59). Models were visualized and prepared for figures using PyMOL.

Acknowledgements

M.A.B.B. acknowledges funding support from a Scientia Fellowship from UNSW, Human Frontiers Science Program Project Grant No. RGY0072/21 and U.S. Navy Office of Naval Research Research Grant No. N62909-22-1-2051.

References

1. Santiveri M, Roa-Eguiara A, Kühne C, Wadhwa N, Hu H, Berg HC, Erhardt M, Taylor NMI. 2020. Structure and Function of Stator Units of the Bacterial Flagellar Motor. Cell 183:244-257.e16.

2. Deme JC, Johnson S, Vickery O, Aron A, Monkhouse H, Griffiths T, James RH, Berks BC, Coulton JW, Stansfeld PJ, Lea SM. 2020. Structures of the stator complex that drives rotation of the bacterial flagellum. 12. Nat Microbiol 5:1553–1564.

3. Hu H, Popp PF, Santiveri M, Roa-Eguiara A, Yan Y, Martin FJO, Liu Z, Wadhwa N, Wang Y, Erhardt M, Taylor NMI. 2023. Ion selectivity and rotor coupling of the Vibrio flagellar sodium-driven stator unit. 1. Nat Commun 14:4411.

4. Hennell James R, Deme JC, Kjær A, Alcock F, Silale A, Lauber F, Johnson S, Berks BC, Lea SM. 2021. Structure and mechanism of the proton-driven motor that powers type 9 secretion and gliding motility. 2. Nature Microbiology 6:221–233.

5. Celia H, Noinaj N, Zakharov SD, Bordignon E, Botos I, Santamaria M, Barnard TJ, Cramer WA, Lloubes R, Buchanan SK. 2016. Structural insight into the role of the Ton complex in energy transduction. Nature 538:60–65.

6. Celia H, Noinaj N, Buchanan SK. 2020. Structure and Stoichiometry of the Ton Molecular Motor. Int J Mol Sci 21:375.

7. Maki-Yonekura S, Matsuoka R, Yamashita Y, Shimizu H, Tanaka M, Iwabuki F, Yonekura K. 2018. Hexameric and pentameric complexes of the ExbBD energizer in the Ton system. eLife 7:e35419.

8. Hu H, Hughes TCD, Popp PF, Roa-Eguiara A, Martin FJO, Rutbeek NR, Hendriks IA, Payne LJ, Yan Y, Sousa VK de, Wang Y, Nielsen ML, Berry RM, Erhardt M, Jackson SA, Taylor NMI. 2023. Structure and mechanism of Zorya anti-phage defense system. bioRxiv https://doi.org/10.1101/2023.12.18.572097.

9. Nandel V, Scadden J, Baker MAB. 2023. Ion-Powered Rotary Motors: Where Did They Come from and Where They Are Going? 13. International Journal of Molecular Sciences 24:10601.

10. Beeby M, Ferreira JL, Tripp P, Albers S-V, Mitchell DR. 2020. Propulsive nanomachines: the convergent evolution of archaella, flagella and cilia. FEMS Microbiology Reviews 44:253–304.

11. Miyata M, Robinson RC, Uyeda TQP, Fukumori Y, Fukushima S, Haruta S, Homma M, Inaba K, Ito M,
Kaito C, Kato K, Kenri T, Kinosita Y, Kojima S, Minamino T, Mori H, Nakamura S, Nakane D,
Nakayama K, Nishiyama M, Shibata S, Shimabukuro K, Tamakoshi M, Taoka A, Tashiro Y, Tulum I,
Wada H, Wakabayashi K. 2020. Tree of motility – A proposed history of motility systems in the tree
of life. Genes to Cells 25:6–21.

12. Armitage JP, Berry RM. 2020. Assembly and Dynamics of the Bacterial Flagellum. Annu Rev
Microbiol 74:181–200.

13. Chang Y, Zhang K, Carroll BL, Zhao X, Charon NW, Norris SJ, Motaleb MA, Li C, Liu J. 2020. Molecular
mechanism for rotational switching of the bacterial flagellar motor. 11. Nat Struct Mol Biol
27:1041–1047.

14. Islam MI, Ridone P, Lin A, Michie KA, Matzke NJ, Hochberg G, Baker MAB. 2023. Ancestral
reconstruction of the MotA stator subunit reveals that conserved residues far from the pore are
required to drive flagellar motility. microLife 4:uqad011.

15. Johnson S, Deme JC, Furlong EJ, Caesar JJE, Chevance FFV, Hughes KT, Lea SM. 2024. Structural basis
of directional switching by the bacterial flagellum. Nat Microbiol 1–11.

16. Asai Y, Kawagishi I, Sockett RE, Homma M. 2000. Coupling ion specificity of chimeras between H+-
and Na+-driven motor proteins, MotB and PomB, in Vibrio polar flagella. EMBO J 19:3639–3648.

17. Asai Y, Yakushi T, Kawagishi I, Homma M. 2003. Ion-coupling Determinants of Na+-driven and H+-
driven Flagellar Motors. Journal of Molecular Biology 327:453–463.

18. Kojima S, Takao M, Almira G, Kawahara I, Sakuma M, Homma M, Kojima C, Imada K. 2018. The helix
rearrangement in the periplasmic domain of the flagellar stator B subunit activates peptidoglycan
binding and ion influx. Structure 26:590-598.e5.

19. Miyamura Y, Nishikino T, Koiwa H, Homma M, Kojima S. 2024. Roles of linker region flanked by
transmembrane and peptidoglycan binding region of PomB in energy conversion of the Vibrio
flagellar motor. Genes to Cells n/a.

20. Nishikino T, Iwatsuki H, Mino T, Kojima S, Homma M. 2020. Characterization of PomA periplasmic
loop and sodium ion entering in stator complex of sodium-driven flagellar motor. The Journal of
Biochemistry 167:389–398.

21. Takekawa N, Isumi M, Terashima H, Zhu S, Nishino Y, Sakuma M, Kojima S, Homma M, Imada K.
2019. Structure of Vibrio FliL, a New Stomatin-like Protein That Assists the Bacterial Flagellar
Motor Function. MBio 10.

22. Zhu S, Nishikino T, Takekawa N, Terashima H, Kojima S, Imada K, Homma M, Liu J. 2020. In Situ
Structure of the Vibrio Polar Flagellum Reveals a Distinct Outer Membrane Complex and Its
Specific Interaction with the Stator. Journal of Bacteriology 202:10.1128/jb.00592-19.

23. Hizukuri Y, Morton JF, Yakushi T, Kojima S, Homma M. 2009. The peptidoglycan-binding (PGB)
Domain of the Escherichia coli Pal Protein can also Function as the PGB Domain in E. coli Flagellar
Motor Protein MotB. The Journal of Biochemistry 146:219–229.

24. Islam MI, Lin A, Lai Y-W, Matzke NJ, Baker MAB. 2020. Ancestral Sequence Reconstructions of MotB
Are Proton-Motile and Require MotA for Motility. Front Microbiol 11.

25. Ridone P, Ishida T, Lin A, Humphreys DT, Giannoulatou E, Sowa Y, Baker MAB. 2022. The rapid
evolution of flagellar ion selectivity in experimental populations of E. coli. Science Advances
8:eabq2492.

26. Ridone P, Winter DL, Baker MAB. 2023. Tuning the stator subunit of the flagellar motor with coiled-
coil engineering. Protein Science 32:e4811.

27. Hosking ER, Manson MD. 2008. Clusters of Charged Residues at the C Terminus of MotA and N
Terminus of MotB Are Important for Function of the Escherichia coli Flagellar Motor. Journal of
Bacteriology 190:5517–5521.

28. Zhu S, Homma M, Kojima S. 2012. Intragenic suppressor of a plug deletion nonmotility mutation in
PotB, a chimeric stator protein of sodium-driven flagella. J Bacteriol 194:6728–6735.

29. Ishida T, Ito R, Clark J, Matzke NJ, Sowa Y, Baker MAB. 2019. Sodium-powered stators of the
bacterial flagellar motor can generate torque in the presence of phenamil with mutations near the
peptidoglycan-binding region. Molecular Microbiology 111:1689–1699.

30. Gouridis G, Muthahari YA, de Boer M, Griffith DA, Tsirigotaki A, Tassis K, Zijlstra N, Xu R,
Eleftheriadis N, Sugijo Y, Zacharias M, Dömling A, Karamanou S, Pozidis C, Economou A, Cordes T.
2021. Structural dynamics in the evolution of a bilobed protein scaffold. Proceedings of the
National Academy of Sciences 118:e2026165118.

31. Salinas VH, Ranganathan R. 2018. Coevolution-based inference of amino acid interactions
underlying protein function. Elife 7:e34300.

32. Teşileanu T, Colwell LJ, Leibler S. 2015. Protein sectors: statistical coupling analysis versus
conservation. PLoS Comput Biol 11:e1004091.

33. Ni B, Ghosh B, Paldy FS, Colin R, Heimerl T, Sourjik V. 2017. Evolutionary Remodeling of Bacterial
Motility Checkpoint Control. Cell Reports 18:866–877.

34. Chaban B, Coleman I, Beeby M. 2018. Evolution of higher torque in *Campylobacter*-type bacterial
flagellar motors. Sci Rep 8:97.

35. Chen S, Beeby M, Murphy GE, Leadbetter JR, Hendrixson DR, Briegel A, Li Z, Shi J, Tocheva EI, Müller
A, Dobro MJ, Jensen GJ. 2011. Structural diversity of bacterial flagellar motors. EMBO J 30:2972–
2981.

36. Nord AL, Gachon E, Perez-Carrasco R, Nirody JA, Barducci A, Berry RM, Pedaci F. 2017. Catch bond
drives stator mechanosensitivity in the bacterial flagellar motor. Proc Natl Acad Sci U S A
114:12952–12957.

37. Leptihn S, Castell OK, Cronin B, Lee E-H, Gross LCM, Marshall DP, Thompson JR, Holden M, Wallace
MI. 2013. Constructing droplet interface bilayers from the contact of aqueous droplets in oil. Nat
Protoc 8:1048–1057.

38. Rosholm KR, Baker MAB, Ridone P, Nakayama Y, Rohde PR, Cuello LG, Lee LK, Martinac B. 2017.
Activation of the mechanosensitive ion channel MscL by mechanical stimulation of supported
Droplet-Hydrogel bilayers. Scientific Reports 7:45180.

39. Ni B, Colin R, Link H, Endres RG, Sourjik V. 2020. Growth-rate dependent resource investment in
bacterial motile behavior quantitatively follows potential benefit of chemotaxis. PNAS 117:595–
601.

40. Islam MI, Bae JH, Ishida T, Ridone P, Lin J, Kelso MJ, Sowa Y, Buckley BJ, Baker M a. B. 2021. Novel
Amiloride Derivatives That Inhibit Bacterial Motility across Multiple Strains and Stator Types. J
Bacteriol 203:e0036721.

41. Kinosita Y, Ishida T, Yoshida M, Ito R, Morimoto YV, Goto K, Berry RM, Nishizaka T, Sowa Y. 2020.
Distinct chemotactic behavior in the original Escherichia coli K-12 depending on forward-and-
backward swimming, not on run-tumble movements. Sci Rep 10:15887.

42. Nishiyama M, Kojima S. 2012. Bacterial Motility Measured by a Miniature Chamber for High-
Pressure Microscopy. 7. International Journal of Molecular Sciences 13:9225–9239.

43. Gurung JP, Navvab Kashani M, Agarwal S, Peralta G, Gel M, Baker MAB. 2021. Separation and
enrichment of sodium-motile bacteria using cost-effective microfluidics. Biomicrofluidics
15:034108.

44. Baker MAB, Hynson RMG, Ganuelas LA, Mohammadi NS, Liew CW, Rey AA, Duff AP, Whitten AE,
Jeffries CM, Delalez NJ, Morimoto YV, Stock D, Armitage JP, Turberfield AJ, Namba K, Berry RM,
Lee LK. 2016. Domain-swap polymerization drives the self-assembly of the bacterial flagellar
motor. Nature Structural and Molecular Biology 23:197–203.

45. Castell OK, Dijkman PM, Wiseman DN, Goddard AD. 2018. Single molecule fluorescence for
membrane proteins. Methods 147:221–228.

46. Weatherill EE, Fahie MA, Marshall DP, Andvig RA, Cheetham MR, Chen M, Wallace MI. 2022. Fast
slow folding of an outer membrane porin. Proc Natl Acad Sci U S A 119:e2121487119.

47. Currin A, Parker S, Robinson CJ, Takano E, Scrutton NS, Breitling R. 2021. The evolving art of creating
genetic diversity: From directed evolution to synthetic biology. Biotechnol Adv 50:107762.

48. Liu L, Huang Y, Wang HH. 2023. Fast and efficient template-mediated synthesis of genetic variants.
Nat Methods 20:841–848.

49. Marsischky G, LaBaer J. 2004. Many Paths to Many Clones: A Comparative Look at High-Throughput
Cloning Methods. Genome Res 14:2020–2028.

50. Quaglia D, Ebert MCCJC, Mugford PF, Pelletier JN. 2017. Enzyme engineering: A synthetic biology
approach for more effective library generation and automated high-throughput screening. PLOS
ONE 12:e0171741.

51. Zhou J, Sharp LL, Tang HL, Lloyd SA, Billings S, Braun TF, Blair DF. 1998. Function of protonatable
residues in the flagellar motor of *Escherichia coli*: a critical role for Asp 32 of MotB. J Bacteriol
180:2729–2735.

52. Parkinson JS. 1978. Complementation analysis and deletion mapping of Escherichia coli mutants
defective in chemotaxis. J Bacteriol 135:45–53.

53. Reisch CR, Prather KLJ. 2015. The no-SCAR (Scarless Cas9 Assisted Recombineering) system for
genome editing in Escherichia coli. Sci Rep 5:15096.

54. Higuchi R, Krummel B, Saiki RK. 1988. A general method of in vitro preparation and specific
mutagenesis of DNA fragments: study of protein and DNA interactions. Nucleic Acids Res 16:7351–
7367.

55. Bush SJ, Foster D, Eyre DW, Clark EL, De Maio N, Shaw LP, Stoesser N, Peto TEA, Crook DW, Walker
AS. 2020. Genomic diversity affects the accuracy of bacterial single-nucleotide polymorphism-
calling pipelines. Gigascience 9:giaa007.

56. Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015. StringTie enables
improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol 33:290–295.

57. Seemann T. 2024. tseemann/snippy. Perl.

58. Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J,
Ronneberger O, Bodenstein S, Zielinski M, Bridgland A, Potapenko A, Cowie A, Tunyasuvunakool K,
Jain R, Clancy E, Kohli P, Jumper J, Hassabis D. 2022. Protein complex prediction with AlphaFold-
Multimer. bioRxiv https://doi.org/10.1101/2021.10.04.463034.

59. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B,
Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M,
Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P,
Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. 7873. Nature
596:583–589.

 Figures

Figure 1. Overview of *E. coli* MotAB and ExbBD complexes.

The Bacterial Flagellar Motor and the TonB-dependent siderophore uptake system of *E.coli* are powered,
respectively, by two homologous heptameric systems, the MotA MotB and the ExbB ExbD complexes.
These two transmembrane systems share their general architecture, subunit stoichiometry and functional
features, as both depend on interaction with the peptidoglycan (PG) layer and the presence of an ion-
motive force (IMF) to operate. The MotAB complex interacts with FliG at the cytoplasmic interface of the
basal body of the BFM (left) while the ExbBD complex is thought to interact with the TM domain of TonB
at the plasma membrane.

Figure 2. Isolation of an ExbD/stator chimera able to drive motility .

A) Representative Alphafold2 models of the TM and plug portions of *Ec* MotB (*Escherichia coli*, blue), *Va*
PomB (*Vibrio alginolyticus,* green), *Ec* ExbD (yellow) and chimera PotB. Schematic diagrams of each B
subunit, their general topology (with alpha helices depicted as cylinder and sections lacking a secondary

structure as connnecting lines) and junction site annotations for the previously reported PotB chimera

and candidate PotB/ExbD chimera PotD 3.3. Coloured sections of the diagrams indicate which original

protein the fragment belongs to in the chimeric construct. Numbers flanking each junction site indicate

the residues belonging to each portion of the chimera and are color-coded according to the scheme above.

B) Representative model of the TM and plug portion (N6-T56) of PotD 3.3, coloured according to the

scheme in A. A sequence similarity alignment is shown to highlight the degree of conservation between

the TM domains of ExbD (residues 23-39) and PotB (22-38), and ExbD and MotB (residues 30-46). Letter

intercalating the alignment indicate identity of homologous residues while the '+' sign indicates similarity.

C). The position of suppressor mutation Q104P in PotD 3.3 is mapped (red) on a composite heptameric

(A_5B_2) model of a PomA (grey) /PotD 3.3 (coloured as in B) stator.

Figure 3. Functional characterization of A-B chimeric interfaces.

(A) Section of the stator complex A-B interface viewed from the periplasmic side. Monomers of MotA, shown in different colors, and blue ovals highlight the pore-facing (interior) and lipid-facing (exterior) residues of the complex. Circular diagrams are used to illustrate the intended arrangement of each protein subdomain in the chimeras tested for their ability to rescue motility. (B) Summary of all motility tests performed in stator-deleted *E. coli* co-transformed with plasmids bearing an A-subunit construct (rows) and a B-subunit construct (columns). The background color of each box indicates a motile (green) or non-motile (red) phenotype. Each A-B combination is schematically represented using the symbology shown in A. (C) Single-cell free-swimming cell velocity measurements for motile strains expressing the stator complexes highlighted in green in (B), expressed from plasmid in *E. coli* Δ*motAB.* Cell velocity was measured in sodium-based (85mM NaCl) and potassium-based (85mM KCl) LB medium. (N = 84 (MotA MotB, Na$^+$), 45 (MotA MotB, K$^+$), 85 (MotA PotB, Na$^+$), 41 (MotA PotB, K$^+$) and 77 (PomA PotB, Na$^+$), 24 (MotA PotB, K$^+$) cells). Error bars indicate the standard deviation of the mean. Statistical comparison was performed using Student's T test, ** P < 0.01; *** P <0.001; **** P < 0.0001.

Figure 4. Functional chimeras require sodium for function.

(A) Comparison of motile phenotypes of *E. coli* strains. Stator-deleted Δ*motAB E. coli* harbouring a proton-powered (MotAMotB) or sodium-powered (PomA/PotB) stator genes are shown as motility controls at the top of each square petri dish, each supplemented with 85 mM NaCl (left) or 85 mM KCl (right). (B) Rotation speed of single tethered cells expressing PotDα and PomA variants engineered with other candidate motility-enhancing mutations detected in directed evolution experiments. The first bar on the left (PomA(H116Y)PotDα genomic) indicated the construct being expressed from the chromosomal *motAB* locus, while all other constructs were expressed from a plasmid in a *E. coli* Δ*motAB* background (N = 22, 0, 19, 11, 39 cells, respectively; **** P < 0.0001; n.s.: no significant difference). (C) Alphafold2 model of

511   the chimeric stator PomA(A86V/H116Y)$_5$PotDα_2 with A-subunits in green and B-subunits in yellow. The

512   position of each mutation is highlighted and labelled in red or purple on the structure and in the inset.