Behavior can be decoded across the cortex when individual differences are considered

Johan Nakuci[a*], Jiwon Yeon[a,b], Ji-Hyun Kim[c], Sung-Phil Kim[c] and Dobromir Rahnev[a]

[a]School of Psychology, Georgia Institute of Technology, Atlanta, Georgia, 30332, USA.
[b]Department of Psychology, Stanford University, Stanford, California, 94305, USA.
[c]Department of Biomedical Engineering, Ulsan National Institute of Science and Technology, Ulsan, South Korea.

*Johan Nakuci is corresponding author

Email: jnakuci@gmail.com

Author Contributions:
J.N. and D.R. designed analysis; J.Y. and D.R. designed study; J.Y., J-H.K., and S-P.K. acquired data; J.N. and J.Y. pre-processed data; J.N. analyzed the data; J.N. and D.R. wrote first draft of paper.

Competing Interest Statement: The authors declare no competing interests.

Keywords: brain-behavior relationship, perceptual decision making, fMRI decoding, individual variability

Acknowledgements: This work was supported by the National Institute of Health (award: R01MH119189) and the Office of Naval Research (award: N00014-20-1-2622).

Abstract

Group-level analyses have typically associated behavioral signatures with a constrained set of brain areas. Here we show that two behavioral metrics – reaction time (RT) and confidence – can be decoded across the cortex when each individual is considered separately. Subjects (N=50) completed a perceptual decision-making task with confidence. We built models decoding trial-level RT and confidence separately for each subject using the activation patterns in one brain area at a time after splitting the entire cortex into 200 regions of interest (ROIs). At the group level, we replicated previous results by showing that both RT and confidence could be decoded from a small number of ROIs (12.0% and 3.5%, respectively). Critically, at the level of the individual, both RT and confidence could be decoded from most brain regions even after Bonferroni correction (90.0% and 72.5%, respectively). Surprisingly, we observed that many brain regions exhibited opposite brain-behavior relationships across individuals, such that, for example, higher activations predicted fast RTs in some subjects but slow RTs in others. These results were further replicated in a second dataset. Lastly, we developed a simple test to determine the robustness of decoding performance, which showed that several hundred trials per subject are required for robust decoding. These results show that behavioral signatures can be decoded from a much broader range of cortical areas than previously recognized and suggest the need to study the brain-behavior relationship at both the group and the individual level.

Introduction

An enduring goal of cognitive neuroscience is to establish the relationship between brain and behavior. For example, studies have linked brain measures to healthy aging[1,2], personality[3], intelligence[4,5], and mood[6]. Additionally, there are considerable efforts to use brain measures for diagnosis[7,8], treatment selection [9], and prediction of patient outcomes in clinical settings[10].

What is the right level at which to study the brain-behavior relationship? It is increasingly evident that this relationship is a complex interplay between group-level factors, which are shared among individuals, and individual-level factors that manifest as unique characteristics[11–13]. The discernible variation in brain activity among individuals has long been recognized for its potential to reveal the intricacies of the differences in behavior between individuals[14,15]. Understanding how the individual differs from the group is critical since individual factors may be crucial for diagnosing and treating pathology[16,17]. Consequently, it is imperative to understand the complex individual factors shaping the brain's role in behavior.

Group-level analyses using both mass univariate statistics or multivariate pattern analysis with searchlight have been used to identify brain behavior signatures across the cortex[18,19]. These analyses typically uncovered behavioral signatures within constrained sets of brain areas. However, in the presence of substantial individual variability, one may expect that many areas of the brain that are not predictive of behavior in the group may nonetheless predict behavior in certain individuals. Nevertheless, it remains unknown how widely across cortex one can decode behavioral signatures in individual subjects. However, the problem is partly statistical since one requires both a lot of power (e.g., high number of trials) within an individual and a large enough sample of individuals for individual differences to manifest.

Here we investigate the extent to which two behavioral signatures – reaction time (RT) and confidence – can be decoded from across the cortex. Each subject (N=50) completed 700 trials from a perceptual decision-making task – a number significantly surpassing standard practice. The high number of trials empowers us to robustly estimate the brain-behavior relationship within individuals. To anticipate, we find that when factoring in individual differences, RT and confidence can be decoded from brain activity across the cortex in stark contrast to group-level

analyses. We replicated these results in a second dataset where subjects (N=36) completed 804

trials of a different perceptual decision-making task. These results demonstrate that behavior can

be predicted from a wider set of brain areas than would be suggested by standard group analyses.

Results

<u>Analytical framework for behavior decoding</u>

We investigated the brain-behavior relationship in a perceptual decision-making task with

confidence where each subject (N=50) completed over 700 trials. On each trial, we recorded

subjects' reaction time (RT) and confidence, and investigated how well they can be decoded

from different parts of the brain. We utilized a recently developed method, GLMsingle, to

estimate the voxel-wise activation on a given trial (i.e., single-trial beta)[20]. For each subject, we

performed the decoding analysis on the average activation within each of the 200 regions of

interest (ROIs) from the Schaefer atlas[21]. Specifically, for each trial, we averaged the beta values

from all voxels within a brain region to obtain a single beta value for each ROI. Individual trials

were then randomly separated into training and testing bins (**Fig. 1A**). A linear model was

trained on half of the trials (**Fig. 1B,** *left*) and used to predict behavioral performance on the

remaining trials from the activation (beta values) separately for each ROI (**Fig. 1B,** *middle*).

Decoding performance for a given ROI and subject was determined by correlating the empirical

and predicted RT and confidence (**Fig. 1B,** *right*). The analysis was repeated 25x to ensure that

model performance did not depend on the initial trial split (**Fig. 1C**).

Analytical Framework

Figure 1. Analytical framework for behavior decoding analysis. (A) For each brain region, individual trials were randomly separated into training (red) and testing (green) bins. (B) The trials in the training bin were used to train a linear model (*left*). The model was used to predict the behavioral performance of the trials in the testing bin using the observed brain activation (*middle*). Model performance was estimated using the Pearson correlation between the empirical and predicted behavioral performance (*right*). (C) The analysis was repeated 25x to ensure that model performance was robust and did not depend on the initial trial split.

RT and confidence can be predicted from across the brain

We first performed group-level analyses to decode RT across all subjects for each of the 200

ROIs. To identify brain regions from which RT and confidence could be significantly decoded at

the group-level, the decoding performance values across individual subjects were aggregated and

tested against zero. We found that RT could be significantly decoded in 12.0% of all ROIs after

correcting for multiple comparisons (P < 0.05, Bonferroni-corrected, one-tailed one sample t-

tests; **Fig. S1A**). These standard group-level analyses appear to suggest that RT can be decoded

from only a handful of ROIs in the brain and that the rest of the ROIs cannot be used for

decoding RT.

Critically, we asked whether this conclusion holds on the level of the individual subject. To

address this question, we determined for how many ROIs there was at least one subject for

whom RT could be decoded. We found that with no correction for multiple comparisons,

decoding was significant for at least one subject in every single one of the 200 ROIs (**Fig. 2A,**

Fig. S2A). Next, we examined the decoding accuracy for each ROI separately and applied

Bonferroni correction for the presence of 50 tests (equal to the number of subjects). We found

that RT could be significantly decoded in at least one subject from 90.0% of all ROIs (180 of

200; **Fig. 2A-B**). These ROIs spanned all seven major brain networks associated with the

Schaefer atlas. Further, when using a more stringent Bonferroni correction for 200 tests (equal to

the number of ROIs), RT could be significantly decoded in at least one subject from 78.5% of all

ROIs (157 of 200; **Fig. 2B**). Even with most the stringent Bonferroni correction for 10,000 tests

(the number of subjects times the number of ROIs), RT could be significantly decoded in at least

one subject from 50.5% of all ROIs (101 of 200, **Fig. 2B**).

Figure 2. RT and confidence can be predicted from across the brain in Dataset 1. (A) Brain
regions from which RT could be significantly decoded from in each subject and region of
interest. Significance is shown at four levels of correction: without any correction (P_{uncor}), and
after correcting for 50 (P_{50}), 200 (P_{200}), and 10,000 (P_{10K}) comparisons. NS, not significant.
Colors indicate the most conservative threshold at which one can significantly decode RT from a

given region (see color legend on the right). (B) Percentage of ROIs where RT could be significantly decoded for at least one subject for four levels of multiple comparison correction. (C) Percentage of ROIs per subject where RT could be significantly decoded at four levels of multiple comparison correction. (D) Brain maps plotting percentage of subjects for whom RT could be significantly decoded for four levels of multiple comparison correction. (E-H) Same as Panels A-D but for confidence.

Importantly, even though most ROIs permitted the decoding of RT in at least one subject, this

decoding was only significant in a few subjects per ROI. Specifically, the average ROI could be

used to decode RT in 20.2% without multiple comparison correction, and in 6.6, 4.5, and 1.9%

of subjects after correcting for 50, 200, and 10,000 comparisons (**Fig. 2C**). In all four cases, the

most predictive ROIs were interspersed across much of the cortical surface (**Fig. 2D**). These

results suggest the presence of strong individual differences, such that each ROI is only

predictive of RT in a handful of subjects.

We found similar results when we attempted to decode confidence instead of RT. At the group

level, confidence could be decoded from only 3.5% of ROIs after correcting for 200 comparisons

(one for each ROI; **Fig. S1B**). Nevertheless, at the individual level, many more ROIs could be

used to significantly decode confidence in at least one subject. Specifically, confidence could be

decoded in 99.5% of all ROIs without multiple comparison correction, and in 72.5, 59.5, and

26.5% of ROIs after correcting for 50, 200, and 10,000 comparisons (**Fig 2E-H, Fig. S2B**). Note

that confidence was overall less decodable than RT. Overall, the decoding results demonstrate

that RT and confidence can be decoded from across the cortex when individual differences are

considered.

<u>Opposite brain-behavior relationship among subjects for the same brain region</u>

Critically, for many ROIs, the relationship between brain activity and both RT and confidence

often went in the opposite direction for different subjects. For example, one set of subjects would

exhibit *higher* RT or confidence with higher ROI activation, whereas a different set of subjects

would exhibit *lower* RT or confidence with higher ROI activation (**Fig. 3A-B**). Indeed, we found

that without multiple comparison corrections, high brain activity predicted higher RT in at least

one subject and lower RT in at least one subject in 67.0% of all ROIs (**Fig. 3C**). This percentage

174 decreased to 16.5, 8.0, and 1.0% after correcting for 50, 200, and 10,000 comparisons,

175 respectively. Interestingly, the dorsal attention network contained the most regions with opposite

176 relationship between brain activity and RT among subjects, with the default mode and visual

177 networks as close second and third, respectively (**Fig. 3D**). We found similar results for

178 confidence with 75.5% of all ROIs showing significant decoding in different direction for at least

179 two subjects, and 15.0, 7.5, and 0% after correcting for 50, 200, and 10,000 comparisons,

180 respectively (**Fig. 3E-G**). Further, the default mode network contained the most regions with

181 opposite relationship between brain activity and confidence among subjects (**Fig. 3H**). In

182 contrast to the findings for RT that lacked large clusters where decoding was consistently in the

183 same direction, we found that a subset of ROIs associated with somatomotor network exhibited

184 consistent relationship between brain activity and confidence across subjects. Except for this

185 relatively small cluster, our results demonstrate that the relationship between brain activity and

186 behavioral outcomes is not universal and that it frequently goes in opposite directions for

187 different subjects.

Figure 3. Opposite relationship between brain activity and both RT and confidence among subjects for the same brain region in Dataset 1. (A) Beta values for each subject and region of interest. Red and blue dots reflect subjects for whom RT could be significantly decoded from a given ROI (P < 0.05, uncorrected for multiple comparison). Gray dots show subjects for whom RT could not be significantly decoded from a given ROI. (B) Brain maps showing the ROIs that contained at least one subject with both positive and negative beta values for RT. Significance is shown at four levels of correction: without any correction (P_{uncor}), and after correcting for 50 (P_{50}), 200 (P_{200}), and 10,000 (P_{10K}) comparisons. (C) Percentage of ROIs that contained at least one subject with both positive and negative beta values for RT for each for four levels of multiple comparison correction. (D) Percentage of ROIs within each brain network that that contained at least one subject with both positive and negative beta values for RT after correcting for 50 comparisons. (E-J) Same as Panels A-D but for confidence. FPN, Frontal Parietal Network; DMN, Default Mode Network; DAN, Dorsal Attention Network; LIM, Limbic Network; VAN, Ventral Attention Network; SOM, Somatomotor Network; VIS, Visual Network.

<u>RT and confidence can be decoded from across the brain in Dataset 2</u>

We replicated these results in a second dataset where subjects completed a different perceptual

decision-making task with confidence[22]. Subjects (N=36) completed 804 trials but only half of

them included confidence ratings, thus substantially decreasing the power for the confidence

analyses.

Similar to Dataset 1, we found that at the group-level, RT could be significantly decoded in

13.0% of all ROIs after correcting for 200 comparisons ($P < 0.05$, Bonferroni-corrected, one-

tailed one sample t-tests; **Fig. S3A**). Crucially, we determined for how many ROIs there was at

least one subject for whom RT could be decoded. At the individual level, RT could be decoded

in 100% of all ROIs without multiple comparison correction, and in 87.0, 72, and 43.5% of ROIs

after correcting for 36 (number of subjects), 200 (number of ROIs), and 7,200 (subjects times

ROIs) comparisons (**Fig 4A-D, Fig. S4A**). Given the lower number of trials that contained

confidence, we found that confidence could be significantly decoded in only 1.5% of all ROIs at

the group-level after correcting for multiple comparisons ($P < 0.05$, Bonferroni-corrected, one-

tailed one sample t-tests; **Fig. S3B**). Despite the lower power, at the individual level, confidence

could still be decoded in 76.5 % of all ROIs without multiple comparison correction, and in 47.0,

32.5, and 22.5% of ROIs after correcting for 36, 200, and 7,200 comparisons (**Fig 4E-H, Fig.**

S4B). Overall, the results from Dataset 2 further support the notion that both the RT and

confidence can be decoded from across the cortex when individual differences are considered.

Figure 4. RT and confidence can be decoded from across the brain in Dataset 2. (A) Brain
regions from which RT could be significantly decoded from in each subject and region of
interest. Significance is shown at four levels of correction: without any correction (P_{uncor}), and
after correcting for 36 (P_{36}), 200 (P_{200}), and 7,200 ($P_{7.2K}$) comparisons. NS, not significant.
Colors indicate the most conservative threshold at which one can significantly decode RT from a

given region (see color legend on the right). (B) Percentage of ROIs where RT could be
significantly decoded for at least one subject for four levels of multiple comparison correction.
(C) Percentage of ROIs per subject where RT could be significantly decoded at four levels of
multiple comparison correction. (D) Brain maps plotting percentage of subjects for whom RT
could be significantly decoded for four levels of multiple comparison correction. (E-H) Same as
Panels A-D but for confidence. Note that the confidence findings show similar pattern as in
Dataset 1 despite being based on half the number of trials per subject.
239

<u>Opposite brain-behavior relationship among subjects for the same brain region in Dataset 2</u>

In Dataset 1, we found that for many ROIs, the relationship between brain activity and both RT

and confidence often went in the opposite direction for different subjects. Here we show that the

same effects also occur in Dataset 2. Similar to Dataset 1, we found that without multiple

comparison corrections, high brain activity predicted higher RT in at least one subject and lower

RT in at least one subject in 50.0% of all ROIs and 16.5, 6.5, 0.5% after correcting for 36, 200

and 7,200 comparisons, respectively (**Fig. 5A-D**). Reflecting the lower number of trials per

subject, confidence showed weaker effects with 12.0% of all ROIs showing significant decoding

in different direction for at least two subjects, and 3.0, 1.5, and 0% after correcting for 36, 200,

and 7,200 comparisons, respectively (**Fig. 5E-H**). Interestingly, differences emerged between

Dataset 1 and 2 when comparing which network contained the most regions with opposite

relationship between brain activity. These results demonstrate that the relationship between brain

activity and behavioral outcomes is not universal, and that this opposite relationship is not

limited to a specific set of regions but instead depends on the individual subject.

Figure 5. Opposite relationship between brain activity and both RT and confidence among subjects for the same brain region in Dataset 2. (A) Beta values for each subject and region of interest. Red and blue dots reflect ROIs from which RT could be significantly decoded (P < 0.05, uncorrected for multiple comparison). Gray dots show ROIs from which RT could not be significantly decoded from. (B) Brain maps showing the ROIs that contained at least one subject with both positive and negative beta values for RT. Significance is shown at four levels of correction: without any correction (P_uncor), and after correcting for 36 (P_36), 200 (P_200), and 7,200 (P_7.2K) comparisons. (C) Percentage of ROIs that contained at least one subject with both positive and negative beta values for RT for each for four levels of multiple comparison correction. (D) Percentage of ROIs within each brain network that that contained at least one subject with both positive and negative beta values for RT after correcting for 50 comparisons. (E-J) Same as Panels A-D but for confidence. Note that the confidence findings show similar pattern as in Dataset 2 despite being based on half of the number of trials per subject. FPN, Frontal Parietal Network; DMN, Default Mode Network; DAN, Dorsal Attention Network; LIM, Limbic Network; VAN, Ventral Attention Network; SOM, Somatomotor Network; VIS, Visual Network.

<u>No relationship between decoding performance and frame displacement</u>

To confirm that these results were not due to motion artifacts, we determined if there was an association between decoding performance and frame displacement (FD). We performed a regression analysis where average FD for a subject was used to predict average decoding performance for that subject. We found no significant association between FD and decoding performance in either Dataset 1 (RT: R^2 = 0.02, P = 0.34; Conf: R^2 = 0.002, P = 0.72; **Fig. S5A-B**) or Dataset 2 (RT: R^2 = 0.05, P = 0.20; Conf: R^2 = 0.04, P = 0.22; **Fig. S5C-D**). These results suggest that decoding performance was not driven by motion artifacts.

<u>Robust decoding performance requires several hundred trials per subject</u>

Our results show that RT and confidence can be decoded across the cortex. Our Datasets 1 and 2 included 700 and 804 trials per subject, respectively, which likely contributed to the high decoding we observed at the individual level. Here we use our data to ask how the number of trials used for training and testing affects decoding performance. We develop a simple test to determine how many trials are needed to obtain robust individual-level decoding.

The test relies on estimating decoding performance using a range of trials for both training and testing. For robustness, we used the data from all 200 ROIs to decode both RT and confidence. We trained the linear regression model on subsets of trials that ranged from 5% to 95% of all trials and tested on the remaining trials. For each percentage of trials used, we randomly selected trials for training and testing 25 times. We then computed both the average decoding performance across the 25 iterations and the variance in decoding performance. We then determined the percentage of that should be devoted to training the decoding model to minimize the variance in the decoding performance.

For Dataset 1, minimum decoding variance was obtained when training/testing each contained 350 of trials (50% of trials) for both RT (**Fig. 6A**) and confidence (**Fig. 6B**), while average decoding performance increased with higher number of training trials. Critically, using fewer than 350 trials for training exhibited poor decoding performance. On the other hand, using more

than 350 trials for training improved performance, but the variance in performance increased

across iterations. The maximum difference between decoding performance and variance was

obtained when training was performed on 525 trials (75% of all trials). We found similar results

for Dataset 2. Minimum decoding variance was obtained when training contained 402 trials

(50% of trials) for RT (**Fig. 6C**) and 301 trials (75% of trials) for confidence (**Fig. 6D**), while

decoding performance increased with higher number of training trials. These results suggest the

need of several hundred trials for robustly training a decoding model for RT or confidence,

implying that many previous studies estimating brain-behavior relationship at the level of the

individual may be underpowered.

313
314
Figure 6. Robust decoding of RT and confidence requires several hundred trials. In Dataset
1, the variance in decoding performance was minimized when using 350 trials (corresponding to
50% of all trials) for both (A) RT and (B) confidence. The black dots present the average
decoding performance across subject after 25 repeats for each subject (*left axis)*. The red dots
present the average subject-level decoding variance across the 25 repeats for each subject (*right*
axis). The green boxes indicate the number of trials that minimize the variance in the decoding
performance. The orange boxes indicate the number of trials that maximize the difference
between decoding performance and the decoding variance. Error bars show SEM. (C-D) Same as
Panels A-B but for Dataset 2. Note that in Dataset 2, confidence was measured on only have the
trials (402) and correspondingly the decoding variance was minimized for a higher percentage of
trials (75%).

Discussion

Traditionally, the brain-behavior relationship has been examined at the group-level to identify the commonalities among individuals. Group-level analyses have typically associated behavioral signatures with a constrained set of brain areas. Here, in contrast to the traditional approach, we focus on the brain-behavior relationship within an individual. We tested how well trial-level RT and confidence can be decoded from the activation in each of the 200 cortical regions of interest obtained using the Schaefer atlas. We showed that RT and confidence can be significantly decoded for at least one subject from brain activity across most of the cortex. These results demonstrate that behavior can be predicted from a wider set of brain areas that would be suggested by standard group analyses.

These findings suggest that individual variability is a major factor in the ability to decode behavioral signatures. In fact, we found that most brain regions could predict RT and confidence in only a small percentage of subjects. Therefore, an analysis focused on subject-level might be more sensitive since group-level analyses aggregate over the large degree of within-subject variability[17,23].

Individual variability in brain-behavior relationship may have many sources. Such variation could emerge from functional degeneracy, the ability of different brain regions to perform the same computation[24–26]. Functional degeneracy could arise from differences in structural connectivity among individuals[27,28], since structural connectivity has been associated with differences in behavioral performance[29].

Despite the heterogeneity among individuals, a few brain regions showed relatively high consistency across subjects. The lateral frontal cortex and the visual cortex were the most consistent regions from which RT could be decoded. Similarly, for confidence, the somatomotor system was the most consistent area across subjects from which confidence could be decoded, suggesting an association between action-related brain signals and confidence[30,31].

In a seminal paper, Marek *et al.*[32] showed that thousands of subjects are required to robustly

estimate the brain-behavior relationship. Our results suggest that beyond the number of subjects,

an arguably even more important factor for brain-behavior relationships is the high degree of

between-subject variability. Specifically, if one brain-region is predictive of behavior in one

direction in some people but in the opposite direction in others, then no matter how many people

a model is trained on, it will always fail to capture many of the individual subjects. It has been

suggested that beta values cannot be interpreted particularly from non-linear decoding models

using multiple voxels because the beta-values may indicate reduction in noise[33]. This may well

be true in some cases, but in the cases where opposite beta values are meaningfully different

between individuals, individual variability becomes a crucial limitation in our understanding of

how behavior arises from brain activity. Specifically, most brain-behavior computational models

assume that there is a uniform relationship between a given region and behavior across all

individuals, but this may simply not be true. Instead, we may need to build models that are more

flexible and can account for opposite beta values for different subjects.

Finally, we developed a simple test that future studies can use to determine the optimal number

of trials in the training and testing bins. This is a crucial step in all decoding analyses, but one

that has received little attention. The test relies on training the model on a subset of trials that

ranges from 5 to 95% of all trials, with testing performed on the remaining trials. In the context

of the current study, we found that several hundred (~300) trials per person are needed to

robustly decode the brain-behavior relationship at the individual level. These results suggest that

many previous studies estimating brain-behavior relationship at the individual level may be

underpowered. We suggest that the optimal number of trials used in training a decoding model

could be based on: (1) minimizing the variance in the decoding performance; or (2) maximizing

the difference between decoding performance and the variance in the performance. In the current

study, we opted to minimize the variance in the decoding performance even though it came at the

cost of a lower decoding performance.

In conclusion, our findings show that behavioral signatures can be decoded from a much broader

range of cortical areas than previously recognized. These results highlight that studying the

brain-behavior relationship should be studied at both the group and the individual level.

Methods

<u>Dataset 1 subjects and task</u>

Dataset 1 has been previously published[34]. All details can be found in the original publication.

Briefly, subjects (N= 50; 25 females; mean age = 26; age range = 19-40) completed a task where

they judged which set of colored dots (red vs. blue) is more frequent in a cloud of dots. The

stimulus was presented for 500 ms and subjects made untimed decision and confidence decisions

using separate button presses. Most subjects performed six runs of 128 trials (a total of 768

trials). Three subjects completed only half of the 6th run and another three subjects completed

only the first 5 runs due to time constraints. All subjects were screened for neurological disorders

and MRI contraindications. The study was approved by Ulsan National Institute of Science and

Technology Review Board and all subjects gave written consent.

<u>Dataset 1 MRI recording</u>

The MRI data was collected on a 64-channel head coil 3T MRI system (Magnetom Prisma;

Siemens). Whole-brain functional data were acquired using a T2*-weighted multi-band

accelerated imaging (FoV = 200 mm; TR = 2000 ms; TE = 35 ms; multiband acceleration factor

= 3; in-plane acceleration factor = 2; 72 interleaved slices; flip angle = 90°; voxel size = 2.0 x 2.0

x 2.0 mm^3). High-resolution anatomical MP-RAGE data were acquired using T1-weighted

imaging (FoV = 256 mm; TR = 2300 ms; TE = 2.28 ms; 192 slices; flip angle = 8°; voxel size =

1.0 x 1.0 x 1.0 mm^3).

<u>Dataset 2 subjects and task</u>

Dataset 2 has been previously published as Experiment 2 in (Yeon et al., 2020). All details can

be found in the original publication. Briefly, subjects (N = 36, 23 females; mean age = 21.5; age

range = 18-28) completed a task where they indicated whether a moving-dots stimulus had an

overall coherent motion (always in downward direction) or not. The stimulus was presented for

500 ms and subjects made untimed decision immediately after the stimulus. All subjects

completed 6 runs of 144 trials (a total of 864 trials). In the first half of the experiment (runs 1-3),

subjects performed the task without providing confidence ratings. In the second half of the

experiment (runs 4-6), subjects reported their confidence level with a separate, untimed button

press immediately after making their perceptual decision. All subjects were screened for

neurological disorders and MRI contraindications. The study was approved by the Georgia Tech

Institutional Review Board and all subjects gave written consent.

<u>Dataset 2 MRI recording</u>

The MRI data were collected on 3 T MRI systems using a 32-channel head coils. Anatomical

images were acquired using T1-weighted sequences with a MEMPRAGE sequence,

FoV $=$ 256 mm; TR $=$ 2530 ms; TE $=$ 1.69 ms; 176 slices; flip angle $=$ 7°; voxel

size $=$ 1.0 $\times$ 1.0 $\times$ 1.0 mm^3. Functional images were acquired using T2*-weighted

gradient echo-planar imaging sequences (FoV $=$ 220 mm; TR $=$ 1200 ms; TE $=$ 30 ms;

51 slices; flip angle $=$ 65°; voxel size $=$ 2.5 $\times$ 2.5 $\times$ 2.5 mm^3).

<u>MRI preprocessing</u>

MRI data were preprocessed with SPM12 (Wellcome Department of Imaging Neuroscience,

London, UK). All images were first converted from DICOM to NIFTI and removed the first

three volumes to allow for scanner equilibration. We then preprocessed with the following steps:

de-spiking, slice-timing correction, realignment, segmentation, coregistration, normalization, and

spatial smoothing with 10 mm full width half maximum. Despiking was done using the

3dDespike function in AFNI. The preprocessing of the T1-weighted structural images involved

skull-removal, normalization into MNI anatomical standard space, and segmentation into gray

matter, white matter, and cerebral spinal fluid, soft tissues, and air and background.

<u>Single-trial beta estimation</u>

Single-trial beta responses were estimated with a general linear model (GLM) using GLMsingle,

a Matlab toolbox for single-trial analyses[20]. The hemodynamic response function was estimated

for each voxel and nuisance regressors were estimated in the same manner as described in Allen

et al. [35]. Additionally, regressors for the global signal and for six motion parameters (three

translation and three rotation) were included. The single-trial betas were estimated in three

batches. In each batch, the betas for every third trial were estimated because the trials in our

study were temporally close together. Also, trials that were within 20 seconds from the end of

run were removed due to the lack of sufficient length of signal from which to estimate the trial-

specific hemodynamic response function. Further, to reduce any differences in the modeling

training and fitting that might arise from difference in the number of trials between subject, when

possible, we opted for a uniform number of trials per subject. In total, the analysis on was based

on 700 trials for Dataset 1 and 804 trials for Dataset 2; for subjects who had more trials than

these, we simply removed trials from the end of the experiment. (Note that in six subjects in

Dataset 1 and two subjects in Dataset 2 had fewer total trials because they did not complete all 6

runs.)

<u>Subject-level brain-behavior decoding analysis</u>

For each trial, the activation within each of the 200 regions of interest (ROIs) from the Schaefer

atlas was estimated by averaging all voxel in the ROI[21]. Individual trials were randomly

separated into training and testing bins each containing 350 trials in Dataset 1 and 402 trials in

Dataset 2. (For the confidence analyses in Dataset 2, the training was done on 301 trials and

testing on the remaining 101 trials.) For each ROI, a linear model was trained on training bin and

used to predict behavioral performance on the testing bin. The linear model was trained using

fitlm.m in Matlab.

Decoding performance for a given ROI was determined by correlating the empirical and

predicted RT and confidence. The analysis was repeated 25x to ensure that model performance

was not dependent on the initial division of trials. The decoding performance values in each of

the 25 iterations were first z-scored using the Fisher transformation and then averaged and

converted back to r-values. Significance was determined by converting r-values to t-values.

Decoding analysis was conducted for each individual subject separately. In Dataset 1, subject-

level decoding performance was evaluated at $P < 0.05$ uncorrected and with four Bonferroni

multiple comparison corrections of 50 tests (equal to the number of subjects), 200 tests (equal to

the number of ROIs), and 10,000 tests (the number of subjects times the number of ROIs).

Similarly, in Dataset 2, subject-level decoding performance was evaluated at $P < 0.05$

uncorrected and with four Bonferroni multiple comparison corrections of 36 tests (equal to the

number of subjects), 200 tests (equal to the number of ROIs), and 7,200 tests (the number of

subjects times the number of ROIs). Brain images were generated with BrainSpace toolbox at

https://github.com/MICA-MNI/BrainSpace.git[36].

Group-level brain-behavior decoding analysis

Group-level decoding performance was estimated by averaging the individual subject decoding performance for each ROI. Group-level decoding was Bonferroni corrected for 200 tests (equal to the number of ROIs).

Decoding performance and trial number

We developed a test to determine how many trials are needed to obtain robust individual-level brain-behavior relationships. The test relies on estimating decoding performance on a range of trials in training and testing bins. Specifically, for each subject, a whole brain multilinear regression model was trained on a subset of trials that ranged from 5% to 95% of trials and tested on the remaining trials. The multilinear regression model was repeated 25 times training on different subset of data. Decoding performance was estimated by averaging across the 25 iterations and decoding performance variance was estimated by calculating the standard deviation across the 25 iterations.

Data and code

Data and scripts necessary to perform the analysis are available at https://osf.io/dc9pa/.

References

1. Geerligs, L., Renken, R. J., Saliasi, E., Maurits, N. M. & Lorist, M. M. A Brain-Wide Study of Age-Related Changes in Functional Connectivity. *Cerebral Cortex* **25**, 1987–1999 (2015).

2. Dosenbach, N. U. F. *et al.* Prediction of Individual Brain Maturity Using fMRI. *Science (1979)* **329**, 1358–1361 (2010).

3. Yarkoni, T. Neurobiological substrates of personality: A critical overview. in *APA handbook of personality and social psychology, Volume 4: Personality processes and individual differences.* 61–83 (American Psychological Association, Washington, DC, US, 2015). doi:10.1037/14343-003.

4. Heuvel, M. P. van den, Stam, C. J., Kahn, R. S. & Pol, H. E. H. Efficiency of Functional Brain Networks and Intellectual Performance. *The Journal of Neuroscience* **29**, 7619 (2009).

5. Finn, E. S. *et al.* Functional connectome fingerprinting: Identifying individuals based on patterns of brain connectivity HHS Public Access. *Nat Neurosci* **18**, 1664–1671 (2015).

6. Smith, S. M. *et al.* A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nat Neurosci* **18**, 1565–1567 (2015).

7. Arbabshirani, M., Kiehl, K., Pearlson, G. & Calhoun, V. Classification of schizophrenia patients based on resting-state functional network connectivity. *Front Neurosci* **7**, (2013).

8. Yahata, N. *et al.* A small number of abnormal brain connections predicts adult autism spectrum disorder. *Nat Commun* **7**, 11254 (2016).

9. Williams, L. M. *et al.* Amygdala Reactivity to Emotional Faces in the Prediction of General and Medication-Specific Responses to Antidepressant Treatment in the Randomized iSPOT-D Trial. *Neuropsychopharmacology* **40**, 2398–2408 (2015).

10. Whelan, R. *et al.* Neuropsychosocial profiles of current and future adolescent alcohol misusers. *Nature* **512**, 185–189 (2014).

11. Nakuci, J. *et al.* Quantifying the contribution of subject and group factors in brain activation. *Cerebral Cortex* bhad348 (2023) doi:10.1093/cercor/bhad348.

12. Gratton, C. *et al.* Functional Brain Networks Are Dominated by Stable Group and Individual Factors, Not Cognitive or Daily Variation. *Neuron* **98**, 439–452 (2018).

13. Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI. *Trends in Cognitive Sciences* Preprint at https://doi.org/10.1016/j.tics.2016.03.014 (2016).

14. van Horn, J. D., Grafton, S. T. & Miller, M. B. Individual variability in brain activity: A nuisance or an opportunity? *Brain Imaging Behav* (2008) doi:10.1007/s11682-008-9049-9.

15. Miller, M. B., Donovan, C. L., Bennett, C. M., Aminoff, E. M. & Mayer, R. E. Individual differences in cognitive style and strategy predict similarities in the patterns of brain activity between individuals. *NeuroImage* Preprint at https://doi.org/10.1016/j.neuroimage.2011.05.060 (2012).

16. Gratton, C. *et al.* Defining Individual-Specific Functional Neuroanatomy for Precision Psychiatry. *Biological Psychiatry* vol. 88 28–39 Preprint at (2020).

17. Lebreton, M., Bavard, S., Daunizeau, J. & Palminteri, S. Assessing inter-individual differences with
task-related functional neuroimaging. *Nature Human Behaviour* vol. 3 897–905 Preprint at
(2019).

18. Friston, K. J., Holmes, A. P., Price, C. J., Büchel, C. & Worsley, K. J. Multisubject fMRI Studies and
Conjunction Analyses. *Neuroimage* **10**, 385–396 (1999).

19. Kriegeskorte, N., Goebel, R. & Bandettini, P. Information-based functional brain mapping.
Proceedings of the National Academy of Sciences **103**, 3863–3868 (2006).

20. Prince, J. S. *et al.* Improving the accuracy of single-trial fMRI response estimates using GLMsingle.
Elife **11**, e77599 (2022).

21. Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic
Functional Connectivity MRI. *Cerebral Cortex* **28**, 3095–3114 (2018).

22. Yeon, J., Shekhar, M. & Rahnev, D. Overlapping and unique neural circuits are activated during
perceptual decision making and confidence. *Sci Rep* **10**, (2020).

23. Fisher, A. J., Medaglia, J. D. & Jeronimus, B. F. Lack of group-to-individual generalizability is a
threat to human subjects research. *Proc Natl Acad Sci U S A* **115**, 6106–6115 (2018).

24. Tononi, G., Sporns, O. & Edelman, G. M. Measures of degeneracy and redundancy in biological
networks. *Proc Natl Acad Sci U S A* (1999) doi:10.1073/pnas.96.6.3257.

25. Price, C. J. & Friston, K. J. Degeneracy and cognitive anatomy. *Trends Cogn Sci* (2002)
doi:10.1016/S1364-6613(02)01976-9.

26. Sajid, N., Parr, T., Hope, T. M., Price, C. J. & Friston, K. J. Degeneracy and Redundancy in Active
Inference. *Cerebral Cortex* (2020) doi:10.1093/cercor/bhaa148.

27. Bansal, K., Nakuci, J. & Muldoon, S. F. Personalized brain network models for assessing structure-
function relationships. *Curr Opin Neurobiol* **52**, 1–13 (2018).

28. Muldoon, S. F. *et al.* Stimulation-Based Control of Dynamic Brain Networks. *PLoS Comput Biol*
(2016) doi:10.1371/journal.pcbi.1005076.

29. Kanai, R. & Rees, G. The structural basis of inter-individual differences in human behaviour and
cognition. *Nature Reviews Neuroscience* Preprint at https://doi.org/10.1038/nrn3000 (2011).

30. Fleming, S. M. *et al.* Action-Specific Disruption of Perceptual Confidence. *Psychol Sci* (2015)
doi:10.1177/0956797614557697.

31. Kiani, R. & Shadlen, M. N. Representation of confidence associated with a decision by neurons in
the parietal cortex. *Science (1979)* (2009) doi:10.1126/science.1169405.

32. Marek, S. *et al.* Reproducible brain-wide association studies require thousands of individuals.
Nature **603**, 654–660 (2022).

33. Kriegeskorte, N. & Bandettini, P. Analyzing for information, not activation, to exploit high-
resolution fMRI. *Neuroimage* **38**, 649–662 (2007).

34.   Nakuci, J. *et al.* Quantifying the idiosyncrasy in brain activations. *bioRxiv* 2022.08.01.502338
572   (2023) doi:10.1101/2022.08.01.502338.

573   35.   Allen, E. J. *et al.* A massive 7T fMRI dataset to bridge cognitive neuroscience and artificial
574   intelligence. *Nat Neurosci* **25**, 116–126 (2022).

575   36.   Vos de Wael, R. *et al.* BrainSpace: a toolbox for the analysis of macroscale gradients in
576   neuroimaging and connectomics datasets. *Commun Biol* **3**, 103 (2020).