1 Full title: **Tracking inflammation resolution signatures in lungs after SARS-CoV-2 omicron**
2 **BA.1 infection of K18-hACE2 mice**
3
4 Short title: Inflammation resolution in lungs of a mouse model of SARS-CoV-2 infection
5

6 Agnes Carolin[1#], Kexin Yan[1#], Cameron R. Bishop[1#], Bing Tang[1], Wilson Nguyen[1], Daniel J.
7 Rawle[1†], Andreas Suhrbier[1,2†].
8
9 [1] QIMR Berghofer Medical Research Institute, Brisbane, Queensland, 4029, Australia.

10 [2] GVN Centre of Excellence, Australian Infectious Disease Research Centre, Brisbane,
11 Queensland 4029 and 4072, Australia.

12
13 [#] should be considered joint first authors
14 [†] should be considered joint last authors
15
16 [*]Corresponding author. Email: Andreas. Suhrbier@qimrberghofer.edu.au or
17 Daniel.Rawle@qimrberghofer.edu
18
19
20
Keywords*: SARS-CoV-2, omicron BA.1, K18-hACE2 mice, protective inflammation,
inflammation resolution

Data Availability Statement: All data is provided in the manuscript and accompanying
supplementary files. Raw sequencing data (fastq files) generated for this publication for RNA-
Seq have been deposited in the NCBI SRA, BioProject: PRJNA1081927 and are publicly
available.
28

29 **Funding:** The authors thank the Brazil Family Foundation (and others) for their generous
30 philanthropic donations that helped set up the PC3 (BSL3) SARS-CoV-2 research facility at
31 QIMR Berghofer MRI, as well as ongoing research into SARS-CoV-2, COVID-19 and long-
32 COVID. A.S. is supported by the National Health and Medical Research Council (NHMRC) of
33 Australia (Investigator grant APP1173880).
34

35 **Competing interests:** The authors declare that they have no known competing interests.

Abstract

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) causes Coronavirus Disease 2019 (COVID-19), which can result in severe disease often characterised by a 'cytokine storm' and the associated acute respiratory distress syndrome. However, many infections with SARS-CoV-2 are mild or asymptomatic throughout the course of infection. Although blood biomarkers of severe disease are well studied, less well understood are the inflammatory signatures in lung tissues associated with mild disease or silent infections, wherein infection and inflammation are rapidly resolved leading to sequelae-free recovery. Herein we described RNA-Seq and histological analyses of lungs over time in an omicron BA.1/K18-hACE2 mouse infection model, which displays these latter features. Although robust infection was evident at 2 days post infection (dpi), viral RNA was largely cleared by 10 dpi. Acute inflammatory signatures showed a slightly different pattern of cytokine signatures compared with severe infection models, but where much diminished 30 dpi and absent by 66 dpi. Cellular deconvolution identified significantly increased abundance scores for a number of anti-inflammatory pro-resolution cell types at 5/10 dpi. These included type II innate lymphoid cells, T regulatory cells, and interstitial macrophages. Genes whose expression trended downwards over 2 – 66 dpi included biomarkers of severe disease and were associated with 'cytokine storm' pathways. Genes whose expression trended upward during this period were associated with recovery of ciliated cells, AT2 to AT1 transition, reticular fibroblasts and innate lymphoid cells, indicating a return to homeostasis. Very few differentially expressed host genes were identified at 66 dpi, suggesting near complete recovery. The parallels between mild or subclinical infections in humans and those observed in this BA.1/K18-hACE2 mouse model are discussed.

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused a global pandemic of Coronavirus Disease 2019 (COVID-19), with over 770 million cases and 7 million deaths [1, 2]. Severe acute disease is primarily characterised by infection of the respiratory tract and pathological hyper-inflammatory responses, often referred to as a "cytokine storm" that can lead to acute respiratory distress syndrome (ARDS) and ultimately mortality [3-5]. Post-acute and chronic sequelae of COVID-19 are now well described, with post-COVID conditions affecting 10-30% of COVID-19 patients [6] and long-COVID at least 10% of patients [7-9].

The inflammatory mediators that are associated with severe COVID-19, as distinct from mild disease, have been the focus of many studies, with such insights having the potential to identify therapeutic anti-disease interventions [3, 5, 10-17]. Elevated mRNA or protein levels of a range of cytokines have been identified in peripheral blood of severe COVID-19 patients, these include *inter alia* IL-6, IL-1, IL-2, IL-17, IL-18, TNF, IFNG and CSF2 (GM-CSF) (*ibid*). Elevated IL-10 in the peripheral blood has also been associated with severe disease, but may be useful later in the infection to suppress the hyper-cytokinemia associated with ARDS [18-20]. Elevated Th2 [10, 21-23] and Th17 responses [10, 24] have been associated with poor outcomes, with Th1/Th2 [25-27] and Th1/Th17 imbalances [28] also reported to be detrimental. T regulatory cells (Tregs) are viewed as providing protection against the cytokine storm [29], whereas neutrophils promote the latter and contribute to disease severity [10]. Monocytes and macrophages play key protective roles against SARS-CoV-2 infection, but depending on their differentiation and activation state, are also important contributors to over-production of inflammatory cytokines [30, 31]. Higher viral loads appear to positively associate with COVID-

19 severity [32, 33], although this may be associated with more prolonged high viral loads in

patients with severe disease [34].

An estimated 25-30% of SARS-CoV-2 infections are asymptomatic throughout the course of

infection [35, 36], with a constellation of factors potentially contributing [37, 38]. Although fatal

disease is associated with higher viral loads, asymptomatic individuals often do not have lower

acute viral loads than symptomatic patients [39-42]. Humans have also been shown to tolerate a

certain level of lower respiratory tract infection (and associated radiographic abnormalities),

without reporting symptoms [43]. Persistent asymptomatics do, nevertheless, appear able to

clear the virus more rapidly [44, 45]. For instance, only minimal PCR-detectable viral RNA was

present in saliva of a small cohort of such individuals by days 9-10 post first positive diagnosis

(after daily saliva testing) [44]. Individuals with silent infections may have lower levels of

circulating proinflammatory cytokines [45], although this is not a universal finding [46]. Silent

or subclinical infections have been associated with powerful antiviral responses [47], early and

robust innate immune responses [48], as well as specific gene expression patterns [49, 50], such

as higher plasma levels of the anti-inflammatory cytokines IL-10, IL-1RA and IL-19 [46].

Although in humans there can be long-term consequences of asymptomatic SARS-CoV-2

infections, these are significantly more prevalent after symptomatic infections [51].

Herein we provide the transcriptional profile of lung infection over time in a model of SARS-

CoV-2 omicron BA.1 infection of K18-hACE2 mice, wherein the majority of mice showed no

significant overt signs or symptoms of disease [52]. The acute phase was characterised by a

significant lung infection and a pattern of proinflammatory cytokine signatures differing only

slightly from those reported previously [53]. By 10 days post infection (dpi), virus had been

largely cleared and a series of cell types associated with damage control, suppression of

inflammation, homeostasis and repair were identified by cellular deconvolution.   Few

differentially expressed genes (DEGs) were identified in lungs by 66 dpi, arguing that the model describes near complete resolution of the infection, without long-term pulmonary sequelae. The term "protective inflammation" has been coined to describe the inflammatory processes that lead to subclinical or mild disease, resulting in sequelae-free outcomes [54]. Herein we provide insights into the transcriptional signatures in lungs that characterise resolution of inflammation and return to homeostasis, and also identify a series of parallels with human studies.

Materials and methods

Ethics statements and regulatory compliance

Collection of nasal swabs from consented COVID-19 patients was approved by the QIMR Berghofer Medical Research Institute Human Research Ethics Committee (P3600).

All mouse work was conducted in accordance with the Australian code for the care and use of animals for scientific purposes (National Health and Medical Research Council, Australia). Mouse work was approved by the QIMR Berghofer MRI Animal Ethics Committee (P3600). All infectious SARS-CoV-2 work was conducted in the BioSafety Level 3 (PC3) facility at the QIMR Berghofer MRI (Department of Agriculture, Fisheries and Forestry, certification Q2326 and Office of the Gene Technology Regulator certification 3445). Breeding and use of GM mice was approved under a Notifiable Low Risk Dealing (NLRD) Identifier: NLRD_Suhrbier_Oct2020: NLRD 1.1(a). Mice were euthanized using carbon dioxide.

The SARS-CoV-2 omicron BA.1 isolate

The BA.1 omicron isolate was obtained at QIMR Berghofer MRI from nasal swabs from a

consented COVID-19 patient by culture in Vero E6 cells (ATCC C1008). The omicron BA.1

isolate, SARS-CoV-2$_{QIMR01}$ (SARS-CoV-2/human/AUS/QIMR01/2022), belongs to the BA.1.17

lineage (GenBank: ON819429 and GISAID EPI_ISL_13414183) [55, 56].

BA.1 viral stocks were propagated in Vero E6 cells as described [57]. UV inactivation of the

BA.1 virus was undertaken using the UVC 500 Ultraviolet Crosslinker (Hoefer) (dose 7650 J/m^2

UVC) as described [57]. Virus stocks were titered using CCID$_{50}$ assays [58].

K18-hACE2 mice, infection and monitoring

Heterozygous K18-hACE2 mice (strain B6.Cg-Tg(K18-ACE2)2Prlmn/J, JAX Stock No:

034860) were purchased from The Jackson Laboratory, USA, and were maintained in-house as

heterozygotes by backcrossing to C57BL/6J mice (Animal Resources Center, Canning Vale WA,

Australia). Heterozygotes were inter-crossed to generate homozygous K18-hACE2 mice (on a

C57BL/6J background [53, 59]) as described [52, 53]. Mice were held under standard animal

house conditions (for details see [55]) and homozygous female mice received intrapulmonary

infections delivered via the intranasal route with 5×10^4 CCID$_{50}$ of virus in 50 μl RPMI 1640

whilst under light anesthesia as described [60]. For each group for each time point, a comparable

mean age and age distribution of mice was used (mean age 26. 3 $\pm$ SD 1.1. weeks). Mice were

weighed and monitored as described [60]. Mice were euthanized using CO_2, lungs were

removed, with the left lung fixed in formalin for histology, the right lung inferior lobe placed in

RNAlater for RNA-Seq and the remaining lobes used for tissue titers determination by CCID$_{50}$

assays using Vero E6 cells as described [57, 60].

CCID$_{50}$ assays

Tissue titers were determined as described [57]. Briefly, 5 fold serial dilutions of clarified tissue

homogenates were applied in duplicates to Vero E6 cells in 96 well plates. After 6 days

cytopathic effects were observed by inverted light microscope. The virus titer was determined by

the method of Spearman and Karber [61].

Histology

Lungs were fixed in 10% formalin, embedded in paraffin, and sections stained with H&E (Sigma

Aldrich). Slides were scanned using Aperio AT Turbo (Aperio, Vista, CA, USA). Areas with

overt leukocyte infiltrates (with high focal densities of dark purple staining nuclei) were

measured manually using the "Pen Tool" in the Aperio ImageScope software v10 (Leica

Biosystems, Mt Waverley, Australia). White space analysis was undertaken using QuPath v0.2.3

[62].

Immunohistochemistry

Immunohistochemistry was undertaken as described using the anti-SARS-CoV-2 spike

monoclonal antibody, SCV2-1E8 [56], except that the monoclonal (IgG2a) was purified using

Protein A affinity chromatography and applied to sections at 4 µg/ml for 1 hr.

RNA-Seq and bioinformatics

RNA-Seq was undertaken as described using Illumina Nextseq 2000 platform generating 75 bp

paired end reads [53]. Mean quality scores were above Q20 for all samples. Mouse RNA-Seq

reads were aligned to a combined mouse (GRCm39, version M27) and SARS-CoV-2 BA.5

reference genome [52] using STAR aligner. Viral read counts were generated using Samtools

v1.16. Read counts for host genes were generated using RSEM v1.3.1. Low count genes were

identified and removed separately for each time point using DESeq2 v1.40.2. Differentially

expressed genes were identified using DESeq2 using a FDR cut-off of 0.05. Pathway analysis

was performed with host DEGs using Ingenuity Pathway Analysis (IPA, v84978992) (QIAGEN)

using the Canonical pathways, Up-Stream Regulators (USR) and Diseases or Functions features

as described [59]. Gene Set Enrichment Analyses (GSEAs) were undertaken using GSEA v4.1.0

with gene sets provided in MSigDB ($\approx$ 45,000 gene sets) and in the Blood Transcription Modules

[63], and gene lists generated using DESeq2 ranked by log_2 fold-change. Relative abundances of

cell types were estimated in R v4.1.0 from RSEM 'expected counts' using SpatialDecon v1.4.3

[64] and cell-type expression matrices obtained from the "Mouse Cell Atlas Lung Cell

expression matrix" and the "NanoString Immune Cell Family expression matrix"

Statistics

186 The t-test (with Welch's correction) was used if the difference in variances was <4 fold, skewness

187 was > - 2 and kurtosis was <2. The t test significance and variance were determined using

188 Microsoft Excel. Skewness and kurtosis were determined using IBM SPSS Statistics for

189 Windows v19.0 (IBM Corp., Armonk, NY, USA). Otherwise, the non-parametric Kolmogorov-

190 Smirnov exact test was performed using GraphPad Prism 10.

Results

SARS-CoV-2 omicron BA.1 provided robust lung infection in K18-hACE2 mice that was cleared by 10 dpi

K18-hACE2 mice were given an intrapulmonary inoculum via the intranasal route of SARS-CoV-2 BA.1 (5×10^4 CCID$_{50}$) or the same inoculum of UV-inactivated BA.1 virus. Lungs were harvested at 2, 5, 10, 30 and 66 days post infection (dpi) and tissue titers (Fig. 1a), RNA-Seq (Fig. 1b and see below) and histology (see below) undertaken. Lung viral titers reached a mean titer of $7.86 \pm$ SD 0.49 log$_{10}$CCID$_{50}$/g on 2 days post infection (dpi) (Fig. 1a). Using the same assay system, infection with an original strain isolate in K18-hACE2 mice provided a comparable titer at this time ($7.19 \pm$ SD 0.44 log$_{10}$CCID$_{50}$/g) [53]. However, at 5 dpi the mean titers in BA.1 infected lungs was $4.04 \pm$ SD 0.8 log$_{10}$CCID$_{50}$/g (Fig. 1a), 1.8 logs lower than those seen after infection with an original strain isolate ($5.8 \pm$ SD 0.9 log$_{10}$CCID$_{50}$/g) [53]. By 10 dpi, 3 of 4 BA.1 infected mice showed no detectable virus titers (Fig. 1a).

Fig 1. Viral loads and immunohistochemistry of viral antigen in lungs of BA.1 infected K18-hACE2 mice. **a** Lung tissue titers at the indicated days post infection with BA.1 (limit of detection ≈ 2 log$_{10}$CCID$_{50}$/g). **b** Viral read counts per million reads, determined by RNA-Seq for K18-hACE2 mice infected with BA.1 or inoculated with UV-inactivated BA.1 virus. Statistics by Kolmogorov Smirnov Exact tests. **c** Viral read counts per million reads, determined by RNA-Seq for K18-hACE2 mice infected with original strain isolates; regraphed from Bishop et al. 2022 [53]. **d** Immunohistochemistry of BA.1-infected lungs at 2 dpi, stained with an anti-spike monoclonal antibody. Staining of bronchial epithelia cells (dark brown) can be seen top left and

bottom right (left image), with enlargements of these two areas shown in the right 2 images. **e**

As for d showing staining of cells surrounding alveoli; morphology consistent with that of

pneumocytes (alveolar epithelial cells). * alveolar air sacs.

Viral read counts of infected mice obtained from RNA-Seq data showed a similar trend (Fig.

1b) to viral titers. No significant read counts were obtained from mice inoculated with UV-

inactivated virus (Fig. 1b). Previously reported RNA-Seq viral read count data for infection of

K18-hACE2 mice by original strain isolates is provided as a comparison (Fig. 1c). This data is

taken from three independent studies [53, 65, 66] and has been presented previously [53], but is

regraphed herein in a slightly different format to facilitate comparison (Fig. 1c). Mean lung viral

RNA levels on 2 dpi with an original strain isolate were $\approx$ 17 fold higher than those seen after

BA.1 infection, and they did not substantially decline thereafter (Fig. 1c). Overall the reduced

lung viral loads in BA.1-infected mice is consistent with previous reports [67, 68].

As described previously [52, 67, 68], most mice in the BA.1/K18-hACE2 model do not

develop a brain infection. Brain infection is associated with >20% weight loss and an ensuing

ethical requirement for euthanasia (S1a Fig). After infection of K18-hACE2 mice with original

strain isolates, mice succumb to brain infections between 4 and 7 dpi [52, 60]. This BA.1 model

thus provides a significant lung infection, and permits analysis of long term responses to SARS-

CoV-2.

234 BA.1 infected bronchial epithelial cells and pneumocytes in K18-

235 hACE2 mouse lungs

Immunohistochemistry (IHC) of lungs taken 2 dpi (peak viral load), using an anti-spike

monoclonal antibody [56], illustrated clear staining of bronchial epithelial cells (Fig. 1d) and

cells surrounding alveoli that have a morphology consistent with pneumocytes (Fig. 1e). The

BA.1/K18-hACE2 model thus recapitulates the main cell tropisms seen in human lung tissues,

with infection of bronchial and alveolar epithelial cells by omicron variants well described [69,

70].

BA.1 induced histopathological lesions in lungs that resolved by 66 dpi

H&E stained sections of lung tissues from BA.1 infected K18-hACE2 mice illustrated a series of

histopathological features (Fig. 2, S2 Fig) previously identified in SARS-CoV-2 mouse models

[71], although they appeared less severe than those seen after infection of K18-hACE2 mice with

original strain isolates, consistent with previous reports [67, 68]. For instance, in the latter,

pronounced sloughing of the bronchial epithelium and fulminant occlusions of bronchi (with

serum and red blood cells) were observed [66, 72].

Fig 2. Histology of lungs after BA.1 infection of K18-hACE2 mice. **a** H&E staining of lung

tissue sections at the indicated days after infection/inoculation. Green dotted lines encircle overt

foci of cellular infiltrates. **b** Quantitation of cellular infiltrates. All cellular infiltrates were

marked as in "a" and the summed areas as a percentage of the total lung section area was

determined for each mouse. Statistics by Kolmogorov Smirnov Exact tests. **c** Whole H&E

stained lung sections from day 5 after infection/inoculation, illustrating reduced white space in

lungs from infected mice. **d** QuPath digital analysis of white space in H&E stained whole lung

sections such as those shown in "c". Statistics by t test.

In BA.1 infected mice (as in other models [55, 71]), focal areas of leucocyte infiltration were

clearly evident, characterized by dense clusters of dark-purple staining leukocyte nuclei;

although the infiltrates were more diffuse at 2 dpi (Fig. 2a, green dashed lines). These areas were

measured manually using the Aperio ImageScope "Pen Tool" (Fig. 2a, green dashed lines),

providing quantitation of leukocyte infiltration. Significantly higher levels of infiltration were

seen for BA.1-infected versus UV-inactivated BA.1-inoculated mice at all time-points except 66

dpi, with infiltrates also substantially reduced by 30 dpi (Fig. 2b) (Aperio Positive Pixel Count

assessment of leukocyte infiltrates also showed significantly higher infiltrate levels at 5 dpi; S1b

Fig). Lung consolidation with resulting loss of white space, primarily as a result of diminished

alveolar air sack volumes, was also clearly seen at 5 dpi (Fig. 2c). Significant loss of white space

at 5 dpi was confirmed by image analysis of scanned lung sections using QuPath (Fig. 2d).

Minor intrusion of red blood cells and serum into bronchi, and some alveolar edema were also

occasionally observed in BA.1-infected mice (S2 Fig). H&E images from PBS and naïve control

lungs are shown in S3 Fig.

In summary, lungs of BA.1 infected K18-hACE2 mice displayed some typical acute

histopathological features described previously for SARS-CoV-2 infections, although less severe

than after infection with original strain isolates. At 30 dpi BA.1 infection-associated

histopathology was clearly resolving, and by 66 dpi had largely resolved.

Cell types associated with resolution of inflammation were evident

on 5 and 10 dpi

BA.1 infected lungs and UV BA.1 inoculated lungs were compared at each time point (2, 5, 10,

30 and 66 dpi) by RNA-Seq, with DESeq2 [73]. Full gene lists and bioinformatic analyses are

provided for 2 dpi (S1 Table), 5 dpi (S2 Table), 10 dpi (S3 Table), 30 dpi (S4 Table), and 66 dpi

(S5 Table).

Ingenuity Pathway analysis (IPA) Diseases or Functions annotations for Activation of

leukocytes, Chemotaxis of leukocytes, Leukocyte migration and Quantity of leukocytes showed a

progressive decline over time, with a slight elevation 10 dpi (Fig. 3a), consistent with Fig. 2b.

To gain insights into the type of cellular infiltrates present in the lungs over time post infection

with BA.1, cellular deconvolution (SpacialDecon) analysis using full gene sets (S1 – S5 Tables)

were used together with the gene expression matrices from the "Mouse Cell Atlas (MCA) Lung

Cell expression matrix" and the "NanoString (NS) Immune Cell Family expression matrix". As

might be expected [74, 75], a series of innate mononuclear leukocyte populations were identified

at 5 dpi, which included macrophages, monocytes, dendritic cells, and NK cells, as well as

gamma delta T cells [76] (Fig. 3b). Increased abundance of nuocytes, ILC2 cells, was also

identified on both 5 and 10 dpi (Fig. 3b,c, nuocytes). These innate effector leukocytes mediate

type-2 immunity [77] and are involved in the initiation of inflammation, but also have key roles

in damage control, homeostasis and repair [78, 79]. Lower frequencies of ILC2 cells have been

associated with severe COVID-19 [80]. An increase abundance score for pericytes was also

identified in infected lungs at 5 dpi (but not other times) (S4 Fig). Pericytes express ACE2 and

binding of spike protein has been shown *in vitro* to induce profound transcriptional changes in

these cells [81, 82], which may explain this result.

Fig 3. Cellular infiltrates and types identified by RNA-Seq of lungs for 2 – 30 dpi. **a** Z

scores for the indicated annotations from Ingenuity Pathway analysis (IPA) Diseases or Function

analysis plotted against dpi. (N.I. not identified; for 66 dpi there were insufficient DEGs for IPA

analysis). **b** Using cellular deconvolution (SpatialDecon) and the gene expression matrices from the "Mouse Cell Atlas (MCA) Lung Cell expression matrix" and the "NanoString (NS) Immune Cell Family expression matrix", cell types whose abundance scores were significantly different on day 5 were identified (q values provided). Blue text indicates cell type associated with inflammation resolution. **c** As for b but for day 10. Blue text indicates cell type associated with inflammation resolution. We assume "colon" T reg are well annotated in the NS matrices (and are thus readily identified), and represent tissue T regs, rather than implying that colonic T regs have migrated to the lungs.

At 10 dpi, in addition to nuocytes, significantly increased abundance of three more cell types associated with repair and return to homeostasis was evident (Fig. 3c, blue text). Inducible T regulatory cells (T regs) characterised by low neurophilin expression (Nrp-1^{lo}) [83] were more abundant in infected lungs. T regs secrete *inter alia* IL-10 and are generally protective [84], not just by dampening proinflammatory responses, but also via prevention of fibrosis and maintenance of tissue homeostasis [85]. Interstitial macrophages were also elevated in infected lungs, with these cells having an inherently anti-inflammatory phenotype, secreting *inter alia* IL-10 (USR z score 1.69, q=1.39E-29, S3 Table) and supporting T regs in the lung [86]. An Ear2+ alveolar macrophage population was also identified, with Nra4a1+, Ear2+ macrophages recently found to have a novel reparative phenotype [87]. Although perhaps proinflammatory, CD8 T cells were also more abundant at 10 dpi (Fig. 3c), with CD8 T cells recently identified as key to protection against SARS-CoV-2 [88].

Cytokine signatures in BA.1 infected lungs and their resolution by 30-66 dpi

Differentially expressed genes (DEGs, q<0.05) (S1 –S5 Tables) were analyzed by IPA, with cytokine up-stream regulator (USR) z scores illustrated by a heat map (Fig. 4a). The acute cytokine responses were similar to those published previously for K18-hACE2 mouse models of SARS-CoV-2 infection [53]. When cytokine USR z scores at 2 dpi from BA.1 infection (Fig. 4a) were compared with those from original isolate infection [53], small differences in z scores (between ≈1 and -1) emerged (Fig. 4b). A series of USRs associated with less severe disease had slightly higher z score in BA.1 infected mice; these included EPO (erythropoietin) [89], IL-15 [90], and increased type I interferon responses [17, 48] (Fig. 4b). The z score was also higher for IL-10, with IL-10 elevated in severe COVID-19, but also able to suppress hyper-cyotokinemia [18-20]. Elevated IL-4, IL-9 and IL-13 have been associated with more severe disease [21, 22, 91], although IL-9, IL-13 (and IL-5) are produced by type 2 innate lymphoid cells (ILC2 cells) [92] (see below), and increased IL-4 and IL-13 signatures may (in this context) reflect a relatively more balanced Th1/Th2 response [25, 26]. Elevated OSM and IL-6 are also associated with more severe disease [17, 93]. Another group of cytokines/chemokines associated with increased disease severity (IL-2, IFNG, IL-17, CSF2, CCL5, CXCL8, CCL2 [10, 94, 95] showed slightly lower z scores for BA.1 infected lungs (Fig. 4b). IL-33 may drive fibrosis [96] and also showed a lower USR z score (Fig. 4b).

Fig 4. Inflammatory pathways identified by RNA-Seq of lungs for 2 – 30 dpi. a Heat map of cytokine UpStream Regulator (USR) z scores determined by IPA. White boxes means no cytokine USR annotation was returned for 30 dpi by IPA. RNA-Seq identified too few DEGs on

66 dpi for IPA analysis. **b** IPA cytokine USRs from BA.1 infected K18-hACE2 mice were

compared with those previously reported for the more severe infection of K18-hACE2 mice by

an original strain isolate. Differences in cytokine z scores for 2 dpi (with a cutoff filter of >0.2

and < -0.2) are plotted. BA.1 infected lungs show slightly increased z scores for some cytokine

signatures associated with reduced disease severity (green) and slightly reduced z scores for

some cytokine signatures associated with increased disease severity (red).

Importantly, the cytokine response signatures had substantially abated by 30 dpi (Fig. 4a),

with the small number of DEGs on day 66 insufficient for meaningful pathway analysis. In

addition, few, if any, upregulated inflammatory DEGs were identified at 66 dpi (see below).

Transcriptional responses 66 dpi reflected adaptive immunity and resolution

The RNA-Seq data sets and bioinformatic analyses comparing lungs from BA.1 infected mice 66

dpi with mice 66 days after inoculation with UV-inactivated virus (S5 Table) are summarized in

Fig. 5. Only 29 DEGs were identified, of which 14 were immunoglobulin genes. This represents

an insufficient number of DEGs for meaningful pathway analyses. DEGs were thus clustered

into several groupings based on function (Fig. 5, described below). A plot of the first two

principle components of normalized Variance Stabilizing Transformation (VST) counts from all

experimental groups indicated that the relatively small number of DEGs for day 66 was due to

limited separation between BA.5 infected and UV control groups (S5 Fig).

Most of the upregulated DEGs in lungs at 66 dpi were immunoglobulin genes (Fig. 5) consistent with a more rigorous development of an adaptive immune response in BA.1 infected mice compared with mice inoculated with UV-inactivated virus. GSEAs also illustrate the development of T cell responses, with down-regulation of Nr4a family members potentially suggesting avoidance of T cell exhaustion [97]. However, the three Nr4a genes have a number of roles in T cell biology [98], and also have a diverse set of other activities in *inter alia* macrophages [87, 99] and epithelial cells [100].

The three upregulated, non-Ig genes, can be associated with wound healing/tissue repair; Grb7 (Growth factor receptor-bound protein 7) [101], Arfgap3 (ADP-ribosylation factor GTPase-activating protein 3) [102, 103], and Cap1 (cyclase-associated protein 1) [104]. In addition, Dusp8 (Dual specificity phosphatase 8) was down-regulated, with Dusp8 protein involved in negative regulation of MAP kinase superfamily members; thus Dusp down-regulation would promote cellular proliferation and differentiation [105].

Many of the remaining down-regulated genes have roles in various immune activities (Fig. 5): Prkd1 (Serine/threonine-protein kinase D1), involved in *inter alia* promoting neutrophil infiltration [106]; S1pr1 (Sphingosine-1-phosphate receptor 1), essential for immune cell trafficking [107]; Coq10b, required for coenzyme Q function [108, 109]; Kcne4 (Potassium voltage-gated channel subfamily E member 4), an immune regulator via modulation of Kv1.3 [110]; Atf3, a stress-inducible transcription factor [111]; Marco, a scavenger receptor [112]; and Thbs1 (thrombospondin), a matricellular protein involved in lung inflammation resolution and

repair [113]. Two stress-inducible genes, Atf3 and Hspa1b (a heat shock protein 70 member)

were down-regulated, with Atf3 and Hsp70 upregulated in a large range of stress responses [100,

111].

In summary, at 66 dpi there were very few DEGs, with no indication of ongoing inflammation

or tissue dysregulation, with identification of only a small number of upregulated genes that

could be associated with wound healing processes.

406 Genes whose expression positively correlated with dpi were

407 associated with ciliated epithelial cells and protein synthesis

Using the complete gene sets (S1 – S5 Tables), genes were identified whose expression showed

significant positive correlation with dpi (Pearson correlations, FDR corrected p, q<0.05). Genes

whose expression also showed significant positive correlation with days post inoculation of UV-

inactivated virus were excluded from the analysis, thereby retaining only genes associated with

active infection. Immunoglobulin genes were also removed as the development of immune

responses is understood and established. The process identified 2786 genes (S6 Table).

Enrichr and IPA analyses of the 2786 genes indicated a significant association with ciliated

epithelial cells and protein synthesis, with 59 of the 2786 representing ribosomal protein genes

(Fig. 6a; S6 Table). Ciliated epithelial cells are a known target of SARS-CoV-2 infection [60,

114], with omicron variants showing a higher propensity to infect these cells [69]. The increase

in ciliated epithelial cell genes thus indicates growth/recovery and differentiation of these cells

over the 2-66 days of the experiment. The increase in signatures associated with protein

translation during this period is consistent with the development of adaptive immune responses

and ongoing tissue repair processes.

Fig 6. Genes whose expression correlated with dpi. **a** Using the whole gene lists, genes whose expression showed a statically significant positive correlation with dpi were identified. After removing genes that also positively correlated with dpi in the UV-inactivated BA.1 inoculated control groups, 2786 genes remained. The 2786 genes were analyzed by Enrichr and IPA (for the latter the r value was entered in the log2 expression column). Significant results clustered into two categories 'Ciliated epithelial cell' and 'Protein synthesis'. Of the 2786 genes, 59 were RPL (ribosomal protein) genes. **b** Using the whole gene lists, genes whose expression showed a statically significant negative correlation with dpi were identified. After removing genes that also negatively correlated with dpi in the UV control groups, 1996 genes remained. The 1996 genes were analyzed by IPA as above, with some top annotations shown. **c** Normalized expression data (VST - Variance Stabilizing Transformation) for five of the 1996 genes are plotted against dpi, with linear regression lines and 95% confidence intervals. Pearson correlation coefficients (r) and corrected significances, FDR (q), are provided for each gene. **d** Using cellular deconvolution (as in Fig. 3b,c), cell types whose abundance scores significantly positive correlated with dpi were identified. Linear regression lines and 95% confidence intervals, and Pearson correlation coefficients (r) with corrected significances, FDR (q), are provided. (Full data sets for this figure are provided in S6 Table).

Genes whose expression negatively correlated with dpi indicated inflammation resolution

Using the complete gene sets, genes were identified whose expression showed significant negative correlation with dpi (Pearson correlations, q<0.05). Genes whose expression also

showed significant negative correlation with days post inoculation of UV-inactivated virus were

excluded from the analysis, thereby retaining only genes associated with active infection. The

process identified 1996 genes (S6 Table).

IPA analyses of the 1996 genes identified reductions in cell movement/migration (Fig. 6b),

consistent with Fig. 2 and 3a. A series of cytokine USRs were also down-regulated (S6 Table),

with the top cytokine USRs (by negative z score) representing cytokines previously identified as

being robustly associated with COVID-19 disease severity [10]. The latter was supported by

identification of "Pathogen Induced Cytokine Storm Signaling Pathway" as a top IPA Canonical

pathway annotation with a negative z score (Fig. 6b). Also informative were decreasing

neutrophil degranulation and fibrosis signatures (Fig. 6b); both phenomena have been associated

with long COVID [115, 116].

Negative correlations between expression and dpi are plotted for five of the 1996 genes (Fig.

6c). These genes were selected as their expression levels have been positively correlated with

COVID-19 disease severity and/or they have been proposed as potential targets for prognosis or

treatment; SerpinE1 [117], Cxcl10 [118], Nfkb1 [119], IL6 [120], and Hspa1b (a Hsp70 family

member) [121, 122].

Cell types whose abundance scores increased with dpi indicated lung regeneration

Cell types whose abundance scores (determined by cellular deconvolution) showed a significant

positive correlation (Pearsons q<0.05) with dpi were identified. The top scoring cell type was

ciliated cells (Fig. 6d); consistent with Fig. 6a. The analysis also identified significant increases

in AT1 cells, which likely reflects the AT2 to AT1 transition during alveolar epithelial cell

regeneration seen after SARS-CoV-2 alveolar damage in mouse models [123]. Increasing

abundance scores for reticular fibroblasts, are also consistent with successful lung repair [49,

124], with no indication of significant fibrosis. Increasing numbers of innate lymphoid cells

(ILC) likely also reflect the process of returning the tissue to homeostasis [125, 126].

Discussion

We describe herein the transcriptional profile of BA.1 omicron lung infections in K18-hACE2

mice, and the post-infection recovery up to 66 dpi. The expression of hACE2 via the keratin 18

promoter in the K18-hACE2 mice did not lead to overtly aberrant cell tropism in the lungs, with

infection of bronchial and alveolar epithelia cells by omicron variants described previously [69,

70]. Compared to infection with original strain isolates, BA.1 viral loads in lungs were lower

and were rapidly cleared, a feature associated in COVID-19 patients with less severe disease [34]

or subclinical infections [44, 45]. By 66 dpi only a small number of DEGs were identified,

arguing that the infection had been resolved and the tissue had largely returned to normal. UV-

inactivated virus was used as a control to exclude responses to viral inoculation and restrict the

analyses to effects arising from replicating virus. Features often associated with post-acute or

long-COVID were not identified, for instance, there were no indications of prolonged infection

[127], persistent inflammation [128], impaired lung regeneration [129], fibrosis [116], prolonged

T cell activation [130], Th17 bias [131], chronic neutrophil activation [115] or clotting

abnormalities [7].

The concept of "protective inflammation" was recently described for COVID-19 [54] and

clearly applies to many other viral infections where subclinical or mild infections are also well

documented [132]. Deleterious or pathological inflammatory disease processes tend to be the

focus of research, with perhaps less attention paid to characterisation of inflammatory responses

that are resolved and lead to subclinical or mild disease and sequelae-free outcomes.

Pathological versus protective responses are likely to have distinct gene expression profiles,

patterns, magnitudes and sequence of events [133, 134], with the current study describing the

responses associated with resolution of lung infection and inflammation. Importantly, at 10 dpi a

series of cell types associated with damage control, suppression of inflammation, homeostasis

and repair showed increased abundance scores (determined using transcription data from lung

tissues and cellular deconvolution). Specifically these cells were, ILC2 cells [78, 79] (also seen

at 5 dpi), inducible T regs [84, 85], interstitial macrophages [86] and Ear2+ macrophages [87].

Restoration of ciliated cells, regeneration of alveolar epithelial cell with AT2 to AT1 transition

[123], and increases in fibroblast activity without fibrosis [49], also appear to characterize lung

recovery in this model. Clearly, prompt clearance of virus [44, 48] (rather than viral persistence

[127, 135]) and development of protective adaptive immune responses [47] are also likely to be

key to such outcomes. The latter has recently been shown to include CD8 T cells [88], which

may be represented by a population of CD8b1^{high} cells which reached significance by 10 dpi in

the BA.1/K18-hACE2 model.

Defining key aspects of "protective inflammation" remains a challenge for human

populations, as individuals with subclinical or mild disease are generally difficult to identify and

recruit promptly. A better understanding of what characterizes rapid, sequelae-free resolution of

inflammation, would clearly facilitate development of interventions for COVID-19 associated

ARDS, with animal models providing ready access to lung tissues to help unravel the complex

mechanisms that underpin such processes.

Acknowledgements

The authors thank the following staff at QIMR Berghofer MRI; Dr I. Anraku for management of the PC3 facility, Dr Viviana Lutzky for proof reading, Ashwini R Potadar and Crystal Chang for histology services, the animal house staff for mouse breeding and agistment, and Dr. Gunter Hartel for assistance with statistics.

Supplemental information

The paper is supported by supplementary figures (S1 – S5 Fig) and Tables (S1 – S6 Table).

S1 Fig. Overview of change in body weight overtime and ratio of purple over red staining in lungs at each timepoint.

S2 Fig. Histopathological features of viral infection in lungs of BA.1 infected mice in comparison to UV-inactivated BA.1 inoculated mice.

S3 Fig. H&E staining of control lungs from K18-hACE2 mice.

S4 Fig. Increased pericytes abundance score at 5 dpi.

S5 Fig. First two principle components of normalized VST-transformed counts from all experimental groups.

S1 Table Differential expression and pathway analysis for 2 dpi

S2 Table Differential expression and pathway analysis for 5 dpi

S3 Table Differential expression and pathway analysis for 10 dpi

S4 Table Differential expression and pathway analysis for 30 dpi

S5 Table Differential expression and pathway analysis for 66 dpi

S6 Table Gene expression vs dpi

Author Contributions

Conceptualization: Andreas Suhrbier, Daniel J. Rawle, Cameron R. Bishop.

Data curation; Cameron R. Bishop, Andreas Suhrbier.

Funding acquisition: Andreas Suhrbier, Daniel J. Rawle.

Investigation: Agnes Carolin, Kexin Yan, Bing Tang, Wilson Nguyen.

Methodology: Daniel J. Rawle, Cameron R. Bishop, Andreas Suhrbier.

Supervision: Daniel J. Rawle, Cameron R. Bishop, Wilson Nguyen, Andreas Suhrbier.

Visualization: Andreas Suhrbier, Cameron R. Bishop, Kexin Yan.

Writing –original draft: Andreas Suhrbier.

Writing – review & editing: Agnes Carolin, Cameron R. Bishop, Daniel J. Rawle.

References

1. WHO COVID-19 dashboard 2023 [cited 2024 9 January]. Available from: https://data.who.int/dashboards/covid19/cases.
2. WHO COVID-19 dashboard. https://data.who.int/dashboards/covid19/deaths?n=c. Accessed Feb 2024.
3. Silva MJ, Ribeiro LR, Gouveia MI, Marcelino BD, Santos CS, Lima KV, et al. Hyperinflammatory Response in COVID-19: A Systematic Review. Viruses. 2023;15(2):553. doi: 10.3390/v15020553.
4. Verma G, Dhawan M, Saied AA, Kaur G, Kumar R, Emran TB. Immunomodulatory approaches in managing lung inflammation in COVID-19: A double-edge sword. Immunity, Inflammation and Disease. 2023;11(9):e1020. doi: https://doi.org/10.1002/iid3.1020.
5. Al-Nesf MAY, Abdesselem HB, Bensmail I, Ibrahim S, Saeed WAH, Mohammed SSI, et al. Prognostic tools and candidate drugs based on plasma proteomics of patients with severe COVID-19 complications. Nature Communications. 2022;13(1):946. doi: 10.1038/s41467-022-28639-4.
6. Mueller MR, Ganesh R, Hurt RT, Beckman TJ. Post-COVID Conditions. Mayo Clinic Proceedings. 2023;98(7):1071-8. doi: https://doi.org/10.1016/j.mayocp.2023.04.007.
7. Altmann DM, Whettlock EM, Liu S, Arachchillage DJ, Boyton RJ. The immunology of long COVID. Nature Reviews Immunology. 2023;23(10):618-34. doi: 10.1038/s41577-023-00904-7.
8. Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. Nature Reviews Microbiology. 2023;21(3):133-46. doi: 10.1038/s41579-022-00846-2.
9. Sick and tired: Casting a long shadow. Report April 2023. https://www.aph.gov.au/Parliamentary_Business/Committees/House/Health_Aged_Care_and_Sport/LongandrepeatedCOVID/Report. Accesses Feb 2024. .
10. Nasrollahi H, Talepoor AG, Saleh Z, Eshkevar Vakili M, Heydarinezhad P, Karami N, et al. Immune responses in mildly versus critically ill COVID-19 patients. Frontiers in Immunology. 2023;14:1077236. doi: 10.3389/fimmu.2023.1077236.
11. Wang J, Jiang M, Chen X, Montaner LJ. Cytokine storm and leukocyte changes in mild versus severe SARS-CoV-2 infection: Review of 3939 COVID-19 patients in China and emerging pathogenesis and therapy concepts. Journal of Leukocyte Biology. 2020;108(1):17-41. doi: 10.1002/jlb.3covr0520-272r.
12. Qin R, He L, Yang Z, Jia N, Chen R, Xie J, et al. Identification of Parameters Representative of Immune Dysfunction in Patients with Severe and Fatal COVID-19 Infection: a Systematic Review and Meta-analysis. Clinical Reviews in Allergy & Immunology. 2023;64(1):33-65. doi: 10.1007/s12016-021-08908-8.
13. Kalinina O, Golovkin A, Zaikova E, Aquino A, Bezrukikh V, Melnik O, et al. Cytokine Storm Signature in Patients with Moderate and Severe COVID-19. International Journal of Molecular Sciences. 2022;23(16):8879. doi: 10.3390/ijms23168879.
14. Patel J, Bass D, Beishuizen A, Bocca Ruiz X, Boughanmi H, Cahn A, et al. A randomised trial of anti-GM-CSF otilimab in severe COVID-19 pneumonia (OSCAR). Eur Respir J. 2023;61(2):2101870. Epub 2022/10/14. doi: 10.1183/13993003.01870-2021. PubMed PMID: 36229048.

15. Ghofrani Nezhad M, Jami G, Kooshkaki O, Chamani S, Naghizadeh A. The Role of
Inflammatory Cytokines (Interleukin-1 and Interleukin-6) as a Potential Biomarker in the
Different Stages of COVID-19 (Mild, Severe, and Critical). J Interferon Cytokine Res.
2023;43(4):147-63. Epub 2023/04/17. doi: 10.1089/jir.2022.0185. PubMed PMID:
37062817.
16. Mohd Zawawi Z, Kalyanasundram J, Mohd Zain R, Thayan R, Basri DF, Yap WB.
Prospective Roles of Tumor Necrosis Factor-Alpha (TNF-α) in COVID-19:
Prognosis, Therapeutic and Management. International Journal of Molecular Sciences.
2023;24(7):6142. doi: 10.3390/ijms24076142.
17. Arunachalam PS, Wimmers F, Mok CKP, Perera R, Scott M, Hagan T, et al. Systems
biological assessment of immunity to mild versus severe COVID-19 infection in humans.
Science. 2020;369(6508):1210-20. Epub 2020/08/14. doi: 10.1126/science.abc6261.
PubMed PMID: 32788292; PubMed Central PMCID: PMCPMC7665312.
18. Habel JR, Chua BY, Kedzierski L, Selva KJ, Damelang T, Haycroft ER, et al. Immune
profiling of SARS-CoV-2 infection during pregnancy reveals NK cell and γδ T cell
perturbations. JCI Insight. 2023;8(7):e167157. Epub 2023/04/11. doi:
10.1172/jci.insight.167157. PubMed PMID: 37036008; PubMed Central PMCID:
PMCPMC10132165.
19. Carlini V, Noonan DM, Abdalalem E, Goletti D, Sansone C, Calabrone L, et al. The
multifaceted nature of IL-10: regulation, role in immunological homeostasis and its
relevance to cancer, COVID-19 and post-COVID conditions. Frontiers in Immunology.
2023;14:1161067. doi: 10.3389/fimmu.2023.1161067.
20. Shih L-J, Yang C-C, Liao M-T, Lu K-C, Hu W-C, Lin C-P. An important call: Suggestion of
using IL-10 as therapeutic agent for COVID-19 with ARDS and other complications.
Virulence. 2023;14(1):2190650. doi: 10.1080/21505594.2023.2190650.
21. Donlan AN, Sutherland TE, Marie C, Preissner S, Bradley BT, Carpenter RM, et al. IL-13 is
a driver of COVID-19 severity. JCI Insight. 2021;6(15):e150107. Epub 2021/06/30. doi:
10.1172/jci.insight.150107. PubMed PMID: 34185704; PubMed Central PMCID:
PMCPMC8410056.
22. Gibellini L, De Biasi S, Meschiari M, Gozzi L, Paolini A, Borella R, et al. Plasma Cytokine
Atlas Reveals the Importance of TH2 Polarization and Interferons in Predicting COVID-19
Severity and Survival. Front Immunol. 2022;13:842150. Epub 2022/04/08. doi:
10.3389/fimmu.2022.842150. PubMed PMID: 35386702; PubMed Central PMCID:
PMCPMC8979161.
23. Sasson J, Moreau GB, Petri WA. The role of interleukin 13 and the type 2 immune pathway
in COVID-19: A review. Annals of Allergy, Asthma & Immunology. 2023;130(6):727-32.
doi: https://doi.org/10.1016/j.anai.2023.03.009.
24. Pourgholaminejad A, Pahlavanneshan S, Basiri M. COVID-19 immunopathology with
emphasis on Th17 response and cell-based immunomodulation therapy: Potential targets and
challenges. Scand J Immunol. 2022;95(2):e13131. Epub 2021/12/23. doi: 10.1111/sji.13131.
PubMed PMID: 34936112.
25. Gupta P, Patel K, Garg G, Mathew BJ, Kale D, Singh J, et al. Imbalance of T Helper Cell
Subset Specific Transcription Factors and Associated Cytokines in Patients with Severe
COVID-19. Gene Expression. 2023;22(3):159-66. doi: 10.14218/GE.2023.00016.
26. Hasegawa T, Hato T, Okayama T, Ikeo K, Miyamoto Y, Iwanaga N, et al. Th1 cytokine
endotype discriminates and predicts severe complications in COVID-19. European Cytokine
Network. 2022;33(2):1-12. doi: 10.1684/ecn.2022.0477.

27.   Pavel AB, Glickman JW, Michels JR, Kim-Schulze S, Miller RL, Guttman-Yassky E.
642   Th2/Th1 Cytokine Imbalance Is Associated With Higher COVID-19 Risk Mortality.
643   Frontiers in Genetics. 2021;12:706902. doi: 10.3389/fgene.2021.706902.
644   28.   Gil-Etayo FJ, Garcinuño S, Utrero-Rico A, Cabrera-Marante O, Arroyo-Sanchez D,
645   Mancebo E, et al. An Early Th1 Response Is a Key Factor for a Favorable COVID-19
646   Evolution. Biomedicines. 2022;10(2):296. doi: 10.3390/biomedicines10020296.
647   29.   Dhawan M, Rabaan AA, Alwarthan S, Alhajri M, Halwani MA, Alshengeti A, et al.
648   Regulatory T Cells (Tregs) and COVID-19: Unveiling the Mechanisms, and Therapeutic
649   Potentialities with a Special Focus on Long COVID. Vaccines. 2023;11(3):699. doi:
650   10.3390/vaccines11030699.
651   30.   Kosyreva A, Dzhalilova D, Lokhonina A, Vishnyakova P, Fatkhudinov T. The Role of
652   Macrophages in the Pathogenesis of SARS-CoV-2-Associated Acute Respiratory Distress
653   Syndrome. Frontiers in Immunology. 2021;12:682871. doi: 10.3389/fimmu.2021.682871.
654   31.   Meidaninikjeh S, Sabouni N, Marzouni HZ, Bengar S, Khalili A, Jafari R. Monocytes and
655   macrophages in COVID-19: Friends and foes. Life Sciences. 2021;269:119010. doi:
656   https://doi.org/10.1016/j.lfs.2020.119010.
657   32.   Dadras O, Afsahi AM, Pashaei Z, Mojdeganlou H, Karimi A, Habibi P, et al. The
658   relationship between COVID-19 viral load and disease severity: A systematic review.
659   Immunity, Inflammation and Disease. 2022;10(3):e580. doi:
660   https://doi.org/10.1002/iid3.580.
661   33.   Waller H, Carmona-Vicente N, James A, Govender M, Hopkins FR, Larsson M, et al. Viral
662   load at hospitalization is an independent predictor of severe COVID-19. Eur J Clin Invest.
663   2023;53(1):e13882. Epub 2022/10/04. doi: 10.1111/eci.13882. PubMed PMID: 36190270;
664   PubMed Central PMCID: PMCPMC9874715.
665   34.   Puhach O, Meyer B, Eckerle I. SARS-CoV-2 viral load and shedding kinetics. Nature
666   Reviews Microbiology. 2023;21(3):147-61. doi: 10.1038/s41579-022-00822-w.
667   35.   Alene M, Yismaw L, Assemie MA, Ketema DB, Mengist B, Kassie B, et al. Magnitude of
668   asymptomatic COVID-19 cases throughout the course of infection: A systematic review and
669   meta-analysis. PLOS ONE. 2021;16(3):e0249090. doi: 10.1371/journal.pone.0249090.
670   36.   Oran DP, Topol EJ. The Proportion of SARS-CoV-2 Infections That Are Asymptomatic.
671   Annals of Internal Medicine. 2021;174(5):655-62. doi: 10.7326/M20-6976.
672   37.   Polyak MJ, Abrahamyan L, Bego MG. Editorial: Immune determinants of COVID-19
673   protection and disease: A focus on asymptomatic COVID and long COVID. Front Immunol.
674   2023;14:1185693. Epub 2023/04/15. doi: 10.3389/fimmu.2023.1185693. PubMed PMID:
675   37056777; PubMed Central PMCID: PMCPMC10086613.
676   38.   Sah P, Fitzpatrick MC, Zimmer CF, Abdollahi E, Juden-Kelly L, Moghadas SM, et al.
677   Asymptomatic SARS-CoV-2 infection: A systematic review and meta-analysis. Proceedings
678   of the National Academy of Sciences. 2021;118(34):e2109229118. doi:
679   10.1073/pnas.2109229118.
680   39.   Zuin M, Gentili V, Cervellati C, Rizzo R, Zuliani G. Viral Load Difference between
681   Symptomatic and Asymptomatic COVID-19 Patients: Systematic Review and Meta-
682   Analysis. Infectious Disease Reports. 2021;13(3):645-53. doi: 10.3390/idr13030061.
683   40.   Tsukagoshi H, Shinoda D, Saito M, Okayama K, Sada M, Kimura H, et al. Relationships
684   between Viral Load and the Clinical Course of COVID-19. Viruses. 2021;13(2):304. doi:
685   10.3390/v13020304.
686   41.   Ra SH, Lim JS, Kim GU, Kim MJ, Jung J, Kim SH. Upper respiratory viral load in
687   asymptomatic individuals and mildly symptomatic patients with SARS-CoV-2 infection.

Thorax. 2021;76(1):61-3. Epub 2020/09/24. doi: 10.1136/thoraxjnl-2020-215042. PubMed PMID: 32963115.

42. Yin S-W, Zhou Z, Wang J-L, Deng Y-F, Jing H, Qiu Y. Viral loads, lymphocyte subsets and cytokines in asymptomatic, mildly and critical symptomatic patients with SARS-CoV-2 infection: a retrospective study. Virology Journal. 2021;18(1):126. doi: 10.1186/s12985-021-01597-x.

43. Ooi EE, Low JG. Asymptomatic SARS-CoV-2 infection. The Lancet Infectious Diseases. 2020;20(9):996-8. doi: https://doi.org/10.1016/S1473-3099(20)30460-6.

44. Bae S, Kim JY, Lim SY, Park H, Cha HH, Kwon J-S, et al. Dynamics of Viral Shedding and Symptoms in Patients with Asymptomatic or Mild COVID-19. Viruses. 2021;13(11):2133. doi: 10.3390/v13112133.

45. Long Q-X, Tang X-J, Shi Q-L, Li Q, Deng H-J, Yuan J, et al. Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. Nature Medicine. 2020;26(8):1200-4. doi: 10.1038/s41591-020-0965-6.

46. De Biasi S, Tartaro DL, Gibellini L, Paolini A, Quong A, Petes C, et al. Endogenous control of inflammation characterizes pregnant women with asymptomatic or paucisymptomatic SARS-CoV-2 infection. Nature Communications. 2021;12(1):4677. doi: 10.1038/s41467-021-24940-w.

47. Le Bert N, Clapham HE, Tan AT, Chia WN, Tham CYL, Lim JM, et al. Highly functional virus-specific cellular immune response in asymptomatic SARS-CoV-2 infection. Journal of Experimental Medicine. 2021;218(5):e20202617. doi: 10.1084/jem.20202617.

48. Le Bert N, Samandari T. Silent battles: immune responses in asymptomatic SARS-CoV-2 infection. Cellular & Molecular Immunology. 2024;21(2):159-70. doi: 10.1038/s41423-024-01127-z.

49. Soares-Schanoski A, Sauerwald N, Goforth CW, Periasamy S, Weir DL, Lizewski S, et al. Asymptomatic SARS-CoV-2 Infection Is Associated With Higher Levels of Serum IL-17C, Matrix Metalloproteinase 10 and Fibroblast Growth Factors Than Mild Symptomatic COVID-19. Front Immunol. 2022;13:821730. Epub 2022/04/29. doi: 10.3389/fimmu.2022.821730. PubMed PMID: 35479098; PubMed Central PMCID: PMCPMC9037090.

50. Sfikakis PP, Verrou KM, Ampatziadis-Michailidis G, Tsitsilonis O, Paraskevis D, Kastritis E, et al. Blood Transcriptomes of Anti-SARS-CoV-2 Antibody-Positive Healthy Individuals Who Experienced Asymptomatic Versus Clinical Infection. Front Immunol. 2021;12:746203. Epub 2021/10/23. doi: 10.3389/fimmu.2021.746203. PubMed PMID: 34675930; PubMed Central PMCID: PMCPMC8523987.

51. Ma Y, Deng J, Liu Q, Du M, Liu M, Liu J. Long-Term Consequences of Asymptomatic SARS-CoV-2 Infection: A Systematic Review and Meta-Analysis. Int J Environ Res Public Health. 2023;20(2):1613. Epub 2023/01/22. doi: 10.3390/ijerph20021613. PubMed PMID: 36674367; PubMed Central PMCID: PMCPMC9863678.

52. Stewart R, Yan K, Ellis SA, Bishop CR, Dumenil T, Tang B, et al. SARS-CoV-2 omicron BA.5 and XBB variants have increased neurotropic potential over BA.1 in K18-hACE2 mice and human brain organoids. Front Microbiol. 2023;14:1320856. Epub 2023/12/11. doi: 10.3389/fmicb.2023.1320856. PubMed PMID: 38075874; PubMed Central PMCID: PMCPMC10706942.

53. Bishop CR, Dumenil T, Rawle DJ, Le TT, Yan K, Tang B, et al. Mouse models of COVID-19 recapitulate inflammatory pathways rather than gene expression. PLoS Pathog.

2022;18(9):e1010867. Epub 2022/09/27. doi: 10.1371/journal.ppat.1010867. PubMed
PMID: 36155667; PubMed Central PMCID: PMCPMC9536645.

54. Mangalmurti N, Hunter CA. Cytokine Storms: Understanding COVID-19. Immunity.
2020;53(1):19-25. Epub 2020/07/02. doi: 10.1016/j.immuni.2020.06.017. PubMed PMID:
32610079; PubMed Central PMCID: PMCPMC7321048.

55. Yan K, Dumenil T, Tang B, Le TT, Bishop CR, Suhrbier A, et al. Evolution of ACE2-
independent SARS-CoV-2 infection and mouse adaption after passage in cells expressing
human and mouse ACE2. Virus Evol. 2022;8(2):veac063. Epub 2022/08/04. doi:
10.1093/ve/veac063. PubMed PMID: 35919871; PubMed Central PMCID:
PMCPMC9338707.

56. Morgan MS, Yan K, Le TT, Johnston RA, Amarilla AA, Muller DA, et al. Monoclonal
Antibodies Specific for SARS-CoV-2 Spike Protein Suitable for Multiple Applications for
Current Variants of Concern. Viruses. 2023;15(1):139. Epub 2023/01/22. doi:
10.3390/v15010139. PubMed PMID: 36680179; PubMed Central PMCID:
PMCPMC9863740.

57. Rawle DJ, Le TT, Dumenil T, Yan K, Tang B, Nguyen W, et al. ACE2-lentiviral
transduction enables mouse SARS-CoV-2 infection and mapping of receptor interactions.
PLoS Pathog. 2021;17(7):e1009723. Epub 2021/07/03. doi: 10.1371/journal.ppat.1009723.
PubMed PMID: 34214142; PubMed Central PMCID: PMCPMC8282004.

58. Yan K, Rawle DJ, Le TT, Suhrbier A. Simple rapid in vitro screening method for SARS-
CoV-2 anti-virals that identifies potential cytomorbidity-associated false positives. Virol J.
2021;18(1):123. Epub 2021/06/11. doi: 10.1186/s12985-021-01587-z. PubMed PMID:
34107996; PubMed Central PMCID: PMCPMC8188739.

59. Rawle DJ, Le TT, Dumenil T, Bishop C, Yan K, Nakayama E, et al. Widespread
discrepancy in Nnt genotypes and genetic backgrounds complicates granzyme A and other
knockout mouse studies. Elife. 2022;11:e70207. Epub 2022/02/05. doi:
10.7554/eLife.70207. PubMed PMID: 35119362; PubMed Central PMCID:
PMCPMC8816380.

60. Dumenil T, Le TT, Rawle DJ, Yan K, Tang B, Nguyen W, et al. Warmer ambient air
temperatures reduce nasal turbinate and brain infection, but increase lung inflammation in
the K18-hACE2 mouse model of COVID-19. Sci Total Environ. 2022;859(Pt 1):160163.
Epub 2022/11/18. doi: 10.1016/j.scitotenv.2022.160163. PubMed PMID: 36395835;
PubMed Central PMCID: PMCPMC9659553.

61. TCID50_calculator_v2_17-01-20_MB - Excel sheet to calculate TCID50 titers (Spearman &
Kärber method). https://www.klinikum.uni-heidelberg.de/zentrum-fuer-
infektiologie/molecular-virology/welcome/downloads. Accessed Feb 2024.

62. Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, et al.
QuPath: Open source software for digital pathology image analysis. Scientific Reports.
2017;7(1):16878. doi: 10.1038/s41598-017-17204-5.

63. Li S, Rouphael N, Duraisingham S, Romero-Steiner S, Presnell S, Davis C, et al. Molecular
signatures of antibody responses derived from a systems biology study of five human
vaccines. Nat Immunol. 2014;15(2):195-204. Epub 2013/12/18. doi: 10.1038/ni.2789.
PubMed PMID: 24336226; PubMed Central PMCID: PMCPMC3946932.

64. Danaher P, Kim Y, Nelson B, Griswold M, Yang Z, Piazza E, et al. Advances in mixed cell
deconvolution enable quantification of cell types in spatial transcriptomic data. Nat
Commun. 2022;13(1):385. Epub 2022/01/21. doi: 10.1038/s41467-022-28020-5. PubMed
PMID: 35046414; PubMed Central PMCID: PMCPMC8770643 performing this work. No

authors had nonfinancial competing interests. J.M.B. is listed on the relevant U.S. patent 10640816, "Simultaneous quantification of gene expression in a user-defined region of a cross-sectioned tissue".

65. Winkler ES, Bailey AL, Kafai NM, Nair S, McCune BT, Yu J, et al. SARS-CoV-2 infection of human ACE2-transgenic mice causes severe lung inflammation and impaired function. Nat Immunol. 2020;21(11):1327-35. Epub 2020/08/26. doi: 10.1038/s41590-020-0778-2. PubMed PMID: 32839612; PubMed Central PMCID: PMCPMC7578095.

66. Mills RJ, Humphrey SJ, Fortuna PRJ, Lor M, Foster SR, Quaife-Ryan GA, et al. BET inhibition blocks inflammation-induced cardiac dysfunction and SARS-CoV-2 infection. Cell. 2021;184(8):2167-82 e22. Epub 2021/04/04. doi: 10.1016/j.cell.2021.03.026. PubMed PMID: 33811809; PubMed Central PMCID: PMCPMC7962543.

67. Lee NY, Lee YW, Hong SM, On D, Yoon GM, An SH, et al. SARS-CoV-2 Omicron variant causes brain infection with lymphoid depletion in a mouse COVID-19 model. Lab Anim Res. 2023;39(1):8. Epub 2023/05/10. doi: 10.1186/s42826-023-00157-4. PubMed PMID: 37161442; PubMed Central PMCID: PMCPMC10169124.

68. Liu S, Selvaraj P, Sangare K, Luan B, Wang TT. Spike protein-independent attenuation of SARS-CoV-2 Omicron variant in laboratory mice. Cell Rep. 2022;40(11):111359. Epub 2022/09/09. doi: 10.1016/j.celrep.2022.111359. PubMed PMID: 36075211; PubMed Central PMCID: PMCPMC9420700.

69. Wu CT, Lidsky PV, Xiao Y, Cheng R, Lee IT, Nakayama T, et al. SARS-CoV-2 replication in airway epithelia requires motile cilia and microvillar reprogramming. Cell. 2023;186(1):112-30 e20. Epub 2022/12/30. doi: 10.1016/j.cell.2022.11.030. PubMed PMID: 36580912; PubMed Central PMCID: PMCPMC9715480.

70. Chiu MC, Li C, Liu X, Yu Y, Huang J, Wan Z, et al. A bipotential organoid model of respiratory epithelium recapitulates high infectivity of SARS-CoV-2 Omicron variant. Cell Discovery. 2022;8(1):57. doi: 10.1038/s41421-022-00422-1.

71. Yinda CK, Port JR, Bushmaker T, Offei Owusu I, Purushotham JN, Avanzato VA, et al. K18-hACE2 mice develop respiratory disease resembling severe COVID-19. PLoS Pathog. 2021;17(1):e1009195. Epub 2021/01/20. doi: 10.1371/journal.ppat.1009195. PubMed PMID: 33465158; PubMed Central PMCID: PMCPMC7875348.

72. Amarilla AA, Sng JDJ, Parry R, Deerain JM, Potter JR, Setoh YX, et al. A versatile reverse genetics platform for SARS-CoV-2 and other positive-strand RNA viruses. Nat Commun. 2021;12(1):3431. Epub 2021/06/10. doi: 10.1038/s41467-021-23779-5. PubMed PMID: 34103499; PubMed Central PMCID: PMCPMC8187723.

73. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. Epub 2014/12/18. doi: 10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PubMed Central PMCID: PMCPMC4302049.

74. Falck-Jones S, Österberg B, Smed-Sörensen A. Respiratory and systemic monocytes, dendritic cells, and myeloid-derived suppressor cells in COVID-19: Implications for disease severity. Journal of Internal Medicine. 2023;293(2):130-43. doi: https://doi.org/10.1111/joim.13559.

75. Lee MJ, Blish CA. Defining the role of natural killer cells in COVID-19. Nature Immunology. 2023;24(10):1628-38. doi: 10.1038/s41590-023-01560-8.

76. von Borstel A, Nguyen THO, Rowntree LC, Ashhurst TM, Allen LF, Howson LJ, et al. Circulating effector γδ T cell populations are associated with acute coronavirus disease 19

in unvaccinated individuals. Immunology & Cell Biology. 2023;101(4):321-32. doi:
https://doi.org/10.1111/imcb.12623.

77.   Neill DR, Wong SH, Bellosi A, Flynn RJ, Daly M, Langford TKA, et al. Nuocytes represent
a new innate effector leukocyte that mediates type-2 immunity. Nature.
2010;464(7293):1367-70. doi: 10.1038/nature08900.

78.   Fonseca W, Lukacs NW, Elesela S, Malinczak CA. Role of ILC2 in Viral-Induced Lung
Pathogenesis. Front Immunol. 2021;12:675169. Epub 2021/05/07. doi:
10.3389/fimmu.2021.675169. PubMed PMID: 33953732; PubMed Central PMCID:
PMCPMC8092393.

79.   Pyung YJ, Park D-J, Kim CG, Yun C-H. Remodeling and Restraining Lung Tissue Damage
Through the Regulation of Respiratory Immune Responses. Tissue Engineering and
Regenerative Medicine. 2023;20(3):329-39. doi: 10.1007/s13770-022-00516-7.

80.   García M, Kokkinou E, Carrasco García A, Parrot T, Palma Medina LM, Maleki KT, et al.
Innate lymphoid cell composition associates with COVID-19 disease severity. Clin Transl
Immunology. 2020;9(12):e1224. Epub 2020/12/22. doi: 10.1002/cti2.1224. PubMed PMID:
33343897; PubMed Central PMCID: PMCPMC7734472.

81.   Khaddaj-Mallat R, Aldib N, Bernard M, Paquette AS, Ferreira A, Lecordier S, et al. SARS-
CoV-2 deregulates the vascular and immune functions of brain pericytes via Spike protein.
Neurobiol Dis. 2021;161:105561. Epub 2021/11/16. doi: 10.1016/j.nbd.2021.105561.
PubMed PMID: 34780863; PubMed Central PMCID: PMCPMC8590447.

82.   Avolio E, Srivastava PK, Ji J, Carrabba M, Tsang CTW, Gu Y, et al. Murine studies and
expressional analyses of human cardiac pericytes reveal novel trajectories of SARS-CoV-2
Spike protein-induced microvascular damage. Signal Transduct Target Ther. 2023;8(1):232.
Epub 2023/06/03. doi: 10.1038/s41392-023-01489-2. PubMed PMID: 37268620; PubMed
Central PMCID: PMCPMC10236384.

83.   Yadav M, Louvet C, Davini D, Gardner JM, Martinez-Llordella M, Bailey-Bucktrout S, et
al. Neuropilin-1 distinguishes natural and inducible regulatory T cells among regulatory T
cell subsets in vivo. J Exp Med. 2012;209(10):1713-22, S1-19. Epub 2012/09/12. doi:
10.1084/jem.20120822. PubMed PMID: 22966003; PubMed Central PMCID:
PMCPMC3457729.

84.   Gladstone DE, D'Alessio F, Howard C, Lyu M-A, Mock JR, Gibbs KW, et al. Randomized,
double-blinded, placebo-controlled trial of allogeneic cord blood T-regulatory cells for
treatment of COVID-19 ARDS. Blood Advances. 2023;7(13):3075-9. doi:
10.1182/bloodadvances.2022009619.

85.   Franco A, Song J, Chambers C, Sette A, Grifoni A. SARS-CoV-2 spike-specific regulatory
T cells (Treg) expand and develop memory in vaccine recipients suggesting a role for
immune regulation in preventing severe symptoms in COVID-19. Autoimmunity.
2023;56(1):2259133. doi: 10.1080/08916934.2023.2259133.

86.   Bain CC, MacDonald AS. The impact of the lung environment on macrophage development,
activation and function: diversity in the face of adversity. Mucosal Immunol.
2022;15(2):223-34. Epub 2022/01/13. doi: 10.1038/s41385-021-00480-w. PubMed PMID:
35017701; PubMed Central PMCID: PMCPMC8749355.

87.   Chen J, Huang XR, Yang F, Yiu WH, Yu X, Tang SCW, et al. Single-cell RNA Sequencing
Identified Novel Nr4a1(+) Ear2(+) Anti-Inflammatory Macrophage Phenotype under
Myeloid-TLR4 Dependent Regulation in Anti-Glomerular Basement Membrane (GBM)
Crescentic Glomerulonephritis (cGN). Adv Sci (Weinh). 2022;9(18):e2200668. Epub

2022/04/30. doi: 10.1002/advs.202200668. PubMed PMID: 35484716; PubMed Central
PMCID: PMCPMC9218767.
88.   Antoun E, Peng Y, Dong T. Vaccine-induced CD8+ T cells are key to protection from
SARS-CoV-2. Nature Immunology. 2023;24(10):1594-6. doi: 10.1038/s41590-023-01621-y.
89.   Yağcı S, Serin E, Acicbe Ö, Zeren Mİ, Odabaşı MS. The relationship between serum
erythropoietin, hepcidin, and haptoglobin levels with disease severity and other biochemical
values in patients with COVID-19. International Journal of Laboratory Hematology.
2021;43(S1):142-51. doi: https://doi.org/10.1111/ijlh.13479.
90.   Kandikattu HK, Venkateshaiah SU, Kumar S, Mishra A. IL-15 immunotherapy is a viable
strategy for COVID-19. Cytokine & Growth Factor Reviews. 2020;54:24-31. doi:
https://doi.org/10.1016/j.cytogfr.2020.06.008.
91.   Sadhu S, Dalal R, Dandotiya J, Binayke A, Singh V, Tripathy MR, et al. IL-9 aggravates
SARS-CoV-2 infection and exacerbates associated airway inflammation. Nature
Communications. 2023;14(1):4060. doi: 10.1038/s41467-023-39815-5.
92.   Kral M, van der Vorst EPC, Surnov A, Weber C, Döring Y. ILC2-mediated immune
crosstalk in chronic (vascular) inflammation. Front Immunol. 2023;14:1326440. Epub
2024/01/05. doi: 10.3389/fimmu.2023.1326440. PubMed PMID: 38179045; PubMed
Central PMCID: PMCPMC10765502.
93.   Ghosn L, Chaimani A, Evrenoglou T, Davidson M, Grana C, Schmucker C, et al.
Interleukin-6 blocking agents for treating COVID-19: a living systematic review. Cochrane
Database Syst Rev. 2021;3(3):CD013881. Epub 2021/03/19. doi:
10.1002/14651858.CD013881. PubMed PMID: 33734435.
94.   Patterson BK, Seethamraju H, Dhody K, Corley MJ, Kazempour K, Lalezari J, et al. CCR5
inhibition in critical COVID-19 patients decreases inflammatory cytokines, increases CD8
T-cells, and decreases SARS-CoV2 RNA in plasma by day 14. International Journal of
Infectious Diseases. 2021;103:25-32. doi: https://doi.org/10.1016/j.ijid.2020.10.101.
95.   Pius-Sadowska E, Niedźwiedź A, Kulig P, Baumert B, Sobuś A, Rogińska D, et al. CXCL8,
CCL2, and CMV Seropositivity as New Prognostic Factors for a Severe COVID-19 Course.
International Journal of Molecular Sciences. 2022;23(19):11338. doi:
10.3390/ijms231911338.
96.   Zizzo G, Cohen PL. Imperfect storm: is interleukin-33 the Achilles heel of COVID-19?
Lancet Rheumatol. 2020;2(12):e779-e90. Epub 2020/10/20. doi: 10.1016/S2665-
9913(20)30340-4. PubMed PMID: 33073244; PubMed Central PMCID: PMCPMC7546716.
97.   Chen J, López-Moyado IF, Seo H, Lio C-WJ, Hempleman LJ, Sekiya T, et al. NR4A
transcription factors limit CAR T cell function in solid tumours. Nature.
2019;567(7749):530-4. doi: 10.1038/s41586-019-0985-x.
98.   Odagiu L, May J, Boulet S, Baldwin TA, Labrecque N. Role of the Orphan Nuclear
Receptor NR4A Family in T-Cell Biology. Front Endocrinol (Lausanne). 2020;11:624122.
Epub 2021/02/19. doi: 10.3389/fendo.2020.624122. PubMed PMID: 33597928; PubMed
Central PMCID: PMCPMC7883379.
99.   Phelan DE, Shigemura M, Aldhafiri S, Mota C, Hall TJ, Sznajder JI, et al. Transcriptional
Profiling of Monocytes Deficient in Nuclear Orphan Receptors NR4A2 and NR4A3 Reveals
Distinct Signalling Roles Related to Antigen Presentation and Viral Response. Frontiers in
Immunology. 2021;12:2447. doi: 10.3389/fimmu.2021.676644.
100. Bishop CR, Yan K, Nguyen W, Rawle DJ, Tang B, Larcher T, et al. Microplastics
dysregulate innate immunity in the SARS-CoV-2 infected lung. bioRxiv.
2023:2023.11.19.567745. doi: 10.1101/2023.11.19.567745.

101. Paudyal P, Shrestha S, Madanayake T, Shuster CB, Rohrschneider LR, Rowland A, et al.
Grb7 and Filamin-a associate and are colocalized to cell membrane ruffles upon EGF
stimulation. J Mol Recognit. 2013;26(11):532-41. Epub 2013/10/04. doi: 10.1002/jmr.2297.
PubMed PMID: 24089360; PubMed Central PMCID: PMCPMC4096567.
102. Liu X, Wang X, Xu T, Liu X, Li L. Effects of Different Delivery Modes on the Expression
of Vesicle Transport-Related Genes in Female Pelvic Floor Muscle Repair After Injury.
Appl Biochem Biotechnol. 2023;196(2):667-78. Epub 2023/05/12. doi: 10.1007/s12010-
023-04510-0. PubMed PMID: 37171760.
103. Li S, Wang Z, Chen M, Xiao Y, Min J, Hu M, et al. ArfGAP3 regulates vesicle transport and
glucose uptake in myoblasts. Cell Signal. 2023;103:110551. Epub 2022/12/09. doi:
10.1016/j.cellsig.2022.110551. PubMed PMID: 36476390.
104. Zhang X, Pizzoni A, Hong K, Naim N, Qi C, Korkhov V, et al. CAP1 binds and activates
adenylyl cyclase in mammalian cells. Proceedings of the National Academy of Sciences.
2021;118(24):e2024576118. doi: 10.1073/pnas.2024576118.
105. Ding T, Zhou Y, Long R, Chen C, Zhao J, Cui P, et al. DUSP8 phosphatase: structure,
functions, expression regulation and the role in human diseases. Cell & Bioscience.
2019;9(1):70. doi: 10.1186/s13578-019-0329-4.
106. Schönfelder J, Seibold T, Morawe M, Sroka R, Schneider N, Cai J, et al. Endothelial Protein
kinase D1 is a major regulator of post-traumatic hyperinflammation. Frontiers in
Immunology. 2023;14:1093022. doi: 10.3389/fimmu.2023.1093022.
107. Rivera J, Proia RL, Olivera A. The alliance of sphingosine-1-phosphate and its receptors in
immunity. Nature Reviews Immunology. 2008;8(10):753-63. doi: 10.1038/nri2400.
108. Tsui HS, Pham NVB, Amer BR, Bradley MC, Gosschalk JE, Gallagher-Jones M, et al.
Human COQ10A and COQ10B are distinct lipid-binding START domain proteins required
for coenzyme Q function. J Lipid Res. 2019;60(7):1293-310. Epub 2019/05/03. doi:
10.1194/jlr.M093534. PubMed PMID: 31048406; PubMed Central PMCID:
PMCPMC6602128.
109. Mantle D, Heaton RA, Hargreaves IP. Coenzyme Q10 and Immune Function: An Overview.
Antioxidants. 2021;10(5):759. doi: 10.3390/antiox10050759.
110. Colomer-Molera M, Sastre D, Sole Codina L, Vallejo-Gracia A, Navarro Perez M, Capera-
Aragones J, et al. Kv1.3-dependent immune system activation is regulated by KCNE4.
Biophysical Journal. 2023;122(3):14a. doi: https://doi.org/10.1016/j.bpj.2022.11.302.
111. Li D, Jing J, Dong X, Zhang C, Wang J, Wan X. Activating transcription factor 3: A
potential therapeutic target for inflammatory pulmonary diseases. Immunity, Inflammation
and Disease. 2023;11(9):e1028. doi: https://doi.org/10.1002/iid3.1028.
112. Palecanda A, Paulauskis J, Al-Mutairi E, Imrich A, Qin G, Suzuki H, et al. Role of the
Scavenger Receptor MARCO in Alveolar Macrophage Binding of Unopsonized
Environmental Particles. Journal of Experimental Medicine. 1999;189(9):1497-506. doi:
10.1084/jem.189.9.1497.
113. Tabary M, Gheware A, Penaloza HF, Lee JS. The matricellular protein thrombospondin-1 in
lung inflammation and injury. Am J Physiol Cell Physiol. 2022;323(3):C857-C65. Epub
2022/08/02. doi: 10.1152/ajpcell.00182.2022. PubMed PMID: 35912991; PubMed Central
PMCID: PMCPMC9467471.
114. Schreiner T, Allnoch L, Beythien G, Marek K, Becker K, Schaudien D, et al. SARS-CoV-2
Infection Dysregulates Cilia and Basal Cell Homeostasis in the Respiratory Epithelium of
Hamsters. Int J Mol Sci. 2022;23(9):5124. Epub 2022/05/15. doi: 10.3390/ijms23095124.
PubMed PMID: 35563514; PubMed Central PMCID: PMCPMC9102945.

115. Woodruff MC, Bonham KS, Anam FA, Walker TA, Faliti CE, Ishii Y, et al. Chronic
inflammation, neutrophil activity, and autoreactivity splits long COVID. Nature
Communications. 2023;14(1):4201. doi: 10.1038/s41467-023-40012-7.
116. Cui L, Fang Z, De Souza CM, Lerbs T, Guan Y, Li I, et al. Innate immune cell activation
causes lung fibrosis in a humanized model of long COVID. Proceedings of the National
Academy of Sciences. 2023;120(10):e2217199120. doi: 10.1073/pnas.2217199120.
117. Kellici TF, Pilka ES, Bodkin MJ. Therapeutic Potential of Targeting Plasminogen Activator
Inhibitor-1 in COVID-19. Trends Pharmacol Sci. 2021;42(6):431-3. Epub 2021/04/20. doi:
10.1016/j.tips.2021.03.006. PubMed PMID: 33867130; PubMed Central PMCID:
PMCPMC7997307.
118. Lorè NI, De Lorenzo R, Rancoita PMV, Cugnata F, Agresti A, Benedetti F, et al. CXCL10
levels at hospital admission predict COVID-19 outcome: hierarchical assessment of 53
putative inflammatory biomarkers in an observational study. Molecular Medicine.
2021;27(1):129. doi: 10.1186/s10020-021-00390-4.
119. Kircheis R, Haasbach E, Lueftenegger D, Heyken WT, Ocker M, Planz O. NF-κB Pathway
as a Potential Target for Treatment of Critical Stage COVID-19 Patients. Frontiers in
Immunology. 2020;11:598444. doi: 10.3389/fimmu.2020.598444.
120. Nikkhoo B, Mohammadi M, Hasani S, Sigari N, Borhani A, Ramezani C, et al. Elevated
interleukin (IL)-6 as a predictor of disease severity among Covid-19 patients: a prospective
cohort study. BMC Infectious Diseases. 2023;23(1):311. doi: 10.1186/s12879-023-08294-w.
121. Zangeneh Z, Khamisipour G. Elevated HSP70 and HSP90 as Predictive Markers of Immune
Activation and Lung Injury in SARS-COV-2 Disease. Iranian Journal of Immunology.
2023;20(3):368-73. doi: 10.22034/iji.2023.97202.2488.
122. Prerna J, Swati G, Shailendra M, Kamini J, Manisha M, Rumaisha S, et al. Host inducible-
HSP70A1A is an irresistible drug target to combat SARS-CoV2 infection and pathogenesis.
bioRxiv. 2023:2023.05.05.539661. doi: 10.1101/2023.05.05.539661.
123. Dinnon KH, Leist SR, Okuda K, Dang H, Fritch EJ, Gully KL, et al. SARS-CoV-2 infection
produces chronic pulmonary epithelial and immune cell dysfunction with fibrosis in mice.
Science Translational Medicine. 2022;14(664):eabo5070. doi:
10.1126/scitranslmed.abo5070.
124. Ghonim MA, Boyd DF, Flerlage T, Thomas PG. Pulmonary inflammation and fibroblast
immunoregulation: from bench to bedside. J Clin Invest. 2023;133(17):e170499. Epub
2023/09/01. doi: 10.1172/JCI170499. PubMed PMID: 37655660; PubMed Central PMCID:
PMCPMC10471178.
125. Monticelli LA, Sonnenberg GF, Abt MC, Alenghat T, Ziegler CG, Doering TA, et al. Innate
lymphoid cells promote lung-tissue homeostasis after infection with influenza virus. Nat
Immunol. 2011;12(11):1045-54. Epub 2011/09/29. doi: 10.1031/ni.2131. PubMed PMID:
21946417; PubMed Central PMCID: PMCPMC3320042.
126. Gomes AMC, Farias GB, Dias-Silva M, Laia J, Trombetta AC, Godinho-Santos A, et al.
SARS-CoV2 pneumonia recovery is linked to expansion of innate lymphoid cells type 2
expressing CCR10. European Journal of Immunology. 2021;51(12):3194-201. doi:
https://doi.org/10.1002/eji.202149311.
127. Chen B, Julg B, Mohandas S, Bradfute SB, Force RMPT. Viral persistence, reactivation, and
mechanisms of long COVID. eLife. 2023;12:e86015. doi: 10.7554/eLife.86015.
128. Yin K, Peluso MJ, Luo X, Thomas R, Shin M-G, Neidleman J, et al. Long COVID manifests
with T cell dysregulation, inflammation and an uncoordinated adaptive immune response to
SARS-CoV-2. Nature Immunology. 2024;25:218–25. doi: 10.1038/s41590-023-01724-6.

129. Zhao F, Ma Q, Yue Q, Chen H. SARS-CoV-2 Infection and Lung Regeneration. Clin
Microbiol Rev. 2022;35(2):e0018821. Epub 2022/02/03. doi: 10.1128/cmr.00188-21.
PubMed PMID: 35107300; PubMed Central PMCID: PMCPMC8809385.
130. Santopaolo M, Gregorova M, Hamilton F, Arnold D, Long A, Lacey A, et al. Prolonged T-
cell activation and long COVID symptoms independently associate with severe COVID-19
at 3 months. eLife. 2023;12:e85009. doi: 10.7554/eLife.85009.
131. De Biasi S, Meschiari M, Gibellini L, Bellinazzi C, Borella R, Fidanza L, et al. Marked T
cell activation, senescence, exhaustion and skewing towards TH17 in patients with COVID-
19 pneumonia. Nature Communications. 2020;11(1):3434. doi: 10.1038/s41467-020-17292-
4.
132. Rawle DJ, Hugo LE, Cox AL, Devine GJ, Suhrbier A. Generating prophylactic immunity
against arboviruses in vertebrates and invertebrates. Nature Reviews Immunology (in press).
2024.
133. Foy BH, Sundt TM, Carlson JCT, Aguirre AD, Higgins JM. Human acute inflammatory
recovery is defined by co-regulatory dynamics of white blood cell and platelet populations.
Nat Commun. 2022;13(1):4705. Epub 2022/08/23. doi: 10.1038/s41467-022-32222-2.
PubMed PMID: 35995789; PubMed Central PMCID: PMCPMC9395541 application related
to this work submitted by Mass General Brigham. No authors have any other competing
interests.
134. Doran AC, Yurdagul A, Tabas I. Efferocytosis in health and disease. Nature Reviews
Immunology. 2020;20(4):254-67. doi: 10.1038/s41577-019-0240-6.
135. Proal AD, VanElzakker MB, Aleman S, Bach K, Boribong BP, Buggert M, et al. SARS-
CoV-2 reservoir in post-acute sequelae of COVID-19 (PASC). Nature Immunology.
2023;24(10):1616-27. doi: 10.1038/s41590-023-01601-2.

1038

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Lungs, BA.1 vs. UV-BA.1, day 66 (29 DEGs, q<0.05)

↑ **Plasma cell development**　　↑ **T cell responses**　　↓ **DEGs associated with immune functions**

DEG	log$_2$FC
Igkv3-10	6.89
Ighg2c	6.39
Igkv1-88	5.92
Ighv8-12	5.64
Igkv3-2	5.42
Igkv4-91	5.02
Igkv6-17	4.86
Ighv1-7	4.76
Igkv12-46	4.56
Igkv8-28	4.50
Igkv10-94	3.67
Igkv6-15	3.54
Ighv1-26	3.02
Igkc	2.40

Plasma cells & B cells, immunoglobulins (M156.0)
BTM GSEA
NES= 2.1 q=0.026

MM-GOBP_ Immunoglobulin production
MSigBD GSEA
NES= 3.31 q<0.001

T cell activation (M7.3)
BTM GSEA
NES= 2.06 q=0.028

↓ **T cell exhaustion?**

DEG	log$_2$FC
Nr4a1	-3.12
(Nr4a2	-1.7*)
Nr4a3	-2.75

DEG	log$_2$FC	Function
Prkd1	-0.60	Promotes neutrophil infiltration
S1pr1	-0.62	Regulates immune reponses
Coq10b	-1.03	Aerobic respiration/antioxidant
Kcne4	-1.41	Immune regulator
Atf3	-1.81	Stress-inducible transcription factor
Marco	-1.83	Scavenger receptor
Thbs1	-1.96	Matricellular protein, damage control

DEGs associated with ↑ tissue repair

DEG	log$_2$FC	Function
Grb7	0.56	Cell migration
Arfgap3	0.34	Vesicle Transport
Cap1	0.28	cAMP proliferation
Dusp8	-1.00	MAPK inhibitor

↓ **Stress response DEGs**

DEG	log$_2$FC
ATF3	-1.81
Hspa1b	-3.12

GSEAs
MSigDB
Blood transcription modules

Figure 6

a Genes whose expression levels positively correlated with dpi (Pearson correlations, q<0.05) n= 2786

Enrichr and IPA analyses

↑ **Ciliated epithelial cell**

Enrichr
Descartes Cell Types and Tissue 2021   Ciliated epithelial cells in Lung   q=1.11e-19, Combined score 134
CellMarker Augmented 2021   Ciliated cell:Lung   q=1.08e-13, Combined score 115

IPA Diseases or Functions   Formation of Cilia   p=7.88e-11   z= 2.58

↑ **Protein synthesis**

Enrichr
GO Biological Process 2023 Cytoplasmic Translation. q= 3.26e-11, Combined score 181

59 ribosomal protein genes (RPL)

b Genes whose expression levels negatively correlated with dpi (Pearson correlations, q<0.05) n=1996

IPA analysis

↓ **Cell movement/migration**   ↓ **Cytokine responses**   **Cytokine storm, fibrosis**
                                                                    ↓ **and neutrophil degranulation**

IPA Diseases or Functions   **IPA Cytokine USRs**   **IPA Canonical (top examples)**
(top annotations)   **(top annotations)**   Neutrophil degranulation $-\log_{10}$p=19.3   z= -9.59
Cell movement   p=1.75e-85 z= -14.71   TNF p=7.3e-73   z= -13.17   Pathogen Induced Cytokine
Migration of cells p=3.11e-85 z= -14.16 IFNG p=1.4e-83 z= -12.42   Storm Signaling Pathway $-\log_{10}$p=22.6   z= -8.79
                                        IL1B p=7.7e-54   z= -9.4   Pulmonary Fibrosis Idiopathic
                                        CSF2 p=2.5e-52   z= -9.33   Signaling pathway $-\log_{10}$p=17.9   z= -8.3

c

| Serpine1 | Cxcl10 | Nfkb1 | IL6 | Hspa1b |
| q=1.33e-4 r= -0.89 | q=2.72e-3 r= -0.75 | q=2.52e-3 r= -0.75 | q=3.1e-2 r= -0.6 | q=4.92e-2 r= -0.56 |

Normalized VST read counts

Days post infection

d Cell types whose abundance scores positively correlated with dpi (Pearson correlations, q<0.05)

| Ciliated cell (MCA) | AT1 cell (MCA) | Fibroblastic reticular (NS) | ILC, innate lymphoid cell (NS) |
| q = 9.16e-05 r = 0.82 | q = 2.66e-02 r = 0.55 | q = 0.018 r = 0.63 | q = 0.018 r = 0.63 |

Abundance score

Days post infection