The molecular basis of pine wilt disease resistance in *Pinus massoniana*

Jinjin Wang[1][2][†], Shouping Cai[1][†], Liqiong Zeng[1], Jie Tong[3], Danling Hang[4], Xinliang Zhang[5], Yu

Fang[6], Jiayu Jia[2], Chenli Li[2], Shunde Su[1][*], Jun Su[2][*]

1 Fujian Academy of Forestry, Key Laboratory of National Forestry and Grassland Administration on
Timber Forest Breeding and Cultivation for Mountainous Areas in Southern China, Fuzhou, China

2 Basic Forestry and Proteomics Research Center, Fujian Provincial Key Laboratory of Haixia Applied
Plant Systems Biology, Fujian Agriculture and Forestry University, Fuzhou, China

3 Silviculture Department, Shaxian Shuinan State-Owned Forest Farm, Sanming, China

4 Silviculture Department, Shaxian Guanzhuang State-Owned Forest Farm, Sanming, China

5 Forest pest control and quarantine station, Sanming, China

6 Institute of Soil Fertilizer, Fujian Academy of Agricultural Sciences, Fuzhou, China

[†] These authors have contributed equally to this work and share first authorship.

[*] **Correspondence:**

Jun Su

junsu@fafu.edu.cn

Shunde Su

ssdforest@163.com

22 Number of words: 4333

23 Number of figures: 5

Keywords: Keywords: Pine wilt disease, Pine wood nematode (*Bursaphelenchus xylophilus*), *Pinus massoniana*, **resistant and tolerance strategy, multi-omics**

Abstract

Pine wilt disease (PWD), caused by the pine wood nematode (PWN) *Bursaphelenchus xylophilus*, results in significant economic and ecological damage to *Pinus* forests and plantations worldwide. *Pinus massoniana* is the primary host of PWD in southern China, but its response to the PWN remains largely unstudied. Previously, we observed PWD in a *P. massoniana* nursery that contained over 71 commonly used cultivars. Through field phenotyping, we identified two groups of PWN resistant cultivars. Resistant cultivars (RC) exhibited very low PWN carrying amounts (PCA) and had relatively low mortality, and tolerant cultivars (TC) had high PCA but low mortality. In this study, we confirmed via PWN inoculation assays that the resistant and tolerant cultivars had lower mortality rates, 10% and 11%, respectively, than other cultivars (which had a mortality rate of 83.3%). The RC had a PCA that was significantly lower than that of other cultivars, while the TC exhibited a higher PCA. To explore the molecular mechanisms underlying the response of *P. massoniana* to PWN, the transcriptome and metabolome of the above cultivars were profiled by high throughput sequencing. As no reference genome is available for *P. massoniana*, we generated a new full-length transcriptome library with iso-seq. Using the transcriptome and metabolome from different *P. massoniana* cultivars inoculated with PWN, we found three major PWD resistance strategies. 1) The common response strategy involved three important molecular pathways. First, synthesis of indole-3-acetic acid (IAA) and abcisic acid (ABA) were suppressed during PWN invasion, thus suppressing the synthesis of polysaccharides (especially myo-inositol and trehalose) and (-)-riboflavin through ABC transporters. Second, by inhibiting aspartic acid, the synthesis of arginine and proline through APS5 (aspartate aminotransferase 5) were suppressed. Third, by reducing cysteine, GERD (germacrene D synthase) was up-regulated, and sesquiterpenoid and triterpenoid metabolites were accumulated to resist PWN invasion. 2) The most effective resistance strategy involved the accumulation of reactive oxygen species (ROS), which enhanced jasmonic acid (JA) accumulation and highly induced the expression of chitinase, thus improving resistance to PWN. 3) The tolerance strategy involved the induction of phosphatidylcholine, which promoted flavonoid and anthocyanin synthesis via LPIN (phosphatidate phosphatase), LOX2S (lipoxygenase), and LOX1_5 (linoleate 9S-lipoxygenase), thus significantly inhibiting the pathogenicity of the PWN but not the PCA. These results illustrate the molecular mechanisms by which *P. massoniana* cultivars resist or tolerate PWN and are of great significance in the prevention and control of PWD.

1   Introduction

Pine wilt disease (PWD) is one of the most destructive diseases in *Pinus* forests worldwide and poses a serious threat to the health of forest ecosystems in China (Xie et al., 2017; Li et al., 2021; Zhang et al., 2023). *Pinus massoniana* is severely affected by PWD in southern China. Since the disease was first discovered, over 5,000,000 m ³of *Pinus* forests have been infected (Cai et al., 2021; An et al., 2023; Li et al., 2023). Due to its hardiness and low maintenance costs, *P. massoniana* is the main cultivated variety of *Pinus* in impoverished areas of southern China, and it is irreplaceable (Yu et al., 2020; Zhu et al., 2022). However, due to the widespread PWD epidemic (Hopf-Biziks et al., 2017; Pajares et al., 2017), planting *P. massoniana* has become a major challenge, and the selection of resistant cultivars is critical to solving this problem.

Cultivating pine wood nematode (PWN) resistant cultivars is currently the most economical and effective approach to reduce PWD and ensure the health of *Pinus* forests (Liu et al., 2022; Hu et al.,

2023). Through genetic engineering, traditional breeding, and molecular marker-assisted selection, cultivars can be screened and planted (Li et al., 2010; Török et al., 2019; Thulasinathan et al., 2023). This limits the spread of PWD and enhances the ecological stability, biodiversity, and sustainable development of pine forest ecosystems. In recent years, a number of PWD resistant cultivars have been developed domestically and internationally, such as GD5, DAC12, and AH1 (Deng et al., 2024; Gao et al., 2024). Foreign cultivars such as Clone number 227, Tanabe 54, and Tosashimizu 63 have also exhibited PWD resistance (Hirao et al., 2019; Guo et al., 2023). However, the above cultivars were naturally selected, and their effectiveness in the field is unstable. Genetic engineering can be used to screen resistant cultivars, improve their stability in field production, and significantly reduce the time required for traditional breeding (Li et al., 2010; Mackelprang and Lemaux, 2020; Rahman et al., 2023). However, the use genetic engineering to improve the economic and ecological traits of *Pinus* spp. remains limited. An improved understanding of cultivar resistance and the breeding of stable and effective cultivars through genetic engineering are crucial for the management of PWD.

Systematic analysis of the molecular mechanisms of PWN pathogenicity and the selection of resistance genes are prerequisites for genetic engineering and lay the foundation for effective proactive control methods and comprehensive prevention strategies (Bláhová et al., 2020; Mackelprang and Lemaux, 2020). Previous research and empirical evidence have revealed that PWN pathogenicity begins with its proliferation in the xylem of pine trees, which causes mechanical damage (Ichihara et al., 2000; Umebayashi et al., 2011; Yazaki et al., 2018). Additionally, PWN is often accompanied by other pathogens (such as fungi), and co-infection exacerbates the damage to pine trees (Xue et al., 2019; Jia et al., 2023). Proteases and polysaccharide enzymes are secreted by PWN, and these toxins disrupt cellular structure, interfere with normal physiological functions, and lead to tree death (Zhao et al., 2007; Faria et al., 2015). Scientists have explored resistance by *P. massoniana* and revealed several key genes related to PWN resistance. Among them, *PmNBS-LRR97*, *PmGPPS1*, *PmTPS4*, and *PmTPS21* participate in defense against PWN (Xie et al., 2023; Liu et al., 2021; Liu et al., 2022). These results indicate that the related resistance genes have significant advantages in long-term pest control. However, the lack of reference genome for *P. massoniana* and a focus on single genes or single-level interactions can inhibit an overall understanding of multidimensional and multivariate relationships, reducing effectiveness in the complex forest environment over the long term.

Previously, we analyzed the spectral performance in the field of the germplasm resource library concurrent with PWN carrying capacity to identify resistant cultivars (RC) and tolerant cultivars (TC) (Liu et al., 2023a). Building on this, we constructed a full-length transcriptome library for *P. massoniana* through iso-seq, analyzed the transcriptomes and metabolomes of the RC and TC, and explored the molecular basis of *P. massoniana* resistance to PWN. An in-depth exploration of the biochemical interactions between different cultivars and PWN enabled us to clarify the strategies used by *P. massoniana* in response to PWN invasion and to determine their molecular basis. Our findings are of great significance for understanding PWD transmission and control.

2   Materials and Methods

2.1 Plant material and growth conditions

The experimental subjects were taken from the nursery at Guanzhuang State-owned Forest Farm in Sanming, Fujian, China (26.5603°N, 117.7455°E), A total of 71 clonal interforest cultivars were obtained. The study area comprised a 76 ha, 16-year-old pure *P. massoniana* plantation with an annual average temperature of 19.9 ℃ and precipitation of 1375.2 mm. In the early stage, three types of clones were screened based on spectral characteristics and PWN carrying amounts (PCA): tolerant cultivars

(TC) (high PCA, spectral data biased towards health, low mortality), resistant cultivars (RC) (extremely
low PCA, spectral data biased towards health, low mortality) and neutral cultivars (NC) (high PCA,
spectral data biased towards death, high mortality) (Liu et al., 2023a). Eight clones (TC numbers 43,
45, and 58; RC numbers 52 and 59; and NC numbers 64, 41, and 51) were measured in the laboratory.

In the above *P. massoniana* cultivars, PCA and mortality were quantified 6 weeks after inoculation
with PWN. Cultivars were placed in a growth chamber (16 h light, 8 h dark, 70% humidity, 28 ℃ in
the light, and 24 ℃ in the dark) for 2 months before experimental treatment.

2.2 Pine wood nematode inoculation assay

123 Pine wood nematode adults were cultured as previously described (Cai et al., 2021). Seedlings of *P.*
124 *massoniana* were inoculated with 1 mL of a PWN solution (5000 individuals / mL), with 15 replicates
125 (individual plants) per cultivar. Double-distilled water served as a negative control.

2.3 Quantification of relative PWN levels

Real-time quantitative polymerase chain reaction (RT-qPCR) was used to quantify PWN in *P.*
massoniana cultivars. Fresh branches (with leaves) collected from the entire young seedling from the
laboratory, were used for PWN quantification. Total genomic DNA was extracted from each plant
sample using a MoBio PowerPlant® Pro DNA Isolation Kit (cat no.12855-50, MoBio, USA),
according to the manufacturer's protocol. DNA quantity and quality were measured with a NanoDrop
2000 photometer (Thermo Fisher Scientific, USA), and DNA integrity was determined by 1% agarose
gel electrophoresis. The extracted DNA was stored at -80 ℃ until further use. Quantitative PCR was
conducted using a PWN-specific BxCW primer (F: 5 '-TTGCATTCTACGGCCAGTCC-3'; R: 5′-
ACTGACTTTCGATGGCTCCG-3') and a host-specific PmEF2 primer (F: 5'-
CTGCGATGTCCCTCATGTTA-3'; R: 5′- AACAAGGTCTTTCCCCTCGT-3') (one transcript from
P. massoniana was used as the internal control) (Xie et al., 2023), with Hieff™ qPCR SYBR Green
Master Mix (Low Rox Plus, cat no. 11202ES08, Yeasen, Shanghai, China) on a QuantStudio 6 Flex
PCR system (ABI). The qPCR signals were normalized to those of the reference gene PST in *Pinus* by
applying the $2^{-\Delta\Delta CT}$ method (Fang et al., 2023). Three biological replicates and three technical replicates
were used. One-way analysis of variance (ANOVA; Tukey test) was used to determine differences
among groups.

2.4 Generation of full-length reference transcripts

The sample preparation, as described in section 2.1, was repeated using three techniques with six weeks
after infection by PWN. Total RNA was extracted from the cultivars using TRIzol (Invitrogen, CA,
USA), eukaryotic mRNA was enriched with magnetic beads with Oligo(dT), and mRNA was reverse
transcribed into cDNA using a SMARTer™ PCR cDNA Synthesis Kit. BluePippin was used to screen
full-length cDNA fragments (Sage Science, Beverly, MA, USA) to generate cDNA libraries of
different lengths (1–2 k, 2–3 k, and 3–6 k). The SMRT bell library was obtained by terminal repair of
full-length cDNA and a SMRT dumbbell connector. The libraries were then sequenced via a PacBio
RS II platform (Menlo Park, CA, USA). Trimmomatic (v0.33) was used for quality control and filtering
of raw reads. Low quality reads and those with short read lengths were removed. Nonchimeric full-
length sequences were identified from reads of inserts based on the poly (A) tail signals, 5′ adapter
sequences, and 3′ adapter sequences. An isoform-level clustering algorithm was used to iteratively
cluster the full-length transcript sequence (Chin et al., 2013). In combination with non-full-length
sequences, consistent sequences were corrected using a corrective (polish) procedure. Finally,

sequence redundancy was removed using CD-HIT (http://weizhongli-lab.org/cd-hit/) (Li and Godzik,
2006).

2.5 RNA-seq and analysis

The sample preparation, as described in section 2.4. The Illumina cDNA library was prepared using a
NEBNext Ultra II RNA Library Prep Kit (NEB, Ipswich, MA, USA) according to the standard
experimental protocol. An Illumina Novaseq 6000 with a PE 150 sequencing strategy was used for
computer sequencing. Read segments sequenced by Illumina were mapped to SMRT cDNA libraries
by Bowtie2 (Langdon, 2015) and quantified by RSEM (Li et al., 2015) using fragments per million
mapped read segments (FPKM) per kilobase transcript. The Non-Redundant Protein Sequence
Database (NR) (ftp://ftp.ncbi.nlm.nih.gov/blast/db/), Gene Ontology (GO) database
(http://geneontology.org/), Swiss-Prot Protein Sequence Database (SwissProt)
(http://www.ebi.ac.uk/uniprot), clusters of euKaryotic Orthologous Groups (KOG) tool
(ftp://ftp.ncbi.nlm.nih.gov/pub/COG/), and the Kyoto Encyclopedia of Genes and Genomes (KEGG)
database (http://www.genome.jp/kegg/) were used for analyses. DESeq (1.10.1) (Love et al., 2014)
was used for differential expression analysis, and R software was used for correlation analysis and
drawing heat maps.

2.6 Metabolomic analysis

Metabolites were extracted from each sample as previously described (De Vos et al., 2007). Ultra high
performance liquid chromatography (UHPLC) separation was performed using an ACQUITY UPLC
BEH amide column (1.7 μm, 2.1 mm × 100 mm, 186004801, Waters, Milford, USA) equipped by
Oebiotech Company (Shanghai, China). The flow rate of the mobile phase was set at 0.5 mL / min and
consisted of a 0.1% formic acid solution (A) (A117-50, Thermo Fisher Scientific, Waltham, MA, USA)
and a 0.1% formic acid in acetonitrile (B) (A998-4, Thermo Fisher Scientific, Waltham, MA, USA).
The column temperature was 25 ℃, the auto-sampler temperature was 4 ℃, and the sample volume
was 1 μL (Chen et al., 2007; Theodoridis et al., 2008). Data were pruned from different samples to
distinguish the metabolites in different resistant cultivars of *P. massoniana*. Next, commercial
databases, including the KEGG database and MetaboAnalyst (https://www.kegg.jp/), were used with
the search term "metabolite pathway" (https://www.genome.jp/kegg/pathway.html).

3   Results

3.1 Nomination of potential PWN-determining genes in *P. massoniana* using a full-length transcriptome

To uncover the pathogenic mechanisms of the PWN in its host pine species, we sequenced the
transcriptome of *P. massoniana* and screened for potential PWN-determining genes (β) by comparing
protein information from *Pinus densiflora* and *Pinus tabuliformis* and predicting the function of β. The
sequencing process involved extracting unigenes from raw data through steps that included filtering,
quality control, classification, clustering, correction (polishing), and redundancy removal, leading to
the prediction of 55,724 coding DNA sequences (**Figure 1A**). The integrity of the transcript data across
three generations was evaluated using BUSCO analysis, which revealed that 93.5% of the isoforms
were complete (**Supplementary Figure S1**). Furthermore, 52,731 (96.63%) genes were annotated in
at least one database (KOG, NR, KEGG, or Swissprot, among others) (**Figure 1B**). To identify
potential PWN-determining genes in *P. massoniana*, 7,497 genes (α) shared between *P. massoniana*
and *P. densiflora*, both of which host PWN, were screened for amino acid sequence similarity greater
than 90%. Subsequently, α was compared with a non-PWN host, *P. tabuliformis*, for amino acid

sequence similarity less than 90%, which resulted in the identification of 1,938 potential PWN-determining genes (β) in *P. massoniana* (**Figure 1C**). KEGG analysis of β in *P. massoniana* revealed their predominant enrichment in MAPK signaling and plant-pathogen interaction pathways. (**Figure 1D**).

3.2 Transcriptome analysis revels PWN carrier determinant genes in *P. massoniana*

Previously, the TC, NC, and RC were identified within the germplasm bank of *P. massoniana*. In this research, all three cultivars were uniformly inoculated with PWN, and the PCA were quantified six weeks post-inoculation. We found that the PCA in the TC was significantly higher than that in the NC, which in turn exhibited a higher PCA than the RC ($P \leq 0.01$). Specifically, the PCA in the TC and RC were 1.68 and 0.34 times that of NC, respectively (**Figure 2A**). Concurrently, we identified genes that were strongly associated with the PCA ($P \leq 0.01$, $|r| \geq 0.8$), resulting in a total of 7,710 genes (**Supplementary Table S1**). These genes were then compared with β to determine the PWN carrier determinant genes (γ) in *P. massoniana*, which amounted to 318 genes (**Figure 2B**). In the TC vs NC and the RC vs NC, 128 and 63 genes were upregulated, respectively, and 116 and 200 genes were downregulated, respectively (**Figure 2C**), indicating that spatial variation in the expression of related genes occurred during PWN invasion of the TC and RC. Furthermore, KEGG functional enrichment analysis was performed to predict the function and pathway of γ and revealed significant enrichment in MAPK signaling and plant-pathogen interaction pathways (**Figure 2D**).

3.3 Molecular strategies of *P. massoniana* in response to PWN invasion

Interforest spectral data previously revealed three distinct types of *P. massoniana* with varying health (Liu et al., 2023a). Consequently, we evaluated the mortality rates of these host plants six weeks after inoculation with PWN. The mortality rate of the *P. massoniana* NC was significantly higher than that of the TC or the RC ($P \leq 0.01$); mortality was 7.4 times or 6.6 times higher in the NC, respectively (**Figure 3A**). For the PCA of each cultivar (**Figure 2A**), we found that the TC exhibited a high PCA yet maintained a low mortality rate, whereas the RC demonstrated both low mortality and PCA. We speculated that these were two different PWN resistant cultivars. After full-length transcriptome sequencing, we performed second-generation transcriptome sequencing for the three cultivars six weeks after PWN inoculation to assess gene expression in each cultivar and ascertain the molecular mechanisms underpinning resistance to PWN by the TC and RC. We identified 274 common related genes (CRGs), 243 tolerance-related genes (TRGs), and 227 resistance-related genes (RRGs) across the β in theTC vs NC, and RC vs NC comparison groups (**Figure 3B**). Concurrently, we evaluated the relative expression of the three gene categories in each cultivar (**Figure 3C**). Among them, 49% of TRGs (118 of 243) exhibited higher relative expression in the TC compared to the other two cultivars. The relative expression of RRGs in the RC was higher in 59% (161 of 274) and 71% (161 of 227) than in the other two cultivars. Additionally, KEGG enrichment analysis of CRGs, TRGs, and RRGs revealed enrichment not only in linoleic acid metabolism, MAPK signaling, and plant-pathogen interaction pathways but also in other pathways. Specifically, TRGs were enriched in the spliceosome pathway (**Figure 3D**), CRGs were enriched in ABC transporter and starch and sucrose metabolism pathways (**Figure 3E**), and RRGs were enriched in ABC transporter pathways (**Figure 3F**).

3.4 *P. massoniana* metabolic responses to PWN invasion

Identifying downstream metabolites is essential to understanding *P. massoniana* resistance to PWN. Non-targeted metabolomic sequencing was performed on three cultivars six weeks after PWN

inoculation. Metabolites that were highly correlated with PCA were selected in different cultivars (P
$\leq$ 0.01, |r| $\geq$ 0.8), resulting in the identification of 465 metabolites (δ) (**Figure 4A**). Among these, 145
metabolites were positively correlated with PWN ($P \leq$ 0.01, $1 \geq r \geq$ 0.8), while 320 were negatively
correlated with PWN ($P \leq$ 0.01, $1 \leq r \leq$ 0.8) (**Figure 4A**). KEGG enrichment analysis indicated that δ
were significantly enriched in the arachidonic acid, arginine, and proline metabolism pathways (**Figure
4B**). Moreover, we analyzed the differential accumulation of metabolites in the TC vs NC and RC vs
NC and identified a total of 302 common related metabolites (CRMs), 531 tolerance-related
metabolites (TRMs), and 222 resistance-related metabolites (RRMs) (**Figure 4C**). The expression of
each type of metabolite in each cultivar was calculated (**Supplementary Figure S2**). KEGG
enrichment analysis of CRMs, TRMs, and RRMs showed that TRMs were primarily enriched in purine
metabolism, flavone and flavonol biosynthesis, and ABC transporter pathways (**Figure 4D**). CRMs
were enriched in phenylpropanoid biosynthesis, purine metabolism, and alkaloid biosynthesis
pathways (**Figure 4E**). RRMs were enriched in ABC transporter and metabolic pathways (**Figure 4F**).

3.5 The genetic landscape of *P. massoniana* following PWN invasion

Integrating transcriptomic and metabolomic analyses can shed light on plant metabolic pathways.
Differentially expressed genes (DEGs) and differentially accumulated metabolites (DAMs) may
reveal the mechanisms behind plant resistance to pathogens. Our analysis of the transcriptomes and
metabolomes of *P. massoniana* cultivars revealed genes and metabolites within the same metabolic
pathways (**Supplementary Figure S3**). Based on the relative contents of pertinent genes and
metabolites (**Figure 5A**), we identified three strategies by which *P. massoniana* responds to PWN
invasion (**Figure 5B**). Initially, the PWN triggers a common response in all cultivars, which
comprises the inhibition of endogenous hormone synthesis (IAA, ABA, aspartic acid, and cysteine),
the downregulation of ABCB1 and ASP5 expression, and the upregulation of ABCC1, ABCC2, and
germacrene D synthase expression. Alterations in metabolites, including L-arginine, glutamic acid,
myo-inositol, (-)-riboflavin, trehalose, and aspartic acid, ultimately restrict the reproductive and
pathogenic processes of the PWN. In contrast to the NC, the RC relies on specialized mechanisms to
curtail PWN reproduction. Upon PWN invasion, the RC initially promotes the accumulation of
linoleic acid and induces the synthesis of JA through AOC, AOC3, and LOX2S (lipoxygenase)
proteins, thereby regulating downstream genes (EGL3, MYC2) and metabolites. Pine wood
nematode promotes the accumulation of ROS and the synthesis of chitinase but also limits the
amount of PWN carried by *P. massoniana*, which protects the tree from PWD. The TC also employs
special mechanisms to limit PWN pathogenicity. Initially, PWN invasion triggers the accumulation
of DPPC (phosphatidylcholine), which, under the action of LPIN (phosphatidate phosphatase),
stimulates the synthesis of linolenic acid. This regulates the production of JA through LOX1_5
(linoleate 9S-lipoxygenase) and LOX2S proteins. Jasmonic acid then promotes the synthesis of
downstream flavonoid metabolites (neohesperidin dihydrochalcone, flavonol base, flavone base,
DL−phenylalanine, and neobavaisoflavone), which limits the pathogenicity of the PWN and protects
P. massoniana from PWD.

4   Discussion

As the primary host for PWN in southern China, it is important to understand how *P. massoniana*
responds to PWN invasion. Indoor tests confirmed that PCA and mortality were low in the RC that
was screened from the forest, while the TC had a high PCA but low mortality. On this basis, we
explored broad-spectrum resistance to PWN and the molecular basis for resistance and tolerance in *P.*
massoniana via transcriptomic and metabolomic analyses (**Figure 5B**).

Initially, we conducted RNA-seq on three cultivars using the reported full-length transcriptome of *P.*
massoniana as a reference, but the mapping rate for the second-generation transcriptome was low (Liu
et al., 2020). Therefore, we performed full-length transcriptome sequencing on 71 varieties in the seed
orchard and obtained 55,724 high-quality transcripts with an average length of 3,343.56 bp, and 93.5%
of the transcripts were annotated in the database. Using this library as a reference genome, the average
splicing ratio of the second-generation transcriptome sequencing results reached 94.73%. Previously,
81,837 high-quality transcripts with an average length of 846.54 bp were reported, and the average
splicing ratio for second-generation transcriptome sequencing was 85.02% (Liu et al., 2020). The full-
length transcriptome sequencing results in this study were superior to previous data in several respects,
with the exception of the number of transcripts. However, considering the average length of transcripts,
the volume of data in this study is larger than that for reference transcripts in previous studies.
Additionally, not all pine trees are hosts for the PWN; therefore, we screened for potential PWN
interaction-related genes in *Pinus* by using comparative genomics. According to genome data for *P.*
densiflora (another PWN host) and *P. tabuliformis* (a non-host), 1,938 common potential PWN-
determining genes (β) differed significantly in host trees relative to non-host trees. These were defined
as potential PWN-host pine tree interaction proteins and were enriched in the MAPK signaling and
plant-pathogen interaction pathways. The MAPK signaling pathway is involved in two major barriers,
pathogen-associated molecular pattern (PAMP)-triggered immunity and effector-triggered immunity,
which help plants resist microbial invasion. By transmitting and amplifying external signals, specific
defense genes are activated and promote plant resistance (Manna et al., 2023; Peng et al., 2023). The
plant-pathogen interaction pathway involves the recognition of specific molecular patterns that trigger
immune responses and prevent microbial invasion and spread (Kunkel and Johnson, 2021; Rubio-
Somoza and Blázquez, 2023). These findings suggest that the formation of PWN hosts is related to
associated microbes.

In field experiments, over 87% of *P. massoniana* cultivars produced a common response but were
unable to restrict the proliferation or pathogenesis of the PWN, which corroborates observations of
severe, widespread PWD in the field. The common response of *P. massoniana* included the ABC
transporter, arginine, and proline pathways. First, ABC transporters were involved in the transport of
secondary metabolites and plant hormones, detoxification, and signal transduction, which indirectly
aided resistance to PWN (Wang et al., 2022; Li et al., 2024; Zhou et al., 2024). Second, arginine and
proline indirectly affected the reproduction and pathogenicity of the PWN by promoting plant growth
and development, enhancing stress resistance, acting as antioxidant defense molecules, participating in
defense responses, and regulating metabolic pathways (Zhang et al., 2022; Theisen et al., 2024; Wang
et al., 2024b). The specific mechanisms by which ABC transporters, arginine, and proline function in
the resistance of *P. massoniana* to PWN require further study. In the common response of *P.*
massoniana, monoterpenes and sesquiterpenes had multiple roles as chemical defense substances. First,
they had a direct toxic effect on the PWN, inhibiting its growth and reproduction by disrupting its
nervous system or metabolic pathways (Hwang et al., 2021, 2022; Sánchez-Martínez et al., 2021).
Second, these substances can act as signaling molecules that induce plants to produce defense enzymes
and secondary metabolites that enhance resistance (Castorina et al., 2020; Paul et al., 2020; Xu et al.,
2021). The antioxidant effects of monoterpenes and sesquiterpenes also alleviate oxidative stress
caused by PWN infection (Farhadi et al., 2020; Son et al., 2023). However, the common response is
insufficient in limiting the PWN, possibly because it may detoxify secondary metabolites (including
terpenoid compounds) (Jia et al., 2023). In future studies, adding monoterpenes and sesquiterpenes to
the diet of the PWN and observing mortality may indicate whether the PWN has a detoxification system.

In this study, chitinase, the JA signaling pathway, and related DEGs and DAMs involved in
maintaining ROS homeostasis were enriched in the RC and served as the main effector signaling

pathways for defending against the PWN (**Figure 5A**). Stress may disrupt ROS homeostasis, leading to ROS accumulation, which in turn promotes the accumulation of JA (Wang et al., 2019; Furuta et al., 2024; Shi et al., 2024) and upregulates the expression of chitinase, ultimately affecting the PCA (Shen et al., 2022; Qiu et al., 2023). The role of chitinase in protecting plants from stress has been well established (Ju et al., 2016; Berini et al., 2018; Jha and Modi, 2018; Lu et al., 2020); chitinase has been shown to degrade chitin in nematode eggshells, which limits nematode reproduction (Chan et al., 2010; Lee and Kim, 2015; Chen et al., 2021). This strategy has been widely applied in the control of agricultural nematodes. Therefore, we speculate that in the RC, chitinase expression is the main mode of action of resistance (**Figure 5B**). In future studies, verifying the direct effect of chitinase on the PWN will allow us to ascertain its importance in defense and its potential as a target for new eco-friendly nematicides (Zhao et al., 2023).

Flavonoids play a key role in tree defense against the PWN through signal transduction and metabolic regulation (Isshiki et al., 2014; Vazquez-Vilar et al., 2023; Wang et al., 2024a). Following PWN invasion, the host recognizes PWN secretions and related microbial products, activates defense signaling pathways, and induces the expression of flavonoids (Park et al., 2020). These substances have antioxidant and antimicrobial functions that limit PWN pathogenicity and microbial infection (Raorane et al., 2019; Liu et al., 2023b). Pine wilt nematode pathogenicity involves numerous symbiotic or associated microbes (Jia et al., 2023). Flavonoids affect the environment for nematodes by inhibiting the growth of certain pathogenic microbes or regulating the structure of beneficial microbial communities (Borah et al., 2018). Therefore, the role of flavonoids in PWN invasion is not limited to direct defense but also involves regulating complex interactions between the host plant and microbes. We found that the tolerance strategy promoted synthesis of the JA substrate linolenic acid, which increased the accumulation of flavonoids (**Figure 5B**). The TC had the highest flavonoid content (**Figure 5A**), suggesting that it regulates PWN-associated microbes by accumulating flavonoids and limiting PWD, which may be the reason for low mortality in the TC (**Figure 3A**).

The molecular strategies of all three varieties of *P. massoniana* cultivars involved the JA signaling pathway. However, due to the unique genetic backgrounds of *P. massoniana* cultivars, there were significant differences in the composition and regulation of the JA signaling pathway. The RC may possess more sensitive JA receptors or more effective signal transduction components, thereby exhibiting higher resistance to PWN invasion (**Figure 1A**). It is worth noting that the JA signaling pathway does not exist in isolation but interacts with other plant hormones (such as auxin and abscisic acid) and environmental signals (such as light and temperature) to form a complex regulatory network (Qiu et al., 2022; Hu et al., 2023b; Sybilska and Daszkowska-Golec, 2023). Different pines may exhibit different regulatory interactions that affect their defense mechanisms. Over the long term, different pines may have developed specific defense mechanisms under different environmental conditions (Kalachova et al., 2023). Epigenetic factors, such as DNA methylation and histone modification, may also affect the activation and expression of the JA signaling pathway, thereby leading to differences in tree defense (Campos-Rivero et al., 2017; Du et al., 2024). Subsequent studies can investigate the impact of different exogenous signals (such as plant hormones or environmental changes) on *P. massoniana* resistance in the field. By exploring the interactions between different signaling pathways, we can better understand *P. massoniana* defense in the natural environment and provide a scientific basis for the management of PWD.

In summary, transcriptome and metabolome analyses provided new insights into the genetic diversity and genetic basis of pine tree resistance to PWN invasion. Our results clarified the mechanisms by which the RC restricted PWN reproduction through chitinase synthesis and the TC protected plant health via flavonoid accumulation. Future research on molecular interactions between different

resistant cultivars and the PWN, the effects of chitinase and flavonoids, and the mechanisms of the
response process will further expand our understanding of PWD and its prevention and control.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

JS designed and managed the research. JW and SC wrote the manuscript. LZ and JT performed all
analyses and drawings. DH, XZ and TF conducted the field test and sampling. SS, JJ and CL
designed the chemical treatment related experiments.

7   Acknowledgments

We are very grateful to the Fujian Forestry Bureau for its help providing detailed information of
sample areas in this study.

8   Reference styles

An, Y., Li, Y., Ma, L., Li, D., Zhang, W., Feng, Y., et al. (2023). Transcriptomic response of Pinus
massoniana to infection stress from the pine wood nematode Bursaphelenchus xylophilus.
Stress Biology 3, 50. doi: 10.1007/s44154-023-00131-z

Berini, F., Katz, C., Gruzdev, N., Casartelli, M., Tettamanti, G., and Marinelli, F. (2018). Microbial
and viral chitinases: Attractive biopesticides for integrated pest management. Biotechnology
Advances 36, 818–838. doi: 10.1016/j.biotechadv.2018.01.002

Bláhová, Z., Harvey, T. N., Pšenička, M., and Mráz, J. (2020). Assessment of Fatty Acid Desaturase
(Fads2) Structure-Function Properties in Fish in the Context of Environmental Adaptations and
as a Target for Genetic Engineering. Biomolecules 10, 206. doi: 10.3390/biom10020206

Borah, B., Ahmed, R., Hussain, M., Phukon, P., Wann, S. B., Sarmah, D. Kr., et al. (2018). Suppression
of root-knot disease in Pogostemon cablin caused by Meloidogyne incognita in a rhizobacteria
mediated activation of phenylpropanoid pathway. Biological Control 119, 43–50. doi:
10.1016/j.biocontrol.2018.01.003

Cai, S., Jia, J., He, C., Zeng, L., Fang, Y., Qiu, G., et al. (2021). Multi-Omics of Pine Wood Nematode
Pathogenicity Associated With Culturable Associated Microbiota Through an Artificial
Assembly Approach. Front Plant Sci 12, 798539. doi: 10.3389/fpls.2021.798539

Campos-Rivero, G., Osorio-Montalvo, P., Sánchez-Borges, R., Us-Camas, R., Duarte-Aké, F., and De-
la-Peña, C. (2017). Plant hormone signaling in flowering: An epigenetic point of view. Journal
of Plant Physiology 214, 16–27. doi: 10.1016/j.jplph.2017.03.018

Castorina, G., Grassi, F., Consonni, G., Vitalini, S., Oberti, R., Calcante, A., et al. (2020).
Characterization of the Biogenic Volatile Organic Compounds (BVOCs) and Analysis of the
PR1 Molecular Marker in Vitis vinifera L. Inoculated with the Nematode Xiphinema index.
International Journal of Molecular Sciences 21, 4485. doi: 10.3390/ijms21124485

Chan, Y.-L., Cai, D., Taylor, P. W. J., Chan, M.-T., and Yeh, K. W. (2010). Adverse effect of the
chitinolytic enzyme PjCHI-1 in transgenic tomato on egg mass production and embryonic
development of Meloidogyne incognita. Plant Pathology 59, 922–930. doi: 10.1111/j.1365-
3059.2010.02314.x

Chen, C., Gonzalez, F. J., and Idle, J. R. (2007). LC-MS-based metabolomics in drug metabolism.
Drug Metab Rev 39, 581–597. doi: 10.1080/03602530701497804

Chen, W., Chen, Q., Kumar, A., Jiang, X., Zhang, K. Y. J., and Yang, Q. (2021). Structure-based
virtual screening of highly potent inhibitors of the nematode chitinase CeCht1. Journal of
Enzyme Inhibition and Medicinal Chemistry 36, 1198–1204. doi:
10.1080/14756366.2021.1931862

Chin, C.-S., Alexander, D. H., Marks, P., Klammer, A. A., Drake, J., Heiner, C., et al. (2013).
Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat
Methods 10, 563–569. doi: 10.1038/nmeth.2474

De Vos, R. C. H., Moco, S., Lommen, A., Keurentjes, J. J. B., Bino, R. J., and Hall, R. D. (2007).
Untargeted large-scale plant metabolomics using liquid chromatography coupled to mass
spectrometry. Nat Protoc 2, 778–791. doi: 10.1038/nprot.2007.95

Deng L., Qinghua L. I. U., Zhichun Z., Kai G. a. O., and Dinghui L. U. O. (2024). Genetic diversity
analysis and core collection of pinewood nematodiasis-resistant Pinus massoniana germplasm
resources. zjnldxxb 41, 67–78. doi: 10.11833/j.issn.2095-0756.20230333

Du, M., Zhu, Y., Nan, H., Zhou, Y., and Pan, X. (2024). Regulation of sugar metabolism in fruits.
Scientia Horticulturae 326, 112712. doi: 10.1016/j.scienta.2023.112712

Fang, X., Liu, Y., Xiao, J., Ma, C., and Huang, Y. (2023). GC-MS and LC-MS/MS metabolomics
revealed dynamic changes of volatile and non-volatile compounds during withering process of
black tea. Food Chem 410, 135396. doi: 10.1016/j.foodchem.2023.135396

Faria, J. M. S., Sena, I., Vieira da Silva, I., Ribeiro, B., Barbosa, P., Ascensão, L., et al. (2015). In vitro
co-cultures of Pinus pinaster with Bursaphelenchus xylophilus: a biotechnological approach to
study pine wilt disease. Planta 241, 1325–1336. doi: 10.1007/s00425-015-2257-9

Furuta, Y., Yamamoto, H., Hirakawa, T., Uemura, A., Pelayo, M. A., Iimura, H., et al. (2024). Petal
abscission is promoted by jasmonic acid-induced autophagy at Arabidopsis petal bases. Nat
Commun 15, 1098. doi: 10.1038/s41467-024-45371-3

Gao, C., Ren, L., Wang, M., Wang, Z., Fu, N., Wang, H., et al. (2022). Full-Length Transcriptome
Sequencing-Based Analysis of Pinus sylvestris var. mongolica in Response to Sirex noctilio
Venom. Insects 13, 338. doi: 10.3390/insects13040338

Gao, J., Xu, L., and Ye, J. (2024). Growth and genetic diversity analysis of clones screened by
phenotypical resistant to pine wilt disease in Pinus massoniana. Journal of Nanjing Forestry
University (Natural Sciences Edition), 67–78.

Guo, J.-Y., Zhu, Z.-H., Chen, Y.-M., and Zhu, L.-H. (2023). Screen of Pinus massoniana for Resistance
to Pinewood Nematode: In Vitro Propagation and Evaluation of Regenerated Microshoots.
Forests 14, 1056. doi: 10.3390/f14051056

Hirao, T., Matsunaga, K., Hirakawa, H., Shirasawa, K., Isoda, K., Mishima, K., et al. (2019).
Construction of genetic linkage map and identification of a novel major locus for resistance to
pine wood nematode in Japanese black pine (Pinus thunbergii). BMC Plant Biology 19, 424.
doi: 10.1186/s12870-019-2045-y

Hopf-Biziks, A., Schröder, T., and Schütz, S. (2017). Long-term survival and non-vector spread of the
pinewood nematode, Bursaphelenchus xylophilus, via wood chips. Forest Pathology 47,
e12340. doi: 10.1111/efp.12340

Hu, J., Gebremariam, T. G., Zhang, P., Qu, Y., Qiu, D., Shi, X., et al. (2023a). Resistance to Powdery
Mildew Is Conferred by Different Genetic Loci at the Adult-Plant and Seedling Stages in
Winter Wheat Line Tianmin 668. Plant Disease 107, 2133–2143. doi: 10.1094/PDIS-11-22-
2633-RE

Hu, S., Yu, K., Yan, J., Shan, X., and Xie, D. (2023b). Jasmonate perception: Ligand–receptor
interaction, regulation, and evolution. Molecular Plant 16, 23–42. doi:
10.1016/j.molp.2022.08.011

Hu, X.-G., Zhuang, H., Lin, E., Borah, P., Du, M., Gao, S., et al. (2022). Full-Length Transcriptome
Sequencing and Comparative Transcriptomic Analyses Provide Comprehensive Insight Into
Molecular Mechanisms of Cellulose and Lignin Biosynthesis in Cunninghamia lanceolata.
Frontiers in Plant Science 13. Available at: https://www.frontiersin.org/journals/plant-
science/articles/10.3389/fpls.2022.883720 (Accessed March 3, 2024).

Hwang, H.-S., Han, J.-Y., and Choi, Y.-E. (2021). Enhanced Emission of Monoterpene 3-Carene in
Pinus densiflora Infected by Pine Wood Nematode and Characterization of 3-Carene Synthase.
Forests 12, 514. doi: 10.3390/f12050514

Hwang, H.-S., Kim, Y.-R., Han, J.-Y., and Choi, Y.-E. (2022). Nematicidal Properties and Chemical
Composition of Pinus rigida Mill. Resin against Pinewood Nematodes. Forests 13, 1131. doi:
10.3390/f13071131

Ichihara, Y., Fukuda, K., and Suzuki, K. (2000). Early Symptom Development and Histological
Changes Associated with Migration of Bursaphelenchus xylophilus in Seedling Tissues of
Pinus thunbergii. Plant Disease 84, 675–680. doi: 10.1094/PDIS.2000.84.6.675

Isshiki, R., Galis, I., and Tanakamaru, S. (2014). Farinose flavonoids are associated with high freezing
tolerance in fairy primrose (Primula malacoides) plants. Journal of Integrative Plant Biology
56, 181–188. doi: 10.1111/jipb.12145

Jha, S., and Modi, H. A. (2018). Statistical optimization of chitinase production by Streptomyces
rubiginosus SP24 and efficacy of purified chitinase to control root-knot nematode infection in
Vigna radiata under controlled conditions. Chemical and Biological Technologies in
Agriculture 5, 20. doi: 10.1186/s40538-018-0133-0

Jia, J., Chen, L., Yu, W., Cai, S., Su, S., Xiao, X., et al. (2023). The novel nematicide chiricanine A suppresses Bursaphelenchus xylophilus pathogenicity in Pinus massoniana by inhibiting Aspergillus and its secondary metabolite, sterigmatocystin. Front Plant Sci 14, 1257744. doi: 10.3389/fpls.2023.1257744

Ju, Y., Wang, X., Guan, T., Peng, D., and Li, H. (2016). Versatile glycoside hydrolase family 18 chitinases for fungi ingestion and reproduction in the pinewood nematode Bursaphelenchus xylophilus. International Journal for Parasitology 46, 819–828. doi: 10.1016/j.ijpara.2016.08.001

Kalachova, T., Jindřichová, B., Burketová, L., Monard, C., Blouin, M., Jacquiod, S., et al. (2023). Controlled natural selection of soil microbiome through plant-soil feedback confers resistance to a foliar pathogen. Plant Soil 485, 181–195. doi: 10.1007/s11104-022-05597-w

Kunkel, B. N., and Johnson, J. M. B. (2021). Auxin Plays Multiple Roles during Plant–Pathogen Interactions. Cold Spring Harb Perspect Biol 13, a040022. doi: 10.1101/cshperspect.a040022

Langdon, W. B. (2015). Performance of genetic programming optimised Bowtie2 on genome comparison and analytic testing (GCAT) benchmarks. BioData Min 8, 1. doi: 10.1186/s13040-014-0034-0

Lee, Y. S., and Kim, K. Y. (2015). Statistical optimization of medium components for chitinase production by Pseudomonas fluorescens strain HN1205: role of chitinase on egg hatching inhibition of root-knot nematode. Biotechnology & Biotechnological Equipment 29, 470–478. doi: 10.1080/13102818.2015.1010702

Li, D., Li, Y., Wang, X., Zhang, W., Wen, X., Liu, Z., et al. (2023). Engineered pine endophytic Bacillus toyonensis with nematocidal and colonization abilities for pine wilt disease control. Frontiers in Microbiology 14. Available at: https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2023.1240984 (Accessed February 24, 2024).

Li, J., Liu, C., Wang, S., and Mao, X. (2024). Staphylococcus aureus enters viable-but-nonculturable state in response to chitooligosaccharide stress by altering metabolic pattern and transmembrane transport function. Carbohydrate Polymers 330, 121772. doi: 10.1016/j.carbpol.2023.121772

Li, J., Ma, W., Zeng, P., Wang, J., Geng, B., Yang, J., et al. (2015). LncTar: a tool for predicting the RNA targets of long noncoding RNAs. Brief Bioinform 16, 806–812. doi: 10.1093/bib/bbu048

Li, T. (YiZhou), Zhang, Y., Liu, H., Wu, Y., Li, W., and Zhang, H. (2010). Stable expression of Arabidopsis vacuolar Na+/H+ antiporter gene AtNHX1, and salt tolerance in transgenic soybean for over six generations. Chin. Sci. Bull. 55, 1127–1134. doi: 10.1007/s11434-010-0092-8

Li, W., and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics 22, 1658–1659. doi: 10.1093/bioinformatics/btl158

Li, Y.-L., Fan, C.-J., Jiang, X.-H., Tian, X.-Y., and Han, Z.-M. (2021). Bursaphelenchus xylophilus: An Important Pathogenic Factor of Pine Wilt Disease and Its Relationship with

Bursaphelenchus mucronatus. Plant Disease 105, 3055–3062. doi: 10.1094/PDIS-02-21-0396-
RE

Liu, B., Liu, Q., Zhou, Z., Yin, H., Xie, Y., and Wei, Y. (2021). Two terpene synthases in resistant
Pinus massoniana contribute to defence against Bursaphelenchus xylophilus. Plant, Cell &
Environment 44, 257–274. doi: 10.1111/pce.13873

Liu, F., Su, S., Chen, J., Xu, Q., Song, G., Dong, Y., et al. (2023a). The nematicide emamectin benzoate
increases ROS accumulation in Pinus massoniana and poison Monochamus alternatus. PLoS
One 18, e0295945. doi: 10.1371/journal.pone.0295945

Liu, M., Fan, F., He, S., Guo, Y., Chen, G., Li, N., et al. (2022). Creation of Elite Rice with High-Yield,
Superior-Quality and High Resistance to Brown Planthopper Based on Molecular Design. Rice
15, 17. doi: 10.1186/s12284-022-00563-7

Liu, R., Tao, Y., Xu, T., Wu, T., Yu, Q., Pan, S., et al. (2023b). Antioxidant activity increased due to
dynamic changes of flavonoids in orange peel during Aspergillus niger fermentation.
International Journal of Food Science & Technology 58, 3329–3336. doi: 10.1111/ijfs.16239

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol 15, 550. doi: 10.1186/s13059-014-0550-8

Lu, X., Wang, B., Cai, X., Chen, S., Chen, Z., and Xin, Z. (2020). Feeding on tea GH19 chitinase
enhances tea defense responses induced by regurgitant derived from Ectropis grisescens.
Physiologia Plantarum 169, 529–543. doi: 10.1111/ppl.13094

Mackelprang, R., and Lemaux, P. G. (2020). Genetic Engineering and Editing of Plants: An Analysis
of New and Persisting Questions. Annu Rev Plant Biol 71, 659–687. doi: 10.1146/annurev-
arplant-081519-035916

Manna, M., Rengasamy, B., and Sinha, A. K. (2023). Revisiting the role of MAPK signalling pathway
in plants and its manipulation for crop improvement. Plant, Cell & Environment 46, 2277–2295.
doi: 10.1111/pce.14606

Pajares, J. A., Álvarez, G., Hall, D. R., Ibarra, N., Hoch, G., Halbig, P., et al. (2017). Attractants for
management of the pine sawyer beetle Monochamus sutor, a potential vector of
Bursaphelenchus xylophilus. Journal of Applied Entomology 141, 97–111. doi:
10.1111/jen.12320

Park, J., Jeon, H. W., Jung, H., Lee, H.-H., Kim, J., Park, A. R., et al. (2020). Comparative
Transcriptome Analysis of Pine Trees Treated with Resistance-Inducing Substances against the
Nematode Bursaphelenchus xylophilus. Genes 11, 1000. doi: 10.3390/genes11091000

Paul, S., Gross, D., Bechtel, A., and Dutta, S. (2020). Preservation of monoterpenoids in Oligocene
resin: Insights into the evolution of chemical defense mechanism of plants in deep-time.
International Journal of Coal Geology 217, 103326. doi: 10.1016/j.coal.2019.103326

Peng, J., Zhang, Y., Liu, X., Zou, Y., Song, H., Wang, S., et al. (2023). Design, synthesis, antibacterial
activity, and mechanism of novel resveratrol derivatives containing an 1,3,4-oxadiazole moiety.
Pesticide Biochemistry and Physiology 193, 105457. doi: 10.1016/j.pestbp.2023.105457

Qiu, J., Xie, J., Chen, Y., Shen, Z., Shi, H., Naqvi, N. I., et al. (2022). Warm temperature compromises JA-regulated basal resistance to enhance Magnaporthe oryzae infection in rice. Molecular Plant 15, 723–739. doi: 10.1016/j.molp.2022.02.014

Qiu, Y.-J., Wu, X.-Q., Wen, T.-Y., Hu, L.-J., Rui, L., Zhang, Y., et al. (2023). The Bursaphelenchus xylophilus candidate effector BxLip-3 targets the class I chitinases to suppress immunity in pine. Molecular Plant Pathology 24, 1033–1046. doi: 10.1111/mpp.13334

Rahman, S. U., Raza, G., Naqvi, R. Z., McCoy, E., Hammad, M., LaFayette, P., et al. (2023). A source of resistance against yellow mosaic disease in soybeans correlates with a novel mutation in a resistance gene. Frontiers in Plant Science 14. Available at: https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2023.1230559 (Accessed February 25, 2024).

Raorane, C. J., Lee, J.-H., Kim, Y.-G., Rajasekharan, S. K., Garc á-Contreras, R., and Lee, J. (2019). Antibiofilm and Antivirulence Efficacies of Flavonoids and Curcumin Against Acinetobacter baumannii. Frontiers in Microbiology 10. Available at: https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2019.00990 (Accessed March 1, 2024).

Rubio-Somoza, I., and Bl ázquez, M. A. (2023). Plant–pathogen interactions: The need to evolve to stay the same. Current Biology 33, R902–R904. doi: 10.1016/j.cub.2023.07.037

S ánchez-Mart ñez, J. D., Bueno, M., Alvarez-Rivera, G., Tudela, J., Iba ñez, E., and Cifuentes, A. (2021). In vitro neuroprotective potential of terpenes from industrial orange juice by-products. Food Funct. 12, 302–314. doi: 10.1039/D0FO02809F

Shen, S., Ding, B., Jiang, X., Yang, M., Yang, Q., and Dong, L. (2022). Discovery of novel inhibitors targeting nematode chitinase CeCht1: Virtual screening, biological evaluation, and molecular dynamics simulation. Frontiers in Chemistry 10. Available at: https://www.frontiersin.org/articles/10.3389/fchem.2022.1021295 (Accessed March 4, 2024).

Shi, A., Xu, J., Guo, Y., Rensing, C., Chang, J., Zhang, T., et al. (2024). Jasmonic acid's impact on Sedum alfredii growth and cadmium tolerance: A physiological and transcriptomic study. Science of The Total Environment 914, 169939. doi: 10.1016/j.scitotenv.2024.169939

Sybilska, E., and Daszkowska-Golec, A. (2023). A complex signaling trio in seed germination: Auxin-JA-ABA. Trends in Plant Science 28, 873–875. doi: 10.1016/j.tplants.2023.05.003

Theisen, F. F., Prestel, A., Elkjær, S., Leurs, Y. H. A., Morffy, N., Strader, L. C., et al. (2024). Molecular switching in transcription through splicing and proline-isomerization regulates stress responses in plants. Nat Commun 15, 592. doi: 10.1038/s41467-024-44859-2

Theodoridis, G., Gika, H. G., and Wilson, I. D. (2008). LC-MS-based methodology for global metabolite profiling in metabonomics/metabolomics. TrAC Trends in Analytical Chemistry 27, 251–260. doi: 10.1016/j.trac.2008.01.008

Thulasinathan, T., Ayyenar, B., Kambale, R., Manickam, S., Chellappan, G., Shanmugavel, P., et al. (2023). Marker Assisted Introgression of Resistance Genes and Phenotypic Evaluation Enabled

Identification of Durable and Broad-Spectrum Blast Resistance in Elite Rice Cultivar, CO 51.
Genes 14, 719. doi: 10.3390/genes14030719

Török, K., Szentmiklóssy, M., Tremmel-Bede, K., Rakszegi, M., and Tömösközi, S. (2019).
Possibilities and barriers in fibre-targeted breeding: Characterisation of arabinoxylans in wheat
varieties and their breeding lines. Journal of Cereal Science 86, 117–123. doi:
10.1016/j.jcs.2019.01.012

Umebayashi, T., Fukuda, K., Haishi, T., Sotooka, R., Zuhair, S., and Otsuki, K. (2011). The
Developmental Process of Xylem Embolisms in Pine Wilt Disease Monitored by Multipoint
Imaging Using Compact Magnetic Resonance Imaging. Plant Physiology 156, 943–951. doi:
10.1104/pp.110.170282

Vazquez-Vilar, M., Fernandez-del-Carmen, A., Garcia-Carpintero, V., Drapal, M., Presa, S., Ricci, D.,
et al. (2023). Dually biofortified cisgenic tomatoes with increased flavonoids and branched-
chain amino acids content. Plant Biotechnology Journal 21, 2683–2697. doi:
10.1111/pbi.14163

Wang, G., Hu, C., Zhou, J., Liu, Y., Cai, J., Pan, C., et al. (2019). Systemic Root-Shoot Signaling
Drives Jasmonate-Based Root Defense against Nematodes. Current Biology 29, 3430-3438.e4.
doi: 10.1016/j.cub.2019.08.049

Wang, T., Gu, X., Guo, L., Zhang, X., and Li, C. (2024a). Integrated metabolomics and transcriptomics
analysis reveals γ-aminobutyric acid enhances the ozone tolerance of wheat by accumulation
of flavonoids. Journal of Hazardous Materials 465, 133202. doi:
10.1016/j.jhazmat.2023.133202

Wang, X., Yi, X., Hou, C., Wang, X., Sun, X., Zhang, Z., et al. (2022). Map-based cloning and
functional analysis revealed ABCC2 is responsible for Cry1Ac toxin resistance in Bombyx
mori. Archives of Insect Biochemistry and Physiology 110, e21886. doi: 10.1002/arch.21886

Wang, Y., Shi, W., Jing, B., and Liu, L. (2024b). Modulation of soil aeration and antioxidant defenses
with hydrogen peroxide improves the growth of winter wheat (Triticum aestivum L.) plants.
Journal of Cleaner Production 435, 140565. doi: 10.1016/j.jclepro.2024.140565

Xie, W., Huang, A., Li, H., Feng, L., Zhang, F., and Guo, W. (2017). Identification and comparative
analysis of microRNAs in Pinus massoniana infected by Bursaphelenchus xylophilus. Plant
Growth Regul 83, 223–232. doi: 10.1007/s10725-016-0221-8

Xie, Y., Liu, B., Gao, K., Zhao, Y., Li, W., Deng, L., et al. (2023). Comprehensive Analysis and
Functional Verification of the Pinus massoniana NBS-LRR Gene Family Involved in the
Resistance to Bursaphelenchus xylophilus. Int J Mol Sci 24, 1812. doi: 10.3390/ijms24031812

Xu, M., Jiang, Y., Chen, S., Chen, F., and Chen, F. (2021). Herbivory-Induced Emission of Volatile
Terpenes in Chrysanthemum morifolium Functions as an Indirect Defense against Spodoptera
litura Larvae by Attracting Natural Enemies. J. Agric. Food Chem. 69, 9743–9753. doi:
10.1021/acs.jafc.1c02637

Xue, Q., Xiang, Y., Wu, X.-Q., and Li, M.-J. (2019). Bacterial Communities and Virulence Associated
with Pine Wood Nematode Bursaphelenchus xylophilus from Different Pinus spp. International
Journal of Molecular Sciences 20, 3342. doi: 10.3390/ijms20133342

Yazaki, K., Takanashi, T., Kanzaki, N., Komatsu, M., Levia, D. F., Kabeya, D., et al. (2018). Pine wilt
disease causes cavitation around the resin canals and irrecoverable xylem conduit dysfunction.
Journal of Experimental Botany 69, 589–602. doi: 10.1093/jxb/erx417

Zhang, X., Huang, T., Wang, Q., Guo, Y., Zhang, P., Xie, H., et al. (2022). Mechanisms of Resistance
to Spot Blotch in Yunnan Iron Shell Wheat Based on Metabolome and Transcriptomics.
International Journal of Molecular Sciences 23, 5184. doi: 10.3390/ijms23095184

Zhang, Y., Gu, X., Huang, C., Zhou, J., Shi, J., and Zhao, L. (2023). Temperature-regulated metabolites
of Serratia marcescens inhibited reproduction of pinewood nematode Bursaphelenchus
xylophilus. iScience 26, 107082. doi: 10.1016/j.isci.2023.107082

Zhao, B. G., Liu, Y., and Lin, F. (2007). Effects of Bacteria Associated with Pine Wood Nematode
(Bursaphelenchus xylophilus) on Development and Egg Production of the Nematode. Journal
of Phytopathology 155, 26–30. doi: 10.1111/j.1439-0434.2006.01188.x

Zhao, Z., Chen, W., Dong, Y., Yang, Q., Lu, H., and Zhang, J. (2023). Discovery of Potent N-
Methylcarbamoylguanidino Insect Growth Regulators Targeting OfChtI and OfChi-h. J. Agric.
Food Chem. 71, 12431–12439. doi: 10.1021/acs.jafc.3c02448

Zhou, Y., Wang, Y., Zhang, D., and Liang, J. (2024). Endomembrane-biased dimerization of ABCG16
and ABCG25 transporters determines their substrate selectivity in ABA-regulated plant growth
and stress responses. Molecular Plant 0. doi: 10.1016/j.molp.2024.02.005

 FIGURE LEGENDS

FIGURE 1 | Nomination of potential PWN carrying determination genes in *Pinus massoniana*
using full-length transcriptome. (**A**) Workflow of full-length transcriptome sequencing and analysis.
(**B**) Venn diagrams of unigenes annotated in different databases. (**C**) To screen the potential PWN-
carrying determination genes in *Pinus massoniana* (β), the common genes (α, >90% similarity of amino
acid sequences) between two PWN host pine (*Pinus massoniana* and *Pinus densiflora*) were compared
with proteins of non-PWN host pine (*Pinus tabuliformis*) to develop the library of non-common (β,
<90% similarity of amino acid sequences) genes. (**D**) A bubble plot diagram presents the KEGG
annotation of genes in β. The top 25 KEGG terms are shown.

FIGURE 2 | Transcriptome analysis revels the PWN carrier determine genes in *P. massoniana*.
(**A**) Relative PWN carrying amount of different *P. massoniana* cultivars (TC, tolerant cultivar; NC,
neutral cultivar; RC, resistant cultivar) were quantified six weeks post inoculation. Each box represents
a mean value, with bars representing one standard deviation (SD). ** indicate statistical differences (*P*
≤ *0.01*) between samples using one-way ANOVAs (Tukey's test). (**B**) Venn diagrams shown the
number of potential PWN-carrying determination genes in *P. massoniana* (β), differentially expressed
genes (DEGs) significantly correlated with PWN amount, and PWN-carrying determination genes in
P. massoniana (γ). (**C**) Differential gene expression analysis revealed 318 genes (γ) up-regulated and
down-regulated in all two clusters. Adjusted *P* ≤ *0.01* are shown in red, and adjusted *P* ≥ *0.01* are
shown in black. (**D**) A bubble plot diagram presents the KEGG annotation of genes in γ. The top 25
KEGG terms are shown.

FIGURE 3 | Different molecular strategies of *Pinus massoniana* in response to PWN invasion. (**A**)
Mortality of different *P. massoniana* cultivars were documented six weeks post inoculating with PWN.
Each box represents a mean value, with bars representing one standard deviation (SD). ** indicate
statistical differences (*P* ≤ 0.01) between samples using one-way ANOVAs (Tukey's test). (**B**) Upset
maps shown the number of potential PWN-carrying determination genes in *Pinus massoniana* (β),
DEGs of tolerance and neutral cultivars (TC vs NC), DEGs of resistance and neutral cultivars (RC vs
NC), Common Related Genes (CRGs), Tolerance Related Genes (TRGs) and Resistant Related Genes
(RRGs) (**C**) Heat maps of relative expression level of TRGs, CRGs and RRGs in different *P.
massoniana* cultivars, data were homogenized by Z-score. Bubble plot diagrams present the KEGG
annotation of TRGs (**D**), CRGs (**E**), and RRGs (**F**), respectively. The top 25 KEGG terms are shown.

FIGURE 4 | Different metabolic strategies of *Pinus massoniana* in response to PWN invasion. (**A**)
The circular heat map shows the relative expression (inner circle) and correlation coefficients (outside
circle) of metabolites highly correlated with PWN (δ) in different *P. massoniana* cultivars, data were
homogenized by Z-score. (**B**) A bubble plot diagram presents the KEGG annotation of metabolites in
δ. The top 25 KEGG terms are shown. (**C**) Upset maps shown the number of different accumulation
metabolites (DAMs) in *P. massoniana* of tolerance and neutral cultivars (TC vs NC), DAMs of
resistance and neutral cultivars (RC vs NC), Common Related Metabolites (CRMs), Tolerance Related
Metabolites (TRMs) and Resistant Related Metabolites (RRMs). Bubble plot diagrams present the
KEGG annotation of TRMs (**D**), CRMs (**E**), and RRMs (**F**), respectively. The top 25 KEGG terms are
shown.

FIGURE 5 | Genetic landscape in *Pinus massoniana* regard to PWN invasion. (**A**) Heat map
represent the relative expression level of key different expressed genes (DEGs) and different

accumulation metabolites (DAMs) involved in different strategies of *P. massoniana* against PWN
invasion, data were homogenized by Z-score. (**B**) Illumination of critical genetic responses in *P.*
massoniana against PWN invasion regard to different ecological cultivar