Discovery of knock-down resistance in the major malaria vector *Anopheles funestus* reveals the legacy of persistent DDT pollution.

Authors

Joel O. Odero[1,2*] ✉, Tristan P. W. Dennis[3*], Brian Polo[4], Joachim Nwezeobi[5], Marilou Boddé[5], Sanjay C. Nagi[3], Anastasia Hernandez-Koutoucheva[5], Ismail H. Nambunga[1], Hamis Bwanary[1], Gustav Mkandawile[1], Nicodem J Govella[1], Emmanuel W. Kaindoa[1], Heather M. Ferguson[2], Eric Ochomo[4], Chris S. Clarkson[5], Alistair Miles[5], Mara K. N. Lawniczak[5], David Weetman[3#], Francesco Baldini[1,2#], Fredros O. Okumu[1,2#] ✉

[*]These authors contributed equally to this work

[#]These authors equally supervised this work

✉ Corresponding author

Affiliations

1. Environmental Health and Ecological Sciences Department, Ifakara Health Institute, Ifakara, Tanzania

2. School of Biodiversity, One Health, and Veterinary Medicine, G12 8QQ, University of Glasgow, Glasgow, UK

3. Department of Vector Biology, Liverpool School of Tropical Medicine, L3 5QA, Liverpool, UK

4. Entomology Section, Centre for Global Health Research, Kenya Medical Research Institute, Kisumu, Kenya

5. Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, CB10 1SA, UK.

Abstract

A major mechanism of insecticide resistance in arthropod pests is knock-down resistance (*kdr*) caused by mutations in the voltage-gated sodium channel (*Vgsc*) gene. Common in most malaria *Anopheles* vector species, *kdr* mutations have never been observed in *Anopheles funestus*, the principal vector in Eastern and Southern Africa. From whole-genome sequencing of 333 *An. funestus* samples from a breadth of populations in Tanzania, we found 8 novel amino acid substitutions in the *Vgsc* gene, including the *kdr* variant, L1014F (L976F in *An. funestus*), in tight linkage disequilibrium with another (P1842S). The mutants were found only at high frequency in one region, with a significant decline between 2017 and 2023. When evaluating the resistance phenotype of these samples, we found a strong association between L976F and survivorship to the exposure to DDT insecticide, but no association with a pyrethroid insecticide (deltamethrin). No DDT products are currently prequalified by WHO for vector control, and the chemical is banned in Tanzania. However, widespread DDT contamination and a legacy of extensive stockpiles in the same region where we found the *kdr* alleles, may have selected for this evolution. Continued monitoring is necessary to confirm the origin of *kdr* in *An. funestus*, and how it may threaten the effectiveness of insecticide-based vector control in Africa.

Main

Chemical insecticides are central to the control of agricultural pests and disease vectors. The control of *Anopheles* mosquitoes through the distribution of over 2.9 billion insecticide-treated bed nets (ITNs)[1] has helped avert an estimated 633 million cases of malaria[2] - a disease that still kills 600,000 yearly[1]. However, the widespread use of insecticides for agricultural pest and disease vector control also has detrimental consequences, including direct lethal and sub-lethal effects on human and animal

health[3,4] and destabilizing effects on ecosystem structure and function. For example,

insecticide exposure is a key stressor affecting the population decline of pollinators,

essential for ecosystem health and food production[5,6]. These impacts are often

unpredictable[7].

Strong selection pressures generated by insecticide-based agricultural pest and

disease vector control activities have resulted in the independent evolution of a

diverse range of mechanisms that confer insecticide resistance (IR) phenotypes in

numerous insect species[8]. One of the earliest described IR mechanisms was the

emergence of knock-down resistance (*kdr*), mediated by mutations in the target-site of

pyrethroid and organochlorine insecticides, located in the voltage-gated sodium

channel gene (*Vgsc*), an essential component of the nervous system[9]. These *kdr*-driven

resistance phenotypes appeared rapidly after the introduction of the organochlorine

dichloro-diphenyl-trichloroethane (DDT) spraying for insect control in the mid-20th

century[10]. Knock-down resistance [11,12], the key ingredient in ITNs - the first line of

defence against malaria. The threat posed by IR to malaria vector control has led to

intense efforts to develop coordinated strategies for IR management that integrate

entomological, molecular, and genomic surveillance data towards proactive strategies

for mitigating the impacts of IR[13,14].

As part of a genomic survey done in Tanzania to understand the evolution and

spread of insecticide resistance in *Anopheles funestus* - the dominant malaria vector in

Eastern and Southern Africa[15], we report the first discovery of *kdr* mutations in *An.*

funestus. Despite a complete ban on DDT use for agriculture and vector control in

Tanzania since 2008[16], we discover that this mutation confers resistance to DDT, and

suggests environmental contamination from extensive DDT stockpiles[17], or unofficial

agricultural use, as possible causes. The emergence of *kdr*, which confers resistance to

insecticides in major crop pests and vectors of disease (such as *An. gambiae* and *Aedes*

aegypti)[18], highlights the potential of chemical insecticide contamination or unofficial

use to exert unexpected and potentially harmful impacts on public health.

We analysed WGS data from 333 mosquitoes sampled from 10 sites across

Tanzania (**Figure 1A**). We performed genome-wide selection scans (GWSS) with the

G123 statistic[26] to test for evidence of selective sweeps in the *An. funestus* genome

associated with known or novel IR loci (**Figure 1B**); grouping samples by

administrative region (see **Supp. Table 1** for per-group sample numbers). We detected

a clear signal of elevated G123 in the region containing the *Vgsc* gene in samples from

the Morogoro region in the southeastern part of the country (**Figure 1B**). In Kagera,

Katavi, and Mwanza regions, there was no visible sign of a selective sweep at or near

the *Vgsc* region. In Dodoma, Lindi, Ruvuma, and Tanga, there were peaks of elevated

G123 near to *Vgsc*, but these appeared within the context of relatively high G123 across

the chromosome **(Figure 1B).**

Figure 1: (A) Map of *An. funestus* collection locations. Points indicate sample

collection locations. Point colour indicates the administrative region from which

samples were collected. (**B**) G123 selection scans of *An. funestus* chromosome 3RL,

coloured and windowed by sample collection region (where n>20 – see **Supp Table**

2). X axis indicates position (in base-pairs (bp)), Y axis indicates the selection statistic

G123. Grey dotted line indicates the location of the *Vgsc* gene. Note Mwanza region

is absent from (**B**) as there were too few samples (n<20) to perform a selection scan.

Mutations in *Vgsc* confer *kdr* in numerous pest and insect taxa[19]. While not previously reported in *An. funestus*[20], *kdr* mutations in major malaria vectors within the *An. gambiae* complex are subject to intense selection[21,22] and confer resistance to pyrethroid and organochlorine insecticides used in ITNs and insecticide sprays[12,19]. We searched our data for mutations in the *Vgsc* gene and found 8 amino acid substitutions occurring at frequencies greater than 5% (**Figure 2A**). Of these, two alleles, L976F and P1842S occurred at the highest frequency (**Figure 2A**). The frequencies of P1824S and L976F were highest in samples collected from Morogoro in 2017 (0.75 and 0.90 respectively) **(Figure 2A)** and declined yearly, reaching their lowest frequency in samples collected in 2023 (0.48 and 0.56 respectively; χ^2 = 12.15, p=0.0005; **Figure 2B).** These mutations occurred at very low frequencies or were absent in all other locations (**Figure 2A).** We aligned the *An. funestus Vgsc* sequence (Gene ID: AFUN2_008728) with that of *Musca domestica* (Gene ID: X96668) and *An. gambiae* (AGAP004707-RD AgamP4.12 gene set)[22]. The amino acid change at *An. funestus* L976F corresponded to L1014F in *M. domestica* and L995F in *An. gambiae* in domain II subunit 6 (IIS6) of the *Vgsc* gene (**Table 1**), which in *An. gambiae* species complex drastically increases IR to DDT and pyrethroids[12,23]. The second variant P1842S corresponded to P1874S in *An. gambiae* and P1879 in *M. domestica* and were all in the C-terminal domain (**Table 1**).

Table 1: Comparative non-synonymous nucleotide variation in the voltage-gated sodium channel gene

Variant					Population allele frequency (%)							
Position[a]	Af[b]	Ag[c]	Md[d]	Domain	Dodoma 2022	Morogoro 2017	Morogoro 2019	Morogoro 2021	Morogoro 2023	Pwani 2019	Ruvuma 2019	Ruvuma 2022
3RL:44,115,564 T>C	I2030V	I2061	P2063	COOH	0.00	0.00	0.00	0.06	0.02	0.00	0.00	0.02

Position	AF	Ag	Md									
3RL:44,115,768 C>A	G1962C	A1994	P1997	COOH	0.00	0.00	0.00	0.00	0.00	0.05	0.06	0.05
3RL:44,116,128 G>A	P1842S	P1874S	P1879	COOH	0.04	0.75	0.69	0.67	0.48	0.10	0.11	0.11
3RL:44,116,334 T>C	N1773S	N1805	N1810		0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00
3RL:44,116,923 A>T	F1638Y	F1670	V1675		0.00	0.05	0.00	0.00	0.04	0.15	0.00	0.00
3RL:44,117,167 A>T	W1557R	W1589	W1594		0.00	0.10	0.14	0.00	0.04	0.00	0.00	0.00
3RL:44,122,391 C>A	G1144C	G1173	G1180		0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.00
3RL:44,125,475 T>A	L976F	L995F	L1014	IIS6	0.04	0.90	0.83	0.67	0.56	0.25	0.11	0.11

Position[a] Position relative to the *Anopheles funestus* strain FUMOZ reference, chromosome arm 3RL.

AF[b] Codon numbering according to *Anopheles funestus* transcript LOC125769886_t4.

Ag[c] Codon numbering according to *Anopheles gambiae* transcript AGAP004707-RD in geneset AgamP4.12.

Md[d] Codon numbering according to *Musca domestica* EMBL accession X96668 Williamson *et al.*[24].

In *An. gambiae*, multiple *kdr* haplotypes have evolved independently[25,26]. To elucidate *Vgsc* haplotype structure in *An. funestus,* we computed pairwise linkage disequilibrium (LD) using the Rogers and Huff method[27], between nonsynonymous variants occurring at a frequency of > 5% in Tanzanian *An. funestus* [**Figure 2C**]. We found that P1824S occurred in tight LD with L976F (D'=0.95) [**Figure 2C**]. Of other non-synonymous polymorphisms, F1638Y and W1557R exhibited only weak LD with L976F [**Figure 2C**]. We constructed a haplotype clustering dendrogram from haplotypes in all 333 individuals, from the *Vgsc* gene [**Figure 3**]. The clustering dendrogram disclosed three major clades and three main combinations of the four

most prevalent *Vgsc* alleles **[Figure 3].** The most striking signal was a subclade of

identical, or near-identical haplotypes containing both L976F and P1842S [**Figure 3],**

indicating a selective sweep on a combined L976F/P1842S haplotype. This combined

haplotype was present at higher frequencies in the Morogoro region relative to the

neighbouring regions of Pwani, Ruvuma, and Dodoma [**Figure 3**]. Most amino acid

substitutions were present in a single clade in samples from Pwani, Dodoma,

Ruvuma, and especially Morogoro [**Figure 3**]. This extremely restricted geographic

hotspot of *kdr* is in stark contrast to its distribution in *An. gambiae* where *kdr* mutations

spread rapidly across vast spatial scales in response to strong selection pressure[21,28].

Figure 2: (A) Heatmap of *Vgsc* allele frequencies. Y axis labels indicate mutation effect,

chromosome position, and nucleotide change. X axis labels indicate collection date, heatmap

intensity indicates frequency where darker = higher, with frequency labelled in each

heatmap facet. Heatmap is panelled by sample collection region. **(B)** L976F and 1842S

frequencies, in Morogoro region, over time. Y axis indicates allele frequency, X axis indicates

date. Line and point colour refer to mutation, specified in the legend. Bars indicate 95%

confidence intervals. **(C)** Heatmap of linkage disequilibrium (LD) (Rogers and Huff R)

between nonsynonymous variants in the *Vgsc* gene at frequency > 5%. LD is indicated by fill

colour. SNP effects and positions are labelled on X and Y axes. (**D**) Denotes the association of

L976F and P1842S with resistance to Deltamethrin and DDT. Colour and panelling are by

mutation, x axis indicates genotype, y axis indicates mortality, point shape indicates mean

for each insecticide and line indicates the 95% CI based on generalised mixed model

prediction.

Figure 3: Clustering of haplotypes at the *Vgsc* gene (LOC125769886, 3RL:44105643-44156624).

Dendrogram branch length corresponds to no. SNPs difference (y axis). Tips correspond to

individual haplotypes (x axis). The coloured Population bar denotes the administrative region

of origin (as described by the legend). Red blocks at the bottom indicate the presence of the

labelled non-synonymous SNPs in the *Vgsc* gene.

Resistance to all major classes of insecticide is common in *An. funestus* and is

primarily mediated through the increased activity of enzymes that bind to and

metabolise insecticides (metabolic resistance)[29,30]. This contrasts with *An. gambiae*

where resistance is mostly conferred by a combination of metabolic and target-site

resistance[8]. We investigated phenotypic resistance (as measured by mosquito survival

following insecticide exposure) in *An. funestus* to the discriminating doses of

deltamethrin (type II pyrethroid) and DDT and tested whether the metabolic enzyme

synergist piperonyl butoxide (PBO), which is increasingly used on ITNs[31], could

restore susceptibility in pyrethroid-resistant populations in Morogoro region. The

mosquitoes were phenotypically resistant to both deltamethrin (24-hr mortality: 46%,

CI 35.6 – 57.1) and DDT (68%, CI 57.8 - 77.9) but deltamethrin susceptibility was

significantly restored in the presence of PBO (95%, CI 85.6 - 98.4). To explore the

possible association between L976F and P1842S alleles with DDT and deltamethrin

resistance, we genotyped surviving (resistant) and dead (susceptible) mosquitoes

from IR bioassays for both L976Fand P1842S loci). Neither locus was associated with

deltamethrin resistance: L976F (χ^2 = 0.04, p = 0.84) and P1842S (χ^2 = 0.59, p = 0.44).

(**Figure 2D**).   We found a strong association with DDT resistance in mosquitoes

carrying the mutant allele of L976F (χ^2 = 9.23, odds ratio = 11.0, p = 0.0024) and a

marginally non-significant positive association for P1842S (χ^2 = 3.75, p = 0.0528)

(**Figure 2D**). DDT is a largely obsolete, banned, pesticide that is no longer widely used

for vector control in Tanzania, or in Africa as a whole, due to its bio-accumulative and

toxic properties. However, the presence of DDT resistance phenotypes in *An. funestus*

in Morogoro in 2015[32] suggests *kdr*-mediated resistance to DDT has been present at

least since then, and the emergence of *kdr* resistance to DDT suggests that future use

of DDT for IRS may become even less favoured. The lack of association of *kdr* with

pyrethroid resistance might be due to the strong metabolic resistance shown to

pyrethroids in *An. funestus,* reducing the benefit of *kdr* [33]**.** However, since our testing

included only type II pyrethroids, additional data are required to assess potential

impacts of *kdr* on other pyrethroids and on commercial products. The association of

kdr with resistance to DDT but not pyrethroids, combined with selection signals and recently declining *kdr* allele frequencies where we have time series, suggests recent-past, rather than contemporary selection, perhaps due to factors other than the current use of public health pesticides.

There is no record of DDT use in the last decade for agriculture or vector control in the Morogoro region, or Tanzania as a whole. Tanzania, as a signatory to the 2009 Stockholm Convention on Persistent Organic Pollutants, banned the production, importation, and usage of DDT[16], except for limited use in malaria vector control. DDT was last used for malaria control in indoor residual spraying (IRS) in 2010[34]. The Africa Stockpiles Programme (ASP) was launched in 2005 to eliminate stockpiles of obsolete pesticides, including DDT. At this time, it was estimated that Tanzania still possessed approx. 1,500 tonnes of obsolete pesticides [35]. Intriguingly, Mahenge-Ulanga in the Morogoro region held a DDT stockpile of 4.5 tons (as of 2004) in [36], approximately 100 km away from where DDT-resistant *An. funestus* were detected in this study. Additionally, a large stockpile of 19 tons of DDT was also held in Morogoro town, where over 300 tonnes of DDT-contaminated soil were removed for disposal in 2012 [17]. The ASP and the Tanzanian Government eliminated 100% of inventoried publicly held DDT stockpiles and conducted extensive environmental remediation by programme close in 2013 [37]. However, extensive DDT contamination remains [38]. For example, 1040 tonnes of DDT contaminated soil were removed in 2014/2015 [36], and DDT remains in use by private individuals. The coincident proximal location of high levels of *kdr* in *An. funestus* with large past DDT stockpiles as well as the presence of widespread DDT contamination and private usage, leads us to hypothesise that the evolution of *kdr* in *An. funestus* may have been driven by DDT contamination of local larval breeding sites. The removal of DDT stockpiles by the ASP, and ongoing environmental remediation, may have contributed to reduced selection pressure on *kdr*, evident from declining frequency in Morogoro. Continued monitoring of allele

frequency and future studies of *kdr* frequencies targeted towards sites of known DDT

contamination will establish whether this hypothesis is correct.

Our discovery of *kdr* resistance mutations in the *Vgsc* gene in *An. funestus* is a

significant and unwelcome development that has the potential to pose a new threat to

vector control in the region. Reassuringly, it seems that the emergence of *kdr* is not

linked to, nor is presently likely to threaten, the mass rollout of PBO-pyrethroid

bednets currently underway in Tanzania as a response to IR. This does not preclude a

role for *kdr* in the *An. funestus* IR arsenal in the future, and the threat posed by *kdr* to

vector control in *An. gambiae* means the presence of *kdr* in *An. funestus,* even if it

evolved due to exposure to persistent organic pollution, requires close monitoring.

In *Silent Spring* (1962), Rachel Carson brought for the first time into the public

eye the unpredictable and often remote impacts of anti-insect chemical agents on

human health and on nature "on one hand delicate and destructible, on the other

miraculously tough and resilient, and capable of striking back in unexpected ways"

[39]. Our findings underscore the legacy of *Silent Spring* by reinforcing the potential for

pesticides and organic pollutants to exert unexpected influences on animal biology. If

the link we draw between *kdr* evolution and the gradual elimination of DDT and DDT-

contaminated material proves definitive, it provides an optimistic outlook for the

successful recovery of animal species, and the environment more broadly, in response

to human efforts.

Materials and Methods

All scripts and Jupyter Notebooks used to analyse genotype and haplotype data, and

produce figures and tables are available from the GitHub repository:

https://github.com/tristanpwdennis/kdr_funestus_report_2023

Mosquito collection

Anopheles funestus samples analyzed in this study were collected from ten

administrative regions in Tanzania: Dodoma, Kagera, Katavi, Lindi, Morogoro,

Mtwara, Mwanza, Pwani, Ruvuma, and Tanga (**Figure. 1A**). The collections were

done as part of a countrywide *Anopheles funestus* surveillance project in Tanzania

and were subsequently incorporated into the MalariaGEN *Anopheles*

funestus genomic surveillance project database

(https://www.malariagen.net/projects/anopheles-funestus-genomic-surveillance-

project). Mosquitoes were collected in households between 2017 and 2023 using

CDC light traps and mechanical aspirators. They were sorted by sex and taxa and

An. funestus group mosquitoes preserved individually in 96-well plates containing

80% ethanol.

Whole genome sequencing and analysis

The samples were processed as part of the *Anopheles funestus* genomics surveillance

MalariaGEN Vector Observatory (VObs) project

(https://www.malariagen.net/mosquito).Briefly, the mosquitoes were individually

whole-genome-sequenced on an Illumina NovaSeq 6000s instrument. Reads were

aligned to the *An. funestus* reference genome AfunGA1 [40] with Burrows-Wheeler

Aligner (BWA) version v0.7.15. Indel realignment was performed using Genome

Analysis Toolkit (GATK) version 3.7-0 RealignerTargetCreator and IndelRealigner.

Single nucleotide polymorphisms were called using GATK version 3.7-0

UnifiedGenotyper. Genotypes were called for each sample independently, in

genotyping mode, given all possible alleles at all genomic sites where the reference

base was not "N".

Complete specifications of the alignment and genotyping pipelines are available

from the malariagen/pipelines GitHub repository

(https://github.com/malariagen/pipelines/). The aligned sequences in BAM format

were stored in the European Nucleotide Archive (ENA).

The identification of high-quality SNPs and haplotypes were conducted using BWA

version 0.7.15 and GATK version 3.7-0. Quality control involved removal of samples

with low mean coverage, removing cross-contaminated samples, running PCA to

identify and remove population outliers, and sex confirmation by calling the sex of

all samples based on the modal coverage ratio between the X chromosome and the

autosomal chromosome arm 3R. Full quality control methods are available on the

MalariaGEN vector data user guide (https://malariagen.github.io/vector-

data/ag3/methods.html).

We used decision-tree filters that identify genomic sites where SNP calling and

genotyping is likely to be less reliable. More information on site filters can be found

on the MalariaGEN vector data user guide.

Genotypes at biallelic SNPs that passed the decision-tree site filtering process were

phased into haplotypes using a combination of read-backed and statistical phasing.

Read-backed phasing was performed for each sample using WhatsHap version 1.0

[https://whatshap.readthedocs.io/]. Statistical phasing was then performed using

SHAPEIT4 version 4.2.1 [https://odelaneau.github.io/shapeit4/].

Complete specifications of the haplotype phasing pipeline are available from the

malariagen/pipelines GitHub repository

(https://github.com/malariagen/pipelines/tree/master/pipelines/phasing-vector).

Identification of SNPs on *Vgsc*

To identify the *An. funestus Vgsc* gene and the variant that confers target-site

resistance we performed alignments between the *An. gambiae* VGSC transcript

AGAP004707-RD in AgamP4.12 geneset from the Ag1000 phase 3 data resource

(https://www.malariagen.net/data/ag1000g-phase3-snp) and AFUN2_008728 from

the *An. funestus* AfunF1.3 dataset. We extracted single nucleotide polymorphism

(SNPs) altering the amino acid of VGSC protein from the *An. funestus* dataset and

computed the allele frequency on the mosquito cohorts defined by the region and

year of collection ((See **Supp. Table 1** for per region/year sample numbers)). Under

selection pressure various alleles are expected to increase in frequency; we therefore

filtered out variant alleles with a frequency lower than 5% resulting in a list of 8

variant alleles. Multiple sequence alignments of *An. funestus Vgsc* against *An. gambaie*

and *M. domestica* were performed using MEGA v11.013.

Population genetic analyses

We searched for signatures of selective sweeps on the *Vgsc* gene using the *G123*

selection statistic [41]**.** G123 selection scans were performed on *An. funestus* genotypes

by collection region where sample *n*>20 **[**see **Figure 1A** and **Supp Table 2**] **]** using

the *g123_gwss* function in the malariagen_data python API

(https://malariagen.github.io/malariagen-data-python/latest/Af1.html). Linkage

disequilibrium (Rogers and Huff's R-squared) [27] between the 8 *Vgsc* alleles was

calculated using the *rogers_huff_r_between* in scikit-allel

(https://zenodo.org/record/4759368). Haplotype clustering was performed by

performing hierarchical clustering on a Hamming distance matrix, inferred from

phased *An. funestus* haplotypes, using the Scipy library (https://scipy.org/citing-

scipy/). Clustering dendrogram, and bar plot of amino acid substitutions, was

plotted using the seaborn library [42].

Association of L976F and P1842S alleles with insecticide resistance

To test for associations between the identified mutations with IR, we exposed wild

non-blood-fed *An. funestus* mosquitoes of unknown ages to standard doses of

deltamethrin and DDT insecticides following the WHO tube assays. For each

insecticide, we randomly separated phenotypically resistant mosquitoes (i.e., alive 24

hours post-exposure) and susceptible (i.e., dead 24 hours post-exposure) and

extracted DNA from individual mosquitoes using DNeasy Blood and Tissue kit

(Qiagen, Germany). The mosquitoes were identified at the species level using

species-specific primers that can distinguish *An. funestus* from the other members of the group [43]. To establish if the two *kdr* variants are associated with insecticide resistance, we designed PCR primers from *An. funestus Vgsc* (Gene ID: LOC125769886) to amplify domain IIS6 (L976F) and C-terminal (P1842S) (see **Supp Table 3** for primer and thermocycler conditions). The DNA fragments were separated on a 1% agarose gel, cut, purified using PureLink™ Quick Gel Extraction Kit (Invitrogen), and commercially Sanger sequenced. Collectively, we sequenced 76 individual mosquitoes: 56 from deltamethrin and the rest from the DDT bioassays. The frequencies of the wild type and mutant alleles were determined and correlated with phenotypes using generalised linear models in R-software v4.1.1.

Data availability

The sequencing data generated in this study have been deposited in the European Nucleotide Archive (https://www.ebi.ac.uk/ena/browser/home) under study number PRJEB2141.

Acknowledgements

This study was supported by funding support was received from the Bill and Melinda Gates Foundation (grant no. INV-002138) to FO, FB, HMF. Howard Hughes Medical Institute-Gates Foundation International Research Scholar Award (grant no. OPP 1099295) to FO, and the Academy Medical Science Springboard Award (ref: SBF007\100094) to FB. The findings and conclusions within this publication are those of the authors and do not necessarily reflect positions or policies of the HHMI, the BMGF or the AMSS.

This study was supported by the MalariaGEN Vector Observatory which is an
international collaboration working to build capacity for malaria vector genomic
research and surveillance and involves contributions by the following institutions
and teams. Wellcome Sanger Institute: Lee Hart, Kelly Bennett, Anastasia
Hernandez-Koutoucheva, Jon Brenas, Menelaos Ioannidis, Chris Clarkson, Alistair
Miles, Julia Jeans, Paballo Chauke, Victoria Simpson, Eleanor Drury, Osama Mayet,
Sónia Gonçalves, Katherine Figueroa, Tom Maddison, Kevin Howe, Mara
Lawniczak; Liverpool School of Tropical Medicine: Eric Lucas, Sanjay Nagi, Martin
Donnelly; Broad Institute of Harvard and MIT: Jessica Way, George Grant; Pan-
African Mosquito Control Association: Jane Mwangi, Edward Lukyamuzi, Sonia
Barasa, Ibra Lujumba, Elijah Juma. The authors would like to thank the staff of the
Wellcome Sanger Genomic Surveillance unit and the Wellcome Sanger Institute
Sample Logistics, Sequencing, and Informatics facilities for their contributions.

The MalariaGEN Vector Observatory is supported by multiple institutes and
funders. The Wellcome Sanger Institute's participation was supported by funding
from Wellcome (220540/Z/20/A, 'Wellcome Sanger Institute Quinquennial Review
2021-2026') and the Bill & Melinda Gates Foundation (INV-001927). The Liverpool
School of Tropical Medicine's participation was supported by the National Institute
of Allergy and Infectious Diseases ([NIAID] R01-AI116811), with additional support
from the Medical Research Council (MR/P02520X/1). The latter grant is a UK-funded
award and is part of the EDCTP2 programme supported by the European Union.
Martin Donnelly is supported by a Royal Society Wolfson Fellowship
(RSWF\FT\180003). The Pan-African Mosquito Control Association's participation
was funded by the Bill and Melinda Gates Foundation (INV-031595).

The findings and conclusions within this publication are those of the authors and do
not necessarily reflect positions or policies of the funders listed above.

Contributions

The project was conceived and supervised by FOO, FB, and DW. Field collection was
performed by JOO, IHN, HB, and GM. Laboratory analysis, data acquisition and
management, and preparing samples for whole genome sequencing were performed

by JOO. Sequence QC, alignments, SNP calling, and haplotype phasing were
performed by AHK, JN, CSC, and AM. JOO, BP, and TPWD analyzed the data and
generated all figures and tables. The manuscript was drafted by JOO and TPWD and
revised by all authors. Throughout the project, all authors have contributed key
ideas that have shaped the work and the final paper.

Competing interests

The authors declare no competing interests.

References

WHO. *World malaria report 2023*, <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2023> (2023).
Bhatt, S. *et al.* The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. *Nature* **526**, 207- 211, doi:10.1038/nature15535 (2015).
Berny, P. Pesticides and the intoxication of wild animals. *J Vet Pharmacol Ther* **30**, 93-100, doi:10.1111/j.1365-2885.2007.00836.x (2007).
Wurster, C. F., Jr., Wurster, D. H. & Strickland, W. N. Bird Mortality after Spraying for Dutch Elm Disease with Ddt. *Science* **148**, 90-91, doi:10.1126/science.148.3666.90 (1965).
Nicholson, C. C. *et al.* Pesticide use negatively affects bumble bees across European landscapes. *Nature*, doi:10.1038/s41586-023-06773-3 (2023).
Douglas, M. R., Sponsler, D. B., Lonsdorf, E. V. & Grozinger, C. M. County-level analysis reveals a rapidly shifting landscape of insecticide hazard to honey bees (Apis mellifera) on US farmland. *Sci Rep* **10**, 797, doi:10.1038/s41598-019-57225-w (2020).
Köhler, H.-R. & Triebskorn, R. Wildlife Ecotoxicology of Pesticides: Can We Track Effects to the Population Level and Beyond? *Science* **16**, 759-765

doi:10.1126/science.1237591 (2013).
8   Hemingway, J. & Ranson, H. Insecticide resistance in insect vectors of human disease. *Annu Rev Entomol* **45**, 371-391, doi:10.1146/annurev.ento.45.1.371 (2000).
9   Davies, T. G., Field, L. M., Usherwood, P. N. & Williamson, M. S. A comparative study of voltage-gated sodium channels in the Insecta: implications for pyrethroid resistance in Anopheline and other Neopteran species. *Insect Mol Biol* **16**, 361-375, doi:10.1111/j.1365-2583.2007.00733.x (2007).
10   Busvine, J. R. Mechanism of resistance to insecticide in houseflies. *Nature* **168**, 193-195, doi:10.1038/168193a0 (1951).

11   Grigoraki, L. *et al.* CRISPR/Cas9 modified An. gambiae carrying kdr mutation L1014F
functionally validate its contribution in insecticide resistance and combined effect with
metabolic enzymes. *PLoS Genet* **17**, e1009556, doi:10.1371/journal.pgen.1009556 (2021).
12   Reimer, L. *et al.* Relationship between kdr mutation and resistance to pyrethroid and DDT
insecticides in natural populations of Anopheles gambiae. *J Med Entomol* **45**, 260-266,
doi:10.1603/0022-2585(2008)45[260:rbkmar]2.0.co;2 (2008).
13   Hemingway, J. *et al.* Averting a malaria disaster: will insecticide resistance derail malaria
control? *Lancet* **387**, 1785-1788, doi:10.1016/S0140-6736(15)00417-1 (2016).
14   Donnelly, M. J., Isaacs, A. T. & Weetman, D. Identification, Validation, and Application of
Molecular Diagnostics for Insecticide Resistance in Malaria Vectors. *Trends Parasitol* **32**, 197-
206, doi:10.1016/j.pt.2015.12.001 (2016).
15   Msugupakulya, B. J. *et al.* Changes in contributions of different Anopheles vector species to
malaria transmission in east and southern Africa from 2000 to 2022. *Parasit Vectors* **16**, 408,
doi:10.1186/s13071-023-06019-1 (2023).
16   UNEP. *Stockholm Convention on persistent organic pollutants (POPs)*,
<https://chm.pops.int/TheConvention/Overview/TextoftheConvention/tabid/2232/Default.
aspx> (2005).
17   WorldBank. *Saying Goodbye to Tanzania's Old Pesticide Stocks*,
<https://www.worldbank.org/en/news/feature/2012/09/12/saying-goodbye-to-tanzania-s-
old-pesticide-stocks> (2012).
18   Soderlund, D. M. & Knipple, D. C. The molecular biology of knockdown resistance to
pyrethroid insecticides. *Insect Biochem Mol Biol* **33**, 563-577, doi:10.1016/s0965-
1748(03)00023-7 (2003).
19   Rinkevich, F. D., Du, Y. & Dong, K. Diversity and Convergence of Sodium Channel Mutations
Involved in Resistance to Pyrethroids. *Pestic Biochem Physiol* **106**, 93-100,
doi:10.1016/j.pestbp.2013.02.007 (2013).
20   Irving, H. & Wondji, C. S. Investigating knockdown resistance (kdr) mechanism against
pyrethroids/DDT in the malaria vector Anopheles funestus across Africa. *BMC Genet* **18**, 76,
doi:10.1186/s12863-017-0539-x (2017).
21   Hancock, P. A. *et al.* Modelling spatiotemporal trends in the frequency of genetic mutations
conferring insecticide target-site resistance in African mosquito malaria vector species. *BMC*
Biol **20**, 46, doi:10.1186/s12915-022-01242-1 (2022).
22   Clarkson, C. S. *et al.* The genetic architecture of target-site resistance to pyrethroid
insecticides in the African malaria vectors Anopheles gambiae and Anopheles coluzzii. *Mol*
Ecol **30**, 5303-5317, doi:10.1111/mec.15845 (2021).
23   Mitchell, S. N. *et al.* Metabolic and target-site mechanisms combine to confer strong DDT
resistance in Anopheles gambiae. *PLoS One* **9**, e92662, doi:10.1371/journal.pone.0092662
(2014).
24   Williamson, M. S., Martinez-Torres, D., Hick, C. A. & Devonshire, A. L. Identification of
mutations in the housefly para-type sodium channel gene associated with knockdown
resistance (kdr) to pyrethroid insecticides. *Mol Gen Genet* **252**, 51-60,
doi:10.1007/bf02173204 (1996).
25   Ranson, H. *et al.* Identification of a point mutation in the voltage-gated sodium channel gene
of Kenyan Anopheles gambiae associated with resistance to DDT and pyrethroids. *Insect Mol*
Biol **9**, 491-497, doi:10.1046/j.1365-2583.2000.00209.x (2000).
26   Martinez-Torres, D. *et al.* Molecular characterization of pyrethroid knockdown resistance
(kdr) in the major malaria vector Anopheles gambiae s.s. *Insect Mol Biol* **7**, 179-184,
doi:10.1046/j.1365-2583.1998.72062.x (1998).
27   Rogers, A. R. & Huff, C. Linkage disequilibrium between loci with unknown phase. *Genetics*
182, 839-844, doi:10.1534/genetics.108.093153 (2009).

28 Lynd, A. *et al.* LLIN Evaluation in Uganda Project (LLINEUP) - Plasmodium infection
prevalence and genotypic markers of insecticide resistance in Anopheles vectors from 48
districts of Uganda. *medRxiv*, doi:10.1101/2023.07.31.23293323 (2023).
29 Weedall, G. D. *et al.* A cytochrome P450 allele confers pyrethroid resistance on a major
African malaria vector, reducing insecticide-treated bednet efficacy. *Sci Transl Med* **11**,
doi:10.1126/scitranslmed.aat7386 (2019).
30 Coetzee, M. & Koekemoer, L. L. Molecular systematics and insecticide resistance in the
major African malaria vector Anopheles funestus. *Annu Rev Entomol* **58**, 393-412,
doi:10.1146/annurev-ento-120811-153628 (2013).
31 WHO. *Guidelines for malaria*, <https://app.magicapp.org/#/guideline/LwRMXj/rec/j7Krlj>
(2023).
32 Kaindoa, E. W. *et al.* Interventions that effectively target Anopheles funestus mosquitoes
could significantly improve control of persistent malaria transmission in south-eastern
Tanzania. *PLoS One* **12**, e0177807, doi:10.1371/journal.pone.0177807 (2017).
33 Weedall, G. D. *et al.* A cytochrome P450 allele confers pyrethroid resistance on a major
African malaria vector, reducing insecticide-treated bednet efficacy. *Sci. Transl. Med* (2019).
34 Oxborough, R. M. Trends in US President's Malaria Initiative-funded indoor residual spray
coverage and insecticide choice in sub-Saharan Africa (2008-2015): urgent need for
affordable, long-lasting insecticides. *Malar J* **15**, 146, doi:10.1186/s12936-016-1201-1
(2016).
35 FAO. *Prevention and disposal of obsolete pesticides*
<https://www.fao.org/agriculture/crops/obsolete-pesticides/where-stocks/africa-
stocks/en/> (2005).
36 UNEP. *National Implementation Plans*,
<https://www.pops.int/Implementation/NationalImplementationPlans/NIPTransmission/tab
id/253/Default.aspx> (2005).
37 WorldBank. *Africa Stockpiles Program Project Performance and Assessment Report*,
<chrome-
https://documents1.worldbank.org/curated/en/168841477341223021/pdf/108524-PPAR-
PUBLIC.pdf> (2016).
38 Elibariki, R. & Maguta, M. M. Status of pesticides pollution in Tanzania - A review.
Chemosphere **178**, 154-164, doi:10.1016/j.chemosphere.2017.03.036 (2017).
39 Carson, R. *Silent Spring*. (Mariner Books Classics; Anniversary edition, 1962).
40 Ayala, D. *et al.* The genome sequence of the malaria mosquito, Anopheles funestus, Giles,
1900. *Wellcome Open Res* **7**, 287, doi:10.12688/wellcomeopenres.18445.2 (2022).
41 Harris, A. M., Garud, N. R. & DeGiorgio, M. Detection and Classification of Hard and Soft
Sweeps from Unphased Genotypes by Multilocus Genotype Identity. *Genetics* **210**, 1429-
1452, doi:10.1534/genetics.118.301502 (2018).
42 Waskom, M. L. seaborn: statistical data visualization *Journal of Open Source Software* **6**,
3021, doi:DOI: 10.21105/joss.03021 (2021).
43 Koekemoer, L. L., Kamau, L., Hunt, R. H. & Coetzee, M. A cocktail polymerase chain reaction
assay to identify members of the Anopheles funestus (Diptera: Culicidae) group. *Am J Trop
Med Hyg* **66**, 804-811, doi:10.4269/ajtmh.2002.66.804 (2002).

517