1 Research article

2 Dimeric [R25C]PTH(1-34) Activates the Parathyroid Hormone-1 Receptor *in vitro*

3 and Stimulates Bone Formation in Osteoporotic Female Mice

Minsoo Noh[1, 2, *], Xiangguo Che[3, *], Xian Jin[3], Dong-Kyo Lee[3], Hyun-Ju Kim[3], Doo Ri Park[4], Soo

Young Lee[4], Hunsang Lee[2], Thomas Gardella[5], Je-Yong Choi[3, #], Sihoon Lee[1, #]

[1]Department of Internal Medicine and Laboratory of Genomics and Translational Medicine, Gachon

University College of Medicine, Incheon 21565, Republic of Korea

[2]Department of Life Sciences, Korea University, Seoul 02841, Korea

[3]Department of Biochemistry and Cell Biology, Cell and Matrix Research Institute, School of Medicine,

Kyungpook National University, Daegu 41944, Republic of Korea.

[4]Department of Life Sciences, Multitasking Macrophage Research Center, Ewha Womans University,

Seoul 03760, Korea

[5]Endocrine Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114,

USA

*These two authors are equally contributed to this manuscript.

#Correspondence:

Je-Yong Choi D.D.S., Ph.D., Department of Biochemistry and Cell Biology, Cell and Matrix Research

Institute, School of Medicine, Kyungpook National University, Daegu 41944, Republic of Korea. Tel.:

+82-53-420-4823; Fax: +82-53-422-1466; E-mail: jechoi@knu.ac.kr

&

Sihoon Lee, M.D., Ph.D., Department of Internal Medicine and Laboratory of Molecular Endocrinology,

Gachon University School of Medicine, Incheon 21565, Republic of Korea. Tel.: +82-32-458-2646;

Fax: +82-32-460-2381; E-mail: shleemd@gachon.ac.kr

Abstract

Osteoporosis, characterized by reduced bone density and strength, increases fracture risk,

pain, and limits mobility. Established therapies of Parathyroid hormone (PTH) analogs effectively

promote bone formation and reduce fractures in severe osteoporosis, their use is limited by potential

adverse effects. In the pursuit of safer osteoporosis treatments, we investigated ^{R25C}PTH, a PTH

variant wherein the native arginine at position 25 is substituted by cysteine. These studies were

prompted by our finding of high bone mineral density in a hypoparathyroidism patient with the R25C

homozygous mutation, and we explored its effects on PTH type-1 receptor (PTH1R) signaling in cells

and bone metabolism in mice. Our findings indicate that ^{R25C}PTH(1-84) forms dimers both

intracellularly and extracellularly, and the synthetic dimeric peptide, ^{R25C}PTH(1-34), exhibits altered

activity in PTH1R-mediated cAMP response. Upon a single injection in mice, dimeric ^{R25C}PTH(1-34)

induced acute calcemic and phosphaturic responses comparable to PTH(1-34). Furthermore,

repeated daily injections increased calvarial bone thickness in intact mice and improved trabecular

and cortical bone parameters in ovariectomized (OVX) mice, akin to PTH(1-34). The overall results

reveal a capacity of a dimeric PTH peptide ligand to activate the PTH1R *in vitro* and *in vivo* as PTH,

suggesting a potential path of therapeutic PTH analog development.

Key words: Parathyroid hormone, PTH type 1 receptor, Bone formation, Osteoporosis

Introduction

Osteoporosis is a prevalent global bone disorder characterized by low bone mineral density (BMD), causing weakened bones prone leading to fragility fractures, particularly in the spine, hip, and wrist. The development of osteoporosis is influenced by factors including gender, being more prevalent in women; hormonal changes, like decreased estrogen levels during menopause; and age, with heightened susceptibility post-menopause in women contributing to bone loss. Recent meta-analysis of previous studies indicates a global osteoporosis prevalence of 23.1 % among women and 11.7 % among men (1, 2). Osteoporosis stands as a noteworthy risk factor that poses challenges to the preservation of autonomous mobility and overall well-being within an aging society. There is thus a pressing need for safe and efficacious therapies for osteoporosis that mitigate fractures, alleviate associated symptoms, and preserve physical functionality.

Anti-resorptive agents (e.g., bisphosphonates, denosumab, and romosozumab) encompass one therapeutic approach that aims to specifically counteract the declines in bone mass by tempering the balance between bone resorption and formation (3-5). It is pertinent to acknowledge, however, that the prolonged use of most of such agents is limited due to potential long-term side effects. Furthermore, most anti-resorptive therapies cannot stimulate new bone formation (6). In contrast, bone anabolic agents, such as parathyroid hormone (PTH) and its analogs, such as teriparatide (recombinant human PTH(1-34)), increase BMD by stimulating bone formation more than bone resorption (7). PTH has an exceptionally short half-life in the blood of approximately 2-4 minutes (8, 9), which helps in avoiding excessive increases in blood calcium levels that can otherwise limit the utility of PTH-related medications, while yet inducing a desired anabolic effect on bone. It is also worth noting that studies in rodents reveal that long-term administration of a PTH anabolic agent can lead to bone overgrowth, osteosarcoma, as well as hypercalcemia (10, 11). Consequently, a goal of ongoing research is to uncover the underlying molecular mechanisms driving the anabolic and catabolic effects of PTH to thereby secure more effective therapeutic alternatives for osteoporosis (12).

PTH is produced and secreted by the parathyroid glands as a straight-chain monomeric polypeptide of 84 amino acids (13-15). It plays a vital role in maintaining calcium and phosphate equilibrium by acting on the PTH1R, a class B G protein-coupled receptor (GPCR) (16-18) that is

expressed primarily in cells of bone and kidney (19). The orchestrated downstream effects of PTH in

target cells act to ensure optimal bone health and the maintenance of the ambient blood calcium and

phosphate concentrations required for proper nerve conduction, muscle activity, and systemic cellular

communication, whereas disturbances in this system can lead to multiple disorders.

Central to the PTH signaling cascade is the activation of intracellular G proteins (20), most

prominently Gsα, which in turn activates transmembrane adenylyl cyclases, leading to the synthesis

of the second messenger cyclic AMP (cAMP), and the activation of cAMP-dependent protein kinase A

(PKA) (21, 22). PTH can also activate other second messenger cascades, including the

Gαq/phospholipase C (PLC) / inositol 1,4,5-trisphosphate (IP3), diacylglycerol (DAG), and protein

kinase C (PKC) signaling pathways (23, 24), highlighting the diverse biology of PTH and the PTH1R

(25). Adding to this, the PTH1R also mediates the actions of parathyroid hormone-related protein

(PTHrP), a development protein that acts in the formation of bones and other tissues. PTHrP shares

homology with PTH in the first 34 amino acids, which encompass the receptor-binding portions of the

two respective ligands. The PTH1R thus has an intrinsic capacity for dual ligand recognition, which

opens possibilities for exploring new modes of therapeutic development for diseases such as

osteoporosis and hypoparathyroidism (19, 20, 26-29). Pharmacologically, the PTH1R can adopt at

least two distinct ligand-binding conformations, RG and R^0, the selectivity for which can lead to altered

modes of signaling *in vitro* and *in vivo* for peptides such as PTH, PTHrP, and various hybrid analogs

(30-34) .

The current study extends our prior investigation in which we identified in a patient with

chronic hypocalcemia and hyperphosphatemia a mutation that changes the arginine at position 25 in

the mature PTH(1-84) polypeptide to cysteine ([R25C]PTH) (35, 36). Antibody assays revealed the

[R25C]PTH mutant protein to be present in the patient's blood at markedly elevated levels. We now have

found that this patient expressing the [R25C]PTH variant has higher-than-normal BMD. We further

characterize the [R25C]PTH protein and find that it can manifest in two distinct molecular forms: as a

monomer and as a dimer, and we demonstrate that a dimeric [R25C]PTH(1-34) synthetic peptide retains

agonistic properties on the PTH1R that are driven by a moderate selectivity for the RG vs. R^0 receptor

conformation. Finally, we demonstrate in mice that [R25C]PTH(1-34) can induce skeletal responses that

are similar to those induced by PTH(1-34), but without triggering an excessive hypercalcemic

response. Considering the proven bone-anabolic capacity of several established PTH agonist ligands,

and the need for safe, long-term treatments for skeletal disorders, our studies on dimeric ^{R25C}PTH(1-

34) suggest alternative strategies to consider in such drug development programs.

Results

Dimerization of ^{R25C}PTH(1-84)

In our previous studies, we showed that synthetic monomeric peptide, ^{R25C}PTH(1-34), as compared to PTH(1-34), exhibits a moderately diminished PTH1R-binding affinity and decreased cAMP signaling potency *in vitro*, and that with long-term infusion in mice, ^{R25C}PTH(1-34) leads to only minimal calcemic and phosphaturic effects, which corroborates the hypocalcemia seen in the original patient, despite the significantly elevated levels of ^{R25C}PTH in the plasma (35). In subsequent analysis of this patient, we found a particularly high BMD (Supplemental figure 1), which prompted us to further characterize the functional properties of ^{R25C}PTH, as described herein. We produced recombinant PTH(1-84) with or without the R25C mutation by expression of the corresponding cDNA in HEK293T cells (Figure 1). We considered the possibility that the introduction of a sole new cysteine within the polypeptide chain of ^{R25C}PTH(1-84) might induce homologous bimolecular dimerization through a disulfide bond involving the thiol functional group in each monomer (37, 38). To specifically investigate this, we designed the cDNA constructs to express either pre-pro-PTH(1-115)-3xFLAG or ^{R56C}pre-pro-PTH(1-115)-3xFLAG, such that after intracellular processing and cleavage of the pre-pro regions, the mature PTH(1-84)-3xFLAG or ^{R25C}PTH(1-84)-3xFLAG peptides would be generated upon transfection in HEK293T cells (Figure 1C) (39-41). We performed western blot analysis of total cell lysates and conditioned culture media collected from the transfected cells to specifically assess the possible presence of a disulfide-bonded dimeric form. Each sample was thus prepared in either reduced or non-reduced form and proteins were detected using an anti-flag antibody. The results demonstrated the presence of both a low molecular weight monomeric, and in the non-reduced samples, a higher molecular weight dimeric form of the ^{R25C}PTH(1-84)-3XFLAG protein in both the cell lysate and extracellular conditioned medium fractions, and the dimer appeared to be at an elevated proportion in the medium relative to the lysate (Figure 1D). To control for potential artifact effects attributed to the 3xFLAG tag, we utilized plasmid constructs pcDNA3.0-(pre-pro-PTH)-IRES and pcDNA3.0-(^{R56C}pre-pro-PTH)-IRES encoding non-tagged PTH variants and an anti-PTH(39-84) antibody for Western blot analysis, which again confirmed the presence of the dimer in total HEK293T cell lysates (Supplemental figure 2).

We observed in the above studies that the expression level of ^{R25C}PTH(1-84) was higher

than that of wild-type PTH(1-84) in both cell lysate and medium, which we considered might be due to

an intrinsic enhancement in protein stability and resistance to protein degradation in the dimeric

molecule. To address this, we treated the cells with the proteasome inhibitor MG132, which acts by

forming a hemiacetal with the hydroxyl groups of active site threonine residues, and compared the

expression levels of wild-type PTH(1-84) and ^{R25C}PTH(1-84) in the treated vs. untreated cells. The

results indicated that while the expression level of wild-type PTH was increased by MG132 treatment,

it did not reach the level of ^{R25C}PTH, suggesting that the difference in expression is not related to a

difference in sensitivity to proteasome-mediated degradation (Supplemental figure 3).

Overall, we have confirmed that ^{R25C}PTH(1-84) can form a dimeric structure, and the

^{R25C}PTH(1-84) secreted outside the cells predominantly exists in dimeric form. Thus, utilizing dimer

^{R25C}PTH(1-84) in the analysis would be more relevant to understanding the actual function of ^{R25C}PTH.

Consequently, we aim to conduct further validation using dimeric ^{R25C}PTH in our subsequent

investigations.

Functional characterization of dimeric ^{R25C}PTH(1-34) *in vitro*

To explore the functional properties of dimeric ^{R25C}PTH, we conducted experiments using

synthetic peptides of PTH(1-34), ^{R25C}PTH(1-34) (monomeric) and disulfide-bonded dimeric ^{R25C}PTH(1-

34). First, we examined the receptor-binding affinity of these ligands by performing competition

experiments using membranes prepared from HEK293-derived GP-2.3 cells that stably express the

human PTH1R and assay formats designed to assess binding to either the G protein-uncoupled R^0 or

G protein-coupled RG receptor conformation. The results revealed that monomeric ^{R25C}PTH(1-34)

bound to both the R^0 and RG conformations with comparable, albeit slightly weaker affinity as

compared to PTH(1-34), while dimeric ^{R25C}PTH(1-34) bound to each conformation with weaker affinity

than did the monomeric form while showing an apparent selectivity for higher binding to the RG vs R^0

conformation of PTH1R (Figure 2A).

To investigate the signaling properties of the ligands, we measured the changes in the

increase in intracellular levels of cAMP induced by each ligand in an osteoblastic SaOS-2 derived cell

line (SGS-72 cells) that stably expresses the Glosensor cAMP reporter. These assays revealed that

each ligand dose-dependently increased the cAMP levels in the cells, detected as an increase in

luminescence in an Envision plate reader, and while the potencies were moderately and more

substantially reduced for the monomeric and dimeric forms of the ligand, respectively, as compared to

PTH(1-34), the maximum response attained by each ligand were comparable (Figure 2B). Dimeric

[R25C]PTH(1-34) thus retains signaling functionality at the PTH1R that is characterized by a potency

approximately commensurate with its affinity for binding to the RG PTH1R conformation.

Effect of single injection of dimeric [R25C]PTH(1-34) on calcium and phosphate regulation in mice

To assess whether dimeric [R25C]PTH can function *in vivo*, we injected the ligand, and in

parallel either vehicle or PTH(1-34) (each peptide at a dose of 50 nmol/kg) into CD1 female mice and

measured levels of ionized calcium (Ca^{2+}) in the blood (n = 6 mice/group), inorganic phosphate (Pi) in

plasma (n = 12 mice/group), and the excretion of Pi into urine (n = 6 mice/group). Blood Ca^{2+} levels

were measured at serial time points of pre-injection, 1, 2, 4, and 6 hours post-injection. Both PTH(1-

34) and dimeric [R25C]PTH(1-34) induced increases in blood Ca^{2+} levels that were significant, relative to

the levels in vehicle-injected mice, at 1 and 2 hours post-injection, and the levels then returned to the

baseline levels of vehicle control mice by 4 hours (Figure 3A). Plasma Pi levels were measured in

samples obtained pre-injection, at 6 minutes, and 1, 2, and 6 hours post-injection. PTH(1-34) induced

a significant decrease in plasma Pi at 1-hour post-injection, and the levels subsequently returned to

baseline by 2 hours. Injection of dimeric [R25C]PTH(1-34) resulted in a slight decrease in plasma Pi at 2

hours post-injection, but the effect was not significant (Figure 3B). Consistent with this trend, Pi levels

in the urine of mice injected with dimeric [R25C]PTH(1-34) were increased significantly at 2 hours post-

injection and then gradually returned to baseline levels (Figure 3C). These results thus indicate that

dimeric [R25C]PTH can elicit calcemic and phosphaturic responses *in vivo* that are fully with those

expected for an injected PTH1R agonist ligand.

Effect of dimeric ^{R25C}PTH(1-34) on bone calvariae in mice

To initially assess the effects that short-term treatment of dimeric ^{R25C}PTH(1-34) can have on

bone, we injected 8-week-old male C57BL/6 mice once a day for six days (days 1-6), with either

dimeric ^{R25C}PTH(1-34), PTH(1-34) or vehicle, and after 10 days without treatment (day 16) followed by

euthanasia, we isolated the calvariae for histological analysis of new bone formation. Specifically, we

examined sections stained with hematoxylin and eosin (H&E) to assess the width of newly formed

bone areas along the edge of each sample. These regions exhibited a more vivid coloration compared

to the surrounding existing bone tissue, demarcated by a dotted line for clarity (Figure 4A).

Measurements were taken below and above the dissection area where new bone had formed, and

these measurements were then utilized to calculate the mean values for further analysis. These

analyses revealed that both PTH(1-34) and dimeric ^{R25C}PTH(1-34) significantly increased the width of

the new bone area by approximately four-fold, as compared to the vehicle group (Figure 4B). These

findings thus support a capacity of dimeric ^{R25C}PTH(1-34) to induce new bone formation *in vivo,*

similar to PTH, despite molecular and structural changes.

Effect of dimeric ^{R25C}PTH on bone mass in osteoporotic mice

To more directly assess the impact of dimeric ^{R25C}PTH(1-34) on bone mass, we administered

it to ovariectomized (OVX) mice, which serve as a well-established model for postmenopausal

osteoporosis. The OVX mice were injected 5 times per week for 4 weeks with either the dimeric ligand

(OVX + dimeric ^{R25C}PTH(1-34), PTH(1-34) (OVX + PTH(1-34)) or vehicle (OVX + vehicle, OVX-

controls), and sham-operated (Sham) mice were used as further controls. Mice were euthanized at

the end of the injection period and tissue samples were isolated for analysis. Quantitative micro-

computed tomography (μ-CT) analysis of the femurs obtained from each group revealed that, as

compared to OVX + vehicle controls, treatment with PTH(1–34) increased femoral trabecular bone

volume fraction (Tb.BV/TV) by 121%, cortical bone volume fraction (Ct.BV/TV) by 128%, cortical

thickness (Ct.Th) by 115%, cortical area (Ct.Ar) by 110%, and cortical area fraction (Ct.Ar/Tt.Ar) by

118%, while decreased total tissue area (Tt.Ar) by 93% (Figure 5A and 5B). Treatment with dimeric

^{R25C}PTH(1-34) had similar effects on the femoral cortical bone parameters, as it increased Ct.BMD by

104%, Ct.BV/TV by 125%, Ct.Th by 107%, and Ct.Ar/Tt.Ar by 116%, while decreased Tt.Ar 86% (Figure 5G). Considering the reduction of Tt.Ar and no change of Ct.Ar compared to the OVX+vehicle controls, the increase of Ct.Ar/Tt.Ar indicates a decrease in bone marrow space. The increase in cortical bone BMD was significant with dimeric [R25C]PTH(1-34) but not with PTH(1-34), whereas an increase in femoral trabecular bone was only observed with PTH(1-34).

The effects of the treatments on bone strength were assessed by conducting a three-point bending test on femurs isolated from the mice. The maximum load parameter was significantly decreased in the femurs from OVX-control versus Sham mice and was significantly increased, relative to the OVX-controls, by treatment with the PTH(1-34), but not dimeric [R25C]PTH(1-34) (Figure 5C). The slope parameter was significantly decreased in the femurs from OVX-control versus Sham mice and tended to increase versus OVX-controls by treatment with either PTH(1-34) or [R25C]PTH(1-34), but the changes were not significant.

We further analyzed the levels of bone metabolism markers in the serum obtained from the mice at the study endpoint (Figure 5D). The levels of serum calcium were within the normal range in all treatment groups, while serum phosphate levels were modestly increased in the OVX mice treated with PTH(1-34) or dimeric [R25C]PTH(1-34) as compared to with vehicle, but the effect was significant only with PTH(1-34). The serum levels of CTX-1, a bone resorption marker (42), were elevated in each of the OVX groups versus the Sham group, and tended to be lower in the OVX-PTH(1-34) treatment group, but the change relative to OVX-vehicle was not significant (Figure 5D). Interestingly, serum P1NP and alkaline phosphatase (ALP) levels were significantly increased in dimeric [R25C]PTH(1-34)-treated group, compared to the OVX-vehicle group (43, 44). Histological staining of proximal tibial sections for tartrate-resistant acid phosphatase (TRAP) activity, a marker of osteoclast-mediated bone resorption, revealed an apparent increase in this activity in bones of the OVX mice, as compared to those in Sham control mice, reflecting a heightened rate of bone turnover, as also suggested by the increased levels of serum CTX-1 in the OVX mice, and the TRAP staining appeared to be reduced in the tibiae of the OVX mice treated with PTH(1-34) (Figure 5D and E). Further histomorphometric analysis confirmed a significant increase in the osteoclast surface area relative to bone surface area (Oc.S/BS) in the proximal tibiae of the OVX-vehicle mice, relative to that in the

Sham-control mice, and this parameter was significantly decreased by treatment with PTH(1-34), but

not with dimeric ^{R25C}PTH(1-34) (Figure 5F).

We then analyzed the bone microstructure in the lumbar vertebrae through von Kossa

staining of histological sections and histomorphometric quantification (Figure 6A). The trabecular bone

volume fraction (Tb BV/TV, %), and trabecular number (Tb N) were significantly reduced in the OVX-

vehicle group, as compared to the Sham group, and treatment with either PTH(1-34) or dimeric

^{R25C}PTH(1-34) resulted in a significant increase in each of these parameters, as well as a concomitant

reduction in trabecular separation (Tb Sp), as compared to the respective parameters in the OVX-

vehicle group (Figure 6B).

Dynamic bone histomorphometry was also performed on the vertebrae to evaluate rates of

bone formation (Figure 6C). The trabecular mineral apposition rate (MAR) and cortical MAR were

significantly increased in both the OVX-PTH(1-34) and OVX-dimeric ^{R25C}PTH(1-34) groups, as

compared to in the OVX-vehicle group, and although there was a tendency for an increase in the

bone formation rate (BFR/BS) in both the trabecular and cortical bone with either PTH(1-34) or

dimeric ^{R25C}PTH(1-34) treatment, the differences were not statistically significant, as compared to the

OVX control (Figure 6D).

In summary, injection of dimeric ^{R25C}PTH into osteoporotic OVX mice resulted in significant

increases in cortical bone in the femurs and trabecular bone in the vertebrae, as well as significant

increases in osteoblast function and serum markers of bone formation, ALP and P1NP, without

inducing excessive bone resorption or hypercalcemia.

Discussion

In this study, we show the introduction of a cysteine mutation at the 25th amino acid position

of mature parathyroid hormone (^{R25C}PTH) facilitates the formation of homodimers comprised of the

resulting dimeric ^{R25C}PTH peptide *in vitro*. This dimerization surprisingly was compatible with receptor

binding affinity and led to relatively minor deviations in functional behavior as assessed in our cell-

based assays and compared to the standard monomeric control PTH peptide. The homozygous

^{R25C}PTH   mutation   was   identified   in   patients   who   presented   with   hypocalcemia   and

hyperphosphatemia, despite elevated PTH levels (35, 45), and the mutation was found to impact the

bioactive region of PTH (35, 46-49). Our initial research on this ^{R25C}PTH mutant focused on the

monomeric state of the peptide, and these studies revealed relatively moderate decreases in PTH1R

binding affinity and cAMP-stimulating potency *in vitro* and moderately impaired calcemic effects upon

infusion in mice. Additional patient observations, however, revealed the patients to have a higher BMD

than anticipated for age-matched averages. This elevated bone mass, coupled with the elevated

serum PTH levels, prompted our further investigations into the properties of the mutant PTH, as

described herein.

Our investigations brought to light the capacity of the cysteine-25 mutation to induce dimer

formation in the otherwise monomeric PTH polypeptide as produced in transfected cells. This result

was established by comparing western blots of transfected cell lysates analyzed under reducing

versus non-reducing conditions of gel electrophoresis. Subsequently, we employed synthetic peptides

to further explore the functional properties of dimeric ^{R25C}PTH(1-34). The results of our current studies

show some divergence from our previous findings obtained using the monomeric counterpart,

^{R25C}PTH (1-34) . Compared to the monomer, dimeric ^{R25C}PTH(1-34) exhibited a more preferential

binding affinity for the RG versus R^0 PTH1R conformation, despite a diminished affinity for either

conformation. We also observed that the potency of cAMP production in cells was lower for dimeric

^{R25C}PTH as compared to the monomeric ^{R25C}PTH, in accordance with a lower PTH1R-binding affinity.

Previous reports indicated that a mutation at the 25th position of PTH results in the loss of calcium ion

allosteric effects on monomeric ^{R25C}PTH, leading to faster ligand dissociation, rapid receptor recycling,

and a shorter cAMP time course (50). Correspondingly, the weaker receptor affinity and reduced

cAMP production observed in dimeric [R25C]PTH suggest a possibility that the formation of a disulfide

bond at the 25th position significantly alters the function of PTH as a PTH1R ligand. While these

structural effects are not yet fully understood and need to be investigated further.

We further pursued *in vivo* applications in mice. We assessed the calcemic and

phosphatemic responses to a single injection of synthetic peptides of either PTH(1-34) or dimeric

[R25C]PTH(1-34) in intact mice. We found the dimer could induce increases in plasma calcium levels

that were at least as robust and as sustained as those induced by PTH(1-34) and were similarly

phosphaturic (Figure 3).

Activation of the canonical Gαs-cAMP-PKA signaling pathway is generally thought to underly

most of the biological responses induced by PTH1R activation, and our studies in SGS-72 cells

confirm that dimeric PTH can activate this pathway, albeit not as efficiently as a monomeric PTH

peptide. Arg25 resides in the 20-34 (C-terminal region) of PTH(1-34), which plays a significant role in

the binding of the ligand to the extracellular domain (ECD) of the PTH1R. In concert with the binding

of the 20-24 region of PTH to the ECD, the N-terminal portion of PTH engages the transmembrane

domain (TMD) of the PTH1R, inducing the conformational changes involved in G protein coupling and

cAMP production (34, 51-54). The precise binding mode used by dimeric [R25C]PTH to the PTH1R is

unknown, but it may be anticipated that it differs to some extent from that used by the monomeric

peptide, due, for example, to the changes in bulk molecular size and display of accessible functional

groups. Consequently, the receptor conformational changes and the modes of coupling to

downstream effectors may differ for the monomeric versus dimeric ligands, which could potentially

lead to altered signaling and biological responses *in vivo*. Whether such changes account for the

increased bone density observed in the patient with the homozygous [R25C]PTH mutation is unknown,

but cannot be presently ruled out.

The results of practical assessments of dimeric [R25C]PTH(1-34) for effects on calvarial bone

after short-term (6-days) injection into normal mice, and for effects on bone mass after long-term (4-

weeks) daily injection in osteoporotic OVX mice demonstrate the dimer can mediate significant

influences on bone metabolism alike to wild-type PTH. In OVX mice, PTH and dimeric [R25C]PTH

significantly increased bone formation markers P1NP and ALP. However, wild-type PTH, but not

dimeric [R25C]PTH, decreased bone resorption markers such as TRAP staining and CTX-1 levels,

suggesting different effects of wild-type monomeric PTH and dimeric [R25C]PTH. In this study, the

increase of bone resorption markers of dimeric [R25C]PTH compared to PTH may be one reason for low

bone strength. In addition, the data showed an increase in cortical bone area (Ct.Ar) in the PTH

treatment group but not in the PTH dimer treatment group. However, both dimeric and monomeric

PTH treatments reduced total tissue area (Tt.Ar), suggesting potential effects on bone growth in the

width of mice or humans. While further investigation is necessary, the current experimental data,

along with patient symptoms showing no bone resorption despite lifelong exposure to high levels of

[R25C]PTH, imply that the dimeric form of [R25C]PTH can provide a new insight into PTH and PTH analogs

with distinct functionalities compared to the wild-type PTH.

This study has several limitations. First, it is urgently necessary to determine whether dimeric

[R25C]PTH is present in human patient serum. Second, TRAP staining showed an inhibitory effect of

PTH treatment on the primary spongiosa area. However, the secondary spongiosa, which more

accurately reflects bone remodeling (55), was not examined due to the barely detectable bone in this

area in OVX-induced osteoporosis mouse models. Third, it is unclear whether similar bone

phenotypes exist between human [R25C]PTH patients and dimeric [R25C]PTH-treated mice, particularly

regarding low bone strength. Although the dimeric [R25C]PTH-treated group showed higher cortical BMD

compared to WT-Sham or PTH groups, there was no difference in bone strength compared to the

osteoporotic mouse model. Fourth, our study demonstrated that PTH or [R25C]PTH treatment decreased

circumferential length, which could affect bone growth in width. However, whether this phenotype is

also observed in patients treated with PTH or [R25C]PTH remains uncertain. Finally, we did not analyze

the [R25C]PTH mutant mouse model, which would allow us to compare phenotypes that most closely

resemble those of human patients.

Interestingly, the recent identification of a young patient in Denmark displaying homozygous

[R25C]PTH has opened avenues for observing the direct impacts of [R25C]PTH within the human biological

system (45). The continual monitoring and observation of patients will contribute to a more profound

comprehension of the long-term consequences associated with [R25C]PTH exposure. This extensive

observation is crucial in delineating the extended effects of this compound on individuals.

Consequently, by conducting thorough investigations to confirm the potential bone anabolic effect of

^{R25C}PTH, we hope to develop a novel bone anabolic agent with a targeted focus on the PTH1R.

Materials and Methods

Plasmid construction

The coding sequences (CDS) of pre-pro-PTH and the mutated form, ^{R56C}pre-pro-

PTH[^{R25C}PTH], were amplified using primers containing the attB site. These CDS fragments were

obtained from pcDNA3.0-(hpre-pro-PTH)-IRES and pcDNA3.0-(h^{R56C}pre-pro-PTH)-IRES, which were

used in the previous research conducted by Lee*, et al.* (35). Both CDS were introduced into donor

vector pDONR223 with Gateway™ BP Clonase™ II Enzyme mix kit (Invitrogen, USA). Then pre-pro-

PTH and ^{R56C}pre-pro-PTH were each cloned into pcDNA3.1-ccdB-3xFLAG-V5 with LR Gateway™ LR

Clonase™ II Enzyme mix (Invitrogen, USA) to construct pcDNA3.1-(pre-pro-PTH)-3xFLAG-V5, and

pcDNA3.1-(^{R56C}pre-pro-PTH)-3xFLAG-V5. All experimental procedures were done with the

manufacturer's instruction. pDONR223 was a gift from Kim Lab (Roswell Park Comprehensive

Cancer Center, USA), and pcDNA3.1-ccdB-3xFLAG-V5 was a gift from Taipale Lab (Donnelly Centre,

University of Toronto, Canada).

Cell culture

All cell lines were grown at 37°C in 5 % CO_2. HEK293T cells were cultured in Dulbecco's

Modified Eagle Medium (Cytiva™, HyClone DMEM/High glucose with L-Glutamine and HEPES, Cat

No. SH30243.01, USA) with 10 % FBS (Gibco™, Fetal Bovine Serum, certified, Origin: USA, Cat No.

16000044, Lot No. 2522247RP, USA) and 1 % penicillin-streptomycin (Cytiva™, HyClone Penicillin-

Streptomycin 100X solution, Cat No. SV30010, USA). Cell lines obtained from commercial sources

were not further authenticated.

Transfection

pcDNA3.1-(pre-pro-PTH)-3xFLAG-V5 and pcDNA3.1-(^{R56C}pre-pro-PTH)-3xFLAG-V5 were

each transfected into HEK293T with Lipofectamine™ 3000 Transfection Reagent (Invitrogen™, Cat

No. L3000001, USA) according to the manufacturer's instruction. After 48 hours of transfection, a

culture medium was collected and used for western blot as a secreted protein sample. The rest of the

cells were lysed by RIPA buffer (Thermo Scientific™, RIPA Lysis and Extraction Buffer, Cat No. 89900,

USA) following the manufacturer's instruction and used for western blot as total cell lysate sample.

Western blot

Protein samples were prepared in two types, reduced sample and non-reduced sample.

Reduced samples were a mixture of protein (secreted protein or cell lysate), sample buffer

(Invitrogen™, NuPAGE™ LDS Sample Buffer (4X), Cat No. NP0007, USA), with reducing agent

(Invitrogen™, 10X Bolt™ Sample Reducing Agent, Cat No. B0009, USA) and heated for 5 min at

95°C. Non-reduced samples were a mixture of protein, and sample buffer, without reducing agent,

and not heated. The protein samples were loaded onto 4 - 12 % Bis-Tris protein gels (GeneSTAR,

StarPAGE Bis-Tris 4-12 %, 15-well, Cat No. GPG4115), and ran with MOPS/SDS running buffer

(GeneSTAR, 20X MOPS/SDS Running Buffer, Cat No. GMB0080). Transfer to the membrane was

done by iBlot™ 2 Dry Blotting System (Invitrogen™, iBlot™ 2 Gel Transfer Device, Cat No. IB21001,

USA) with PVDF transfer stack (Invitrogen™, iBlot™ 2 Transfer Stacks-PVDF-mini, Cat No. IB24002,

USA) according to manufacturer's instruction. The membranes were blocked for 1 hour at room

temperature (RT) in 5 % skim-milk solution in Tris-buffered saline (TBS; 20 mM Tris-base, 500 mM

NaCl, pH 7.5) and then washed three times for 10 min each with tris-buffered saline with 0.05 %

tween-20 (TBST). The membranes were incubated with primary antibody for 1 hour at RT, then

washed three times for 10 min each with TBST. If needed, the membranes were incubated with

horseradish peroxidase (HRP)-conjugated secondary antibody for 1 hour at RT after primary antibody

incubation, then washed three times for 10 min each with TBST. After washing, the membranes were

rinsed and soaked in TBS. To develop a blot image, the membranes were treated with a

chemiluminescent substrate solution (Merck Millipore, Immobilon ECL Ultra Western HRP Substrate,

Cat No. WBULS0500, USA) according to the manufacturer's instruction. The blot images were

obtained by LAS 4000 mini (Cytiva, ImageQuant™ LAS 4000 mini, USA). The dilution condition for

the anti-FLAG with hHRP conjugated antibody (Sigma-Aldrich, Monoclonal ANTI-FLAG® M2-

Peroxidase (HRP) antibody produced in mouse, Cat No. A8592, USA) was 1:2,000. The dilution

condition for the anti-HSP90 primary antibody (Santa Cruz Biotechnology Inc., HSP 90 Antibody (AC-

16), Cat No. sc-101494, USA) was 1:5,000. The dilution condition for the anti-mouse secondary

antibody was 1:5,000. Each antibody was diluted in TBST with 1 % BSA solution.

Proteasome inhibition assay

HEK293T cells were seeded in culture dishes at approximately 60 % confluence, and they

were allowed to grow for about 20 to 24 hours prior to transfection. The transfection of pcDNA3.1-

(pre-pro-PTH)-3xFLAG-V5 and pcDNA3.1-(^{R56C}pre-pro-PTH)-3xFLAG-V5 was conducted following the

method mentioned earlier. After 24 hours of transfection, MG132, dissolved in DMSO, was added to

the cells to achieve a final concentration of 10 μM. For the mock treatment, DMSO alone was added.

The cells were then incubated for an additional 24 hours after MG132 treatment. Both the culture

medium and cell lysate were prepared for western blot analysis to assess the restored protein levels.

The western blot procedure was carried out as described in the previous section.

Peptide synthesis and quantification

419   Human PTH(1-34), ^{R25C}PTH(1-34), and dimeric ^{R25C}PTH(1-34) were chemically synthesized

420   by Anygen (Gwangju, Republic of Korea). The purity and mass of each peptide were analyzed by

421   high-performance liquid chromatography (HPLC) and matrix-assisted laser desorption ionization-time

422   of flight mass spectrometry (MALDI-TOF MS) in the manufacturer.

PTH1R competition binding assay

The binding of PTH and its analogs to G protein-uncoupled PTH1R (R^0 conformation) and G

protein-coupled PTH1R (RG conformation) was assessed using a competition method with

membranes prepared from GP-2.3 cells (HEK-293 cells stably expressing the hPTH1R). For tracer

radioligands, we utilized ^{125}I-PTH(1-34) and ^{125}I-MPTH(1-15). The unlabeled ligands tested were

PTH(1–34), ^{R25C}PTH(1-34), and dimeric ^{R25C}PTH(1-34). Binding to the R^0 conformation was assessed

using ^{125}I-PTH(1-34) as the tracer radioligand, while binding to the RG conformation was assessed

using ^{125}I-MPTH(1-15). The addition of unlabeled ligands PTH(1–34), ^{R25C}PTH(1-34), or dimeric

^{R25C}PTH(1-34) caused dissociation of the tracer radioligand from each receptor if it had affinity to the

receptors. Measurement of the dissociated ratio of the radioligand indicated the binding affinity

between PTH1R and each unlabeled ligand. Each experiment was replicated four times.

cAMP assay

437   To measure intracellular cAMP production, SGS-72 cells, derived from SaOS-2 cells and

438   stably expressing the Glosensor cAMP reporter, were utilized to measure intracellular cAMP

439   production. The detection of cAMP-dependent expression was performed using an Envision plate

440   reader (PerkinElmer, Waltham, MA, USA), based on luciferase-based luminescence, as previously

441   described by Maeda, *et al.* (56). Each measurement was replicated four times.

Animal model used in the study

CD1 female mice were purchased from Charles River Laboratories (Massachusetts, USA),

and all animal care and experimental procedures were conducted under the guidelines set by the

Institutional Animal Care and Use Committee (IACUC) at Massachusetts General Hospital (MGH).

The mice were housed in a specific pathogen-free environment, with 4-5 mice per cage, under a 12-

hour light cycle at a temperature of 22±2°C.

Eight-week-old C57BL/6N female mice were purchased from KOATECH (Gyeonggi-do,

Republic of Korea), and stabilized mice for 2 weeks. All animal care and experimental procedures

were conducted under the guidelines set by the Institutional Animal Care and Use Committees of

Kyungpook National University (KNU-2021-0101). The mice were housed in a specific pathogen-free

environment, with 4-5 mice per cage, under a 12-h light cycle at 22 ± 2°C. They were provided with

standard rodent chow and water ad libitum.

An ovariectomizing (OVX) osteoporosis mouse model was established using 10-week-old

C57BL/6N female mice. Following surgery, mice were divided into the following four groups (n = 6

mice/group) as follows: Sham, OVX control group, OVX + PTH (1–34) treated group (40 µg/kg/day),

and OVX + dimeric ^{R25C}PTH treated group (40-80 µg/kg/day). OVX mice were allowed to recover for 4

weeks after surgery. Afterward, PTH (1–34) or ^{R25C}PTH was injected subcutaneously 5 times a week

for 4 weeks. Micro-computed tomography (µ-CT) and histological analyses were performed on 4

groups at 18 weeks of age.

Acute injections

The peptides PTH(1-34) and dimeric ^{R25C}PTH(1-34) were diluted in a solution comprising

0.05% Tween 80, 10 mM citric acid, and 150 mM NaCl at a pH of 5.0. Intravenous injections of these

peptides were administered at doses ranging from 50-100 µg/kg into 9-week-old CD1 female mice. As

a control, mice received only the vehicle. Levels of ionized calcium (Ca2+) in the blood (n = 6

mice/group) were measured at serial time points of pre-injection, 1, 2, 4, and 6 hours post-injection.

Inorganic phosphate (Pi) in plasma (n = 12 mice/group), and the excretion of Pi into urine (n = 6

mice/group) were measured in samples obtained at serial time points of pre-injection, at 6 minutes,

and 1, 2, and 6 hours post-injection.

Calvarial injection mouse model

C57BL/6 male mice (8-week-old) were divided into the following three groups (n = 6

mice/group): control, PTH (1-34) treated group (80 µg/kg/day), and ^{R25C}PTH treated group (160

µg/kg/day). Subcutaneous injections of the respective drugs were administered once daily for 6 days.

On the sixteenth day, the mice were sacrificed, and their bone tissues were harvested and fixed in 10 %

formaldehyde at 4°C. The fixed bone tissues were then decalcified in PBS (pH 7.4) containing 0.5

moles of ethylenediaminetetraacetic acid (EDTA). Following decalcification, the tissues were

embedded in paraffin, and paraffinized tissues were sectioned to a thickness of 5-7 µm. Histological

analysis was performed using the sectioned tissue slides stained with hematoxylin and eosin (H&E).

The area of new bone formation, which displays a more intense coloration compared to the existing

bone tissue, was examined.

μ-CT analysis

Mouse femurs were fixed in a 4 % paraformaldehyde solution for 24 hours at 4°C. In μ-CT,

we used the Quantum FX μ-CT (Perkin Elmer, Waltham, MA, USA). The images were acquired at a

9.7 μm voxel resolution, with settings of 90 kV and 200 μA, a 10 mm field of view, and a 3-minute

exposure time. Serial cross-sectional images were reconstructed using the Analyze 12.0 software

(Overland Park, KS, USA). To ensure consistent analysis, identical regions of interest (ROIs) were

selected for the trabecular and cortical bones. Bone parameters and density were analyzed in the

region between 0.3–1.755 mm (150 slices) from the bottom of the growth plate. Analysis of bone

structure was performed using adaptive thresholding in CT Analyser. Thresholds for analysis were

determined manually based on grayscale values for each experimental group: trabecular bone: 3000;

cortical bone: 5000 for all samples. All bone parameters were evaluated according to the guidelines of

the American Society for Bone and Mineral Research (57).

Three-point bending test

The left femur of the mouse was immersed in 0.9 % NaCl solution, wrapped in gauze, and

stored at –20°C until ready for a three-point bending test. In this test, we placed the mouse femurs

horizontally with the anterior surface facing upwards, centered on the supports, and the compressive

force was applied vertically to the mid-shaft. The pressure sensor was positioned at a distance that

allowed maximum allowable pressure (200N) without interfering with the test (20.0 mm for the femur).

A miniature material testing machine (Instron, MA, U.S.A.) was used for this test. The crosshead

speed was decreased to 1 mm/min until failure. During the test, force-displacement data were

collected to determine the maximum load and slope of the bones.

Serum biochemistry analysis

Serum bone resorption and osteogenesis marker levels, specifically the C-terminal

telopeptide of type I collagen (CTX) and procollagen type I N-terminal propeptide (P1NP), were

assessed in mice from the Sham, OVX-control, PTH(1-34), and dimeric ^{R25C}PTH(1-34) groups by

using the enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instructions.

Additionally, their concentrations were determined using specific mouse CTX-1 and P1NP ELISA kits

(Cloud Clone, Wuhan, China) respectively.

Bone histological analyses

The tibiae were initially fixed in 4% paraformaldehyde at 4°C overnight. The following day,

samples were decalcified using 10% ethylenediaminetetraacetic acid (EDTA) solution (pH 7.4) for 4

weeks at 4°C. The decalcified tibiae were then embedded in paraffin and sectioned at 3 μm thick. For

TRAP staining, dehydrated paraffin sections were fixed in an acetone/ethanol mixture (1:1) for 1

minute, followed by complete air-drying at RT for 20 minutes. Thereafter, the sections were immersed

in TRAP reagent for 30 minutes at 37°C. In the histomorphometry analysis for TRAP staining, we

used the primary spongiosa for the trabecular ROI because of the barely detectable in the secondary

spongiosa of OVX model. Osteoclast surface per bone surface (Oc.S/BS) and Osteoclast number per

bone surface (Oc.N/BS) analysis followed the ASBMR guidelines (58).

Dynamic bone histomorphometric analysis

To conduct dynamic histomorphometry analysis, we injected the mice with 25 mg/kg body

weight of calcein (Sigma-Aldrich) or Alizarin Red S (Sigma-Aldrich) intraperitoneally before sacrifice,

with 3- or 10-day intervals between injections as previously described (59, 60). Briefly, femurs or

vertebrates were fixed in 4 % paraformaldehyde solution for 24 hours at 4°C. The following day, the

samples were washed with phosphate-buffered saline (PBS) solution and then dehydrated using a

gradient of ethanol (50 %, 70 %, 85 %, 90 %, and 100 %). Subsequently, we embedded the

dehydrated femurs or vertebrates in methyl methacrylate (Sigma) to prepare resin blocks. The resin

blocks were sectioned at 6 μm thick by using a Leica SP1600 microtome (Leica Microsystems,

Germany). The fluorescence signals of calcein (green) and Alizarin Red S (red) from ROIs were

captured using a fluorescence microscope (Leica, Wetzlar, Germany). For vertebral bone analysis,

bone mineralization was evaluated by von Kossa staining. The sections were placed in 2-

methoxyethyl acetate (Sigma-Aldrich) for 20 minutes, followed by rehydration with serial ethanol

solutions (100 %, 90 %, 80 %, 70 %, and 50 %) and distilled water for 2 minutes each. The sections

were subsequently dipped in a 1 % $AgNO_3$ solution (Sigma-Aldrich) for 5 minutes under ultraviolet

(UV) light photons, washed in distilled water for 5 minutes, and dipped in 5 % sodium thiosulfate

solution for 5 minutes to remove nonspecific binding. Finally, we covered the sections with mounting

solution and captured images by using a Leica microscope. The parameters of dynamic bone

histomorphometry were analyzed using the Bioquant Osteo 2019ME program (Bioquant Osteo,

Nashville, TN, USA).

Statistical analysis

Statistical analysis was performed in GraphPad Prism 10.1.2. The data are presented as the

mean ± standard error of the means (SEM). Statistically significant differences between the two

groups were determined using an analysis of variance (ANOVA). A p-value less than 0.05 was

considered statistically significant.

Author contributions

Conceptualization, Methodology: M.N., X.C., X.J., D-K.L., H-J.K., D.P., S.Y.L., H.L., T.J.G., J.-Y.C.,

and S.L.; Validation, Formal Analysis: M.N., and X.C.; Investigation, Resource: M.N., X.C., X.J., D.-

K.L., and D.P.; writing - Original Draft Preparation: M.N., and X.C.; Writing – Review & Editing: M.N.,

X.C., T.J.G., H.L., J.-Y.C., and S.L.; Visualization: M.N., and X.C.; Supervision: H.-J.K., H.L., J.-Y.C.,

and S.L.; Project Administration: J.-Y.C., and S.L.; Funding Acquisition: S.Y.L., H.L., J.-Y.C., and S.L.

Acknowledgments

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the

Korea government (MSIT) (2022R1A2C3006002 to S.L.; 2022R1F1A1074610 and

2022R1A4A1025913 to H.L.; RS-2023-00217798 and 2021R1A2C3003675 S.Y.L.;

2022R1A2C1006105 to J.-Y.C.), Gachon University Gil Medical Center (FRD2023-12 to S.L.), and

Korea University Grants.

References

1.   T. Sozen, L. Ozisik, N. C. Basaran, An overview and management of osteoporosis. *Eur J*
Rheumatol **4**, 46-56 (2017).

2.   N. Salari *et al.*, The global prevalence of osteoporosis in the world: a comprehensive
systematic review and meta-analysis. *J Orthop Surg Res* **16**, 609 (2021).

3.   S. Papapoulos *et al.*, Five years of denosumab exposure in women with postmenopausal
osteoporosis: results from the first two years of the FREEDOM extension. *J Bone Miner Res*
27, 694-701 (2012).

4.   D. M. Reid *et al.*, Zoledronic acid and risedronate in the prevention and treatment of
glucocorticoid-induced osteoporosis (HORIZON): a multicentre, double-blind, double-dummy,
randomised controlled trial. *Lancet* **373**, 1253-1263 (2009).

5.   M. R. McClung *et al.*, Romosozumab in postmenopausal women with low bone mineral
density. *N Engl J Med* **370**, 412-420 (2014).

6.   F. Cosman *et al.*, Romosozumab Treatment in Postmenopausal Women with Osteoporosis. *N*
Engl J Med **375**, 1532-1543 (2016).

7.   T. J. Martin, N. A. Sims, E. Seeman, Physiological and pharmacological roles of PTH and
PTHrP in bone using their shared receptor, PTH1R. *Endocrine reviews* **42**, 383-406 (2021).

8.   C. Bieglmayer, G. Prager, B. Niederle, Kinetic analyses of parathyroid hormone clearance as
measured by three rapid immunoassays during parathyroidectomy. *Clinical Chemistry* **48**,
1731-1738 (2002).

9.   G. W. Maier *et al.*, Parathyroid hormone after adenectomy for primary hyperparathyroidism. A
study of peptide hormone elimination kinetics in humans. *J Clin Endocrinol Metab* **83**, 3852-
3856 (1998).

10.   J. L. Vahle *et al.*, Skeletal changes in rats given daily subcutaneous injections of recombinant
human parathyroid hormone (1-34) for 2 years and relevance to human safety. *Toxicol Pathol*
30, 312-321 (2002).

11.   N. M. Appelman-Dijkstra, S. E. Papapoulos, From disease to treatment: from rare skeletal
disorders to treatments for osteoporosis. *Endocrine* **52**, 414-426 (2016).

12.   S. Khosla, L. C. Hofbauer, Osteoporosis treatment: recent developments and ongoing

challenges. *Lancet Diabetes Endocrinol* **5**, 898-907 (2017).

13.   E. Moallem, R. Kilav, J. Silver, T. Naveh-Many, RNA-Protein binding and post-transcriptional

regulation of parathyroid hormone gene expression by calcium and phosphate. *J Biol Chem*

273, 5253-5259 (1998).

14.   J. Silver, J. Russell, L. M. Sherwood, Regulation by vitamin D metabolites of messenger

ribonucleic acid for preproparathyroid hormone in isolated bovine parathyroid cells. *Proc Natl*

Acad Sci U S A **82**, 4270-4273 (1985).

15.   R. A. Chen, W. G. Goodman, Role of the calcium-sensing receptor in parathyroid gland

physiology. *Am J Physiol Renal Physiol* **286**, F1005-1011 (2004).

16.   J. F. Habener, M. Rosenblatt, J. T. Potts, Jr., Parathyroid hormone: biochemical aspects of

biosynthesis, secretion, action, and metabolism. *Physiol Rev* **64**, 985-1053 (1984).

17.   V. Veldurthy *et al.*, Vitamin D, calcium homeostasis and aging. *Bone Res* **4**, 16041 (2016).

18.   P. R. Kiela, F. K. Ghishan, Recent advances in the renal-skeletal-gut axis that controls

phosphate homeostasis. *Lab Invest* **89**, 7-14 (2009).

19.   H. Jüppner *et al.*, AG protein-linked receptor for parathyroid hormone and parathyroid

hormone-related peptide. *Science* **254**, 1024-1026 (1991).

20.   T. J. Gardella, H. Ju¨ ppner, Interaction of PTH and PTHrP with their receptors. *Reviews in*

Endocrine and Metabolic Disorders **1**, 317-329 (2000).

21.   L. S. Weinstein, S. Yu, D. R. Warner, J. Liu, Endocrine manifestations of stimulatory G protein

alpha-subunit mutations and the role of genomic imprinting. *Endocr Rev* **22**, 675-705 (2001).

22.   T. N. Feinstein *et al.*, Retromer terminates the generation of cAMP by internalized PTH

receptors. *Nat Chem Biol* **7**, 278-284 (2011).

23.   A. Iida-Klein *et al.*, Mutations in the second cytoplasmic loop of the rat parathyroid hormone

(PTH)/PTH-related protein receptor result in selective loss of PTH-stimulated phospholipase

C activity. *J Biol Chem* **272**, 6882-6889 (1997).

24.   R. Dunlay, K. Hruska, PTH receptor coupling to phospholipase C is an alternate pathway of

signal transduction in bone and kidney. *Am J Physiol* **258**, F223-231 (1990).

25.   A.-B. Abou-Samra *et al.*, Expression cloning of a common receptor for parathyroid hormone

and parathyroid hormone-related peptide from rat osteoblast-like cells: a single receptor

stimulates intracellular accumulation of both cAMP and inositol trisphosphates and increases

intracellular free calcium. *Proceedings of the National Academy of Sciences* **89**, 2732-2736

(1992).

26.   H. G. Bone *et al.*, Ten years' experience with alendronate for osteoporosis in postmenopausal

women. *N Engl J Med* **350**, 1189-1199 (2004).

27.   M. R. Rubin *et al.*, Therapy of Hypoparathyroidism With PTH(1-84): A Prospective Six Year

Investigation of Efficacy and Safety. *J Clin Endocrinol Metab* **101**, 2742-2750 (2016).

28.   K. K. Winer, J. A. Yanovski, G. B. Cutler, Jr., Synthetic human parathyroid hormone 1-34 vs

calcitriol and calcium in the treatment of hypoparathyroidism. *JAMA* **276**, 631-636 (1996).

29.   K. K. Winer *et al.*, Long-Term Parathyroid Hormone 1-34 Replacement Therapy in Children

with Hypoparathyroidism. *J Pediatr* **203**, 391-399 e391 (2018).

30.   M. Okazaki *et al.*, Prolonged signaling at the parathyroid hormone receptor by peptide ligands

targeted to a specific receptor conformation. *Proceedings of the National Academy of*

Sciences **105**, 16525-16530 (2008).

31.   A. A. Pioszak, N. R. Parker, T. J. Gardella, H. E. Xu, Structural basis for parathyroid hormone-

related protein binding to the parathyroid hormone receptor and design of conformation-

selective peptides. *J Biol Chem* **284**, 28382-28391 (2009).

32.   R. W. Cheloha, S. H. Gellman, J. P. Vilardaga, T. J. Gardella, PTH receptor-1 signalling-

mechanistic insights and therapeutic prospects. *Nat Rev Endocrinol* **11**, 712-724 (2015).

33.   S. R. Hoare, T. J. Gardella, T. B. Usdin, Evaluating the signal transduction mechanism of the

parathyroid hormone 1 receptor. Effect of receptor-G-protein interaction on the ligand binding

mechanism and receptor conformation. *J Biol Chem* **276**, 7741-7753 (2001).

34.   T. Dean, J. P. Vilardaga, J. T. Potts, Jr., T. J. Gardella, Altered selectivity of parathyroid

hormone (PTH) and PTH-related protein (PTHrP) for distinct conformations of the PTH/PTHrP

receptor. *Mol Endocrinol* **22**, 156-166 (2008).

35.   S. Lee *et al.*, A Homozygous [Cys25]PTH(1-84) Mutation That Impairs PTH/PTHrP Receptor

Activation Defines a Novel Form of Hypoparathyroidism. *J Bone Miner Res* **30**, 1803-1813

(2015).

36.   C. H. Bae *et al.*, A novel human PTH analog [Cys25] hPTH (1–34) restores bone mass in

ovariectomized mice. *The Journal of Clinical Endocrinology & Metabolism* **101**, 3700-3708 (2016).

37. R. R. Banerjee, M. A. Lazar, Dimerization of resistin and resistin-like molecules is determined by a single cysteine. *J Biol Chem* **276**, 25970-25973 (2001).

38. M. V. Trivedi, J. S. Laurence, T. J. Siahaan, The role of thiols and disulfides on protein stability. *Curr Protein Pept Sci* **10**, 614-625 (2009).

39. B. Kemper, J. F. Habener, R. C. Mulligan, J. T. Potts, Jr., A. Rich, Pre-proparathyroid hormone: a direct translation product of parathyroid messenger RNA. *Proc Natl Acad Sci U S A* **71**, 3731-3735 (1974).

40. T. J. Vasicek *et al.*, Nucleotide sequence of the human parathyroid hormone gene. *Proc Natl Acad Sci U S A* **80**, 2127-2131 (1983).

41. K. M. Wiren *et al.*, Mutations in signal sequence cleavage domain of preproparathyroid hormone alter protein translocation, signal sequence cleavage, and membrane-binding properties. *Mol Endocrinol* **3**, 240-250 (1989).

42. H. N. Rosen *et al.*, Serum CTX: a new marker of bone resorption that shows treatment effect more often than other markers because of low coefficient of variability and large changes with bisphosphonate therapy. *Calcif Tissue Int* **66**, 100-103 (2000).

43. F. Pagani, C. M. Francucci, L. Moro, Markers of bone turnover: biochemical and clinical perspectives. *J Endocrinol Invest* **28**, 8-13 (2005).

44. G. Wheater, M. Elshahaly, S. P. Tuck, H. K. Datta, J. M. van Laar, The clinical utility of bone marker measurements in osteoporosis. *J Transl Med* **11**, 201 (2013).

45. S. L. Andersen, A. L. Frederiksen, A. B. Rasmussen, M. Madsen, A. R. Christensen, Homozygous missense variant of PTH (c.166C>T, p.(Arg56Cys)) as the cause of familial isolated hypoparathyroidism in a three-year-old child. *J Pediatr Endocrinol Metab* **35**, 691-694 (2022).

46. A. Arnold *et al.*, Mutation of the signal peptide-encoding region of the preproparathyroid hormone gene in familial isolated hypoparathyroidism. *J Clin Invest* **86**, 1084-1087 (1990).

47. D. B. Parkinson, R. V. Thakker, A donor splice site mutation in the parathyroid hormone gene is associated with autosomal recessive hypoparathyroidism. *Nat Genet* **1**, 149-152 (1992).

683   48.   T. Sunthornthepvarakul, S. Churesigaew, S. Ngowngarmratana, A novel mutation of the signal
684   peptide of the preproparathyroid hormone gene associated with autosomal recessive familial
685   isolated hypoparathyroidism. *J Clin Endocrinol Metab* **84**, 3792-3796 (1999).

686   49.   D. A. Ertl, S. Stary, B. Streubel, A. Raimann, G. Haeusler, A novel homozygous mutation in
687   the parathyroid hormone gene (PTH) in a girl with isolated hypoparathyroidism. *Bone* **51**, 629-
688   632 (2012).

689   50.   A. D. White *et al.*, Ca(2+) allostery in PTH-receptor signaling. *Proc Natl Acad Sci U S A* **116**,
690   3294-3299 (2019).

691   51.   L. J. Clark *et al.*, Allosteric interactions in the parathyroid hormone GPCR-arrestin complex
692   formation. *Nat Chem Biol* **16**, 1096-1104 (2020).

693   52.   A. Kumar, M. Baumann, J. Balbach, Small Molecule Inhibited Parathyroid Hormone Mediated
694   cAMP Response by N-Terminal Peptide Binding. *Sci Rep* **6**, 22533 (2016).

695   53.   L. H. Zhao *et al.*, Structure and dynamics of the active human parathyroid hormone receptor-1.
696   *Science* **364**, 148-153 (2019).

697   54.   X. Zhai *et al.*, Molecular insights into the distinct signaling duration for the peptide-induced
698   PTH1R activation. *Nat Commun* **13**, 6276 (2022).

699   55.   R. L. Jilka, Molecular and cellular mechanisms of the anabolic effect of intermittent PTH.
700   *Bone* **40**, 1434-1446 (2007).

701   56.   A. Maeda *et al.*, Critical role of parathyroid hormone (PTH) receptor-1 phosphorylation in
702   regulating acute responses to PTH. *Proc Natl Acad Sci U S A* **110**, 5864-5869 (2013).

703   57.   M. L. Bouxsein *et al.*, Guidelines for assessment of bone microstructure in rodents using
704   micro-computed tomography. *J Bone Miner Res* **25**, 1468-1486 (2010).

705   58.   D. W. Dempster *et al.*, Standardized nomenclature, symbols, and units for bone
706   histomorphometry: a 2012 update of the report of the ASBMR Histomorphometry
707   Nomenclature Committee. *J Bone Miner Res* **28**, 2-17 (2013).

708   59.   K. E. Lim *et al.*, Core binding factor β of osteoblasts maintains cortical bone mass via
709   stabilization of Runx2 in mice. *Journal of Bone and Mineral Research* **30**, 715-722 (2015).

710   60.   X. Jin *et al.*, Silk sericin-based functional food for osteoporosis prevention and therapy.
711   *Journal of Functional Foods* **116**, 106148 (2024).

Figures

Figure 1. Formation of R25C mutant PTH(1-84) dimer.

(A) Schematic representation of PTH gene structure and expression. (B) Schematic representation of

[R56C]pre-pro-PTH(1-115) (In mature form, [R25C]PTH(1-84)) gene structure and expression (C) Schematic

representation of recombinant PTH proteins (D) *In vitro* dimerization of [R25C]PTH. Recombinant protein

constructs were transfected into HEK293T cells, and expression of PTH-3xFLAG and

[R25C]PTH3xFLAG in culture medium or cell lysate was demonstrated by western blot. The result

confirms the presence of dimeric [R25C]PTH. (*bp: base pairs; *AA: amino acids; *RA: reducing agent)

Figure 2. Effect of PTH, monomeric [R25C]PTH, and dimeric [R25C]PTH to the PTH1R *in vitro*.

(A) The binding of PTH(1-34), monomeric [R25C]PTH(1-34), and dimeric [R25C]PTH(1-34) to the PTH1R in

R^0 conformation of RG conformation was assessed by competition methods using [125]I-PTH(1-34) and

[125]I-MPTH(1-15) as radioligand (n = 4). (B) Ligand potency for cAMP signaling was assessed in SGS-

72 cells, which were derived from SaOS-2 cells modified to express Glosensor cAMP reporter (n = 4).

The cells were preloaded with luciferin and treated with varying concentrations of PTH(1-34),

monomeric [R25C]PTH(1-34), and dimeric [R25C]PTH(1-34). Error bars represent mean ± standard error.

Figure 3. Calcemic and phosphatemic responses by PTH injection in CD1 female mice.

(A) Plasma Calcemic Response after Injection (n = 6). Both PTH(1-34) and dimeric [R25C]PTH(1-34)

significantly elevate ionized calcium levels in plasma at 1 to 2 hours post-injection. After 2 hours post-

injection, plasma ionized calcium level gradually restored to baseline levels similar to those of the

vehicle group. (B) Plasma Phosphatemic Response after Injection (n = 12). Following PTH(1-34)

injection, plasma phosphate levels significantly decrease at 1-hour post-injection, subsequently

returning to baseline akin to those of the vehicle group. Conversely, dimeric [R25C]PTH(1-34) injection

shows no significant alteration in phosphatemic response but demonstrates a tendency towards a

slight decrease in phosphate levels, gradually restoring to baseline levels akin to those of the vehicle

group. (C) Urine Phosphatemic Response after Injection (n = 6). The urine phosphate levels markedly

increased at 1 hour post-injection for both PTH(1-34) and dimeric [R25C]PTH(1-34), followed by a return

to baseline levels akin to those of the vehicle group. This analysis was conducted using 9-week-old

female CD1 mice. The mice were administered PTH(1-34) and dimeric [R25C]PTH(1-34) at a

concentration of 50 nmol/kg for each compound. Error bars represent mean ± standard error. p-values

were determined using the two-way ANOVA to compare the mean of each test cell with the mean of

the control cell at the same time point. * denotes p-value < 0.05 for PTH(1-34) compared to vehicle, **

denotes p-value < 0.01 for PTH(1-34) compared to vehicle, # denotes p-value < 0.05 for dimeric

[R25C]PTH(1-34) compared to vehicle, ## denotes p-value < 0.01 for dimeric [R25C]PTH(1-34) compared to

vehicle.

Figure 4. Effect of dimeric ^{R25C}PTH(1-34) in calvarial injection model.

(A) Dissections of the calvarial bones. Calvarial injections were performed on eight-week-old male

C57BL/6 mice (N = 6 per group) that received daily administrations of vehicle, PTH(1-34), or dimeric

^{R25C}PTH(1-34) for six days. Following a 10-day treatment period, histological sections of calvariae,

stained with hematoxylin (blue-purple; indicating cell nuclei) and eosin (pink; representing bone

matrix), were obtained. The area of new bone formation, with more intense staining compared to the

existing bone tissue, is denoted by the dotted line. The bar indicates 50 μm. (B) Quantification of new

bone width in calvarial injection model. The result showed a significant increase in new bone width

following injections of both PTH(1-34) and dimeric ^{R25C}PTH(1-34) compared to the vehicle group. p-

values were obtained using the one-way ANOVA to compare the mean of each column with the mean

of a control column. ** indicates p-value <0.01 against vehicle, *** indicates p-value <0.001 against

vehicle.

Figure 5. Impact of ^{R25C}PTH(1-34) on Bone Turnover.

The effects of Sham, OVX-Control (OVX + vehicle), OVX treated with PTH(1-34) (OVX + PTH(1-34)),

and OVX treated with dimeric ^{R25C}PTH(1-34) (OVX + dimeric ^{R25C}PTH(1-34)) on bone turnover in mice.

(A) Femurs obtained from mice in each group were subjected to μCT analyses for the assessment of

bone mass. (B) Several parameters of (A) were quantified using μCT measurements, including

trabecular bone mineral density (Tb.BMD), trabecular bone volume to tissue volume (Tb.BV/TV),

trabecular bone thickness (Tb.Th), trabecular number (Tb.N), trabecular separation (Tb.Sp), cortical

bone mineral density (Ct.BMD), cortical bone volume to tissue volume (Ct.BV/TV), cortical thickness

(Ct.Th), cortical area (Ct.Ar), total tissue area (Tt.Ar), and cortical area to total tissue area (Ct.Ar/Tt.Ar).

(C) A 3D-point bending test was conducted with femurs obtained from mice in each group. The left

panel describes a schematic model of a 3D-point bending test. The middle and right panels each

indicate the maximum bending load (kgf) and slope (kgf/mm). (D) Serum levels of calcium,

phosphorus, CTX, P1NP, and ALP were measured for each group using an ELISA assay. (E) TRAP

staining of histological sections of proximal tibias was carried out to visualize osteoclast activity. The

scale bars represent 100 μm (F) Quantification of osteoclast number per bone surface (Oc.N/BS), and

osteoclast surface per bone surface (Oc.S/BS) was performed. Each group consisted of six samples

(n = 6). The error bars indicate mean ± standard error. p-values were obtained using the one-way

ANOVA to compare the mean of each column with the mean of a control column. * indicates p-value <

0.05, ** indicates p-value < 0.01, *** indicates p-value < 0.001, **** indicates p-value < 0.0001.

Figure 6. Impact of [R25C]PTH(1-34) on Osteoblast Function in Vertebrae.

The effects of Sham, OVX-Control (OVX + vehicle), OVX treated with PTH(1-34) (OVX + PTH(1-34)), and OVX treated with dimeric [R25C]PTH(1-34) (OVX + dimeric [R25C]PTH(1-34)) on osteoblast function in mice. (A) Mineralization of vertebrae obtained from each group was assessed through Von Kossa staining. (B) Quantification of trabecular bone parameters including trabecular bone volume to tissue volume (Tb.BV/TV), trabecular number (Tb.N), and trabecular separation (Tb.Sp) was performed using the Bioquant Osteo 2019 v19.9.60 program. Each bar represents 500 µm. (C) Fluorescent microscopic observations of trabecular and cortical bone sections from each group demonstrate the apposition of xylenol (red) and calcein (green) labels. (D) Quantification of trabecular bone parameters such as trabecular bone mineral apposition rate (Tb.MAR), trabecular bone formation rate to bone surface (Tb.BFR/BS), cortical bone MAR (Ct.MAR), and trabecular MAR (Tb.MAR) was

802   carried out using the Bioquant Osteo 2019 v19.9.60 program. Each group consisted of five samples

803   (n = 5). The error bars indicate mean ± standard error. p-values were obtained using the one-way

804   ANOVA compare the mean of each column with the mean of a control column. * indicates p-value <

805   0.05, ** indicates p-value < 0.01, *** indicates p-value < 0.001, **** indicates p-value < 0.0001.

A

R⁰ conformation of PTH1R

RG conformation of PTH1R

- PTH(1-34)
- Monomeric ^{R25C}PTH(1-34)
- Dimeric ^{R25C}PTH(1-34)

B

SGS-72 (SaOS2/glosensor)

- PTH(1-34)
- Monomeric ^{R25C}PTH(1-34)
- Dimeric ^{R25C}PTH(1-34)

A

Vehicle

PTH(1-34)

Dimeric ^R25C^PTH(1-34)

B

○ Vehicle
● PTH(1-34)
■ Dimeric ^R25C^PTH(1-34)

New bone width (μm)

***
**

A

Sham | OVX +Vehicle | OVX + PTH(1-34) | OVX + dimeric ^{R25C}PTH(1-34)

B

Tb.BV/TV (%) | Tb.N | Tb.Sp (mm)

- Sham
- OVX + Vehicle
- OVX + PTH(1-34)
- OVX + Dimeric ^{R25C}PTH(1-34)

C

Sham | OVX +Vehicle | OVX + PTH(1-34) | OVX + dimeric ^{R25C}PTH(1-34)

Trabecular bone

Cortical bone

D

Tb.MAR (mcm/day) | Tb.BFR/BS (mcm/day) | Ct.MAR (mcm/day) | Ct.BFR/BS (mcm/day)

- Sham
- OVX + Vehicle
- OVX + PTH(1-34)
- OVX + Dimeric ^{R25C}PTH(1-34)