Evolution of Virulence in Emerging Epidemics: From Theory to Experimental Evolution and Back

Wakinyan Benhamou[iD,1,†,⋆], François Blanquart[iD,2,†], Marc Choisy [iD,3,4], Thomas W. Berngruber[5], Rémi Choquet[iD,1,‡], and Sylvain Gandon[iD,1,‡]

[1] *CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France*
[2] *Centre for Interdisciplinary Research in Biology (CIRB), Collège de France, CNRS, INSERM, PSL Research University, Paris, France*
[3] *Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford OX3 7BN, UK*
[4] *Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam*
[5] *Reinhard-zu-Rhynernstrasse 7, 59069 Hamm, Germany*

[†] *These authors contributed equally to this work*
[‡] *These authors also contributed equally to this work*

[⋆] *E-mail:* wakinyan.benhamou@cefe.cnrs.fr

March 13, 2024

Abstract

The experimental validation of theoretical predictions is a significant step in the demonstration of the predictive power of a model. In a previous study, we monitored the evolution of the temperate phage λ spreading in continuous cultures of *Escherichia coli*. This experimental work confirmed the influence of the epidemiological dynamics on the evolution of transmission and virulence of the virus. A variant with larger propensity to lyse bacterial cells was favoured in emerging epidemics (when the density of susceptible cells was large), but counter-selected when most cells were infected. Although this approach qualitatively validated an important theoretical prediction, no attempt was made to fit the model to the data nor to further develop the model to improve the goodness of fit. Here, we show how theoretical analysis and model fitting can be used to estimate key parameters of the phage life cycle and yield new insights on the evolutionary epidemiology of the phage λ. First, we show that adding an additional stage to model the phage life cycle improves the fit of the model to the data. Second, we carry out a theoretical analysis that yields useful approximations that capture the effects of epidemiological dynamics on selection and differentiation across distinct compartments. Finally, we infer the parameters of our new model and, in particular, we estimate key phenotypic traits characterizing the two strains of the virus used in our experiment. This work illustrates how the interplay between experimental and theoretical approaches can help enhance our understanding of evolutionary epidemiology of infectious diseases.

1 Introduction

Evolutionary epidemiology theory predicts that the evolution of pathogen transmission is driven by the availability of susceptible hosts. At the onset of an epidemic, when the density of susceptible hosts is high, more transmission is favoured by natural selection. When a positive covariance exists between transmission and virulence (i.e. a virulence-transmission trade-off) (Alizon & Michalakis, 2015; Alizon et al., 2009; Anderson & May, 1982), this selection for higher transmission can indirectly select for higher virulence (Bull, 1994; Day, 2002; Day & Proulx, 2004; Frank, 1996; Gandon & Day, 2007; Lenski & May, 1994). Yet, an experimental validation of this prediction is needed to demonstrate the relevance of this result for the evolution of pathogens in emerging epidemics.

A previous study used experimental evolution of the temperate bacteriophage — or phage, for short — λ to put this prediction to the test (Berngruber et al., 2013). Phages are viruses that infect bacteria and phage λ is the archetypal temperate phage, which can switch between a lytic and a lysogenic life style. Upon infection, the virus may commit to the lytic pathway by hijacking the host's replication machinery to produce new virions (viral particles) and eventually release them in the environment after the lysis of the host cell. Alternatively, the virus may commit to the lysogenic pathway by integrating its genome into the bacterial chromosome where it will lay in a dormant state as a prophage and be hereditarily transmitted to daughter cells at the pace of lysogen divisions. The prophage may also regain virulence by excising itself from the genome, switching to a lytic cycle (reactivation, also called induction) and, thus, shifting from vertical to horizontal transmission. (Echols, 1972; Gandon, 2016; Lwoff, 1953; Ptashne, 1992) The evolution experiment designed in (Berngruber et al., 2013) monitored the competition between two strains of phage λ with distinct life-history strategies in continuous cultures of *E. coli*. The first strain is the wildtype, which is known to have a relatively large lysogenization rate and low reactivation rate. The second strain is the λcI857 variant, which carries a point mutation in the transcriptional repressor protein cI (St-Pierre & Endy, 2008; Sussman & Jacob, 1962); the cI mutant is known to be more virulent and transmitted mostly horizontally through lytic cycles. (Berngruber et al., 2013) developed a mathematical model tailored to the life cycle of phage λ. Numerical simulations of this model using parameter values from previous experimental studies led to three theoretical predictions: (i) the virulent strain outcompetes the wildtype when susceptible hosts are abundant, but the direction of selection is reversed as soon as the epidemic reaches high prevalence, (ii) the lower the initial prevalence, the higher the increase

in virulence during the epidemic and (iii) the virulent strain is always more frequent among viral particles than among prophages. Tracking both the epidemiology (prevalence) and the evolution of the virus (frequency of the virulent strain among viral particles and among infected bacteria), all three predictions were confirmed experimentally (Berngruber et al., 2013). Yet, the data was only used as a qualitative validation of the theory and no attempt was made to explore the quantitative match between the predicted and the observed dynamics of the virus.

In the present work, we show how the quantitative analysis of the experimental results from (Berngruber et al., 2013) improves our understanding of the evolutionary epidemiology of phage λ. First, we modified the structure of the epidemiological model to better capture the evolutionary dynamics of the virus among infected bacteria. Second, we carry out a theoretical analysis of this model to provide useful approximations to predict the evolutionary dynamics of the virus at different stages of the epidemic. Finally, we develop a statistical inference approach to obtain quantitative estimates of the parameters of the model and, especially, the life-history traits of the different strains of phage λ.

2 Materials and methods

Following (Berngruber et al., 2013), we first model the competition between the wildtype strain – hereafter denoted by the subscript w – and the mutant strain λcI857 – hereafter denoted by the subscript m – of phage λ in a chemostat with a well-mixed continuous culture of its bacterial host *E. coli* (**Fig. 1**). We summarize the notations in **Table 1**. We then recall the experimental data generated in the original study and we describe how we generate simulated data to validate our ability to estimate model parameters. In the last section, we detail the statistical inference approach we developed.

2.1 A model coupling epidemiology and evolution of phage virulence

2.1.1 Epidemiology

Bacteria are either susceptible (S) or infected with phage λ (we do not consider resistant bacteria); infected bacteria may be lysogenic (L), following phage integration (lysogenization), or carrying phages replicating in their cytoplasm prior to lysis (Y), following either lytic infection or prophage reactiva-

tion. Unlike the original model in (Berngruber et al., 2013), adding an extra stage Y allows us to take into account the lysis time and we show below that this new model improves the goodness of fit to the data (see **Results §3.2.2**). The free virus stage (V) corresponds to the free viral particles (virions) that are in the culture medium – "free" meaning "extracellular" here. Throughout, for each compartment, for instance S, we denote $S(t)$ its density at the current time t and $\dot{S}(t)$ its differentiation with respect to time. Hence, $L(t) = L_w(t) + L_m(t)$ is the total density at time t of L cells, $Y(t) = Y_w(t) + Y_m(t)$, of Y cells, $V(t) = V_w(t) + V_m(t)$, of free viral particles, and $N(t) = S(t) + L(t) + Y(t)$, of bacteria.

Susceptible and lysogenic bacteria grow at a logistic rate $r(1 - N(t)/K)$, where r is the intrinsic growth rate and K the carrying capacity. Bacteria and virions are removed from the chemostat at a dilution rate δ. Free viral particles adsorb onto bacterial cells at a rate a; adsorption is non-reversible, that is the fate of adsorbed viruses is only to infect or die out. Infection of a susceptible host requires in addition the injection of the phage's genetic material into the bacterial cytoplasm (fusion, with probability b). The force of infection – i.e., the *per capita* infection rate – is therefore given by $abV(t)$. We assume that superinfection (including co-infection with both strains) does not occur. Notably, prophage establishment of phage λ is known to provide cellular immunity, or superinfection inhibition (Berngruber et al., 2010; Gandon, 2016; Lwoff, 1953; Ptashne, 1992). Upon infection, the wildtype and virulent strains of phage λ may either be integrated as prophages into the host genome (lysogenic cycle) with probabilities ϕ_w and ϕ_m, respectively, such that $\Delta\phi = \phi_m - \phi_w < 0$, or start the biosynthesis and assembly of viral copies (lytic cycle) with complementary probabilities. Once integrated, reactivations of wildtype and virulent prophages occur at a rate α_w and α_m, respectively, such that $\Delta\alpha = \alpha_m - \alpha_w > 0$. Following either lytic infections or prophage reactivations, host cells are lysed at a rate τ and eventually release B viral particles upon lysis (burst size). We assume that parameters a, b, τ and B are the same for the wildtype and the mutant which yields the following system of ordinary differential equations (ODEs):

$$
\begin{cases}
\dot{S}(t) &= \underbrace{rS(t)\left(1 - \dfrac{N(t)}{K}\right)}_{\text{Growth}} - \underbrace{abV(t)S(t)}_{\text{Infection}} - \underbrace{\delta S(t)}_{\text{Removal}} \\[2em]
\dot{L}(t) &= \underbrace{rL(t)\left(1 - \dfrac{N(t)}{K}\right)}_{\substack{\text{Growth \&} \\ \text{vertical transmission}}} + \underbrace{\overline{\phi}(t)abV(t)S(t)}_{\text{Lysogenization}} - \underbrace{\overline{\alpha}(t)L(t)}_{\substack{\text{Prophage} \\ \text{reactivation}}} - \underbrace{\delta L(t)}_{\text{Removal}} \\[2em]
\dot{Y}(t) &= \underbrace{\left(1 - \overline{\phi}(t)\right)abV(t)S(t)}_{\text{Lytic infection}} + \underbrace{\overline{\alpha}(t)L(t)}_{\substack{\text{Prophage} \\ \text{reactivation}}} - \underbrace{\tau Y(t)}_{\text{Lysis}} - \underbrace{\delta Y(t)}_{\text{Removal}} \\[2em]
\dot{V}(t) &= \underbrace{\tau Y(t)B}_{\text{Virion release}} - \underbrace{aN(t)V(t)}_{\text{Adsorption}} - \underbrace{\delta V(t)}_{\text{Removal}}
\end{cases}
\tag{1}
$$

where $\overline{\phi}(t)$ is the mean probability of lysogenisation among free phage particles and $\overline{\alpha}(t)$ is the mean rate of reactivation among prophages:

$$
\begin{cases}
\overline{\phi}(t) &= q(t)\phi_m + \left(1 - q(t)\right)\phi_w \\[1em]
\overline{\alpha}(t) &= p(t)\alpha_m + \left(1 - p(t)\right)\alpha_w
\end{cases}
\tag{2}
$$

with $q(t) = V_m(t)/V(t)$ and $p(t) = L_m(t)/L(t)$, the frequencies of the virulent strain at time t among compartments V and L, respectively.

2.1.2   Evolution

Along with these epidemiological dynamics, we also track the evolutionary dynamics of phage λ. We recall that $p(t)$ and $q(t)$ refer to frequencies of the virulent strain in compartment L and V, respectively. In addition, we also denote $f(t) = Y_m(t)/Y(t)$, the frequency of the virulent strain at time t among Y cells, and $g(t)$, the frequency of the virulent strain in infected cells (either Y or L) such that:

$$
g(t) = \frac{Y_m(t) + L_m(t)}{Y(t) + L(t)} = f(t)\left(\frac{Y(t)}{Y(t) + L(t)}\right) + p(t)\left(\frac{L(t)}{Y(t) + L(t)}\right).
\tag{3}
$$

Using (1) and (2), one may calculate the ODEs of each of these frequencies (see **SI Appendix §S1.2**). More conveniently, we will then focus on logit-frequencies instead, that is the log odds ln(frequency of the mutant / frequency of the wildtype). Taken together, the equations of the temporal

dynamics of the frequencies and the model (1) yield the coupled evolutionary-epidemiological dynamics of this phage-bacteria system. The analysis of this model can provide key insight on the evolutionary forces acting on the virus; it may also provide useful approximations for the change in mutant frequency at different stages of the epidemic.

2.2 Time series datasets

2.2.1 Experimental data

We use experimental time series obtained in the first evolution experiment of (Berngruber et al., 2013). Briefly, this experiment started with eight independent chemostats (5 mL chamber volume, maintained at 35°C) of well-mixed bacterial cultures of *E. coli* MG1655 (RecA+) at carrying capacity. Bacteria were initially infected by both the wildtype and virulent strain λcI857 of phage λ (prophage stage with initial ratio 1:1). Two treatments were considered using 4 chemostats each: (i) an epidemic treatment – low initial prevalence, around 1% – and (ii) an endemic treatment – high initial prevalence, around 99%. Throughout the course of the experiment, several quantities were monitored: the prevalence, the frequency infected by each strain and the strain frequency in the culture medium (free virus stage). Samplings in each chemostat were performed hourly, from 1 to 60 h maximum. The prevalence and the frequency of infected hosts were tracked using flow cytometry (FACS) with fluorescent protein marker colors (CFP and YFP) while the strain frequency in the free virus stage was tracked by qPCR.

2.2.2 Simulated data

Alongside experimental data, we also carry out an analysis based on simulated data in order to validate our ability to infer parameters from experimental data. For this purpose, we start with: $\alpha_w = 7 \times 10^{-3}$, $\alpha_m = 2 \times 10^{-2}$, $\phi_w = 0.2$, $\phi_m = 2 \times 10^{-2}$, $a = 3 \times 10^{-9}$, $b = 0.1$, $B = 80$, $r = 1.4$, $\tau = 1.5$, $K = 10^9$ and $\delta = 0.8$. At $t = 0$, bacteria are at carrying capacity K with initial prevalence 1% (epidemic treatment) or 99% (endemic treatment) and the initial prophage ratio for the two strains is 1:1. We simulate the deterministic model (1) for the epidemic and endemic treatment (**Fig. S1**). I.i.d. Gaussian noise is then added at each time point to mimic measurement errors on the logit-prevalence $\mathrm{logit}\big(P(t)\big)$, the logit-frequency of hosts infected by the virulent phage $\mathrm{logit}\big(g(t)\big)$ and the logit-frequency of the virulent phage in the free virus stage $\mathrm{logit}\big(q(t)\big)$ – this is independently repeated four times for each simulation to obtain four replicates (chemostats) per treatment. We modulate data quantity through

two sampling frequencies: $0.1\ h^{-1}$ vs. $1\ h^{-1}$, and we modulate data quality through two standard deviations (SD) of measurement errors: 0.01 vs. 0.5. We thus end up with four combinations of data quantity and quality (see example in **Fig. S5**) from which we then try to recover parameter values.

2.3 Maximum likelihood estimation

For the estimation process, we used a two-step approach. Using theoretical results, we first compute point estimates of the rates of prophage reactivation α_w and α_m. Then, we fix the latter to estimate the remaining parameters of the model using non-linear optimizations. Sieve bootstrap were used to compute the joint distributions of all these estimated parameters.

2.3.1 Estimation of the rates of prophage reactivation

From the analysis of the model (see **Results §3.1**), we show that when the system reaches high prevalence the selection gradient $\mathscr{S}$ of the mutant – i.e., the rate at which it grows or declines in frequency on the logit scale – is simply given by: $\mathscr{S} = \alpha_w - \alpha_m = -\Delta\alpha$ (**Results §3.1.2**). Furthermore, the differentiation $\mathcal{Q}^{VL}$ of the virulent strain between free phages and prophages converges towards approximately: $\mathcal{Q}^{VL} = \alpha_m/\alpha_w = 1 + \Delta\alpha/\alpha_w$ (**Results §3.1.3**). Combining these two expressions enables us to estimate separately both reactivation rates:

$$
\begin{cases}
\alpha_w & = \dfrac{\mathscr{S}}{1 - \mathcal{Q}^{VL}} \\
\alpha_m & = \dfrac{\mathscr{S} \times \mathcal{Q}^{VL}}{1 - \mathcal{Q}^{VL}}
\end{cases}
$$

For each chemostat, we therefore only keep the data from the time point the prevalence has reached 95%. We fit a linear model on the logit-frequency infected by the virulent phage logit $(g(t))$ to estimate the slope $\mathscr{S}$, and we estimate $\mathcal{Q}^{VL}$ by calculating the geometric mean of $\frac{q(t)(1-g(t))}{(1-q(t))g(t)}$ (see details in **SI Appendix §S3.1**). Note that we substitute $p(t)$ by $g(t)$ for the calculation of the differentiation because we only have access to $g(t)$ in the experiment, and $g(t)$ is almost identical to $p(t)$ towards the end of the epidemic (see **Results §3.1.2** and **Fig. S1**).

2.3.2 Estimation of the remaining parameters

We have three response variables: (i) $\text{logit}(P(t))$, the logit-prevalence, (ii) $\text{logit}(g(t))$, the logit-frequency of hosts infected by the virulent phage and (iii) $\text{logit}(q(t))$, the logit-frequency of the virulent phage in the free virus stage. Let $P_0^{epidemic}$ and $P_0^{endemic}$ be the initial conditions (at $t = 0$) of the prevalence in the epidemic and endemic treatments, respectively, and p_0, the initial condition of the frequency infected by the virulent phage (identical for both treatments). Let $\theta = \left(\phi_w, \phi_m, a, b, r, \tau, P_0^{epidemic}, P_0^{endemic}, p_0\right)$ be the vector of model parameters to estimate. Several parameter values are fixed: $K = 10^9$ $cells$ and $\delta = 0.8$ h^{-1} (Berngruber et al., 2013); α_w and α_m are fixed to their previous point estimates (cf. **Methods §2.3.1**); and $B = 80$ $virus\,.\,cell^{-1}$ (Wang, 2006) as we show in **Results §3.2.1** that the burst size B is not separately identifiable from the parameter b. We estimate as well σ_P, σ_g and σ_q, the SD of the measurement errors for each of the three response variables, respectively. We assume that variation around the deterministic dynamics stems solely from measurement errors and we thereby neglect any additional process stochasticity. This is justified in particular by the very controlled conditions of the experiment and by the large population sizes of both the bacteria and the phage in the chemostats. For each response variable, measurement errors are assumed to be normally distributed and i.i.d. across all treatments, replicates and time points. The corresponding log-likelihoods $(\ln(\mathcal{L}))$ are thence respectively given by:

$$\ln(\mathcal{L}_P(\theta, \sigma_P)) = \sum_{i,j,t} \ln\left[\varphi\left(\text{logit}\left(P_{i,j}(t)\right)^{data} \mid \text{logit}\left(P_{i,j}(\theta, t)\right)^{sim}, \sigma_P^2\right)\right]$$

$$\ln(\mathcal{L}_g(\theta, \sigma_g)) = \sum_{i,j,t} \ln\left[\varphi\left(\text{logit}\left(g_{i,j}(t)\right)^{data} \mid \text{logit}\left(g_{i,j}(\theta, t)\right)^{sim}, \sigma_g^2\right)\right]$$

$$\ln(\mathcal{L}_q(\theta, \sigma_q)) = \sum_{i,j,t} \ln\left[\varphi\left(\text{logit}\left(q_{i,j}(t)\right)^{data} \mid \text{logit}\left(q_{i,j}(\theta, t)\right)^{sim}, \sigma_q^2\right)\right]$$

where i refers to treatments (epidemic vs. endemic), j to the jth replicate (chemostat), t to time points and $\varphi(.\mid \mu, \sigma^2)$ to the PDF of the Normal distribution with mean μ and variance σ^2; sim indicates model outputs. We then denote $\widehat{\theta}$, the MLE estimator of θ such that:

$$\widehat{\theta} = \underset{\theta}{\text{argmax}}\left(\ln(\mathcal{L}_P(\theta, \sigma_P)) + \ln(\mathcal{L}_g(\theta, \sigma_g)) + \ln(\mathcal{L}_q(\theta, \sigma_q))\right).$$

In practice, we minimize the negative overall log-likelihood using the Nelder-Mead (aka downhill simplex) algorithm. Parameter bounds (reported in **Table S2**) are imposed through parameter transformations. Due to the presence of local minima, optimizations are repeated for 2,000 sets of uniformly drawn starting points to ensure convergence to a global minimum. The best MLE set of estimates $\hat{\theta}$ corresponds to the fit associated with the lowest negative overall log-likelihood with successful completion.

2.3.3 Confidence intervals

We generate bootstrapped data to compute 95% CIs of our parameters using sieve bootstrap (Bühlmann, 1997) on the residuals between experimental data and the best fit of our model. For this purpose, autoregressive moving-average (ARMA) models are fit to the time series of centred residuals of each chemostat independently. We then use ARMA models to simulate new residuals from which we reconstruct new datasets. We eventually reiterate the above estimation procedure (cf. **§2.3.1-2.3.2**), but starting non-linear optimizations only from the best MLE estimates we obtained with the original data. By repeating this for 10,000 bootstrapped datasets, we compute the joint distributions of estimated parameters.

2.4 Details of the implementation

Numerical simulations and data analyses were carried out using R (R Core Team, 2022) version 4.2.0 (2022-04-22). ODEs were solved numerically by the function '*ode*' – with method '*lsoda*' – from the package '*deSolve*' (Soetaert et al., 2010). Non-linear optimizations for MLE were tackled with the function '*nmk*' (Kelley, 1999), from the package '*dfoptim*'. Fit and selection of ARMA models for sieve bootstrap were carried out using the function '*auto.arima*', from the package '*forecast*' (Hyndman & Khandakar, 2008).

3 Results

3.1 Theoretical analysis

When $r > \delta$, the virus-free system converges to an equilibrium where $S(\infty) = K(1 - \delta/r)$. When a single strain $k \in \{w, m\}$ of the virus (with phenotypes α_k and ϕ_k) is introduced in the bacterial

population (fully susceptible, with density S_0), the fate of the phage-bacteria system depends on the basic reproduction number of the pathogen $\mathscr{R}_{0,k}$ which is given by:

$$\mathscr{R}_{0,k} = \frac{A + \sqrt{A^2 - 4r\left(1 - \frac{S_0}{K}\right)(1 - \phi_k)abS_0\tau B(aS_0 + \delta)(\alpha_k + \delta)(\tau + \delta)}}{2(aS_0 + \delta)(\alpha_k + \delta)(\tau + \delta)}, \tag{4}$$

with $A = r\left(1 - \frac{S_0}{K}\right)(aS_0 + \delta)(\tau + \delta) + abS_0\tau B(\alpha_k + (1 - \phi_k)\delta)$ (see **SI Appendix §S2.1** for the construction of the next-generation matrix, following (Diekmann et al., 2010)). When $\mathscr{R}_{0,k} < 1$, the virus goes extinct and the bacterial population converges to the previous virus-free equilibrium. Alternatively, when $\mathscr{R}_{0,k} > 1$, an epidemic breaks out and eventually stabilises to an endemic equilibrium where all the cells are infected by the virus (**SI Appendix §S2.1**).

In the following, we analyse the evolutionary dynamics (i) at the beginning of an epidemic where $S(t) = S_0$ and (ii) when the system stabilizes towards the endemic equilibrium where $S(t) = 0$.

3.1.1 Evolution in an emerging epidemic

At the onset of the epidemic, susceptible cells are highly abundant. For the sake of simplicity, we analyse the dynamics of the epidemic when the host density is assumed to be constant over time ($S(t) = S_0$). As in the experimental design, we start with a bacterial population at carrying capacity ($S_0 \approx K$). Density dependence reduces cell reproduction and vertical transmission of the virus. Consequently, the epidemic is mainly driven by the lytic pathway. At $t = 0$ though, only lysogens are introduced at very low density. After a very short time, the phage-bacteria system reaches its new dynamical regime, following prophage reactivations, lyses and releases of virions. From there, we assume that $L(t)/Y(t) \approx 0$ and we focus on the dynamics of the mutant in the compartment Y. We derive an approximation of the selection gradient $\mathscr{S}$ of the virulent phage, which corresponds to the rate at which it increases in frequency on the logit scale, and we show in **SI Appendix §2.2** that:

$$\mathscr{S} \propto -\Delta\phi, \tag{5}$$

meaning that it is selected for ($\Delta\phi < 0$, cf. **Fig. S3-A**). This is consistent with the experimental data where the virulent phage transiently outcompetes the wildtype at the early stage of the epidemic treatment (Berngruber et al., 2013). Our prediction is relatively accurate when the density of susceptible hosts remains effectively constant over time; otherwise, however, our prediction deviates from

the simulation whose rate slows down because the density of susceptible cells is rapidly decreased by the spread of the epidemic (**Fig. 2-A** and **Fig. S3-C**).

3.1.2 Evolution at the end of the epidemic

At the end of the epidemic we expect that all the cells will be infected by a prophage and there will be no susceptible cells anymore. Consequently, no horizontal transmission takes place and, in contrast with the previous scenario, we can neglect the density of Y cells relative to the density of lysogens. Indeed, since $1/\alpha_w \gg 1/\tau$ (time elapsed between phage integration and reactivation is much longer than lysis time), then $Y(t)/L(t) \approx 0$ (see details in **SI Appendix §S2.3**). Note that this also means that the frequency of the virulent strain is mainly driven by its frequency in L and $g(t) \approx p(t)$. We show in **SI Appendix §S2.3** that the selection gradient of the virulent phage is given by:

$$\mathscr{S} = -\Delta\alpha. \tag{6}$$

As a result, the virulent phage is counter-selected in the long-term ($\Delta\alpha > 0$) and, in each compartment, linearly declines in frequency at a rate $\mathscr{S}$ (negative slope) on the logit scale (**Fig. 2-B** and **Fig. S1-B**).

3.1.3 Differentiation across compartments

The two previous subsections deal with the dynamics of evolution over time; let us now look at the dynamics of selection between different compartments. We define here the differentiation $\mathcal{Q}^{VL}$ between the free virus stage and lysogenic cells as:

$$\mathcal{Q}^{VL}(t) = \frac{q(t)}{1 - q(t)} \frac{1 - p(t)}{p(t)}, \tag{7}$$

such that: $\ln\left(\mathcal{Q}^{VL}(t)\right) = \text{logit}(q(t)) - \text{logit}(p(t))$.

We show in **SI Appendix §S2.4** that, at the end of the epidemic, the differentiation $\mathcal{Q}^{VL}(t)$ converges approximately towards:

$$\mathcal{Q}^{VL} \approx \frac{\alpha_m}{\alpha_w} = 1 + \frac{\Delta\alpha}{\alpha_w} \tag{8}$$

(**Fig. 2-C**), a value of 1 corresponding to no differentiation. We obtain the same expression for the

convergence of the differentiation between Y and L cells and around 1 between the free virus stage V and Y cells (cf. **SI Appendix §2.4**). Interestingly, these results are also valid at $t = 0^+$ (**Fig. S4**) as there is no free phages yet and thereby no horizontal transmission yet which, within a very short space of time, resembles the end of the epidemic. When the system stabilizes at the end of the epidemic, the virulent strain is therefore more frequent among free viruses and Y cells than among L cells (prophages) but we also expect almost no differentiation between free viruses and Y cells (**Fig. S4**).

3.2 Statistical inference

3.2.1 Inference from simulated data

We first conduct a simulation study. We start by validating our estimation method of the rates of prophage reactivation α_w and α_m. Combining equations (6) and (8), we compute point estimates of both parameters (cf. **Methods §2.3.1**) for a large number of simulated datasets. Overall, estimated values show a very good match with those used in the original simulations (**Fig. S5**), especially when the SD of measurement errors is small.

We then show that parameters b and B may not be separately identifiable (**Fig. S2-A**, with sampling effort $= 10\ h^{-1}$ and SD of measurement error $= 0.01$). We fix K and δ to their known values, as well as α_w and α_m, as though the rates of prophage reactivation have been correctly estimated beforehand. Setting $(b,\ B)$ to many combinations of values $((b,\ B) \in [0,\ 0.2] \times [0,\ 100])$ and maximizing over the remaining parameters, the resulting landscape of the overall log-likelihood suggests that only the product $b \times B$ is identifiable (**Fig. S6**). Such compensatory effect may be understood as the impossibility here to discriminate whether phage λ infect fewer bacteria (lower probability b of fusion) but release more viral copies upon lysis (larger burst size B) or vice-versa. This is the reason why we fix B in the following and only estimate b.

Fixing B to its true value used in the original simulations, point estimates of parameters are quite close to their true values (**Fig. S7**). However, some parameters – like r, τ or b – seem more difficult to estimate with a lower sampling effort.

3.2.2 Inference from experimental data

The original model used in (Berngruber et al., 2013) failed to properly capture the evolutionary dynamics among infected hosts at the early state of the epidemic. Such discrepancy was an opportunity to go back from experimental data to theory and to challenge the structure of the model. We noticed that adding an extra stage Y in the infected compartment allowed us to better fit the experimental data (AIC=2901) than the original model (AIC=3002). This significant improvement strongly supported the new version of the model developed in this study and for which we present the estimation results below.

First, we estimate: $\Delta\alpha = 9.31 \times 10^{-3}\ h^{-1}$ and $\mathcal{Q}^{VL} = 4.60$ (see **Fig. 3** for the latter). Point estimates of the rates of prophage reactivation (expressed in h^{-1}) are thus: $\hat{\alpha}_w = 2.58 \times 10^{-3}$ and $\hat{\alpha}_m = 1.19 \times 10^{-2}$.

Second, we perform non-linear optimizations to estimate the remaining parameters. Overall, computed dynamics of the logit-prevalence and logit-frequencies of the virulent phage closely fit experimental data (**Fig. 4**). Best MLE estimates – with an overall log-likelihood equals to -1436.44 – are listed in **Table 2**, along with their 95% bootstrap-based CIs (see **Fig. S8** for the density distributions of estimated parameters). We show pairwise correlation coefficients in **Fig. S9**. Some parameters are correlated positively, in particular: α_w with α_m, p_0 with both reactivation rates, ϕ_m with $P_0^{epidemic}$ and r with b; or negatively, in particular: τ with b and r. In **Fig. S10**, we plot the 95% distributions of fitted values. To test the impact of the fixed value used for the burst size B, we studied how perturbations of the original value (80 $virus\,.\,cell^{-1}$) affected the estimation of the other parameter values (new point estimates from non-linear optimizations). In **Fig. S11**, we show that all parameters are extremely robust to these perturbations, except for parameter b as expected from **Fig. S6** (but the product $b \times B$ is always around 3.95). It is also worth noting that the basic reproduction number is not affected by the choice of the value of B, as it always appears as a product with b in (4). We estimate the basic reproduction number of the wildtype strain $\mathcal{R}_{0,w}$ to be 1.48 (95% CI [1.04, 2.15]) and of the virulent strain $\mathcal{R}_{0,m}$ to be 2.20 (95% CI [1.58, 3.05]). Interestingly, the basic reproduction number of the virulent mutant $\mathcal{R}_{0,m}$ is higher than that of the wildtype $\mathcal{R}_{0,w}$, but the virulent mutant always loses the competition in the long term. Indeed, the basic reproduction number gives the expected number of secondary infections in a fully susceptible host population. At the early stage of the epidemic, the basic reproduction number provides a good predictor of the fitness

of the two strains: the mutant outcompetes the wildtype because $\mathscr{R}_{0,m} > \mathscr{R}_{0,w}$. When the density of susceptible host cells drops, however, the basic reproduction number becomes a poor predictor of fitness.

We compared our estimates of model parameters with previous studies (see reported values in **Table S3**), focusing in particular on the parameters of the phage. Note that previous estimations of these life-history traits are obtained from experimental assays using monomorphic phage cultures. Our estimate of the rate of prophage reactivation of the wildtype strain α_w falls in the range of previous *in vitro* estimates, which is very large (between 10^{-7} and $10^{-2}\,h^{-1}$ (De Paepe et al., 2016; Little et al., 1999; Zong et al., 2010)), which is probably due to the variability in experimental methods. For the cI variant, the orders of magnitude of α_m are more consistent (De Paepe et al., 2016; Zong et al., 2010). Other studies, using single-cell monitoring of infected cells to estimate the probability of lysogenization of the wildtype strain ϕ_w, obtained very similar results when the multiplicity of infection (MOI) is low, around 0.35 on average (Zeng et al., 2010) – but see the **Discussion** for the effect of MOI. We estimate a 12-fold decrease for the probability of lysogenization of the virulent strain ϕ_m and, to our knowledge, this parameter has not been estimated elsewhere. The lysis time $1/\tau$, estimated close to $1\,h$ (though with larger uncertainty), is also consistent with the biology of phage λ (De Paepe et al., 2016; Lindberg et al., 2014; Shao & Wang, 2008; Zong et al., 2010). On the other hand, the adsorption rate a is poorly estimated. Although this issue did not arise with simulated data, estimates based on experimental data are mostly at their upper bound (10^{-6}), several orders of magnitude higher than expected (De Paepe et al., 2016; Lindberg et al., 2014; Shao & Wang, 2008). To investigate the sensitivity of the other estimates to this, we reiterate non-linear optimizations fixing a to different values ranging from 10^{-9} to 1 to compute new point estimates. Most parameters show high robustness and no relative variation higher than 50% is found for values of a between 10^{-8} to 1 (**Fig. S12-A**). We carry out the same sensitivity analysis for the intrinsic growth rate of the host r which was also poorly estimated (its distribution being mainly flat): with values of r ranging between 1 and 5, the probabilities of lysogenization are little affected by the perturbations in r (relative variation contained between -7% and $+15\%$) while parameters b and τ – the most correlated with r – are the most sensitive (**Fig. S12-B**).

4 Discussion

In a previous study, (Berngruber et al., 2013) carried out an evolution experiment with the phage λ to validate several theoretical predictions on the evolution of virulence and transmission. This study used a two-step approach: (i) a mathematical model tailored to the biology of phage λ was used to show how epidemiological dynamics feeds back on the evolution of the pathogen; (ii) tracking the variation of the frequency of a viral mutant across time and across compartments (infected host and free virus) confirmed the impact of the density of susceptible cells on the transient evolution of the virus at different stages of its life cycle. Yet, this experimental validation was limited to a qualitative match with the theoretical predictions of numerical simulations of the model. In the present study, we adopt a reverse approach where we start from the data and we try to improve the theoretical model developed to describe the evolutionary epidemiology of phage λ.

First, we noticed that the previous model failed to properly capture the transient evolutionary dynamics in the infected compartment. We improved the goodness of fit with the experimental data by distinguishing two types of infected cells: lysogenic cells (L) and cells committed to the lytic pathway (Y). Indeed, adding extra stages allowed us to observe a transitory peak in the frequency of hosts infected by the virulent phage (**Fig. S13-A**, middle panel), similar to what we observed in the experimental data. In this model, the peak is due to the frequency of the virulent phage among Y bacteria that, unlike L bacteria, transiently overshoots during the acute phase of the epidemic. Crucially, lysis was assumed to be instantaneous in the original model (Berngruber et al., 2013) and adding this additional stage in the phage life cycle allowed us to take into account the lag between infection and lysis. For the sake of simplicity and parsimony, we only kept a single extra stage Y, yielding exponentially distributed lysis time.

Second, we performed a theoretical analysis of this new model to go beyond the numerical approach used in (Berngruber et al., 2013). In particular, we obtained analytic approximations for the spread of the virulent phage at the beginning and at the end of the epidemic: (i) the virulent mutant increases in frequency at the onset of the epidemic (when the density of susceptible hosts is high) with a speed approximately proportional to $-\Delta\phi$; (ii) the virulent mutant decreases in frequency at the end of the epidemic (when the density of susceptible hosts is low) with a negative selection gradient $-\Delta\alpha$, which is consistent with the prediction that in the long-term, and in the absence of an influx of susceptible hosts, the temperate phage should evolve a fully lysogen strategy where $\alpha = 0$ and $\phi = 1$ (Bruce et al.,

2021; Wahl et al., 2019); (iii) the virulent phage is always more frequent in the the free virus stage than among prophages and this differentiation is approximated by $1 + \Delta\alpha/\alpha_w$ at the end of the epidemic. The first approximation only held for a very limited period of time because the density of susceptible cells drops very rapidly and this epidemiological feedback affects the selection on transmission and virulence. In contrast, the final two predictions were valid as soon as the epidemic reaches high prevalence and, interestingly, yielded a novel way to estimate the rates of prophage reactivations of both strains.

Third, we developed an inference approach to estimate the parameters of the model. We implemented a two-step MLE procedure: (i) we first obtained point estimates of the reactivation rates α_w and α_m of both viral strains, (ii) then we fixed α_w and α_m to their point estimates and we ran non-linear optimizations to infer the remaining parameters of the model (except the burst size B which had to be fixed because of an identifiability issue). Sieve bootstraps were used to generate joint distributions for all the estimated parameters (**Table 2**). We showed that our estimates of the key phenotypic traits α_w, α_m and ϕ_w were consistent with existing literature (**Table S3**).

Crucially, our inference approach is very different from previous studies. First, we analyse the dynamics of a polymorphic viral population. Second, we use three different types of data to infer the parameter values of the model: (i) epidemiological data (i.e. the prevalence of the infection), (ii) temporal variation of variant frequencies and (iii) differentiation of the variant frequency across compartments. Each data carries complementary information and allows us to jointly estimate the life-history traits of both strains of the phage. This novel method is particularly well suited to estimate their rates of prophage reactivation, for which only the endemic treatment is needed – see **SI Appendix §S3.2** where we propose a Bayesian counterpart to easily obtain posterior distributions and credible intervals for these two parameters (applied to our experimental data, we show results in **Fig. S14-A**). The estimation of the probabilities of lysogenization is however more difficult as it requires to monitor the transient dynamics of the epidemic treatment. The rapid epidemiological feedback that occurred in emerging epidemics makes it necessary to consider both the epidemiology and the evolution of the infection. It might thus be relevant to carry out shorter experiments for the epidemic treatment – i.e., focusing on the the early state of the epidemic – but with a more frequent sampling effort to monitor more precisely the change in frequency. Alternatively, the use of two-stage chemostats, where the influx of susceptible hosts could be maintained experimentally (Bonachela &

Levin, 2014; Husimi et al., 1982), might also be an interesting option worth investigating.

Even though we have improved the original model, many features of the present model could be challenged. For example, we assume all phenotypic traits to be constant across time and across chemostats. Yet, key life-history traits of phage λ are known to vary with the environment (Wegrzyn & Wegrzyn, 2005) (i.e., phenotypic plasticity). In particular, the probability of lysogenization and the reactivation rate of the λ mutant used in the evolution experiment is temperature-sensitive (Berngruber et al., 2013; St-Pierre & Endy, 2008; Zong et al., 2010). Interestingly, the estimated rates of reactivation varied among chemostats (**Fig. S14-B**, using a Bayesian approach). This variation may have been driven by small variations in temperature among the chemostats that could have affected the reactivation rate of the mutant (higher rates of reactivation would be expected with higher temperature). The probability of lysogenization of phage λ is also known to be more likely to occur at high multiplicity of infection (MOI) (Kourilsky & Knapp, 1975; Kourilsky, 1973; Zeng et al., 2010). MOI-dependant phenotypic plasticity may also affect the rate of reactivation of temperate phages (Bruce et al., 2021), or the lysis time of virulent phages (Rutberg & Rutberg, 1965). Yet, our model has the advantage of the parsimony, while already convincingly reproducing the qualitative patterns observed in the data.

To conclude, our study emphasizes the benefits of combining theoretical, statistical and experimental approaches. Combining effectively these different perspectives requires an iterative process. Each step is an opportunity to challenge and improve our understanding of the biological model, to elucidate microbial life cycles and to provide support and guidance during experimentation. While experimental life-history assays focus on only one strain at a time, we demonstrate the relevance of evolution experiments where different strains are put in competition and where tracking their relative frequencies over time may yield novel and efficient ways to estimate some dimensions of their phenotypes, especially when it is difficult to access absolute densities. This approach could be used under different conditions and in particular *in vivo* (De Paepe et al., 2016). The present study may thus help characterize the phenotypic traits of microbial organisms used in experimental evolution.

Abbreviations (alphabetical order)

AIC:	Akaike Information Criterion
ARMA:	AutoRegressive-Moving-Average
CFP:	Cyan Fluorescent Protein
CI:	Confidence Interval
FACS:	Fluorescence-Activated Cell Sorting
i.i.d.:	independent and identically distributed
MLE:	Maximum Likelihood Estimation
MOI:	Multiplicity Of Infection
ODE:	Ordinary Differential Equation
PDF:	Probability Density Function
qPCR:	quantitative Polymerase Chain Reaction
SD:	standard deviation
YFP:	Yellow Fluorescent Protein

Supplementary material

Supplementary Figures and Tables (PDF)

Supplementary Information (SI Appendix) (PDF)

Data accessibility

Data that were used in this study, along with the scripts for the analyses (in R), are available in this GitHub repository.

Funding

We acknowledge the French Ministry of Higher Education and Research and the École Normale Supérieure Paris-Saclay for the PhD scholarship of WB.

Conflict of interest

The authors declare no conflict of interest.

Figure 1: **Flow chart of the phage-bacteria system.** The subscripts w and m refer to the wildtype (in green) and mutant (or virulent, in red) strain of phage λ, respectively. In the bacterial population ($E.\ coli$) of size N, bacteria are either susceptible (S), lysogenic (L) – i.e., carrying a prophage in their chromosome (small ellipses) – or carrying phages (small pentagons) replicating in their cytoplasm prior to lysis (Y); V is the free virus stage (culture medium). Width of the arrows between S, L and Y reflects the relative rates of different events: the wildtype strain is transmitted mostly vertically (high probability of lysogenization and low rate of prophage reactivation) while the virulent strain is transmitted mostly horizontally (low probability of lysogenization and high rate of prophage reactivation). Dashed arrows symbolize the role of the free viral particles in the force of infection (epidemiological feedback). See notations in **Table 1**.

Figure 2: **Theoretical predictions**. At $t = 0$, bacteria are at carrying capacity K with initial prevalence 1% (epidemic treatment) or 99% (endemic treatment). The initial prophage ratio for the two strains is 1:1. See **Table S1** for parameter values. (A) The virulent phage is selected for at the early state of the epidemic. Our approximation for the trajectory of the logit-frequency of the virulent phage (light red shading, see **SI Appendix §S2.2**) predicts the dynamics of the virulent phage when the density of susceptible cells is kept constant (dashed red line). The slope of the upper bound of our approximation is proportional to $-\Delta\phi = \phi_w - \phi_m$ (equation (5)). As only lysogens are introduced in the susceptible bacterial population at $t = 0$, we let the phage-bacteria system reach its new dynamical regime and only start our approximation at $t = 1$ (white dot). However, the rapid drop in the availability of susceptible hosts due to the viral epidemic reduces the increase of the virulent phage relative to our approximation (compare the full red line and the dashed red line). (B) The virulent phage is counter-selected when the epidemic reaches high prevalence (indicated with the white dots in the epidemic and endemic treatments). At the end of the epidemic, the logit-frequency of the virulent phages decreases linearly with negative slope $-\Delta\alpha = \alpha_w - \alpha_m$ (equation (6)). (C) The virulent strain is more frequent among free viruses (V) than among prophages (L). When the system reaches high prevalence, we predict the differentiation between these two compartments converges towards an equilibrium that is approximately equal to $1 + \Delta\alpha/\alpha_w = \alpha_m/\alpha_w$ (equation (8)).

Figure 3: **Differentiation of the virulent strain between free phages and infected cells**. We compute $q(t)(1 - g(t))/\big((1 - q(t))g(t)\big)$ for each chemostat (thin lines) and the corresponding geometric mean for each treatment (thick lines). The value 1 (horizontal grey line) corresponds to no differentiation. For each chemostat, values before the time point the system has reached a prevalence of 95% are shown in transparent. With high prevalence, we have $g(t) \approx p(t)$ and we thereby obtain the differentiation of the virulent strain between free phages (V) and prophages (L). In these conditions, such differentiation is expected to reach an equilibrium – the horizontal dashed line indicates the corresponding geometric mean (4.60) across all chemostats and time points – with theoretical value $1 + \Delta\alpha/\alpha_w = \alpha_m/\alpha_w$ (cf. **Fig. 2-C**).

Figure 4: **Fitted values on experimental data**. Fitted values (thick dark lines) were obtained from the simulation based on the best MLE estimations (see **Table 2**). We estimate model parameters using experimental data (light-colored thin lines) from the evolution experiment designed in (Berngruber et al., 2013); initial prevalence is either low (endemic condition, in blue) or high (epidemic condition, in red). Light-colored thick lines correspond to mean logit-values across all replicates per treatment. (A) Logit-prevalence of infection; (B) logit-frequency of cells infected (involved either in a lysogenic or a lytic cycle) by the the mutant (virulent) phage; (C) logit-frequency of the mutant (virulent) in the culture medium (free virus stage).

Table 1: **Notations.** The subscripts w and m refer to the wildtype strain and the mutant (or virulent) strain λcI857 of phage λ, respectively. Overlines refer to mean values of life-history traits across all genotypes. SD stands for 'standard deviation'.

Term	Definition
N	Bacterial population
S	Susceptible bacteria
L, L_w, L_m	Lysogenic bacteria (bacteria carrying a prophage)
Y, Y_w, Y_m	Bacteria with phages replicating in their cytoplasm prior to lysis
V, V_w, V_m	Free viral particles (virions)
P	Prevalence ($1 - S/N$)
p	Frequency of L cells infected by the mutant strain m
q	Frequency of the mutant strain m in the free virus stage (V)
f	Frequency of Y cells infected by the mutant strain m
g	Frequency of cells infected (either Y or L) by the mutant strain m
$\overline{\alpha}$, α_w, α_m	Rates of prophage reactivation; $\alpha_w < \alpha_m$
$\overline{\phi}$, ϕ_w, ϕ_m	Probabilities of lysogenization (phage integration into the host genome upon infection); $\phi_w > \phi_m$
$\Delta\alpha$, $\Delta\phi$	Phenotypic differences between the mutant and wildtype strain; $\Delta\alpha = \alpha_m - \alpha_w$, $\Delta\phi = \phi_m - \phi_w$
a	Adsorption rate of free viral particles onto the surface of bacteria
b	Probability of fusion (injection of the phage genome) upon adsorption
τ	Lysis rate; $1/\tau$ corresponds to the mean lysis time
B	Burst size (number of released viral particles upon lysis)
r	Bacterial intrinsic growth rate
K	Carrying capacity for the bacterial population: $K = 10^9\ cells$
δ	Dilution rate of the continuous culture: $\delta = 0.8\ h^{-1}$
$\mathscr{R}_{0,w}$, $\mathscr{R}_{0,m}$	Basic reproduction numbers of the phage
$\mathscr{S}$	Selection gradient of the virulent phage (rate at which it grows or declines in frequency on the logit scale)
$\mathcal{Q}^{VL}$	Differentiation of the virulent phage between free phages and prophages
σ_P, σ_g, σ_q	SD of measurement errors for logit(P), logit(g) and logit(q), respectively

Table 2: **MLE parameter estimations.** In a first step, parameters α_w and α_m are estimated directly using analytic approximations and linear models. In a second step, we fix α_w and α_m to their previous point estimates and, starting from 2,000 sets of initial values, non-linear optimizations maximizing the overall log-likelihood with the experimental data are run to estimate the remaining parameters. 95% CIs are based on sieve bootstrap (6,686 sets of estimates). Throughout, fixed parameters are: $K = 10^9\ cells$, $\delta = 0.8\ h^{-1}$ and $B = 80\ virus\,.\,cell^{-1}$.

Parameter	Estimated value	95% CI	Unit
	Parameters of the phage		
α_w	2.58×10^{-3}	$[1.94 \times 10^{-4},\ 7.87 \times 10^{-3}]$	h^{-1}
α_m	1.19×10^{-2}	$[1.03 \times 10^{-3},\ 2.85 \times 10^{-2}]$	h^{-1}
ϕ_w	0.347	$[0.198,\ 0.501]$	$-$
ϕ_m	2.91×10^{-2}	$[7.70 \times 10^{-3},\ 6.94 \times 10^{-2}]$	$-$
a	10^{-6}	$[4.34 \times 10^{-7},\ 10^{-6}]$	$h^{-1}\,.\,cell^{-1}$
b	4.94×10^{-2}	$[2.44 \times 10^{-2},\ 7.51 \times 10^{-2}]$	$-$
τ	1.08	$[0.83,\ 3.95]$	h^{-1}
	Parameter of the bacteria		
r	1.60	$[0.77,\ 5]$	h^{-1}
	Initial conditions		
$P_0^{epidemic}$	4.24×10^{-3}	$[1.00 \times 10^{-3},\ 1.13 \times 10^{-2}]$	$-$
$P_0^{endemic}$	99.61×10^{-2}	$[0.988,\ 0.998]$	$-$
p_0	0.416	$[0.4,\ 0.52]$	$-$
	SD of measurement errors		
σ_P	1.51	$[1.13,\ 2.00]$	$-$
σ_g	0.52	$[0.52,\ 0.75]$	$-$
σ_q	0.83	$[0.74,\ 1.16]$	$-$

References

Alizon, S., Hurford, A., Mideo, N., & Van Baalen, M. (2009). Virulence evolution and the trade-off hypothesis: History, current state of affairs and the future. *Journal of evolutionary biology*, *22*(2), 245–259. https://doi.org/10.1111/j.1420-9101.2008.01658.x

Alizon, S., & Michalakis, Y. (2015). Adaptive virulence evolution: The good old fitness-based approach. *Trends in ecology & evolution*, *30*(5), 248–254. https://doi.org/10.1016/j.tree.2015.02.009

Anderson, R. M., & May, R. M. (1982). Coevolution of hosts and parasites. *Parasitology*, *85*(2), 411–426. https://doi.org/10.1017/S0031182000055360

Berngruber, T. W., Froissart, R., Choisy, M., & Gandon, S. (2013). Evolution of virulence in emerging epidemics. *PLoS pathogens*, *9*(3), e1003209. https://doi.org/10.1371/journal.ppat.1003209

Berngruber, T. W., Weissing, F. J., & Gandon, S. (2010). Inhibition of superinfection and the evolution of viral latency. *Journal of Virology*, *84*(19), 10200–10208. https://doi.org/10.1128/jvi.00865-10

Bonachela, J. A., & Levin, S. A. (2014). Evolutionary comparison between viral lysis rate and latent period. *Journal of Theoretical Biology*, *345*, 32–42. https://doi.org/10.1016/j.jtbi.2013.12.006

Bruce, J. B., Lion, S., Buckling, A., Westra, E. R., & Gandon, S. (2021). Regulation of prophage induction and lysogenization by phage communication systems. *Current Biology*, *31*(22), 5046–5051. https://doi.org/10.1016/j.cub.2021.08.073

Bühlmann, P. (1997). Sieve bootstrap for time series. *Bernoulli*, 123–148. https://doi.org/10.2307/3318584

Bull, J. J. (1994). Virulence. *Evolution*, *48*(5), 1423–1437. https://doi.org/10.1111/j.1558-5646.1994.tb02185.x

Day, T. (2002). On the evolution of virulence and the relationship between various measures of mortality. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *269*(1498), 1317–1323. https://doi.org/10.1098/rspb.2002.2021

Day, T., & Proulx, S. R. (2004). A general theory for the evolutionary dynamics of virulence. *The American Naturalist*, *163*(4), E40–E63. https://doi.org/10.1086/382548

De Paepe, M., Tournier, L., Moncaut, E., Son, O., Langella, P., & Petit, M.-A. (2016). Carriage of λ latent virus is costly for its bacterial host due to frequent reactivation in monoxenic mouse intestine. *PLoS genetics*, *12*(2), e1005861. https://doi.org/10.1371/journal.pgen.1005861

Diekmann, O., Heesterbeek, J., & Roberts, M. G. (2010). The construction of next-generation matrices for compartmental epidemic models. *Journal of the royal society interface*, *7*(47), 873–885. https://doi.org/10.1098/rsif.2009.0386

Echols, H. (1972). Developmental pathways for the temperate phage: Lysis vs lysogeny. *Annual review of genetics*, *6*(1), 157–190. https://doi.org/10.1146/annurev.ge.06.120172.001105

Frank, S. A. (1996). Models of parasite virulence. *The Quarterly review of biology*, *71*(1), 37–78. https://doi.org/10.1086/419267

Gandon, S. (2016). Why be temperate: Lessons from bacteriophage λ. *Trends in microbiology*, *24*(5), 356–365. https://doi.org/10.1016/j.tim.2016.02.008

Gandon, S., & Day, T. (2007). The evolutionary epidemiology of vaccination. *Journal of the Royal Society Interface*, *4*(16), 803–817. https://doi.org/10.1098/rsif.2006.0207

Husimi, Y., Nishigaki, K., Kinoshita, Y., & Tanaka, T. (1982). Cellstat—A continuous culture system of a bacteriophage for the study of the mutation rate and the selection process at the DNA level. *Review of Scientific Instruments*, *53*(4), 517–522. https://doi.org/10.1063/1.1137002

Hyndman, R. J., & Khandakar, Y. (2008). Automatic time series forecasting: The forecast package for r. *Journal of statistical software*, *27*, 1–22. https://doi.org/10.18637/jss.v027.i03

Kelley, C. T. (1999). *Iterative methods for optimization*. SIAM.

Kourilsky, P., & Knapp, A. (1975). Lysogenization by bacteriophage lambda: III.-multiplicity dependent phenomena occuring upon infection by lambda. *Biochimie*, *56*(11-12), 1517–1523. https://doi.org/10.1016/S0300-9084(75)80275-6

Kourilsky, P. (1973). Lysogenization by bacteriophage lambda: I. multiple infection and the lysogenic response. *Molecular and General Genetics MGG*, *122*(2), 183–195. https://doi.org/10.1007/BF00435190

Lenski, R. E., & May, R. M. (1994). The evolution of virulence in parasites and pathogens: Reconciliation between two competing hypotheses. *Journal of theoretical biology*, *169*(3), 253–265. https://doi.org/10.1006/jtbi.1994.1146

Lindberg, H. M., McKean, K. A., & Wang, I.-N. (2014). Phage fitness may help predict phage therapy efficacy. *Bacteriophage*, *4*(4), e964081. https://doi.org/10.4161/21597073.2014.964081

Little, J. W., Shepley, D. P., & Wert, D. W. (1999). Robustness of a gene regulatory circuit. *The EMBO journal*, *18*(15), 4299–4307. https://doi.org/10.1093/emboj/18.15.4299

Lwoff, A. (1953). Lysogeny. *Bacteriological reviews*, *17*(4), 269–337. https://doi.org/10.1128/br.17.4.269-337.1953

Ptashne, M. (1992). *A genetic switch: Phage λ and higher organisms.* Oxford: Blackwell Publishers.

R Core Team. (2022). *R: A language and environment for statistical computing.* R Foundation for Statistical Computing. Vienna, Austria. https://www.R-project.org/

Rutberg, B., & Rutberg, L. (1965). Role of superinfecting phage in lysis inhibition with phage t4 in escherichia coli. *Journal of bacteriology*, *90*(4), 891–894. https://doi.org/10.1128/jb.90.4.891-894.1965

Shao, Y., & Wang, I.-N. (2008). Bacteriophage adsorption rate and optimal lysis time. *Genetics*, *180*(1), 471–482. https://doi.org/10.1534/genetics.108.090100

Soetaert, K., Petzoldt, T., & Setzer, R. W. (2010). Solving differential equations in r: Package desolve. *Journal of statistical software*, *33*, 1–25. https://doi.org/10.18637/jss.v033.i09

St-Pierre, F., & Endy, D. (2008). Determination of cell fate selection during phage lambda infection. *Proceedings of the National Academy of Sciences*, *105*(52), 20705–20710. https://doi.org/10.1073/pnas.0808831105

Sussman, R., & Jacob, F. (1962). [On a thermosensitive repression system in the escherichia coli lambda bacteriophage]. *Comptes rendus hébdomadaires des séances de l'Académie des sciences*, *254*, 1517–1519.

Wahl, L. M., Betti, M. I., Dick, D. W., Pattenden, T., & Puccini, A. J. (2019). Evolutionary stability of the lysis-lysogeny decision: Why be virulent? *Evolution*, *73*(1), 92–98. https://doi.org/10.1111/evo.13648

Wang, I.-N. (2006). Lysis timing and bacteriophage fitness. *Genetics*, *172*(1), 17–26. https://doi.org/10.1534/genetics.105.045922

Wegrzyn, G., & Wegrzyn, A. (2005). Genetic switches during bacteriophage λ development. *Progress in nucleic acid research and molecular biology*, *79*, 1–48. https://doi.org/10.1016/S0079-6603(04)79001-7

Zeng, L., Skinner, S. O., Zong, C., Sippy, J., Feiss, M., & Golding, I. (2010). Decision making at a subcellular level determines the outcome of bacteriophage infection. *Cell*, *141*(4), 682–691. https://doi.org/10.1016/j.cell.2010.03.034

Zong, C., So, L.-h., Sepúlveda, L. A., Skinner, S. O., & Golding, I. (2010). Lysogen stability is determined by the frequency of activity bursts from the fate-determining gene. *Molecular systems biology*, *6*(1), 440. https://doi.org/10.1038/msb.2010.96