Spatial integration of sensory input and motor output in *Pseudomonas aeruginosa* chemotaxis through colocalized distribution

Zhengyu Wu[1,2,4,5], Maojin Tian[2,3,5], Sanyuan Fu[2,5], Min Chen[2], Rongjing Zhang[2,*], Junhua Yuan[2,*]

[1]Research Center of Translational Medicine, Central Hospital Affiliated to Shandong First Medical University, Jinan, Shandong 250013, China.

[2]Hefei National Research Center for Physical Sciences at the Microscale and Department of Physics, University of Science and Technology of China, Hefei, Anhui 230026, China.

[3] Center of Translational Medicine, Zibo Central Hospital Affiliated to Binzhou Medical University, Zibo, Shandong 255036, China.

[4]Research Center of Translational Medicine, Jinan Central Hospital, Shandong University, Jinan, Shandong 250013, China.

[5]These authors contributed equally

*Corresponding authors:

Junhua Yuan or Rongjing Zhang, Department of Physics, University of Science and Technology of China, Hefei, Anhui, 230026, China, tel +86-551-63600851, email: jhyuan@ustc.edu.cn; rjzhang@ustc.edu.cn.

Abstract

The opportunistic pathogen *Pseudomonas aeruginosa* serves as a model organism for studying multiple signal transduction pathways. The chemoreceptor cluster, a core component of the chemotaxis pathway, is assembled from hundreds of proteins. The unipolar distribution of receptor clusters has long been recognized, yet the precise mechanism governing their assembly remains elusive. Here, we directly observed the relative positions of the flagellar motor and chemoreceptor cluster using flagellar filament labeling and gene editing techniques. Surprisingly, we found that both are located at the same cell pole, with the distribution pattern controlled by the polar anchor protein FlhF. Additionally, the efficient assembly of the chemoreceptor cluster is partially dependent on the integrity of the motor structure. Furthermore, we discovered that overexpression of the chemotaxis regulatory protein CheY leads to high intracellular levels of the second messenger c-di-GMP, triggering cell aggregation. Therefore, the colocalization of the chemoreceptor cluster and flagellum in *P. aeruginosa* serves to avoid cross-pathway signaling interference, enabling cells to conduct various physiological activities in an orderly manner.

Introduction

In biological systems, the process by which proteins self-assemble into organized complex structures is ubiquitous[1,2]. Pole-to-pole protein oscillations in the *Min* system ensure that the FtsZ ring, a crucial component of cell division, is placed precisely in the middle of the cell body[3,4]. In response to antibiotic treatment and heat stress, some bacteria generate multiple protein foci throughout the cytoplasm[5,6]. The polar localization of the secretion system, such as the type VI secretion system, is mediated by targeting proteins and potentially facilitates host-pathogen interactions[7,8]. Thus, spontaneous assembly is accompanied by the formation of spatial patterns that order the intracellular environment.

To grow and colonize within the host, bacteria have evolved a mechanism for

migrating towards more favorable environments. The locomotion organ (the flagellar

motor) and the chemotaxis pathway play crucial roles in achieving this goal. The

former provides physical drive, while the latter offers directional guidance. The

initiation of cell reorientation is controlled by motor switching, which is governed by

the chemotaxis two-component signaling pathway. This pathway comprises

transmembrane chemoreceptors, a histidine kinase CheA, an adaptor protein CheW, a

response regulator CheY, and two adaptation proteins, CheB and CheR. Receptors

collaborate with CheA and CheW to form chemosensory arrays in a hexagonal

pattern[9,10]. Upon phosphorylation of CheY by the kinase at the array, it freely diffuses

to the motor and binds to FliM, a component of the motor switch complex, to affect

motor switching[11-13]. It is worth noting that there exists diversity among different

bacterial species in the placement of flagellar motors[14], whereas chemoreceptor

clusters are typically found at one or both ends of the cell[15]. These two structural units

are functionally connected. However, it remains unclear whether there is an

interaction in their spatial distribution and what the specific regulatory mechanism

might be.

Pseudomonas aeruginosa, a common human opportunistic pathogen, possesses four

chemosensory pathways that perform distinct functions and are stimulated by signal

binding to 26 chemoreceptors[16]. Among them, F6 pathway-related receptors that

mediate chemotaxis, as well as the flagellum are localized at the same cell pole[17,18].

Furthermore, the polar anchor protein FlhF has been reported to guide the flagellum to

grow at the cell pole[19,20]. However, the mechanism by which chemoreceptor

array-related proteins aggregate and self-assemble at the cell pole is controversial.

The distribution of the chemosensory arrays in the peritrichous bacterium *Escherichia*

coli has been extensively studied and attributed to several mechanisms, including

stochastic self-assembly[21,22], membrane curvature sorting[23,24], and inefficient

clustering in the lateral region[25]. However, these mechanisms cannot explain the

unipolar distribution pattern observed in *P. aeruginosa*. Unlike *E. coli*, specific genes

responsible for the placement of the chemosensory array have been identified in several bacterial species. In *Caulobacter crescentus*, chemosensory arrays assemble at the new cell pole with the help of TipN and TipF proteins[26]. A tripartite ParC-ParP-CheA interaction network was reported to promote polar localization of chemosensory array in *Vibrio parahaemolyticus*[27,28]. However, as there are no related genes in *P. aeruginosa*, the mechanism of its chemoreceptor array distribution remains a mystery.

Here, we combined gene editing and in vivo fluorescence imaging of flagellar filaments to directly observe the distribution of chemoreceptor arrays and flagellar motors, proposing a cooperative construction model of chemotaxis network and flagella during the entire division cycle of *P. aeruginosa*. The core focus of this study is to clarify the regulatory mechanism of its chemoreceptor array distribution. We found a substantial association between the assembly of the flagellar motor and the chemoreceptor array. The complete assembly of the motor serves as a partial prerequisite for the assembly of the receptor array, and its assembly site is also regulated by the polar anchor protein FlhF. Furthermore, by introducing exogenously expressed CheY protein, we found that this triggers the expression of c-di-GMP at a high level. From this, we infer that the colocalization of the chemoreceptor array and the flagellum in *P. aeruginosa* avoids the cross-pathway interference of signaling molecules, thus providing a guarantee for the coexistence of multiple chemosensory pathways.

Results

Robust generation of daughter cell with both chemotaxis network and flagellar motor

CheY has been shown to colocalize with chemoreceptors[29]. To visualize the distribution of chemoreceptor arrays in cells, we fused the gene encoding the enhanced yellow fluorescent protein (eYFP) to the *cheY* gene in the *P. aeruginosa*

chromosome (Fig. 1A). All genetic modifications were carried out at the native

position on the chromosome, allowing the expression of chemotaxis proteins in

precise stoichiometry from their natural promoters (Fig. 1B). The mutant strain with

cheY-eyfp was able to moderately expand on soft agar plates[17], proving that its

chemotaxis behavior was not notably affected, so it was selected for subsequent

experiments. As shown in Fig. 1C, CheY-eYFP was mainly located at the cell pole,

suggesting that CheA, CheW and CheY in *P. aeruginosa* form a signal transduction

complex that was primarily distributed at the cell pole, as described previously[17].

From our measurements, nearly 90% (332/372) of cells contain obvious receptor

clusters, and factors such as fluorescence bleaching may have caused the loss of

fluorescent spots in some cells. In addition, we observed the presence of fluorescent

spots at both ends of some cells with a large aspect ratio (probably approaching cell

division), indicating that the newly generated progeny cells will have a complete

chemotaxis network. Next, we sought to observe the distribution of the flagella in

cells simultaneously to understand the motility of future progeny.

Our recent study demonstrated the successful labeling of the flagellar filament with

thiol-reactive fluorescent dye by introducing cysteine mutations into the flagellin

FliC[30,31] (Fig. 1B). To avoid fluorescence interference, we utilized a dye with an

excitation peak near 568 nm, considering that the excitation peak of eYFP is 513 nm,

and the emission peak is 527 nm. We employed a xenon lamp as the excitation light

source and switched filters to enable simultaneous fluorescence imaging of flagellar

filaments and chemosensory arrays within the same cell. Remarkably, our findings

revealed a surprising colocalization of chemosensory arrays and flagellar filaments at

the same end of the cell body, and this colocalization was consistent, as shown in Fig.

1D. Similar to the chemoreceptor arrays, we observed the phenomenon of double

flagella symbiosis in cells with a large aspect ratio, suggesting that *P. aeruginosa* has

evolved a cell division mechanism with precise timing regulation. Before cell division,

both poles of the mother cell assembled chemoreceptor arrays and flagellar motors.

This ensures that daughter cells possess complete motility and chemotaxis, thereby

greatly enhancing the environmental adaptability of the population.

FlhF mediates proper unipolar localization of the chemoreceptor array

Given that the physiological and physical environments of both cell poles are nearly

identical, it is unlikely that the unipolar distribution of the chemoreceptor array can be

attributed to passive regulatory factors. The number and distribution of flagellar

motors and chemoreceptor arrays vary among different bacterial species[14,15], and the

localization system for flagella has been well studied. Specifically, the FlhF-FlhG

system, discovered in several monotrichous bacterial species, has been shown to

control the location and number of flagella[32,33]. In *P. aeruginosa*, a knockout of *flhF*

leads to mis-localized flagellar assembly[19,20]. Considering the consistent

colocalization pattern between chemosensory arrays and flagellar motors in *P.*

aeruginosa, we speculate that the distribution of receptor clusters is also regulated by

similar molecular mechanisms.

To investigate the role of FlhF in the localization of receptor arrays, we constructed a

Δ*flhF* strain for fluorescence observation. The results revealed that the chemoreceptor

arrays no longer grow robustly at the cell pole (Fig. 2A), and the assembly positions

of the flagellar motor change accordingly, colocalizing with the receptor arrays (Fig.

2B). We also quantified the proportion of individuals with obvious receptor clusters,

which was more than 80% (182/221), similar to the wild-type strain. These

experimental results suggest that the polar anchoring protein FlhF has a minimal

impact on the assembly efficiency of *P. aeruginosa*'s chemosensory array, but it

strictly controls its unipolar distribution. Furthermore, a consistent colocalization

between the flagellar motor and chemosensory array was observed, independent of

FlhF. Thus, it is crucial to determine whether there is a causal relationship in the

assembly order of these two structural units.

Motor structural integrity is a prerequisite for chemoreceptor self-assembly

Cryo-electron microscopy has successfully revealed the complete flagella structure in

various bacterial species. This structure includes multiple components such as the

inner-membrane MS ring, the cytoplasmic C ring, and the internal protein secretion

system, all of which exhibit strong similarities[34]. This suggests that the core structure

of the flagellar motor is highly conserved. The MS ring, composed of 26 FliF proteins,

forms the foundation of the flagellar structure[35]. In addition to serving as a mounting

platform for the C ring, the MS ring also acts as a protective shell for the internal

protein secretion machinery[36]. FliG, the main component protein of the C ring, binds

directly to the cytoplasmic surface of FliF in a 1:1 ratio[37]. Thus, the efficient assembly

of the C ring requires preassembly of the MS ring. To observe the distribution of the

chemosensory array following the disruption of the C ring and MS ring, we

constructed Δ*fliG* and Δ*fliF* mutants.

We performed fluorescence observation on the Δ*fliG* mutant, which has a disrupted

flagellar motor C ring. The results showed that the proportion of Δ*fliG* cells with

obvious chemoreceptor clusters decreased significantly compared to the wild-type

cells (59.8%, 140/234), although the chemosensory arrays remained at a single cell

pole. We also examined the Δ*fliF* mutant, which has a disrupted flagellar motor MS

ring. Similar to the Δ*fliG* mutant, the proportion of individuals with obvious

chemoreceptor clusters decreased significantly (62.5%, 202/323). Subsequently, we

constructed a Δ*flhF*Δ*fliF* mutant and observed that the proportion of individuals with

obvious chemoreceptor clusters further decreased (50.7%, 341/672). To ascertain

whether it is motor integrity rather than functionality that influences the efficiency of

chemosensory array assembly, we constructed a stator mutant (Δ*motA*Δ*motCD*). In

this mutant, the motor is completely stalled while the structure remains intact. We

found that the mutant performed similarly to the wild-type strain in terms of

chemosensory array assembly (84.3% of individuals with obvious clusters, 204/242).

The fluorescence imaging of receptor clusters for multiple mutants in this section is

shown in Figure. 3A. We utilized Western blotting to measure the expression levels of

CheY, which were found to be similar across the different strains (Fig. 3B). This

further substantiated that the observed phenomenon is based on the structural integrity

of the motor rather than the protein expression level. Additionally, we quantified the proportion of cells with receptor clusters and the absolute fluorescence intensity of individual clusters (Fig. 3C-D). Overall, our findings suggest that the polar anchor protein FlhF and the structural integrity of the motor are crucial for the formation of chemosensory arrays. The former guides them to the appropriate site, while the latter influences the assembly efficiency of the chemoreceptor complex.

Colocalization of chemosensory array and flagellar motor avoids cross-pathway interference in *P. aeruginosa* signal transduction.

The distribution patterns of receptors associated with various signal transduction pathways in *P. aeruginosa* vary considerably. For instance, the biofilm formation-related receptor WspA, unlike the chemotaxis receptors discussed here, is distributed throughout the cell. This distribution is thought to enhance its sensitivity to mechanical perturbations of the cell membrane[38]. Previous studies speculated that the highly consistent position of chemoreceptor arrays and flagellar motors enhances bacterial chemotaxis performance[28]. However, a spatial separation between the chemoreceptors and flagellar motors should not significantly impact the temporal comparison in bacterial chemotaxis. The diffusion time of the phosphorylated chemotaxis regulatory protein CheY-P across the longest distance in a bacterial cell body (along the cell's long axis) is approximately 100 ms[39,40], while the timescale for the chemotaxis temporal comparison is on the order of seconds[41]. Consequently, the potential significance of the colocalization of chemoreceptors and flagellar motors remains unclear.

The signal transduction pathways in *E. coli* are relatively simple, and the chemotaxis response regulator CheY-P affects only the regulation of motor switching[11]. In contrast, the intracellular signal transduction pathways in *P. aeruginosa* are more complex. We thus hypothesized that the co-localization of chemoreceptors and flagellar motors in *P. aeruginosa* ensures locally distributed CheY-P molecules, thereby eliminating the need for a high level of intracellular CheY-P and avoiding

potential side effects on other signaling pathways.

To test the potential effect of an increased intracellular CheY-P level, we constructed

the CheY expression plasmid *cheY*-pJN105, transformed it into the wild-type *P.*

aeruginosa strain, and induced it with varying concentrations of arabinose. We

observed that the higher the inducer concentration, the more pronounced the cell

aggregation became in the field of view (Fig. 4A). The transition from planktonic

individuals to aggregated communities is similar to biofilm formation, which is

known to be accompanied by a significant increase in intracellular c-di-GMP levels.

To investigate whether a higher level of intracellular CheY-P increased the

intracellular c-di-GMP level, we sought to detect the c-di-GMP level. We introduced

the plasmid pCdrA-gfp into *P. aeruginosa*, using it as a c-di-GMP biosensor with the

fluorescence intensity proportional to the c-di-GMP level[42]. To ensure the coexistence

of the plasmids, the intracellular CheY concentration was controlled by the plasmid

cheY-pME6032. We found that the average single-cell fluorescence intensity of the

CheY overexpression strain was 71.8% higher than that of the wild-type strain, while

the c-di-GMP level of the *cheY* deletion strain was 58.6% lower than that of the

wild-type (Fig. 4B).

Therefore, the colocalization of chemosensory array and flagellar motor facilitates the

precise regulation of cell swimming direction within the low intracellular CheY-P

concentration threshold, using locally distributed CheY-P molecules. This helps to

avoid the occurrence of the aforementioned cross-pathway interference.

Summary and Discussion

P. aeruginosa harbors multiple signal transduction systems that regulate

flagella-mediated swimming motility (Che pathway), pili-mediated interface

twitching motility (Pil pathway), and biofilm formation (Wsp pathway)[16]. These

systems allow it to thrive in various ecological niches within complex external

environments. Here, employing a combination of chromosomal fluorescent protein

fusion and flagellar labeling techniques in living cells, we directly observed that the

chemosensory array of *P. aeruginosa* Che pathway and the flagellar motor share a

high degree of consistency in their assembly sites. We found that the chemosensory

array, comprising Che proteins, was consistently located at the cell pole where the

flagellum was positioned. Based on these observations, we deduced the construction

mode of the chemotactic network and flagellar motor during the entire cell growth

cycle (Fig. 5A). As the cell body matures and elongates, a new flagellum grows at the

opposing cell pole, accompanied by the assembly of a fresh chemosensory array. The

cell then undergoes division from the middle, generating two daughter cells, each

equipped with fully functional motility and chemotaxis capabilities. This ensures

robust inheritance of these chemotaxis and motility-related macromolecular machines.

Based on our understanding of the regulation mechanism of flagellar position in *P.*

aeruginosa, we constructed a Δ*flhF* mutant and found that the position distribution of

its chemosensory array is actively regulated by FlhF-mediated molecular mechanisms,

rather than being passively restricted by membrane curvature factors as seen in *E. coli*.

A similar active regulation phenomenon was previously reported in *Vibrio cholerae*,

with the corresponding molecular regulation mechanism detailed extensively[28]. The

flagella and chemosensory arrays of newborn *V. cholerae* cells are located at the old

cell pole, with the polar anchor protein HubP distributed at both cell poles. As the cell

grows, HubP recruits the ParA homologue ParC. Subsequently, ParC recruits ParP to

assemble a new chemosensory array at the new cell pole. Upon cell division, HubP

relocates to the middle of the cell, ensuring its robust presences at both poles of the

daughter cells. Here, HubP recruits ParA homologues including FlhG, which

influence flagellar motor assembly. However, previous experiments confirmed that

the cell cycle-dependent polar localization of ParC and chemotactic proteins is

independent of the flagellar position regulatory protein FlhF. This suggests that the

localization of flagellar motors and chemosensory arrays in *V. cholerae* is controlled

separately through different pathways. In contrast, flagellar motors were no longer robustly distributed at the cell pole after *flhF* was knocked out in *P. aeruginosa*, whereas the assembly site of chemosensory array and flagellar motor remained remarkably consistent. This suggests that FlhF regulates the distribution patterns of both and implies an interaction between them.

We further constructed Δ*fliG* and Δ*fliF* mutants, which disrupted the assembly of the flagellar motor. Surprisingly, we observed that the assembly of chemosensory array was impaired under these conditions, with a significant decrease in the proportion of individuals exhibiting obvious Che protein fluorescent bright spots. In contrast, the chemosensory array assembly in the stator mutant (Δ*motA*Δ*motCD*) was similar to that of the wild-type strain. This suggests that the integrity rather than functionality of the flagellar motor in *P. aeruginosa* is critical for the assembly of its chemosensory array, and its assembly efficiency is greatly reduced in the absence of flagellar motors. To eliminate the potential confounding effects of FlhF and flagellar motors, we constructed a Δ*flhF*Δ*fliF* mutant strain. Under this condition, the proportion of individuals with obvious Che protein fluorescent bright spots dropped to ~50%, indicating that passive regulation based on membrane curvature may still exist in *P. aeruginosa*, albeit to a lesser degree.

Is the remarkable consistency in the spatial distribution of these two independent structural units an unintentional outcome or a hidden mystery? By introducing a c-di-GMP monitoring system at the single-cell level, we discovered that overexpressing CheY substantially increases the intracellular c-di-GMP concentration. Previous studies generally believed that CheY-P, as a chemotactic regulatory protein, only influences flagellar motor switching. Here, we found that CheY-P, as a signaling molecule, can accomplish threshold-limited cross-pathway regulation. At low CheY-P concentrations, it is conservatively involved in motor switching regulation, as confirmed by previous studies. At high CheY-P concentrations, cell motility is suppressed, and c-di-GMP is triggered to express, though its specific mechanism

needs further exploration. The intracellular environment is highly crowded[43], and the

proximal growth pattern shown in Fig. 5B has multiple physiological significances.

On one hand, chemotaxis-related protein can precisely regulate cell motility with

minimal synthesis costs. On the other hand, the chemotactic network and the flagellar

motor are spatially integrated within the cell, ensuring their robust existence within

the cell body and preventing mutual interference with collateral pathways.

Many bacteria possess multiple signal transduction pathways. The orderly operation

of each pathway within the micrometer-scale space presents a fascinating scientific

problem. Here, we utilized the well-known chemotaxis network as a starting point to

shed light on this problem, providing inspiration for future in-depth understanding of

related issues.

Materials and methods

Strains and cell culture

The strains and plasmids used in this study are listed in Table 1. The *Escherichia coli*

TOP10 strain was used for standard genetic manipulations. A single-colony isolate

was grown in 3 ml of LB broth (1% Bacto tryptone, 0.5% yeast extract and 1% NaCl)

overnight to saturation on a rotary shaker (250 rpm) at 37°C. An aliquot was diluted

1:100 into 10 ml of LB broth and grew to exponential phase. Appropriate antibiotics

were added if necessary to prevent plasmid loss: for *E. coli*, 15 μg/ml Gentamicin and

25 μg/ml Tetracycline; for *P. aeruginosa*, 30 μg/ml Gentamicin and 50 μg/ml

Tetracycline. To induce protein expression, Isopropyl β-D-1-thiogalactopyranoside

(IPTG, 1.0 mM) was added to strains with pME6032-derivative vectors, and

arabinose (0.005%, 0.01%, 0.05%, 0.1%) was added to strains with

pJN105-derivative vectors. 2 ml of cells were harvested by centrifugation at 2000×g

for 2 min, washed twice in an equal volume of motility buffer (MB) [50 mM

potassium phosphate, 15 μM EDTA, 0.15 M NaCl, 5 mM Mg^{2+} and 10 mM lactic acid

(pH 7.0)][44], and resuspended in 5 ml MB for subsequent fluorescence observation.

Construction of in-frame deletion mutants

We used polymerase chain reaction (PCR) to generate ~1000 bp DNA fragments with upstream (Up) and downstream (Dn) sequences flanking the target gene (including *flhF*, *fliF*, *motA* and *fliG*) to be deleted. The Up and Dn DNA sequences were fused with the linearized pex18gm vector using Gibson assembly[45]. The resulting vectors were transformed into *P. aeruginosa* by electroporation, and the desired knockout mutants were obtained by double selections on gentamicin plates (LB plates with 30 μg/ml gentamicin) and sucrose plates (NaCl-free LB plates with 15% sucrose) at 37°C[46].

Generation of chromosomal fusions of *yfp* to *P. aeruginosa* gene

To generate *eyfp* fusion at the N-terminus of *cheY*, we first used PCR to generate upstream and downstream DNA sequences flanking the *cheY* gene. A subsequent PCR reaction aimed to yield the *cheY-eyfp* fusion with a 3× glycine linker (GGCGGAGGA), with pVS88 serving as the source of the enhanced yellow fluorescent (*eyfp*). The three DNA sequences, along with the linearized vector, were fused to *cheY-eyfp*-pex18gm using Gibson assembly. Finally, gene replacement was used to transfer the fusion constructs into the *P. aeruginosa* chromosome by homologous recombination.

Flagellar staining and fluorescence imaging

Flagellar filaments were labeled by following the protocol described previously[47]. Cells (1 ml of exponential-phase culture) were harvested by centrifugation at 2000×*g* for 10 min and washed twice in 1 ml MB. The final pellet was adjusted to a volume of ~100 μl that concentrated the bacterial 10-fold. Alexa Fluor 568 maleimide (Invitrogen-Molecular Probes) was added to a final concentration of 20 μg/ml, and labeling was allowed to proceed for 30 min at room temperature with gyration at 80 rpm. Unused dye was then removed by washing the cells with MB three times, and the final pellet was resuspended in MB.

For fluorescence imaging, 50 μl of cells were added to the chamber (constructed with two double-sticky tapes, a glass slide, and a poly-L-Lysine coated coverslip), incubated for 3 min, and rinsed with 100 μl MB. The boundary of the chamber was then sealed with Apiezon vacuum grease. The chamber was placed on a Nikon Ti-E inverted fluorescence microscope with a 100× oil-immersion objective and a sCMOS camera (Primer95B, Photometrics). The flagellar filament and chemosensory array of cells were observed separately with corresponding filter-set, using a 200-ms exposure time. To quantify the fluorescence intensity of the receptor cluster, we took the intensity maximum point as the center and extract a 3×3 pixel matrix around it. The mean value of the elements in this matrix represents the final fluorescence intensity.

Western Blotting

The total protein of *P. aeruginosa* strain cells in each treatment group was extracted and its concentration was measured with Bicinchoninic acid (BCA Protein Quantification Kit, Yeasen Biotechnology Co., Ltd) when the cells grew to exponential phase with same reduced turbidity (OD_{600} between 0.9 and 1.0). The exact amount of protein was subjected to SDS-PAGE electrophoresis, then transferred to a nitrocellulose membrane, which was blocked with 50 mg/mL skim milk in TBST buffer (20 mM Tris, 150 mM NaCl [pH7.4], 0.1%Tween 20) for 1h at room temperature. Since the N-terminal sequence of EYFP is almost identical to that of GFP, a rabbit anti-GFP antibody (1:2000, Yeasen Biotechnology Co., Ltd) and a Mouse anti-β-actin antibody (1:2000, Invitrogen) were added and incubated at 4°C overnight. The nitrocellulose membrane was gently washed with TBST for 20 min, three times. Subsequently, HRP labeled goat anti-rabbit IgG (1:4000, Abcam) and rabbit anti-mouse IgG (1:4000, Abcam) were added and incubated at room temperature for 1 h. The nitrocellulose membrane was again gently washed with TBST for 20 min, three times. Then, the membrane was developed using the ECL chemiluminescence detection system (Beyotime P0018FS) and visualized with the ChemiDoc XRS+ system (Bio-Rad). Three independent experiments were conducted

for reproducibility.

Monitoring c-di-GMP signal at the single-cell level

For measurements of c-di-GMP signaling, the validated reporter plasmid pCdrA-*gfp*,

which produces green fluorescent protein (GFP) in response to an increase in

c-di-GMP, was introduced into the experimental system. A 1000× diluted culture was

inoculated into the apparatus described in the previous section and allowed to adhere

to the surface for ~10 min, with the slight difference that the coverslips were not

coated with poly-L-Lysine to avoid interference from background fluorescence.

Bacteria were imaged using a Nikon Ti-E inverted microscope, which was equipped

with an EMCCD camera (DU897, iXon3, Andor Technology). We used a 100×

oil-immersion objective. Bacteria were illuminated with a 488-nm laser

(Sapphire 488−200 mW, Coherent), using a standard GFP filter sets in the

fluorescence imaging system with an exposure time of 500 ms. To ensure the

consistency of culture and observation conditions, the data of wild-type and

cheY-pME6032 carried strains were collected at the same time. More than 200

independent individuals of each strain were used for data analysis.

Single-cell c-di-GMP concentration was characterized by calculating fluorescence

intensity. The average fluorescence intensity of a single cell was calculated by

dividing the total fluorescence intensity of a single cell by the cell volume. Bacterial

were simplified to a hollow cylindrical trunk with hollow hemispherical caps at both

ends. The cell volume can be calculated by taking the length of the short axis of the

cell body as the diameter of the sphere and the length of the long axis as the sum of

the height of the cylinder and the diameter of the sphere. The mean fluorescence

intensity of the cell was obtained by subtracting the average background fluorescence

intensity from the total fluorescence intensity of the cell, and then dividing by the cell

volume.

References

Halatek, J., Brauns, F. & Frey, E. Self-organization principles of intracellular pattern formation. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373**, 20170107 (2018).

Thanbichler, M. & Shapiro, L. Getting organized--how bacterial cells move proteins and DNA. *Nat. Rev. Microbiol.* **6**, 28-40 (2008).

Loose, M., Kruse, K. & Schwille, P. Protein self-organization: lessons from the min system. *Annu. Rev. Biophys.* **40**, 315-336 (2011).

Rothfield, L., Taghbalout, A. & Shih, Y. L. Spatial control of bacterial division-site placement. *Nat. Rev. Microbiol.* **3**, 959-968 (2005).

Fay, A. & Glickman, M. S. An essential nonredundant role for mycobacterial DnaK in native protein folding. *PLoS Genet.* **10**, e1004516 (2014).

Vaubourgeix, J. *et al.* Stressed mycobacteria use the chaperone ClpB to sequester irreversibly oxidized proteins asymmetrically within and between cells. *Cell Host Microbe.* **17**, 178-190 (2015).

Wang, J., Brodmann, M. & Basler, M. Assembly and Subcellular Localization of Bacterial Type VI Secretion Systems. *Annu. Rev. Microbiol.* **73**, 621-638 (2019).

Low, H. H. *et al.* Structure of a type IV secretion system. *Nature* **508**, 550-553 (2014).

Briegel, A. *et al.* Bacterial chemoreceptor arrays are hexagonally packed trimers of receptor dimers networked by rings of kinase and coupling proteins. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 3766-3771 (2012).

Zhang, P., Khursigara, C. M., Hartnell, L. M. & Subramaniam, S. Direct visualization of Escherichia coli chemotaxis receptor arrays using cryo-electron microscopy. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3777-3781 (2007).

Cluzel, P., Surette, M. & Leibler, S. An ultrasensitive bacterial motor revealed by monitoring signaling proteins in single cells. *Science* **287**, 1652-1655 (2000).

Sourjik, V. & Berg, H. C. Binding of the Escherichia coli response regulator CheY to its target measured in vivo by fluorescence resonance energy transfer. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 12669-12674 (2002).

Welch, M., Oosawa, K., Aizawa, S. & Eisenbach, M. Phosphorylation-dependent binding of a signal molecule to the flagellar switch of bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 8787-8791 (1993).

Schuhmacher, J. S., Thormann, K. M. & Bange, G. How bacteria maintain location and number of flagella? *FEMS Microbiol. Rev.* **39**, 812-822 (2015).

Jones, C. W. & Armitage, J. P. Positioning of bacterial chemoreceptors. *Trends Microbiol.* **23**, 247-256 (2015).

Matilla, M. A., Martin-Mora, D., Gavira, J. A. & Krell, T. Pseudomonas aeruginosa as a Model To Study Chemosensory Pathway Signaling. *Microbiol. Mol. Biol. Rev.* **85**, e00151-00120 (2021).

17   Guvener, Z. T., Tifrea, D. F. & Harwood, C. S. Two different Pseudomonas
aeruginosa chemosensory signal transduction complexes localize to cell poles
and form and remould in stationary phase. *Mol. Microbiol.* **61**, 106-118
(2006).
18   Amako, K. & Umeda, A. Flagellation of Pseudomonas aeruginosa during the
cell division cycle. *Microbiol. Immunol.* **26**, 113-117 (1982).
19   Schniederberend, M., Abdurachim, K., Murray, T. S. & Kazmierczak, B. I.
The GTPase activity of FlhF is dispensable for flagellar localization, but not
motility, in Pseudomonas aeruginosa. *J. Bacteriol.* **195**, 1051-1060 (2013).
20   Murray, T. S. & Kazmierczak, B. I. FlhF is required for swimming and
swarming in Pseudomonas aeruginosa. *J. Bacteriol.* **188**, 6995-7004 (2006).
21   Greenfield, D. *et al.* Self-organization of the Escherichia coli chemotaxis
network imaged with super-resolution light microscopy. *PLoS Biol.* **7**,
e1000137 (2009).
22   Thiem, S. & Sourjik, V. Stochastic assembly of chemoreceptor clusters in
Escherichia coli. *Mol. Microbiol.* **68**, 1228-1236 (2008).
23   Strahl, H. *et al.* Transmembrane protein sorting driven by membrane curvature.
Nat. Commun. **6**, 8728 (2015).
24   Draper, W. & Liphardt, J. Origins of chemoreceptor curvature sorting in
Escherichia coli. *Nat. Commun.* **8**, 14838 (2017).
25   Koler, M., Peretz, E., Aditya, C., Shimizu, T. S. & Vaknin, A. Long-term
positioning and polar preference of chemoreceptor clusters in E. coli. *Nat.*
Commun. **9**, 4444 (2018).
26   Huitema, E., Pritchard, S., Matteson, D., Radhakrishnan, S. K. & Viollier, P. H.
Bacterial birth scar proteins mark future flagellum assembly site. *Cell* **124**,
1025-1037 (2006).
27   Ringgaard, S., Schirner, K., Davis, B. M. & Waldor, M. K. A family of
ParA-like ATPases promotes cell pole maturation by facilitating polar
localization of chemotaxis proteins. *Genes Dev.* **25**, 1544-1555 (2011).
28   Ringgaard, S. *et al.* ParP prevents dissociation of CheA from chemotactic
signaling arrays and tethers them to a polar anchor. *Proc. Natl. Acad. Sci. U. S.*
A. **111**, E255-E264 (2014).
29   Sourjik, V. & Berg, H. C. Localization of components of the chemotaxis
machinery of *Escherichia coli* using fluorescent protein fusions. *Mol.*
Microbiol. **37**, 740-751 (2000).
30   Wu, Z., Tian, M., Zhang, R. & Yuan, J. Dynamics of the Two Stator Systems
in the Flagellar Motor of Pseudomonas aeruginosa Studied by a Bead Assay.
Appl. Environ. Microbiol. **87**, e0167421 (2021).
31   Tian, M., Wu, Z., Zhang, R. & Yuan, J. A new mode of swimming in singly
flagellated Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* **119**,
e2120508119 (2022).
32   Altegoer, F., Schuhmacher, J., Pausch, P. & Bange, G. From molecular
evolution to biobricks and synthetic modules: a lesson by the bacterial
flagellum. *Biotechnol. Genet. Eng. Rev.* **30**, 49-64 (2014).

Kazmierczak, B. I. & Hendrixson, D. R. Spatial and numerical regulation of flagellar biosynthesis in polarly flagellated bacteria. *Mol. Microbiol.* **88**, 655-663 (2013).

Minamino, T. & Imada, K. The bacterial flagellar motor and its structural diversity. *Trends Microbiol.* **23**, 267-274 (2015).

Suzuki, H., Yonekura, K. & Namba, K. Structure of the rotor of the bacterial flagellar motor revealed by electron cryomicroscopy and single-particle image analysis. *J. Mol. Biol.* **337**, 105-113 (2004).

Minamino, T., Imada, K. & Namba, K. Molecular motors of the bacterial flagella. *Curr. Opin. Struct. Biol.* **18**, 693-701 (2008).

Levenson, R., Zhou, H. & Dahlquist, F. W. Structural insights into the interaction between the bacterial flagellar motor proteins FliF and FliG. *Biochemistry* **51**, 5052-5060 (2012).

O'Connor, J. R., Kuwada, N. J., Huangyutitham, V., Wiggins, P. A. & Harwood, C. S. Surface sensing and lateral subcellular localization of WspA, the receptor in a chemosensory-like system leading to c-di-GMP production. *Mol. Microbiol.* **86**, 720-729 (2012).

Sagawa, T. *et al.* Single-cell E. coli response to an instantaneously applied chemotactic signal. *Biophys. J.* **107**, 730-739 (2014).

Terasawa, S. *et al.* Coordinated reversal of flagellar motors on a single Escherichia coli cell. *Biophys. J.* **100**, 2193-2200 (2011).

Segall, J. E., Block, S. M. & Berg, H. C. Temporal comparisons in bacterial chemotaxis. *Proc. Natl. Acad. Sci. U. S. A.* **83**, 8987-8991 (1986).

Rybtke, M. T. *et al.* Fluorescence-based reporter for gauging cyclic di-GMP levels in Pseudomonas aeruginosa. *Appl. Environ. Microbiol.* **78**, 5060-5069 (2012).

Zhou, H., Rivas, G. & Minton, A. P. Macromolecular crowding and confinement: biochemical, biophysical, and potential physiological consequences. *Annu. Rev. Biophys.* **37**, 375-397 (2008).

Cai, Q., Li, Z., Ouyang, Q., Luo, C. & Gordon, V. D. Singly Flagellated Pseudomonas aeruginosa Chemotaxes Efficiently by Unbiased Motor Regulation. *mBio* **7**, e00013-00016 (2016).

Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**, 343-345 (2009).

Hmelo, L. R. *et al.* Precision-engineering the Pseudomonas aeruginosa genome with two-step allelic exchange. *Nat. Protoc.* **10**, 1820-1841 (2015).

Turner, L., Zhang, R., Darnton, N. C. & Berg, H. C. Visualization of Flagella during bacterial Swarming. *J. Bacteriol.* **192**, 3259-3267 (2010).

Author Contributions

J.Y. and R.Z. designed the work; Z.W., M.T. and S.F. performed the measurements with help from M.C.; Z.W and S.F analyzed the data; Z.W., J.Y. and R.Z. wrote the paper. Z.W., M.T. and S.F. contributed equally to this work.

Conflicts of interest
The authors declare no competing interests.

Acknowledgments
This work was supported by National Natural Science Foundation of China Grants (11925406, 12090053, 12304241, and 12304251), a Grant from the Ministry of Science and Technology of China (2019YFA0709303), Grants from the Natural Science Foundation of Shandong Province No. ZR2023QA111 and ZR2023QC168, and a Grant from the Tai Shan Young Scholar Foundation of Shandong Province (tsqn.2024.T.M.).

582 **Table 1.** Strains and plasmids used in this study.

Strain，Plasmid	Genotype, phenotype, and Description	Source
Strains		
P. aeruginosa		
PAO1	wild-type strain	Fan Jin Group
PAO1 *fliC*[T394C]	Replacement of chromosomal *fliC* in PAO1	Ref 31
PAO1 *fliC*[T394C] *cheY-eyfp*	*yfp* fusions at the N-terminus of *cheY* in PAO1 *fliC*[T394C]	This work
PAO1 *fliC*[T394C] *cheY-eyfp* Δ*flhF*	Nonpolar *flhF* deletion in PAO1 *fliC*[T394C] *cheY-eyfp*	This work
PAO1 *fliC*[T394C] *cheY-eyfp* Δ*fliF*	Nonpolar *fliF* deletion in PAO1 *fliC*[T394C] *cheY-eyfp*	This work
PAO1 *fliC*[T394C] *cheY-eyfp* Δ*fliG*	Nonpolar *fliG* deletion in PAO1 *fliC*[T394C] *cheY-eyfp*	This work
PAO1 *fliC*[T394C] *cheY-eyfp* Δ*flhF* Δ*fliF*	Nonpolar *fliF* deletion in PAO1 *fliC*[T394C] *cheY-eyfp* Δ*flhF*	This work
PAO1 *fliC*[T394C] *cheY-eyfp* Δ*motA* Δ*motCD*	Nonpolar *motAB* and *motCD* deletion in PAO1 *fliC*[T394C] *cheY-eyfp*	This work
E. coli		
Top10	F-*mcrA* Δ(*mrr-hsRMS-mcrBC*)Φ80*lacZ*Δ*M15* Δ*lacX74 recA1* *ara*D139 Δ*(araleu)*7697 *galU galK rpsL(*Nal[R]*) endA1 nupG*	Invitrogen
Plasmids		
pex18gm	oriT[+]sacB[+]; gene replacement vector with MCS from pUC18; Gm[r]	Fan Jin Group
flhF-pex18gm	In-frame deletion of *flhF* cloned into pex18gm; Gm[r]	This work
fliF-pex18gm	In-frame deletion of *fliF* cloned into pex18gm; Gm[r]	This work
fliG-pex18gm	In-frame deletion of *fliG* cloned into pex18gm; Gm[r]	This work
motA-pex18gm	In-frame deletion of *motA* cloned into pex18gm; Gm[r]	This work
motCD-pex18gm	In-frame deletion of *motCD* cloned into pex18gm; Gm[r]	Ref 30
cheY-eyfp-pex18gm	*yfp* fusions *cheY* cloned into pex18gm; Gm[r]	This work
cheY-pJN105	*cheY* overexpression vector in pJN1005, *cheY* expression is controlled by P[BAD] promoter; Gm[r]	This work
cheY-pME6032	*cheY* overexpression vector in pME6032, *cheY* expression is controlled by P[lac] promoter; Tet[r]	This work
pCdrA-*gfp*	pUCP22-NotI based cyclic di-GMP level reporter, Gm[r]	Ref 42

Fig. 1. **A**. Schematic diagram of the chemotaxis signal transduction network and flagellar motor of *P. aeruginosa*. Fusion protein CheY-EYFP and fluorescently labeled flagellar filaments were used as markers to indicate the position of the chemosensory array and motor, respectively. **B.** The labeling mechanism of flagellar filaments and chemotaxis regulatory protein CheY. Filaments (with cysteine point mutation FliC^{T394C}) were labeled through sulfhydryl-maleimide conjugation, and *cheY-eyfp* fusion with a 3× glycine linker was used to visualize chemosensory array positions. **C.** Subcellular localization of CheY-EYFP in the wild-type strain of *P. aeruginosa*. CheY-EYFP is mainly located at the single cell pole, and the white arrow points to individuals with obvious chemosensory arrays at both cell poles, which generally have a large aspect ratio of the cell body. The yellow dashed box marks the cell outline. **D.** The merged imaging of flagellar filaments and CheY-EYFP in the wild-type strain of *P. aeruginosa*, where flagellar motor and chemosensory array

601   colocalize in cells. White arrows point to individuals about to be divided, and the

602   yellow dashed box marks the cell outline. The scale bar is 1 μm.

Fig. 2. **A**. Subcellular localization of CheY-EYFP in the Δ*flhF* strain of *P. aeruginosa*. CheY-EYFP is no longer robustly distributed at the single cell pole. The yellow dashed box marks the cell outline. **B.** The merged imaging of flagellar filaments and CheY-EYFP in the Δ*flhF* strain of *P. aeruginosa*, flagellar motor and chemosensory array still colocalize in cells. The yellow dashed box marks the cell outline. The scale bar is 1 μm.

Fig. 3. **A**. Subcellular localization of CheY-EYFP in various *P. aeruginosa* strains. The scale bar is 10 μm. **B.** Western blot analysis was performed to detect CheY expression in various *P. aeruginosa* strains. β-actin was used as the reference protein. **C.** Occurrence probability of obvious chemosensory arrays in *P. aeruginosa* wild-type and several mutant strains. The proportion of individuals with obvious chemosensory arrays decreased significantly in the motor-incomplete strains (Δ*fliF* and Δ*fliG*), and this value was further reduced after *flhF* knockout (Δ*fliF*Δ*flhF*). The Δ*fliF* and Δ*motA*Δ*motCD* strains have a similar chemosensory array occurrence probability as the wild-type strain. **D.** Absolute fluorescence brightness of the chemosensory array in *P. aeruginosa* wild-type and several mutant strains.

Fig. 4. **A**. The evolution of cell aggregation as the intracellular CheY concentration increases by induction with higher concentrations of arabinose. The scale bar is 10 μm. **B.** Quantitative characterization of intracellular c-di-GMP levels at different CheY concentrations. From top to bottom, they correspond to the Δ*cheY* strain (N=198), wildtype strain (N=249) and CheY overexpression strain (N=228), respectively.

Polar flagellar
Receptor cluster

Chemoattractant
MCP   Methyl-accepting chemotaxis proteins
Flagellar motor
High concentration CheY-P region

Fig. 5. **A**. The construction mode of the chemotaxis network and flagellar motor throughout the complete cell growth cycle. **B**. The proximal growth mode of the *P. aeruginosa* flagellar motor and receptor clusters will effectively regulate the spatial range of CheY action, thus avoiding unintentional cross-pathway regulation.