Spatial integration of sensory input and motor output in *Pseudomonas aeruginosa* chemotaxis through colocalized distribution

Zhengyu Wu[1,2,4,5], Maojin Tian[2,3,5], Sanyuan Fu[2,5], Min Chen[2], Rongjing Zhang[2,*], Junhua Yuan[2,*]

[1]Research Center of Translational Medicine, Central Hospital Affiliated to Shandong First Medical University, Jinan, Shandong 250013, China.

[2]Hefei National Research Center for Physical Sciences at the Microscale and Department of Physics, University of Science and Technology of China, Hefei, Anhui 230026, China.

[3]Department of Critical Care Medicine, Zibo Central Hospital Affiliated to Binzhou Medical University, Zibo, Shandong 255036, China.

[4]Research Center of Translational Medicine, Jinan Central Hospital, Shandong University, Jinan, Shandong 250013, China.

[5]These authors contributed equally

*Corresponding authors:

Junhua Yuan or Rongjing Zhang, Department of Physics, University of Science and Technology of China, Hefei, Anhui, 230026, China, tel +86-551-63600851, email: jhyuan@ustc.edu.cn; rjzhang@ustc.edu.cn.

Abstract

The opportunistic pathogen *Pseudomonas aeruginosa* serves as a model organism for studying multiple signal transduction pathways. The chemoreceptor cluster, a core component of the chemotaxis pathway, is assembled from hundreds of proteins. The unipolar distribution of receptor clusters has long been recognized, yet the precise mechanism governing their assembly remains elusive. Here, we directly observed the relative positions of the flagellar motor and chemoreceptor cluster using flagellar filament labeling and gene editing techniques. Surprisingly, we found that both are located at the same cell pole, with the distribution pattern controlled by the polar anchor protein FlhF. Additionally, the efficient assembly of the chemoreceptor cluster is partially dependent on the integrity of the motor structure. Furthermore, we discovered that overexpression of the chemotaxis regulatory protein CheY leads to high intracellular levels of the second messenger c-di-GMP, triggering cell aggregation. Therefore, the colocalization of the chemoreceptor cluster and flagellum in *P. aeruginosa* serves to avoid cross-pathway signaling interference, enabling cells to conduct various physiological activities in an orderly manner.

Introduction

In biological systems, the process by which proteins self-assemble into organized complex structures is widespread[1,2]. Pole-to-pole protein oscillations in the *Min* system ensure that the FtsZ ring, a crucial component of cell division, is placed precisely in the middle of the cell body[3,4]. In response to antibiotic treatment and heat stress, some bacteria generate multiple protein foci throughout the cytoplasm[5,6]. The polar localization of the secretion system, such as the type □ secretion system, is mediated by targeting proteins and potentially facilitates host-pathogen interactions[7,8]. Thus, spontaneous assembly is accompanied by the formation of spatial patterns that organize the intracellular environment.

To grow and colonize in various ecological niches, bacteria have evolved a

mechanism for migrating towards more favorable environments. The locomotion

organ (the flagellar motor) and the chemotaxis pathway play crucial roles in achieving

this goal. The former provides physical drive, while the latter offers directional

guidance. The initiation of cell reorientation is controlled by motor switching, which

is governed by the chemotaxis two-component signaling pathway. This pathway

comprises transmembrane chemoreceptors, a histidine kinase CheA, an adaptor

protein CheW, a response regulator CheY, and two adaptation proteins, CheB and

CheR. Receptors collaborate with CheA and CheW to form chemotaxis complex in a

hexagonal pattern[9,10]. Upon phosphorylation of CheY by the kinase at the complex, it

freely diffuses to the motor and binds to FliM, a component of the motor switch

complex, to affect motor switching[11-13]. It is worth noting that there exists diversity

among different bacterial species in the placement of flagellar motors[14], whereas

chemoreceptor clusters are typically found at one or both ends of the cell[15]. These two

structural units are functionally connected. However, it remains unclear whether there

is an interaction in their spatial distribution and what the specific regulatory

mechanism might be.

Pseudomonas aeruginosa, a common human opportunistic pathogen, possesses four

chemosensory pathways that perform distinct functions and are stimulated by signal

binding to 26 chemoreceptors[16]. Among them, the proteins encoded by

chemotaxis-related genes collectively constitute the F6 pathway. Previous studies

have suggested that the receptors involved in this pathway localize to the old cell

pole[17], similar to the flagellum[18], implying colocalization of the receptors and the

flagellum at the same pole; however, direct evidence for this colocalization is still

lacking. Furthermore, the polar anchor protein FlhF has been reported to guide the

flagellum to grow at the cell pole[19,20]. However, the mechanism by which chemotaxis

complex-related proteins aggregate and self-assemble at the cell pole is controversial.

The distribution of the chemotaxis complex in the peritrichous bacterium *Escherichia*

coli has been extensively studied and attributed to several mechanisms, including

stochastic self-assembly[21,22], membrane curvature sorting[23,24], and inefficient

clustering in the lateral region[25]. However, these mechanisms cannot explain the

unipolar distribution pattern observed in *P. aeruginosa*. Unlike *E. coli*, specific genes

responsible for the placement of the chemotaxis complex have been identified in

several bacterial species. In *Caulobacter crescentus*, chemotaxis complex assemble at

the new cell pole with the help of TipN and TipF proteins[26]. A tripartite

ParC-ParP-CheA interaction network was reported to promote polar localization of

chemotaxis complex in *Vibrio parahaemolyticus*[27,28]. However, as there are no related

genes in *P. aeruginosa*, the mechanism of its chemotaxis complex distribution

remains a mystery.

Here, we combined gene editing and in vivo fluorescence imaging of flagellar

filaments to directly observe the distribution of chemotaxis complex and flagellar

motors, proposing a cooperative construction model of chemotaxis network and

flagella during the entire division cycle of *P. aeruginosa*. The core focus of this study

is to clarify the regulatory mechanism of its chemotaxis complex distribution. We

found a substantial association between the assembly of the flagellar motor and the

chemotaxis complex. The complete assembly of the motor serves as a partial

prerequisite for the assembly of the chemotaxis complex, and its assembly site is also

regulated by the polar anchor protein FlhF. Furthermore, by introducing exogenously

expressed CheY protein, we found that this triggers the expression of c-di-GMP at a

high level. From this, we infer that the colocalization of the chemotaxis complex and

the flagellum in *P. aeruginosa* avoids the cross-pathway interference of signaling

molecules, thus providing a guarantee for the coexistence of multiple chemosensory

pathways.

Results

Robust generation of daughter cell with both chemotaxis network and flagellar

motor

CheY has been shown to colocalize with chemoreceptors[17,29]. To visualize the distribution of chemotaxis complex in cells, we fused the gene encoding the enhanced yellow fluorescent protein (eYFP) to the *cheY* gene in the *P. aeruginosa* chromosome (Fig. 1A). All genetic modifications were carried out at the native position on the chromosome, allowing the expression of chemotaxis proteins in precise stoichiometry from their natural promoters (Fig. 1B). The mutant strain with *cheY-eyfp* was able to moderately expand on soft agar plates[17], proving that its chemotaxis behavior was not notably affected, so it was selected for subsequent experiments. As shown in Fig. 1C, CheY-eYFP was mainly located at the cell pole, suggesting that CheA, CheW and CheY in *P. aeruginosa* form a signal transduction complex that was primarily distributed at the cell pole, as described previously[17]. Representative large-field images containing many cells are shown in Fig. S1. Additionally, we constructed a plasmid expressing CheA-CFP and electroporated it into the *cheY-eyfp* strain. Fluorescence imaging revealed a high degree of spatial overlap between CheA-CFP and CheY-EYFP (Fig. S2), confirming that CheY-EYFP accurately marks the location of the chemoreceptor complex. From our measurements, nearly 90% (332/372) of cells contain obvious receptor clusters, and factors such as fluorescence bleaching may have caused the loss of fluorescent spots in some cells. In addition, we observed the presence of fluorescent spots at both ends of some cells with a large aspect ratio (probably approaching cell division), indicating that the newly generated progeny cells will have a complete chemotaxis network. Next, we sought to observe the distribution of the flagella in cells simultaneously to understand the motility of future progeny.

Our recent study demonstrated the successful labeling of the flagellar filament with thiol-reactive fluorescent dye by introducing cysteines into the flagellin FliC[30,31] (Fig. 1B). To avoid fluorescence interference, we utilized a dye with an excitation peak near 568 nm, considering that the excitation peak of eYFP is 513 nm, and the emission peak is 527 nm. We employed a xenon lamp as the excitation light source and switched filters to enable simultaneous fluorescence imaging of flagellar

filaments and chemotaxis complex within the same cell.

Remarkably, our findings revealed a surprising colocalization of chemotaxis complex

and flagellar filaments at the same end of the cell body, and this colocalization was

consistent, as shown in Fig. 1D. A typical large field was shown in Fig. S3. Similar to

the chemotaxis complex, we observed the phenomenon of double flagella symbiosis

in cells with a large aspect ratio, suggesting that *P. aeruginosa* has evolved a cell

division mechanism with precise timing regulation. Before cell division, both poles of

the mother cell assembled chemotaxis complex and flagellar motors. This ensures that

daughter cells possess complete motility and chemotaxis, thereby greatly enhancing

the environmental adaptability of the population.

FlhF mediates proper unipolar localization of the chemotaxis complex

Despite potential differences in the physical and especially physiological

environments at the two cell poles, it is unlikely that the unipolar distribution of the

chemotaxis complex can be attributed to passive regulatory factors. The number and

distribution of flagellar motors and chemotaxis complex vary among different

bacterial species[14,15], and the localization system for flagella has been well studied.

Specifically, the FlhF-FlhG system, discovered in several monotrichous bacterial

species, has been shown to control the location and number of flagella[32,33]. In *P.*

aeruginosa, a knockout of *flhF* leads to mis-localized flagellar assembly[19,20].

Considering the consistent colocalization pattern between chemotaxis complex and

flagellar motors in *P. aeruginosa*, we speculate that the distribution of chemotaxis

complex is also regulated by similar molecular mechanisms.

To investigate the role of FlhF in the localization of chemotaxis complex, we

constructed a Δ*flhF* strain for fluorescence observation. The results revealed that the

chemotaxis complex no longer grow robustly at the cell pole (Fig. 2A), and the

assembly positions of the flagellar motor change accordingly, colocalizing with the

chemotaxis complex (Fig. 2B). Representative large fields containing many cells are

shown in Fig. S1. Additionally, we categorized receptor cluster distribution patterns into three types: precise polar, near polar, and mid-cell. The relevant statistics for the wild-type strain and the Δ*flhF* mutant are presented in Fig. S4, which show a significant increase in the proportion of non-precise polar chemoreceptors following *flhF* knockout. We also quantified the proportion of individuals with obvious receptor clusters, which was more than 80% (182/221), similar to the wild-type strain. These experimental results suggest that the polar anchoring protein FlhF has a minimal impact on the assembly efficiency of *P. aeruginosa*'s chemotaxis complex, but it strictly controls its unipolar distribution. Furthermore, a consistent colocalization between the flagellar motor and chemotaxis complex was observed, independent of FlhF. Thus, it is crucial to determine whether there is a causal relationship in the assembly order of these two structural units.

Motor structural integrity is a prerequisite for chemoreceptor self-assembly

Cryo-electron microscopy has successfully revealed the complete flagella structure in various bacterial species. This structure includes multiple components such as the inner-membrane MS ring, the cytoplasmic C ring, and the internal protein secretion system, all of which exhibit strong similarities[34]. This suggests that the core structure of the flagellar motor is highly conserved. The MS ring, composed of 26 FliF proteins, forms the foundation of the flagellar structure[35]. In addition to serving as a mounting platform for the C ring, the MS ring also acts as a protective shell for the internal protein secretion machinery[36]. FliG, the main component protein of the C ring, binds directly to the cytoplasmic surface of FliF in a 1:1 ratio[37]. Thus, the efficient assembly of the C ring requires preassembly of the MS ring. To observe the distribution of the chemotaxis complex following the disruption of the C ring and MS ring, we constructed Δ*fliG* and Δ*fliF* mutants.

We performed fluorescence observation on the Δ*fliG* mutant, which has a disrupted flagellar motor C ring. The results showed that the proportion of Δ*fliG* cells with obvious chemoreceptor clusters decreased significantly compared to the wild-type

cells (59.8%, 140/234), although the chemotaxis complex remained at a single cell pole. We also examined the Δ*fliF* mutant, which has a disrupted flagellar motor MS ring. Similar to the Δ*fliG* mutant, the proportion of individuals with obvious chemotaxis complex decreased significantly (62.5%, 202/323). Subsequently, we constructed a Δ*flhF*Δ*fliF* mutant and observed that the proportion of individuals with obvious chemoreceptor clusters further decreased (50.7%, 341/672). To ascertain whether it is motor integrity rather than functionality that influences the efficiency of chemotaxis complex assembly, we constructed a stator mutant (Δ*motA*Δ*motCD*). In this mutant, the motor is completely stalled while the structure remains intact. We found that the mutant performed similarly to the wild-type strain in terms of chemotaxis complex assembly (84.3% of individuals with obvious clusters, 204/242). The fluorescence imaging of receptor clusters for multiple mutants in this section is shown in Figure. 3A, and corresponding large-field images are provided in Fig. S1. We utilized Western blotting to measure the expression levels of CheY, which were found to be similar across the different strains (Fig. 3B). These further substantiated that the observed phenomenon is based on the structural integrity of the motor rather than the protein expression level. In contrast, the *cheA* knockout strain was constructed to disrupt the assembly of the chemoreceptor complex. We fluorescently labeled its flagellar filaments and found that its phenotype was no different from that of the wild-type strain (Fig. S5), indicating that the assembly of the chemotactic receptor complex does not affect the flagellar assembly efficiency. The P ring (mainly composed of FlgI) is thought to act as a bushing for the peptidoglycan layer, and its absence results in partial assembly of the motor structure[38]. We constructed Δ*flgI* mutant and found that the proportion of cells with distinct chemotactic complexes was similar to that of the wild-type (Fig. S6). Additionally, we quantified the proportion of cells with receptor clusters (Fig. 3C). Overall, our findings suggest that the polar anchor protein FlhF and the structural integrity of the motor are crucial for the formation of chemotaxis complex. The former guides them to the appropriate site, while the latter influences the assembly efficiency of the chemoreceptor complex.

Colocalization of chemotaxis complex and flagellar motor avoids cross-pathway interference in *P. aeruginosa* signal transduction.

The distribution patterns of receptors associated with various signal transduction pathways in *P. aeruginosa* vary considerably. For instance, the biofilm formation-related receptor WspA, unlike the chemotaxis receptors discussed here, is distributed throughout the cell. This distribution is thought to enhance its sensitivity to mechanical perturbations of the cell membrane[39]. Previous studies speculated that the highly consistent position of chemotaxis complex and flagellar motors enhances bacterial chemotaxis performance[28]. However, the diffusion time of the phosphorylated chemotaxis regulatory protein CheY-P across the longest distance in a bacterial cell body (along the cell's long axis) is approximately 100 ms[40,41], whereas the timescale for the chemotaxis temporal comparison is on the order of seconds[42]. Additionally, a study by Fukuoka and colleagues reported that intracellular chemotaxis signal transduction requires approximately 240 ms beyond CheY or CheY-P diffusion time[41]. Moreover, the CCW/CW interval of the *P. aeruginosa* flagellar motor under normal conditions is 1-2 s, as determined by bead assays[30] or tethered cell assays[43]. Taken together, these indicate that for *P. aeruginosa*, which moves via a run-reverse mode, the potential 100 ms reduction in response time due to co-localization of the chemotaxis complex and motor has a limited effect on overall chemotaxis timing. Consequently, the physiological significance of the colocalization of chemoreceptors and flagellar motors remains unclear.

E. coli mediates chemotaxis through a single pathway involving five types of chemoreceptors[44]. As a core regulatory protein for chemotaxis, CheY participates in multiple processes such as adaptation and chemotaxis response, with CheY-P binds to the motor to regulate switching[11]. In contrast, *P. aeruginosa* has four distinct chemosensory pathways that perform different functions, involving different CheY homologs, and are stimulated by signals binding to 26 types of chemoreceptors. Consequently, *P. aeruginosa* possesses more complex chemosensory pathways[16]. We thus hypothesized that the co-localization of chemoreceptors and flagellar motors in *P.*

aeruginosa ensures locally distributed CheY-P molecules, thereby eliminating the

need for a high level of intracellular CheY-P and avoiding potential side effects on

other signaling pathways.

To test the potential effect of an increased intracellular CheY-P level, we constructed

the CheY expression plasmid *cheY*-pJN105, transformed it into the wild-type *P.*

aeruginosa strain, and induced it with varying concentrations of arabinose. We

observed that the higher the inducer concentration, the more pronounced the cell

aggregation became in the field of view (Fig. 4A). The transition from planktonic

individuals to aggregated communities is similar to biofilm formation, which is

known to be accompanied by a significant increase in intracellular c-di-GMP levels.

To investigate whether a higher level of intracellular CheY-P increased the

intracellular c-di-GMP level, we sought to detect the c-di-GMP level. We introduced

the plasmid pCdrA-gfp into *P. aeruginosa*, using it as a c-di-GMP biosensor with the

fluorescence intensity proportional to the c-di-GMP level[45]. To ensure the coexistence

of the plasmids, the intracellular CheY concentration was controlled by the plasmid

cheY-pME6032. The intracellular CheY levels induced by varying arabinose

concentrations were measured, showing clear differences across the concentration

gradient (Fig. S7). We found that the average single-cell fluorescence intensity of the

CheY overexpression strain was 71.8% higher than that of the wild-type strain, while

the c-di-GMP level of the *cheY* deletion strain was 58.6% lower than that of the

wild-type (Fig. 4B).

The colocalization of both the signaling source (the kinase) and sink (the phosphatase)

at the chemoreceptor complex at the cell pole results in a rapid decay of CheY-P

concentration within approximately 0.2 µm from the cell pole, leading to a nearly

uniform distribution elsewhere in the cell, as demonstrated by Vaknin and Berg[46]. This

spatial arrangement effectively confines high CheY-P levels to the pole region. When

the motor is also localized at the cell pole, the need for elevated CheY-P

concentrations throughout the cytoplasm is reduced. Therefore, the colocalization of chemotaxis complex and flagellar motor facilitates the precise regulation of cell swimming direction within the low intracellular CheY-P concentration threshold, using locally distributed CheY-P molecules. This helps to avoid the occurrence of the aforementioned cross-pathway interference. To further confirm that CheY overexpression promotes aggregation through increased c-di-GMP levels, we performed additional experiments co-overexpressing CheY and a phosphodiesterase (PDE) from *E. coli* to reduce intracellular c-di-GMP. These experiments showed that PDE expression mitigates cell aggregation caused by CheY overexpression in *P. aeruginosa* (Fig. S8).

Summary and Discussion

P. aeruginosa harbors multiple signal transduction systems that regulate flagella-mediated swimming motility (Che pathway), pili-mediated interface twitching motility (Pil pathway), and biofilm formation (Wsp pathway)[16]. These systems allow it to thrive in various ecological niches within complex external environments. Here, employing a combination of chromosomal fluorescent protein fusion and flagellar labeling techniques in living cells, we directly observed that the chemotaxis complex of *P. aeruginosa* Che pathway and the flagellar motor share a high degree of consistency in their assembly sites. We found that the chemotaxis complex, comprising Che proteins, was consistently located at the cell pole where the flagellum was positioned. Based on these observations, we deduced the construction mode of the chemotactic network and flagellar motor during the entire cell growth cycle (Fig. 5A). As the cell body matures and elongates, a new flagellum grows at the opposing cell pole, accompanied by the assembly of a fresh chemotaxis complex. The cell then undergoes division from the middle, generating two daughter cells, each equipped with fully functional motility and chemotaxis capabilities. This ensures robust inheritance of these chemotaxis and motility-related macromolecular machines.

Based on our understanding of the regulation mechanism of flagellar position in *P.*

aeruginosa, we constructed a Δ*flhF* mutant and found that the position distribution of

its chemotaxis complex is actively regulated by FlhF-mediated molecular mechanisms,

rather than being passively restricted by membrane curvature factors as seen in *E. coli*.

A similar active regulation phenomenon was previously reported in *Vibrio cholerae*,

with the corresponding molecular regulation mechanism detailed extensively[28]. The

flagella and chemotaxis complex of newborn *V. cholerae* cells are located at the old

cell pole, with the polar anchor protein HubP distributed at both cell poles. As the cell

grows, HubP recruits the ParA homologue ParC. Subsequently, ParC recruits ParP to

assemble a new chemotaxis complex at the new cell pole. Upon cell division, HubP

relocates to the middle of the cell, ensuring its robust presences at both poles of the

daughter cells. Here, HubP recruits ParA homologues including FlhG, which

influence flagellar motor assembly. However, previous experiments confirmed that

the cell cycle-dependent polar localization of ParC and chemotactic proteins is

independent of the flagellar position regulatory protein FlhF. This suggests that the

localization of flagellar motors and chemotaxis complex in *V. cholerae* is controlled

separately through different pathways. In contrast, flagellar motors were no longer

robustly distributed at the cell pole after *flhF* was knocked out in *P. aeruginosa*,

whereas the assembly site of chemotaxis complex and flagellar motor remained

remarkably consistent. This suggests that FlhF regulates the distribution patterns of

both and implies an interaction between them.

We further constructed Δ*fliG* and Δ*fliF* mutants, which disrupted the assembly of the

flagellar motor. Surprisingly, we observed that the assembly of chemotaxis complex

was impaired under these conditions, with a significant decrease in the proportion of

individuals exhibiting obvious Che protein fluorescent bright spots. In contrast, the

chemotaxis complex assembly in the stator mutant (Δ*motA*Δ*motCD*) was similar to

that of the wild-type strain. It should be noted that *P. aeruginosa* has a four-tiered

transcriptional regulatory circuit controlling flagellar biogenesis, in which FliG and

FliF belong to class II and are regulated by the master transcriptional regulator FleQ[47].

Chemotaxis complex-related genes (*cheA*, *cheW*, etc.) are regulated by intracellular free FliA proteins, and there is no evidence that FliA is regulated by proteins such as FliG. To further confirm that the expression of core chemotaxis proteins was not affected by the knockout of *fliG*, we performed Western blotting to measure CheY expression levels in wild-type, Δ*fliF* and Δ*fliG* strains, and found no significant differences (Fig. 3B). These results suggested that the integrity rather than functionality of the flagellar motor in *P. aeruginosa* is critical for the assembly of its chemotaxis complex, and its assembly efficiency is greatly reduced in the absence of flagellar motors. To eliminate the potential confounding effects of FlhF and flagellar motors, we constructed a Δ*flhF*Δ*fliF* mutant strain. Under this condition, the proportion of individuals with obvious Che protein fluorescent bright spots dropped to ~50%, indicating that passive regulation based on membrane curvature may still exist in *P. aeruginosa*, albeit to a lesser degree.

Is the remarkable consistency in the spatial distribution of these two independent structural units an unintentional outcome or a hidden mystery? Given that Kulasekara et al. previously proposed that CheA and its phosphorylation state can cause c-di-GMP heterogeneity among individual *P. aeruginosa* cells through a specific PDE[48,49], we sought to verify whether the phenotypic differences induced by CheY here are closely related to changes in the overall population level of c-di-GMP. We introduced a c-di-GMP monitoring system at the single-cell level and discovered that overexpression CheY substantially increases the intracellular c-di-GMP concentration. Previous studies generally believed that CheY-P, as a chemotactic regulatory protein, only influences flagellar motor switching. Here, we found that CheY-P, as a signaling molecule, can accomplish threshold-limited cross-pathway regulation. At low CheY-P concentrations, it is conservatively involved in motor switching regulation, as confirmed by previous studies. At high CheY-P concentrations, cell motility is suppressed, and c-di-GMP is triggered to express, though its specific mechanism needs further exploration. Unlike CheA, a structural protein of the chemotaxis complex, CheY is a protein that can shuttle in the cytoplasm and can also regulate the

intracellular c-di-GMP level. This finding expands our understanding of the connection between the internal chemotaxis network and the c-di-GMP pathway. The intracellular environment is highly crowded[50], and the proximal growth pattern shown in Fig. 5B has multiple physiological significances. On one hand, chemotaxis-related protein can precisely regulate cell motility with minimal synthesis costs. On the other hand, the chemotactic network and the flagellar motor are spatially integrated within the cell, ensuring their robust existence within the cell body and preventing mutual interference with collateral pathways.

Many bacteria possess multiple signal transduction pathways. The orderly operation of each pathway within the micrometer-scale space presents a fascinating scientific problem. Here, we utilized the well-known chemotaxis network as a starting point to shed light on this problem, providing inspiration for future in-depth understanding of related issues.

Materials and methods

Strains and cell culture

The strains and plasmids used in this study are listed in Table 1. The *Escherichia coli* TOP10 strain was used for standard genetic manipulations. A single-colony isolate was grown in 3 ml of LB broth (1% Bacto tryptone, 0.5% yeast extract and 1% NaCl) overnight to saturation on a rotary shaker (250 rpm) at 37°C. An aliquot was diluted 1:100 into 10 ml of LB broth and grew to exponential phase. Appropriate antibiotics were added if necessary to prevent plasmid loss: for *E. coli*, 15 μg/ml Gentamicin and 25 μg/ml Tetracycline; for *P. aeruginosa*, 30 μg/ml Gentamicin and 50 μg/ml Tetracycline. To induce protein expression, Isopropyl β-D-1-thiogalactopyranoside (IPTG, 1.0 mM) was added to strains with pME6032-derivative vectors, and arabinose (0.005%, 0.01%, 0.05%, 0.1%) was added to strains with pJN105-derivative vectors. 2 ml of cells were harvested by centrifugation at 2000×g for 2 min, washed twice in an equal volume of motility buffer (MB) [50 mM

409 potassium phosphate, 15 µM EDTA, 0.15 M NaCl, 5 mM Mg^{2+} and 10 mM lactic acid

410 (pH 7.0)][51], and resuspended in 5 ml MB for subsequent fluorescence observation.

Construction of in-frame deletion mutants

We used polymerase chain reaction (PCR) to generate ~1000 bp DNA fragments with

upstream (Up) and downstream (Dn) sequences flanking the target gene (including

flhF, *fliF*, *motA* and *fliG*) to be deleted. The Up and Dn DNA sequences were fused

with the linearized pex18gm vector using Gibson assembly[52]. The resulting vectors

were transformed into *P. aeruginosa* by electroporation, and the desired knockout

mutants were obtained by double selections on gentamicin plates (LB plates with 30

µg/ml gentamicin) and sucrose plates (NaCl-free LB plates with 15% sucrose) at

37°C[53].

Generation of chromosomal fusions of *yfp* to *P. aeruginosa* gene

To generate *eyfp* fusion at the N-terminus of *cheY*, we first used PCR to generate

upstream and downstream DNA sequences flanking the *cheY* gene. A subsequent PCR

reaction aimed to yield the *cheY-eyfp* fusion with a 3× glycine linker

(GGCGGAGGA), with pVS88 serving as the source of the enhanced yellow

fluorescent (*eyfp*). The three DNA sequences, along with the linearized vector, were

fused to *cheY-eyfp*-pex18gm using Gibson assembly. Finally, gene replacement was

used to transfer the fusion constructs into the *P. aeruginosa* chromosome by

homologous recombination.

Flagellar staining and fluorescence imaging

Flagellar filaments were labeled by following the protocol described previously[54].

Cells (1 ml of exponential-phase culture) were harvested by centrifugation at 2000×g

for 10 min and washed twice in 1 ml MB. The final pellet was adjusted to a volume of

~100 µl that concentrated the bacterial 10-fold. Alexa Fluor 568 maleimide

(Invitrogen-Molecular Probes) was added to a final concentration of 20 µg/ml, and

labeling was allowed to proceed for 30 min at room temperature with gyration at 80

rpm. Unused dye was then removed by washing the cells with MB three times, and

the final pellet was resuspended in MB.

For fluorescence imaging, 50 μl of cells were added to the chamber (constructed with

two double-sticky tapes, a glass slide, and a poly-L-Lysine coated coverslip),

incubated for 3 min, and rinsed with 100 μl MB. The boundary of the chamber was

then sealed with Apiezon vacuum grease. The chamber was placed on a Nikon Ti-E

inverted fluorescence microscope with a 100× oil-immersion objective and a sCMOS

camera (Primer95B, Photometrics). The flagellar filament and chemotaxis complex of

cells were observed separately with corresponding filter-set, using a 200-ms exposure

time. To quantify the fluorescence intensity of the receptor cluster, we took the

intensity maximum point as the center and extract a 3×3 pixel matrix around it. The

mean value of the elements in this matrix represents the final fluorescence intensity.

Western Blotting

The total protein of *P. aeruginosa* strain cells in each treatment group was extracted

and its concentration was measured with Bicinchoninic acid (BCA Protein

Quantification Kit, Yeasen Biotechnology Co., Ltd) when the cells grew to

exponential phase with same reduced turbidity (OD_{600} between 0.9 and 1.0). The

exact amount of protein was subjected to SDS-PAGE electrophoresis, then transferred

to a nitrocellulose membrane, which was blocked with 50 mg/mL skim milk in TBST

buffer (20 mM Tris, 150 mM NaCl [pH7.4], 0.1%Tween 20) for 1h at room

temperature. Since the N-terminal sequence of EYFP is almost identical to that of

GFP, a rabbit anti-GFP antibody (1:2000, Yeasen Biotechnology Co., Ltd) and a

Mouse anti-β-actin antibody (1:2000, Invitrogen) were added and incubated at 4°C

overnight. The nitrocellulose membrane was gently washed with TBST for 20 min,

three times. Subsequently, HRP labeled goat anti-rabbit IgG (1:4000, Abcam) and

rabbit anti-mouse IgG (1:4000, Abcam) were added and incubated at room

temperature for 1 h. The nitrocellulose membrane was again gently washed with

TBST for 20 min, three times. Then, the membrane was developed using the ECL

chemiluminescence detection system (Beyotime P0018FS) and visualized with the
ChemiDoc XRS+ system (Bio-Rad). Three independent experiments were conducted
for reproducibility.

Monitoring c-di-GMP signal at the single-cell level

For measurements of c-di-GMP signaling, the validated reporter plasmid pCdrA-*gfp*,
which produces green fluorescent protein (GFP) in response to an increase in
c-di-GMP, was introduced into the experimental system. A 1000× diluted culture was
inoculated into the apparatus described in the previous section and allowed to adhere
to the surface for ~10 min, with the slight difference that the coverslips were not
coated with poly-L-Lysine to avoid interference from background fluorescence.
Bacteria were imaged using a Nikon Ti-E inverted microscope, which was equipped
with an EMCCD camera (DU897, iXon3, Andor Technology). We used a 100×
oil-immersion objective. Bacteria were illuminated with a 488-nm laser
(Sapphire 488–200 mW, Coherent), using a standard GFP filter sets in the
fluorescence imaging system with an exposure time of 500 ms. To ensure the
consistency of culture and observation conditions, the data of wild-type and
cheY-pME6032 carried strains were collected at the same time. More than 200
independent individuals of each strain were used for data analysis.

Single-cell c-di-GMP concentration was characterized by calculating fluorescence
intensity. The average fluorescence intensity of a single cell was calculated by
dividing the total fluorescence intensity of a single cell by the cell volume. Bacterial
were simplified to a hollow cylindrical trunk with hollow hemispherical caps at both
ends. The cell volume can be calculated by taking the length of the short axis of the
cell body as the diameter of the sphere and the length of the long axis as the sum of
the height of the cylinder and the diameter of the sphere. The mean fluorescence
intensity of the cell was obtained by subtracting the average background fluorescence
intensity from the total fluorescence intensity of the cell, and then dividing by the cell
volume.

References

Halatek, J., Brauns, F. & Frey, E. Self-organization principles of intracellular pattern formation. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373**, 20170107 (2018).

Thanbichler, M. & Shapiro, L. Getting organized--how bacterial cells move proteins and DNA. *Nat. Rev. Microbiol.* **6**, 28-40 (2008).

Loose, M., Kruse, K. & Schwille, P. Protein self-organization: lessons from the min system. *Annu. Rev. Biophys.* **40**, 315-336 (2011).

Rothfield, L., Taghbalout, A. & Shih, Y. L. Spatial control of bacterial division-site placement. *Nat. Rev. Microbiol.* **3**, 959-968 (2005).

Fay, A. & Glickman, M. S. An essential nonredundant role for mycobacterial DnaK in native protein folding. *PLoS Genet.* **10**, e1004516 (2014).

Vaubourgeix, J. *et al.* Stressed mycobacteria use the chaperone ClpB to sequester irreversibly oxidized proteins asymmetrically within and between cells. *Cell Host Microbe.* **17**, 178-190 (2015).

Wang, J., Brodmann, M. & Basler, M. Assembly and Subcellular Localization of Bacterial Type VI Secretion Systems. *Annu. Rev. Microbiol.* **73**, 621-638 (2019).

Low, H. H. *et al.* Structure of a type IV secretion system. *Nature* **508**, 550-553 (2014).

Briegel, A. *et al.* Bacterial chemoreceptor arrays are hexagonally packed trimers of receptor dimers networked by rings of kinase and coupling proteins. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 3766-3771 (2012).

Zhang, P., Khursigara, C. M., Hartnell, L. M. & Subramaniam, S. Direct visualization of Escherichia coli chemotaxis receptor arrays using cryo-electron microscopy. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3777-3781 (2007).

Cluzel, P., Surette, M. & Leibler, S. An ultrasensitive bacterial motor revealed by monitoring signaling proteins in single cells. *Science* **287**, 1652-1655 (2000).

Sourjik, V. & Berg, H. C. Binding of the Escherichia coli response regulator CheY to its target measured in vivo by fluorescence resonance energy transfer. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 12669-12674 (2002).

Welch, M., Oosawa, K., Aizawa, S. & Eisenbach, M. Phosphorylation-dependent binding of a signal molecule to the flagellar switch of bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **90**, 8787-8791 (1993).

Schuhmacher, J. S., Thormann, K. M. & Bange, G. How bacteria maintain location and number of flagella? *FEMS Microbiol. Rev.* **39**, 812-822 (2015).

Jones, C. W. & Armitage, J. P. Positioning of bacterial chemoreceptors. *Trends Microbiol.* **23**, 247-256 (2015).

Matilla, M. A., Martin-Mora, D., Gavira, J. A. & Krell, T. Pseudomonas aeruginosa as a Model To Study Chemosensory Pathway Signaling. *Microbiol.*

Mol. Biol. Rev. **85**, e00151-00120 (2021).

17   Guvener, Z. T., Tifrea, D. F. & Harwood, C. S. Two different Pseudomonas
aeruginosa chemosensory signal transduction complexes localize to cell poles
and form and remould in stationary phase. *Mol. Microbiol.* **61**, 106-118
(2006).

18   Amako, K. & Umeda, A. Flagellation of Pseudomonas aeruginosa during the
cell division cycle. *Microbiol. Immunol.* **26**, 113-117 (1982).

19   Schniederberend, M., Abdurachim, K., Murray, T. S. & Kazmierczak, B. I.
The GTPase activity of FlhF is dispensable for flagellar localization, but not
motility, in Pseudomonas aeruginosa. *J. Bacteriol.* **195**, 1051-1060 (2013).

20   Murray, T. S. & Kazmierczak, B. I. FlhF is required for swimming and
swarming in Pseudomonas aeruginosa. *J. Bacteriol.* **188**, 6995-7004 (2006).

21   Greenfield, D. *et al.* Self-organization of the Escherichia coli chemotaxis
network imaged with super-resolution light microscopy. *PLoS Biol.* **7**,
e1000137 (2009).

22   Thiem, S. & Sourjik, V. Stochastic assembly of chemoreceptor clusters in
Escherichia coli. *Mol. Microbiol.* **68**, 1228-1236 (2008).

23   Strahl, H. *et al.* Transmembrane protein sorting driven by membrane curvature.
Nat. Commun. **6**, 8728 (2015).

24   Draper, W. & Liphardt, J. Origins of chemoreceptor curvature sorting in
Escherichia coli. *Nat. Commun.* **8**, 14838 (2017).

25   Koler, M., Peretz, E., Aditya, C., Shimizu, T. S. & Vaknin, A. Long-term
positioning and polar preference of chemoreceptor clusters in E. coli. *Nat.*
Commun. **9**, 4444 (2018).

26   Huitema, E., Pritchard, S., Matteson, D., Radhakrishnan, S. K. & Viollier, P. H.
Bacterial birth scar proteins mark future flagellum assembly site. *Cell* **124**,
1025-1037 (2006).

27   Ringgaard, S., Schirner, K., Davis, B. M. & Waldor, M. K. A family of
ParA-like ATPases promotes cell pole maturation by facilitating polar
localization of chemotaxis proteins. *Genes Dev.* **25**, 1544-1555 (2011).

28   Ringgaard, S. *et al.* ParP prevents dissociation of CheA from chemotactic
signaling arrays and tethers them to a polar anchor. *Proc. Natl. Acad. Sci. U. S.*
A. **111**, E255-E264 (2014).

29   Sourjik, V. & Berg, H. C. Localization of components of the chemotaxis
machinery of *Escherichia coli* using fluorescent protein fusions. *Mol.*
Microbiol. **37**, 740-751 (2000).

30   Wu, Z., Tian, M., Zhang, R. & Yuan, J. Dynamics of the Two Stator Systems
in the Flagellar Motor of Pseudomonas aeruginosa Studied by a Bead Assay.
Appl. Environ. Microbiol. **87**, e0167421 (2021).

31   Tian, M., Wu, Z., Zhang, R. & Yuan, J. A new mode of swimming in singly
flagellated Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* **119**,
e2120508119 (2022).

32   Altegoer, F., Schuhmacher, J., Pausch, P. & Bange, G. From molecular
evolution to biobricks and synthetic modules: a lesson by the bacterial

flagellum. *Biotechnol. Genet. Eng. Rev.* **30**, 49-64 (2014).

33   Kazmierczak, B. I. & Hendrixson, D. R. Spatial and numerical regulation of
flagellar biosynthesis in polarly flagellated bacteria. *Mol. Microbiol.* **88**,
655-663 (2013).

34   Minamino, T. & Imada, K. The bacterial flagellar motor and its structural
diversity. *Trends Microbiol.* **23**, 267-274 (2015).

35   Suzuki, H., Yonekura, K. & Namba, K. Structure of the rotor of the bacterial
flagellar motor revealed by electron cryomicroscopy and single-particle image
analysis. *J. Mol. Biol.* **337**, 105-113 (2004).

36   Minamino, T., Imada, K. & Namba, K. Molecular motors of the bacterial
flagella. *Curr. Opin. Struct. Biol.* **18**, 693-701 (2008).

37   Levenson, R., Zhou, H. & Dahlquist, F. W. Structural insights into the
interaction between the bacterial flagellar motor proteins FliF and FliG.
Biochemistry **51**, 5052-5060 (2012).

38   Hizukuri, Y., Yakushi, T., Kawagishi, I. & Homma, M. Role of the
intramolecular disulfide bond in FlgI, the flagellar P-ring component of
Escherichia coli. *J. Bacteriol.* **188**, 4190-4197 (2006).

39   O'Connor, J. R., Kuwada, N. J., Huangyutitham, V., Wiggins, P. A. & Harwood,
C. S. Surface sensing and lateral subcellular localization of WspA, the
receptor in a chemosensory-like system leading to c-di-GMP production. *Mol.*
Microbiol. **86**, 720-729 (2012).

40   Terasawa, S. *et al.* Coordinated reversal of flagellar motors on a single
Escherichia coli cell. *Biophys. J.* **100**, 2193-2200 (2011).

41   Sagawa, T. *et al.* Single-cell E. coli response to an instantaneously applied
chemotactic signal. *Biophys. J.* **107**, 730-739 (2014).

42   Segall, J. E., Block, S. M. & Berg, H. C. Temporal comparisons in bacterial
chemotaxis. *Proc. Natl. Acad. Sci. U. S. A.* **83**, 8987-8991 (1986).

43   Qian, C., Wong, C. C., Swarup, S. & Chiam, K. H. Bacterial tethering analysis
reveals a "run-reverse-turn" mechanism for Pseudomonas species motility.
Appl. Environ. Microbiol. **79**, 4734-4743 (2013).

44   Sourjik, V. Receptor clustering and signal processing in E. coli chemotaxis.
Trends Microbiol. **12**, 569-576 (2004).

45   Rybtke, M. T. *et al.* Fluorescence-based reporter for gauging cyclic di-GMP
levels in Pseudomonas aeruginosa. *Appl. Environ. Microbiol.* **78**, 5060-5069
(2012).

46   Vaknin, A. & Berg, H. C. Single-cell FRET imaging of phosphatase activity in
the *Escherichia coli* chemotaxis system. *Proc. Natl. Acad. Sci. U. S. A.* **101**,
17072-17077 (2004).

47   Dasgupta, N. *et al.* A four-tiered transcriptional regulatory circuit controls
flagellar biogenesis in Pseudomonas aeruginosa. *Mol. Microbiol.* **50**, 809-824
(2003).

48   Kulasekara, B. R. *et al.* c-di-GMP heterogeneity is generated by the
chemotaxis machinery to regulate flagellar motility. *eLife* **2**, e01402 (2013).

49   Kulasekara, B. R. *Insight into a Mechanism Generating Cyclic di-GMP*

Heterogeneity in Pseudomonas aeruginosa PhD thesis, University of Washington, (2013).

50 Zhou, H., Rivas, G. & Minton, A. P. Macromolecular crowding and confinement: biochemical, biophysical, and potential physiological consequences. *Annu. Rev. Biophys.* **37**, 375-397 (2008).

51 Cai, Q., Li, Z., Ouyang, Q., Luo, C. & Gordon, V. D. Singly Flagellated Pseudomonas aeruginosa Chemotaxes Efficiently by Unbiased Motor Regulation. *mBio* **7**, e00013-00016 (2016).

52 Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**, 343-345 (2009).

53 Hmelo, L. R. *et al.* Precision-engineering the Pseudomonas aeruginosa genome with two-step allelic exchange. *Nat. Protoc.* **10**, 1820-1841 (2015).

54 Turner, L., Zhang, R., Darnton, N. C. & Berg, H. C. Visualization of Flagella during bacterial Swarming. *J. Bacteriol.* **192**, 3259-3267 (2010).

Author Contributions

J.Y. and R.Z. designed the work; Z.W., M.T. and S.F. performed the measurements with help from M.C.; Z.W and S.F analyzed the data; Z.W., J.Y. and R.Z. wrote the paper. Z.W., M.T. and S.F. contributed equally to this work.

Conflicts of interest

The authors declare no competing interests.

Acknowledgments

This work was supported by National Natural Science Foundation of China Grants (11925406, 12090053, 12304241, and 12304251), a Grant from the Ministry of Science and Technology of China (2019YFA0709303), Grants from the Natural Science Foundation of Shandong Province No. ZR2023QA111 and ZR2023QC168, a special Funding for High-level talents in the Medical and Health Industry in Jinan, a Grant from the Tai Shan Young Scholar Foundation of Shandong Province (tsqnz20231257) and the Science and Technology Development Program of Jinan Municipal Health Commission (2023-1-3).

664 **Table 1.** Strains and plasmids used in this study.

Strain, Plasmid	Genotype, phenotype, and Description	Source
Strains		
P. aeruginosa		
PAO1	wild-type strain	Fan Jin Group
PAO1 *fliC*^{T394C}	Replacement of chromosomal *fliC* in PAO1	Ref 31
PAO1 *fliC*^{T394C} *cheY*-*eyfp*	*yfp* fusions at the N-terminus of *cheY* in PAO1 *fliC*^{T394C}	This work
PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*flhF*	Nonpolar *flhF* deletion in PAO1 *fliC*^{T394C} *cheY*-*eyfp*	This work
PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*fliF*	Nonpolar *fliF* deletion in PAO1 *fliC*^{T394C} *cheY*-*eyfp*	This work
PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*fliG*	Nonpolar *fliG* deletion in PAO1 *fliC*^{T394C} *cheY*-*eyfp*	This work
PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*flhF* Δ*fliF*	Nonpolar *fliF* deletion in PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*flhF*	This work
PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*motAB* Δ*motCD*	Nonpolar *motAB* and *motCD* deletion in PAO1 *fliC*^{T394C} *cheY*-*eyfp*	This work
PAO1 *fliC*^{T394C} Δ*cheA*	Nonpolar *cheA* deletion in PAO1 *fliC*^{T394C}	This work
PAO1 *fliC*^{T394C} *cheY*-*eyfp* Δ*flgI*	Nonpolar *flgI* deletion in PAO1 *fliC*^{T394C} *cheY*-*eyfp*	This work
E. coli		
Top10	F-*mcrA* Δ(*mrr-hsRMS-mcrBC*)Φ80*lacZ* ΔM15 Δ*lacX74 recA1* *ara*D139 Δ(*araleu*)7697 *galU galK rpsL*(Nal^R) *endA1 nupG*	Invitrogen
Plasmids		
pex18gm	oriT⁻sacB⁻; gene replacement vector with MCS from pUC18; Gm^r	Fan Jin Group
flhF-pex18gm	In-frame deletion of *flhF* cloned into pex18gm; Gm^r	This work
fliF-pex18gm	In-frame deletion of *fliF* cloned into pex18gm; Gm^r	This work
fliG-pex18gm	In-frame deletion of *fliG* cloned into pex18gm; Gm^r	This work
motA-pex18gm	In-frame deletion of *motA* cloned into pex18gm; Gm^r	This work
motCD-pex18gm	In-frame deletion of *motCD* cloned into pex18gm; Gm^r	Ref 30
cheA-pex18gm	In-frame deletion of *cheA* cloned into pex18gm; Gm^r	This work
flgI-pex18gm	In-frame deletion of *flgI* cloned into pex18gm; Gm^r	This work
cheY-*eyfp*-pex18gm	*eyfp* fusions *cheY* cloned into pex18gm; Gm^r	This work
cheY-pJN105	*cheY* overexpression vector in pJN105, *cheY* expression is controlled by P$_{BAD}$ promoter; Gm^r	This work
cheY-pME6032	*cheY* overexpression vector in pME6032, *cheY* expression is controlled by P$_{lac}$ promoter; Tet^r	This work
yhjH-pME6032	*yhjH* overexpression vector in pME6032, *yhjH* expression is controlled by P$_{lac}$ promoter; Tet^r	This work
pCdrA-*gfp*	pUCP22-NotI based cyclic di-GMP level reporter, Gm^r	Ref 45
cheY-*eyfp*-pJN105	*eyfp* fusions *cheY* cloned into pJN105, *cheY*-*eyfp* expression is controlled by P$_{BAD}$ promoter; Gm^r	This work
cheA-*ecfp*-pJN105	*ecfp* fusions *cheA* cloned into pJN105, *cheA*-*ecfp* expression is controlled by P$_{BAD}$ promoter; Gm^r	This work

665
666

Fig. 1. **A**. Schematic diagram of the chemotaxis signal transduction network and flagellar motor of *P. aeruginosa*. Fusion protein CheY-EYFP and fluorescently labeled flagellar filaments were used as markers to indicate the position of the chemotaxis complex and motor, respectively. **B.** The labeling mechanism of flagellar filaments and chemotaxis regulatory protein CheY. Filaments (with cysteine point mutation FliC^{T394C}) were labeled through sulfhydryl-maleimide conjugation, and *cheY-eyfp* fusion with a 3× glycine linker was used to visualize chemotaxis complex positions. **C.** Localization of CheY-EYFP in the wild-type strain of *P. aeruginosa*. CheY-EYFP is mainly located at the single cell pole, and the white arrow points to individuals with obvious chemotaxis complex at both cell poles, which generally have a large aspect ratio of the cell body. The yellow dashed box marks the cell outline. **D.** The merged imaging of flagellar filaments and CheY-EYFP in the wild-type strain of *P. aeruginosa*, where flagellar motor and chemotaxis complex colocalize in cells. 145 cells with labeled flagella were observed, all of which exhibited consistent

682 colocalization. White arrows point to individuals about to be divided, and the yellow

683 dashed box marks the cell outline. The scale bar is 1 μm.

Fig. 2. **A**. Localization of CheY-EYFP in the Δ*flhF* strain of *P. aeruginosa*. CheY-EYFP is no longer robustly distributed at the single cell pole. The yellow dashed box marks the cell outline. **B.** The merged imaging of flagellar filaments and CheY-EYFP in the Δ*flhF* strain of *P. aeruginosa*, flagellar motor and chemotaxis complex still colocalize in cells. 101 cells with labeled flagella were observed, all of which exhibited consistent colocalization. The yellow dashed box marks the cell outline. The scale bar is 1 μm.

Fig. 3. **A**. Localization of CheY-EYFP in various *P. aeruginosa* strains. The scale bar

is 10 μm. **B.** Western blot analysis was performed to detect CheY expression in

various *P. aeruginosa* strains. β-actin was used as the housekeeping protein. **C.**

Occurrence probability of obvious chemotaxis complex in *P. aeruginosa* wild-type

and several mutant strains. The proportion of individuals with obvious chemotaxis

complex decreased significantly in the motor-incomplete strains (Δ*fliF* and Δ*fliG*),

and this value was further reduced after *flhF* knockout (Δ*fliF*Δ*flhF*). The Δ*fliF* and

Δ*motA*Δ*motCD* strains have a similar chemotaxis complex occurrence probability as

the wild-type strain. The number of cells analyzed for each strain (from left to right)

were: 372, 221, 234, 323, 672, and 242. '***': significant difference (P-value <

0.0001), 'ns.': no significant difference (P-value > 0.05).

Fig. 4. **A**. The evolution of cell aggregation as the intracellular CheY concentration
increases by induction with higher concentrations of arabinose. The scale bar is 10 μm.
B. Quantitative characterization of intracellular c-di-GMP levels at different CheY
concentrations. From top to bottom, they correspond to the Δ*cheY* strain (N=198),
wildtype strain (N=249) and CheY overexpression strain (N=228), respectively.

715 **Fig. 5**. **A**. The construction mode of the chemotaxis network and flagellar motor

716 throughout the complete cell growth cycle. **B**. The proximal growth mode of the *P.*

717 *aeruginosa* flagellar motor and receptor clusters will effectively regulate the spatial

718 range of CheY action, thus avoiding unintentional cross-pathway regulation.

Fig. S1. Distribution of chemotaxis complexes in multiple strains within representative large fields. Red dotted circles indicate cells where the chemotaxis complex is located at the mid-cell position.

727 **Fig. S2**. Co-localization of CheY-EYFP and CheA-ECFP in *P. aeruginosa*.

728 CheA-ECFP is shown in blue (left), CheY-EYFP in yellow (center), and the merged

729 image is shown in the right panel.

Fig. S3. Simultaneous observation of the chemotaxis complex and flagellar filaments
in the wild-type strain, shown in a representative large field.

wild-type (N=367)

- precise-polar
- near-polar

Δ*flhF* (N=703)

- precise-polar
- near-polar
- mid-cell

Δ*flhF*Δ*fliF* (N=298)

- precise-polar
- near-polar

Fig. S4. Distribution statistics of the chemotaxis complex in the wild-type strain and

the *flhF* mutant. The distribution patterns are categorized into three types:

precise-polar, near-polar, and mid-cell localization. In the wild-type strain, 98.1% of

cells exhibit precise-polar localization of the chemotaxis complex, with the remainder

showing near-polar distribution. In the *flhF* mutant, the proportion of cells with

precise-polar localization decreases to 85.5%, and mid-cell localization appears in

approximately 5% of cells. The *flhF-fliF* double mutant displays a distribution pattern

similar to that of the wild-type strain.

Fig. S5. Knockout of *cheA* did not affect lagellar assembly efficiency of *P.*

aeruginosa.

Fig. S6. Localization of CheY-EYFP in the Δ*flgI* strain of *P. aeruginosa*. The
phenotype is similar to that of the wild-type strain.

753 **Fig. S7**. Intracellular CheY levels induced by different arabinose concentrations,

754 measured by western blot analysis, showing clear differences across the concentration

755 gradient. β-actin was used as the housekeeping protein.

758 **Fig. S8**. Co-overexpressing CheY and the phosphodiesterase (PDE) YhjH from *E. coli*

759 mitigates cell aggregation caused by CheY overexpression in *P. aeruginosa*.

760