Title: A regulatory B cell subpopulation expressing CD301b lectin promotes breast cancer

growth

Authors: Amy V. Paschall[1]†, Zahra Nawaz[1]†, and Fikri Y Avci[1*]

Affiliations: [1]Department of Biochemistry, Emory Vaccine Center, Winship Cancer Institute,

Emory University School of Medicine, Atlanta, GA 30322

†These authors contributed equally.

*Corresponding author. Email: favci@emory.edu

Abstract

Tumor-associated carbohydrate antigens (TACAs) can promote tumor growth by regulating the

anti-tumor immune response. The accumulation of immune suppressor cells within the tumor in

response to TACAs suggests a critical pathway to suppress immune targeting of the tumor.

Employing murine breast cancer models, we isolated a regulatory B cell subpopulation in the breast

tumor microenvironment that displays an immune suppressive phenotype through its Tn TACA-

binding lectin, CD301b. We then demonstrated that depleting CD301b[+] cells facilitated tumor

control and mouse survival, whereas increasing Tn antigen expression decreased mouse survival.

As tumor cells use Tn expression to overcome immune targeting, interfering with or blocking the

Tn-CD301 axis may unleash the immune system, specifically within the aberrantly glycosylated

tumor microenvironment, offering new immunotherapy for breast and other cancers.

20
21

MAIN TEXT

Introduction

A major setback observed in many cancers arises from the modulation of the immune system to prevent effective anti-cancer responses. Vaccines, CAR T cells, and checkpoint blockade prime the immune system against cancer-associated antigens, promoting cancer cell destruction via immune cytotoxicity (*1*). Breast cancer is a prime example of a "cold tumor" that doesn't trigger a robust immune response (*2*). Immune therapies for breast cancer remain largely ineffective due to the ability of the tumor to grow unimpeded and eventually metastasize. Immune regulation can occur through cytokine secretion (such as IL-10) (*3, 4*), signaling mechanisms, or checkpoints (such as through PD-1) (*5*). During the malignant transformation of a mammalian cell, a dramatic and aberrant modification of cellular glycosylation is observed. Tumor-associated carbohydrate antigens, or TACAs, can induce immune suppression, allowing cancer cells to evade immune cells (*6-8*). Lectins, such as galectins, are carbohydrate-binding proteins that function as immune cell receptors and can activate immune regulatory pathways through their interactions with TACAs (*9, 10*). Similarly, modulating immune profiles in the tumor microenvironment (TME) by engaging sialic-acid-binding immunoglobulin-like lectins (Siglecs) is a known TACA mechanism (*11-13*). Thus, elucidating the protein-carbohydrate interactions in regulating the immune system within the TME is becoming increasingly important.

Tn antigen is a TACA ranked as a high-priority cancer-associated antigen based on its antigenicity and oncogenicity (*14-18*). Tn is a truncated form of the normal, extended cell surface O-glycan. Tn is the terminal O-linked N-acetylgalactosamine (GalNAc) on serine or threonine residues. The aberrant glycosylation associated with Tn may occur when the enzyme responsible for O-glycan elongation—T-synthase—or its associated chaperone—cosmc (C1GALT1C1)—becomes functionally inhibited (*19, 20*). Tn-expressing mucin 1 (MUC1) has been associated with breast

cancer cells (*21, 22*), especially in triple-negative breast cancer (*23, 24*) , and modulating MUC1

expression or Tn glycosylation can inhibit tumor growth (*25-27*). However, developing immune

responses against Tn-MUC1 has been problematic (*27, 28*).

CD301, also known as macrophage galactose-type lectin, or MGL, is a C-type lectin receptor (CLR)

that binds Tn antigen on the surface proteins (*29*). In mice, the Tn-recognizing homolog to the

human CD301 is CD301b, or MGL2 (*30*). CD301b has been considered to be expressed exclusively

by myeloid cells such as dendritic cells (DCs) and macrophages (*17*). Previously, MGL2-expressing

myeloid cell populations have been linked with immunosuppressive responses (*16, 17, 31-33*). DCs

and macrophages were shown to suppress the proliferation of CD4$^+$ effector T lymphocytes through

the interaction of MGL with terminal GalNAc residues on CD45 expressed by T cells (*17*). This

cell-specific glycosylation of CD45 provided an immunoregulatory pathway of MGL-mediated

control of T$_{eff}$ cell function. In another study, CD301b$^+$ dendritic cells were shown to suppress T

follicular helper cells and antibody responses to protein antigens (*31*). Recently, an

immunosuppressive DC subset expressing CD301b was shown to accumulate at secondary sites

and promote metastasis in pancreatic cancer (*33*) and lung cancer (*34*).

Here, we demonstrate the accumulation of CD301b$^+$ immune cells in the tumor microenvironment

as a response to Tn-expression by breast cancer cells. We characterize these cells into populations

of cells expressing markers associated with regulatory innate-like B cells and produce suppressive

cytokines. As a subset of B lymphocytes, regulatory B cells (Bregs) exhibit immunosuppressive

functions in homeostasis (*35*) and numerous disease states, including colitis (*36*), arthritis (*37*), HIV

(*38, 39*), and multiple cancers (*40-43*). Thus, these cells are broadly relevant in multiple tissues,

systems, and microenvironments throughout the body (*35*). Breg cells are typically characterized

by their production of IL-10 (*37, 44*), a cytokine most frequently implicated in immune suppression.

Inhibiting IL-10-producing Bregs has been shown to disrupt normal anti-inflammatory processes and can result in chronic inflammation (*45*). Bregs are often identified by key cell surface markers, including CD1d(*46*), CD24 (*47*), and/or CD5 (*46*). A heterogeneous subpopulation of B cells expressing innate cell markers such as CD11b and demonstrating immunoregulatory functions has been identified (*37, 45, 48-54*). Of note, a subpopulation of innate-like B cells may produce high levels of IgM as one mechanism for decreasing inflammatory responses (*55*).

This work offers a method of immune suppression similar to checkpoint regulation, which has represented critical targets in the rapidly advancing field of cancer immunotherapy. Immune interactions of TACAs have been severely understudied. Elucidating immune suppression mechanisms through protein-carbohydrate interactions has important implications for discovering new immune therapies to boost immune anti-cancer function through overcoming regulation mechanisms.

Results

CD301b$^+$ cells accumulate in a murine breast cancer model

We first sought to establish the relationship between CD301b$^+$ immune cells and Tn glycoantigen-expressing breast cancer cells. We used a CRISPR-Cas9 gene editing model to knockout cosmc chaperone expression in AT3 murine breast cancer cells. Cosmc is a chaperone essential in elongating the core O-glycan beyond the truncated Tn form of α-GalNAc (*19, 20, 56*). When cosmc function/expression is disrupted, elongation of the O-glycan is not observed; instead, the Tn antigen is observed at significantly higher levels. After disrupting cosmc expression in these cell lines, we confirmed increased Tn cell surface expression through flow cytometry using both a reBaGs6 IgM antibody (Supplementary Fig. 1A) (*57*) as well as a complementary biotinylated VVL lectin coupled with fluorescent streptavidin (Supplementary Fig. 1B). Both staining methods indicated

significantly higher Tn expression on the cosmc KO cell line. We also isolated RNA from each cell line and confirmed decreased cosmc expression in the Tn^{hi} cell line through qPCR using cosmc primers (Supplementary Fig. 1C). We then tested whether knocking out cosmc expression changes the proliferation rate of the AT3 breast cancer cells *in vitro*. After culturing both cell lines at the same concentrations for three days, we observed no significant differences in cell proliferation between the two lines (Supplementary Fig. 1D), indicating that Tn expression alone does not promote cancer cell growth.

To determine the *in vivo* tumor cell growth, we injected AT3 (Tn^{lo}) and AT3 cosmc KO (Tn^{hi}) murine breast tumor cells into the mammary pads of C57BL/6 mice (n=25) and monitored tumor growth for up to 23 days. We observed that Tn^{hi} tumors grew significantly faster than Tn^{lo} tumors (Fig. 1A), indicating that Tn expression impacts tumor growth rate. Tumor-bearing and tumor-free control mice were then euthanized to assess CD301b on multiple immune cell subsets in the spleen, draining lymph nodes (or analogous lymph nodes in tumor-free mice), bone marrow, or breast tumors by flow cytometry (for gating strategy, please see Supplementary Fig. 2). A significantly higher accumulation of $CD45^+CD301b^+$ immune cells was observed in the Tn^{hi} tumor microenvironments compared to the Tn^{lo} tumor tissue (Fig. 1B-C). Low levels of cell surface CD301b were observed on immune cell subsets in the spleens or lymph nodes of tumor-free mice (Fig. 1C). However, a significant population of $CD45^+CD301b^+$ cells was observed in the tumor tissues and the bone marrow of the tumor-bearing mice. $CD301b^+$ immune cell accumulation in the TME correlates with tumor Tn expression.

Tumor-infiltrating $CD301b^+$ cells display a regulatory B-cell phenotype

We next observed that essentially all tumor-infiltrating $CD45^+CD301b^+$ immune cells express the B-cell markers B220 and CD19, and they do not express markers associated with other major immune cell types CD3, NK1.1 and CD11c (Fig. 2A-D). CD11b expression on the tumor-

infiltrating CD45$^+$CD301b$^+$ varied (Fig. 2B, 2D). Since CD301b function has primarily been

attributed to myeloid cells, and particularly to DCs, we assessed and observed that the tumor-

infiltrating CD45+CD301b+B220+CD19+ immune cells do not express CD11c (Fig. 2E). The

absolute number of B220+CD19+ cells per mg of tumor tissue is highly increased in Tn^{hi} tumor

tissues compared to Tn^{lo} tumors, with approximately a third of them expressing CD301b+ in the

Tn^{hi} tumors (Fig. 2F).

The tumor-infiltrating B220$^+$CD19$^+$CD301b$^+$ cells express cell surface IgM, a B cell-specific

surface receptor (Fig. 3A). Next, we observed the CD301b$^+$ B cells express surface proteins

associated with regulatory B cells, such as CD5, CD24, Gr1 with varying levels, while other surface

markers traditionally associated with Bregs (CD1d, CD25, CD21, Ly6C and Ly6G) were minimally

expressed on these B220$^+$CD19$^+$CD301b$^+$ B cells (Fig. 3B-F). The tumor-infiltrating CD301b$^+$ cells

may represent a previously uncharacterized subset of B cells with a phenotype indicating an innate-

like regulatory B cell population (*48-55*).

Tn-Expressing Cancer Cells Induces Expression of CD301b on B cells

CD301b mean fluorescence intensity (MFI) was significantly higher in Tn^{hi} tumor-infiltrating cells

than Tn^{lo} (Fig. 3G), indicating that CD301b expression is tumor Tn-associated. To further elucidate

if B cells can express CD301b in response to Tn expression, we obtained splenocytes from healthy

WT C57Bl/6 mice and used magnetic cell sorting protocols to positively select for CD19$^+$ cells or

isolate pan B cell populations through negative selection, compared to unselected total splenic

mononuclear cells (Supplementary Fig. 3). These splenocyte populations were then co-cultured

with either AT3 Tn^{lo} or AT3 Tn^{hi} cancer cells, compared to cultured alone (unstimulated). After

three days of co-culture, cells were analyzed using flow cytometry: viable singlet

CD45$^+$B220$^+$CD19$^+$ cells were gated out, and CD301b MFI was assessed in each group. We

observed significantly higher CD301b expression in all splenocyte groups co-cultured with Tn^{hi}

cancer cells compared with unstimulated or Tn^{lo} cancer cell co-cultures (Fig. 3H). These results

indicate an important role for Tn in the expression of CD301b on B cell populations.

B Cell Deficient Mice did not Exhibit CD301b+ Cell Accumulation in Response to Tn^{hi} Tumor

Towards elucidating the role of CD301b+ B cells in the context of Tn-expressing tumor growth,

we used RAG1 KO (B6.129S7-Rag1tm1Mom/J) mice that lack functional T or B cells. We injected

Tn^{hi} tumor cells into the mammary pads of RAG1 KO mice alongside a WT control group. We

observed significantly decreased tumor growth (Fig. 4A) and no observable CD301b+ immune cells

infiltrating the tumors (Fig. 4B-C) compared to WT mice. No observable B220+CD19+ cells were

present within the RAG1 KO mouse tumors (Fig. 4D). These results further demonstrate that the

tumor-infiltrating CD301b+B220+CD19+IgM+ immune cells constitute a B cell population as they

are absent when mice lack the lymphocyte-specific *rag1* gene. Our results also suggest that

CD301b+ B cells play a regulatory function in promoting tumor growth. However, along with the

CD301b+ B cells, all lymphocytes, including T cells and CD301b- B cells, are absent in RAG1 KO

mice. Therefore, attributing such an immunoregulatory role to tumor-infiltrating CD301b+ B cells

requires elucidating their functional properties specifically, as laid out in the next two sections.

Tumor-infiltrating CD301b$^+$ B cells exhibit immunosuppressive characteristics

To better understand the functional suppressive characteristics of tumor-infiltrating CD301b+ B

cells, we used intracellular staining of Tn^{hi} breast tumor-bearing mouse tumor tissue to stain for

regulatory cytokines. We gated and compared the following populations: CD45$^+$B220$^+$CD19$^+$ B

cells in tumor-free spleens and CD45$^+$CD301b$^+$B220$^+$CD19$^+$ from Tn^{hi} tumors. Of note, we

observed that tumor-associated CD19$^+$B220$^+$CD301b$^+$ produce significantly higher levels of IL10,

TGFβ, and subunits of IL35 than splenic-derived B cells from tumor-free mice as controls (Fig.

5A-B, Supplementary Fig. 4), suggesting a role for immune suppression in this tumor-infiltrating B cell subset.

CD301b/Tn Axis Impacts Tumor Growth and Mouse Survival

To illuminate how CD301b immune cells contribute to tumor growth, we employed a mouse model in which CD301b immune cells can be depleted: B6(FVB)-*Mgl2*$^{tm1.1(HBEGF/EGFP)Aiwsk}$/J mice (MGL2-DTR)([31, 32]). We injected AT3 Tn^{lo} or Tn^{hi} tumor cells (Fig. 6A) into MGL2-DTR mice that were/ were not CD301b-depleted with diphtheria toxin treatment. We measured tumor growth and observed a significant decrease in tumor size when CD301b$^+$ immune cells were depleted in the Tn^{hi} group. We also injected AT3 Tn^{hi} cells into mammary pads of CD301b null MGL2-DTR homozygous mice and observed significantly lower breast tumor growth than WT mice (Fig. 6B).

In a complementary experiment, we used a mouse model expressing human MUC1: C57BL/6-Tg(MUC1)79.24Gend/J ([58]). These mice display aberrant glycosylation on MUC1, such as Tn antigen, unlike wild-type mouse MUC1 glycosylation ([59-62]). We used methylcholanthrene (MCA) as a carcinogen to induce spontaneous tumors in the mammary pads of both WT and Muc1.Tg mice to assess how Tn expression affects tumor growth. We observed significantly increased tumor growth rates to endpoint tumor sizes over 21 weeks in the Muc1.Tg mice (Fig. 6C), indicating that higher Tn expression on spontaneous tumors—not just cancer cell lines—can impact breast cancer prognosis. To our point, tumor-infiltrating B cells from MUC1.Tg spontaneous tumors showed significantly higher percentages of CD301b$^+$ cells than the corresponding B cell populations from WT MCA-induced tumors (Fig. 6D). These data together indicate an important role for the CD301b-Tn axis in breast tumor development and the high therapeutic potential for modulating it to improve the anti-tumor immune response in breast cancer. However, other factors, such as differences in tumorigenicity, cell-intrinsic tumor growth, or metastatic activity, can potentially contribute to the difference in survival in conjunction with or

200   independent from the interaction of MUC1-Tn with CD301b+ cells. Modulating CD301b and Tn

201   expression in this mouse model would enable further evaluation of the CD301b-Tn axis in tumor

202   immunoregulation.

Discussion

Numerous recent research studies have established the importance of Tn as a key antigen in relation

to human cancer development and progression (*20, 21, 63-65*). Additionally, CD301 plays an

immune regulatory role in human infection (*31, 32*) and the progression of human cancer (*33*). In a

recent study, CD301b[+] DCs were shown to accumulate in response to higher levels of Tn in lung

tumor cells, and the Tn-CD301b interactions influence the development of pro-angiogenic

programs (*34*). The link between Tn and CD301-mediated immune responses thus needs further

elucidation, particularly in cancer models known to express aberrant glycosylation, such as breast

cancer. Overcoming immune regulation in breast cancer can potentially turn these "cold" tumors

into immunotherapeutic-responsive "hot" tumors, potentially improving clinical responses to

checkpoint blockade and tumor antigen vaccines. This study shows an accumulation of CD301b[+]

immune cells in response to high Tn antigen expression by developing breast tumors in mice. These

CD301b[+] immune cells phenotypically and functionally display an innate-like B cell phenotype

with regulatory functions, including producing immunoregulatory cytokines. Innate-like B cells

may express CD11b and CD5 but low levels of CD11c (*53, 66*). Innate-like B cells are also

associated with high levels of IgM and low levels of IgD (*55*) These cells have been shown to

produce natural IgMs in response to tumor-associated gangliosides (*67*). Of note, extensive research

studies have shown that B cells with regulatory function cannot currently be defined based on any

specific combination of extracellular markers or transcription factors; rather, the heterogeneity of

regulatory B cells suggests that these cells arise in response to environmental stimuli (*35*). We

observe suppressive CD301b[+] cells accumulating in the tumor microenvironment specifically, and

particularly in response to higher Tn expression by the developing breast tumor. However, as some

accumulation is observed in bone marrow in tumor-bearing mice, studies investigating the origin of these cell populations would be warranted. MGL2/CD301b function has primarily and predominantly been attributed to myeloid cells in the past decades. Our results indicating that a previously uncharacterized regulatory CD301b+ lymphoid cell population plays an important role in the tumor immune escape warrants further exploration of the function and regulation of *mgl2* gene expression. Particularly, the results obtained in the RAG1 KO mouse model would suggest that CD301b+ B cells promote tumor growth. However, the direct role of B cell-specific CD301b and/or CD301b+ B cells could be further assessed by modulating CD301b expressing cells and B cells independently, such as developing a muMt-/Mgl2-DTR chimeric mouse model.

The MUC1.Tg mouse model is limited because it may not account for CD301b-Tn interaction alone and likely exhibits distinct tumor microenvironment development. Understanding the innate-like characteristics of these cells could have implications for their mechanisms of suppressive functions in the context of Tn^{hi} tumors. Analysis (e.g., via transcriptomics) of these tumor-infiltrating cell populations in relation to the tumor immune microenvironment could thus describe the novel population of suppressive cells and their mechanisms of action. Eliminating these cells can improve tumor response in mice, which has important implications for targeting the CD301b-Tn axis to improve immunotherapy. Further characterization of how these cells modulate immune responses in the TME will help determine optimal therapies to target and overcome their accumulation and immunosuppressive mechanisms. Furthermore, therapies targeting CD301 and/or Tn could improve breast cancer response to breakthrough therapies, such as PD-1 or CTLA-4 checkpoint blockade. Future investigation of these therapeutic strategies will thus be a priority in this research. Understanding how these $B220^{+}CD19^{+}IgM^{+}CD301b^{+}$ immune cells modulate the immune pathways in TME and impact disease progression could have important implications for cancer immunotherapy.

Materials and Methods

Mice

Eight-week-old female C57BL/6 mice were obtained from Jackson Laboratories (Bar Harbor, ME) and housed at Emory University Whitehead Biomedical Research Building. MGL2-DTR mutant mice (*Mgl2*$^{tm1.1(HBEGF/EGFP)Aiwsk}$) were a generous gift from Akiko Iwasaki at Yale University. Muc1.Tg mice (C57BL/6-Tg(MUC1)79.24Gend/J) were a generous gift from Sandra Gendler at Mayo Clinic. Mice were kept in microisolator cages and handled under biosafety level 2 (BSL2) hoods. For tissue processing and subsequent flow cytometry, mice were euthanized by carbon dioxide inhalation in accordance with IACUC guidelines. For applicable experiments, serum samples, spleens, lymph nodes, bone marrow, and/or tumors were harvested. Where applicable, cell suspensions were generated through mechanical tissue disruption and collagenase D digestion. Red blood cells were lysed, and samples were filtered through 60 μm nylon filters to obtain single-cell suspensions. For depletion of CD301b$^+$ cells, heterozygous MGL2-DTR mice were treated with diphtheria toxin (0.5μg/mouse) in sterile PBS intraperitoneally every two to three days, starting at day -1 before tumor injection. Effective MGL2 cell depletion was confirmed using flow cytometry in bone marrow and tumor samples (Supplementary Fig. 5A-B). For RAG1 KO experiments, 6-8 weeks old RAG1 KO mice (B6.129S7-Rag1tm1Mom/J) were obtained from Jackson Laboratory. The lack of B cell populations was confirmed using flow cytometry in bone marrow and lymph node samples (Supplementary Fig. 5C).

All mouse experiments were in compliance with the Emory University Institutional Animal Care and Use Committee under an approved animal use protocol. Our animal use protocol adheres to the principles outlined in *U.S. Government Principles for the Utilization and Care of Vertebrate Animals Used in Testing*, *Research and Training*, the Animal Welfare Act, the *Guide for the Care and Use of Laboratory Animals*, and the *AVMA Guidelines for the Euthanasia of Animals*.

Generation of Tn^{hi} breast cancer cells

To express Tn glycans at high levels in tumor cells, we used a CRISPR/Cas9 methodology to stably

silence cosmc expression in AT3 cells using established protocols and reagents. Mouse cosmc guide

RNA and CRISPR/Cas9 plasmid were obtained from Santa Cruz Technology (sc-425587). AT3

murine breast cancer cells were a generous gift from the Kebin Liu lab at Augusta University. AT3

cells were transfected with cosmc CRISPR/Cas9 KO plasmid according to the manufacturer's

protocol. Puromycin was used to select transfected cells. We then used flow cytometry to confirm

higher expression of Tn on the AT3 cell surfaces using the ReBaG6 antibody (generously provided

by Richard Cummings at Harvard University) (Supplementary Fig. 1A) (*57*) and VVL lectin

(Vector Laboratories) (Supplementary Fig. 1B). The proliferation of each transfected cell line (AT3

and AT3 Tn^{hi}) was tested in an MTT proliferation assay for three days, and colorimetric analysis

was performed with a CytoTek plate reader according to protocol; no significant differences in

proliferation compared to parent cells were observed (Supplementary Fig. 1D). Cell lines were

maintained in RPMI media supplemented with 10% FBS, sodium pyruvate, HEPES buffer, NEAA,

β-mercaptoethanol, and penicillin/streptomycin at 37°C, 5% CO_2.

AT3 Tn^{lo} and/or AT3 Tn^{hi} Tumor Challenge

Breast tumor cells were harvested and washed in sterile PBS. Cells were suspended in a final

concentration of 2.5E6/ml sterile PBS. Cells were subcutaneously injected into the mammary pads

of mice at 2.5E5/100μl/mouse. Mice were observed and euthanized at the tumor endpoint

(maximum tumor dimension between 0.9 cm and 1.2 cm, typically after Day 17). For instance, Tn^{lo}

tumor-bearing mice analyzed for Fig. 2 were sacrificed on Day 23, and Tn^{hi} tumor-bearing mice

were sacrificed on Day 17. RAG1 KO mice in Fig. 4 were sacrificed on Day 23. In cases where

tumors were observed to ulcerate or reach maximum growth sooner than the earliest endpoint

(before Day 17), the mice were euthanized immediately, and samples were excluded. Tumor

volumes were calculated as ((length x width x width)/2) in mm^3.

Flow Cytometry

Cells were stained in PBS with TruStain fcX (BioLegend, Cat. No. 101320) to reduce non-specific

antibody binding. Cell samples were stained with the following antibodies and stains (in multiple

sets to prevent fluorophore overlap): B220-FITC (BioLegend, clone RA3-6B2), CD11c-PacBlue

(BioLegend, clone N418), CD11b-FITC (BioLegend, clone M1/70), CD3e-APC (BioLegend, 145-

2C11), CD45-APCCy7 (BioLegend, clone 30-F11), CD19-PE (BioLegend, clone 6D5), NK1.1-

PerCP (BioLegend, PK136), Gr1-PECy5 (BioLegend), CD19-APC (BioLegend, clone 1D3),

Ly6C-BV605 (BioLegend, clone HK1.4), CD11b-PacBlue (BioLegend M1/70), LycG-BV605

(BioLegend clone RTK2758), CD1d-PacBlue (BioLegend, clone 1B1), CD5-PECy5 (BioLegend,

clone 53-7.3), CD24-PECy7 (BioLegend, clone 30-F1), CD301b-APC (BioLegend, clone URA-1),

CD301b-PE (BioLegend, clone URA-1), CD25-BV605 (BioLegend, clone PC61), CD21-BV510

(BioLegend), IL10 PECy7 (BioLegend, clone JES5-16E3), TGFb-BV421 (BioLegend, clone TW7-

16B4), IgM-PECy5 (BioLegend, clone RMM-1), IgD-PECy7 (BioLegend, clone 11-26c.2a)

IL35:EBI3 subunit – PerCP (R&D, clone 355022), IL35:p35 subunit – AF594 (R&D, clone 27537),

and Ghost Red 710 (Tonbo Biosciences). For intracellular staining, cell samples were treated with

GolgiStop for four hours prior to the staining protocol. Intracellular staining was performed

according to the protocol (BD Biosciences). All isotype controls were obtained from BioLegend

except isotypes for IL35 subunits, which were obtained from R&D. Samples were washed and

analyzed with flow cytometry (Cytek Aurora or Agilent Novocyte Penteon). Fluorescence minus

one (FMO) plus specific isotype control antibody-stained samples were used as negative staining

controls, and single stains were used for compensation. Flow cytometry data were analyzed using

FlowJo Single Cell Analysis Software (Treestar, Inc., Ashland, Oregon) with gating strategies

shown in Supplementary Fig. 2 (example showing tumor tissue from Tn[hi]-tumor bearing mice).

Splenocyte B Cell Selection and Co-Culture

Splenocytes were obtained from healthy WT C57Bl/6 mice. BioLegend MojoSort antibody kits

were used to positively select CD19 cells or negatively isolate Pan B cells, according to

manufacturer protocols. Flow cytometry was used to determine the B cell purity of each sample

(Supplementary Fig. 3). Viable cells were counted and cultured in 96-well plates at 1E5 cells/100 µl

complete RPMI (supplemented with Ca+) /well. Tumor cells were harvested, and viable cells were

counted. Indicated cell populations were then plated in each splenocyte sample at 1E5 cells/100 µl

complete RPMI (for the "Unstimulated" sample, 100 µl complete RPMI was used without tumor

cells), for a final volume of 200 µl /well. Plates were cultured for 3 days at 37°C, 5% CO_2. Samples

were washed and prepared for flow cytometry analysis.

Statistical Analysis

GraphPad Prism v8 was used for statistical analyses. Two-way ANOVA with Tukey's multiple

comparisons test was used to determine statistical significance between experimental groups in

each of the applicable experimental models (Fig. 1C, Fig. 2F, Fig. 4D, Fig. 6A). Ordinary one-way

ANOVA with Tukey's multiple comparisons test was used to analyze Fig. 5. Mouse survival in

Fig. 6C was compared using Log-rank (Mantel-Cox) test. An unpaired parametric two-tailed t-test

was used for Figs. 1A, 3G, 4A, 4C, 6B, 6D. Significance is indicated on each graph based on p-

value: >0.05 = ns; $<.05$ = *; <0.01 = **; <0.001 = ***; <0.0001 = ****.

References

1. Y. Zhang, Z. Zhang, The history and advances in cancer immunotherapy: understanding the
characteristics of tumor-infiltrating immune cells and their therapeutic implications. *Cell Mol*
Immunol **17**, 807-821 (2020).

2. A. Swoboda, R. Nanda, Immune Checkpoint Blockade for Breast Cancer. *Cancer Treat Res* **173**,
155-165 (2018).

3. D. F. Fiorentino, M. W. Bond, T. R. Mosmann, Two types of mouse T helper cell. IV. Th2 clones
secrete a factor that inhibits cytokine production by Th1 clones. *J Exp Med* **170**, 2081-2095
(1989).

4. S. E. Acuner Ozbabacan, A. Gursoy, R. Nussinov, O. Keskin, The structural pathway of interleukin 1
(IL-1) initiated signaling reveals mechanisms of oncogenic mutations and SNPs in inflammation
and cancer. *PLoS Comput Biol* **10**, e1003470 (2014).

5. G. J. Freeman *et al.*, Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family
member leads to negative regulation of lymphocyte activation. *J Exp Med* **192**, 1027-1034 (2000).

6. N. Rodrigues Mantuano, M. Natoli, A. Zippelius, H. Laubli, Tumor-associated carbohydrates and
immunomodulatory lectins as targets for cancer immunotherapy. *J Immunother Cancer* **8**, (2020).

7. H. Laubli, A. Varki, Sialic acid-binding immunoglobulin-like lectins (Siglecs) detect self-associated
molecular patterns to regulate immune responses. *Cell Mol Life Sci* **77**, 593-605 (2020).

8. S. S. Pinho, C. A. Reis, Glycosylation in cancer: mechanisms and clinical implications. *Nat Rev*
Cancer **15**, 540-555 (2015).

9. K. V. Marino, A. J. Cagnoni, D. O. Croci, G. A. Rabinovich, Targeting galectin-driven regulatory
circuits in cancer and fibrosis. *Nat Rev Drug Discov* **22**, 295-316 (2023).

10. A. J. Cagnoni *et al.*, Galectin-1 fosters an immunosuppressive microenvironment in colorectal
cancer by reprogramming CD8(+) regulatory T cells. *Proc Natl Acad Sci U S A* **118**, (2021).

11. R. Beatson *et al.*, The mucin MUC1 modulates the tumor immunological microenvironment
through engagement of the lectin Siglec-9. *Nat Immunol* **17**, 1273-1281 (2016).

12. N. T. Marcos *et al.*, Role of the human ST6GalNAc-I and ST6GalNAc-II in the synthesis of the
cancer-associated sialyl-Tn antigen. *Cancer Res* **64**, 7050-7057 (2004).

13. M. J. Schultz *et al.*, The Tumor-Associated Glycosyltransferase ST6Gal-I Regulates Stem Cell
Transcription Factors and Confers a Cancer Stem Cell Phenotype. *Cancer Res* **76**, 3978-3988
(2016).

14. A. V. M. Paschall, D. R.; Avci, F. Y, in *Encyclopedia of Cell Biology, Second Edition,* R. A. H.
Bradshaw, G. W.; Stahl, P. D., Ed. (Elsevier, 2023), pp. 404-413.

15. T. Freire, R. Lo-Man, S. Bay, C. Leclerc, Tn glycosylation of the MUC6 protein modulates its
immunogenicity and promotes the induction of Th17-biased T cell responses. *J Biol Chem* **286**,
7797-7811 (2011).

16. E. Saeland *et al.*, The C-type lectin MGL expressed by dendritic cells detects glycan changes on
MUC1 in colon carcinoma. *Cancer Immunol Immunother* **56**, 1225-1236 (2007).

17. S. J. van Vliet, S. I. Gringhuis, T. B. Geijtenbeek, Y. van Kooyk, Regulation of effector T cells by
antigen-presenting cells via interaction of the C-type lectin MGL with CD45. *Nat Immunol* **7**, 1200-
1208 (2006).

18. M. A. Cheever *et al.*, The prioritization of cancer antigens: a national cancer institute pilot project
for the acceleration of translational research. *Clin Cancer Res* **15**, 5323-5337 (2009).

19. Y. Wang *et al.*, Cosmc is an essential chaperone for correct protein O-glycosylation. *Proc Natl*
Acad Sci U S A **107**, 9228-9233 (2010).

20. T. Ju, R. P. Aryal, M. R. Kudelka, Y. Wang, R. D. Cummings, The Cosmc connection to the Tn
antigen in cancer. *Cancer Biomark* **14**, 63-81 (2014).

21. R. Beatson *et al.*, The Breast Cancer-Associated Glycoforms of MUC1, MUC1-Tn and sialyl-Tn, Are
Expressed in COSMC Wild-Type Cells and Bind the C-Type Lectin MGL. *PLoS One* **10**, e0125994
(2015).

22. R. E. Beatson, J. Taylor-Papadimitriou, J. M. Burchell, MUC1 immunotherapy. *Immunotherapy* **2**,
305-327 (2010).

23. A. Siroy *et al.*, MUC1 is expressed at high frequency in early-stage basal-like triple-negative breast
cancer. *Hum Pathol* **44**, 2159-2166 (2013).
24. J. M. Burchell, R. Beatson, R. Graham, J. Taylor-Papadimitriou, V. Tajadura-Ortega, O-linked
mucin-type glycosylation in breast cancer. *Biochem Soc Trans* **46**, 779-788 (2018).
25. D. Chen *et al.*, Immunotherapy of spontaneous mammary carcinoma with fusions of dendritic
cells and mucin 1-positive carcinoma cells. *Immunology* **109**, 300-307 (2003).
26. P. Mukherjee *et al.*, MUC1-specific immune therapy generates a strong anti-tumor response in a
MUC1-tolerant colon cancer model. *Vaccine* **25**, 1607-1618 (2007).
27. V. Lakshminarayanan *et al.*, Immune recognition of tumor-associated mucin MUC1 is achieved by
a fully synthetic aberrantly glycosylated MUC1 tripartite vaccine. *Proc Natl Acad Sci U S A* **109**,
261-266 (2012).
28. L. R. Loureiro *et al.*, Challenges in Antibody Development against Tn and Sialyl-Tn Antigens.
Biomolecules **5**, 1783-1809 (2015).
29. E. Rodriguez *et al.*, Fasciola hepatica Immune Regulates CD11c+ Cells by Interacting with the
Macrophage Gal/GalNAc Lectin. *Front Immunol* **8**, 264 (2017).
30. S. K. Singh *et al.*, Characterization of murine MGL1 and MGL2 C-type lectins: distinct glycan
specificities and tumor binding properties. *Mol Immunol* **46**, 1240-1249 (2009).
31. Y. Kumamoto, T. Hirai, P. W. Wong, D. H. Kaplan, A. Iwasaki, CD301b+ dendritic cells suppress T
follicular helper cells and antibody responses to protein antigens. *Elife* **5**, (2016).
32. Y. Kumamoto *et al.*, CD301b(+) dermal dendritic cells drive T helper 2 cell-mediated immunity.
Immunity **39**, 733-743 (2013).
33. J. A. Kenkel *et al.*, An Immunosuppressive Dendritic Cell Subset Accumulates at Secondary Sites
and Promotes Metastasis in Pancreatic Cancer. *Cancer Res* **77**, 4158-4170 (2017).
34. V. da Costa *et al.*, The Tn antigen promotes lung tumor growth by fostering immunosuppression
and angiogenesis via interaction with Macrophage Galactose-type lectin 2 (MGL2). *Cancer Lett*
518, 72-81 (2021).
35. E. C. Rosser, C. Mauri, Regulatory B cells: origin, phenotype, and function. *Immunity* **42**, 607-612
(2015).
36. A. Mizoguchi, E. Mizoguchi, R. N. Smith, F. I. Preffer, A. K. Bhan, Suppressive role of B cells in
chronic colitis of T cell receptor alpha mutant mice. *J Exp Med* **186**, 1749-1756 (1997).
37. C. Mauri, D. Gray, N. Mushtaq, M. Londei, Prevention of arthritis by interleukin 10-producing B
cells. *J Exp Med* **197**, 489-501 (2003).
38. Y. Jiao *et al.*, Regulatory B cells correlate with HIV disease progression. *Microbiol Immunol* **58**,
449-455 (2014).
39. J. Liu *et al.*, IL-10-producing B cells are induced early in HIV-1 infection and suppress HIV-1-
specific T cell responses. *PLoS One* **9**, e89236 (2014).
40. Y. Zhang, R. Morgan, E. R. Podack, J. Rosenblatt, B cell regulation of anti-tumor immune
response. *Immunol Res* **57**, 115-124 (2013).
41. X. Zhou, Y. X. Su, X. M. Lao, Y. J. Liang, G. Q. Liao, CD19(+)IL-10(+) regulatory B cells affect survival
of tongue squamous cell carcinoma patients and induce resting CD4(+) T cells to CD4(+)Foxp3(+)
regulatory T cells. *Oral Oncol* **53**, 27-35 (2016).
42. M. Shen, J. Wang, X. Ren, New Insights into Tumor-Infiltrating B Lymphocytes in Breast Cancer:
Clinical Impacts and Regulatory Mechanisms. *Front Immunol* **9**, 470 (2018).
43. L. Ye *et al.*, Tumor-derived exosomal HMGB1 fosters hepatocellular carcinoma immune evasion
by promoting TIM-1(+) regulatory B cell expansion. *J Immunother Cancer* **6**, 145 (2018).
44. F. Burke, A. J. Stagg, P. A. Bedford, N. English, S. C. Knight, IL-10-producing B220+CD11c- APC in
mouse spleen. *J Immunol* **173**, 2362-2372 (2004).
45. S. Fillatreau, C. H. Sweenie, M. J. McGeachy, D. Gray, S. M. Anderton, B cells regulate
autoimmunity by provision of IL-10. *Nat Immunol* **3**, 944-950 (2002).
46. K. Yanaba *et al.*, A regulatory B cell subset with a unique CD1dhiCD5+ phenotype controls T cell-
dependent inflammatory responses. *Immunity* **28**, 639-650 (2008).

47.   F. Flores-Borja *et al.*, CD19+CD24hiCD38hi B cells maintain regulatory T cells while limiting TH1
and TH17 differentiation. *Sci Transl Med* **5**, 173ra123 (2013).
48.   X. Liu *et al.*, B cells expressing CD11b effectively inhibit CD4+ T-cell responses and ameliorate
experimental autoimmune hepatitis in mice. *Hepatology* **62**, 1563-1575 (2015).
49.   S. A. Geherin *et al.*, IL-10+ Innate-like B Cells Are Part of the Skin Immune System and Require
alpha4beta1 Integrin To Migrate between the Peritoneum and Inflamed Skin. *J Immunol* **196**,
2514-2525 (2016).
50.   E. E. Ghosn, Y. Yang, J. Tung, L. A. Herzenberg, L. A. Herzenberg, CD11b expression distinguishes
sequential stages of peritoneal B-1 development. *Proc Natl Acad Sci U S A* **105**, 5195-5200 (2008).
51.   J. B. Wong *et al.*, B-1a cells acquire their unique characteristics by bypassing the pre-BCR
selection stage. *Nat Commun* **10**, 4768 (2019).
52.   T. Kobayashi *et al.*, Regulatory B1a Cells Suppress Melanoma Tumor Immunity via IL-10
Production and Inhibiting T Helper Type 1 Cytokine Production in Tumor-Infiltrating CD8(+) T
Cells. *J Invest Dermatol* **139**, 1535-1544 e1531 (2019).
53.   D. O. Griffin, T. L. Rothstein, Human "orchestrator" CD11b(+) B1 cells spontaneously secrete
interleukin-10 and regulate T-cell activity. *Mol Med* **18**, 1003-1008 (2012).
54.   M. Kanayama *et al.*, Myeloid-like B cells boost emergency myelopoiesis through IL-10 production
during infection. *J Exp Med* **220**, (2023).
55.   X. Zhang, Regulatory functions of innate-like B cells. *Cell Mol Immunol* **10**, 113-121 (2013).
56.   T. Ju, R. D. Cummings, A unique molecular chaperone Cosmc required for activity of the
mammalian core 1 beta 3-galactosyltransferase. *Proc Natl Acad Sci U S A* **99**, 16613-16618 (2002).
57.   Y. Matsumoto *et al.*, Identification of Tn antigen O-GalNAc-expressing glycoproteins in human
carcinomas using novel anti-Tn recombinant antibodies. *Glycobiology* **30**, 282-300 (2020).
58.   G. J. Rowse, R. M. Tempero, M. L. VanLith, M. A. Hollingsworth, S. J. Gendler, Tolerance and
immunity to MUC1 in a human MUC1 transgenic murine model. *Cancer Res* **58**, 315-321 (1998).
59.   S. Cascio, A. M. Farkas, R. P. Hughey, O. J. Finn, Altered glycosylation of MUC1 influences its
association with CIN85: the role of this novel complex in cancer cell invasion and migration.
Oncotarget **4**, 1686-1697 (2013).
60.   O. J. Finn *et al.*, Importance of MUC1 and spontaneous mouse tumor models for understanding
the immunobiology of human adenocarcinomas. *Immunol Res* **50**, 261-268 (2011).
61.   D. K. Kadayakkara *et al.*, Inflammation driven by overexpression of the hypoglycosylated
abnormal mucin 1 (MUC1) links inflammatory bowel disease and pancreatitis. *Pancreas* **39**, 510-
515 (2010).
62.   S. O. Ryan, A. M. Vlad, K. Islam, J. Gariepy, O. J. Finn, Tumor-associated MUC1 glycopeptide
epitopes are not subject to self-tolerance and improve responses to MUC1 peptide epitopes in
MUC1 transgenic mice. *Biol Chem* **390**, 611-618 (2009).
63.   M. D. Burdick, A. Harris, C. J. Reid, T. Iwamura, M. A. Hollingsworth, Oligosaccharides expressed
on MUC1 produced by pancreatic and colon tumor cell lines. *J Biol Chem* **272**, 24198-24202
(1997).
64.   B. T. Hofmann *et al.*, COSMC knockdown mediated aberrant O-glycosylation promotes oncogenic
properties in pancreatic cancer. *Mol Cancer* **14**, 109 (2015).
65.   C. B. Madsen *et al.*, Glycan elongation beyond the mucin associated Tn antigen protects tumor
cells from immune-mediated killing. *PLoS One* **8**, e72413 (2013).
66.   N. Baumgarth, B-1 Cell Heterogeneity and the Regulation of Natural and Antigen-Induced IgM
Production. *Front Immunol* **7**, 324 (2016).
67.   N. Rodriguez-Zhurbenko, T. D. Quach, T. L. Rothstein, A. M. Hernandez, Human B-1 cells are
important contributors to the naturally-occurring IgM pool against the tumor-associated
ganglioside Neu5GcGM3. *Front Immunol* **13**, 1061651 (2022).

504
505

ACKNOWLEDGEMENTS

We acknowledge the following individuals for contributing transgenic mice, cell lines, and monoclonal antibodies: Dr. Akiko Iwasaki, Dr. Richard Cummings, Dr. Kebin Liu, Dr. Sarah Gendler. National Institutes of Health grants R01AI123383, R01AI152766, and R41AI157287 supported this work.

Figure legends:

Fig. 1. Tumor growth and CD301b expression in Tn^{lo} and Tn^{hi} breast cancer models. AT3 (Tn^{lo}) and AT3 cosmc KO cells (Tn^{hi}) (2.5E5/mouse) were injected into the mammary pads of C57BL/6 mice (n = 25). Tumor sizes were monitored. **A.** Tumor sizes were calculated based on (length times width^2)/2 for volumes in mm^3. **B-C.** Tumor-free and tumor-bearing mice were euthanized, and tissues were obtained from spleens, lymph nodes, bone marrow, and tumors. Flow cytometry was performed to assess immune profiles. CD45+ cells were gated from total mononuclear cells (FSC/SSC) and viable cells according to the gating strategy in Supplementary Fig. 2. CD301b+ cells were then gated from CD45+ cells and assessed in tumor-free and tumor-bearing tissues.

Fig. 2. Characterization of CD45+CD301b+ immune cells. Tn^{hi} AT3 tumor-bearing mice were euthanized, and tumor tissues were harvested. Single-cell suspensions were stained for indicated immune cell markers. CD45+CD301b+ immune cells were gated according to fluorescence minus one (FMO) plus isotype staining controls (Supplementary Fig. 2) in: **A-C.** CD45+CD301b+ cells were gated in Tn^{hi} tumor tissues, and immune cell populations (B220, CD19, CD11c, CD11b, CD3, NK1.1) were assessed compared to indicated controls. **D.** Quantification of A-C. **E.** Gated $CD45^+CD301b^+B220^+CD19^+$ cells were assessed for CD11c and CD11b. F. Comparison between the total numbers of $CD45^+B220^+CD19^+$ cells, $CD45^+B220^+CD19^+CD301b^+$ cells, and all other $CD45^+$ cells in Tn^{hi} and Tn^{lo} tumors per mg of tumor.

Fig. 3. Phenotypes of Tn^{hi} tumor-infiltrating CD45$^+$CD301b$^+$ B cells. Tn^{hi} AT3 tumor-bearing
mice were euthanized, and tumor tissues were harvested. Single-cell suspensions were stained for
indicated immune cell markers. **A-E.** CD45$^+$CD301b$^+$ B cells (B220$^+$CD19$^+$) were gated according
to FMO) plus isotype staining controls (Supplementary Fig. 2) and assessed for the presence of the
surface proteins associated with regulatory B cells **F.** Quantification of **A-E**. **G.** Mean fluorescent
intensity (MFI) of CD301b was assessed in gated CD45$^+$B220$^+$CD19$^+$ cells infiltrating Tn^{lo} and
Tn^{hi} tumors **H**. Splenocytes were obtained from tumor-free C57BL/6 mice. Cells were unselected,
positively selected for CD19$^+$ cells, or negatively isolated based on pan B cell markers (please see
Supplementary Fig. 3 for cell purity). Each splenocyte sample was co-cultured with or without Tn^{hi}
or Tn^{lo} AT3 tumor cells for three days. Flow cytometry was used to analyze CD301b$^+$ expression
in CD45$^+$B220$^+$CD19$^+$ cells. Indicated values represent the mean fluorescent intensity of CD301b$^+$.

Fig. 4. **Tumor growth is attenuated in a B cell-deficient mouse model.** Tn^{hi} tumors were injected
into the mammary pads of WT or RAG KO mice. Tumor growth was **A.** monitored. When tumor
sizes reached a maximum dimension of 1.2 cm, tumor tissues were collected, and **B-C** analyzed for
CD301b+ immune cell infiltration. **D.** Total numbers of CD45+B220+CD19+ cells in Tn^{hi} tumors
in WT C57BL/6 and RAG KO mice per mg of the tumor.

Fig. 5. Tn-associated Regulation by CD45$^+$CD301b$^+$ tumor-infiltrating immune cells. A-B
AT3 Tn^{hi} (cosmc KO) cells (2.5E5/mouse) were injected into the mammary pad in 4-6 mice per
group. Mice were euthanized, and tumors were processed for single-cell suspensions. Cells were
stained for extracellular markers, and intracellular staining was used to assess **A.** Mean Fluorescent
Intensity (MFI) and **B.** % positive of indicated parent populations of cytokines IL10, IL35 (EBI3

subunit), IL35 (p35 subunit), and TGFβ in gated cell populations (please see our gating strategy in

Supplementary Fig.4).

Fig. 6. Tumor growth in CD301b-depleted and Tn^{hi} breast cancer models. A. Tn^{lo} (AT3) and

Tn^{hi} (AT3 cosmc KO) tumor cells (2.5x10^5/mouse) were injected into heterozygous MGL2-DTR

mice with or without CD301b$^+$ cells depleted (n= 4 or 5) mice per group into the mammary pad.

Tumor sizes were monitored. **B.** Tn^{hi} tumor cells (2.5x10^5/mouse) were injected into the mammary

pads of homozygous MGL2-DTR mice (n=5). Tumor sizes were monitored. **C**. Methylcholanthrene

(MCA) was injected into the mammary pads of WT C57BL/6 and MUC1.Tg (Tn^{hi}) mice to generate

spontaneous breast tumors. Survival of mice (tumor dimensions > 1.2cm) were monitored. **D.**

MCA-induced tumor tissues were obtained from mice, and flow cytometry was used to assess

CD301b$^+$ expression on tumor-infiltrating B cells.

Tn^hi Tumor-infiltrating Gated on CD45+ CD301b+

A   CD19+ B220+

B   CD11c+ CD11b+

C   CD3+ NK1.1+

D

E Tn^hi Tumor-infiltrating Gated on CD45+ CD301b+ B220+ CD19+

F

F. Tumor-Infiltrating CD301b+ B Cell Phenotype

H. MFI of CD301b-APC on B220+CD19+ Gated Cells from CD45+

A

Tumor Growth in CD301b Depletion

- Tn^lo
- Tn^lo CD301b-depleted
- Tn^hi
- Tn^hi CD301b-depleted

B

Breast Tumor Growth

- WT
- MGL2-DTR homozygous

C

Survival Curve - Spontaneous Breast Tumor

- WT
- MUC1.Tg

p = 0.03

D

CD301b+

Supplementary Materials for

A regulatory B cell subpopulation expressing CD301b lectin promotes breast cancer growth

Authors: Amy V. Paschall[1]†, Zahra Nawaz[1]†, and Fikri Y Avci[1*]

Affiliations: [1]Department of Biochemistry, Emory Vaccine Center, Winship Cancer Institute, Emory University School of Medicine, Atlanta, GA 30322

†These authors contributed equally.

*Corresponding author. Email: favci@emory.edu

This PDF file includes:

Figs. S1 to S5

Generation of Tn^{hi} murine breast cancer cells. A-B. CRISPR-Cas9 KO of cosmc chaperone increases Tn cell surface expression in AT3 murine breast cancer cells. AT3 (Tn^{lo}) and AT3 cosmc KO (Tn^{hi}) cells were stained with A. mouse anti-Tn (reBaGs6 IgM) primary and anti-IgM secondary antibodies, with anti-IgM fluorescent antibody only as a control, and B. biotinylated VVL followed by fluorescently-conjugated streptavidin, with fluorescent streptavidin only as a control. **C.** RNA was isolated from AT3 and AT3 Tn^{hi} (cosmc KO) cell lines. qPCR was performed in triplicate wells using validated C1GALT1 (cosmc) qPCR primers and using β-actin as a control for each sample. Fold changes were normalized against the average of Tn^{hi} group. **D.** AT3 (Tn^{hi})and AT3 cosmc KO cells (Tn^{hi}) were cultured in 96 well plates (n=5) at 1E4 cells/200mL for 3 days at 37 degrees. MTT proliferation assay was performed to determine cell growth.

Fig. S2.

Gating strategy and FMO isotype control representation for flow cytometry analysis of mouse tissues. Bone marrow, lymph nodes, spleens, and tumors were harvested from mice. Single-cell suspensions were generated and stained for flow cytometry using an Fc blocker to minimize background staining. Shown is a representative tumor sample. Immune cells were gated based on the forward and side scatter areas. Singlets were further gated based on forward scatter height. Viable cells were further gated based on GhostRed710 negative cells. CD45+ cells were then gated, and immune populations were gated on CD45+ cells as described.

Fig. S3.

B Cell purification from Mouse Spleen. Spleens were obtained from healthy, tumor-free C57BL/6 mice (n=3). Spleens were processed into single-cell suspensions under sterile conditions. Spleen samples were then aliquoted for unpurified, pan B isolation (according to the BioLegend Pan B Isolation kit for mice) or CD19+ selection (according to the BioLegend CD19+ Selection kit for mice). After magnetic sorting, cell samples were stained with anti-CD45 and anti-B220 fluorescently conjugated antibodies. B cell populations were assessed using flow cytometry to analyze B220+ populations out of CD45+ gated live immune singlets.

Fig. S4.

Intracellular staining of regulatory cytokines. Cell samples were obtained from tissues of Tn^hi tumor-bearing mice. Single-cell suspensions were obtained and stained for extracellular markers. Cell samples were then fixed, permeabilized, and stained for indicated intracellular cytokines. Flow cytometry was performed. CD45+ CD301b+ B220+ CD19+ from tumor tissues of tumor-bearing mice were compared to CD45+ B220+ CD19+ gated cells from spleens of tumor-free C57BL/6 mice. Shown is the representative gating strategy along with the appropriate controls.

Fig. S5.

A-B. Depletion of CD301b+ cells following DT treatment in MGL2-DTR mice. Tn^{hi} **(**AT3 cosmc KO cells) tumor cells (2.5×10^5/mouse) were injected into the mammary pads of mice. DT was injected intraperitoneally (I.P.) at day -1 before tumor injection and administered every 2-3 days during tumor growth. Mice were euthanized and indicated tissues were processed for single-cell suspensions. Flow cytometry assessed CD301b expression from CD45+ immune cells in bone marrow and tumor. **C.** Validation of B cell absence in RAG1 KO mice. RAG1 KO mice (B6.129S7-Rag1tm1Mom/J) were obtained from Jackson Laboratory. Spleens were obtained, and flow cytometry was used to confirm the absence of CD45+ B220+ CD19+ B cells. <insert page break here>