Title: CD301b lectin expression in the breast tumor microenvironment augments tumor growth

Authors: Ahmet Ozdilek[1]†, Amy V. Paschall[1]†, Zahra Nawaz[1], Afaq M. M. Niyas[1], and Fikri Y. Avci[1]*

Affiliations: [1]Department of Biochemistry, Emory Vaccine Center, Winship Cancer Institute, Emory University School of Medicine, Atlanta, GA 30322

†These authors contributed equally.

*Corresponding author. Email: favci@emory.edu

Abstract

Aberrant tumor glycosylation can alter immune recognition; however, the specific influence of glycan-lectin interactions on tumor progression remains poorly understood. Here, we identify the C-type lectin receptor CD301b (encoded by *Mgl2*) as a regulator of immune activity within the breast tumor microenvironment (TME). Using a murine triple-negative breast cancer model, we demonstrate that tumors expressing the Tn glycoantigen grow more rapidly, and this growth is facilitated by CD301b+ immune cells. Depletion or genetic loss of CD301b markedly suppressed tumor growth, indicating that CD301b promotes tumor progression through myeloid-tumor interactions. Phenotypic analyses revealed that CD301b+ cells within tumors are type 2 conventional dendritic cells (cDC2s), a subset known to influence immune polarization. Single-cell RNA sequencing of human breast cancers showed that the human ortholog CLEC10A is expressed in cDC2-like dendritic cells and select macrophage subsets, suggesting a conserved role for CD301+ myeloid populations. Transcriptomic profiling of tumors developed in *Mgl2*-deficient mice revealed a shift toward an inflammatory, immune-activated state consistent with enhanced antitumor immunity. Together, these findings establish a link between tumor glycosylation and lectin signaling of myeloid cells, highlighting CD301b as a potential target for reprogramming the tumor immune microenvironment in breast cancer.

Introduction

A major setback observed in many cancers arises from immune modulation within the tumor microenvironment (TME), preventing effective anti-cancer responses. Vaccines, CAR T cells, and checkpoint blockade prime the immune system against cancer-associated antigens, promoting cancer cell destruction through immune cytotoxicity [1]. Immune therapies for breast cancer remain largely ineffective due to the ability of the tumor to grow unimpeded and eventually metastasize. Immune regulation can occur through cytokine secretion (such as IL-10), signaling mechanisms, or checkpoints (such as through PD-1) [2-5]. During the malignant transformation of a mammalian cell, a dramatic and aberrant modification of cellular glycosylation is observed. Tumor-associated carbohydrate antigens, or TACAs, can induce immune suppression, allowing cancer cells to evade immune cells [6-8]. Lectins are carbohydrate-binding proteins that function as receptors for immune cells and can activate immune regulatory pathways through their interactions with TACAs [9-11]. Similarly, modulating immune profiles in the TME by engaging sialic-acid-binding immunoglobulin-like lectins (Siglecs) is a known TACA mechanism [12-14]. Thus, elucidating the roles of lectins in regulating the immune response within the TME is becoming increasingly important.

Tn antigen is a TACA ranked as a high-priority cancer-associated antigen based on its antigenicity and oncogenicity [11,15-19]. Tn is a truncated form of the cell surface O-glycan, consisting of the terminal O-linked N-acetylgalactosamine (GalNAc) attached to serine or threonine. The aberrant glycosylation associated with Tn may occur when the enzyme responsible for O-glycan elongation, T-synthase or its associated chaperone, Cosmc (C1GALT1C1), becomes functionally inhibited [20,21]. Tn-expressing mucin 1 (MUC1) has been associated with breast cancer cells [22,23], especially in triple-negative breast cancer [24,25]. Modulating MUC1 expression or Tn glycosylation can inhibit tumor growth [26-28]. However, developing immune responses against Tn-MUC1 has been problematic [29].

CD301, also known as macrophage galactose-type lectin (MGL) or CLEC10A, is a C-type lectin receptor (CLR) that binds to Tn antigen on the surface proteins in humans [19,30]. In mice, the Tn-recognizing homolog of the human CD301 is CD301b, also known as MGL2 [19,31]. CD301b is primarily expressed by myeloid cells such as dendritic cells (DCs) and macrophages [17]. Previously, CD301b-expressing myeloid cell populations have been linked with immunosuppressive responses [16,17,19,32-34]. DCs and macrophages can suppress the proliferation of CD4$^+$ effector T lymphocytes through the interaction of MGL with terminal GalNAc residues on CD45 expressed by T cells [17]. This cell-specific glycosylation of CD45 provided an immunoregulatory pathway, mediated by MGL, thereby controlling effector T cell function. In another study, CD301b$^+$ dendritic cells suppressed T follicular helper cells and antibody responses to protein antigens [32]. Recently, an immunosuppressive DC subset expressing CD301b was shown to accumulate at secondary sites and promote metastasis in pancreatic cancer [34] and lung cancer [35].

In this study, we investigated the role of CD301b in a murine model of triple-negative breast cancer and found that the loss of CD301b expression significantly restricted tumor growth. Within the tumor microenvironment, CD301b-expressing immune cells were identified as type 2 conventional dendritic cells. Analysis of publicly available single-cell RNA sequencing (scRNA-seq) datasets revealed similar CD301-expressing myeloid cell populations in human breast cancer tissues, suggesting that this regulatory axis may extend beyond the murine model. To further explore the mechanisms underlying the observed tumor growth restriction, we performed bulk RNA sequencing (bulk RNA-seq) on murine tumors, which showed heightened inflammatory immune responses when CD301b was absent. Together, these findings link CD301 expression to the control of tumor-associated inflammation and point to its potential as a target for new breast cancer therapies.

Results

CD301b/Tn Axis Impacts Tumor Growth.

We first aimed to investigate the relationship between CD301b[+] immune cells and Tn glycoantigen-

expressing breast cancer cells. We used a CRISPR-Cas9 gene editing model to knock out Cosmc

expression in AT3 murine breast cancer cells. Cosmc is a chaperone essential in elongating the core

O-glycan beyond the truncated Tn form of α-GalNAc [20,21,36]. When the Cosmc function/expression

is disrupted, elongation of the O-glycan is not observed; instead, the Tn antigen is observed at

significantly higher levels. After disrupting Cosmc expression in these cell lines, we confirmed

increased Tn cell surface expression through flow cytometry using both a reBaGs6 IgM antibody

(Suppl. Fig. 1A) [37] as well as a complementary biotinylated VVL lectin coupled with fluorescent

streptavidin (Suppl. Fig. 1B). Both staining methods indicated significantly higher Tn expression

on the *Cosmc* KO cell line. We also isolated RNA from each cell line and confirmed decreased

Cosmc expression in the Tn[hi] cell line through qPCR using *Cosmc* primers (Suppl. Fig. 1C). We

then tested whether knocking out Cosmc expression changes the proliferation rate of the AT3 breast

cancer cells *in vitro*. After culturing both cell lines at the same concentrations for three days, we

observed no significant differences in cell proliferation between the two lines (Suppl. Fig. 1D),

indicating that Tn expression alone does not promote cancer cell growth.

To determine *in vivo* tumor cell growth, we injected AT3 (Tn[lo]) and AT3 *Cosmc* KO (Tn[hi]) murine

breast tumor cells into the mammary pads of C57BL/6 mice and monitored tumor growth. We

observed that Tn[hi] tumors grew significantly faster than Tn[lo] tumors (Fig. 1A), indicating that Tn

expression impacts tumor growth rate.

To examine the contribution of CD301b-expressing immune cells to tumor growth, we employed

heterozygous *Mgl2[+/DTReGFP]* mice (Mgl2-DTR), in which CD301b+ immune cells can be selectively

depleted by diphtheria toxin (DT) treatment [32,33]. In the first experiment (Fig. 1B), AT3 Tn[low] or

Tn[hi] tumor cells were injected into Mgl2-DTR mice with or without DT administration. Tumor

growth was significantly reduced in DT-treated mice of the Tn[hi] group, indicating that CD301b[+]

immune cells promote tumor progression. In a complementary experiment (Fig. 1C), we injected

AT3 Tn^{hi} cells into homozygous *Mgl2$^{DTReGFP/DTReGFP}$* mice (CD301b-null, *Mgl2* KO), which lack

surface expression of CD301b due to the disruption of both alleles but retain the immune cell

populations. These mice also exhibited markedly reduced tumor growth compared with wild-type

controls. Together, these experiments indicate that the observed phenotype is associated with tumor

Tn expression and facilitated by the CD301b protein.

Tumor-infiltrating CD301b$^+$ cells display a type 2 conventional dendritic cell (cDC2) phenotype.

We next characterized tumor-infiltrating CD45$^+$CD301b$^+$ immune cells in the murine triple-

negative breast cancer model. These cells expressed CD11c, a canonical dendritic cell (DC)

marker [38,39] (Fig. 2A) and were strongly positive for MHCII, confirming their DC identity (Fig.

2B). CD301b$^+$ cells also expressed CD11b, a defining marker of mouse type 2 conventional

dendritic cells (cDC2s) [40] (Fig. 2B). Mouse cDC2s can be distinguished from cDC1s by CD103

and SIRP-alpha expressions [38,41]. We examined the expression of these markers by tumor-

infiltrating CD301b$^+$ cells, which display a cDC2s phenotype (Fig. 2C). Although CD301b$^+$ cells

are primarily cDC2s, not all DCs or cDC2s express CD301b in the TME (Fig. 2D and 2E).

cDC2s constitute a subset of antigen-presenting cells that play key roles in coordinating adaptive

immune responses. In contrast to cDC1s, which specialize in cross-presentation and cytotoxic T

cell activation, cDC2s promote CD4$^+$ T cell priming and modulate immune polarization within

tissues [42-44]. Recent studies have shown that cDC2s exhibit remarkable plasticity in the tumor

microenvironment, where they can adopt either immunostimulatory or tolerogenic phenotypes in

response to local cues [44].

CD301$^+$ immune cells in the human breast cancer TME include both dendritic cells and

macrophages

Since CD301b$^+$ cells in the murine tumor microenvironment were identified as cDC2s, we then

investigated whether CD301 expression in human breast cancers similarly correlated with dendritic

cells or extended to other myeloid subsets. To address this, we analyzed publicly available single-

cell RNA sequencing (scRNA-seq) data (GSE161529) from 20 patients, including triple-negative (n = 8), ER[+] (n = 6), and HER2[+] (n = 6) tumors [45]. Following quality control, data integration and annotation, we focused on CD45[+] immune cells to map CLEC10A (human CD301) expression across myeloid populations (Fig. 3A; Suppl. Fig. 3A). CLEC10A expression was most prominent in dendritic cells and was also detectable in macrophage/monocyte populations, with negligible expression in other immune cells (Fig. 3B–C). Across breast cancer subtypes, dendritic cells consistently showed higher CLEC10A expression than macrophages (Suppl. Fig. 3B). Within the dendritic cell compartment, we identified four subsets—cDC1, cDC2, cDC-LAMP3[+], and plasmacytoid DC (pDC)—and found that CLEC10A expression was highest in cDC2 (56.9%) and moderate in cDC-LAMP3[+] (11.3%), but low in cDC1 (1.9%) and absent in pDCs (0%) (Fig. 3D–F; Suppl. Fig. 3C–D).

We next examined tumor-associated macrophages (TAMs). TAMs are key regulators of tumor inflammation, tissue remodeling, and immune suppression [46]. TAMs were subdivided into transcriptionally defined subsets reflecting distinct functional programs: C1QC[+] macrophages, associated with immunosuppression [47-49]; NLRP3[+] macrophages, associated with poor prognosis and tumor growth [50,51]; and INHBA[+] macrophages, linked to angiogenesis, matrix remodeling, and tumor progression [52] (Fig. 3G; Suppl. Fig. 3E). Among these subsets, *CLEC10A* transcription distributed similarly in C1QC[+] macrophages (11%), NLRP3[+] macrophages (9%), and INHBA[+] macrophages (7%) (Fig. 3H–I). Interestingly, CLEC10A-positive NLRP3[+] macrophages were detected exclusively in triple-negative breast cancers, where expression levels were comparable to those in INHBA[+] macrophages (Suppl. Fig. 3F). This enrichment suggests that CD301 expression extends beyond dendritic cells to select macrophage populations, particularly those engaged in inflammatory and tissue-remodeling responses within aggressive tumor subtypes.

Taken together, these results indicate that CD301 expression in the human breast cancer microenvironment is concentrated within myeloid lineages—encompassing both cDC2-like

dendritic cells and specialized macrophage subsets. This pattern mirrors the cellular distribution

observed in the mouse TME, suggesting that CD301 marks a conserved myeloid program

potentially involved in coordinating immune regulation and tissue remodeling during tumor

progression.

The lack of CD301b is associated with a strong inflammatory immune signature in the breast

TME

Since CD301b directly associates with tumor growth and identifies cDC2 and macrophage

populations within the breast tumor microenvironment (TME), we then examined how its loss

impacts immune signaling and tumor–immune interactions. To this end, we performed bulk RNA

sequencing (bulk RNA-seq) on tumors derived from wild-type (WT) and *Mgl2* knockout (*Mgl2*

KO) mice following injection of AT3 Tn^{hi} tumor cells. The global transcriptomic heatmap (Fig.

4A) revealed distinct clustering and clear separation between WT and *Mgl2* KO mouse tumors,

indicating a strong transcriptional divergence associated with *Mgl2* loss. This observation provided

the foundation for downstream pathway and gene-level analyses to elucidate how CD301b

influences immune regulation within the TME.

To dissect these transcriptomic differences, Gene Set Enrichment Analysis (GSEA) was performed

using the Hallmark and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment gene set

databases (Fig. 4B–C). *Mgl2* KO tumors displayed significant enrichment of immune and

inflammatory pathways, including TNF-α/NF-κB, IL-6/JAK-STAT3, interferon-α/γ, and IL-17

signaling—pathways broadly associated with myeloid activation, cytokine production, and tumor

immunosurveillance [53,54]. Despite the slower tumor growth observed in the knockout mice,

complement/coagulation, hypoxia, and glycolytic pathways were also upregulated, suggesting a

compensatory increase in metabolic and inflammatory activity within the TME. Collectively, these

data indicate that CD301b functions as an immunoregulatory node that tempers cytokine and

interferon responses, whereas its loss enhances pro-inflammatory signaling. The dominance of

interferon- and NF-κB–driven programs aligns with the slower tumor progression observed in KO

mice, implying that CD301b deficiency reprograms the TME toward a functionally immune-active,

tumor-controlling state.

The pathway-focused volcano plot (Fig. 4D) highlights key upregulated genes underpinning these

responses, including *Cxcl3, Il1a, Il23a, Il36g, Csf3, Nos2, S100a8, S100a9,* and *Lcn2*. These

represent canonical NF-κB and IL-17 targets known to mediate myeloid recruitment, nitric oxide

production, and acute-phase inflammation—hallmarks of innate immune activation [55-57]. The

upregulation of the serine protease inhibitor and matrix metalloproteinase genes *Serpinb2* and

Mmp10 in the tumor microenvironment of knockout mice suggests a tumor-suppressive function

through the regulation of extracellular matrix remodeling and modulation of immune responses

[58,59]. Thus, CD301b loss produces a highly activated, cytokine-rich TME that maintains elements

of tissue repair while fostering effector cell response and immune surveillance.

The volcano plot of all differentially expressed genes (DEGs) (Fig. 4E) contextualizes these

changes within the full transcriptome. Upregulated genes largely mirrored those driving enriched

pathways, confirming that *Cxcl3, Il1a, Il23a, Nos2, S100a8/a9,* and *Csf3* constitute the core *Mgl2*

KO transcriptional program rather than isolated pathway artifacts. On the other hand,

downregulation of the TGF-β family growth differentiation factor 3 and the apelin receptor genes

Gdf3 and *Aplnr* is associated with promoting tumor growth, angiogenesis, and metastasis [60,61].

Together, these results suggest that CD301b deficiency reprograms the TME toward a cytokine-

driven, inflammatory, and interferon-dominant state that enhances immune activation and tumor

control. The loss of CD301b lifts an immunoregulatory brake within the TME, unleashing broad

inflammatory signaling that, despite introducing metabolic stress, creates a net anti-tumor

environment characterized by immune activation and delayed tumor growth.

Discussion

Aberrant glycosylation is a defining hallmark of malignant transformation, and our findings identify CD301b as a key immunoregulatory lectin linking breast tumor-associated Tn antigens to myeloid immune modulation. CD301b+ cells, primarily cDC2s, promoted breast tumor growth, whereas their depletion or genetic loss limited progression. These observations align with previous studies demonstrating that TACAs interact with lectins to influence myeloid differentiation and immune regulation [19,21,22,62-64].

Transcriptomic profiling of tumors developed in *Mgl2*-deficient mice revealed broad activation of NF-κB, IL-6–JAK–STAT3, and interferon pathways, consistent with a shift toward a pro-inflammatory, "immune-hot" microenvironment [65]. These data position CD301b as an immunoregulatory node that tempers innate activation, similar in concept to a checkpoint-like mechanism operating within the myeloid compartment [9,10,66,67]. While this study does not define the signaling circuitry involved, it indicates that CD301b expression in cDC2s and macrophages contributes to a previously unrecognized immunoregulatory phenotype in breast cancer TME.

The complex and pleiotropic nature of CD301b's activity may explain its influence on the immune landscape of cancer [66-68]. CD301b's modulatory behavior mirrors that of other C-type lectin receptors [69]. CD301b-mediated restraint may protect against chronic inflammation, but in tumors, it can inadvertently favor immune escape [70,71]. Conversely, CD301b loss triggers inflammation driven by NF-κB and interferons, which enhances effector recruitment but imposes metabolic and hypoxic stress on the TME [72]. Together, these findings support a model in which CD301b functions as a molecular regulator balancing immune activation and tolerance in breast cancer.

Mechanistically, CD301b–Tn interactions may regulate antigen processing, cytokine release, or costimulatory signaling within dendritic cells, thereby shaping T cell activation thresholds [69]. These possibilities remain speculative but highlight the potential of CD301b as a new immune-regulatory axis distinct from canonical checkpoints, such as PD-1 or CTLA-4. Future work should investigate how TACA ligands and intracellular adaptors regulate CD301b signaling and consequent immune

programming in the TME. Further single-cell and spatial analyses of breast TME transcriptome in

the presence and absence of CD301b will help define how CD301b⁺ subsets integrate into existing

immune networks across tumor stages.

From a translational perspective, targeting the CD301–Tn interaction offers a promising route to

remodel the breast cancer immune microenvironment. Pharmacologic blockade or glycomimetic

interference could complement checkpoint inhibitors by dismantling glycan-mediated myeloid

suppression, whereas selective induction of this pathway may have therapeutic relevance in

autoimmune disease [73]. Ultimately, elucidating the intricate mechanisms that govern immune

modulation through CD301b will be essential to selectively induce these properties in disease-

specific contexts, enabling the development of knowledge-based, precision immunotherapies.

In summary, this study identifies an immunomodulatory CD301b⁺ myeloid phenotype that

contributes to breast cancer growth and whose loss induces a robust inflammatory program in the

breast TME. While the molecular mechanisms remain to be defined, our data indicate that CD301b

acts as a glycan-sensitive, checkpoint-like regulator of myeloid activity. Elucidating its ligand

specificity and downstream signaling will clarify how breast tumor glycosylation reshapes TME

and may reveal new strategies to enhance immunotherapy efficacy.

Materials and Methods

Mice

Eight-week-old female C57BL/6 mice were obtained from Jackson Laboratories (Bar Harbor, ME)

and housed at Emory University Whitehead Biomedical Research Building. *Mgl2^{DTReGFP/DTReGFP}*

mice were a generous gift from Akiko Iwasaki at Yale University. To obtain heterozygous

Mgl2^{+/DTReGFP} mice, C57BL/6 were bred with *Mgl2^{DTReGFP/DTReGFP}* mice. Mice were kept in

microisolator cages and handled under biosafety level 2 (BSL2) hoods. For tissue processing and

subsequent flow cytometry, mice were euthanized by carbon dioxide inhalation in accordance with

IACUC guidelines. Where applicable, cell suspensions were generated through mechanical tissue

disruption and collagenase D digestion. Red blood cells were lysed, and samples were filtered

through 60 μm nylon filters to obtain single-cell suspensions. For depletion of CD301b[+] cells,

heterozygous Mgl2-DTR mice were treated with diphtheria toxin (0.5μg/mouse) in sterile PBS

intraperitoneally every two to three days, starting at day -1 before tumor injection.

All mouse experiments were in compliance with the Emory University Institutional Animal Care

and Use Committee under an approved animal use protocol. Our animal use protocol adheres to the

principles outlined in *U.S. Government Principles for the Utilization and Care of Vertebrate*

Animals Used in Testing, *Research and Training*, the Animal Welfare Act, the *Guide for the Care*

and Use of Laboratory Animals, and the *AVMA Guidelines for the Euthanasia of Animals*.

Generation of Tn^hi breast cancer cells

To express Tn glycans at high levels in tumor cells, we used a CRISPR/Cas9 methodology to stably

silence Cosmc expression in AT3 cells using established protocols and reagents. Mouse *Cosmc*

guide RNA and CRISPR/Cas9 plasmid were obtained from Santa Cruz Technology (sc-425587).

AT3 murine breast cancer cells were a generous gift from the Kebin Liu lab at Augusta University.

AT3 cells were transfected with *Cosmc* CRISPR/Cas9 KO plasmid according to the manufacturer's

protocol. Puromycin was used to select transfected cells. We then used flow cytometry to confirm

higher expression of Tn on the AT3 cell surfaces using the ReBaG6 antibody (generously provided

by Richard Cummings at Harvard University) (Suppl. Fig. 1A) [37] and VVL lectin (Vector

Laboratories) (Suppl. Fig. 1B). The proliferations of transfected and untransfected cell lines (AT3

Tn^hi and AT3 Tn^low) were tested in an MTT proliferation assay for three days, and colorimetric

analysis was performed with a CytoTek plate reader according to protocol; no significant

differences in proliferation compared to parent cells were observed (Suppl. Fig. 1D). Cell lines were

maintained in RPMI media supplemented with 10% FBS, sodium pyruvate, HEPES buffer, NEAA,

β-mercaptoethanol, and penicillin/streptomycin at 37°C, 5% CO2.

AT3 Tn^low and/or AT3 Tn^hi Tumor Challenge

AT3 cells were harvested and washed in sterile PBS. Cells were suspended in a final concentration

of 2.5E6/ml sterile PBS. Cells were subcutaneously injected into the mammary pads of mice at

2.5E5/100μl/mouse. Mice were observed and euthanized at the tumor endpoint (maximum tumor

dimension between 0.9 cm and 1.2 cm. Tumor volumes were calculated as ((length x width x

width)/2) in mm^3.

Flow Cytometry

Cells were stained in PBS with TruStain fcX (BioLegend, Cat. No. 101320) to reduce non-specific

antibody binding. Cell samples were stained with the following antibodies and stains: CD11c-

PacBlue (BioLegend, clone N418), CD11b-Alexa Fluor 488 (BioLegend, clone M1/70), SIRPα-

Alexa Fluor 700 (BioLegend, clone P84), CD103-PE/Dazzle 594 (BioLegend, clone 2E7), MHCII-

BV785 (BioLegend, clone M5/114.15.2), CD45 Alexa Fluor 647 (BioLegend, clone 30-F11),

CD301b-PE (BioLegend, clone URA-1), and LIVE/DEAD Fixable Blue (Invitrogen). All isotype

controls were obtained from BioLegend. Samples were washed and analyzed with flow cytometry

(Cytek Aurora). Fluorescence minus one (FMO) plus specific isotype control antibody-stained

samples were used as negative staining controls, and single stains were used for compensation.

Flow cytometry data were analyzed using FlowJo Single Cell Analysis Software with gating

strategies shown in Suppl. Fig. 2.

Bulk RNA-seq and Bioinformatics

Tumors were harvested from wild-type C57BL/6J and *Mgl2* KO mice (n = 3 per group). Total RNA

was extracted using the MagMAX™ mirVana™ Total RNA Isolation Kit in combination with the

KingFisher Apex system (Thermo Fisher Scientific). RNA integrity and concentration were

assessed with a Qubit 3.0 fluorometer (Thermo Fisher Scientific). High-quality RNA samples were

used for library preparation, followed by quality control and sequencing using Novogene's standard

protocol. Libraries were sequenced on the Illumina NovaSeq X Plus platform to generate paired-

end 150 bp reads (PE150) at Novogene Inc. Raw FASTQ files were retrieved and subjected to

quality control with FastQC [74]. Reads were aligned to the *Mus musculus* reference genome

(GRCm39/mm39) using HISAT2 [75]. A gene-level count matrix was generated with featureCounts

[76]. The count matrix was imported to the downstream differential expression analysis using the

DESeq2 R package [77]. Significantly differentially expressed genes were defined by adjusted p-

value < 0.05. Gene Set Enrichment Analysis (GSEA) was performed on the RNA-seq dataset using

the clusterProfiler R package [78] to identify significantly enriched KEGG and hallmark pathways

between *Mgl2* KO and wild-type tumors. To focus on cancer-relevant biology, enrichment results

were refined to include immune- and cancer-related pathways.

Analyzing the scRNA-seq data

The scRNA-seq data generated using the 10x Genomics Chromium platform from human breast

cancer tumors (GSE161529) [45] were obtained from Gene Expression Omnibus (GEO) [79]. Data were

processed into the Seurat R package (version 5), and low-quality cells were removed based on

established Seurat quality control parameters [80]. After QC, putative doublets were removed using

the *DoubletFinder* R package [81]. Datasets were integrated, and non-immune cells were removed by

subsetting the CD45$^+$ cells with detectable *PTPRC* expression (*PTPRC* > 0). Following

preprocessing, cell annotation was contacted with scATOMIC [82]. The subclasses of the dendritic

cells and macrophages were further validated with canonical lineage markers [83,84].

Statistical Analysis

GraphPad Prism v8 was used for statistical analyses. Two-way ANOVA with Tukey's multiple

comparisons test was used to determine statistical significance between experimental groups in

each of the applicable experimental models (Fig. 1B). An unpaired parametric two-tailed t-test was

used for Fig. 1A, 1C, 2A, 2B, 2C, 2D, 2E, and Suppl. Fig. 1C. Significance is indicated on each

graph based on p-value: >0.05 = ns; <.05 = *; <0.01 = **; <0.001 = ***; <0.0001 = ****.

Acknowledgements

We acknowledge the following individuals for their contributions of transgenic mice, cell lines, and

monoclonal antibodies: Dr. Akiko Iwasaki, Dr. Kebin Liu, and Dr. Richard Cummings. National

Institutes of Health grants R01AI123383 and R01AI152766 supported this work.

References
1   Zhang, Y. & Zhang, Z. The history and advances in cancer immunotherapy: understanding the
     characteristics of tumor-infiltrating immune cells and their therapeutic implications. *Cell Mol
     Immunol* **17**, 807-821 (2020). https://doi.org:10.1038/s41423-020-0488-6
2   Swoboda, A. & Nanda, R. Immune Checkpoint Blockade for Breast Cancer. *Cancer Treat Res* **173**,
     155-165 (2018). https://doi.org:10.1007/978-3-319-70197-4_10
3   Fiorentino, D. F., Bond, M. W. & Mosmann, T. R. Two types of mouse T helper cell. IV. Th2 clones
     secrete a factor that inhibits cytokine production by Th1 clones. *J Exp Med* **170**, 2081-2095
     (1989). https://doi.org:10.1084/jem.170.6.2081
4   Acuner Ozbabacan, S. E., Gursoy, A., Nussinov, R. & Keskin, O. The structural pathway of
     interleukin 1 (IL-1) initiated signaling reveals mechanisms of oncogenic mutations and SNPs in
     inflammation and cancer. *PLoS Comput Biol* **10**, e1003470 (2014).
     https://doi.org:10.1371/journal.pcbi.1003470
5   Freeman, G. J. *et al.* Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family
     member leads to negative regulation of lymphocyte activation. *J Exp Med* **192**, 1027-1034 (2000).
6   Rodrigues Mantuano, N., Natoli, M., Zippelius, A. & Laubli, H. Tumor-associated carbohydrates
     and immunomodulatory lectins as targets for cancer immunotherapy. *J Immunother Cancer* **8**
     (2020). https://doi.org:10.1136/jitc-2020-001222
7   Laubli, H. & Varki, A. Sialic acid-binding immunoglobulin-like lectins (Siglecs) detect self-
     associated molecular patterns to regulate immune responses. *Cell Mol Life Sci* **77**, 593-605
     (2020). https://doi.org:10.1007/s00018-019-03288-x
8   Pinho, S. S. & Reis, C. A. Glycosylation in cancer: mechanisms and clinical implications. *Nat Rev
     Cancer* **15**, 540-555 (2015). https://doi.org:10.1038/nrc3982
9   Marino, K. V., Cagnoni, A. J., Croci, D. O. & Rabinovich, G. A. Targeting galectin-driven regulatory
     circuits in cancer and fibrosis. *Nat Rev Drug Discov* **22**, 295-316 (2023).
     https://doi.org:10.1038/s41573-023-00636-2
10   Cagnoni, A. J. *et al.* Galectin-1 fosters an immunosuppressive microenvironment in colorectal
     cancer by reprogramming CD8(+) regulatory T cells. *Proc Natl Acad Sci U S A* **118** (2021).
     https://doi.org:10.1073/pnas.2102950118
11   Paschall, A. V. M., D. R.; Avci, F. Y. in *Encyclopedia of Cell Biology, Second Edition* (ed R. A.; Hart
     Bradshaw, G. W.; Stahl, P. D.) 404-413 (Elsevier, 2023).
12   Beatson, R. *et al.* The mucin MUC1 modulates the tumor immunological microenvironment
     through engagement of the lectin Siglec-9. *Nat Immunol* **17**, 1273-1281 (2016).
     https://doi.org:10.1038/ni.3552
13   Marcos, N. T. *et al.* Role of the human ST6GalNAc-I and ST6GalNAc-II in the synthesis of the
     cancer-associated sialyl-Tn antigen. *Cancer Res* **64**, 7050-7057 (2004).
     https://doi.org:10.1158/0008-5472.CAN-04-1921
14   Schultz, M. J. *et al.* The Tumor-Associated Glycosyltransferase ST6Gal-I Regulates Stem Cell
     Transcription Factors and Confers a Cancer Stem Cell Phenotype. *Cancer Res* **76**, 3978-3988
     (2016). https://doi.org:10.1158/0008-5472.CAN-15-2834

15 Freire, T., Lo-Man, R., Bay, S. & Leclerc, C. Tn glycosylation of the MUC6 protein modulates its
immunogenicity and promotes the induction of Th17-biased T cell responses. *J Biol Chem* **286**,
7797-7811 (2011). https://doi.org:10.1074/jbc.M110.209742
16 Saeland, E. *et al.* The C-type lectin MGL expressed by dendritic cells detects glycan changes on
MUC1 in colon carcinoma. *Cancer Immunol Immunother* **56**, 1225-1236 (2007).
https://doi.org:10.1007/s00262-006-0274-z
17 van Vliet, S. J., Gringhuis, S. I., Geijtenbeek, T. B. & van Kooyk, Y. Regulation of effector T cells by
antigen-presenting cells via interaction of the C-type lectin MGL with CD45. *Nat Immunol* **7**, 1200-
1208 (2006). https://doi.org:10.1038/ni1390
18 Cheever, M. A. *et al.* The prioritization of cancer antigens: a national cancer institute pilot project
for the acceleration of translational research. *Clin Cancer Res* **15**, 5323-5337 (2009).
https://doi.org:10.1158/1078-0432.CCR-09-0737
19 Tumoglu, B., Keelaghan, A. & Avci, F. Y. Tn antigen interactions of macrophage galactose-type
lectin (MGL) in immune function and disease. *Glycobiology* **33**, 879-887 (2023).
https://doi.org:10.1093/glycob/cwad083
20 Wang, Y. *et al.* Cosmc is an essential chaperone for correct protein O-glycosylation. *Proc Natl*
Acad Sci U S A **107**, 9228-9233 (2010). https://doi.org:10.1073/pnas.0914004107
21 Ju, T., Aryal, R. P., Kudelka, M. R., Wang, Y. & Cummings, R. D. The Cosmc connection to the Tn
antigen in cancer. *Cancer Biomark* **14**, 63-81 (2014). https://doi.org:10.3233/CBM-130375
22 Beatson, R. *et al.* The Breast Cancer-Associated Glycoforms of MUC1, MUC1-Tn and sialyl-Tn, Are
Expressed in COSMC Wild-Type Cells and Bind the C-Type Lectin MGL. *PLoS One* **10**, e0125994
(2015). https://doi.org:10.1371/journal.pone.0125994
23 Beatson, R. E., Taylor-Papadimitriou, J. & Burchell, J. M. MUC1 immunotherapy. *Immunotherapy*
2, 305-327 (2010). https://doi.org:10.2217/imt.10.17
24 Siroy, A. *et al.* MUC1 is expressed at high frequency in early-stage basal-like triple-negative breast
cancer. *Hum Pathol* **44**, 2159-2166 (2013). https://doi.org:10.1016/j.humpath.2013.04.010
25 Burchell, J. M., Beatson, R., Graham, R., Taylor-Papadimitriou, J. & Tajadura-Ortega, V. O-linked
mucin-type glycosylation in breast cancer. *Biochem Soc Trans* **46**, 779-788 (2018).
https://doi.org:10.1042/BST20170483
26 Chen, D. *et al.* Immunotherapy of spontaneous mammary carcinoma with fusions of dendritic
cells and mucin 1-positive carcinoma cells. *Immunology* **109**, 300-307 (2003).
27 Mukherjee, P. *et al.* MUC1-specific immune therapy generates a strong anti-tumor response in a
MUC1-tolerant colon cancer model. *Vaccine* **25**, 1607-1618 (2007).
https://doi.org:10.1016/j.vaccine.2006.11.007
28 Lakshminarayanan, V. *et al.* Immune recognition of tumor-associated mucin MUC1 is achieved by
a fully synthetic aberrantly glycosylated MUC1 tripartite vaccine. *Proc Natl Acad Sci U S A* **109**,
261-266 (2012). https://doi.org:10.1073/pnas.1115166109
29 Loureiro, L. R. *et al.* Challenges in Antibody Development against Tn and Sialyl-Tn Antigens.
Biomolecules **5**, 1783-1809 (2015). https://doi.org:10.3390/biom5031783
30 Rodriguez, E. *et al.* Fasciola hepatica Immune Regulates CD11c+ Cells by Interacting with the
Macrophage Gal/GalNAc Lectin. *Front Immunol* **8**, 264 (2017).
https://doi.org:10.3389/fimmu.2017.00264
31 Singh, S. K. *et al.* Characterization of murine MGL1 and MGL2 C-type lectins: distinct glycan
specificities and tumor binding properties. *Mol Immunol* **46**, 1240-1249 (2009).
https://doi.org:10.1016/j.molimm.2008.11.021
32 Kumamoto, Y., Hirai, T., Wong, P. W., Kaplan, D. H. & Iwasaki, A. CD301b+ dendritic cells suppress
T follicular helper cells and antibody responses to protein antigens. *Elife* **5** (2016).
https://doi.org:10.7554/eLife.17979
33 Kumamoto, Y. *et al.* CD301b(+) dermal dendritic cells drive T helper 2 cell-mediated immunity.
Immunity **39**, 733-743 (2013). https://doi.org:10.1016/j.immuni.2013.08.029

Kenkel, J. A. *et al.* An Immunosuppressive Dendritic Cell Subset Accumulates at Secondary Sites and Promotes Metastasis in Pancreatic Cancer. *Cancer Res* **77**, 4158-4170 (2017). https://doi.org:10.1158/0008-5472.CAN-16-2212

da Costa, V. *et al.* The Tn antigen promotes lung tumor growth by fostering immunosuppression and angiogenesis via interaction with Macrophage Galactose-type lectin 2 (MGL2). *Cancer Lett* **518**, 72-81 (2021). https://doi.org:10.1016/j.canlet.2021.06.012

Ju, T. & Cummings, R. D. A unique molecular chaperone Cosmc required for activity of the mammalian core 1 beta 3-galactosyltransferase. *Proc Natl Acad Sci U S A* **99**, 16613-16618 (2002). https://doi.org:10.1073/pnas.262438199

Matsumoto, Y. *et al.* Identification of Tn antigen O-GalNAc-expressing glycoproteins in human carcinomas using novel anti-Tn recombinant antibodies. *Glycobiology* **30**, 282-300 (2020). https://doi.org:10.1093/glycob/cwz095

Breed, E. R. *et al.* Type 2 cytokines in the thymus activate Sirpalpha(+) dendritic cells to promote clonal deletion. *Nat Immunol* **23**, 1042-1051 (2022). https://doi.org:10.1038/s41590-022-01218-x

Kumamoto, Y., Denda-Nagai, K., Aida, S., Higashi, N. & Irimura, T. MGL2 Dermal dendritic cells are sufficient to initiate contact hypersensitivity in vivo. *PLoS One* **4**, e5619 (2009). https://doi.org:10.1371/journal.pone.0005619

Izumi, G. *et al.* CD11b(+) lung dendritic cells at different stages of maturation induce Th17 or Th2 differentiation. *Nat Commun* **12**, 5029 (2021). https://doi.org:10.1038/s41467-021-25307-x

Bottcher, J. P. & Reis e Sousa, C. The Role of Type 1 Conventional Dendritic Cells in Cancer Immunity. *Trends Cancer* **4**, 784-792 (2018). https://doi.org:10.1016/j.trecan.2018.09.001

Haniffa, M. *et al.* Human tissues contain CD141hi cross-presenting dendritic cells with functional homology to mouse CD103+ nonlymphoid dendritic cells. *Immunity* **37**, 60-73 (2012). https://doi.org:10.1016/j.immuni.2012.04.012

Liu, P., Zhao, L., Kroemer, G. & Kepp, O. Conventional type 1 dendritic cells (cDC1) in cancer immunity. *Biol Direct* **18**, 71 (2023). https://doi.org:10.1186/s13062-023-00430-5

Saito, Y., Komori, S., Kotani, T., Murata, Y. & Matozaki, T. The Role of Type-2 Conventional Dendritic Cells in the Regulation of Tumor Immunity. *Cancers (Basel)* **14** (2022). https://doi.org:10.3390/cancers14081976

Pal, B. *et al.* A single-cell RNA expression atlas of normal, preneoplastic and tumorigenic states in the human breast. *EMBO J* **40**, e107333 (2021). https://doi.org:10.15252/embj.2020107333

Mantovani, A. *et al.* The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol* **25**, 677-686 (2004). https://doi.org:10.1016/j.it.2004.09.015

Zhang, S. *et al.* C1q(+) tumor-associated macrophages contribute to immunosuppression through fatty acid metabolic reprogramming in malignant pleural effusion. *J Immunother Cancer* **11** (2023). https://doi.org:10.1136/jitc-2023-007441

Yang, J. *et al.* Integrated genomic and transcriptomic analysis reveals unique characteristics of hepatic metastases and pro-metastatic role of complement C1q in pancreatic ductal adenocarcinoma. *Genome Biol* **22**, 4 (2021). https://doi.org:10.1186/s13059-020-02222-w

Yang, Q. *et al.* Single-Cell RNA Sequencing Reveals the Heterogeneity of Tumor-Associated Macrophage in Non-Small Cell Lung Cancer and Differences Between Sexes. *Front Immunol* **12**, 756722 (2021). https://doi.org:10.3389/fimmu.2021.756722

Chen, L. *et al.* NLRP3 in tumor-associated macrophages predicts a poor prognosis and promotes tumor growth in head and neck squamous cell carcinoma. *Cancer Immunol Immunother* **72**, 1647-1660 (2023). https://doi.org:10.1007/s00262-022-03357-4

Daley, D. *et al.* NLRP3 signaling drives macrophage-induced adaptive immune suppression in pancreatic carcinoma. *J Exp Med* **214**, 1711-1724 (2017). https://doi.org:10.1084/jem.20161707

Li, F. L. *et al.* INHBA promotes tumor growth and induces resistance to PD-L1 blockade by suppressing IFN-gamma signaling. *Acta Pharmacol Sin* **46**, 448-461 (2025). https://doi.org:10.1038/s41401-024-01381-x

Mantovani, A., Allavena, P., Sica, A. & Balkwill, F. Cancer-related inflammation. *Nature* **454**, 436-444 (2008). https://doi.org:10.1038/nature07205

54   Grivennikov, S. I., Greten, F. R. & Karin, M. Immunity, inflammation, and cancer. *Cell* **140**, 883-899
(2010). https://doi.org:10.1016/j.cell.2010.01.025

55   Liu, T., Zhang, L., Joo, D. & Sun, S. C. NF-kappaB signaling in inflammation. *Signal Transduct*
Target Ther **2**, 17023- (2017). https://doi.org:10.1038/sigtrans.2017.23

56   Qian, Y., Kang, Z., Liu, C. & Li, X. IL-17 signaling in host defense and inflammatory diseases. *Cell*
Mol Immunol **7**, 328-333 (2010). https://doi.org:10.1038/cmi.2010.27

57   Medzhitov, R. Origin and physiological roles of inflammation. *Nature* **454**, 428-435 (2008).
https://doi.org:10.1038/nature07201

58   Harris, N. L. E. *et al.* SerpinB2 regulates stromal remodelling and local invasion in pancreatic
cancer. *Oncogene* **36**, 4288-4298 (2017). https://doi.org:10.1038/onc.2017.63

59   Shuman Moss, L. A., Jensen-Taubman, S. & Stetler-Stevenson, W. G. Matrix metalloproteinases:
changing roles in tumor progression and metastasis. *Am J Pathol* **181**, 1895-1899 (2012).
https://doi.org:10.1016/j.ajpath.2012.08.044

60   Yang, Y. *et al.* The Apelin/APLNR system modulates tumor immune response by reshaping the
tumor microenvironment. *Gene* **834**, 146564 (2022). https://doi.org:10.1016/j.gene.2022.146564

61   Ehira, N. *et al.* An embryo-specific expressing TGF-beta family protein, growth-differentiation
factor 3 (GDF3), augments progression of B16 melanoma. *J Exp Clin Cancer Res* **29**, 135 (2010).
https://doi.org:10.1186/1756-9966-29-135

62   Burdick, M. D., Harris, A., Reid, C. J., Iwamura, T. & Hollingsworth, M. A. Oligosaccharides
expressed on MUC1 produced by pancreatic and colon tumor cell lines. *J Biol Chem* **272**, 24198-
24202 (1997).

63   Hofmann, B. T. *et al.* COSMC knockdown mediated aberrant O-glycosylation promotes oncogenic
properties in pancreatic cancer. *Mol Cancer* **14**, 109 (2015). https://doi.org:10.1186/s12943-015-
0386-1

64   Madsen, C. B. *et al.* Glycan elongation beyond the mucin associated Tn antigen protects tumor
cells from immune-mediated killing. *PLoS One* **8**, e72413 (2013).
https://doi.org:10.1371/journal.pone.0072413

65   Medzhitov, R. The spectrum of inflammatory responses. *Science* **374**, 1070-1075 (2021).
https://doi.org:10.1126/science.abi5200

66   Bruni, D., Angell, H. K. & Galon, J. The immune contexture and Immunoscore in cancer prognosis
and therapeutic efficacy. *Nat Rev Cancer* **20**, 662-680 (2020). https://doi.org:10.1038/s41568-
020-0285-7

67   Thorsson, V. *et al.* The Immune Landscape of Cancer. *Immunity* **48**, 812-830 e814 (2018).
https://doi.org:10.1016/j.immuni.2018.03.023

68   van der Meijs, N. L., Travecedo, M. A., Marcelo, F. & van Vliet, S. J. The pleiotropic CLEC10A:
implications for harnessing this receptor in the tumor microenvironment. *Expert Opin Ther*
Targets **28**, 601-612 (2024). https://doi.org:10.1080/14728222.2024.2374743

69   Geijtenbeek, T. B. & Gringhuis, S. I. C-type lectin receptors in the control of T helper cell
differentiation. *Nat Rev Immunol* **16**, 433-448 (2016). https://doi.org:10.1038/nri.2016.55

70   Biswas, S. K. & Mantovani, A. Macrophage plasticity and interaction with lymphocyte subsets:
cancer as a paradigm. *Nat Immunol* **11**, 889-896 (2010). https://doi.org:10.1038/ni.1937

71   Mills, C. D., Lenz, L. L. & Harris, R. A. A Breakthrough: Macrophage-Directed Cancer
Immunotherapy. *Cancer Res* **76**, 513-516 (2016). https://doi.org:10.1158/0008-5472.CAN-15-
1737

72   Leone, R. D. & Powell, J. D. Metabolism of immune cells in cancer. *Nat Rev Cancer* **20**, 516-531
(2020). https://doi.org:10.1038/s41568-020-0273-y

73   Macauley, M. S., Crocker, P. R. & Paulson, J. C. Siglec-mediated regulation of immune cell
function in disease. *Nat Rev Immunol* **14**, 653-666 (2014). https://doi.org:10.1038/nri3737

74   Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data. Available online
at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc. (2010).

75 Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915 (2019).
https://doi.org:10.1038/s41587-019-0201-4
76 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).
https://doi.org:10.1093/bioinformatics/btt656
77 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org:10.1186/s13059-014-0550-8
78 Xu, S. *et al.* Using clusterProfiler to characterize multiomics data. *Nat Protoc* **19**, 3292-3320
(2024). https://doi.org:10.1038/s41596-024-01020-z
79 Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res* **30**, 207-210 (2002).
https://doi.org:10.1093/nar/30.1.207
80 Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat*
Biotechnol **42**, 293-304 (2024). https://doi.org:10.1038/s41587-023-01767-y
81 McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell
RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-337 e324 (2019).
https://doi.org:10.1016/j.cels.2019.03.003
82 Nofech-Mozes, I., Soave, D., Awadalla, P. & Abelson, S. Pan-cancer classification of single cells in
the tumour microenvironment. *Nat Commun* **14**, 1615 (2023). https://doi.org:10.1038/s41467-
023-37353-8
83 Hu, C. *et al.* CellMarker 2.0: an updated database of manually curated cell markers in
human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**, D870-D876 (2023).
https://doi.org:10.1093/nar/gkac947
84 Cheng, S. *et al.* A pan-cancer single-cell transcriptional atlas of tumor infiltrating myeloid cells.
Cell **184**, 792-809 e723 (2021). https://doi.org:10.1016/j.cell.2021.01.010

550
551

Figure legends:

Fig. 1. CD301b/Tn Axis Impacts Tumor Growth

A. AT3 (Tn^{low}) and AT3 *Cosmc* KO cells (Tn^{hi}) (2.5E5/mouse) were injected into the mammary

pads of C57BL/6 mice (n = 25). Tumor sizes were monitored. Tumor sizes were calculated based

on (length times width^2)/2 for volumes in mm^3. **B.** AT3 (Tn^{low}) and AT3 *Cosmc* KO cells (Tn^{hi})

tumor cells (2.5x10^5/mouse) were injected into the mammary pads of WT mice and heterozygous

Mgl2-DTR mice (n= 4 or 5) with or without CD301b$^+$ cells depleted. Tumor sizes were monitored.

C. AT3 *Cosmc* KO cells (Tn^{hi}) (2.5x10^5/mouse) were injected into the mammary pads of *Mgl2* KO

mice (n=5). Tumor sizes were monitored.

Fig. 2. Characterization of CD45$^+$CD301b$^+$ immune cells in murine breast cancer TME.

Single cell suspensions from tumors of WT mice injected with AT3 *Cosmc* KO cells (Tn^{hi}) were

stained, and expressions of surface markers were analyzed with flow cytometry (gated as in Suppl.

Fig. 2). **A.** Among live CD45$^+$ cells, CD301b$^+$ cells are CD11c$^+$. **B.** CD301b$^+$ and CD11c$^+$ cells are

MHCII$^+$ and CD11b$^+$. **C.** CD301b$^+$ and CD11c$^+$ cells are cDC2. DCs and cDC2s in the TME, **D**

and E, respectively, consist of Cd301b-negative and positive cells.

Fig. 3. Characterization of CD45$^+$CD301b$^+$ immune cells in human breast cancer TME.

A. UMAP of all immune cells colored by major lineage (B/Plasma cell, Dendritic cell,

Macrophage/Monocyte, NK cell, T cell). **B.** Immune cells highlighting CLEC10A$^+$ cells (colored)

over all cells (gray). **C.** Bar plot showing the fraction of CLEC10A$^+$ cells within each major lineage.

D. UMAP of dendritic cell compartment colored by subset (cDC1, cDC2, cDC-LAMP3$^+$, pDC). **E.**

Dendritic cell subset highlighting CLEC10A$^+$ cells (colored) over all DCs (gray). **F.** Bar plot

showing the fraction of CLEC10A$^+$ cells within each DC subset. **G.** UMAP of macrophage

compartment colored by subset: C1QC$^+$ macrophages, INHBA$^+$ macrophages, and NLRP3$^+$

macrophages. **H.** Macrophage subset highlighting CLEC10A$^+$ cells (colored) over all macrophages

(gray). **I.** Bar plot showing the fraction of CLEC10A$^+$ cells within each macrophage subset.

UMAP axes indicate the first two dimensions. Percentages in bar plots are calculated as

(CLEC10A$^+$ cells / total cells) within the indicated group.

Fig. 4. The lack of CD301b is associated with a strong inflammatory immune signature in the

breast TME

A. Heatmap showing differentially expressed genes (DEGs) identified by RNA-seq from tumors in

Mgl2 KO and WT mice. The blue and red bands indicate low and high gene expression quantity,

respectively. Biological replicates showed the highest degree of correlation. **B.** Gene Set

Enrichment Analysis (GSEA) of Hallmark pathways reveals immune-related and cancer-associated

pathways, all of which are upregulated in Mgl2 KO. **C.** GSEA of KEGG pathways shows immune-

related and cancer-associated pathways, and they are all upregulated in Mgl2 KO. **D.** Volcano plot

showing the pathway-associated genes (from panels B and C). The genes with a $\log_2$ fold change of 2 or greater are highlighted in purple. The X-axis represents $\log_2$-transformed fold change, and the Y-axis represents $-\log_{10}$-transformed significance. The VENN diagram shows the distribution of genes between Hallmark and KEGG gene sets. **E.** Volcano plot of DEGs between tumors from *Mgl2* KO and WT mice. Red points indicate upregulated DEGs, blue points indicate downregulated DEGs, gray points represent non-significant genes, and purple points indicate the pathway-associated genes (from panel D). All the genes with a $\log_2$ fold change ≥ 2 are labeled.

Supplementary Fig. 1.

Generation of AT3 *Cosmc* KO (Tn^{hi}) murine breast cancer cells. **A. and B.** CRISPR-Cas9 KO of Cosmc chaperone increases Tn cell surface expression in AT3 murine breast cancer cells. AT3 (Tn^{low}) and AT3 *Cosmc* KO (Tn^{hi}) cells were stained with **A.** mouse anti-Tn (reBaGs6 IgM) primary and anti-IgM secondary antibodies, with secondary antibody-only as a control, and **B.** biotinylated VVL followed by fluorescently-conjugated streptavidin, with fluorescently-conjugated streptavidin-only as a control. **C.** RNA was isolated from AT3 (Tn^{low}) and AT3 *Cosmc* KO cells (Tn^{hi}). qPCR was performed in triplicate wells using validated *Cosmc* qPCR primers and using b-actin as a control for each sample. Fold changes were normalized against the average of AT3 *Cosmc* KO cells (Tn^{hi}). **D.** AT3 (Tn^{low}) and AT3 *Cosmc* KO cells (Tn^{hi}) were cultured in 96 well plates (n=5) at 1E4 cells/200mL for 3 days at 37oC. MTT proliferation assay was performed to determine cell growth.

Supplementary Fig. 2.

Flow cytometry gating strategy for identifying dendritic cells and their subsets. DCs were identified by expressions of MHCII and CD11c (bottom, middle, right) among CD45$^+$ cells (top, right). Isotype control of CD11c antibody was used to locate CD11c$^+$ cells (bottom, right). cDC1s and cDC2s were identified by expressions of CD103 and SIRP-alpha (bottom, middle, left). Total CD45$^+$ cells were used to locate CD103$^{-/+}$, and SIRP-alpha$^{-/+}$ cells (bottom, left).

Supplementary Fig. 3.

A. Heatmap showing canonical markers used to define major immune cell lineages (B/Plasma cell,

Dendritic cell, Macrophage/Monocyte, NK cell, and T cell). **B.** Bar plots showing the frequency of

$CLEC10A^+$ cells across major immune lineages in ER^+, $HER2^+$, and triple-negative (TN) breast

cancers. **C.** Heatmap displaying canonical dendritic cell subset markers used to define cDC1, cDC2,

$cDC-LAMP3^+$, and pDC populations. **D.** Corresponding bar plots showing $CLEC10A^+$ proportions

within each DC subset across ER^+, $HER2^+$, and TN tumors. **E.** Heatmap showing markers used to

define macrophage polarization and functional states, including $C1QC^+$ macrophages, $INHBA^+$

macrophages, and $NLRP3^+$ macrophages. **F.** Bar plots showing the proportion of $CLEC10A^+$ cells

within each macrophage subtype across ER^+, $HER2^+$, and TN breast cancers.

Heatmaps display scaled average expression per cell group. Bar plots represent the percentage of

$CLEC10A^+$ cells within each indicated population.