1 **Defects in exosome biogenesis are associated with sensorimotor defects in zebrafish**
2 ***vps4a* mutants**
3
4
5 Running title: Sensorimotor defects in zebrafish *vps4a* mutants
6
7

8 Anna Shipman[1], Yan Gao[1], Desheng Liu[2], Shan Sun[2], Jingjing Zang[3], Zoha Syed[1], Amol
9 Bhagavathi[1], Eliot Smith[1,4], Timothy Erickson[5], Matthew Hill[1,6], Stephan Neuhauss[3], Sen-Fan
10 Sui[2], Teresa Nicolson[1]
11
12
[1]Department of Otolaryngology, Stanford University, Stanford, CA, United States
[2]State Key Laboratory of Membrane Biology, Beijing Frontier Research Center for Biological
Structure, Beijing Advanced Innovation Center for Structural Biology, School of Life Sciences,
Tsinghua University, Beijing, China
[3]Department of Molecular Life Sciences, University of Zurich, Zurich, Switzerland
[4]Present address: Biotechne Corporation, Newark, CA United States 94304
[5]Department of Biology, University of New Brunswick, Fredericton, Canada
[6]Present address: Department of Otolaryngology, University of Colorado, Aurora, CO, United
States
22
23
24
25 Corresponding author: Teresa Nicolson
26 Email: tnicolso@stanford.edu

27
28
29 **Conflict of Interest Statement:** The authors declare no competing financial interests.
30
31
32

Abstract

Mutations in human *VPS4A* are associated with neurodevelopmental defects, including motor delays and defective muscle tone. *VPS4A* encodes a AAA-ATPase that is required for membrane scission, but how mutations in *VPS4A* lead to impaired control of motor function is not known. Here we identified a mutation in zebrafish *vps4a*, T248I, that affects sensorimotor transformation. In biochemical experiments we show that the T248I mutation reduces the ATPase activity of Vps4a and disassembly of its substrate, ESCRT filaments, which mediate membrane scission. Consistent with the established role for Vps4a in the endocytic pathway and exosome biogenesis, *vps4a*^{T248I} mutants have enlarged endosomal compartments in the CNS and decreased numbers of circulating exosomes. Resembling the central form of hypotonia in human *VPS4A* patients, motor neurons and muscle cells are unaffected in mutant zebrafish as they react robustly to touch. Unlike somatosensory function, optomotor responses, vestibulospinal (VS), and acoustic startle reflexes are severely impaired in *vps4a*^{T248I} mutants, indicating a greater sensitivity of these circuits to the T248I mutation. ERG recordings indicate that visual ability is largely reduced in the mutants, however, *in vivo* imaging of tone-evoked responses in the inner ear and ascending auditory pathway show comparable activity. Further investigation of central pathways in *vps4a*^{T248I} mutants revealed that sensory cues failed to fully activate neurons in the VS and medial longitudinal fasciculus (MLF) nuclei that directly innervate motor neurons. Our results suggest that a defect in sensorimotor transformation underlies the profound yet selective effects on motor reflexes resulting from the loss of membrane scission mediated by Vps4a.

Introduction

Neurodevelopmental disorders are a collection of conditions that affect central nervous system growth and development and can have a wide variety of associated clinical signs, such as impaired cognition, difficulties in communication, and motor defects. Many neurodevelopmental disorders are the result of genetic mutations and often target molecular pathways that impact protein synthesis, transcriptional or epigenetic regulation or synaptic signaling [1–3]. An early feature that is common to several neurodevelopmental defects is central hypotonia (low muscle tone), however, it remains unclear how the underlying biochemical and sensorimotor mechanisms can result in this feature [4].

De novo heterozygous missense mutations in human Vacuolar protein sorting-associated protein 4A (*VPS4A*) have been associated with neurodevelopmental disorders characterized by pronounced delays in development, motor skills and speech, and severe intellectual disabilities and visual dysfunction [5]. Symptoms arising from problems with muscle tone such as hypotonia and dystonia are also thought to be central in nature [6]. In addition, mutations in *VPS4A* are associated with Congenital Dyserythropoietic Anemia [7,8]. *VPS4A* encodes a type I AAA-ATPase that was originally identified in yeast. In multicellular organisms, a paralogue, *VPS4B*, is also expressed and both genes are highly conserved from plants to humans [9–12]. VPS4A binds to the ESCRT-III complex to promote dissociation of the complex via ATP hydrolysis, resulting in membrane scission events that are key for proper viral bud release, cytokinesis, neurite severing, plasma membrane repair and multivesicular body formation [13–16]. In animal studies, it has been previously shown that RNAi knockdown of *vps4* in fruit fly larvae results in defective pruning of neurites and knock out of *Vps4a* in mice results in lethality at preweaning stages [17,18]. In line with the known function of VPS4A, fibroblasts from *VPS4A* human patients display defects in endosomal morphology and cell division [5]. How these defects lead to motor system impairment in probands is not fully understood.

Here, we identified a missense mutation in zebrafish *vps4a*, T248I, which was

generated in a small scale ENU mutagenesis screen for mutants that display auditory/vestibular

defects [19]. Zebrafish have become a model for studying human disease due to the genetic

conservation with humans, as well as the relatively short development and life cycle compared

to mammalian models, and the large number of available genetic tools and behavioral assays.

Brain development and structures within the zebrafish CNS use similar signaling pathways and

serve similar functions as those in mammals, making them particularly useful for studying

neurodevelopmental diseases [20–25].

We show that the zebrafish *vps4a* mutants display endosomal abnormalities and

reduced numbers of exosomes in the CNS. Both defects are consistent with the induction of

stress response genes in mid- and hindbrain regions of *vps4a^{T248I}* mutants. Using *in vivo*

calcium imaging, we detected strongly reduced activity in the motor circuits associated with

movements evoked by sensory inputs. These findings support a model where the reduced

enzymatic activity of Vps4a results in a disruption in late endosome function and exosome

release, consequently reducing cellular homeostasis within neural circuits mediating

sensorimotor transformation.

Results

A missense mutation in *vps4a* reduces ATPase activity and ESCRT-III filament

disassembly

raumschiff mutants were generated from an ENU mutagenesis screen for genes that affect

larval auditory and vestibular function [19]. Mutant larvae fail to respond to auditory cues and

maintain an upright posture. The mutation is recessive and fully penetrant at 5 days

postfertilization (dpf) with lethality occurring around 9-10 dpf. To identify the causative gene of

the *raumschiff* phenotype, we used whole-genome sequencing and identified a threonine to

isoleucine missense mutation in *vps4a* using SNPTrack [26]. The critical interval for *raumschiff*

was also mapped to a region approximately 700 kb (or 1 cM) south of the polymorphic marker

z5669, placing the locus at 34.33 Mb on Chromosome 25. *vps4a* is located at 34.93Mb on

Chromosome 25, which is within the predicted region for the causative gene. No other

polymorphisms were identified by SNPTrack in this region (Fig 1A).

Vps4a is a hexameric type I AAA-ATPase member of the ESCRT-III complex, whose

role is to promote the dissociation of the ESCRT-III subunits via ATP hydrolysis [11,13,27,28].

The threonine to isoleucine mutation (T248I) we identified is present in the AAA ATPase domain

in a region that is highly conserved between species (Fig 1B). To determine whether the T248I

mutation was comparable to a nonsense allele of *vps4a*, we edited the *vps4a* locus using

multiple CRISPR/Cas9 guides [29,30]. In our experiments, multi-guide injections resulted in a

G0 phenotype in 16.9% of injected larvae that was indistinguishable from the auditory/vestibular

phenotype observed in *vps4a*^{T248I} mutants (400 embryos were injected at < 1 hour past

fertilization; of the larvae that survived to 5 dpf, 56/331 larvae showed absence of acoustic

startle reflexes and abnormal posture). These results suggest that the T248I mutation is

comparable to a null allele of Vps4a. In addition, the *vps4b* paralogue is also ubiquitously

expressed [31], therefore we generated *vps4a/vps4b* G0 double mutants using CRISPR-based

editing to test whether *vps4b* could compensate for the loss of *vps4a* during development. A

previous study using morpholinos against *vps4b* reported a tooth phenotype analogous to

human dentin dysplasia I [31]. We observed that embryos injected with multiple CRISPR

guides against *vps4b* alone, or *vps4a* and *vps4b* together, developed normally until 3 dpf. The

lack of embryonic lethality is presumably due to the presence of maternal RNA during early

developmental stages. However, in contrast to the morpholino study, *vps4b* and *vps4a/4b* multi-

guide injected larvae exhibited progressive necrosis and by 5 dpf, there was complete lethality

(558 embryos were injected with *vps4b* guides; 468 larvae survived to 1 dpf, and 2 survived to 5

dpf. 298 embryos were injected with *vps4a/4b* guides; 236 larvae survived to 1 dpf, and 1

survived to 5 dpf). Our results suggest that *vps4b* is required for global cellular homeostasis

whereas *vps4a* selectively affects the nervous system. A similar effect was reported for the Nsfa

and Nsfb AAA-ATPases that mediate vesicle fusion; mutations in zebrafish *nsfa* largely affect

the nervous system, while the loss of *nsfb* functions leads to general necrosis at 4 dpf [32–34].

Based on the Cryo-EM structure of the yeast Vps4 AAA-ATPase [35], threonine 248 is

located at the interface between Vps4a hexamer subunits (Fig 1C). To assess the effect of the

isoleucine substitution on ATPase activity, we generated an equivalent mutation, T254I, in the

AAA ATPase domain of yeast Vps4. When tested for ATPase activity *in vitro*, Vps4^{T254I} had

severely reduced enzymatic activity compared to wild-type Vps4a, although some activity

remained ($p < 0.0001$, $n_{WT} = 4$, $n_{vps4(T254I)} = 4$, $t = 47.92$, df = 6) (Fig 1D).

To determine whether the residual function of the T254I protein could mediate filament

disassembly, we incubated Vps4^{T254I} with the ESCRT-III filaments formed by Vps2, Vps24 and

Snf7 *in vitro* and used sedimentation to assess disassembly. In our assay, disassembled

ESCRT-III filament proteins are present in the supernatant whereas assembled filaments

sediment to the pellet fraction [35]. We observed that wild-type Vps4a completely disassembled

the ESCRT-III filaments, with all proteins found in the supernatant (Fig 1E). In contrast, mutant

Vps4^{T254I} only partially disassembled the ESCRT filaments (Fig 1E). Using negative stain

electron microscopy (EM), assembled ESCRT-III filaments appear as rings or have corkscrew

structures (Fig 1F, upper left panel). When incubated with wild-type Vps4, filament structures

were absent, indicating full disassembly (Fig 1F, upper right panel). In the presence of Vps4^{T254I},

assembled filaments were still detectable albeit there were fewer filaments compared to

samples where no Vps4 protein was added (Fig 1F, lower panels). These results show that the

T245I mutation at the interface between Vps4a subunits reduces but does not fully abolish the

ATPase activity of the protein, and this reduction in activity is reflected in the incomplete

disassociation of ESCRT-III filaments.

The *vps4a^T248I* mutation results in enlarged endosomal membrane compartments in the

CNS

Using RNA-FISH, we found that the expression of *vps4a* transcripts is ubiquitous at 5 dpf, with

higher levels expressed in the CNS and retina (Fig 2A). In light of the higher expression of

vps4a in the CNS and the neurological disorder in human patients, we focused our attention on

the larval brain. As stated previously, Vps4a mediates the formation of multivesicular bodies

from late endosomes. To determine if the mutation affects this function in the nervous system,

we performed *in vivo* imaging of LysoTracker, a vital dye that accumulates in acidic membrane

compartments such as low pH endosomal compartments. As shown in Fig 2B, endocytic

compartments in *vps4a^T248I* mutants labeled more brightly with LysoTracker compared to wild-

type siblings. The average size and pixel density of acidic membrane compartments in the

hindbrain were quantified and we found that *vps4a^T248I* mutants have significantly larger acidic

membrane compartments ($p = 0.0015$, $n_{WT} = 7$, $n_{vps4a(T248I)} = 10$, $t = 3.873$, $df = 15$) (Fig 2C, left

panel) that take up more dye ($p = 0.0020$, $n_{WT} = 7$, $n_{vps4a(T248I)} = 10$, $t = 3.744$, $df = 15$) (Fig 2C,

right panel). Together, these results imply that the *vps4a^T248I* mutation has a detrimental effect

on endocytic compartment processing in the CNS.

In vivo* imaging reveals a reduction of circulating exosomes in the CNS of *vps4a^T248I

mutants

Given the known role of Vps4a in multivesicular body formation, we examined whether the

mutation affects the number of exosomes that are released upon fusion of multivesicular bodies

to the plasma membrane. To detect circulating exosomes in live fish, we used CD63-pHluorin,

a pH-sensitive GFP reporter that is quenched in intraluminal vesicles in low pH multivesicular

bodies and fluoresces when these vesicles are released into a higher pH extracellular milieu as

exosomes [36]. We microinjected a plasmid for ubiquitous expression of CD63-pHluorin into

single-cell stage embryos, and selected for larvae that transiently expressed the CD63-pHluorin

based on signal in the muscle cells, which is readily detectable by 3 dpf (Fig 3A). The plasma

membrane of muscle cells was commonly labeled in our injected fish as was previously reported

[36] and served as a positive control for expression. We detected a strong reduction in CD63-

phluorin signal in the CNS blood vessels on 3 dpf (Fig 3B), as well as in the midbrain ventricle of

4 dpf $vps4a^{T248I}$ mutants (Figs 3C,D). We quantified the CD63-pHlorin signal in the midbrain

ventricle and normalized to background signal from the surrounding CNS tissue. The

relationship between CD63-pHluorin signal in the midbrain ventricle is moderately correlated to

signal in the muscle cells in wild-type siblings but not in mutants (Fig 3E). We then calculated a

ratio of the normalized pixel intensity of the midbrain ventricle CD63-pHluorin signal to the

muscle cell CD63-pHlourin signal and found that the ventricle/muscle cell intensity ratio is

significantly decreased in $vps4a^{T248I}$ mutants compared to wild-type siblings (p = 0.0382 (exact),

n_{WT} = 16, $n_{vps4a(T248I)}$ = 8, Mann-Whitney test) (Fig 3F). This result indicates that there is a

decrease in the number of exosomes present in the brain ventricle of mutants, supporting the

hypothesis that biogenesis of these extracellular vesicles is reduced by the $vps4a^{T248I}$ mutation.

Transcriptional changes in the $vps4a^{T248I}$ mutant

Considering the broad changes in endosomal morphology and the reduction in circulating

exosomes in the CNS of $vps4a^{T248I}$ mutants, we next examined the gross morphology of the

brain. Using whole mount immunohistochemistry, we labelled the axonal fibers of mutants and

siblings with acetylated tubulin antibody to visualize axon morphology. No obvious changes in

gross morphology between wild-type siblings and $vps4a^{T248I}$ mutants were apparent (Fig 4A).

However, we noted the presence of pyknotic nuclei (Fig 4B), which are associated with cell

death [37]. We quantified the number of these abnormal cells and found that $vps4a^{T248I}$ mutants

have significantly more abnormal cells in the CNS compared to wild-type siblings (p < 0.0001

(exact), n_{WT} = 9, $n_{vps4a(T248I)}$ = 11, Mann-Whitney Test).

As exosomes carry cargo containing microRNA and DNA and have been implicated in

the regulation of DNA transcription [38], we assayed for transcriptional changes in the $vps4a^{T248I}$

mutant by performing a differential gene expression analysis on a bulk RNAseq data set that

was initially used to find the mutation. We then used HCR RNA-FISH to validate the top $\log_2$

fold changes in gene expression. We found that the immediate early response genes *atf3* and

jun, and the axonal regeneration gene *gap43* were dramatically upregulated in mid- and

hindbrain regions in the $vps4a^{T248I}$ larvae (Fig 4C, D). These striking changes in gene

expression are consistent with a failure in intercellular communication by exosomes, and/or

endosomal processing within neurons.

Impaired vision in $vps4a^{T248I}$ mutants

The clinical profile of human patients with missense mutations in *VPS4A* includes visual deficits

with a range of symptoms such as retinal dystrophy and cataract formation [5,7]. Although we

did not observe expanded melanophores in the $vps4a^{T248I}$ mutant, which are indicative of

blindness in larvae, we did find pronounced changes in visual function and gene expression in

the retina. The optomotor response (OMR) in zebrafish is a reflex that can be evoked by a

visual stimulus such as moving stripes, which causes the fish to swim in the direction of

perceived motion, resulting in stabilization relative to the environment [39]. Compared to wild-

type siblings, $vps4a^{T248I}$ mutant larvae have a greatly reduced OMR, suggesting they are blind

($p < 0.0001$ (exact), n_{WT} = 12, $n_{vps4a(T248I)}$ = 10, Mann-Whitney test) (Fig 5A). To investigate this

defect further, we recorded electroretinograms (ERG) to assess the light induced changes in the

electrical potential of the whole retina. The b-wave amplitude generated from an ERG can

provide insight into the function of the outer retina [40]. A range of light intensities (from the

dimmest at log-4, to the brightest at log0) were tested and we found that the b-wave amplitude

in $vps4a^{T248I}$ mutants is significantly decreased compared to wild-type siblings (log 0, $p <$

0.0001, n_{WT} = 23, $n_{vps4a(T248I)}$ = 34; log -1, p < 0.0001, n_{WT} = 23, $n_{vps4a(T248I)}$ = 33; log -2, p <

0.0001, n_{WT} = 23, $n_{vps4a(T248I)}$ = 34; log -3, p = 0.0101, n_{WT} = 23, $n_{vps4a(T248I)}$ = 34; log -4, p =

0.9527, n_{WT} = 23, $n_{vps4a(T248I)}$ = 34, Dunn's multiple comparisons test) (Fig 5B). In addition, we

found that *jun* and *gap43* are upregulated in the retina of *vps4a^T248I^* mutants, specifically in the

photoreceptor and retinal ganglion layers (Fig 5C). We also detected reduced *opn1sw1*

expression in the photoreceptors of mutants. Together, these analyses indicate that the

*vps4a^T248I^* mutants have pronounced visual defects originating in the retina. Notably, the

decrease in b-wave amplitude is variable among individual mutants and may not fully explain

the absence of the OMR.

Selective vestibular defects in *vps4a^T248I^* mutants

At the free-swimming stage of development (5 dpf), *vps4a^T248I^* larvae do not maintain an upright

posture, swim sideways or upside down, and do not inflate their swim bladder, which is

classically associated with functional deficits in vestibular hair-cells [41]. To further investigate

this phenotype in *vps4a^T248I^* mutant larvae, we tested vestibular function by assessing vestibular

induced eye movements (VIEM) [42]. Larvae with loss of hair-cell function mutations have

reduced eye movements in response to head rotation [41], however, we did not observe a

significant difference between the *vps4a^T248I^* mutants and wild-type siblings (p = 0.4632 (exact),

n_{WT} = 18, $n_{vps4a(T248I)}$ = 16, Mann-Whitney test) (Fig 6A). As mutant larvae have pronounced

postural defects, we also tested the vestibulospinal reflex (VSR), which assesses vestibular

induced motor function in the trunk [43]. We compared the maximum tail angle between wild-

type siblings and mutants and found the *vps4a^T248I^* mutants had a significantly reduced tail angle

(p = 0.0012 (exact), n_{WT} = 7, $n_{vps4a(T248I)}$ = 10, Mann-Whitney test) (Fig 6B). The normal VIEM

along with the reduced VSR result suggests that *vps4a^T248I^* mutants have selective defects in

vestibular pathways.

264   In zebrafish, acceleration of the head is detected by utricular hair cells [42,44,45], which

265 then transmit signals to the utricular afferents of the VIIIth cranial nerve (Fig 6C). The VIIIth

266 nerve innervates the superior vestibular nucleus (SVN), vestibulospinal cell (VS) and tangential

267 nucleus (TAN), as well as the Mauthner cell in the hindbrain [46–49]. Signals from the SVN and

268 TAN are relayed by the oculomotor and trochlear nuclei to oculomotor muscles to produce eye

269 movements associated with the VIEM (Fig 6C;[46,48,50]). VS neurons directly innervate motor

270 neurons in the trunk to produce tail movements [48,50]. To assess the function of the VSR

271 circuit, we used a pan-neuronal GCaMP7a calcium indicator to visualize neuronal activity [51].

272 Recently it was demonstrated that utricular hair cells in zebrafish larvae respond to high

273 frequency tones, analogous to the responses of vestibular hair cells to bone-conducted vibration

274 in mammals [52,53]. To stimulate the vestibular circuitry mediating the VIEM and VSR, we

275 delivered pulses of 600 Hz at 110 dB using a mini-speaker attached to the glass slide on which

276 larvae were mounted for calcium imaging. Electron microscopy data has shown that the lateral

277 dendrite of the Mauthner cell is adjacent to the VS nucleus [48]. To visualize this dendrite, we

278 performed backfill injections of rhodamine dextran prior to calcium imaging, enabling us to use

279 the Mauthner cell as a landmark for the approximate locations of the SVN, VS and TAN (Fig

280 6D). Upon tone stimulation, neurons in all three nuclei responded with an increase in

281 fluorescence (Fig 6D'). Nevertheless, the average raw fluorescence trace for the VS region has

282 a slightly lower peak amplitude in the $vps4a^{T248I}$ mutant compared to wild-type siblings (Fig 6E,

283 right). Analysis of the group data from individual fish indicates that the percentage change in

284 normalized fluorescence in the VS region is reduced in $vps4a^{T248I}$ mutants (p = 0.0337 (exact),

285 n_{WT} = 11, $n_{vps4a(T248I)}$ = 15, t = 2.252, df = 24)(Fig. 6F, right). As predicted by our behavioral

286 analysis, we did not observe a significant difference in the raw fluorescence trace or percentage

287 change in normalized fluorescence in the TAN or SVN between $vps4a^{T248I}$ mutants and wild-

288 type siblings (TAN p = 0.6224 (exact), SVN p = 0.1405 (exact), n_{WT} = 11, $n_{vps4a(T248I)}$ = 15, t =

289 0.4988 (SVN), df = 24 (SVN), t = 1.524 (TAN), df = 24 (TAN)) (Figs 6E,F, left and center).

Overall, our calcium imaging data confirm that the VIEM circuit is intact and suggest that

reduced activity in the VS neurons, which directly innervate motor neurons, contributes in part to

the reduction in tail movements in response to vestibular cues in the $vps4a^{T248I}$ mutants.

Intact ascending auditory pathway in $vps4a^{T248I}$ mutants

In zebrafish, auditory signals excite saccular hair cells, which are innervated by the afferent

neurons of the statoacoustic ganglion (or VIIIth nerve). If the auditory cues are sufficiently loud

enough to startle, the signal is transmitted to the giant escape fiber neuron known as the

Mauthner cell, which directly activates motor neurons[54–56]. Less intense sounds activate

hindbrain neurons within the medial octavolateralis nuclei region, which then communicate with

the torus semicircularis for further auditory processing [45,57–59] (Fig 7A). $vps4a^{T248I}$ mutant

larvae are largely unresponsive to tapping on the petri dish, suggesting that hearing is

profoundly impaired. We assessed the severity of this defect by quantifying the auditory evoked

behavioral response (AEBR), which is a startle response to loud, pure tone stimuli. $vps4a^{T248I}$

mutant larvae responded to pure tones significantly less compared to wild-type siblings,

indicating that mutants have pronounced defects ($p < 0.0001$ (exact), $n_{WT} = 15$, $n_{vps4a(T248I)} = 20$,

Mann-Whitney test) (Fig 7B). The pan-neuronal GCaMP7a calcium indicator was used to

examine neuronal activity in the saccular hair cells and statoacoustic ganglion in response to

the same 110 dB 600 Hz stimulus (Figs 7C,D). In saccular hair cells, we observed robust

responses in average raw fluorescence traces (Fig 7E) and comparable calcium transients in

wild-type or mutant individual fish ($p = 0.4162$ (exact), $n_{WT} = 20$, $n_{vps4a(T248I)} = 13$, $t = 0.8228$, df =

35) (Fig 7F). Comparable responses were also seen in the statoacoustic ganglion ($p = 0.8748$

(exact), $n_{WT} = 20$, $n_{vps4a(T248I)} = 13$, $t = 0.1588$, df = 31) (Figs 7E,F). These results are consistent

with LysoTracker labeling, which shows no significant difference in dye uptake between wild-

type siblings and $vps4a^{T248I}$ mutant in hair cells in the saccule (S1 Figs A-D; $p = 0.0934$ (exact),

n_{WT} = 6, $n_{vps4a(T248I)}$ = 9, t = 1.810, df = 13), utricle (p = 0.6314 (exact), n_{WT} = 6, $n_{vps4a(T248I)}$ = 9, t =

0.4913, df = 13) or anterior cristae (p = 0.3233 (exact), n_{WT} = 10, $n_{vps4a(T248I)}$ = 10, t = 1.016, df =

18). We also examined the saccular hair cells and statoacoustic ganglion for changes in gene

misexpression but did not detect transcriptional changes in these tissues (S1 Fig E).

The auditory circuitry within the hindbrain for non-startle responses is less well defined

in larvae, therefore we focused on a midbrain region of the ascending auditory pathway and

examined neuronal activity in the torus semicircularis, which is akin to the inferior colliculus in

mammals and key for auditory processing [57,60,61]. This midbrain structure is located

beneath the optic tectum at the level of the semicircular canals and saccule of the inner ear and

the medial longitudinal fasciculus (MLF) (Fig 7G). We observed a visible increase in GCaMP7a

reporter activity in the torus semicircularis of both wild-type siblings and $vps4a^{T248I}$ mutants after

pure tone stimulation (Fig 7H). No significant differences were detected in the average raw

fluorescence traces or percentage change in normalized fluorescence between wild-type

siblings and $vps4a^{T248I}$ mutants for the torus semicircularis (p = 0.9672 (exact), n_{WT} = 11,

$n_{vps4a(T248I)}$ = 13, t = 0.04157, df = 22) (Figs 7I,J). Together, these data indicate that the

ascending auditory pathway of $vps4a^{T248I}$ mutants including the inner ear and central circuitry is

functional.

Defective sensorimotor transformation in $vps4a^{T248I}$ mutants

Although $vps4a^{T248I}$ mutants have a greatly reduced acoustic startle response, our data

demonstrate that auditory cues activate the ascending auditory pathway. Notably, mutant larvae

are not paralyzed, however, they are largely inactive at the free-swimming stage. Therefore, we

assessed whether $vps4a^{T248I}$ mutants have a general motor system defect that prevents motor

responses to sensory cues.

339   To test whether startle reflexes are globally affected, we measured the ability of

340 *vps4a^{T248I}* mutants to produce a deep C-bend of the trunk in response to contact of a stiff probe

341 with the dorsal region of the head. We observed that both wild-type siblings and *vps4a^{T248I}*

342 mutants displayed robust C-bend motions of their trunks in response to touch (Figs 8B,C; p =

343 0.1079 (exact), n$_{WT}$ = 10, n$_{vps4a(T248I)}$ = 11, t = 1.687, df = 19). This result suggests that touch

344 receptors, along with Mauthner cells and other vital components of motor output such as spinal

345 cord neurons, motor neurons, and muscle cells are functional. In addition, axonal fibers in

346 superficial layers of the skin, the hindbrain, and anterior trunk regions did not show any changes

347 in gross morphology in *vps4a^{T248I}* mutants (Fig 8D). These data point to a deficit within central

348 circuitry in *vps4a^{T248I}* mutants as downstream motor components of the startle reflex are

349 unaffected.

350   As described above, auditory signals from the statoacoustic ganglion are directly

351 transmitted to the Mauthner cell, which innervates trunk motor neurons to generate an escape

352 reflex. However, the 110 dB intensity of our tone stimulus did not result in detectable activation

353 of the Mauthner cell soma, which is consistent with a previous imaging study [62]. Instead, we

354 observed a reliable response in the midbrain nucleus of the medial longitudinal fasciculus

355 (nMLF) (Figs 8A, E-H). The nMLF is a vital component of the descending motor system, serving

356 as a command center for forward locomotion [63–70]. The nMLF contains subpopulations of

357 vGlut1 and vGlut2 positive neurons, which control both fast escape-swimming behaviors as well

358 as slow forward locomotion [68,71]. Neurons in the nMLF were recently shown to respond to

359 hindbrain cranial relay neurons (CRN; Fig 8A), which can be activated by aversive auditory

360 stimulation of Mauthner cells, leading to smooth swimming after an initial C-bend response [63].

361 Although the auditory circuitry to the nMLF activated by non-startle stimuli has not been

362 characterized, the nMLF has been implicated in fine postural control via inputs from the TAN of

363 the ascending vestibuloocular pathway [50]. In *vps4a^{T248I}* mutants, activation of neurons in the

364 TAN is comparable to wild-type responses.

Considering that the nMLF is central to motor function in response to sensory cues, we

compared GCaMP7a signals in this cluster of neurons between wild-type siblings and $vps4a^{T248I}$

mutants in response to the 600 Hz stimulus. These midbrain clusters can be identified based on

their prominent axonal tracts that descend near the midline (Fig 8E). We found that the peak

amplitudes of calcium transients in the average raw fluorescence trace in the mutant were

markedly decreased (Figs 8F-G). In addition, the percentage change in normalized

fluorescence in the nMLF was significantly reduced in all individual $vps4a^{T248I}$ mutants tested (p

< 0.0001 (exact), n_{WT} = 14, $n_{vps4a(T248I)}$ = 11, t = 6.937, df = 23) (Fig 8H).

We next investigated the neuronal activity of the spinal cord in response to tone

stimulation using the pan neuronal GCaMP7a reporter (Fig 8I). The average raw fluorescence

trace of the $vps4a^{T248I}$ mutants show lower peaks in response to tone stimulation (Fig 8K). In the

examples shown in Figure 8J, a second reporter (GCaMP7a expressed under a muscle cell

promoter, [72]) in the trunk muscles shows activation in wild-type larva but none in the mutant.

Overall, the percentage change in normalized fluorescence is strongly reduced in $vps4a^{T248I}$

mutants (p < 0.0001 (exact), n_{WT} = 11, $n_{vps4a(T248I)}$ = 9, Mann-Whitney test) (Fig 8L). Together,

our results indicate that tone stimulation fails to trigger responses in the nMLF descending

motor pathway and downstream trunk motor neurons in $vps4a^{T248I}$ mutants.

Discussion

Here we present a T248I mutation in $vps4a$, which causes sensorimotor defects in zebrafish

larvae. Spanning biochemical to systems level analyses, our study indicates that a reduction in

Vps4a enzymatic activity leads to endosomal defects and reduced exosome biogenesis,

resulting in pronounced deficits in vision and sensorimotor transformation of auditory and

vestibular cues. We suggest that the mechanisms underlying this type of dysfunction in

zebrafish may also contribute to the condition seen in human patients with *de novo* mutations in

the human *VPS4A* orthologue.

Defective disassembly of membrane scission filaments and decreased exosome

biogenesis

395   Vps4a plays a key role in ESCRT-III filament disassembly, and the energy required for

this function is produced through the ATP hydrolysis activity of Vps4a [14,27,28,73]. Using an

equivalent mutation in yeast, T254I, the threonine to isoleucine substitution in the ATPase

domain of yeast *Vps4* results strongly reduced ATPase activity and ESCRT-III filament

disassembly. Studies of an ATP hydrolysis deficient mutant, Vps4^{E233Q}, reveal that a complete

loss of ATP hydrolysis abolishes filament disassembly [74]. In contrast, the yeast protein with

the T254I mutation still retains a low level of activity. Nevertheless, our *vps4a* CRISPR

experiments suggest that the missense mutation is comparable to a null allele, where we

observe a similar auditory/vestibular phenotype in multi-gRNA injected fish without gross

developmental defects. These observations suggest that select neural circuits are highly

sensitive to perturbation of Vps4a activity.

406   ESCRT-III filament disassembly is a key step in membrane scission, which is vital to

cellular mechanisms of multivesicular body formation, exosome biogenesis, cytokinesis and

neurite severing [75]. Multivesicular bodies form when late endosomal membrane buds inward,

and membrane scission that allows the separation of these buds from the membrane results in

intraluminal vesicles (ILVs) [14,38,76,77]. The resulting multivesicular bodies can subsequently

either fuse to the plasma membrane to release the ILVs into the extracellular matrix, or fuse with

lysosomes to promote degradation of its contents [78,79]. Both fates of ILVs are key for

maintaining cell homeostasis and signal transduction [80,81]. An early study of protein sorting

mutants yeast proposed a model where in Class E *vps* mutant cells, there is a pre-vacuolar or

endosome-like organelle that accumulates a quinacrine vital dye because of defective protein

trafficking [82]. It has since been established that expression of a dominant negative Vps4, a

class E *vps*, or depletion of other ESCRT components, inhibits the formation of multivesicular

bodies and the proper sorting of cargo into intraluminal vesicles [83–86]. Using *in vivo* imaging,

we observed an increase in LysoTracker labeling in the central nervous system of $vps4a^{T248I}$

mutants. This accumulation of dye and enlargement of acidic membrane compartments

suggests a failure of invagination of the membrane of endosomes, resulting in endosomal

compartments with larger surface areas, a similar phenotype to what was reported in yeast

mutants and in human patient fibroblasts [5].

ILVs in multivesicular bodies serve as a source of exosomes and are secreted from

nearly all cell types and contain a variety of signaling molecules, such as proteins, lipids, and

nucleic acids [87–90]. Exosomes have been implicated in a variety of diseases such as various

cancers and neurodegenerative diseases due to the vital role exosomes play in intercellular

communication [77,91–95]. A previous study on erythroid cells, which found that transferrin

receptor trafficking, a process dependent on proper endosomal sorting and exosome secretion,

is disrupted in human patients with *de novo VPS4A* mutations [7]. Using a transiently

expressed CD63-pHluorin marker to visualize circulating exosomes in live fish, we found a

significant reduction in CD63-pHluorin positive exosomes in the CNS blood vessels and

hindbrain ventricle of $vps4a^{T248I}$ mutants, suggesting a defect in biogenesis. Disruptions to

intercellular signaling through exosomes can manifest as transcriptional changes [79,96,97].

Indeed, we found that several genes, either associated with transcription (*atf3* and *jun*) or

regeneration (*gap43*), are highly upregulated in the hindbrain, midbrain and retina of $vps4a^{T248I}$

mutants. Both *atf3* and *jun* are known to be stress response genes, whereas upregulation of

gap43 may be a compensatory response to disruption of axonal processes. Although exosomes

mediate cell-to-cell signaling and are important for cellular homeostasis, it is not clear whether

the transcriptional changes in $vps4a^{T248I}$ mutants are due to (*i*) reduced exosome signaling, or

(*ii*) disruptions in endosomal trafficking, or (*iii*) a combination of both defects. Alternatively, other

disruptions to membrane scission in the CNS may be involved as well. The expression of stress

response genes in the CNS in zebrafish mutants during larval development is consistent with

the microcephaly seen in human patients. While we did not observe changes in gross

morphology of the CNS, cell death in the brain was evident yet sparse at the larval stage shown

here. We speculate that more pronounced defects in cellular homeostasis would be evident in

older animals, however, we limited our analyses to early stages of larval development during

which we could discern early onset defects.

Sensory and motor deficits

Patients with *VPS4A* mutations are reported to have sensory deficits relating to vision

[5]. Poor vision is a common feature that can be associated with retinal dystrophy and

cataracts. With respect to visual function in the zebrafish mutant, larvae fail to swim in response

to moving visual cues and outer retinal responses to light are significantly reduced, signifying

defects in vision that align with transcriptional changes in the retina. Nevertheless, the

reduction in outer retinal function was variable among mutants, yet the optomotor response was

severely reduced in all *vps4a*^{T248I} larvae tested. These results suggests that the loss of

photoreceptor function provides only a partial explanation of the absence of motor reflexes

associated with vision and is indicative of an inability to transform visual cues into motor output

in a subset of mutants. Notably, human patients are reported to have difficulties in fixing and

following a moving object. This issue with visual fixation may be due to poor visual acuity or a

combination of poor vision and oculomotor pathway dysfunction.

Loss of central control of muscle tone, ataxia and motor delays are common traits found

in patients with neurodevelopmental disorders, including *VPS4A* patients [5,7]. The *vps4a*^{T248I}

phenotype shares some features with these disorders in that mutant larvae lack control over

posture and have sensorimotor defects. Vestibulospinal reflexes, which result in corrective

movements of the trunk, are greatly diminished in mutants. However, not all vestibular function

in lost; $vps4a^{T248I}$ mutants still produce eye movements in response to rotation of the head.

Consistent with the behavioral data, neurons of the TAN and SVN, which are part of

vestibuloocular pathway [46,48], show comparable activity in the $vps4a^{T248I}$ mutant to wild-type

neurons. This data indicates that perturbation of Vps4a activity does not result in global defects

in vestibular function, or in brain function in general, but rather that specific circuits are more

sensitive to mutations in $vps4a$. Normal vestibular induced eye movements and activation of

TAN and SVN neurons also indicate that excitation of utricular hair cells and vestibular afferent

neurons is unaffected in $vps4a^{T248I}$ mutants. In contrast to the vestibuloocular pathway, the

vestibulospinal circuit is impaired when $vps4a$ is mutated. Hindbrain VS neurons, which directly

innervate trunk motor neurons, have decreased neuronal activity in mutants. Recently it was

shown that ablation of VS neurons leads to modest defects in posture in zebrafish larvae [98],

suggesting that other vestibular circuits contribute to postural control. The reduced activation of

the vestibulospinal pathway in $vps4a^{T248I}$ mutants may therefore only partially account for the

inability to maintain an upright posture.

Further evidence for a loss in central control of motor responses emerged from our

analysis of the absence of an acoustic startle reflex in $vps4a^{T248I}$ mutants. Despite the lack of

response to aversive acoustic stimuli, $vps4a^{T248I}$ mutants have comparable activation of the

ascending auditory pathway as seen in wild-type siblings. Deficits in the descending motor

pathways or the spinal cord are also not the cause for the absence of the motor output in

response to visual or tone stimulation. $vps4a^{T248I}$ mutants are not paralyzed and aversive touch

stimuli activate escape responses, demonstrating that mutant larvae can generate robust tail

movements. The above experiments indicate that both saccular afferent neurons and Mauthner

cells are activated by sensory cues in $vps4a^{T248I}$ mutants, implying that a second pathway may

be defective in generating acoustic startle reflexes. One such pathway may involve hindbrain

spiral fiber neurons, which have been shown to be essential for Mauthner cell-dependent

escape responses [99]. Nevertheless, our *in vivo* imaging experiments using a non-startle tone

stimulus revealed a pronounced reduction in the activity of midbrain nMLF neurons and

downstream spinal cord neurons of $vps4a^{T248I}$ mutants. Neurons of the nMLF play a key role in

receiving sensory input and sending motor commands to the spinal cord and are thought to play

an analogous role to the interstitial nucleus of Cajal in the mammalian midbrain, which controls

eye and head movements [66,68,100]. In zebrafish, both visual and vestibular systems provide

sensory input to the nMLF [46,101,102]. Interestingly, TAN neurons of the vestibuloocular

pathway, also play a dual role of controlling eye and body movements in zebrafish. TAN

neurons, which project to the nMLF [46], were recently shown to transmit signals about the roll

axis of the body to the nMLF, implicating this circuit in fine control of posture [50]. In $vps4a^{T248I}$

mutants, TAN neurons are also activated by the tone stimulus, yet postural defects are

pronounced, further bolstering the notion that sensory input such as vestibular cues fail to

trigger sensorimotor transformation within the nMLF.

In summary, our combined biochemical and *in vivo* analyses have revealed that loss of

Vps4a function generates sensorimotor processing defects in the CNS of zebrafish larvae. Our

study provides a better understanding of the cause of motor dysfunction when *vps4a* is mutated

and has implications for potential therapies as well. In emerging work on exosomes as

therapeutical reagents, these extracellular vesicles, which can cross the blood brain barrier, can

serve as both a diagnostic tool and as a potential vehicle for delivery [103]. Recent studies

have shown that exosome therapy can aid in sensorimotor recovery after injury or resulting from

acute or chronic neurodegenerative diseases [104–106], thus there is potential for exosomes to

be used in early neurodevelopmental disorders such as those associated with *VPS4A*.

Materials and Methods

Zebrafish care and use

We maintained zebrafish lines for the $vps4a^{T248I}$ allele in the Top Long Fin (TLF) wild-type (WT) or $mitfa^{b692}$ nacre backgrounds. The $vps4a^{T248I}$ Tg(α-tubulin-gal4;MP-gal4;cmlc2-GFP;UAS-GCaMP7a) line was generated by crossing into a $mitfa^{b692}$ Tg(α-tubulin-gal4;MP-gal4;cmlc2-GFP;UAS-GCaMP7a) fish line (a gift from Philippe Mourrain). The MP-gal4 transgene was present in most fish, marking the muscle cells as well. Breeding stocks were housed at 28.5°C and animal husbandry followed standard zebrafish methods for laboratory utilization [107], as approved and overseen by the Institutional Animal Care and Use Committees at both Oregon Health and Sciences University and Stanford University. The experiments used zebrafish larvae 3-7 days post fertilization (dpf) before gender differentiation occurs. Embryos and larvae were raised in E3 medium (0.33 mM CaCl2, 0.17 mM KCl, 0.33 mM MgSO4, and 5 mM NaCl) incubated at 28.5°C. When appropriate, larvae were anesthetized in E3 with 0.03% 3-amino benzoic acid ethylester (MESAB, Western Chemical) to minimize pain and distress.

Genotyping Methods

For the $vps4a^{T248I}$ allele, we genotyped fish and larvae using kompetitive allele-specific PCR (KASP) genotyping (LGC Biosearch Technologies) per the manufacturer's instructions.

Identification of raumschiff mutant allele

Positional cloning was performed using simple sequence length polymorphism (SSLP) markers mapped to the zebrafish genome [108]. 5 dpf zebrafish larvae were sorted by phenotype.

For Whole Genome Sequencing, 5 dpf zebrafish larvae were sorted into separate pools of 20 homozygous mutants and 20 heterozygous and homozygous wild-type siblings. Genomic DNA was collected using a DNeasy Blood & Tissue Kit (Qiagen) and samples were processed by Azenta Life Sciences (formerly GENEWIZ) for library preparation and Illumina HiSeq

sequencing. The resulting .gz files were uploaded to SNPTrack

(http://genetics.bwh.harvard.edu/snptrack/) to identify the mutation.

CRISPR injections

Single guide RNAs against *vps4a* and *vps4b* were designed using the CHOPCHOP website

(https://chopchop.cbu.uib.no/) and synthesized (Alt-R CRISPR-Cas9 System, IDT). Three

guides were chosen based on if they targeted the ATPase domain or additional exons common

to all isoforms, low number of off target sites and efficiency. Multi-guide CRISPR injections

were performed as described previously [30]. Briefly, crRNA and tracrRNA guides were

combined in equal molar amounts in Duplex Buffer (IDT), and this RNA Duplex mix was

combined with 1 µg/µL Cas9-NLS protein (PNA Bio). 1 nL of RNA guide-CAS9 mixture was

injected into single cell stage TLF wildtype embryos. Embryos were checked daily for viability

and auditory/vestibular phenotypes scored on 5 dpf.

Protein expression and purification

Saccharomyces cerevisiae Vps4 (wild-type), and mutant Vps4(T254I) were cloned into the

pET28a vector with an N-terminal His6 tag. The two proteins were expressed in Escherichia coli

BL21 (DE3) strains and induced with 1 mM IPTG at 20°C overnight. Sonication was used to lyse

cells and centrifuged the lysates. The supernatant was loaded into Ni–NTA beads (GE

Healthcare) and the eluates were incubated with 5 U mL−1 apyrase (Sigma-Aldrich) at 4°C

overnight. All proteins were further purified by SEC using a Superdex 200 10/300 GL column

(GE Healthcare) with 20 mM Tis-HCl at pH 8.0, 150 mM NaCl buffer. The purity and quality of all

the proteins were tested by SDS–PAGE and Nano-drop (Implen).

Saccharomyces cerevisiae Snf7 mutant (R52E) and Vps24 were cloned into the pET28a

vector with an N-terminal His6 tag. The two proteins were expressed in Escherichia coli BL21

(DE3) strains and induced with 1 mM IPTG at 20°C overnight. Sonication was used to lyse cells

and centrifuged the lysates. The supernatant was loaded to a Ni–NTA beads (GE Healthcare)

and the elute was further purified by SEC using a Superdex 200 10/300 GL column (GE

Healthcare) with 20 mM Tis-HCl at pH 8.0, 150 mM NaCl buffer. The purity and quality of all the

proteins were tested by SDS–PAGE and Nano-drop (Implen).

Saccharomyces cerevisiae Vps2 were cloned into the pET28a vector with an N-terminal

MBP-His6 -TEV-tag. We expressed the resulting Vps2 construct in E.coli BL21 (DE3) at 20°C

for 16 hours with 1 mM IPTG. We purified the protein using Ni–NTA resin (GE Healthcare). We

mixed the eluted fractions from the His column with TEV (tobacco etch virus) protease for 3

hours at 4°C. We then loaded the cleaved samples onto a 5 mL HiTrap Q-sepharose FF column

(GE Healthcare) and eluted the material with a gradient of 0%-100% buffer (20 mM Tris-HCl, pH

8.0, 500 mM NaCl) in 50 column volumes.

Sedimentation Analysis of Filament Disassembly

Yeast ESCRT III filaments were assembled by mixing Snd7(R52E), Vps24 and Vps2 in a

molar ratio of 2:1:1 at room temperature for 1 hour. Then, the formed filaments were centrifuged

at 50,000 r.p.m. for 30 minutes in a TLA-100 ultracentrifuge rotor (Beckman Coulter). The pellet

was resuspended in the buffer containing 20 mM Tris-HCl at pH 8.0, 150 mM NaCl. The pellet

(P) and supernatant (S) were analyzed by SDS–PAGE.

For sedimentation analysis of ESCRT III filament disassembly, filaments were mixed

with 5 μM Vps4 or Vps4(T254I) in 20 mM Tris-HCl at pH 8.0, 150 mM NaCl, 2 mM MgCl2 buffer,

separately. Reactions were started by adding ATP to a final concentration of 5 mM. After

incubation at room temperature for 10 minutes, reactions were stopped by adding EDTA to a

final concentration of 50 mM. Samples were then subjected to ultracentrifugation and analyzed

via SDS-PAGE and negative staining.

ATPase activity assay

ATPase activities were determined for wild-type and mutant Vps4 proteins using the

Quantichrome ATPase/GTPase Assay Kit (BioAssay Systems) at room temperature. Briefly,

5 µM Vps4 or mutant Vps4(T254I) protein was added to ATPase assay buffer and 4 mM ATP for

30 minutes, and then stopped using reagent from the kit. The solution was incubated at room

temperature for 30 minutes, and immediately the absorbance at 620 nm was detected using a

plate reader (Enspire, Perkin Elmer). The released phosphate was calculated based on the

absorbance standard curve established using KH_2PO_4 standards.

Auditory-evoked behavioral response (AEBR)

AEBR was tested as described previously [109,110] using the Zebrabox system (ViewPoint Life

Sciences). 5 dpf wild-type siblings or *vps4a*^{T248I} mutant larvae were confined to wells of a 96-

well plate in 200 µL E3 per well, in the dark. Larvae were subjected to 100 millisecond, 1kHz

pure tones at 157 decibel sound pressure (dB SPL, relative to 1mPA) every 2 seconds for 1

minute. An infrared camera recorded larval movements and ZebraLab tracking software

documented pixel changes, which represent movement over time for each larva. To distinguish

between evoked responses and spontaneous background movement, movements within 1

second before a stimulus were excluded from a given trial. For each larva, the positive

responses were calculated as a percentage out of total possible response within a minute, and

the best of three trials was used for analysis.

Vestibulospinal Reflex (VSR)

The device and method used to test the VSR was done as previously described [43,44]. Briefly,

5 dpf larvae were embedded dorsal-up in 2% prewarmed low melt agarose in E3 media without

anesthetic on a mounting chamber. The agarose around the trunk and pectoral fins was

carefully removed and replaced with 20 µL E3 media, and the mounting chamber was placed in

the device. A motor rocks the platform from −75° to + 75° at 0.53 Hz, and an infrared camera

recorded video of the reflex at 30 fps (1280 frames).

Modified ZebraZoom software [111] tracks the movements of the trunk at regular

intervals and calculates the real tail angle over time between the X axis and the axis formed by

the tip of the tail and the center of the swimming bladder.

Vestibular-induced eye movement (VIEM)

The VIEM experiment was done similarly as described previously using a modified device

[42,44]. 5 dpf larvae were embedded on a small cover slip in prewarmed 2% low melt agarose

in E3 buffer, without anesthetic. Agarose surrounding the eyes was gently removed and

replaced with approximately 10 µL of E3 media to permit free eye moment. The mounted larvae

were positioned vertically, head down, between an infrared camera and an infrared LED array

on a programmable rocking platform. Larvae kept in darkness during the experiment. The

platform rocked (-45° to 45°) at 0.25 Hz frequency for 60 seconds while larval eye movements

were video recorded in infrared.

Software-based video analyses quantified the amplitudes of angular eye movements as

a function of their frequency of repetition. Larvae with normal VIEM show peak movement

amplitude at 0.25 Hz, in synchrony with platform rotation. Baseline spontaneous eye movement,

taken as the average amplitude across all quantified frequencies, was subtracted to determine

the VIEM amplitude at 0.25 Hz. We calculated the normalized amplitude of the eye ratio for

each larva and used the two-tailed unpaired t test with Welch's correction to compare mutants

to their corresponding siblings.

Touch Startle Reflex

Touch Startle was performed by mounting 7 dpf zebrafish larvae similarly to the VSR mounting

method. A micromanipulator holding a borosilicate glass capillary tube (Sutter, #BF150-86-10)

pulled into a needle was used to touch the head of the larvae. Startle movements were

recorded by a Hamamatsu C11440 ORCA-Flash 2.8 camera at 100 fps. Quantification of C-

bends was performed as described in [112].

Optomotor Reflex (OMR)

To test for the optomotor reflex, we adapted the assay from Neuhauss et al., 1999. A set of 4

tracks (each lane measuring at 12 cm x 0.6 cm) and comb that fit between the lanes, was

constructed using 3D printing. The lanes were placed in a 15 cm petri dish (VWR) and filled

partially with E3 buffer. The comb was inserted to block the starting end of the lanes, and 5-6

dpf larvae were placed in the starting area of the lanes. The petri dish was placed over a

screen that plays a video of 0.5 cm stripes. To record the movement of the larva along the

lanes, a camera was mounted above the petri dish. To perform the experiment, the video of

stripes was played, and the comb was removed, allowing larva to swim against the movement

of stripes to the end of the lane. To quantify this reflex, screenshots of the video were taken of

the larva in the starting region of the lane prior to comb removal and then 30 seconds after the

comb was removed. The distance between the location of each larva prior to comb removal

and 30 seconds after was measured in ImageJ in millimeters.

Electroretinogram (ERG)

ERG recordings were performed on 5 - 6 dpf mutant and wild-type sibling larvae, as described

in [40,113]. b-wave amplitudes were plotted and analyzed using Dunn's multiple comparisons.

Immunohistochemistry

Larvae were fixed with 4% PFA, dehydrated with Methanol, rehydrated using a graded series of

PBSDT (1x PBS with 0.25% Tween-20 and 1% DMSO): Methanol and then treated with 150

mM Tris-HCl for antigen retrieval. Samples were then washed and permeabilized with ice cold

100% acetone, followed by more washes. Samples were blocked for 2 hours at room

temperature with 6% goat serum and 3% BSA, then incubated overnight with 1:1000 anti-

acetylated tubulin rabbit primary antibodies (ThermoFisher). Samples were then incubated

overnight with 1:1000 anti-rabbit Alexa 488 (Invitrogen) secondary antibodies, protected from

light.

Lysotracker

5 dpf PTU-treated larva were sorted and placed in mutant-sibling pairs in a 2 mL Eppendorf

tube. LysoTracker Red DND-99 (Invitrogen) was diluted to 1:100 to examine brains or 1:200 to

examine inner ear hair cells. The larvae were incubated in Lysotracker for 40 minutes at 28.5°C

with the tube left open but protected from light. The larvae were then washed with E3 buffer for

10 minutes and immediately imaged.

Hindbrain lysotracker uptake was quantified by selecting a single Z-stack slice of the

hindbrain region and thresholding it to highlight particles in the size range of endosomal

compartments (size = 0.02-2 μm). The integrated density and area of particles from two equally

sized regions of interest from hindbrain sections and a single region of neuropil were measured

from each slice using the Analyze Particles function of Fiji. The integrated density

measurements were averaged and normalized to the integrated density of the neuropil.

CD63-phluorin Injections and Analysis

pUbi-CD63-pHluorin was a gift from Frederik Verweij and was generated as described in [114].

Nacre-background $vps4a^{T248I}$ mutant and sibling embryos were injected at the one cell stage

with 100 ng/μL pUbi-CD63-pHluorin. Damaged embryos were sorted out, and only larvae with

CD63-pHluorin expression on muscle cell membranes were selected for live imaging on 3-4 dpf.

Time lapse videos were recorded at 50 fps for 2 minutes at 63x with a 2x optical zoom. For

analysis, three frames were combined, and the mean pixel intensity was measured from regions

of the ventricular intracranial space, surrounding tissue as background, and muscle cells. The

mean pixel intensity of ventricular intracranial space was normalized to background and then

divided by muscle cell pixel intensity.

Bulk Tissue RNASeq

Total RNA was isolated from 5 dpf wild type and vps4a mutant larvae (n = 15-20 each

genotype) using TRIzol reagent (Thermo Fisher, 15596026) according to the manufacturer's

protocol. RNA-seq library construction and single-end sequencing were performed by the OHSU

Massively Parallel Sequencing Shared Resource (RRID SCR_009984, Oregon Health &

Science University, Portland, OR, USA). Sequencing reads were mapped to the Danio rerio

GRCz10 genome using Tophat v2.1.1 [115] from within an OSHU instance of the Galaxy

sequence analysis platform [116]. The resulting BAM files were imported to SeqMonk v1.42.0

(www.bioinformatics.babraham.ac.uk/projects/seqmonk/) for data visualization and statistical

analysis using the Intensity Difference filter for unreplicated data sets. Genes with an adjusted

p-value < 0.05 were considered differentially expressed.

Hybridization Chain Reaction (HCR)

For HCR, custom probes for *vps4a*, *atf3*, *jun*, *gap43, opn1sw1* and the corresponding

fluorescent B-amplifier hairpins were ordered from Molecular Instruments

(https://www.molecularinstruments.com/). Probes were chosen based on their log2(fold

change) and number of reads from bulk RNA sequencing analysis. Probes were assigned B-

amplifiers with the intention of doing multi-plex experiments. Probe sequences are considered

confidential and proprietary information of Molecular Instruments based on their Terms and

Conditions, so further inquiries may be directed to their technical team.

PTU-treated larvae were sorted at 5 dpf and fixed overnight in 4% paraformaldehyde

(PFA) at 4°C with rocking. After fixation, larvae were washed in PBST (1X PBS with 0.1%

Tween-20), dehydrated with a methanol:PBS series (25%, 50% and 75% methanol in PBS),

then stored at least overnight at -20°C. Larvae were subsequently rehydrated with a reverse

methanol:PBST series, washed several times with PBST, permeabilized for 15 minutes with 20

μg/mL Proteinase K, post-fixed with 4% PFA and then washed again. Probe detection and

amplification with fluorescent hairpins was performed as recommended by the MI protocol for

whole mount zebrafish larva with extended pre-hybridization and pre-amplification times (4

hours each) and probe concentrations adjusted based on the reads of each probe (5 nM for

gap43, 10 nM for *vps4a*).

Calcium Imaging

Larvae at 3 dpf *vps4a^{T248I} Tg(α-tubulin-gal4:UAS-GCaMP7a)* were sorted for strong GCaMP

expression in the brain. 5 dpf GCaMP positive larvae were mounted in 2% low melt agarose on

a custom depression slide fitted with a mini-speaker (DAEX-9-45M Skinny Mini Exciter 9 mm 1

W 4 Ohm). Using an AIYIMA A07 amplifier, tone stimuli were played through the mini speaker

using a custom script in MATLAB (R2020b, MathWorks). 600 Hz (0.35 V) or 450 Hz (0.2 V)

tones were played for 0.6 seconds or 0.1 seconds, respectively, five times with 10 second

intervals. Time lapse images and fluorescence values were recorded using LAS-X software

(Ver 3.7.2.22383). To account for variability in GCaMP7a expression between larvae, all

fluorescence values were normalized by subtracting background measurements. Traces were

generated by averaging the normalized fluorescence value (F) for each time point across all

larvae and plotting this value over time. To calculate percentage change in fluorescence

(ΔFt/F0%), 2-3 frames, containing image artifacts (due to the tone stimulation shaking the slide)

were identified for each stimulus, and fluorescence values corresponding to these frames were

discarded. The F0 measurement was selected from the frame prior to tone stimulation, based

on the time lapse video and presence of artifact frames, and Ft was selected as the highest

frame immediately after tone stimulation. (ΔFt/F0)/F0 was used to calculate the change in

GCaMP7a fluorescence and was multiplied by 100 to convert into a percentage to generate

"(ΔFt/F0%)". The average percentage change in fluorescence was then calculated for each

larva and differences between wild-type siblings and *vps4a* mutants was analyzed using

unpaired t tests.

Retrograde Labeling of Mauthner Cells

Retrograde labeling was adapted from previously published methods [117,118]. Larvae were

anesthetized with 0.3% MESAB prior to spinal injection of 10mg/mL 10,000 MW

Tetramethylrhodamine dextran (10,000 MW, D1866, Thermo Fisher) in PBS. Custom glass

injection needles were prepared from borosilicate glass capillaries (TW100F-4, World Precision

Instruments) using a micro-needle puller (Suttner Instruments Co, Model P-87) using heat cycle

517, pulling cycle 75, velocity 75 and time 90, and then the tip was broken using forceps.

Larvae were mounted laterally in 2% low melt agarose for injection, and then incubated in E3

media overnight to recover before imaging.

Microscopy and Image Processing

Images for LysoTracker, CD63pHluorin, and HCR were take using an upright LSM700 laser-

scanning confocal microscope with Zeiss Zen software (Carl Zeiss). For live imaging of larva,

they were anesthetized in 0.03% MESAB in E3 media and mounted in 2% prewarmed low melt

agarose. Images were collected using 488 nm, 555 nm and 647 nm lasers, depending on the

requirements of the individual experiment (Acetylated Tubulin and CD63pHluorin used 488 nm,

LysoTracker used 555 nm, HCR used up to all 3 depending on the fluorescent hairpins used).

We used a Leica HC PL Fluotar 10x/0.3 lens for Acetylated Tubulin, HCR and LysoTracker

images, a Plan Apochromat 40x/1.0DIC lens for LysoTracker quantification images and an

Acroplan 63x/0.95W water immersion lens for higher magnification HCR images. Laser power

and gain were determined independently for each fluorophore and kept consistent within each

experiment. The FIJI build of ImageJ Version 2.1.0/1.53t was used to process all raw image

files. For Calcium Signaling, scans were taken using a Leica SP-8 Confocal Microscope

equipped with a 20x (NA 1.0) objective, using LAS-X software. Images were collected using

488 nm white laser light.

Experimental Design and Statistical Analysis

The number of zebrafish larvae used is denoted in the figure legends. G*Power software was

used to ensure there were sufficient samples tested. All statistical analyses were performed

using GraphPad Prism software (version 9). The data was tested for normal distribution using

Shapiro-Wilk test. Data is presented as mean $\pm$ SD, and significance between mutant and wild-

type sibling groups was determined using unpaired t test unless otherwise noted. The image in

figure 1C was adapted from Sun et al., 2017 using Mol* [119] and RCSB PDB (PDB ID:5XMI)

software.

Acknowledgements: The authors would like to thank the past members of the Genetics

Department of the Max Planck Institute in Tuebingen, Germany, particularly Tatjana Piotrowski

and Stefan Schulte-Merker for their assistance in a small-scale allele screen. We would also like

to thank Peng Sun for his technical assistance, and Matthew Esqueda and Sivan Brodo-Abo for

their help with animal care. This work was supported in part by a National Institute on Deafness

and Other Communication Disorders award (DC0170046) and SICHL funding from the OHNS

department of Stanford University to T.N.; a grant from the National Natural Science Foundation

of China to S.S. (92254306); Swiss National Science Foundation (310030_204648) to S.N.

References

1. Lewis EM, Kroll KL. Development and disease in a dish: the epigenetics of

neurodevelopmental disorders. Epigenomics. 2018;10: 219–231. doi:10.2217/epi-2017-

0113

2. Cardoso AR, Lopes-Marques M, Silva RM, Serrano C, Amorim A, Prata MJ, et al. Essential

genetic findings in neurodevelopmental disorders. Hum Genomics. 2019;13: 31.

doi:10.1186/s40246-019-0216-4

3. Parenti I, Rabaneda LG, Schoen H, Novarino G. Neurodevelopmental Disorders: From

Genetics to Functional Pathways. Trends Neurosci. 2020;43: 608–621.

doi:10.1016/j.tins.2020.05.004

4. Dan B. Developmental central hypotonia: implications for counselling, prognosis, and

management. Dev Med Child Neurol. 2022;64: 4. doi:10.1111/dmcn.15085

5. Rodger C, Flex E, Allison RJ, Sanchis-Juan A, Hasenahuer MA, Cecchetti S, et al. De Novo

VPS4A Mutations Cause Multisystem Disease with Abnormal Neurodevelopment. Am J

Hum Genet. 2020;107: 1129–1148. doi:10.1016/j.ajhg.2020.10.012

6. Ganguly J, Kulshreshtha D, Almotiri M, Jog M. Muscle Tone Physiology and Abnormalities.

Toxins. 2021;13: 282. doi:10.3390/toxins13040282

7. Seu KG, Trump LR, Emberesh S, Lorsbach RB, Johnson C, Meznarich J, et al. VPS4A

Mutations in Humans Cause Syndromic Congenital Dyserythropoietic Anemia due to

Cytokinesis and Trafficking Defects. Am J Hum Genet. 2020;107: 1149–1156.

doi:10.1016/j.ajhg.2020.10.013

8. Lunati A, Petit A, Lapillonne H, Gameiro C, Saillour V, Garel C, et al. VPS4A mutation in

syndromic congenital hemolytic anemia without obvious signs of dyserythropoiesis. Am J

Hematol. 2021;96: E121–E123. doi:10.1002/ajh.26099

9. Scheuring S, Röhricht RA, Schöning-Burkhardt B, Beyer A, Müller S, Abts HF, et al.

Mammalian cells express two VPS4 proteins both of which are involved in intracellular

protein trafficking. J Mol Biol. 2001;312: 469–480. doi:10.1006/jmbi.2001.4917

10. Strausberg RL, Feingold EA, Grouse LH, Derge JG, Klausner RD, Collins FS, et al. Generation

and initial analysis of more than 15,000 full-length human and mouse cDNA sequences.

Proc Natl Acad Sci U S A. 2002;99: 16899–16903. doi:10.1073/pnas.242603899

11. Babst M, Sato TK, Banta LM, Emr SD. Endosomal transport function in yeast requires a novel

AAA-type ATPase, Vps4p. EMBO J. 1997;16: 1820–1831. doi:10.1093/emboj/16.8.1820

12. Beyer A, Scheuring S, Müller S, Mincheva A, Lichter P, Köhrer K. Comparative sequence and

expression analyses of four mammalian VPS4 genes. Gene. 2003;305: 47–59.

doi:10.1016/S0378-1119(02)01205-2

13. Monroe N, Han H, Gonciarz MD, Eckert DM, Karren MA, Whitby FG, et al. The oligomeric

state of the active Vps4 AAA ATPase. J Mol Biol. 2014;426: 510–525.

doi:10.1016/j.jmb.2013.09.043

14. Alonso Y Adell M, Migliano SM, Teis D. ESCRT-III and Vps4: a dynamic multipurpose tool for

membrane budding and scission. FEBS J. 2016;283: 3288–3302. doi:10.1111/febs.13688

15. Babst M, Katzmann DJ, Estepa-Sabal EJ, Meerloo T, Emr SD. Escrt-III: an endosome-

associated heterooligomeric protein complex required for mvb sorting. Dev Cell. 2002;3:

271–282. doi:10.1016/s1534-5807(02)00220-4

16. Ott C, Nachmias D, Adar S, Jarnik M, Sherman S, Birnbaum RY, et al. VPS4 is a dynamic

component of the centrosome that regulates centrosome localization of γ-tubulin,

centriolar satellite stability and ciliogenesis. Sci Rep. 2018;8: 3353. doi:10.1038/s41598-

018-21491-x

17. Cacheiro P, Muñoz-Fuentes V, Murray SA, Dickinson ME, Bucan M, Nutter LMJ, et al. Human

and mouse essentiality screens as a resource for disease gene discovery. Nat Commun.

2020;11: 655. doi:10.1038/s41467-020-14284-2

18. Loncle N, Agromayor M, Martin-Serrano J, Williams DW. An ESCRT module is required for

neuron pruning. Sci Rep. 2015;5: 8461. doi:10.1038/srep08461

19. Haffter P, Granato M, Brand M, Mullins MC, Hammerschmidt M, Kane DA, et al. The

identification of genes with unique and essential functions in the development of the

zebrafish, Danio rerio. Development. 1996;123: 1–36. doi:10.1242/dev.123.1.1

20. Vaz R, Hofmeister W, Lindstrand A. Zebrafish Models of Neurodevelopmental Disorders:

Limitations and Benefits of Current Tools and Techniques. Int J Mol Sci. 2019;20: 1296.

doi:10.3390/ijms20061296

21. de Abreu MS, Genario R, Giacomini ACVV, Demin KA, Lakstygal AM, Amstislavskaya TG, et

al. Zebrafish as a Model of Neurodevelopmental Disorders. Neuroscience. 2020;445: 3–11.

doi:10.1016/j.neuroscience.2019.08.034

22. Fetcho JR. The utility of zebrafish for studies of the comparative biology of motor systems. J

Exp Zoolog B Mol Dev Evol. 2007;308B: 550–562. doi:10.1002/jez.b.21127

23. Bandmann O, Burton EA. Genetic zebrafish models of neurodegenerative diseases.

Neurobiol Dis. 2010;40: 58–65. doi:10.1016/j.nbd.2010.05.017

24. Kalueff AV, Stewart AM, Gerlai R. Zebrafish as an emerging model for studying complex

brain disorders. Trends Pharmacol Sci. 2014;35: 63–75. doi:10.1016/j.tips.2013.12.002

25. Sakai C, Ijaz S, Hoffman EJ. Zebrafish Models of Neurodevelopmental Disorders: Past,

Present, and Future. Front Mol Neurosci. 2018;11. Available:

https://www.frontiersin.org/articles/10.3389/fnmol.2018.00294

26. Leshchiner I, Alexa K, Kelsey P, Adzhubei I, Austin-Tse CA, Cooney JD, et al. Mutation

mapping and identification by whole-genome sequencing. Genome Res. 2012;22: 1541–

1548. doi:10.1101/gr.135541.111

27. Davies BA, Azmi IF, Payne J, Shestakova A, Horazdovsky BF, Babst M, et al. Coordination of

substrate binding and ATP hydrolysis in Vps4-mediated ESCRT-III disassembly. Mol Biol

Cell. 2010;21: 3396–3408. doi:10.1091/mbc.E10-06-0512

28. McCullough J, Frost A, Sundquist WI. Structures, Functions, and Dynamics of ESCRT-III/Vps4

Membrane Remodeling and Fission Complexes. Annu Rev Cell Dev Biol. 2018;34: 85–109.

doi:10.1146/annurev-cellbio-100616-060600

29. Wu RS, Lam II, Clay H, Duong DN, Deo RC, Coughlin SR. A Rapid Method for Directed Gene

Knockout for Screening in G0 Zebrafish. Dev Cell. 2018;46: 112-125.e4.

doi:10.1016/j.devcel.2018.06.003

30. Kroll F, Powell GT, Ghosh M, Gestri G, Antinucci P, Hearn TJ, et al. A simple and effective F0

knockout method for rapid screening of behaviour and other complex phenotypes. Ekker

SC, Stainier DY, Balciunas D, editors. eLife. 2021;10: e59683. doi:10.7554/eLife.59683

31. Yang Q, Chen D, Xiong F, Chen D, Liu C, Liu Y, et al. A splicing mutation in VPS4B causes

dentin dysplasia I. J Med Genet. 2016;53: 624–633. doi:10.1136/jmedgenet-2015-103619

32. Hanovice NJ, Daly CMS, Gross JM. N-Ethylmaleimide–Sensitive Factor b (nsfb) Is Required

for Normal Pigmentation of the Zebrafish Retinal Pigment Epithelium. Invest Ophthalmol

Vis Sci. 2015;56: 7535–7544. doi:10.1167/iovs.15-17704

33. Woods IG, Lyons DA, Voas MG, Pogoda H-M, Talbot WS. nsf Is Essential for Organization of

Myelinated Axons in Zebrafish. Curr Biol. 2006;16: 636–648.

doi:10.1016/j.cub.2006.02.067

34. Gao Y, Khan YA, Mo W, White KI, Perkins M, Pfuetzner RA, et al. Sensory deficit screen

identifies nsf mutation that differentially affects SNARE recycling and quality control. Cell

Rep. 2023;42: 112345. doi:10.1016/j.celrep.2023.112345

35. Sun S, Li L, Yang F, Wang X, Fan F, Yang M, et al. Cryo-EM structures of the ATP-bound

Vps4E233Q hexamer and its complex with Vta1 at near-atomic resolution. Nat Commun.

2017;8: 16064. doi:10.1038/ncomms16064

36. Verweij FJ, Bebelman MP, Jimenez CR, Garcia-Vallejo JJ, Janssen H, Neefjes J, et al.

Quantifying exosome secretion from single cells reveals a modulatory role for GPCR

signaling. J Cell Biol. 2018;217: 1129–1142. doi:10.1083/jcb.201703206

37. Hou L, Liu K, Li Y, Ma S, Ji X, Liu L. Necrotic pyknosis is a morphologically and biochemically

distinct event from apoptotic pyknosis. J Cell Sci. 2016;129: 3084–3090.

doi:10.1242/jcs.184374

38. Kalluri R, LeBleu VS. The biology, function, and biomedical applications of exosomes.

Science. 2020;367: eaau6977. doi:10.1126/science.aau6977

39. Kist AM, Portugues R. Optomotor Swimming in Larval Zebrafish Is Driven by Global Whole-

Field Visual Motion and Local Light-Dark Transitions. Cell Rep. 2019;29: 659-670.e3.

doi:10.1016/j.celrep.2019.09.024

40. Niklaus S, Cadetti L, Berg-Maurer CM vom, Lehnherr A, Hotz AL, Forster IC, et al. Shaping of

Signal Transmission at the Photoreceptor Synapse by EAAT2 Glutamate Transporters.

eNeuro. 2017;4. doi:10.1523/ENEURO.0339-16.2017

41. Nicolson T. The genetics of hair-cell function in zebrafish. J Neurogenet. 2017;31: 102–112.

doi:10.1080/01677063.2017.1342246

42. Mo W, Chen F, Nechiporuk A, Nicolson T. Quantification of vestibular-induced eye

movements in zebrafish larvae. BMC Neurosci. 2010;11: 110. doi:10.1186/1471-2202-11-

110

43. Gao Y, Nicolson T. Temporal Vestibular Deficits in synaptojanin 1 (synj1) Mutants. Front Mol

Neurosci. 2021;13: 604189. doi:10.3389/fnmol.2020.604189

44. Sun P, Zhang Y, Zhao F, Wu J-P, Pun SH, Peng C, et al. An Assay for Systematically

Quantifying the Vestibulo-Ocular Reflex to Assess Vestibular Function in Zebrafish Larvae.

Front Cell Neurosci. 2018;12: 257. doi:10.3389/fncel.2018.00257

45. Tanimoto M, Ota Y, Horikawa K, Oda Y. Auditory input to CNS is acquired coincidentally

with development of inner ear after formation of functional afferent pathway in zebrafish.

J Neurosci Off J Soc Neurosci. 2009;29: 2762–2767. doi:10.1523/JNEUROSCI.5530-08.2009

46. Bianco IH, Ma L-H, Schoppik D, Robson DN, Orger MB, Beck JC, et al. The Tangential Nucleus

Controls a Gravito-inertial Vestibulo-ocular Reflex. Curr Biol. 2012;22: 1285–1295.

doi:10.1016/j.cub.2012.05.026

47. Jia Y, Bagnall MW. Monosynaptic targets of utricular afferents in the larval zebrafish. Front

Neurol. 2022;13. doi:10.3389/fneur.2022.937054

48. Liu Z, Hildebrand DGC, Morgan JL, Jia Y, Slimmon N, Bagnall MW. Organization of the

gravity-sensing system in zebrafish. Nat Commun. 2022;13: 5060. doi:10.1038/s41467-

022-32824-w

49. Hamling KR, Harmon K, Schoppik D. The Nature and Origin of Synaptic Inputs to

Vestibulospinal Neurons in the Larval Zebrafish. eNeuro. 2023;10: ENEURO.0090-23.2023.

doi:10.1523/ENEURO.0090-23.2023

50. Sugioka T, Tanimoto M, Higashijima S. Biomechanics and neural circuits for vestibular-

induced fine postural control in larval zebrafish. Nat Commun. 2023;14: 1217.

doi:10.1038/s41467-023-36682-y

51. Muto A, Ohkura M, Abe G, Nakai J, Kawakami K. Real-Time Visualization of Neuronal

Activity during Perception. Curr Biol. 2013;23: 307–311. doi:10.1016/j.cub.2012.12.040

52. Privat M, Romano SA, Pietri T, Jouary A, Boulanger-Weill J, Elbaz N, et al. Sensorimotor

Transformations in the Zebrafish Auditory System. Curr Biol. 2019;29: 4010-4023.e4.

doi:10.1016/j.cub.2019.10.020

53. Tanimoto M, Watakabe I, Higashijima S. Tiltable objective microscope visualizes selectivity

for head motion direction and dynamics in zebrafish vestibular system. Nat Commun.

2022;13: 7622. doi:10.1038/s41467-022-35190-9

54. Kohashi T, Oda Y. Initiation of Mauthner- or Non-Mauthner-Mediated Fast Escape Evoked

by Different Modes of Sensory Input. J Neurosci. 2008;28: 10641–10653.

doi:10.1523/JNEUROSCI.1435-08.2008

55. Sillar KT. Escape Behaviour: Reciprocal Inhibition Ensures Effective Escape Trajectory. Curr

Biol. 2009;19: R697–R699. doi:10.1016/j.cub.2009.07.015

56. López-Schier H. Neuroplasticity in the acoustic startle reflex in larval zebrafish. Curr Opin

Neurobiol. 2019;54: 134–139. doi:10.1016/j.conb.2018.10.004

57. Vanwalleghem G, Heap LA, Scott EK. A profile of auditory-responsive neurons in the larval

zebrafish brain. J Comp Neurol. 2017;525: 3031–3043. doi:10.1002/cne.24258

58. Inoue M, Tanimoto M, Oda Y. The role of ear stone size in hair cell acoustic sensory

transduction. Sci Rep. 2013;3: 2114. doi:10.1038/srep02114

59. Echteler SM. Connections of the auditory midbrain in a teleost fish, Cyprinus carpio. J Comp

Neurol. 1984;230: 536–551. doi:10.1002/cne.902300405

60. Edds-Walton PL, Fay RR. Directional and frequency response characteristics in the

descending octaval nucleus of the toadfish (Opsanus tau). J Comp Physiol A. 2008;194:

1013–1029. doi:10.1007/s00359-008-0373-8

61. McCormick CA. Anatomy of the Central Auditory Pathways of Fish and Amphibians. In: Fay

RR, Popper AN, editors. Comparative Hearing: Fish and Amphibians. New York, NY:

Springer; 1999. pp. 155–217. doi:10.1007/978-1-4612-0533-3_5

62. Marsden KC, Granato M. In vivo Ca2+ imaging reveals that decreased dendritic excitability

drives startle habituation. Cell Rep. 2015;13: 1733–1740. doi:10.1016/j.celrep.2015.10.060

63. Xu L, Guan NN, Huang C-X, Hua Y, Song J. A neuronal circuit that generates the temporal

motor sequence for the defensive response in zebrafish larvae. Curr Biol. 2021;31: 3343-

3357.e4. doi:10.1016/j.cub.2021.06.054

64. Shimazaki T, Tanimoto M, Oda Y, Higashijima S. Behavioral Role of the Reciprocal Inhibition

between a Pair of Mauthner Cells during Fast Escapes in Zebrafish. J Neurosci. 2019;39:

1182–1194. doi:10.1523/JNEUROSCI.1964-18.2018

65. Koyama M, Kinkhabwala A, Satou C, Higashijima S, Fetcho J. Mapping a sensory-motor

network onto a structural and functional ground plan in the hindbrain. Proc Natl Acad Sci.

2011;108: 1170–1175. doi:10.1073/pnas.1012189108

66. Thiele TR, Donovan JC, Baier H. Descending Control of Swim Posture by a Midbrain Nucleus

in Zebrafish. Neuron. 2014;83: 679–691. doi:10.1016/j.neuron.2014.04.018

67. Migault G, van der Plas TL, Trentesaux H, Panier T, Candelier R, Proville R, et al. Whole-Brain

Calcium Imaging during Physiological Vestibular Stimulation in Larval Zebrafish. Curr Biol.

2018;28: 3723-3735.e6. doi:10.1016/j.cub.2018.10.017

68. Severi KE, Portugues R, Marques JC, O'Malley DM, Orger MB, Engert F. Neural Control and

Modulation of Swimming Speed in the Larval Zebrafish. Neuron. 2014;83: 692–707.

doi:10.1016/j.neuron.2014.06.032

69. Wang W-C, McLean DL. Selective Responses to Tonic Descending Commands by Temporal

Summation in a Spinal Motor Pool. Neuron. 2014;83: 708–721.

doi:10.1016/j.neuron.2014.06.021

70. Sankrithi NS, O'Malley DM. Activation of a multisensory, multifunctional nucleus in the

zebrafish midbrain during diverse locomotor behaviors. Neuroscience. 2010;166: 970–993.

doi:10.1016/j.neuroscience.2010.01.003

71. Berg EM, Mrowka L, Bertuzzi M, Madrid D, Picton LD, El Manira A. Brainstem circuits

encoding start, speed, and duration of swimming in adult zebrafish. Neuron. 2023;111:

372-386.e4. doi:10.1016/j.neuron.2022.10.034

72. Leung LC, Wang GX, Madelaine R, Skariah G, Kawakami K, Deisseroth K, et al. Neural

signatures of sleep in zebrafish. Nature. 2019;571: 198–204. doi:10.1038/s41586-019-

1336-7

73. Schöneberg J, Pavlin MR, Yan S, Righini M, Lee I-H, Carlson L-A, et al. ATP-dependent force

generation and membrane scission by ESCRT-III and Vps4. Science. 2018;362: 1423–1428.

doi:10.1126/science.aat1839

74. Su M, Guo EZ, Ding X, Li Y, Tarrasch JT, Brooks CL, et al. Mechanism of Vps4 hexamer

function revealed by cryo-EM. Sci Adv. 2017;3: e1700325. doi:10.1126/sciadv.1700325

75. Babst M, Davies BA, Katzmann DJ. Regulation of Vps4 during MVB sorting and cytokinesis.

Traffic Cph Den. 2011;12: 1298–1305. doi:10.1111/j.1600-0854.2011.01230.x

76. Piper RC, Katzmann DJ. Biogenesis and Function of Multivesicular Bodies. Annu Rev Cell Dev

Biol. 2007;23: 519–547. doi:10.1146/annurev.cellbio.23.090506.123319

77. Bebelman MP, Janssen E, Pegtel DM, Crudden C. The forces driving cancer extracellular

vesicle secretion. Neoplasia N Y N. 2020;23: 149–157. doi:10.1016/j.neo.2020.11.011

78. Meldolesi J. Exosomes and Ectosomes in Intercellular Communication. Curr Biol. 2018;28:

R435–R444. doi:10.1016/j.cub.2018.01.059

79. Dobrowolski R, De Robertis EM. Endocytic control of growth factor signalling: multivesicular

bodies as signalling organelles. Nat Rev Mol Cell Biol. 2012;13: 53–60.

doi:10.1038/nrm3244

80. Palmulli R, van Niel G. To be or not to be... secreted as exosomes, a balance finely tuned by

the mechanisms of biogenesis. Stahl P, Raposo G, editors. Essays Biochem. 2018;62: 177–

191. doi:10.1042/EBC20170076

81. Cocucci E, Racchetti G, Meldolesi J. Shedding microvesicles: artefacts no more. Trends Cell

Biol. 2009;19: 43–51. doi:10.1016/j.tcb.2008.11.003

82. Raymond CK, Howald-Stevenson I, Vater CA, Stevens TH. Morphological classification of the

yeast vacuolar protein sorting mutants: evidence for a prevacuolar compartment in class E

vps mutants. Mol Biol Cell. 1992;3: 1389–1402. doi:10.1091/mbc.3.12.1389

83. Frankel E, Shankar R, Moresco J, Yates J, Volkmann N, Audhya A. Ist1 regulates ESCRT-III

assembly and function during multivesicular endosome biogenesis in Caenorhabditis

elegans embryos. Nat Commun. 2017;8. doi:10.1038/s41467-017-01636-8

84. Bishop N, Woodman P. ATPase-defective Mammalian VPS4 Localizes to Aberrant

Endosomes and Impairs Cholesterol Trafficking. Mol Biol Cell. 2000;11: 227–239.

85. González AE, Muñoz VC, Cavieres VA, Bustamante HA, Cornejo V-H, Januário YC, et al.

Autophagosomes cooperate in the degradation of intracellular C-terminal fragments of

the amyloid precursor protein via the MVB/lysosomal pathway. FASEB J. 2017;31: 2446–

2459. doi:10.1096/fj.201600713R

86. Willén K, Edgar JR, Hasegawa T, Tanaka N, Futter CE, Gouras GK. Aβ accumulation causes

MVB enlargement and is modelled by dominant negative VPS4A. Mol Neurodegener.

2017;12: 61. doi:10.1186/s13024-017-0203-y

87. Zhang Y, Liu Y, Liu H, Tang WH. Exosomes: biogenesis, biologic function and clinical

potential. Cell Biosci. 2019;9: 19. doi:10.1186/s13578-019-0282-2

88. Waqas MY, Javid MA, Nazir MM, Niaz N, Nisar MF, Manzoor Z, et al. Extracellular vesicles

and exosome: insight from physiological regulatory perspectives. J Physiol Biochem.

2022;78: 573–580. doi:10.1007/s13105-022-00877-6

89. Colombo M, Raposo G, Théry C. Biogenesis, Secretion, and Intercellular Interactions of

Exosomes and Other Extracellular Vesicles. Annu Rev Cell Dev Biol. 2014;30: 255–289.

doi:10.1146/annurev-cellbio-101512-122326

90. Gurunathan S, Kang M-H, Kim J-H. A Comprehensive Review on Factors Influences

Biogenesis, Functions, Therapeutic and Clinical Implications of Exosomes. Int J

Nanomedicine. 2021;16: 1281–1312. doi:10.2147/IJN.S291956

91. Han W, Zhang H, Feng L, Dang R, Wang J, Cui C, et al. The emerging role of exosomes in

communication between the periphery and the central nervous system. MedComm.

2023;4: e410. doi:10.1002/mco2.410

92. Duarte-Silva E, Oriá AC, Mendonça IP, de Melo MG, Paiva IHR, Maes M, et al. Tiny in size,

big in impact: Extracellular vesicles as modulators of mood, anxiety and

neurodevelopmental disorders. Neurosci Biobehav Rev. 2022;135: 104582.

doi:10.1016/j.neubiorev.2022.104582

93. Dilsiz N. Role of exosomes and exosomal microRNAs in cancer. Future Sci OA. 2020;6:

FSO465. doi:10.2144/fsoa-2019-0116

94. Gomes AR, Sangani NB, Fernandes TG, Diogo MM, Curfs LMG, Reutelingsperger CP.

Extracellular Vesicles in CNS Developmental Disorders. Int J Mol Sci. 2020;21: 9428.

doi:10.3390/ijms21249428

95. Lim Y-J, Lee S-J. Are exosomes the vehicle for protein aggregate propagation in

neurodegenerative diseases? Acta Neuropathol Commun. 2017;5: 64.

doi:10.1186/s40478-017-0467-z

96. Budnik V, Ruiz-Cañada C, Wendler F. Extracellular vesicles round off communication in the

nervous system. Nat Rev Neurosci. 2016;17: 160–172. doi:10.1038/nrn.2015.29

97. Gurung S, Perocheau D, Touramanidou L, Baruteau J. The exosome journey: from

biogenesis to uptake and intracellular signalling. Cell Commun Signal CCS. 2021;19: 47.

doi:10.1186/s12964-021-00730-1

98. Hamling KR, Harmon K, Kimura Y, Higashijima S, Schoppik D. The vestibulospinal nucleus is a

locus of balance development. bioRxiv. 2023; 2023.12.06.570482.

doi:10.1101/2023.12.06.570482

99. Lacoste AMB, Schoppik D, Robson DN, Haesemeyer M, Portugues R, Li JM, et al. A

convergent and essential interneuron pathway for Mauthner-cell-mediated escapes. Curr

Biol CB. 2015;25: 1526–1534. doi:10.1016/j.cub.2015.04.025

100. Fukushima K. The interstitial nucleus of Cajal and its role in the control of movements of

head and eyes. Prog Neurobiol. 1987;29: 107–192. doi:10.1016/0301-0082(87)90016-5

101. Matsuda K, Kubo F. Circuit Organization Underlying Optic Flow Processing in Zebrafish.

Front Neural Circuits. 2021;15. Available:

https://www.frontiersin.org/articles/10.3389/fncir.2021.709048

102. Liu Z, Bagnall MW. Organization of vestibular circuits for postural control in zebrafish. Curr

Opin Neurobiol. 2023;82: 102776. doi:10.1016/j.conb.2023.102776

103. Huda MN, Nafiujjaman M, Deaguero IG, Okonkwo J, Hill ML, Kim T, et al. Potential use of

exosomes as diagnostic biomarker and in targeted drug delivery: progress in clinical and

preclinical application. ACS Biomater Sci Eng. 2021;7: 2106–2149.

doi:10.1021/acsbiomaterials.1c00217

104. Chen Y, Li J, Ma B, Li N, Wang S, Sun Z, et al. MSC-derived exosomes promote recovery

from traumatic brain injury via microglia/macrophages in rat. Aging. 2020;12: 18274–

18296. doi:10.18632/aging.103692

105. Yari H, Mikhailova MV, Mardasi M, Jafarzadehgharehziaaddin M, Shahrokh S, Thangavelu

L, et al. Emerging role of mesenchymal stromal cells (MSCs)-derived exosome in

neurodegeneration-associated conditions: a groundbreaking cell-free approach. Stem Cell

Res Ther. 2022;13: 423. doi:10.1186/s13287-022-03122-5

106. Zhang X, Jiang W, Lu Y, Mao T, Gu Y, Ju D, et al. Exosomes combined with biomaterials in

the treatment of spinal cord injury. Front Bioeng Biotechnol. 2023;11: 1077825.

doi:10.3389/fbioe.2023.1077825

107. Westerfield M. The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish (Danio

rerio), 4th Edition. University of Oregon Press; 2000.

108. Knapik EW, Goodman A, Ekker M, Chevrette M, Delgado J, Neuhauss S, et al. A

microsatellite genetic linkage map for zebrafish (Danio rerio). Nat Genet. 1998;18: 338–

343. doi:10.1038/ng0498-338

109. Erickson T, Morgan CP, Olt J, Hardy K, Busch-Nentwich E, Maeda R, et al. Integration of

Tmc1/2 into the mechanotransduction complex in zebrafish hair cells is regulated by

Transmembrane O-methyltransferase (Tomt). Ginty DD, editor. eLife. 2017;6: e28474.

doi:10.7554/eLife.28474

110. Smith ET, Pacentine I, Shipman A, Hill M, Nicolson T. Disruption of tmc1/2a/2b Genes in

Zebrafish Reveals Subunit Requirements in Subtypes of Inner Ear Hair Cells. J Neurosci.

2020;40: 4457–4468. doi:10.1523/JNEUROSCI.0163-20.2020

111. Mirat O, Sternberg JR, Severi KE, Wyart C. ZebraZoom: an automated program for high-

throughput behavioral analysis and categorization. Front Neural Circuits. 2013;7: 107.

doi:10.3389/fncir.2013.00107

112. Budick SA, O'Malley DM. Locomotor repertoire of the larval zebrafish: swimming, turning

and prey capture. J Exp Biol. 2000;203: 2565–2579. doi:10.1242/jeb.203.17.2565

113. Makhankov YV, Rinner O, Neuhauss SCF. An inexpensive device for non-invasive

electroretinography in small aquatic vertebrates. J Neurosci Methods. 2004;135: 205–210.

doi:10.1016/j.jneumeth.2003.12.015

114. Verweij FJ, Revenu C, Arras G, Dingli F, Loew D, Pegtel DM, et al. Live Tracking of Inter-

organ Communication by Endogenous Exosomes In Vivo. Dev Cell. 2019;48: 573-589.e4.

doi:10.1016/j.devcel.2019.01.004

115. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment

of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol.

2013;14: R36. doi:10.1186/gb-2013-14-4-r36

116. Afgan E, Baker D, van den Beek M, Blankenberg D, Bouvier D, Čech M, et al. The Galaxy

platform for accessible, reproducible and collaborative biomedical analyses: 2016 update.

Nucleic Acids Res. 2016;44: W3–W10. doi:10.1093/nar/gkw343

117. Gahtan E, Sankrithi N, Campos JB, O'Malley DM. Evidence for a widespread brain stem

escape network in larval zebrafish. J Neurophysiol. 2002;87: 608–614.

doi:10.1152/jn.00596.2001

118. Wehner D, Becker T, Becker CG. Restoration of anatomical continuity after spinal cord

transection depends on Wnt/β-catenin signaling in larval zebrafish. Data Brief. 2017;16:

65–70. doi:10.1016/j.dib.2017.10.068

119. Sehnal D, Bittrich S, Deshpande M, Svobodová R, Berka K, Bazgier V, et al. Mol* Viewer:

modern web app for 3D visualization and analysis of large biomolecular structures. Nucleic

Acids Res. 2021;49: W431–W437. doi:10.1093/nar/gkab314

1132

1133

1134   FIGURE LEGENDS

1135

Figure 1. The T248I mutation in *vps4a* reduces ATPase activity and ESCRT-III filament

disassembly. A, *vps4a* is located on Chromosome 25 in a location corresponding to where

classical mapping methods have identified the location of the gene responsible for the

raumschiff phenotypes. **B,** Bar diagram of Vps4a protein showing the location of the T248I

mutation. **C,** Atomic model structure of the Vps4^{E233Q} hexamer, with T248 marked in magenta

(Image adapted from the RCSB PDB ID:5XMI, using Mol*). **D,** Mutant Vps4 (T254I) yeast

protein has decreased ATP activity compared to wild-type Vps4 protein. **E,** Sedimentation

analysis of Vps4 and Vps4(T254I)-mediated disassembly of ESCRT-III filaments. Pellet (P) and

supernatant (S) fractions were analyzed by SDS-PAGE. Marker (M). **F,** Negative staining EM

micrograph of sedimentation analysis of ESCRT-III filament disassembly.

1146

Figure 2. *vps4a*^{T248I} mutants have enlarged endosomal compartments in the CNS. A,

Confocal Z-projections of a 5 dpf wild-type zebrafish larva labeled with HCR RNA-FISH probes

for *vps4a.* fb, forebrain; ot, optic tectum; tpz, tectal proliferation zone, cb, cerebellum; hb,

hindbrain. **B,** LysoTracker staining of 5 dpf wild-type siblings and *vps4a*^{T248I} mutants show an

increased staining in the mutant. **C,** Acidic membrane compartment size in the hindbrain of

vps4a^{T248I} mutants (N = 10) is significantly larger than those in wild-type siblings (N = 7).

LysoTracker vital dye taken up by *vps4a*^{T248I} mutants (N = 10), measured as normalized pixel

integrated density, is significantly higher compared to wild-type siblings (N =7).

1155

Figure 3. *vps4a*^{T248I} mutants release fewer exosomes. A, Schematic of injecting 1-cell stage

nacre-background embryos with pUbi:CD63-pHluorin. Healthy larvae were sorted on 1 dpf, and

larvae expressing CD63-pHluorin in muscle cells was sorted on 3 dpf, for imaging on 4 dpf. **B,**

Decreased CD63-pHluorin expression in 3 dpf *vps4a*^{T248I} mutant CNS blood vessels. **C,** Dorsal

view of 4 dpf zebrafish larva showing the midbrain ventricle. **D,** Decreased CD63-pHluorin

expression in *vps4a*^{T248I} mutant brain ventricles, but trunk muscles express comparable levels to

wild-type siblings. **E,** Pixel intensity of muscles cells is correlated with normalized pixel intensity

of CD63-pHluorin signal in the midbrain ventricle in wild-type siblings (N = 16). **F,** Quantification

of CD63-pHluorin fluorescence in wild-type siblings (N = 16) and *vps4a*^{T248I} mutants (N = 8).

1165

Figure 4. *vps4a*^{T248I} mutants upregulate stress response and regeneration genes in the

CNS. A, Axons labeled with acetylated tubulin antibody staining of *vps4a*^{T248I} mutants show no

gross morphological defects. Fb, forebrain; ot, optic tectum; mb, midbrain; cb, cerebellum; hb,

hindbrain. **B,** *vps4a*^{T248I} mutants have an increase of morphologically abnormal, pyknotic cells

in the CNS. Two examples are highlighted with black dashed circles. The number of abnormal

cells in the *vps4a*^{T248I} mutants (N = 11) was quantified and found to be significantly higher than

in wild-type siblings (N = 9). **C,** Maximum intensity projections of *atf3* and *jun* mRNA expression

in the hindbrain of 5 dpf wild-type sibling and *vps4a*^{T248I} mutants. **D,** Maximum intensity

projections of *gap43* and *jun* mRNA expression in the hindbrain of 5 dpf wild-type sibling and

vps4a^T248I mutants.

1176

Figure 5. Impaired vision in *vps4a^T248I* mutants. **A,** 5 dpf *vps4a^T248I* mutants (N = 10) have a

reduced optomotor response compared to wild-type siblings (N = 12). **B,** Decreased b wave

amplitude in electroretinograms of 5-6 dpf *vps4a^T248I* mutants (N = 34) in comparison to wild-

type siblings (N = 23) at brighter light intensities (log -4 to log 0). **C,** HCR RNA-FISH of *gap43*,

jun and *opn1sw1* in the eye of *vps4a^T248I* mutants compared to wild-type siblings. Pr,

photoreceptors; Rg, retinal ganglion cells, lens.

1183

Figure 6. *vps4a^T248I* mutants have selective defects in vestibular circuitry. **A,** 5 dpf

vps4a^T248I mutants (N = 16) do not have a significantly different VIEM compared to wild-type

siblings (N = 18). **B,** 5 dpf *vps4a^T248I* mutants (N = 10) have a reduced VSR compared to wild-

type siblings (N = 7). **C,** Simplified schematic showing the flow of information evoked by head

rotation from the inner ear to hindbrain nuclei, vestibulo-ocular neurons, and motor neurons.

UHC, utricular hair cells; CN VIII, VIIIth nerve; SVN, superior vestibular nucleus; VS,

vestibulospinal cells; TAN, tangential neurons; M, Mauthner cell; CN IV, trochlear nucleus; CN

III, oculomotor nucleus; OM, oculomotor muscles. **D,** 5 dpf larva showing basal GCamP7a

expression and backfill injected Rhodamine-Dextran marking the Mauthner cell. White dotted

lines mark the TAN, SVN and VS. **D'** Pre- and post-stimulus GCaMP7a levels in vestibular

region in a representative wild-type sibling and *vps4a^T248I* mutant. The TAN, SVN and VS ROIs

are outlined by dotted lines. **E,** Average trace of normalized fluorescence (F) for the TAN, SVN

and VS of wild-type siblings and *vps4a^T248I* mutants during auditory stimulation at 600Hz. **F,**

Percentage change in fluorescence of the TAN, SVN and VS of wild-type siblings and *vps4a^T248I*

mutants during auditory stimulation at 600Hz. The VS in *vps4a^T248I* mutants (N = 11) have a

reduced change in fluorescence compared to wild-type siblings (N = 15). There is no significant

difference between mutants and wild-type siblings in the TAN or SVN.

1201

Figure 7. Activation of the ascending auditory pathway in *vps4a^{T248I}* mutants. A, simplified

schematic showing the flow of information from an auditory stimulus to motor neurons

(responsible for startle reflexes) or to the torus semicircularis (key for auditory processing). S,

saccular otolith; SHC, saccular hair cells; CN VIII, VIIIth nerve; MON, medial octavolateralis

nucleus; M, Mauthner cell; TS, torus semicircularis. **B,** 5 dpf *vps4a^{T248I}* mutants (N = 20) have a

greatly reduced AEBR compared to wild-type siblings (N = 15). **C,** Basal levels of GCaMP7a

expression in the inner ear of 5 dpf larvae, focusing on the saccular hair cells (dotted line) and

statoacoustic ganglion (SAG) (dashed line). **D,** Pre- and post-stimulus GCaMP7a levels in

saccular hair cells and SAG in a representative wild-type sibling and *vps4a^{T248I}* mutant larvae.

E, Average traces of normalized fluorescence (F) for the saccular hair cells and SAG of wild-

type siblings and *vps4a^{T248I}* mutants during auditory stimulation at 600Hz. **F,** Percentage

change in fluorescence of the saccular hair cells and SAG of wild-type siblings and *vps4a^{T248I}*

mutants during auditory stimulation at 600Hz. There is no significant difference between

mutants (N = 13) and wild-type siblings (N = 20) in the saccular hair cells or the SAG. **G,**

Representative image of 5 dpf zebrafish basal levels of GCaMP7a expression in the torus

semicircularis (TS). **H,** Pre-and post-stimulus GCaMP7a levels in the TS in representative wild-

type sibling and *vps4a^{T248I}* mutant larvae. **I,** Average trace of normalized fluorescence (F) for

the TS of wild-type siblings and *vps4a^{T248I}* mutants during auditory stimulation at 600Hz. **J,**

There is no significant difference in percentage change fluorescence between *vps4a^{T248I}*

mutants (N = 13) and wild-type siblings (N = 11) in the TS.

1222

1223   **Figure 8. *vps4a^{T248I}* mutants display defects in sensorimotor transformation**. **A,** simplified

1224   schematic showing the flow of information from a tone stimulus from the inner ear to motor

1225 neurons via Mauthner cell and nMLF pathways. S, saccular otolith; SHC, saccular hair cells;

1226 CN VIII, VIIIth nerve; M, Mauthner cell; HB, hindbrain; CRN, cranial relay neurons; nMLF,

1227 nucleus of the medial longitudinal fasciculus. **B,** 7 dpf *vps4a^{T248I}* mutant larvae have an intact

1228 startle reflex when touched. **C,** 7 dpf *vps4a^{T248I}* mutants (N = 11) do not have a significantly

1229 different C-bend angle compared to wild-type siblings (N = 10). **D,** Acetylated tubulin antibody

1230 staining of *vps4a^{T248I}* mutants shows no gross morphological defects in hindbrain/spinal cord or

1231 somatosensory axons. **E,** Representative image of 5 dpf zebrafish, showing location of nMLF

1232 expressing basal levels of GCaMP7a. **F,** Pre- and post-stimulus GCaMP7a levels in the nMLF

1233 in a representative wild-type sibling and *vps4a^{T248I}* mutant. **G,** Average trace of normalized

1234 fluorescence (F) for the nMLF of wild-type siblings and *vps4a^{T248I}* mutants during tone

1235 stimulation at 600Hz. **H,** *vps4a^{T248I}* mutants (N = 11) have a significantly reduced percentage

1236 change fluorescence in the nMLF compared to wild-type siblings (N = 14). **I,** Representative

1237 image of 5 dpf zebrafish trunk showing spinal cord neurons expressing basal levels of

1238 GCaMP7a. **J,** Pre-and post-stimulus GCaMP7a levels in the spinal cord motor neurons of

1239 representative wild-type sibling and *vps4a^{T248I}* mutant. **K,** Average trace of normalized

1240 fluorescence (F) for the spinal cord motor neurons of wild-type siblings and *vps4a^{T248I}* mutants

1241 during tone stimulation at 450Hz. **L,** *vps4a^{T248I}* mutants (N = 9) have a significantly reduced

1242 percentage change fluorescence in spinal cord neurons compared to wild-type siblings (N = 11).

1243

1244 **Figure S1. *vps4a^{T248I}* mutant hair cells do not display changes in LysoTracker labeling or**

1245 **gene expression. A,** Representative maximum projection images of wild-type siblings and

1246 *vps4a^{T248I}* mutant larvae showing LysoTracker staining of saccule and utricle. Hb, hindbrain. **B,**

1247 LysoTracker vital dye taken up by the utricle and saccule of *vps4a^{T248I}* mutants (N = 9),

1248 measured as normalized pixel integrated density, is not significantly higher compared to wild-

1249 type siblings (N =6). **C,** Maximum projection of LysoTracker staining of anterior cristae hair

1250 cells. Anterior cristae hair cells were imaged due to similarity to hair cells and convenience. **D,**

1251   LysoTracker vital dye uptake in apical endosomes in *vps4a^T248I* mutant anterior cristae hair cells

1252   (N = 10), measured as normalized pixel integrated density, is not significantly different

1253   compared to wild-type siblings (N = 10). Nuc, nucleus. **E,** Maximum intensity projections of *jun*

1254   mRNA expression in the saccular hair cells and statoacoustic ganglion of 5 dpf wild-type

1255   siblings and *vps4a^T248I* mutants.

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

Fig 1

A 7/353 recombinant larvae = 0.99 cM or 700 kb

18/353 recombinant larvae = 2.8cM

z5669 34.33 Mb

z1213 35.87 Mb

rpgrip1l 36.04 Mb

vps4a 34.93 Mb

sall1b 35.98 Mb

B

MIT   AAA ATPase Domain   β-domain   final helix

1   123   *   353   437

T248I

		*	
H. sapiens	240-	ESEAARRIK**T**EFLVQM	-255
D. rerio	239-	ESEAARRIK**T**EFLVQM	-254
S. cerevisiae	245-	ESEASRRIK**T**EFLVQM	-260
A. thaliana	245-	ESEASRRIK**T**EFLVQM	-260

C

T248

D

ATPase Activity

<0.0001

Enzyme Activity (U/L)

WT   Vps4(T254I)

E

Filament   Vps4(WT)   Vps4(T254I)   Vps4(WT) + Filament   Vps4(T254I) + Filament

S P S P S P S P S P M

180 kD
150 kD
100 kD
70 kD

Vps4 -   55 kD

Snf7(R52E) -   40 kD
Vps2/Vps24 -   35 kD

25 kD

15 kD
10 kD

F

Filament   Filament + Vps4(WT)

Filament + Vps4(T254I)

100 nm

Fig 2

A *vps4a* expression

WT Sibling

fb
ot
tpz
cb
hb
eye
ear
100 µm

ot   tpz
eye
ear

B LysoTracker Label

WT Sibling

ot
cb
ear
hb
100 µm
5 µm

*vps4a^T248I^*

C

Compartment Size
CNS

0.0015

Average Size of
Endocytic Compartments (µm²)

WT Sibling   *vps4a^T248I^*

LysoTracker Label
CNS

0.0020

Normalized
Integrated Density(A.U.)

WT Sibling   *vps4a^T248I^*

Fig 3

Fig 4

A   Axon Labeling

B   Abnormal Cells

Fig 5

A

Optomotor Response

Distance travelled in 30 seconds (mm)

<0.0001

WT Sibling $vps4a^{T248I}$

B

Electroretinogram

b-wave Amplitude (mV)

0.20   0.0014   <0.0001   <0.0001   <0.0001

log -4   log -3   log -2   log -1   log 0

Light Intensity

● WT Sibling
○ $vps4a^{T248I}$

C

	gap43	*jun*	*opn1sw1*	overlay
WT Sibling	pr / lens / rg			*gap43* *jun* *opn1sw1*
$vps4a^{T248I}$				

50 µm

Fig 6

Fig 7

Fig 8

Fig S1

A LysoTracker Label

WT Sibling

eye

saccule · utricle

ear · hb

100 µm · 50 µm

$vps4a^{T248I}$

B

Utricle
0.63

Normalized Integrated Density(A.U.)

WT Sibling · $vps4a^{T248I}$

Saccule
0.093

Normalized Integrated Density(A.U.)

WT Sibling · $vps4a^{T248I}$

C LysoTracker · DIC · overlay

WT Sibling

$vps4a^{T248I}$

10 µm

D

$vps4a^{T248I}$

apical endosomes

nuc

1 µm

Anterior Cristae
0.32

Normalized Integrated Density(A.U.)

WT Sibling · $vps4a^{T248I}$

E Saccular Hair Cells · Statoacoustic Ganglion

jun · *jun*

WT Sibling · WT Sibling

$vps4a^{T248I}$ · $vps4a^{T248I}$

5 µm · 5 µm