1 **Functional assessment of cell entry and receptor use for merbecoviruses**

3 Michael Letko[1#]

5 1. Paul G. Allen School for Global Health, Washington State University, Pullman, WA,

6 99163.

[#]**Corresponding author:**

michael.letko@wsu.edu

ABSTRACT:

The merbecovirus subgenus of coronaviruses includes Middle East Respiratory Syndrome

Coronavirus (MERS-CoV), which is a zoonotic respiratory pathogen that transmits from

dromedary camels to humans and causes severe respiratory disease. Viral discovery efforts

have uncovered hundreds of merbecoviruses in different species across multiple continents, but

few of these viruses have been isolated or studied under laboratory conditions, leaving basic

questions regarding their threat to humans unresolved. Viral entry into host cells is considered

an early and critical step for transmission between hosts. In this study, a scalable approach to

assessing novel merbecovirus cell entry was developed and used to measure receptor use

across the entire merbecovirus subgenus. Merbecoviruses are sorted into four clades based on

the receptor binding domain of the spike glycoprotein. Receptor tropism is clade-specific, with

only one clade using DPP4 and multiple clades using ACE2, including the entire HKU5 cluster

of bat coronaviruses.

INTRODUCTION:

Middle East Respiratory Syndrome Coronavirus (MERS-CoV) was first discovered in the middle east in 2012 and can cause severe respiratory illness in humans[1]. Field studies have shown that MERS-CoV is endemic in dromedary camels in the region and routinely transmits to humans, sustaining a slow but continual regional outbreak[2,3]. MERS-CoV is the prototypic member of the merbecovirus subgenus of beta-coronaviruses. Virus discovery efforts have uncovered hundreds of merbecoviruses circulating among diverse wildlife across multiple continents[4-13]. As is true with most of the global virome, the majority of all sequenced coronaviruses have never been isolated or molecularly characterized, leaving many questions regarding their threat of spillover into humans unknown.

In order to transmit to between species, coronaviruses must be able to enter the cells of their hosts. MERS-CoV cell entry is mediated by the viral spike protein and occurs in three steps: (1) spike attaches virus to the cell surface, (2) binds to the host cell receptor molecule, (3) and exposes structural elements in the base of the spike protein that mediate membrane fusion and subsequent cell entry[14-16]. Each function is performed by a separate region of the spike protein, with the Receptor Binding Domain (RBD) forming the core interaction with the host receptor[17-20]. While MERS-CoV uses the receptor dipeptidyl peptidase IV (DPP4) to infect cells, several studies have shown that some bat merbecoviruses do not use DPP4 and instead use angiotensin converting enzyme 2 (ACE2) to infect their hosts[21-23].

Similar to other beta-coronaviruses, the receptor binding domain (RBD) in merbecovirus spike proteins is a single, contiguous domain located in the N-terminal half of the spike sequence, and folds as a structurally stable unit that engages with the host receptor (Fig. 1a-b). Gene synthesis is required to produce fragments of viral genomes that have not been isolated, with sequence length dictating labor and cost. To reduce cost of viral gene synthesis while maintaining the ability to study viral protein function, we previously employed a method to replace the RBD of the sarbecovirus, SARS-CoV, with RBDs from any other sarbecovirus[24]. The resulting chimeric

spike gene was used in viral pseudotype assays to measure the entry efficiency in cell culture of

a panel of sarbecovirus RBDs[24]. Using this approach (called "SarbecoType"), we previously

characterized the cell entry for the majority of all published sarbecovirus RBDs, which lead to

isolation of new sarbecoviruses, identification of understudied cell entry pathways, and improved

understanding for molecular determinants of sarbecovirus cell entry[24-28].

Here, a similar approach called "MerbecoType" was taken with the merbecovirus

subgenus of betacoronaviruses. Similar to sarbecoviruses, RBDs of merbecoviruses can be

classified into four different "clades" based on the presence or absence of indels, and exhibit

clade-specific trends in receptor use and cell entry. Surprisingly, at least half of the merbecovirus

RBD clades can use ACE2, with the entire group of HKU5 viruses exhibiting clear preference for

ACE2 from their natural host species: *Pipistrellus ambramus* bats. These findings will be important

for future efforts to identify looming coronavirus threats and in the development of interventions

such as broad-spectrum antivirals and universal coronavirus vaccines.

RESULTS:

Mapping the functional receptor binding domain

In order to test a large panel of receptor binding domains from diverse merbecoviruses in viral pseudotype entry assays, the first objective was to produce a MERS-CoV spike expression construct that would allow for easy replacement of the receptor binding domain (RBD) with the RBD from other merbecoviruses (Fig. 1a-c). MERS-CoV was selected as the spike backbone because of its extensive structural characterization and compatibility with cell-culture based models of merbecovirus entry. Similar to our prior sarbecovirus study, the MERS-CoV spike was codon optimized for human cells, appended with a C-terminal FLAG tag and silent mutations were introduced outside of the RBD (Fig. 1b)[24]. While MERS-CoV uses DPP4 for entry, ACE2 has been shown to be the receptor for some other merbecoviruses including PDF2180 (Fig. 1d)[22]. Thus the RBD from PDF2180 made an ideal candidate to develop a chimeric MERS-CoV spike: a successful exchange of DPP4 use in MERS-CoV spike to ACE2 was considered indicative of a functional spike that would allow for further exploration of merbecovirus RBD receptor preferences. Unfortunately, there are very few amino acid motifs that are conserved across all merbecovirus spike sequences, complicating the task of identifying a functional region common to all merbecoviruses. However, closer inspection of published spike sequences revealed several highly conserved glycine residues flanking the RBD (MERS-CoV spike positions 372 and 610; Fig. 1a, d, e). These two residues are just beyond the RBD region of spike used for co-crystal structures with host receptor molecules. While the RBD has been determined structurally for some merbecoviruses, regions flanking the RBD are flexible, difficult to resolve with x-ray crystallography and not included in RBD:receptor based co-structures (Fig. 1a, c; pink shaded region). Several chimeras between MERS-CoV and PDF2180 were produced: version 1 included the region of PDF2180 spike between the conserved glycines, version 2 was based on structure and version 3 was based on the PDF2180-RBD as defined previously[22] (Fig. 1e). DNA fragments encoding the different RBD fragments were cloned into the MERS-CoV spike backbone and used

to generate vesicular stomatitis virus (VSV) single-cycle pseudotypes (Fig. 1c). While Baby

Hamster Kidney cells (BHKs) are not naturally susceptible to coronavirus entry but can support

entry if provided the receptor, BHKs expressing human ACE2 do not always recapitulate natural

viral entry for ACE2-dependent viruses with low affinity for the receptor[28,29]. Therefore, 293T,

which exhibit low susceptibility to coronavirus entry, were stably transduced to express human

ACE2 or DPP4. Infecting these stable cell lines with pseudotypes bearing chimeric spikes,

revealed that the chimeric spike containing the entire region of PDF2180 spike between the

conserved glycines was able to use human ACE2, indicating a successful receptor exchange for

MERS-CoV spike (Fig. 1d, f; PDF2180 V1). The regions of PDF2180 spike between residues 376-

381 and 586-609 were essential for receptor-use exchange, as two other spikes with shorter

regions of the PDF2180 RBD were unable to use ACE2 (Fig. 1a, d-f; PDF2180 V2, V3). Addition

of exogenous protease (trypsin) enhanced viral entry, allowing more sensitive detection of low

affinity interactions. Importantly, trypsin-enhanced entry was still receptor dependent (Fig. 1f).

Production of a chimeric spike containing an analogous region from the DPP4-dependent

HKU4/SM3A virus showed a retention in DPP4 use in MERS-CoV spike further suggesting this

region would allow for assessment of diverse receptor preferences (Fig. 1e-f).

Optimizing MERS-CoV spike for pseudotyping

With a working receptor-exchange in the spike construct, the next goal was to optimize the spike

backbone for pseudotype efficiency, which would allow for improved sensitivity to low-level viral

entry. After engaging with the host receptor, coronavirus spikes are processed by host-cell

proteases to release the spike fusion peptide and mediate membrane fusion. While the TMPRSS-

family of cell-surface proteases and endosomal cathepsins have been shown to cleave many

different coronavirus spikes, including MERS-CoV, SARS-CoV and SARS-CoV-2, furin is another

protease that plays a role both spike-biogenesis and cell entry, depending on the availability of an

appropriate cleavage site[30-33]. MERS-CoV spike contains two furin sites, at the S1/S2 and S2'

junctions, although the role of furin in entry is less clear as removal of these sites has no effect

on viral replication *in vitro*[34,35]. Because previous studies have shown that removal of the furin site

from SARS-CoV-2 spike results in increased pseudotype efficiency, several versions of spike were

produced with either one or both sites mutated[36-38]. Additionally, C-terminal truncations in MERS-

CoV, SARS-CoV and SARS-CoV-2 spikes have shown to increase cell-surface expression of

spike in transfected cells, resulting in improved pseudotype efficiency[39-41]. Therefore, the protease

site mutants were also tested with or without a 16-amino acid truncation in the C-terminal end of

MERS-CoV spike (Extended Data Fig. 1a-c). Surprisingly, these modifications to spike had very

little effect on entry in 293T cells for MERS-CoV WT, PDF2180 RBD or HKU4/SM3A RBD

(Extended Data Fig. 1b-c). While not statistically significant, spikes containing a C-terminal

truncation and removal of the S2' furin site performed slightly better than the other spikes,

therefore, this set of modifications was included for further experiments.

A panel of merbecoviruses encompassing total RBD diversity

The 2012 MERS-CoV outbreak lead to an influx of coronavirus sequencing and discovery

research, with sequences from hundreds of MERS-CoV patients as well as dromedary camels

after they were discovered as the intermediate species[42-45]. Genbank has over 2000 entries for

"merbecovirus" and "MERS-like" viruses; of these 2000 entries, only approximately 842 entries

contain full-length spike and of those entries, only approximately half are from species other than

humans and camels (Figure 2A). Unfortunately, coronavirus genbank listings do not always

contain universal identifiers, which left many entries for "MERS-CoV" even after excluding human

and camel viruses. Further eliminating viruses titled as "MERS-CoV" from the list resulted 65

spike sequences which encompassed 36 unique RBD sequences (Fig. 2a; table 1). Some

merbecoviruses are likely missing from this list as coronavirus nomenclature from 10-15 years

ago when the majority of these merbecovirus genomes were published was not well defined.

Phylogenetic analysis of the RBD amino acid sequences (as defined between the two glycine

residues established in figure 1) revealed several genetically clustered groups based on

characteristic indels and the sequences surrounding them (Fig. 2b, c). The groups were labeled

as clades 1 through 4, based on the order of their discovery: clade 1 includes MERS and HKU4

discovered between 2006-2012[1,46,47], clade 2 includes HKU5 also discovered around 2006[46],

clade 3 incudes the African merbecoviruses discovered in 2017[13] and clade 4 includes the

contemporary hedgehog merbecoviruses discovered in europe[13] (Fig. 2b, c, table 1). The RBDs

for these merbecoviruses was synthesized and cloned into the MERS-CoV spike backbone

plasmid. All chimeric spike plamsmids were stably expressed in 293T cells used to produce

pseudotyped viral-like particles for downstream experiments (Fig. 2d).

Merbecovirus entry in common laboratory cell lines

Vero cells are common laboratory cell lines in coronavirus research and are used in attempts to

isolate and propagate viruses, including merbecoviruses. VeroE6 cells were poorly susceptible to

the merbecovirus RBD panel, with addition of trypsin only mildly improving entry for some of the

clade 2 and 4 viruses (Fig. 3a). In contrast, Huh-7.5 cells were susceptible to clade 1

merbecovirus RBDs (Fig. 3b). Notably, addition of trypsin improved entry for some of the clade 4

viruses for both cell lines (eg, ErinCov-12-19) (Fig. 3).

Merbecovirus compatibility with known human coronavirus receptors in human cells

The diverse panel of merbecovirus pseudotypes was tested on 293T cell lines stably transduced

to express the known human coronavirus receptors: amino peptidase N (APN), dipeptidyl

peptidase IV (DPP4), or Angiotensin converting enzyme-2 (ACE2). In the absence of trypsin,

untransduced 293T cells exhibited poor susceptibility to the entire panel and APN cells were only

susceptible to entry with the positive control: human coronavirus 229E, which is known to use

APN (Fig. 4a, Extended Data Fig. 2a-b). Cells transduced with human DPP4 exhibited clear

susceptibility to clade 1 merbecovirus RBDs but not the other clades (Figure 4A, Extended Data

Fig. 2c), while cells expressing human ACE2 exhibited low level susceptibility to SARS-CoV-2

spike and clade 3 merbecovirus RBDs (Fig. 4a, Extended Data Fig. 2d). Exogenous trypsin

moderately increased entry in untransduced 293Ts for human coronaviruses MERS-CoV, which

uses human DPP4 and SARS-CoV-2, which uses human ACE2, suggesting low levels of the

human receptors were present on 293Ts (Extended Data Fig. s2a). To reduce the background

signal and asses only the increase in entry resulting from the transduced receptor, the relative

entry signal was normalized against the no-receptor cell line (Fig. 4). In the presence of trypsin,

DPP4 only mediated clade 1 entry, while ACE2 cells exhibited low susceptibility to clade 1, 2, and

4 and high susceptibility to clade 3 merbecovirus RBDs (Fig. 4b). APN cells remained susceptible

to HCoV-229E entry in the presence of trypsin. While entry signal under 10x is not considered as

robust in this assay, there were a large number of viruses that seemed to have low level ACE2

compatibility (Fig. 4b; Extended Data Fig. 2d).

Species-specific receptor use across the merbecovirus RBD clades

Human cells expressing human ACE2 exhibited low level susceptibility to a wide range of

merbeocviruses. To investigate if these viruses are compatible with receptors from their cognate

host species, ACE2 and DPP4 was synthesized from European hedgehogs (*Erinaceus*

europaeus), the natural host species for the ErinCoVs (table 1; merbecovirus RBD clade 4), and

ACE2 was synthesized from Japanese house bats (*Pipistrellus abramus*): the natural host species

for the HKU5 viruses (table 1; merbecovirus RBD clade 2). 293T cells transfected with hedgehog

DPP4 or ACE2 were not particularly susceptible to any of the clades, with or without trypsin

(Figure 5). However, 293T transfected with ACE2 from Japanese house bats exhibited clear

susceptibility to the entire HKU5 cluster of clade 2 merbecovirus RBDs – with or without trypsin

(Fig. 5; Extended data Fig. 2). In a confirmatory assay, *Pipstrellus abramus* ACE2 specifically

mediated only HKU5 RBD entry in BHK cells, which are otherwise not susceptible to entry with

any ACE2 or DPP4-compatible betacoronavirus (Fig. 6). Taken together, this clear and HKU5-

specificific entry strongly suggest ACE2 is the natural receptor for the HKU5 viruses (Fig. 5 and

6).

DISCUSSION:

Compared to sarbecoviruses, which share 90% nucleotide sequence similarity and above,

the diversity in merbecoviruses is higher, initially complicating the chimeric spike approach utilized

in this study (Fig. 1). Wide variability between merbecovirus host species (hedgehogs, bats,

camels, humans and pangolins) also poses unforeseen challenges when studying cross-species

interactions. The chimeric spike approach presented here aims to eliminate some of the

interspecies challenges because using the human-compatible MERS-CoV spike in experiments

performed in human cells should reduce off-target phenotypes relating to cell attachment and

protease processing. Addition of exogenous protease further reduces entry phenotypes not

related to the RBD. Importantly, multiple studies have now shown that chimeric coronavirus spikes

faithfully re-create entry of both full spikes and wild-type virus[24-29,48].

Merbecovirus RBDs can be classified into four clades based on sequence, characteristic

indels, and entry phenotype (Fig. 2; Extended Data Fig. 4, table 1): clade 1 viruses can use DPP4

from multiple species, clade 2 viruses have species-specific ACE2 preference, and clade 3 can

use ACE2 from a wider range of host species. No clear entry was observed with clade 4 viruses

and known receptors, but nevertheless, human-cell entry was observed with ErinCoVs in the

presence of trypsin (Fig. 3-5, Extended Data Fig. 2-3). Because clade 4 entry seemed most

apparent when ACE2 was provided, it is possible these viruses use ACE2 from their host species

and that the approach taken in this study is not suitable for detecting their entry. For example,

hedgehogs have at least two reported transcript variants for ACE2 and three transcript variants

for DPP4 – only one of each was tested here (see methods). Additionally, while the chimeric spike

helps to reduce unintended entry barriers, barriers likely remain, including receptor compatibility

with other critical host cell-membrane factors, which is difficult to anticipate or control for[49].

Alternatively, clade 4 merbecovirus RBDs may use a receptor that is neither DPP4 nor ACE2,

analogous to sarbecovirus RBD clade 2 viruses, which have also can infect human cells through

an unidentified route (Extended Data Fig. 4)[24-27].

MERS-CoV spike can rapidly adapt to sub-optimal interactions with DPP4[50] and the

ongoing SARS-CoV-2 pandemic underscores the ability of spike to adapt and correct sub-optimal

ACE2 interactions while simultaneously escaping the immune response[51-54]. Other research has

shown that clade 3 merbecoviruses gain even broader host ACE2 tropism with single point

mutations at 305 and 510 – with the mutation T510F significantly increasing tropism for human

ACE2[21,22]. Thus, even low-level receptor compatibility observed for the merbecoviruses is

concerning and underscores the need for broadly protective coronavirus vaccines.

HKU5 viruses were first sequenced in 2006, shown to be DPP4-independent in 2014, and

demonstrated to replicate in vero cells in the presence of exogenous trypsin in 2019[12,46,55]. A

previous study identified ACE2 as the receptor for the clade 3 merbecoviruses such as PDF2180

and closely-related NeoCoV, but was unable to identify the receptor for the clade 2 HKU5

viruses[22]. However, this earlier study assessed the ability of HKU5 to interact with ACE2 from

Pipistrellus pipistrellus bats and several other Yangochiropteran bats, which are not the true host-

species for HKU5. The clear preference of HKU5 for *Pipistrellus abramus* ACE2, and the

simultaneous poor use of *Pipistrrellus abramus* ACE2 use from other clade 2 RBDs further

demonstrates the tight species specificity found within clade 2 RBDs (Fig. 5 and 6). *Pipistrellus*

abramus ACE2 varies from previously tested *P. pipistrellus* and *P. khuli* ACE2s at residues 328

and 329, in close proximity to the interaction previously reported with NeoCoV/PDF2180 RBDs[22].

Similarly, some sarbecoviruses are capable of using ACE2 from diverse species, while other

sarbecoviruses can only interact with ACE2 from their own cognate species (Extended Data Fig.

4)[29,48].

A clade 1 RBD pangolin merbecovirus has been shown to use human DPP4 and was

isolated in Vero81 cells[56]. While Vero cells are frequently used in virus isolation attempts, data

from this study suggests that Huh-7.5 cells may be a more suitable cell line for isolating DPP4-

dependent merbecoviruses and perhaps even some clade 4 merbecoviruses. Further, MERS-

CoV grown in Vero cells has been shown to quickly accumulate tissue-culture adaptations in the

spike protein driving the need for alternative cell culture models[57].

Merbecoviruses are yet another type of beta-coronavirus found in wildlife with potential

human cell compatibility and a possibility to cause severe disease in humans. Japanese house

bats (*Pipistrellus abramus*) are the natural host for HKU5 coronaviruses and are synanthropic,

roosting in human dwellings and structures, and therefore pose repeat opportunities for human

exposure[46,58]. Luckily, clade 2 merbecovirus RBDs exhibit low human ACE2 compatibility,

although efforts should still be made to reduce contact with these animals (Fig. 4, Extended Data

Fig. 2). Developing universal vaccines capable of thwarting multiple viruses is a daunting but

essential challenge. The work presented here will serve as a guide to understanding

merbecovirus cell entry and subsequent spillover risk.

METHODS:

Cells

293T, VeroE6, Huh-7.5 and BHK-21 cells were maintained in DMEM supplemented with 10% FBS, penicillin-streptomycin, and L-glutamine and maintained at 37°C, 5% CO_2. Cells stably expressing coronavirus receptors were previously described[28].

Plasmids

MERS-CoV/EMC12 spike (accession number JX869059) was codon optimized, appended with C-terminal FLAG tag, and silent mutations were introduced near spike amino acids 341 and 617 to generate AflII and NotI restriction digest sites. RBDs were codon optimized and synthesized (IDT DNA) as previously described for sarbecoviruses[24]. Lyophilized RBD fragments were resuspended in water and used in downstream Infusion based cloning (Takara Bio) reactions to assemble full-length chimeric spike proteins. *Erinaceus europaeus* ACE2, transcript variant 1 (genbank XM_060183012.1), *Pipistrellus abramus* ACE2 (genbank GQ262782.1) and *Erinaceus europaeus* DPP4 (genbank XM_060177812.1) were codon optimized, split in half and synthesized for Infusion based DNA assembly. Splitting genes into multiple parts reduces synthesis time, costs and overall synthesized product complexity. A complete list of accession numbers for the spike panel can be found in table 1.

Westernblot

As described previously, 293T cells transfected with spike plasmids were lysed in 1% SDS lysis buffer and stored at -80°C until use. Lysates were clarified, boiled, reduced and separated on 10% Bis-Tris gels in MOPS buffer (Invitrogen NuPage).[24] Membranes were probed with FLAG:HRP conjugated monoclonal antibody (ThermoFisher) or GAPDH monoclonal (ThermoFisher) and reporbed with a secondary goat-anti-mouse:HRP poloyclonal mix (ThermoFisher).

Pseudotype entry assay

Pseudotype assays were performed as previously described[25,26]. For human ACE2, APN, and

DPP4 experiments, 293T cells stably transduced with coronavirus receptors were seeded in black

96-well plates and subsequently infected 24 hours with equal volumes of viral pseudotypes.

Pseudotypes were combined with either HBSS or HBSS containing trypsin (not TPCK-treated) to

achieve a final concentration of 100μg/mL trypsin. Cells were washed once with cold PBS,

centrifuged with pseudotypes at 4°C, 1200xG for 1 hour and incubated at 37°C overnight.

Luciferase was measured using Brioght-Glo reagent (Promega) approximately 18 hours post-

infection. Relative entry was calculated by dividing raw lucifefrase values for each spike by the

signal for no-spike pseudotypes. Normalized entry was calculated for the receptors by dividing

the relative entry values for each spike and receptor by the signal for cells without receptor.

Phylogenetic analysis

As described previously, the amino acid sequences for spike receptor binding domains aligned

using Clustal multiple sequence alignment software and default parameters. A maximum

likelihood phylogenetic tree was produced with PhyML v. 3.0[59]. The 'WAG' matrix +G model of

amino acid substitution was selected by the Smart Model Selection method with 1000 bootstrap

replicates[60]. A tree was then visualized as a cladogram with FigTree v1.4.4

(https://github.com/rambaut/figtree).

Structural depictions

MERS-CoV spike RBD bound to DPP4 (PDB ID: 4L72)[61] and the MERS-CoV spike trimer (PDB

ID: 5X59)[62] were visualized in PyMol (version 2.4.0).

Acknowledgements

I would like to thank Dr. Stephanie Seifert, Dr. Pamela Bjorkman, Dr. Alex Cohen, Ziyan Wu and Dr. Tyler Starr for providing valuable insights and comments on the work and manuscript. Research reported in this publication was supported by the National Institute Of Allergy And Infectious Diseases of the National Institutes of Health (NIAID/NIH) under Award Number R01AI179720. The content is solely the responsibility of the author and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES:

Zaki, A. M., van Boheemen, S., Bestebroer, T. M., Osterhaus, A. D. & Fouchier, R. A. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**, 1814-1820, doi:10.1056/NEJMoa1211721 (2012).

Dudas, G., Carvalho, L. M., Rambaut, A. & Bedford, T. MERS-CoV spillover at the camel-human interface. *Elife* **7**, doi:10.7554/eLife.31257 (2018).

Dudas, G. & Rambaut, A. MERS-CoV recombination: implications about the reservoir and potential for adaptation. *Virus Evol* **2**, vev023, doi:10.1093/ve/vev023 (2016).

Moreno, A. *et al.* Detection and full genome characterization of two beta CoV viruses related to Middle East respiratory syndrome from bats in Italy. *Virol J* **14**, 239, doi:10.1186/s12985-017-0907-1 (2017).

Luo, C. M. *et al.* Discovery of Novel Bat Coronaviruses in South China That Use the Same Receptor as Middle East Respiratory Syndrome Coronavirus. *J Virol* **92**, doi:10.1128/JVI.00116-18 (2018).

Lau, S. K. *et al.* Genetic characterization of Betacoronavirus lineage C viruses in bats reveals marked sequence divergence in the spike protein of pipistrellus bat coronavirus HKU5 in Japanese pipistrelle: implications for the origin of the novel Middle East respiratory syndrome coronavirus. *J Virol* **87**, 8638-8650, doi:10.1128/JVI.01055-13 (2013).

Ithete, N. L. *et al.* Close relative of human Middle East respiratory syndrome coronavirus in bat, South Africa. *Emerg Infect Dis* **19**, 1697-1699, doi:10.3201/eid1910.130946 (2013).

He, W. T. *et al.* Virome characterization of game animals in China reveals a spectrum of emerging pathogens. *Cell* **185**, 1117-1129 e1118, doi:10.1016/j.cell.2022.02.014 (2022).

De Sabato, L. *et al.* Can Coronaviruses Steal Genes from the Host as Evidenced in Western European Hedgehogs by EriCoV Genetic Characterization? *Viruses* **12**, doi:10.3390/v12121471 (2020).

Chen, J. *et al.* A bat MERS-like coronavirus circulates in pangolins and utilizes human DPP4 and host proteases for cell entry. *Cell* **186**, 850-863 e816, doi:10.1016/j.cell.2023.01.019 (2023).

Wang, Q. *et al.* Bat origins of MERS-CoV supported by bat coronavirus HKU4 usage of human receptor CD26. *Cell Host Microbe* **16**, 328-337, doi:10.1016/j.chom.2014.08.009 (2014).

Yang, Y. *et al.* Receptor usage and cell entry of bat coronavirus HKU4 provide insight into bat-to-human transmission of MERS coronavirus. *Proc Natl Acad Sci U S A* **111**, 12516-12521, doi:10.1073/pnas.1405889111 (2014).

Anthony, S. J. *et al.* Further Evidence for Bats as the Evolutionary Source of Middle East Respiratory Syndrome Coronavirus. *MBio* **8**, doi:10.1128/mBio.00373-17 (2017).

Raj, V. S. *et al.* Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. *Nature* **495**, 251-254, doi:10.1038/nature12005 (2013).

Park, J. E. *et al.* Proteolytic processing of Middle East respiratory syndrome coronavirus spikes expands virus tropism. *Proc Natl Acad Sci U S A* **113**, 12262-12267, doi:10.1073/pnas.1608147113 (2016).

Li, W. *et al.* Identification of sialic acid-binding function for the Middle East respiratory syndrome coronavirus spike glycoprotein. *Proc Natl Acad Sci U S A* **114**, E8508-E8517, doi:10.1073/pnas.1712592114 (2017).

Li, F. Receptor recognition and cross-species infections of SARS coronavirus. *Antiviral Res* **100**, 246-254, doi:10.1016/j.antiviral.2013.08.014 (2013).

Li, F. Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol* **3**, 237-261, doi:10.1146/annurev-virology-110615-042301 (2016).

19   Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike receptor-
binding domain complexed with receptor. *Science* **309**, 1864-1868,
doi:10.1126/science.1116480 (2005).
20   Li, W. *et al.* Receptor and viral determinants of SARS-coronavirus adaptation to human
ACE2. *EMBO J* **24**, 1634-1643, doi:10.1038/sj.emboj.7600640 (2005).
21   Ma, C. *et al.* Broad host tropism of ACE2-using MERS-related coronaviruses and
determinants restricting viral recognition. *Cell Discov* **9**, 57, doi:10.1038/s41421-023-
00566-8 (2023).
22   Xiong, Q. *et al.* Close relatives of MERS-CoV in bats use ACE2 as their functional
receptors. *Nature* **612**, 748-757, doi:10.1038/s41586-022-05513-3 (2022).
23   Ma, C. *et al.* Identification of ACE2 as the Entry Receptor for Two Novel European Bat
Merbecoviruses. *biorrXiv*.
24   Letko, M., Marzi, A. & Munster, V. Functional assessment of cell entry and receptor
usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nature Microbiology*,
doi:10.1038/s41564-020-0688-y (2020).
25   Guo, H. *et al.* Isolation of ACE2-dependent and -independent sarbecoviruses from
Chinese horseshoe bats. *J Virol* **97**, e0039523, doi:10.1128/jvi.00395-23 (2023).
26   Guo, H. *et al.* ACE2-Independent Bat Sarbecovirus Entry and Replication in Human and
Bat Cells. *mBio* **13**, e0256622, doi:10.1128/mbio.02566-22 (2022).
27   Khaledian, E. *et al.* Sequence determinants of human-cell entry identified in ACE2-
independent bat sarbecoviruses: A combined laboratory and computational network
science approach. *EBioMedicine* **79**, 103990, doi:10.1016/j.ebiom.2022.103990 (2022).
28   Seifert, S. N. *et al.* An ACE2-dependent Sarbecovirus in Russian bats is resistant to
SARS-CoV-2 vaccines. *PLoS Pathog* **18**, e1010828, doi:10.1371/journal.ppat.1010828
(2022).
29   Roelle, S. M., Shukla, N., Pham, A. T., Bruchez, A. M. & Matreyek, K. A. Expanded ACE2
dependencies of diverse SARS-like coronavirus receptor binding domains. *PLoS Biol* **20**,
e3001738, doi:10.1371/journal.pbio.3001738 (2022).
30   Glowacka, I. *et al.* Evidence that TMPRSS2 activates the severe acute respiratory
syndrome coronavirus spike protein for membrane fusion and reduces viral control by
the humoral immune response. *J Virol* **85**, 4122-4134, doi:10.1128/JVI.02232-10 (2011).
31   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278,
doi:10.1016/j.cell.2020.02.052 (2020).
32   Shirato, K., Kawase, M. & Matsuyama, S. Middle East respiratory syndrome coronavirus
infection mediated by the transmembrane serine protease TMPRSS2. *J Virol* **87**, 12552-
12561, doi:10.1128/JVI.01890-13 (2013).
33   Shirato, K., Kawase, M. & Matsuyama, S. Wild-type human coronaviruses prefer cell-
surface TMPRSS2 to endosomal cathepsins for cell entry. *Virology* **517**, 9-15,
doi:10.1016/j.virol.2017.11.012 (2018).
34   Millet, J. K. & Whittaker, G. R. Host cell entry of Middle East respiratory syndrome
coronavirus after two-step, furin-mediated activation of the spike protein. *Proc Natl Acad
Sci U S A* **111**, 15214-15219, doi:10.1073/pnas.1407087111 (2014).
35   Matsuyama, S. *et al.* Middle East Respiratory Syndrome Coronavirus Spike Protein Is
Not Activated Directly by Cellular Furin during Viral Entry into Target Cells. *J Virol* **92**,
doi:10.1128/JVI.00683-18 (2018).
36   Wang, Z. *et al.* Loss of furin site enhances SARS-CoV-2 spike protein pseudovirus
infection. *Gene* **856**, 147144, doi:10.1016/j.gene.2022.147144 (2023).
37   Hoffmann, M., Kleine-Weber, H. & Pohlmann, S. A Multibasic Cleavage Site in the Spike
Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. *Mol Cell* **78**, 779-
784 e775, doi:10.1016/j.molcel.2020.04.022 (2020).

38   Bestle, D. *et al.* TMPRSS2 and furin are both essential for proteolytic activation of
SARS-CoV-2 in human airway cells. *Life Sci Alliance* **3**, doi:10.26508/lsa.202000786
(2020).
39   Fukuma, A. *et al.* Inability of rat DPP4 to allow MERS-CoV infection revealed by using a
VSV pseudotype bearing truncated MERS-CoV spike protein. *Arch Virol* **160**, 2293-
2300, doi:10.1007/s00705-015-2506-z (2015).
40   Petit, C. M. *et al.* Genetic analysis of the SARS-coronavirus spike glycoprotein functional
domains involved in cell-surface expression and cell-to-cell fusion. *Virology* **341**, 215-
230, doi:10.1016/j.virol.2005.06.046 (2005).
41   Chen, H. Y. *et al.* Cytoplasmic Tail Truncation of SARS-CoV-2 Spike Protein Enhances
Titer of Pseudotyped Vectors but Masks the Effect of the D614G Mutation. *J Virol* **95**,
e0096621, doi:10.1128/JVI.00966-21 (2021).
42   Adney, D. R. *et al.* Replication and shedding of MERS-CoV in upper respiratory tract of
inoculated dromedary camels. *Emerg Infect Dis* **20**, 1999-2005,
doi:10.3201/eid2012.141280 (2014).
43   Alagaili, A. N. *et al.* Middle East respiratory syndrome coronavirus infection in dromedary
camels in Saudi Arabia. *MBio* **5**, e00884-00814, doi:10.1128/mBio.00884-14 (2014).
44   Chu, D. K. *et al.* MERS coronaviruses in dromedary camels, Egypt. *Emerg Infect Dis* **20**,
1049-1053, doi:10.3201/eid2006.140299 (2014).
45   Chu, D. K. *et al.* Middle East respiratory syndrome coronavirus (MERS-CoV) in
dromedary camels in Nigeria, 2015. *Euro Surveill* **20**, doi:10.2807/1560-
7917.ES.2015.20.49.30086 (2015).
46   Woo, P. C. *et al.* Molecular diversity of coronaviruses in bats. *Virology* **351**, 180-187,
doi:10.1016/j.virol.2006.02.041 (2006).
47   van Boheemen, S. *et al.* Genomic characterization of a newly discovered coronavirus
associated with acute respiratory distress syndrome in humans. *mBio* **3**,
doi:10.1128/mBio.00473-12 (2012).
48   Starr, T. N. *et al.* ACE2 binding is an ancestral and evolvable trait of sarbecoviruses.
Nature, doi:10.1038/s41586-022-04464-z (2022).
49   Earnest, J. T. *et al.* The tetraspanin CD9 facilitates MERS-coronavirus entry by
scaffolding host cell receptors and proteases. *PLoS Pathog* **13**, e1006546,
doi:10.1371/journal.ppat.1006546 (2017).
50   Letko, M. *et al.* Adaptive Evolution of MERS-CoV to Species Variation in DPP4. *Cell Rep*
24, 1730-1737, doi:10.1016/j.celrep.2018.07.045 (2018).
51   Zhang, S. *et al.* Bat and pangolin coronavirus spike glycoprotein structures provide
insights into SARS-CoV-2 evolution. *Nat Commun* **12**, 1607, doi:10.1038/s41467-021-
21767-3 (2021).
52   Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform
on virus evolution and furin-cleavage effects. *Nat Struct Mol Biol* **27**, 763-767,
doi:10.1038/s41594-020-0468-7 (2020).
53   Wrobel, A. G. *et al.* Evolution of the SARS-CoV-2 spike protein in the human host. *Nat*
Commun **13**, 1178, doi:10.1038/s41467-022-28768-w (2022).
54   Calvaresi, V. *et al.* Structural dynamics in the evolution of SARS-CoV-2 spike
glycoprotein. *Nat Commun* **14**, 1421, doi:10.1038/s41467-023-36745-0 (2023).
55   Menachery, V. D. *et al.* Trypsin Treatment Unlocks Barrier for Zoonotic Bat Coronavirus
Infection. *J Virol* **94**, doi:10.1128/JVI.01774-19 (2020).
56   Xia, L. Y. *et al.* Isolation and characterization of a pangolin-borne HKU4-related
coronavirus that potentially infects human-DPP4-transgenic mice. *Nat Commun* **15**,
1048, doi:10.1038/s41467-024-45453-2 (2024).

57   Scobey, T. *et al.* Reverse genetics with a full-length infectious cDNA of the Middle East
respiratory syndrome coronavirus. *Proc Natl Acad Sci U S A* **110**, 16157-16162,
doi:10.1073/pnas.1311542110 (2013).
58   Wang, W. *et al.* Airport noise disturbs foraging behavior of Japanese pipistrelle bats. *Ecol*
Evol **12**, e8976, doi:10.1002/ece3.8976 (2022).
59   Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood
phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307-321,
doi:10.1093/sysbio/syq010 (2010).
60   Lefort, V., Longueville, J. E. & Gascuel, O. SMS: Smart Model Selection in PhyML. *Mol*
Biol Evol **34**, 2422-2424, doi:10.1093/molbev/msx149 (2017).
61   Wang, N. *et al.* Structure of MERS-CoV spike receptor-binding domain complexed with
human receptor DPP4. *Cell Res* **23**, 986-993, doi:10.1038/cr.2013.92 (2013).
62   Yuan, Y. *et al.* Cryo-EM structures of MERS-CoV and SARS-CoV spike glycoproteins
reveal the dynamic receptor binding domains. *Nat Commun* **8**, 15092,
doi:10.1038/ncomms15092 (2017).

Fig. 1. Identification of the functional RBD in merbecoviruses. (a) The receptor binding

domain in merbecoviruses engages the host receptor, DPP4 (PDB ID: 4L72, 5X59). Conserved

glycines marking the functional RBD boundaries are indicated in yellow spheres. **(b)** Overview

of MERS-CoV spike design for RBD replacement. **(c)** Chimeric spikes are used to generate

single-cycle pseudotyped VSV reporter particles. **(d)** Overview of chimeric spike designs. **(e)**

Amino acid sequence alignment of region flanking the RBD in merbecoviruses. **(f)** 293T cell

lines expressing indicated receptors were infected with pseudotypes and luciferase was

measured for entry. Cells were infected in quadruplicate.

Fig. 2. Diversity in merbecovirus RBDs. (a) Total sequence diversity in merbecoviruses

encompasses approximately 34 RBD sequences **(b)** Cladogram showing the relationship of

RBD amino acid sequences **(c)** Schematic overview of RBD clades, indicating the approximate

size and location of indels characteristic to each merbecovirus RBD clade. **(d)** Westernblot for

FLAG (spike) expression of chimeric spikes.

Fig. 3. Susceptibility of common cell lines to merbecovirus entry. (a) VeroE6 cells or **(b)**

Huh-7.5 cells were infected with VSV pseudotype panel bearing indicated chimeric spikes, in

quadruplicate. Luciferase was measured for cell entry. Dashed line indicates arbitrary reliability

cut-off at 10-fold signal over background.

Fig. 4. Clade-specific use of human coronavirus receptors 293T cells transduced with

known human coronavirus receptors were infected in **(a)** the absence of trypsin or **(b)** in the

presence of 100μg/mL trypsin. Entry relative to no spike was compared to entry in cells without

receptor to produce normalized entry values. Spike sequences for MERS-CoV/EMC12, SARS-

CoV-2 and HCoV-229E are full-length, S1/S2 domains from these viruses, all other spikes are

MERS-CoV spike chimeras with the indicated RBD.

601
602

Fig. 5. ACE2 from Japanese house bats mediates HKU5 entry in human cells. 293T cells

transiently transfected with species' orthologues of coronavirus receptors were infected in **(a)**

the absence of trypsin or **(b)** in the presence of 100µg/mL trypsin. Entry relative to no spike was

compared to entry in cells without receptor to produce normalized entry values. Spike

sequences for MERS-CoV/EMC12, SARS-CoV-2 and HCoV-229E are full-length, S1/S2

domains from these viruses, all other spikes are MERS-CoV spike chimeras with the indicated

RBD.

Fig. 6. ACE2 from Japanese house bats specifically mediates HKU5 entry. BHK-21 cells were transiently transfected with empty vector or *Pipistrellus abramus* ACE2 and infected in the absence of exogenous trypsin. Spike sequences for MERS-CoV/EMC12 and SARS-CoV-2 are full-length, S1/S2 domains from these viruses, all other spikes are MERS-CoV spike chimeras with the indicated RBD. Dashed line indicates arbitrary reliability cut-off at 10-fold signal over background.

Table 1. Diverse merbecovirus panel encompassing global diversity

	Virus	Accession number	Receptor Clade	Host species	Location
1	ErinCoV-19-18	MW246796	4	*Erinaceus europaeus*	Italy
2	ErinCoV-88-1-18	MW246795	4	*Erinaceus europaeus*	Italy
3	ErinCoV-1-18	MW246797	4	*Erinaceus europaeus*	Italy
4	ErinCoV-12-19	MW246800	4	*Erinaceus europaeus*	Italy
5	ErinCoV-15-19	MW246802	4	*Erinaceus europaeus*	Italy
6	HKU31-HeB-MO1	OM451213	4	*Erinaceus amurensis*	China
7	ErinCoV-17-18	MW246798	4	*Erinaceus europaeus*	Italy
8	ErinCoV-11-19	MW246799	4	*Erinaceus europaeus*	Italy
9	PML-PHE1-2011	KC869678	3	*Neoromicia capensis*	South Africa
10	PDF2180	KX574227	3	*Pipistrellus hesperidus*	Uganda
11	HKU5-17s	AGP04930	2	*Pipistrellus abramus*	Hong Kong
12	HKU5-24s	AGP04936	2	*Pipistrellus abramus*	Hong Kong
13	HKU5-33s	KC522104	2	*Pipistrellus abramus*	Hong Kong
14	HKU5-27s	AGP04938	2	*Pipistrellus abramus*	Hong Kong
15	HKU5-20s	AGP04933	2	*Pipistrellus abramus*	Hong Kong
16	HKU5-28s	AGP04939	2	*Pipistrellus abramus*	Hong Kong
17	HKU5-32s	AGP04942	2	*Pipistrellus abramus*	Hong Kong
18	HKU5-21s	AGP04934	2	*Pipistrellus abramus*	Hong Kong
19	HKU5-19s	AGP04932	2	*Pipistrellus abramus*	Hong Kong
20	HKU5-1-LMH03f	NC009020	2	*Pipistrellus abramus*	China
21	HKU5-31s	AGP04941	2	*Pipistrellus abramus*	Hong Kong
22	HKU5-22s	AGP04935	2	*Pipistrellus abramus*	Hong Kong
23	HKU5-30s	AGP04940	2	*Pipistrellus abramus*	Hong Kong
24	HSavi-2011	MG596802	2	*Hypsugo savii*	Italy
25	PKhuli-2011	MG596803	2	*Pipistrellus kuhlii*	Italy
26	Bt-CoV-SC2013	KJ473821	2	*Vespertilio superans*	China
27	MjHKU4r-CoV-1	OQ786862	1	*Manis javanica*	China (Smuggled Malayan pangolins)
28	HKU4-2s	AGP04916	1	*Tylonycteris pachypus*	Hong Kong
29	HKU4-1-B04f	NC009019	1	*Tylonycteris pachypus*	Hong Kong
30	HKU4-10s	AGP04924	1	*Tylonycteris pachypus*	Hong Kong
31	HKU4/SM3A	MW218395	1	*Tylonycteris pachypus*	Hong Kong
32	HKU4-3s	AGP04917	1	*Tylonycteris pachypus*	Hong Kong
33	NL13845	MG021451	1	*Tylonycteris pachypus*	China
34	Bt-CoV 422	MG021452	1	*Tylonycteris pachypus*	China
35	MERS/EMC12	JX869059	1	*Homo sapiens*	Saudi Arabia

Extended Data Figures

 Extended Data Fig. 1. Optimizing MERS-CoV spike for MerbecoType. (a) Overview of S1/S2

and S2' furin sites in MERS-CoV spike. **(b)** 293T cells transduced with human ACE2 or (**c**)

human DPP4 were infected with indicated spikes and luciferase was measured for entryCells

were infected in quadruplicate.

Extended Extended Data Fig. 2. Relative data used in graphs for figure 4 – human

receptors. 293Tcells transduced to express **(a)** no receptor **(b)** human APN **(c)** human DPP4 or

(d) human ACE2 were infected in the absence or presence of 100µg/mL trypsin, in

quadruplicate. Luciferase was measured for entry and relative entry was calculated by

comparing to entry signal from bald pseudotypes. Spike sequences for MERS-CoV/EMC12,

SARS-CoV-2 and HCoV-229E are full-length, S1/S2 domains from these viruses, all other

spikes are MERS-CoV spike chimeras with the indicated RBD. Dashed line indicates arbitrary

reliability cut-off at 10-fold signal over background.

Extended Data Fig. 3. Relative data used in graphs for figure 5 – animal receptors. 293T

cells transfected to express **(a)** *Erinaceus europaeus* ACE2 **(b)** *Erinaceus europaeus* DPP4 or

(c) *Pipistrellus abramus* ACE2 were infected in the absence or presence of 100µg/mL trypsin, in

quadruplicate. Luciferase was measured for entry and relative entry was calculated by

comparing to entry signal from bald pseudotypes. Spike sequences for MERS-CoV/EMC12,

SARS-CoV-2 and HCoV-229E are full-length, S1/S2 domains from these viruses, all other

spikes are MERS-CoV spike chimeras with the indicated RBD. Dashed line indicates arbitrary

reliability cut-off at 10-fold signal over background.

658
659
660
661 **Extended Data Fig. 4. Overview of beta-coronavirus subgenus RBD clades.** RBDs from

662 different beta-coronavirus subgenera share similar trends of clade assortment based on indels

663 and clade-specific entry phenotypes.

664
665