1   **_Eed_ controls craniofacial osteoblast differentiation and mesenchymal proliferation from**
2   **the neural crest**
3

4   Tim Casey-Clyde[1,2,3]*, S. John Liu[1,2,3], Juan Antonio Camara Serrano[4], Camilla Teng[5], Yoon-Gu
5   Jang[5], Harish N. Vasudevan[1,2], Jeffrey O. Bush[5], David R. Raleigh[1,2,3#]
6
[1]Department of Radiation Oncology, University of California San Francisco, San Francisco, CA,
USA
[2]Department of Neurosurgery, University of California San Francisco, San Francisco, CA, USA
[3]Department of Pathology, University of California San Francisco, San Francisco, CA, USA
[4]Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San
Francisco, CA, USA
[5]Department of Cell and Tissue Biology, University of California San Francisco, San Francisco,
CA, USA
15
16 [#]Author for correspondence: david.raleigh@ucsf.edu

17
18 _Running title_
19   _Eed_ and craniofacial development
20

21 _Summary statement_
22   Here we show the epigenetic regulator _Eed_ drives craniofacial osteoblast differentiation and
23 mesenchymal proliferation after induction of the neural crest, a multipotent progenitor cell
24 population   that   gives   rise   to   diverse   cell   types.

Abstract

The histone methyltransferase Polycomb repressive complex 2 (PRC2) is required for specification of the neural crest, and mis-regulation of the neural crest can cause severe congenital malformations. PRC2 is required for induction of the neural crest, but the embryonic, cellular, and molecular consequences of PRC2 activity after neural crest induction are incompletely understood. Here we show that *Eed*, a core subunit of PRC2, is required for craniofacial osteoblast differentiation and mesenchymal proliferation after induction of the neural crest. Integrating mouse genetics with single-cell RNA sequencing, our results reveal that conditional knockout of *Eed* after neural crest cell induction causes severe craniofacial hypoplasia, impaired craniofacial osteogenesis, and attenuated craniofacial mesenchymal cell proliferation that is first evident in post-migratory neural crest cell populations. We show that *Eed* drives mesenchymal differentiation and proliferation *in vivo* and in primary craniofacial cell cultures by regulating diverse transcription factor programs that are required for specification of post-migratory neural crest cells. These data enhance understanding of epigenetic mechanisms that underlie craniofacial development, and shed light on the embryonic, cellular, and molecular drivers of rare congenital syndromes in humans.

Introduction

The embryonic neural crest is a multipotent progenitor cell population that gives rise to peripheral neurons, glia, Schwann cells, melanocytes, and diverse mesenchymal cells such as osteoblasts, chondrocytes, fibroblasts, cardiac mesenchyme, and cardiomyocytes (Bronner and LeDouarin, 2012; Simões-Costa and Bronner, 2015). Following induction in the neural tube, cranial neural crest cells undergo dorsolateral migration to the pharyngeal arches and differentiate to form the craniofacial bones and cartilage (Minoux and Rijli, 2010). To do so, cranial neural crest cells express transcription factors such as *Sox9*, *Sox10,* and *Twist1* that specify cell fate decisions in derivates of the neural crest (Bertol et al., 2022; Cheung et al., 2005). Pre-migratory neural crest cell specification is partially controlled by the master epigenetic regulator Polycomb repressive complex 2 (PRC2), an H3K27 histone methyltransferase that is broadly responsible for chromatin compaction and transcriptional silencing (Margueron and Reinberg, 2011). The catalytic activity of PRC2 requires four core subunits: (1) enhancer of zeste homologue 1 (Ezh1) or Ezh2, (2) suppressor of zeste 12 (Suz12), (3) RBBP4 or RBBP7, and (4) embryonic ectoderm development (Eed) (Piunti and Shilatifard, 2021). Eed binds to H3K27 trimethylation peptides (H3K27me3) and stabilizes Ezh2 for allosteric activation of methyltransferase activity and on-chromatin spreading of H3K27

methylation. *Eed* is required for stem cell plasticity, pluripotency, and maintaining cell fate
decisions, but all PRC2 core subunits are required for embryonic development and loss of any
individual subunit is embryonic lethal around gastrulation(Faust et al., 1995; O'Carroll et al.,
2001; Pasini et al., 2004). *Eed* null embryos can initiate endoderm and mesoderm induction but
have global anterior-posterior patterning defects in the primitive streak (Faust et al., 1995;
Schumacher et al., 1996) and genome-wide defects in H3K27 methylation (Montgomery et al.,
2007).
Ezh2 has been studied in the context of pre-migratory neural crest development. In mice,
Wnt1-Cre Ezh2$^{Fl/Fl}$ embryos fail to develop skull and mandible structures due to de-repression of
Hox transcription factors in cranial neural crest cells (Kim et al., 2018; Schwarz et al., 2014),
and loss of *Ezh2* in mesenchymal precursor cells causes skeletal defects and craniosynostosis
in *Prrx1-Cre Ezh2$^{Fl/Fl}$* embryos (Dudakovic et al., 2015). Ezh2 also regulates neural crest
specification in xenopus (Tien et al., 2015), and *Col2-Cre Eed$^{Fl/Fl}$* mice have kyphosis and
accelerated hypertrophic differentiation due to de-repression of Wnt and TGF-β signaling in
chondrocytes (Mirzamohammadi et al., 2016). With respect to the other PRC2 core subunits,
genetic and molecular interactions allow *Eed* to repress Hox genes to maintain vertebral body
identity during mouse development (Kim et al., 2006), but the embryonic, cellular, and molecular
consequences of *Eed* activity in craniofacial development are unknown. To address this gap in
our understanding of epigenetic mechanisms that may contribute to craniofacial development,
we conditionally deleted *Eed* from the migratory neural crest and its derivatives. Our results
suggest that *Eed* is required for craniofacial osteoblast differentiation from post-migratory neural
crest mesenchymal stem cells, and that *Eed* regulates diverse transcription factor programs that
are required for mesenchymal cell proliferation, differentiation, and osteogenesis. More broadly,
these data show that *Eed* is required at early post-migratory stages in neural crest progenitors
of craniofacial structures.
84
Results and discussion
Loss of Eed after neural crest induction causes severe craniofacial malformations
To determine if *Eed* is required for the development of neural crest derivatives, homozygous
floxed *Eed* alleles (*Eed$^{Fl/Fl}$*) (Yu et al., 2009) were used to conditionally delete *Eed* following
neural crest cell induction using Cre under control of the *Sox10* promoter, which is expressed in
the migratory neural crest at embryonic day E8.75 and results in complete recombination in
post-migratory neural crest cells by E10.5 (Matsuoka et al., 2004; Niethamer et al., 2020).
Sox10-Cre Eed$^{Fl/Fl}$ mice were not recovered past postnatal day 0, suggesting that loss of *Eed*

following induction of the neural crest is embryonic lethal (Fig. 1A). *Sox10-Cre Eed^{Fl/Fl}* embryos

were recovered at expected genotypic frequencies from embryonic day E9.5 to E17.5 (Fig. 1A),

and there were no differences in the penetrance or severity of *Sox10-Cre Eed^{Fl/Fl}* phenotypes

whether Cre was maternally or paternally inherited, as has been reported for other phenotypes

arising in cells expressing *Sox10-Cre* (Crispino et al., 2011; Luo et al., 2020). No craniofacial

phenotypes could be overtly discerned at early post-migratory neural crest cell stages E10.5 or

E11.5, but craniofacial abnormalities were identified in *Sox10-Cre Eed^{Fl/Fl}* embryos starting at

E12.5 that increased in severity throughout the remainder of embryonic development compared

to controls (Fig. 1B). *Sox10-Cre Eed^{Fl/Fl}* phenotypes were broadly consistent with impaired

development of craniofacial structures (Deckelbaum et al., 2012; Ting et al., 2009), including

frontonasal and mandibular hypoplasia, a prominent telencephalon and exencephaly resulting

from underdevelopment of the viscerocranium and frontal calvarium, collapsed nasal cavity

bones, and microtia compared to controls (Fig. 1C). *Sox10-Cre Eed^{Fl/Fl}* embryos had increased

pupillary distance and craniofacial width (Fig. S1A,B) but did not have cardiac outflow tract

defects or other heart abnormalities that can arise from impaired neural crest differentiation

(Lindsay et al., 1999; Lindsay et al., 2001; Nakamura et al., 2009) (Fig. S1C-E; Movie 1, 2).

Whole mount DAPI imaging of *Sox10-Cre Eed^{Fl/Fl}* embryos revealed frontonasal dysplasia with

underdeveloped midface and mandible, and an irregular and corrugated facial structure

compared to controls (Fig. 1D).

Eed specifies craniofacial mesenchymal derivatives from the early post-migratory neural crest

Skeletal stains, computed tomography (microCT), histology, and immunofluorescence were

used to shed light on how loss of *Eed* impacts craniofacial development. Whole mount alcian

blue/alizarin red staining of *Sox10-Cre Eed^{Fl/Fl}* embryos revealed hypoplasia of the

viscerocranium, including reduced frontal, temporal, maxillary, and mandibular bones, and

complete loss of the tympanic ring and premaxillary and nasal bones compared to controls (Fig.

2A,B; Fig. S2A). MicroCT validated these findings, showing fragmentation and hypoplasia of the

frontal, temporal, maxillary, and mandibular bones, loss of frontal calvarial fusion, and absence

of tympanic ring and nasal bones (Fig. 2C; Movie 3, 4). H&E histology showed that frontal

calvarium reduction resulted in anteriorly displaced brain structures in *Sox10-Cre Eed^{Fl/Fl}*

embryos compared to controls, with the midbrain and cortex in the same coronal plane as the

nasal cavity and developing mandible (Fig. 2D; Fig. S2B). Most of the midfacial and mandible

were absent in *Sox10-Cre Eed^{Fl/Fl}* embryos, and the tongue and masseter muscles were

underdeveloped (Fig. 2D). Immunofluorescence demonstrated decreased staining for Runx2, a

transcription factor that is required for osteoblast differentiation and proliferation (Kawane et al.,

2018; Komori, 2009), and disorganized ALPL, a marker of mineralized bone (Liu et al., 2018), in

craniofacial tissues from E15.5 *Sox10-Cre Eed^{Fl/Fl}* embryos compared to controls (Fig. 2E). In

contrast to *Wnt1-Cre Ezh2^{Fl/Fl}* embryos (Schwarz et al., 2014), the craniofacial region of *Sox10-*

Cre Eed^{Fl/Fl} embryos had decreased immunofluorescence staining for Ki67 (Fig. 2F), a marker of

cell proliferation (Gerdes et al., 1984). E15.5 *Sox10-Cre Eed^{Fl/Fl}* embryos also had decreased

immunofluorescence staining for Vimentin (Vim), a marker of mesenchymal cells (Mendez et al.,

2010) (Fig. 2F), and heterogeneous immunofluorescence staining for Sox9, a marker of

chondrocytes that are derived from the neural crest (Mori-Akiyama et al., 2003; Yan et al., 2002)

(Fig. S2C,D), compared to controls. These results suggest craniofacial mesenchymal

differentiation, proliferation, and osteogenesis are impaired in *Sox10-Cre Eed^{Fl/Fl}* embryos

(Honoré et al., 2003; Kim et al., 2003).

Eed regulates craniofacial differentiation and proliferation in vivo and in primary cell cultures

To elucidate early molecular mechanisms underlying craniofacial phenotypes in *Sox10-Cre*

Eed^{Fl/Fl} embryos, immunofluorescence was performed on heads and primary craniofacial cell

cultures from E12.5, the earliest timepoint when craniofacial malformations were identified (Fig.

1B,D). Primary craniofacial cells were microdissected from the anterior and ventral layers of the

E12.5 mandible, dissociated into single-cell suspensions, and cultured for up to 1 week for

experimentation. Results were compared to primary craniofacial cell cultures that were derived

from E15.5 embryos using the same technique, and immunofluorescence of E12.5 and E15.5

heads.

In support of immunofluorescence results from E15.5 heads (Fig. 2E,F; Fig. S2C,D), staining

for Runx2 and Sox9 was decreased in craniofacial tissues from E12.5 *Sox10-Cre Eed^{Fl/Fl}* heads

compared to controls (Fig. 3A). Runx2, Sox9, and Ki67 were also decreased in primary *Sox10-*

Cre Eed^{Fl/Fl} craniofacial cell cultures from E12.5 compared to controls (Fig. 3B,C), and

immunofluorescence on E12.5 heads showed BrdU was decreased in *Sox10-Cre Eed^{Fl/Fl}*

samples compared to controls (Fig. S3A). These differences persisted in primary craniofacial

cell cultures from E15.5, at which point ALPL was also found to be suppressed in primary

Sox10-Cre Eed^{Fl/Fl} craniofacial cell cultures compared to controls (Fig. S3B,C). In contrast,

immunofluorescence intensity for Sox6, Sox10, and the collagen marker Col1a2 (Bowers et al.,

2023) were equivalent across genotypes in E12.5 and E15.5 primary craniofacial cell cultures

(Fig. S3B,C). Immunofluorescence intensity for Sox10 was also equivalent across genotypes in

E12.5 heads compared to controls, but Sox10 immunofluorescence was heterogeneous in

E12.5 *Sox10-Cre Eed^{Fl/Fl}* heads compared to controls (Fig. S3D).

Eed specifies craniofacial mesenchymal stem cell, osteoblast, and proliferating mesenchymal

cell fate from the neural crest

To define cell types and gene expression programs underlying craniofacial phenotypes in

Sox10-Cre Eed^{Fl/Fl} embryos, single-cell RNA sequencing was performed on litter-matched E12.5

Sox10-Cre Eed^{Fl/WT} and *Sox10-Cre Eed^{Fl/Fl}* heads (n=3 biological replicates per genotype).

Uniform manifold approximation and projection (UMAP) analysis of 63,730 single-cell

transcriptomes revealed 23 cell clusters (C0-C22) that were defined using automated cell type

classification (Ianevski et al., 2022), cell signature genes, cell cycle analysis, and differentially

expressed cluster marker genes (Fig. 4A; Fig. S4, S5; Table S1, S2). Differentiating osteoblasts

marked by *Runx2* (C0) and proliferating mesenchymal cells marked by Ki67 (C7) were enriched

in *Sox10-Cre Eed^{Fl/WT}* samples (Fig. 4B-D). Mesenchymal stem cells marked by *Col6a3* (C4) or

Dcn (C5) were enriched in *Sox10-Cre Eed^{Fl/Fl}* samples (Jang et al., 2016; Lamandé et al., 2006)

(Fig. 4B-D). There were subtle differences in the number of interneurons (C17, 1.4% versus

1.9% of cells, p=0.01), Schwann cells (C19, 1.0 versus 1.3% of cells, p=0.03), pericytes (C20,

0.7% vs 0.9%, p=0.05), and spinal neurons (C21, 0.5% vs 0.9% of cells, p=0.003) in *Sox10-Cre*

Eed^{Fl/WT} versus *Sox10-Cre Eed^{Fl/Fl}* samples, but each of these comprised a small minority of

recovered cell types (Student's t tests) (Table S1). There were no differences between

genotypes in the number of chondrocytes (C10, 3.1% versus 2.7%, p=0.09), fibroblasts (C11,

2.6% versus 2.7%, p=0.24), endothelia (C16, 1.9% vs 1.6%, p=0.22), hematopoietic cells, or

other cell types recovered at E12.5 (Student's t tests) (Table S1).

Differential expression analysis of single cell transcriptomes from differentiating osteoblasts

(C0), which were enriched in *Sox10-Cre Eed^{Fl/WT}* samples (Fig. 4c), compared to mesenchymal

stem cells marked by *Col6a3* (C4), which were enriched in *Sox10-Cre Eed^{Fl/Fl}* samples (Fig. 4C),

showed *Sox5* and *Sox6* were reduced in mesenchymal stem cells (Fig. 4E,F; Table S3),

consistent with the known role of Sox transcription factors in craniofacial specification and

development (Smits et al., 2001). Iroquois homeobox (Irx) transcription factors 3 and 5, which

contribute to craniofacial osteogenesis and mineralization (Bonnard et al., 2012; Cain et al.,

2016; Tan et al., 2020), were also suppressed in mesenchymal stem cells compared to

differentiating osteoblasts (Fig. 4E,F; Table S3). Differential expression analysis of single cell

transcriptomes from differentiating osteoblasts compared to mesenchymal stem cells marked by

Dcn (C5), which were also enriched in *Sox10-Cre Eed^{Fl/Fl}* samples (Fig. 4C), showed *Mecom*,

Trps1, and *Ptch1* were suppressed and *Sparc* and *Tnmd* were enriched in mesenchymal stem

cells (Fig. 4G,H; Table S4). Loss of *Mecom* causes craniofacial malformations in mice and

zebrafish (Shull et al., 2020), *Ptch1* is a key component of the Hedgehog pathway that is crucial

for craniofacial osteogenesis (Jeong et al., 2004; Metzis et al., 2013), and *Trps1* regulates

secondary palate and vibrissa development (Cho et al., 2019; Fantauzzo and Christiano, 2011).

Consistently, *Sox10-Cre Eed$^{Fl/Fl}$* samples had decreased vibrissa compared to controls (Fig. 1D;

Fig. S2B). *Sparc* drives morphogenesis of the pharyngeal arches and inner ear (Rotllant et al.,

2008), and *Tnmd* regulates mesenchymal differentiation (Shukunami et al., 2006). *Pax3* and

Pax7, key regulators of neural crest migration and development of skeletal structures

(Maczkowiak et al., 2010), were also suppressed in mesenchymal stem cells compared to

differentiating osteoblasts (Fig. 4G,H; Table S4), and differential expression analysis of single

cell transcriptomes from the two clusters of mesenchymal stem cells that were marked by

Col6a3 (C4) versus *Dcn* (C5) showed *Twist2*, *Irx3*, and *Irx5*, which regulate mesenchymal

differentiation to osteoblasts (Lee et al., 2000; Tamamura et al., 2017; Tan et al., 2020),

distinguished mesenchymal stem cell populations that were found in *Sox10-Cre Eed$^{Fl/WT}$* and

Sox10-Cre Eed$^{Fl/Fl}$ samples (Fig. S6A,B; Table S5) but were both enriched in *Sox10-Cre Eed$^{Fl/Fl}$*

samples compared to controls (Fig. 4C).

To determine if there were differences in the gene expression programs from differentiating

osteoblasts (C0), mesenchymal stem cells (C4, C5), or proliferating mesenchymal cells in

Sox10-Cre Eed$^{Fl/WT}$ versus *Sox10-Cre Eed$^{Fl/Fl}$* samples, differential expression analysis was

performed on single-cell transcriptomes from these clusters across genotypes (Table S6).

Consistent with craniofacial findings after loss of *Ezh2* (Kim et al., 2018; Schwarz et al., 2014)

and vertebral body findings after loss of *Eed* (Kim et al., 2006), Hox transcription factors were

broadly de-repressed in *Sox10-Cre Eed$^{Fl/Fl}$* samples compared to controls (Fig. S6C). PRC2

targets Sp7 in bone marrow stroma cells (Liu et al., 2013) and regulates Wnt signaling in dental

stem cells (Jing et al., 2016), and differential expression analysis showed that *Sp7* was enriched

in mesenchymal single-cell transcriptomes from *Sox10-Cre Eed$^{Fl/WT}$* samples while the Wnt and

stem cell regulator *Gata4* was enriched in mesenchymal single-cell transcriptomes from *Sox10-*

Cre Eed$^{Fl/Fl}$ samples (Fig. S6C). *Mki67*, *Pax3*, and *Pax7* were enriched in mesenchymal single-

cell transcriptomes from *Sox10-Cre Eed$^{Fl/WT}$* samples, while the cell cycle inhibitors *Cdkn2a* and

Cdkn2b were enriched in mesenchymal single-cell transcriptomes from *Sox10-Cre Eed$^{Fl/Fl}$*

samples (Fig. S6C). In sum, these data show that *Eed* specifies craniofacial osteoblast

differentiation and mesenchymal cell proliferation by regulating diverse transcription factor

programs that are required for specification of post-migratory neural crest cells, and that loss of

Eed inhibits mesenchymal stem cell differentiation and mesenchymal cell proliferation by de-
repressing transcription factors and inhibitors of the cell cycle.

Discussion

Epigenetic regulation of the neural crest, which differentiates into diverse mesenchymal
derivatives, is incompletely understood. Here we identify the PRC2 core subunit *Eed* as a potent
regulator of craniofacial development after induction of the neural crest. Conditional deletion of
Eed using *Sox10-Cre* causes severe craniofacial malformations that are consistent with
impaired differentiation and proliferation of cells arising from the pharyngeal arches (Frisdal and
Trainor, 2014). In support of this hypothesis, we observed gross defects in the development of
mesenchyme-derived tissues (Fig. 1, 2) that were consistent with our molecular and cellular
findings from *Sox10-Cre Eed^{Fl/Fl}* embryos (Fig. 2-4). Expression of key regulators of craniofacial
osteogenesis such as *Runx2, Irx3, Irx5, Mecom, Trps1, Ptch1, Pax3,* and *Pax7* were reduced in
Sox10-Cre Eed^{Fl/Fl} heads and primary craniofacial cell cultures compared to controls, and single-
cell RNA sequencing showed reduced craniofacial osteoblast differentiation and reduced
proliferation of mesenchymal cells in *Sox10-Cre Eed^{Fl/Fl}* heads compared to controls.

By using *Sox10-Cre*, which marks the migratory neural crest (Matsuoka et al., 2004), we
bypass potential roles of *Eed* in neural crest cell induction to examine its function in neural crest
cell development. The requirement of PRC2 for craniofacial development has been studied in
the context of *Ezh2* loss using *Wnt1-Cre* (Schwarz et al., 2014), but to our knowledge, our study
is the first report directly linking *Eed* to differentiation of neural crest mesenchymal derivatives.
To date, fifteen human missense mutations in *EED* have been reported as pathogenic for
Cohen-Gibson syndrome (Cohen and Gibson, 2016; Cohen et al., 2015; Goel et al., 2024). Like
Weaver syndrome, which arises from missense mutations in *EZH2* (Gibson et al., 2012;
Tatton-Brown et al., 2013), and Imagawa-Matsumoto syndrome, which arises from missense
mutations in *SUZ12* (Imagawa et al., 2018), Cohen-Gibson syndrome is a rare congenital
syndrome associated with craniofacial malformations, advanced bone age, intellectual disability,
and developmental delay (Spellicy et al., 2019). *EED* mutations in individuals with Cohen-
Gibson syndrome cluster in WD40 domains 3 through 5 and are predicted to (1) abolish
interaction with EZH2, (2) prevent histone methyltransferase activity, and (3) inhibit H3K27
trimethylation peptide binding. In support of these hypotheses, functional investigations have
shown that single nucleotide variants in the WD40 domain of *EED* abolish binding to *EZH2 in*
vitro (Denisenko et al., 1998). Mouse models encoding pathogenic *EZH2* missense variants,
which are predicted to result in loss of function of the PRC2 complex, phenocopy Weaver

syndrome and cause excess osteogenesis and skeletal overgrowth (Gao et al., 2023). In this
study, we show that loss of *Eed* after neural crest induction causes the opposite phenotype,
resulting in craniofacial hypoplasia due to impairments in osteogenesis, mesenchymal cell
proliferation, and mesenchymal stem cell differentiation. The discrepancy of missense variants
in *EED* causing gain in craniofacial osteogenic function in humans versus loss of *Eed* causing
loss in craniofacial osteogenic function in mice warrants further study, including investigation
using alternative Cre drivers to target different developmental stages of the neural crest. Despite
these discrepancies, it is notable that many of the phenotypes we observed after loss of *Eed* in
the neural crest of mice were similar to previously reported phenotypes after loss of *Ezh2* in the
neural crest cell in mice, with the exception of proliferation deficits after loss of *Eed* but not after
loss of *Ezh2* (Schwarz et al., 2014).

Materials and methods

Mice

Mice were maintained in the University of California San Francisco (UCSF) pathogen-free
animal facility in accordance with the guidelines established by the Institutional Animal Care and
Use Committee (IACUC) and Laboratory Animal Resource Center (LARC) protocol AN191840.
Mice were maintained in a 70°F, 50% humidity temperature-controlled barrier facilities under a
12-12h light cycle with access to food and water *ad libitum*. *Sox10-Cre* and *Eed^{Fl}* mice were
obtained from the Jackson Laboratory (B6;CBA-Tg(Sox10-cre)1Wdr/J, 025807 and B6;129S1-
Eed^{tm1Sho}/J, 022727, respectively). The presence of the floxed *Eed* allele was determined
through standard PCR genotyping using the following primers: 5'
GGGACGTGCTGACATTTTCT 3' (forward) and 5' CTTGGGTGGTTTGGCTAAGA 3' (reverse).
To generate embryos at specific time points, *Sox10-Cre^{tg+} Eed$^{Fl/WT}$* mice were bred overnight
with *Eed$^{Fl/Fl}$ mice.* Females were checked for copulation plugs in the morning and the presence
of a vaginal plug was designated as E0.5.

Mouse fetal echocardiography

Ultrasound studies were acquired with a Fujifilm Vevo 2100 Imaging system, and instrument
specifically designed for lab animal imaging studies. Pregnant females were anesthetized using
an isoflurane/oxygen mixturewith an isoflurane concentration of 3% during imaging. The
pregnancy date (at least p17) was checked to observe the correct development of fetal hearts.
Isoflurane was increased until 5% and the abdomen of pregnant females was open surgically to
expose the gravid uterus. The number of fetuses was counted, and craniofacial architecture was

evaluated for phenotypic confirmation. Once each fetus was defined as mutant or wild type,
echocardiography was performed to evaluate left ventricle anatomy and physiology, including
left ventricle size and contractility. Three different measurements were obtained in B and M
modes, and functional parameters, including heart rate, fraction shortening, ejection fraction,
and left ventricle mass were obtained.

Whole-mount embryo staining

Embryos were stained as previously described (Sandell et al., 2018). In brief, embryos were
harvested, and heads were fixed overnight in 4% PFA. Embryos were dehydrated through a
methanol series up to 100% before being treated with Dent's bleach (4:1:1, MeOH, DMSO, 30%
H_2O_2) for 2hr at 25°C. Embryos were serially rehydrated in methanol up to 25%, washed with
PBST, and stained with DAPI solution (1:40,000, Thermo Scientific, cat# 62248) overnight at
4°C. Samples were mounted in low melt agarose for imaging.

Whole-mount skeletal staining

Embryos were stained as previously described (Rigueur and Lyons, 2013). In brief, embryos
were dissected in cold PBS and scalded in hot water to facilitate the removal of eyes, skin, and
internal organs. Embryos were fixed in 95% ethanol overnight then subsequently incubated in
acetone overnight. Embryos were stained with Alcian Blue solution (Newcomer Supply, cat#
1300B) for 8hr, washed in 70% ethanol, and destained in 95% ethanol. Embryos were cleared in
95% ethanol and 1% KOH, and stained in Alizarin red S solution (SCBT, cat#130-22-3) for 4hr.
Embryos were stored in glycerol for imaging.

Micro-computed tomography (microCT)

microCT studies were acquired with a Perkin Elmer´s Quantum GX2 microCT Imaging
system, an instrument specifically designed for lab animal imaging studies. The samples were
imaged in Eppendorf tubes containing preservative media, which were placed on the scanning
bed and fixed with tape to minimize movement. Acquisition parameters were set as follows:
FOV 10mm, acquisition time 14 minutes, current voltage 70KV, amperage 114uA. Each study
was composed by a 512x512x512 voxels matrix with a spatial resolution of $0.018mm^3$. Study
reconstruction was based on Feldkamp´s method using instrument dedicated software. For
image analysis, tridimensional renders were obtained using 3D SLICER software. The threshold
range applied for bone segmentation was 300-1400 Hounsfield Units.

Cryosectioning and H&E staining

Embryos were fixed in 4% PFA overnight at 4°C and subjected to a sucrose gradient from
15% to 30% before embedding in OCT (Tissue-Tek, cat#4583) and storage at -80°C. Embryos
were sectioned at 25µm on a Leica CM1850 cryostat. Slides were immediately subjected to H&E
staining as previously described (Cardiff et al., 2014), or stored at -80°C for immunofluorescence.

Immunofluorescence

Immunofluorescence of cryosections was performed on glass slides. Immunofluorescence for
primary cells was performed on glass cover slips. Primary cells were fixed in 4% PFA for 10
minutes at room temperature. Antibody incubations were performed in blocking solution (0.1%
Triton X-100, 1% BSA, 10% donkey serum) for 2hr at room temperature or overnight at 4°C.
Slides or cover slips were mounted in ProLong Diamond Antifade Mountant (Thermo Scientific,
cat# P36965). Primary antibodies used in this study were anti-Sox9 (1:500, Proteintech, cat#
67439-1-Ig), anti-Runx2 (1:500, Cell Signaling, cat# D1L7F), anti-Ki67 (1:2000, Abcam, cat#
ab15580), anti-Vimentin (1:1000, Abcam, cat# ab8069), anti-BrdU (1:1000, Abcam, cat# ab6326),
anti-Sox10 (1:500, Cell Signaling, cat# D5V9L), anti-ALPL (1:1000, Invitrogen, cat# PA5-47419),
anti-Col1a2 (1:500, Proteintech, cat# 66761-1-Ig), anti-Sox6 (1:1000, Proteintech, cat# 14010-1-
AP), and anti-Eed (1:1000, Cell Signaling, cat# E4L6E). Secondary antibodies used in this study
were Donkey anti-Rabbit IgG (H+L) Alexa Fluor Plus 555 (1:1000, Thermo Scientific, cat#
A32794), Donkey anti-Rabbit IgG (H+L) Alexa Fluor 488 (1:1000, Thermo Scientific, cat#
A21206), and Donkey anti-Rabbit IgG (H+L) Alexa Fluor Plus 568 (1:1000, Thermo Scientific,
cat# A10042). DAPI solution (1:5000, Thermo Scientific, cat# 62248) was added to secondary
antibody solutions.

BrdU staining

Timed mating dams were subjected to intraperitoneal injection of 100mg/kg sterile BrdU in
PBS (Abcam, cat# ab142567). After 2hr, mice were euthanized and embryos were harvested in
cold PBS. Embryos were fixed in 4% PFA overnight at 4°C and cryosectioned. Sections were
incubated with 1M HCl for 2hr then 0.1M sodium borate buffer for 15min before being subjected to
standard immunofluorescence.

Microscopy and image analysis

Whole-mount skeletal and H&E stains were imaged using a Zeiss Stemi 305 Stereo Zoom
microscope running Zeiss Blue v2.0. Whole-mount embryo DAPI stains were imaged using an

upright Zeiss Axio Imager Z2 running AxioVision v4.0. Immunofluorescence images were
obtained using a Zeiss LSM800 confocal laser scanning microscope running Zen Blue v2.0.
Quantification of embryo measurements and immunofluorescence staining intensities was
performed in ImageJ using standard thresholding and measurement of signal integrated density.
Signal to nuclei normalization was performed by dividing signal intensity by DAPI signal per
image. Quantification was performed using n>6 images per embryo.

Primary craniofacial cell culture

Timed mating embryos were dissected on ice cold PBS and heads were removed. Brains
and eyes were removed from each head, and the remaining craniofacial structures were minced
with a sterile blade on a glass surface. Tissue was enzymatically dissociated in 3mg/ml
Collagenase Type 7 (Worthington, cat# CLS-6) in HBSS for 2-3hr at 37°C with frequent
trituration using an Eppendorf ThermoMixer. Dissociated cells were filtered through a 70uM
MACS smart strainer (Miltenyi Biotec, cat#130-110-916) and either plated on cover slips or cell
culture dishes. Primary craniofacial cells were grown in DMEM with 10% FBS. For RNA
extraction and immunofluorescence, cells were harvested one day following initial dissociation.

Single cell RNA-seq

Matched littermate timed embryos were dissected on ice cold PBS. Embryo heads were
removed and minced using sterile razor blades on a glass surface. Tissue was enzymatically
dissociated in 3mg/ml Collagenase Type 7 (Worthington, cat# CLS-6) in HBSS for 2-3hr at 37°C
with frequent trituration using an Eppendorf ThermoMixer. The quality of the dissociation was
frequently monitored by looking at the cell suspension on an automated cell counter. Cells were
spun down at 350rcf, resuspended in MACS BSA stock solution (Miltenyi Biotec, cat# 130-091-
376), and serially filtered through 70μm and 40μm MACS smart strainers (Miltenyi Biotec, cat#
130-110-916). Dissociation quality checks, cell viability, and counting was performed using an
Invitrogen Countess 3 automated cell counter. 10,000 cells were loaded per single-cell RNA
sequencing sample. Single-cell RNA sequencing libraries were generated using the Chromium
Single Cell 3' Library & Gel Bead Kit v3.1 on a 10x Genomics Chromium controller using the
manufacturer recommended default protocol and settings.

Single-cell RNA sequencing analysis

Library demultiplexing, read alignment to the mouse reference genome mm10, and unique
molecular identifier (UMI) quantification was performed in Cell Ranger v7.2.0. Cells with greater

than 200 unique genes were retained for analysis. Data were normalized and variance

stabilized by SCTransform in Seurat v5.0 (Hao et al., 2024). UMAP and cluster analysis were

performed using the Seurat function RunUMAP with parameters of mindist=0.7, res=0.4,

dims=1:30. Cluster markers were identified using Seurat function FindAllMarkers with

parameters min.pct = 0.25, thresh.use = 0.25. Differential gene expression analysis was

performed using DElegate v1.1.0 (Hafemeister and Halbritter, 2023) using the DEseq2 method,

with biological replicate animals (n=3 per genotype) as the replicate_column parameter.

Featureplots were generated using SCPubr v1.1.2 (Blanco-Carmona, 2022) and scale bars

represent log2 UMI's corrected by SCTransform.

Statistics

All experiments were performed with independent biological replicates and repeated, and

statistics were derived from biological replicates. Biological replicates are indicated in each

figure panel or figure legend. No statistical methods were used to predetermine sample sizes,

but sample sizes in this study are similar or larger to those reported in previous publications.

Data distribution was assumed to be normal, but this was not formally tested. Investigators were

blinded to conditions during data collection and analysis. Bioinformatic analyses were performed

blind to clinical features, outcomes, and molecular characteristics. The samples used in this

study were nonrandomized with no intervention, and all samples were interrogated equally.

Thus, controlling for covariates among samples was not relevant. No data points were excluded

from the analyses.

Competing interests statement

The authors declare no competing interests.

Author contributions

All authors made substantial contributions to the conception or design of the study; the

acquisition, analysis, or interpretation of data; or drafting or revising the manuscript. All authors

approved the manuscript. All authors agree to be personally accountable for individual

contributions and to ensure that questions related to the accuracy or integrity of any part of the

work are appropriately investigated and resolved and the resolution documented in the

literature. T-C.C. and D.R.R. conceived and designed the study. Experiments were performed

by T.C-C., C.T., J.A.C.S., and Y-G.J. Data analysis was performed by T.C-C., S.J.L., and D.R.R.

S.J.L. performed bioinformatic analyses. The study was supervised by H.N.V., J.O.B., and
D.R.R. The manuscript was prepared by T.C-C. and D.R.R. with input from all authors.
433

Funding

435　　This study was supported by funding from NIH grant T32 CA151022 to T.C-C., NIH grant
436　R35 DE031926 to J.O.B, and NIH grant R21 HD106238 and DOD grant NFRP NF200021 to
437　D.R.R. Sequencing was performed at the UCSF CAT, supported by UCSF PBBR, RRP IMIA,
438　and NIH 1S10OD028511-01 grants.
439

Data availability

441　　Single-cell RNA sequencing data that were reported in this manuscript have been deposited
442　in the NCBI Sequence Read Archive under PRJNA1077750
443　(https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1077750).

Code availability

446　　No custom software, tools, or packages were used. The open-source software, tools, and
447　packages used for data analysis in this study are referenced in the methods where applicable
448　and include Cell Ranger v7.2.0, Seurat v5.0, DElegate v1.1.0, and SCPubr v1.1.2.
449

Reference list

Bertol, J. W., Johnston, S., Ahmed, R., Xie, V. K., Hubka, K. M., Cruz, L., Nitschke, L., Stetsiv,
M., Goering, J. P., Nistor, P., et al. (2022). Twist1 interacts with beta/delta-Catenins during
neural tube development and regulates fate transition in cranial neural crest cells.
Development 149,.

Blanco-Carmona, E. (2022). Generating publication ready visualizations for Single Cell
transcriptomics using SCpubr. *bioRxiv* 2022.02.28.482303.

Bonnard, C., Strobl, A. C., Shboul, M., Lee, H., Merriman, B., Nelson, S. F., Ababneh, O. H., Uz,
E., Güran, T., Kayserili, H., et al. (2012). Mutations in IRX5 impair craniofacial development
and germ cell migration via SDF1. *Nat. Genet.* 44, 709–713.

Bowers, S. L. K., Meng, Q., Kuwabara, Y., Huo, J., Minerath, R., York, A. J., Sargent, M. A.,
Prasad, V., Saviola, A. J., Galindo, D. C., et al. (2023). Col1a2-Deleted Mice Have Defective
Type I Collagen and Secondary Reactive Cardiac Fibrosis with Altered Hypertrophic
Dynamics. *Cells* 12, 2174.

Bronner, M. E. and LeDouarin, N. M. (2012). Development and evolution of the neural crest: An
overview. *Dev. Biol.* 366, 2–9.

Cain, C. J., Gaborit, N., Lwin, W., Barruet, E., Ho, S., Bonnard, C., Hamamy, H., Shboul, M.,
Reversade, B., Kayserili, H., et al. (2016). Loss of Iroquois homeobox transcription factors 3
and 5 in osteoblasts disrupts cranial mineralization. *Bone Rep.* 5, 86–95.

Cardiff, R. D., Miller, C. H. and Munn, R. J. (2014). Manual Hematoxylin and Eosin Staining of
Mouse Tissue Sections. *Cold Spring Harb. Protoc.* 2014, pdb.prot073411.

Cheung, M., Chaboissier, M.-C., Mynett, A., Hirst, E., Schedl, A. and Briscoe, J. (2005). The
Transcriptional Control of Trunk Neural Crest Induction, Survival, and Delamination. *Dev.*
Cell 8, 179–192.

Cho, K. Y., Kelley, B. P., Monier, D., Lee, B., Szabo-Rogers, H. and Napierala, D. (2019). Trps1
Regulates Development of Craniofacial Skeleton and Is Required for the Initiation of Palatal
Shelves Fusion. *Front. Physiol.* 10, 513.

Cohen, A. S. and Gibson, W. T. (2016). EED-associated overgrowth in a second male patient.
J. Hum. Genet. 61, 831–834.

Cohen, A. S. A., Tuysuz, B., Shen, Y., Bhalla, S. K., Jones, S. J. M. and Gibson, W. T. (2015). A
novel mutation in EED associated with overgrowth. *J. Hum. Genet.* 60, 339–342.

Crispino, G., Pasquale, G. D., Scimemi, P., Rodriguez, L., Ramirez, F. G., Siati, R. D. D.,
Santarelli, R. M., Arslan, E., Bortolozzi, M., Chiorini, J. A., et al. (2011). BAAV Mediated
GJB2 Gene Transfer Restores Gap Junction Coupling in Cochlear Organotypic Cultures
from Deaf Cx26Sox10Cre Mice. *PLoS ONE* 6, e23279.

Deckelbaum, R. A., Holmes, G., Zhao, Z., Tong, C., Basilico, C. and Loomis, C. A. (2012).
Regulation of cranial morphogenesis and cell fate at the neural crest-mesoderm boundary by
engrailed 1. *Development* 139, 1346–1358.

Denisenko, O., Shnyreva, M., Suzuki, H. and Bomsztyk, K. (1998). Point Mutations in the WD40
Domain of Eed Block Its Interaction with Ezh2. *Mol. Cell. Biol.* 18, 5634–5642.

Dudakovic, A., Camilleri, E. T., Xu, F., Riester, S. M., McGee-Lawrence, M. E., Bradley, E. W.,
Paradise, C. R., Lewallen, E. A., Thaler, R., Deyle, D. R., et al. (2015). Epigenetic Control of
Skeletal Development by the Histone Methyltransferase Ezh2*. *J. Biol. Chem.* 290, 27604–
27617.

Fantauzzo, K. A. and Christiano, A. M. (2011). Trps1 activates a network of secreted Wnt
inhibitors and transcription factors crucial to vibrissa follicle morphogenesis. *Development*
139, 203–214.

Faust, C., Schumacher, A., Holdener, B. and Magnuson, T. (1995). The eed mutation disrupts anterior mesoderm production in mice. *Development* 121, 273–285.

Frisdal, A. and Trainor, P. A. (2014). Development and evolution of the pharyngeal apparatus. *Wiley Interdiscip. Rev.: Dev. Biol.* 3, 403–418.

Gao, C. W., Lin, W., Riddle, R. C., Kushwaha, P., Boukas, L., Björnsson, H. T., Hansen, K. D. and Fahrner, J. A. (2023). Novel mouse model of Weaver syndrome displays overgrowth and excess osteogenesis reversible with KDM6A/6B inhibition. *bioRxiv* 2023.06.23.546270.

Gerdes, J., Lemke, H., Baisch, H., Wacker, H. H., Schwab, U. and Stein, H. (1984). Cell cycle analysis of a cell proliferation-associated human nuclear antigen defined by the monoclonal antibody Ki-67. *J. Immunol. (Baltim., Md : 1950)* 133, 1710–5.

Gibson, W. T., Hood, R. L., Zhan, S. H., Bulman, D. E., Fejes, A. P., Moore, R., Mungall, A. J., Eydoux, P., Babul-Hirji, R., An, J., et al. (2012). Mutations in EZH2 Cause Weaver Syndrome. *Am. J. Hum. Genet.* 90, 110–118.

Goel, H., O'Donnell, S. and Edwards, M. (2024). EED related overgrowth: First report of multiple members in a single family. *Am. J. Méd. Genet. Part A* 194, 374–382.

Hafemeister, C. and Halbritter, F. (2023). Single-cell RNA-seq differential expression tests within a sample should use pseudo-bulk data of pseudo-replicates. *bioRxiv* 2023.03.28.534443.

Hao, Y., Stuart, T., Kowalski, M. H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., et al. (2024). Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat. Biotechnol.* 42, 293–304.

Honoré, S. M., Aybar, M. J. and Mayor, R. (2003). Sox10 is required for the early development of the prospective neural crest in Xenopus embryos. *Dev. Biol.* 260, 79–96.

Ianevski, A., Giri, A. K. and Aittokallio, T. (2022). Fully-automated and ultra-fast cell-type identification using specific marker combinations from single-cell transcriptomic data. *Nat Commun* 13, 1246.

Imagawa, E., Albuquerque, E. V. A., Isidor, B., Mitsuhashi, S., Mizuguchi, T., Miyatake, S., Takata, A., Miyake, N., Boguszewski, M. C. S., Boguszewski, C. L., et al. (2018). Novel SUZ12 mutations in Weaver-like syndrome. *Clin. Genet.* 94, 461–466.

Jang, Y. O., Cho, M.-Y., Yun, C.-O., Baik, S. K., Park, K.-S., Cha, S.-K., Chang, S. J., Kim, M. Y., Lim, Y. L. and Kwon, S. O. (2016). Effect of Function-Enhanced Mesenchymal Stem Cells Infected With Decorin-Expressing Adenovirus on Hepatic Fibrosis. *Stem Cells Transl. Med.* 5, 1247–1256.

Jeong, J., Mao, J., Tenzen, T., Kottmann, A. H. and McMahon, A. P. (2004). Hedgehog
signaling in the neural crest cells regulates the patterning and growth of facial primordia.
Genes Dev. 18, 937–951.

Jing, H., Liao, L., An, Y., Su, X., Liu, S., Shuai, Y., Zhang, X. and Jin, Y. (2016). Suppression of
EZH2 Prevents the Shift of Osteoporotic MSC Fate to Adipocyte and Enhances Bone
Formation During Osteoporosis. *Mol. Ther.* 24, 217–229.

Kawane, T., Qin, X., Jiang, Q., Miyazaki, T., Komori, H., Yoshida, C. A., Matsuura-Kawata, V. K.
dos S., Sakane, C., Matsuo, Y., Nagai, K., et al. (2018). Runx2 is required for the
proliferation of osteoblast progenitors and induces proliferation by regulating Fgfr2 and Fgfr3.
Sci. Rep. 8, 13551.

Kim, J., Lo, L., Dormand, E. and Anderson, D. J. (2003). SOX10 Maintains Multipotency and
Inhibits Neuronal Differentiation of Neural Crest Stem Cells. *Neuron* 38, 17–31.

Kim, S. Y., Paylor, S. W., Magnuson, T. and Schumacher, A. (2006). Juxtaposed Polycomb
complexes co-regulate vertebral identity. *Development* 133, 4957–4968.

Kim, H., Langohr, I. M., Faisal, M., McNulty, M., Thorn, C. and Kim, J. (2018). Ablation of Ezh2
in neural crest cells leads to aberrant enteric nervous system development in mice. *PLoS*
ONE 13, e0203391.

Komori, T. (2009). Osteoimmunology, Interactions of the Immune and skeletal systems II. *Adv.*
Exp. Med. Biol. 658, 43–49.

Lamandé, S. R., Mörgelin, M., Adams, N. E., Selan, C. and Allen, J. M. (2006). The C5 domain
of the collagen VI alpha3(VI) chain is critical for extracellular microfibril formation and is
present in the extracellular matrix of cultured cells. *J. Biol. Chem.* 281, 16607–14.

Lee, M. S., Lowe, G., Flanagan, S., Kuchler, K. and Glackin, C. A. (2000). Human dermo-1 has
attributes similar to twist in early bone development. *Bone* 27, 591–602.

Lindsay, E. A., Botta, A., Jurecic, V., Carattini-Rivera, S., Cheah, Y.-C., Rosenblatt, H. M.,
Bradley, A. and Baldini, A. (1999). Congenital heart disease in mice deficient for the
DiGeorge syndrome region. *Nature* 401, 379–383.

Lindsay, E. A., Vitelli, F., Su, H., Morishima, M., Huynh, T., Pramparo, T., Jurecic, V., Ogunrinu,
G., Sutherland, H. F., Scambler, P. J., et al. (2001). Tbx1 haploinsufficiency in the DiGeorge
syndrome region causes aortic arch defects in mice. *Nature* 410, 97–101.

Liu, Y., Strecker, S., Wang, L., Kronenberg, M. S., Wang, W., Rowe, D. W. and Maye, P.
(2013). Osterix-Cre Labeled Progenitor Cells Contribute to the Formation and Maintenance
of the Bone Marrow Stroma. *PLoS ONE* 8, e71318.

Liu, W., Zhang, L., Xuan, K., Hu, C., Liu, S., Liao, L., Li, B., Jin, F., Shi, S. and Jin, Y. (2018). Alpl prevents bone ageing sensitivity by specifically regulating senescence and differentiation in mesenchymal stem cells. *Bone Res.* 6, 27.

Luo, L., Ambrozkiewicz, M. C., Benseler, F., Chen, C., Dumontier, E., Falkner, S., Furlanis, E., Gomez, A. M., Hoshina, N., Huang, W.-H., et al. (2020). Optimizing Nervous System-Specific Gene Targeting with Cre Driver Lines: Prevalence of Germline Recombination and Influencing Factors. *Neuron* 106, 37-65.e5.

Maczkowiak, F., Matéos, S., Wang, E., Roche, D., Harland, R. and Monsoro-Burq, A. H. (2010). The Pax3 and Pax7 paralogs cooperate in neural and neural crest patterning using distinct molecular mechanisms, in Xenopus laevis embryos. *Dev. Biol.* 340, 381–396.

Margueron, R. and Reinberg, D. (2011). The Polycomb complex PRC2 and its mark in life. *Nature* 469, 343–349.

Matsuoka, T., Ahlberg, P. E., Kessaris, N., Iannarelli, P., Dennehy, U., Richardson, W. D., McMahon, A. P. and Koentges, G. (2004). Neural crest origins of the neck and shoulder. *Nature* 436, 347–55.

Mendez, M. G., Kojima, S. and Goldman, R. D. (2010). Vimentin induces changes in cell shape, motility, and adhesion during the epithelial to mesenchymal transition. *FASEB J.* 24, 1838–1851.

Metzis, V., Courtney, A. D., Kerr, M. C., Ferguson, C., Galeano, M. C. R., Parton, R. G., Wainwright, B. J. and Wicking, C. (2013). Patched1 is required in neural crest cells for the prevention of orofacial clefts. *Hum. Mol. Genet.* 22, 5026–5035.

Minoux, M. and Rijli, F. M. (2010). Molecular mechanisms of cranial neural crest cell migration and patterning in craniofacial development. *Development* 137, 2605–2621.

Mirzamohammadi, F., Papaioannou, G., Inloes, J. B., Rankin, E. B., Xie, H., Schipani, E., Orkin, S. H. and Kobayashi, T. (2016). Polycomb repressive complex 2 regulates skeletal growth by suppressing Wnt and TGF-β signalling. *Nat. Commun.* 7, 12047.

Montgomery, N. D., Yee, D., Montgomery, S. A. and Magnuson, T. (2007). Molecular and Functional Mapping of EED Motifs Required for PRC2-Dependent Histone Methylation. *J. Mol. Biol.* 374, 1145–1157.

Mori-Akiyama, Y., Akiyama, H., Rowitch, D. H. and Crombrugghe, B. de (2003). Sox9 is required for determination of the chondrogenic cell lineage in the cranial neural crest. *Proc. Natl. Acad. Sci.* 100, 9360–9365.

Nakamura, T., Gulick, J., Colbert, M. C. and Robbins, J. (2009). Protein tyrosine phosphatase
activity in the neural crest is essential for normal heart and skull development. *Proc. Natl.*
Acad. Sci. 106, 11270–11275.

Niethamer, T. K., Teng, T., Franco, M., Du, Y. X., Percival, C. J. and Bush, J. O. (2020).
Aberrant cell segregation in the craniofacial primordium and the emergence of facial
dysmorphology in craniofrontonasal syndrome. *PLoS Genet.* 16, e1008300.

O'Carroll, D., Erhardt, S., Pagani, M., Barton, S. C., Surani, M. A. and Jenuwein, T. (2001). The
polycomb-group gene Ezh2 is required for early mouse development. *Mol. Cell. Biol.* 21,
4330–6.

Pasini, D., Bracken, A. P., Jensen, M. R., Denchi, E. L. and Helin, K. (2004). Suz12 is essential
for mouse development and for EZH2 histone methyltransferase activity. *EMBO J.* 23, 4061–
4071.

Piunti, A. and Shilatifard, A. (2021). The roles of Polycomb repressive complexes in mammalian
development and cancer. *Nat. Rev. Mol. Cell Biol.* 22, 326–345.

Rigueur, D. and Lyons, K. M. (2013). Skeletal Development and Repair, Methods and Protocols.
Methods Mol. Biol. 1130, 113–121.

Rotllant, J., Liu, D., Yan, Y.-L., Postlethwait, J. H., Westerfield, M. and Du, S.-J. (2008). Sparc
(Osteonectin) functions in morphogenesis of the pharyngeal skeleton and inner ear. *Matrix*
Biol. 27, 561–572.

Sandell, L., Inman, K. and Trainor, P. (2018). DAPI Staining of Whole-Mount Mouse Embryos or
Fetal Organs. *Cold Spring Harb. Protoc.* 2018, pdb.prot094029.

Schumacher, A., Faust, C. and Magnuson, T. (1996). Positional cloning of a global regulator of
anterior–posterior patterning in mice. *Nature* 383, 250–253.

Schwarz, D., Varum, S., Zemke, M., Schöler, A., Baggiolini, A., Draganova, K., Koseki, H.,
Schübeler, D. and Sommer, L. (2014). Ezh2 is required for neural crest-derived cartilage and
bone formation. *Development* 141, 867–877.

Shukunami, C., Takimoto, A., Oro, M. and Hiraki, Y. (2006). Scleraxis positively regulates the
expression of tenomodulin, a differentiation marker of tenocytes. *Dev. Biol.* 298, 234–247.

Shull, L. C., Sen, R., Menzel, J., Goyama, S., Kurokawa, M. and Artinger, K. B. (2020). The
conserved and divergent roles of Prdm3 and Prdm16 in zebrafish and mouse craniofacial
development. *Dev. Biol.* 461, 132–144.

Simões-Costa, M. and Bronner, M. E. (2015). Establishing neural crest identity: a gene
regulatory recipe. *Development* 142, 242–257.

Smits, P., Li, P., Mandel, J., Zhang, Z., Deng, J. M., Behringer, R. R., Crombrugghe, B. de and
Lefebvre, V. (2001). The Transcription Factors L-Sox5 and Sox6 Are Essential for Cartilage
Formation. *Dev. Cell* 1, 277–290.

Spellicy, C. J., Peng, Y., Olewiler, L., Cathey, S. S., Rogers, R. C., Bartholomew, D., Johnson,
J., Alexov, E., Lee, J. A., Friez, M. J., et al. (2019). Three additional patients with EED-
associated overgrowth: potential mutation hotspots identified? *J. Hum. Genet.* 64, 561–572.

Tamamura, Y., Katsube, K., Mera, H., Itokazu, M. and Wakitani, S. (2017). Irx3 and Bmp2
regulate mouse mesenchymal cell chondrogenic differentiation in both a Sox9-dependent
and -independent manner. *J. Cell. Physiol.* 232, 3317–3336.

Tan, Z., Kong, M., Wen, S., Tsang, K. Y., Niu, B., Hartmann, C., Chan, D., Hui, C. and Cheah,
K. S. E. (2020). IRX3 and IRX5 Inhibit Adipogenic Differentiation of Hypertrophic
Chondrocytes and Promote Osteogenesis. *J. Bone Miner. Res.* 35, 2444–2457.

Tatton-Brown, K., Murray, A., Hanks, S., Douglas, J., Armstrong, R., Banka, S., Bird, L. M.,
Clericuzio, C. L., Cormier-Daire, V., Cushing, T., et al. (2013). Weaver syndrome and EZH2
mutations: Clarifying the clinical phenotype. *Am. J. Méd. Genet. Part A* 161, 2972–2980.

Tien, C.-L., Jones, A., Wang, H., Gerigk, M., Nozell, S. and Chang, C. (2015). Snail2/Slug
cooperates with Polycomb repressive complex 2 (PRC2) to regulate neural crest
development. *Development* 142, 722–731.

Ting, M.-C., Wu, N. L., Roybal, P. G., Sun, J., Liu, L., Yen, Y. and Maxson, R. E. (2009). EphA4
as an effector of Twist1 in the guidance of osteogenic precursor cells during calvarial bone
growth and in craniosynostosis. *Development* 136, 855–864.

Yan, Y.-L., Miller, C. T., Nissen, R. M., Singer, A., Liu, D., Kirn, A., Draper, B., Willoughby, J.,
Morcos, P. A., Amsterdam, A., et al. (2002). A zebrafish sox9 gene required for cartilage
morphogenesis. *Dev. (Camb., Engl.)* 129, 5065–79.

Yu, M., Riva, L., Xie, H., Schindler, Y., Moran, T. B., Cheng, Y., Yu, D., Hardison, R., Weiss, M.
J., Orkin, S. H., et al. (2009). Insights into GATA-1-Mediated Gene Activation versus
Repression via Genome-wide Chromatin Occupancy Analysis. *Mol. Cell* 36, 682–695.

Figure legends

Fig. 1. Sox10-Cre Eed^{Fl/Fl} embryos develop craniofacial malformations in utero. (A) Genotyping
frequencies of postnatal postnatal (top) and embryonic (bottom) *Sox10-Cre Eed^{Fl/Fl}* mice. (B)
Sagittal brightfield images of *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre Eed^{Fl/Fl}* embryos from E10.5 to
E17.5 showing craniofacial abnormalities appearing at E12.5 that were observed with 100%
penetrance. Scale bar, 1mm. (C) Coronal brightfield images of E16.5 *Sox10-Cre Eed^{Fl/WT}* or

Sox10-Cre Eed^{Fl/Fl} embryos showing craniofacial abnormalities. Scale bar, 1mm. (D) Whole-

mount fluorescence microscopy of DAPI-stained *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre Eed^{Fl/Fl}*

heads at E12.5, E14.5, or E16.5. Scale bar, 1mm. All images are representative of ≥3 biological

replicates.

Fig. 2. Eed is required for specification of craniofacial mesenchymal derivatives from the neural

crest. (A) Sagittal brightfield images of whole-mount skeletal stains of E18.5 *Sox10-Cre Eed^{Fl/WT}*

or *Sox10-Cre Eed^{Fl/Fl}* embryos. Alcian blue and alizarin red identify cartilage and bone,

respectively. Scale bar, 1mm. (B) Magnified sagittal brightfield images of whole-mount skeletal

stains of E18.5 *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre Eed^{Fl/Fl}* embryos. Bones structures are

annotated (I, interparietal; P, parietal; Mx, maxilla; Mb, mandible; Tr, tympanic ring; F, frontal; N,

nasal; T, temporal; S, supraoccipital). Scale bar, 1mm. (C) Micro-CT images of E19.5 *Sox10-Cre*

Eed^{Fl/WT} or *Sox10-Cre Eed^{Fl/Fl}* heads. Scale bar, 1um. (D) Brightfield coronal H&E images of

E15.5 *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre Eed^{Fl/Fl}* heads. Anatomic structures are annotated (Ct,

cortex; CP, cartilage primordium; CS, conjunctival sac; Gg, genioglossus muscle; HC, hyaloid

cavity; Hg, hyoglossus muscle; L, lens; LV, lateral ventricle; Mb, midbrain; MM, masseter

muscle; Np, nasopharynx; NC, nasal cavity; OB, olfactory bulb; OE, olfactory epithelium; OM,

ocular muscle; RNL, retina neural layer; Sm/St, sublingual and submandibular ducts; T, tongue;

TV, third ventricle; VF, vibrissa follicles). Scale bar, 1mm. (E) Coronal immunofluorescence

images of E15.5 *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre Eed^{Fl/Fl}* heads stained for ALPL (green,

mineralized bone) and Runx2 (red, early osteoblasts). DNA is marked by DAPI. Scale bars,

1mm and 100μm. (F) Coronal immunofluorescence images of E15.5 *Sox10-Cre Eed^{Fl/WT}* or

Sox10-Cre Eed^{Fl/Fl} heads stained for Ki67 (green, proliferating cells) and Vimentin (red,

mesenchymal cells). DNA is marked by DAPI. Scale bars, 1mm and 100μm. All images are

representative of ≥3 biological replicates.

Fig. 3. Eed regulates craniofacial differentiation and proliferation in vivo and in primary cell

cultures. (A) Coronal immunofluorescence images of E12.5 *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre*

Eed^{Fl/Fl} heads stained for Runx2 (red, early osteoblasts) and Sox9 (green). Intracranial tissues

demonstrate *ex vacuo* dilation and anterior herniation in *Sox10-Cre Eed^{Fl/Fl}* heads. DNA is

marked by DAPI. Scale bars, 500μm and 100μm. (B) Immunofluorescence images of primary

craniofacial cell cultures from E12.5 *Sox10-Cre Eed^{Fl/WT}* or *Sox10-Cre Eed^{Fl/Fl}* embryos stained

for Eed, neural crest transcription factors (Sox6, Sox9, Sox10) and markers of early osteoblasts

(Runx2) or cell proliferation (Ki67). Scale bar, 100μm. (C) Quantification of immunofluorescence

imaging intensity from primary craniofacial cell cultures. Lines represent means and error bars

represent standard error of the means. Student's t-tests, ***p≤0.0001. All images are

representative of ≥3 biological replicates.

Fig. 4. Eed specifies craniofacial mesenchymal stem cell, osteoblast, and proliferating

mesenchymal cell fate from the neural crest. (A) Integrated UMAP of 63,730 transcriptomes

from single-cell RNA sequencing of litter-matched E12.5 *Sox10-Cre Eed$^{Fl/WT}$* (n=3) or *Sox10-Cre*

Eed$^{Fl/Fl}$ (n=3) heads. (B) Integrated UMAP overlaying cell cluster distribution from single-cell

RNA sequencing. (C) Quantification of mesenchymal cell cluster distribution from single-cell

RNA sequencing. Lines represent means and error bars represent standard error of the means.

Student's t-tests, **p≤0.01, ***p≤0.0001. (D) Gene expression feature plots of mesenchymal

lineages from single-cell RNA sequencing. (E) Volcano plot showing differentially expressed

genes between C0 (osteoblasts) and C4 (mesenchymal stem cells). (F) Gene expression

feature plots for differentially expressed genes between C0 and C4. (G) Volcano plot showing

differentially expressed genes between C0 and C5 (mesenchymal stem cells). (H) Gene

expression feature plots for differentially expressed genes between C0 and C5.