**A rapid and efficient in vivo inoculation method for introducing tree stem canker**

**pathogens onto leaves, suitable for large-scale assessment of resistance in poplar breeding**

**progeny**

**Zheng Li [a, 1], Bingyu Zhang [b, 1], Yuchen Fu [a], Yutian Suo [a], Yinan Zhang [a], Jinxia Feng [a], Long Pan [a], Wanna**

**Shen [a], Huixiang Liu [c], Xiaohua Su [b] and Jiaping Zhao [a, *]**

a. State Key Laboratory of Tree Genetics and Breeding, Institute of Ecological Conservation and Restoration, Chinese Academy of Forestry,
Beijing, China
b. State Key Laboratory of Tree Genetics and Breeding, Research Institute of Forestry, Chinese Academy of Forestry, Beijing, China
c. Shandong Research Center for Forestry Harmful Biological Control Engineering and Technology, College of Plant Protection, Shandong
Agricultural University, Taian, China

1. The authors contributed equally to this manuscript.
* Correspondence: zhaojiaping@caf.ac.cn; ORCID: https://orcid.org/0000-0002-2054-6989; Tel.: 86-10-62889457.

**Abstract**

Hybrid breeding is the most direct and efficient method of controlling and managing tree

diseases. However, "in vitro stem inoculation" can not be used for rapid, efficient, and low-cost

screening of resistant clones in the early stage of stem canker infection. Therefore, we inoculated

stem canker pathogens on poplar leaves to evaluate the resistance of hybrid clones in the poplar-

*Valsa sordida* pathosystem. "In vivo leaf inoculation" showed that: 1) *V. sordida* induces

extended necrotic lesions and conidia on leaves. 2) The upper leaves exhibited higher resistance

than the middle leaves. 3) The shading conditions induced more severe symptoms on leaves than

the lighting conditions. 4) The susceptibility of poplar leaves to juvenile mycelium was higher

than old mycelium's. 5) The resistance of 48 poplar clones against *V. sordida* was distributed

normally. 6) The efficacy of "in vivo leaf inoculation" was consistent with "in vitro stem

inoculation". In summary, "in vivo leaf inoculation" has the advantages of a wide application

range, simple operation, short experimental process, and host safety. This method is of great

significance for poplar breeding, the study of pathology and molecular biology, pathogenic

differentiation, and sporulation of fungal canker pathogens.

**Keywords:** poplar hybrid breeding, tree diseases, stem canker, in vivo leaf inoculation, *Valsa*

*sordida*

**1. Introduction**

Poplar species are always affected by different kinds of pests, especially two stem canker

pathogens: *Botryosphaeria dothidea* (*B. dothidea*) and *Valsa sordida* (*V. sordida*) canker. These

are the two major forest diseases in China (Zhang & Luo, 2003), which can ruin all the seedlings

in poplar plantations or severely damage mature poplar forests. Hybridization breeding is the

most direct and efficient pathway to manage and control forestry diseases.

For breeding resistant clones to plant disease, accurately and quickly getting the phenotypic

characteristics, including disease incidence, onset time, area of the necrotic, and disease index,

has become a critical pathological problem affecting breeding efficiency. There are three

methods, field investigation assays, callus inoculation (Zhang *et al.*, 1989) and in vitro stem

segment inoculation method, to obtain the disease prototype for the breeding of poplar canker;

however, the In vitro stem segment inoculation is the most commonly used method (Chen *et al.*,

1985; Shi *et al.*, 2014; Zhang *et al.*, 2017). It used eight to ten stem segments 30-40 cm long as

inoculation materials. Each stem segment was inoculated with 5 points, observing the incidence

rate and recorded every five days within 60 days of inoculation, and the disease index was

calculated. Thus, the cost of breeding time and land, as well as the cost of labor and expenditure

that accompanied the former, will be huge.

During the growing season, abundant leaves were continuously produced on trees. Then,

should we carry out an "in vivo leave inoculation" to detect the host resistance to stem canker

disease? The first and most crucial step of the disease onset is the invasion and colonization of

pathogens in host tissues. Therefore, whether the stem canker pathogens reside in poplar leaves

has become the prerequisite for this method. *B. dothidea* not only induces canker disease on

stems and branches of plants but also causes latent infection as endophytic pathogens or lives in

leaves, fruits, and other places (Reviewed by Marsberg *et al.*, 2017). The excised leaves of *Malus*

*spectabilis* were used to detect the pathogenicity of the mutant isolates of *B. dothidea* (Zheng *et*

*al.*, 2017). The mycelium and conidial suspension solution of *V. ceratosperma* (stem canker

pathogen of apple trees) induced disease symptoms on the excised leaves, shoots, and fruits of

apple trees (Wei *et al.*, 2010; Xie, 2019). In addition, *Septoria musiva* Peck, an important fungal

pathogen of hybrid poplars in the north-central and northeastern regions of North America,

causes a leaf spot and stem canker disease of hybrid poplars (review by Feau *et al.*, 2010;

Dunnell & LeBoldus, 2017). Based on the above research, we speculated that leaf disease may

be induced by inoculation of stem canker pathogens. In addition, Zhang *et al.* (2018) proposed

inoculating rice blight pathogens with isolated leaves to detect pathogenicity. Inoculation of

isolated leaves and shoots of *V. ceratosperma* can be used as an accurate and rapid method to

evaluate the pathogenicity of stem canker pathogen *V. ceratosperma* in apple trees. However, no

specific method for disease resistance detection was proposed in this paper, and it has not been

applied to cultivating disease-resistant varieties in tree trunks (Wei *et al.,* 2010). As described

above, acquiring pathotypes and selecting resistance clones are still technical problems in the

hybrid breeding of poplar stem canker diseases.

This study proposed a novel method for the resistance detection of poplar clones based on

the in vivo leaf inoculation of stem canker pathogens. Then, using the poplar/*V. sordida* disease

system, the effects of the leaf development stage (leaf position on the branch), light condition of
leaves, and cultured time of the fungal pathogen were assayed. The differentiations of plant
resistance to *V. sordida* were assayed in a small, randomly selected population of hybrid poplar
clones and then were verified using stem segment inoculation methods. Finally, the diversity of
pathogenicity to poplar in some *B. dothidea* isolates was also detected in this study. It is expected
to develop a rapid, effective, and low-cost method for the inoculation of stem canker pathogens
and a new strategy of hybrid breeding for the canker diseases of poplars and other tree species.

**2. Materials and methods**

**2.1. Fungal and plant materials**

The fungi used in this study are poplar canker pathogen *V. sordida* isolate CZC and 13
poplar blister canker (*B. dothidea*) isolates (Table 1). All fungal isolates were deposited in the
plant physiology laboratory at the Institute of Ecology Conservation and Restoration, Chinese
Forestry Academy (CAF), Beijing. The pathogenic isolates were grown on PDA medium (20%
potato, 2% dextrose, 2% agar; pH 6.0) at 28°C in dark conditions for 4 or 7 days after inoculation,
and then cut into square mycelium plugs; the side length was about 1.0-1.2 cm.
One or 2-year-old poplar saplings of *Populus euramericana* cv. 'Bofeng 3' (Huang *et al.*,
2014), 6-year-old *P. alba var. pyramidalis* saplings, and 2-year-old saplings of the hybrid clones
(*P. deltoids×P. deltoids*, 48 genotypes) were used in this study. All poplar saplings were
cultivated in the experimental field of CAF, with no pest infestation on leaves and stem, and well
irrigated during the experiments. The day before inoculation, 3-4 mature and neighbored leaves
were washed using sterilized water and then sterilized with 75% alcohol (vol/vol) (Figure 1-A1).
After drying, all sterilized leaves were marked for inoculation.

**2.2. Inoculation methods**

Two or four square mycelium plugs were first placed on the selected leaves upper surface
according to the size of leaves, about 1-3 cm apart from the central veins, and avoiding the
secondary veins (Figure 1-A2). The mycelium faced the upper surface of the poplar leaves. One
leaf of each poplar genotype near the pathogen-inoculated leaves was inoculated by 2% WA
(water agar) square plugs as the control. After the pathogen inoculation, the leaves were carefully
covered and sealed with 5 cm wide household scotch tape to fix and prevent the water loss of the
mycelium or WA plugs (Figure 1-A3). Then, the leaves were pierced into 5 points from the
lower surface with a needle, at the center of the inoculated mycelium or WA square plugs or just
at the corners of the square (about 1-2mm away from the corners) (Figure 1-A4). After
inoculation, the sealed condition of the scotch tape and the development of disease symptoms
that appeared on the inoculated leaves were observed and recorded daily. Remove the household

scotch tape after three days (Figure 1-A5). All the inoculated leaves were picked from poplar branches or twigs five days after inoculation (DAI), placed in dark plastic bags, and stored at 4℃. After the experiment, all picked leaves were immediately taken into the laboratory, and scotch tapes, mycelium, and WA plugs were carefully removed from the leaves. Then, all leaves were photoed or scanned as JPEG images, and the images were at least 1500 pixels. The necrotic spots around the prick sites on the leaves were identified, and their area was calculated automatically in the software ImageJ (https://imagej.nih.gov/ij/) or was manual adjusted if necessary. The results were kept in Microsoft EXCEL.

In this study, the lesion shape, color, areas, and whether the hypha-like structure formed around the prick sites inoculated by pathogen or WA were set as the determining criterion for leaf disease. After pathogen inoculation, the leaf necrotic lesions were always extended, the necrotic area was more than 2.0 $mm^2$; the lesion color was always grey or dark; the hypha-like structure was consistently observed, or sometimes, the pycnidia-like structure was observed on the surface of the necrotic spot. On the contrary, after WA inoculation, the leaf necrotic lesions were always brown; the lesion area was average of 1.5 $mm^2$, not extended, and no hypha-like structure was observed on the leaves. For simplicity, the necrotic areas on the inoculated leaves, $A = 2.0$ $mm^2$, were set as the quantitative criteria to determine the occurrence of leaf disease (Table 2). If $A > 2.0$ $mm^2$, the prick sites are diseased, or vice versa. The disease index of each poplar clone was calculated according to the criteria and the ranking level of disease development (Table 2).

In this study, the disease incidence and disease index was calculated using the following formula:

Formula 1: Disease incidence rate (IR) = Number of the diseased prick sites/Total Number of the efficient prick sites × 100%.

Formula 2: Disease index (DI) = ∑ (Number of prick sites by level × Values of severity at all levels)/(Value of the highest severity level × Total Number of the efficient prick sites) × 100.

**2.3. Comparison of the resistance to canker pathogen among the different stages of leaves**

The square mycelium plugs of *V. sordida* isolate CZC, which were cultivated in darkness for seven days, were inoculated on the 5-7[th] newly matured leaves (from the top to the bottom of the poplar saplings, the same applies from now on) and 18-20[th] mature leaves on the same saplings of *P. euramericana* cv. 'Bofeng 3'. The inoculation method was described in 2.2; two square mycelium plugs inoculated every leaf, and three biological replicates (saplings) were used. In total, 120 pathogen-inoculated prick sites were created, 60 for the upper and 60 for the lower leaves. One additional poplar sapling was inoculated by WA plugs as the control and a total of 40 prick sites were created.

**2.4. Comparison of the resistance between lighting and shading leaves**

The lighting and shading leaves on the different branches of 6-year-old *P. alba var. pyramidalis* were inoculated by *V. sordida* isolate CZC (cultivated in the darkness at 28°C for seven days). Besides the light condition, the branches' location and base diameter were identical. Two square pathogen mycelium plugs inoculated the 5-7$^{th}$ newly matured leaves. Two branches (shading and lighting) of three poplar replicates were inoculated in this experiment. The inoculation method is described in 2.2. Therefore, 120 pathogen-inoculated prick sites were created: 60 for the lighting and 60 for the shading groups. In addition, two shading and lighting leaves were also inoculated by WA plugs as the control.

**2.5. Comparison of the pathogenicity of canker pathogen with different cultured time**

To detect the effect of the culture time of the pathogen on its pathogenicity, *V. sordida* isolates CZC cultivated in darkness for 4 and 7 days were inoculated on the leaves of *P. euramericana* cv. 'Bofeng 3'. In this experiment, six poplar saplings were divided into two groups: one inoculated by a 4-days-cultured pathogen (D4) and another inoculated by a 7-days-cultured pathogen (D7). Two square pathogen mycelium plugs inoculated the 5-7$^{th}$ newly matured leaves. The inoculation method is described in 2.2. Finally, 60 pathogen-inoculated prick sites were created for group 4 days, 60 for group 7 days, and 30 prick sites as the control.

**2.6. In vivo assessment of resistance to *Valsa* canker in hybrid poplar clones using leaf inoculation method**

The *Valsa* pathogen isolate CZC was cultured on a PDA medium in the darkness at 28°C for seven days. The 5-7$^{th}$ newly matured leaves, located on one or two hybrid clones, were inoculated by two or four square pathogen mycelium plugs. Three leaves were inoculated with mycelium plugs, and one leaf was inoculated with WA plugs. After all 48 hybrid clones were inoculated, needles immediately pricked the leaves. Finally, pathogen plugs inoculated a total of 30 or 60 prick sites, and WA plugs inoculated 30 prick sites as the control. The detailed inoculation method was the same as the methods described in 2.2. The disease symptoms were observed and photoed five days after inoculation (5 DAI). According to the symptoms and necrotic areas around the prick sites determine the disease incidence and index.

**2.7. In vitro validation of poplar resistance to *Valsa* canker pathogen using stem segments**

To validate the effectiveness of the leaf inoculation method in assessing popular resistance to stem canker pathogen, the response of 15 *P. deltoids×P. deltoids* hybrid clones (5 resistance/high resistance, five non-resistance/non-susceptible, and five susceptible/high

susceptible to canker pathogen determined by in vivo leaf inoculation in Section 2.7) were
detected using an excised stem segment inoculation method. This study used ten dormant poplar
stem segments (2-3 cm in diameter, 25-30 cm in length) derived from the base region (50-60 cm)
of the 1-year-old branches of the selected poplar clones. Before inoculation, the 25-30 cm long
poplar segments were all washed with sterilized water and then sterilized with 75% alcohol, and
the upper end of the segments using plant callus ointment (a commercial product promoting the
regeneration of callus around the wound) after dry (Figure 1B-1). The segments were soaked in
water for three days at room temperature and then inoculated with *V. sordida* isolate CZC
(cultured in 2% PDA at 28°C for seven days) (Figure 1B-2). Zhang (2017) described the
inoculation method, each segment was inoculated with five mycelium plugs (0.5 cm in diameter)
every 3-4 cm per inoculation site from the beginning site (10 cm height from the base end of the
segment) to the upper back. Every inoculation site was pricked using a needle as a regular
triangle (the side length is about 1-2mm). After inoculation, all inoculation sites were covered by
household PE film to preserve moisture, and then all inoculated stem segments were cultured in
water. Ten segments were inoculated in this study, and two were mock inoculated.
The disease development around the inoculation sites was observed and recorded according
to whether the necrotic produced every five days; the necrotic sites were marked, and the total
number of disease sites of every poplar clone was also recorded until 40 days after inoculation
(Yang *et al.*, 1989; Shi *et al.*, 2014). According to the time the disease symptoms appeared, the
disease severity of each poplar clone was ranked (Table S1) (Yang *et al.*, 1989; Zhang *et al.*,
1989), and the disease index of each poplar clone according to the formula (2). At the last
observation, every lesion's longitudinal extension length (a, mm) and the horizontal length (b,
mm) were measured. Because the lesion shapes were mainly subelliptic, the necrotic area of
disease spots was calculated according to the elliptic area formula ($A = a b \pi/4$). The severity
of the disease was ranked according to the criterion in Table S2, and the disease index was
calculated according to the formula (2).

**2.8. Determination of pathogenicity of stem canker pathogens using in vivo leaf inoculation method**

The pathogenicity of a total of 12 *B. dothidea* isolates, which came from different regions
and different, was determined in this study using the leaf inoculation method. All fungal isolates
were inoculated on the 5-7th newly matured leaves of 13 different twigs of one 2-year-old poplar
'Bofeng 3' hybrid clone. Each isolate inoculated three leaves, and two or four square mycelium
plugs were used. Therefore, 30 or 60 pathogen-inoculated prick sites were created after needle
piercing. Moreover, WA plugs inoculated two other twigs or just needle piercing as the two
control treatments, Ctrl-1 and Ctrl-2, respectively. The leaf inoculation method was described in
2.2. Five days after inoculation, the symptoms were observed, photoed, and then automatically

identified using ImageJ software, which was adjusted in the handle if necessary. The disease

incidence, area of necrotic spots, and disease index were determined using results derived from

ImageJ software.

**2.9. Data statistics**

In this study, the t-test, ANOVA (analysis of variance), and LSD (least significant

difference) methods were used to test the significance of the necrotic area; the chi-square method

was used to test the importance of the disease incidence rate, and the Shapiro-Wilk method was

used to test the normal distribution of the numbers of the poplar clones in different resistance

levels.

**3. Results**

**3.1. Stem canker pathogen *Valsa sordida* induced necrotic symptoms on poplar leaves**.

In this study, the control (WA inoculation) and *V. sordida* inoculation induced necrotic

spots around the prick sites on the poplar leaves at 5 DAI (Figure 2A-B). All the control necrotic

spots are unextended; their average areas are $1.5\pm 0.5$ mm$^2$ (Figure 2A, 2C-F), while the

pathogen inoculation can produce extended necrotic spots on poplar leaves (Figure 2B, 2G-K).

No hyphen structure was observed in the stereomicroscope on the surfaces of the control poplars.

In contrast, hyphen structures were observed in the pathogen-inoculated poplars (Figure 2G-I),

even some pycnidium structures (Figure 2J-K), and some extruded milky white spores were

observed on the leaves of some hybrid poplar clones B246 (Figure 2K), B73, B14, B18, B76,

B181 and B1. In some severed diseased poplar clones, such as clone B246, almost all five

necrotic spots overlapped. Based on the average area size of the necrotic spots (NA) in the

control poplar leaves, NA > 2.0mm$^2$ was set as the quantity criterion of the disease onset on the

pathogen-inoculated poplar leaves (Table 2).

**3.2. The influence of leaf ontogeny on disease severity**

This study shows 5-7th newly matured leaves (upper leaves) and 18-20th mature leaves

(lower leaves) of *P. euramericana* cv. 'Bofeng 3' was used to detect the influence of leaf

ontogeny on disease severity. As shown in Figure 2, five days after inoculation (DAI), the

disease severity of the upper leaves is significantly lower than the lower leaves (chi-square test, n

= 60, p < 0.05), the percentage of disease incidence (P) of the upper and lower leaves are 28.33%

and 70.00%, respectively. After discarding the undiseased inoculation points, results showed that

the disease area of the necrotic spots of the upper leaves is also significantly lower than that of

lower leaves ($2.49\pm 0.13$ (n = 17) vs $2.94\pm 0.13$ mm$^2$ (n = 42); t-test, p < 0.05). Moreover, the
disease index (DI) of the upper and lower leaves are 5.67 and 14.67, respectively (Figure 3A).

**3.3. The influence of light condition of leaves on disease severity**

In this study, the 5-7$^{th}$ leaves grew on three lighting branches (were exposed to the sun all
day) and on three shading branches (were not exposed to direct sunlight) of the same two-year-
old saplings were used to compare the influence of light conditions of leaves on disease severity.
Results showed no differences in the disease incidence rate between the lighting and shading
leaves (IR$_{lighting}$ 58.33% vs. IR$_{shading}$ 55.00%, respectively; chi-square test, n = 60, p > 0.05).
However, results suggested that shading promoted the disease severity in poplar leaves: the
average necrotic area of the lighting and shading leaves are $3.87\pm 0.29$ (n = 35) and $7.40\pm 1.20$
mm$^2$ (n = 33), respectively (t-test, p < 0.05). In addition, the disease index of lighting leaves
(DI$_{lighting}$) is 18.00, while the disease index of shading leaves (DI$_{shading}$) is 26.00 (Figure 3B).

**3.4. The culture time of fungal inoculum altered the disease severity on leaves.**

Results showed that the culture time of mycelium inoculums also affected the disease
development in poplar leaves. Their disease incidence rate induced by 4-day-cultured *V. sordida*
mycelium (IR$_{4day}$) is 83.33%, significantly IR$_{7day}$ (65.00%) (chi-square test, n = 60, p < 0.05),
suggesting that the juvenile pathogen inoculums had with higher pathogenicity than the older.
After discarding the undiseased inoculation points, results showed that the necrotic area induced
by 4-day-cultured mycelium plugs (NA$_{4day}$ = $8.51\pm 0.57$ mm$^2$, n = 50) is significantly higher than
that caused by 7-day-cultured mycelium plugs (NA$_{7day}$ = $3.37\pm 0.20$mm$^2$, n = 39) (t-test, p <
0.05). Moreover, the disease indexes, DI$_{4day}$ and DI$_{7day}$, are 50.67 and 13.33, respectively (Figure
3C).

**3.5. In vivo inoculation of stem canker pathogen *Valsa sordida* on leaves revealed the
resistance diversity in poplar hybrid clones.**

In this study, every hybrid clone produced 60 prick sites, and all 48 clones produced 2,880
sites; after discarding the inefficient inoculation sites (the mycelium plugs moved away or dried
in 1-3 DAI), a total of 2,715 efficient inoculation data were obtained occupying 94.27% of all
schemed inoculation dataset. Taking the criteria of the necrotic area (NA) > 2.0 mm$^2$ as the
determining criteria for the occurrence of disease on poplar leaves, a total of 2,145 inoculation
sites exhibited definite necrotic features, indicating that the authentic leaf disease incidence in 40
poplar hybrid clones 79.00%.
Results revealed high resistance diversity to the *Valsa* canker pathogen among 48 poplar
hybrid clones using a novel leaf inoculation method. As shown in Table S3, the disease

incidence rate in 48 poplar clones varied from 31.67% (poplar clone B133) to 100.00 (B1 and B104) (Figure 4A), the size of necrotic area varied from $2.72\pm 0.13\text{mm}^2$ (B168) to $15.28\pm 1.26\text{mm}^2$ (B14) (Figure 4B), and the disease index ranged from 6.67 (B168) to 83.67 (B246 and B135) (Figure 4C).

This study used the Shapiro-Wilk model to test the distribution of the poplar numbers in different resistance levels. Though the p-value in the Shapiro-Wilk test did not support a normal distribution, the absolute kurtosis of the normality test is 1.05 (<10.0). The absolute skewness is 0.44 (3.0), and the resistance among these 48 poplar clones is generally accepted as a normal distribution (Kline *et al.*, 2023) and supported in the P-P plot analysis (Figure 5).

According to the disease index, all 48 poplar hybrid clones were divided into seven disease resistance levels, from the high resistance group to the very high susceptible group (Figure 4C). For example, hybrid B133 was identified as a high resistance (HR) clone, hybrid B168, and other seven clones were identified as resistance (R) clones, and hybrid B135, B246, and B104 were susceptible (S) to *Valsa* canker pathogen CZC, using these leaf disease severity assessment methods (Table S3).

**3.6. In vitro stem segment inoculation verified the effectiveness of the leaf inoculation method**.

As shown in Table 3 and Table S4, the disease incidence rate (IR) of 15 poplar hybrid clones varied from 18.00% (B133) to 98.00% (B246), the $\text{DI}_{stem\text{-}time}$ varied from 15.71 (B133) to 92.86 (B246). The $\text{DI}_{stem\text{-}area}$ of 15 poplar hybrid clones varied from 5.00 (B133) to 78.00 (B246) (Figure 6A-C). The values and ranking of $\text{DI}_{leaf\text{-}area}$, $\text{IR}_{stem}$, $\text{DI}_{stem\text{-}time}$, and $\text{DI}_{stem\text{-}area}$ were compared to assess the consistency of the leaf and stem inoculation methods. As shown in Table 3 and Figure 6D, besides clone B145 in the R/HR group and clone B76 in the S/LS group, the resistance rankings of the other eight poplar clones that were assessed by leaf and stem inoculation method are the same, in particular, the ranking of the highest resistance clone B133 and lowest resistance clone B246 were identically in these two methods. Therefore, the results suggested an alternative pathway, the leaf inoculation method, for acquiring pathotypes and assessing resistance clones in tree pathological research and poplar breeding.

**3.7. In vivo leaf inoculation revealed the diversity of pathogenicity in stem canker pathogen *B. dothidea.***

Results showed that the pathogenicity of 12 *B. dothidea* isolates on poplar hybrid 'Bofeng 3' varied hugely. *B. dothidea* isolate SD47 induced the highest necrotic area in poplar leaves ($6.57\pm0.41\text{mm}^2$), while the size of necrotic area induced by isolate SD60 and CZ1218 are $4.53\pm 0.68$ and $3.20\pm 0.19 \text{ mm}^2$, respectively. Though there was no significant difference between

isolate SD47 and SD60, and also between isolate SD60 and CZ1218, the necrotic area of isolate

SD47 was significantly higher than isolate CZ1218 (ANOVA, $p<0.05$) (Figure 7). In addition,

the sizes of the necrotic area of 9 isolates (CZ1070c, CZ1070b, CZ1010, CZ1055, etc.) were

smaller than 2.0 mm$^2$, the quantitative criteria to determine disease occurrence in leaf in this

study indicated these nine isolates had with low pathogenicity on poplar hybrid 'Bofeng 3' (Table

4). These results suggested that the leaf inoculation method can be used to identify pathogen

pathogenicity.

**4. Discussion**

Selection based on phenotypic is the crucial step of hybrid breeding. Due to their large

crown, slow growth, and long maturity period characteristics in tree species, phenotypic

acquisition of tree stem canker diseases always requires more time, space, and human resources

than herb diseases and tree leaf diseases. Because the necrotic spots are dispersed around the

surface of the branches at different heights, the phenotypic acquisition of stem canker diseases is

more tedious, complicated, and complex than leaf diseases.

Studies illustrated that stem canker pathogens can infect host leaves or live in leaves as

endophytes (Marsberg *et al.*, 2017; Feau *et al.*, 2010; LeBoldus *et al.*, 2010; Dunnell &

LeBoldus, 2017; Zheng *et al.*, 2017; Wei *et al.*, 2010), provided a feasible pathway to detect the

resistance of stem canker disease through leaf inoculation method. However, the consistency of

the leaf inoculation method with the stem inoculation method remains controversial. Wei *et al.*

(2010) and this study in *Valsa* canker disease illustrated the consistency between these two

methods, and the leaf inoculation methods can be used to select resistance clones or the assays of

pathogenicity differentiation in the pathogen population. However, there was no correlation in

the disease severity between leaf spot and stem canker in the poplar-*Septoria* pathosystem

(Dunnell & LeBoldus, 2017).

The disease severity after leaf inoculation is directly related to the resistance of leaves. The

correlation between the leaf age (and the leaf position on branches) and resistance has been

observed for more than 100 years (reviewed by Gusberti *et al.*, 2013). This pattern of increasing

resistance with tissue age is called ontogenic or age-related resistance and can be specified as

leaf-stage resistance in leaf tissues (Asalf *et al.,* 2014; Dunnell & LeBoldus, 2017; Xu *et al.*,

2018). In physiological research, since the active and stable physiological performance, the

newly matured leaves at the top of the branches or saplings (always 5-7$^{th}$ leaves) were taken as

the ideal materials used in the study of photosynthesis, chlorophyll fluorescence, stomatal

movement, water status, physiology and biochemistry, and also gene expression regulation of

tree species (Zhao *et al.*, 2017; Xing *et al.*, 2019; Li *et al.*, 2019; Shen *et al.*, 2023; Li *et al.*,

2024). In powdery mildew of strawberry and *Malus-Venturia* interactions, the leaf resistance

increased rapidly from the first unfurled and expanding leaf to lower leaves, and the resistance

reached the top at the 5-6[th] matured expanding leaves (Asalf *et al.*, 2014; Gessler & Stumm,
1984; MacHardy *et al.*, 1996), in *Malus-V. ceratosperma* system, the inoculated leaves were
described as "fully extended at the top of the stems" (Wei *et al.*, 2010). This study illustrated that
the resistance of 5-7th leaves (always newly matured) is significantly higher than the 18-20[th]
leaves (32-35 leaves grow on the 1-year-old saplings). Moreover, this study illustrated that the
resistance results from the 5-7th leaves were consistent with those from the excised stem
segments (Table 3, Figure 6D). However, besides the differences in disease type (*Valsa* canker
vs. *Septoria* canker) and inoculation method (wounding mycelium vs. non-wounding spore
inoculation), the difference in the leaf position (5-6[th] newly matured leaves vs. the five middle
leaves in the branches of 3 or 4-month-old hybrid poplar clones) might be the main reason for
the inconsistency between the leaf and stem resistance inpoplar-*Septoria* pathosystem (Dunnell
& LeBoldus, 2017). Based on the above analysis, the top 5-7 leaves were suggested as the
optimal inoculation materials for poplar leaf inoculation in assessing and selecting the resistance
clones in poplar breeding.
Solar UVB radiation is a positive modulator of plant defense (Ballaré, 2014). Evidence from
several plant-pathogen systems indicates that shading increases infection by a range of pathogens
(Roberts & Paul, 2006). In addition, high population density (which is usually accompanied by
reduced light exposure) has been shown to markedly increase the severity of plant disease in
natural and agricultural systems (Augspurger & Kelly, 1984; Bell *et al.*, 2006; Burdon &
Chilvers, 1982; Jurke & Fernando, 2008). In this study, we use the leaf inoculation method in the
resistance detection of perennial poplars or the pathogenicity detection of different pathogen
isolates using the leaves on the various branches of the same poplar tree. The light conditions of
different leaves are remarkably different, especially for the backlit and light-oriented leaves.
Then, can the difference be detected among these leaves? The current study provided a positive
answer to the above question: The susceptibility to the pathogen CZC significantly increased in
the shading leaves in hybrid poplar "Bofeng 3" (Figure 3B). Shading is a regular method to
promote the disease onset in a controlled environment. For example, in the hybrid-Septoria
system, LeBoldus and his colleagues covered and sealed the black plastic garbage bags (with two
pieces of moist paper towel) for 48 h to promote the germination and invasion of spore
inoculants and the appearance of disease symptoms (LeBoldus *et al.*, 2010; Dunnell & LeBoldus,
2017). In this study, the leaf inoculation experiment showed that the smaller size of disease
necrotic areas was detected on the lighting leaves (Figure 3B). Moreover, another study showed
that high density (or shading) induced more severe canker disease symptoms on the truck of 6-
year-old *P. alba var. pyramidalis* (Shen *et al.*, 2024). Therefore, in the assays of the poplar
resistance or the pathogen's pathogenicity using the leaf inoculation method, the poplar leaves
with an identical light condition must be carefully considered.

In the Petri plate, the morphology, colors, and growth rate of the colony and mycelium of the pathogens varied during the culture period. Since plants can exhibit age-related resistance, such as leaf-stage resistance, pathogens will likely have some culture of age-related pathogenicity. For example, the pathogenicity and infection rate of the conidia of *Dactylaria higginsii*, which were harvested from 15 days of culturing on PDA, was higher than that of the conidia harvested from other culture ages (Kadir *et al.*, 2011). Conidiospores of the three entomogenous fungi (*Metarhizium anisopliae*, *Paecilomyces fumoso-roseus,* and *Verticillium lecanii*) harvested from young (2–3d old) Petri plate cultures germinated more rapidly than those taken from older cultures. However, conidia of *Beauveria bassiana* all germinated at the same rates, irrespective of culture and age (Richard *et al.*, 1994). This study illustrated that the pathogenicity of the young culture (4 days old in PDA medium) of *V. sordida* isolate CZC to poplars was higher than that of the old culture (7 days old). However, some fungal isolates cannot grow over a 9 cm diameter Petri dish in 7 days; for example, some *V. sordida* isolates, to harvest more mycelium inoculants, the culture conditions of canker pathogens suggested as follows: cultured on PDA medium, at 28°C, in dark condition, for seven days.

Some researchers considered that the most aggressive or virulent strains of a pathogen that causes the disease should be used in the selection and assays of plant germplasm and breeding lines (Balendres *et al.*, 2019). This selection strategy may be beneficial to the funding of the authentic highest resistance clones in the breeding population; however, it also may be a disaster for the whole breeding program for the disease or even the death of all offspring, or it has no breeding significance for the other poor production phenotypes. As shown in Figure 5, the numbers of poplar clones in 7 different pathogenicity levels were near normal distribution, suggesting that the pathogen *V. sordida* CZC is the optimal test strain for the current poplar hybrid population.

If another isolate of *V. sordida* was adopted, whose pathogenicity is higher than that of *V. sordida* CZC, then the number of these 48 poplar clones might be right skewed distributed (almost clones susceptible to the pathogen), and if a weak pathogenicity isolate was adopted, the number of poplar clones might exhibit a left-skewed distribution. Therefore, the optimal pathogen strains must preselected in a new breeding program before final resistance assays. The side length of the square, 1.0-1.2 cm, is an experience value based on our preliminary experiment; if resistance assays of the 48 poplar clones were against a high pathogenicity strain, larger mycelium plugs might be an optimal option. Finally, besides three isolates of of *B. dothidea*, the other nine fungal isolates exhibited no or very low pathogenicity to hybrid poplar "Bofeng 3". The reason for this phenomenon might come from the high disease-resistance characteristics of the hybrid poplar. Then, the pre-selection of poplar clones, which have an appropriate resistance level, is needed in the assays of large-scale pathogen populations.

This study illustrated that poplar stems canker pathogen *V. sordida* can induce lesion symptoms on leaf tissue using a leaf inoculation method. The efficiency of this method in hybrid breeding was validated using a small poplar hybrid population in this study. The process is simple and easy to operate; using this method, a team of three people can complete the inoculation of more than 200 clones in a day. For the rapid development of disease symptoms, the pathotypes can acquired in 3-5 days after inoculation; then, breeders can acquire the pathotypes of a large-scale population (1,000 genotypes or much more) in one month. For short periods of disease stresses (removed from the stems at 3-5 days after inoculation), the pathogen did not or less affected the growth and physiology of the inoculated poplars and then did not or less change the other phenotypes (such as the heights, stem diameter and mass. Therefore, the breeders can quickly achieve multi-objective breeding of canker disease and fast growth, high yield, etc.

In short, compared to the time-consuming and high-cost in vitro stem segment inoculation method, the in vivo leaf inoculation method is fast, efficient, low-cost, and high-throughput, suitable for the large-scale resistance assays of hybrid poplar clones, can be used in the seedling selection of resistance clones in the breeding of stem canker diseases (and some leaf diseases) of poplar or other tree species. Combined with the new breeding technology, such as genomic selection, the breeding resources can excavate deeply for the new resistance-related genes, gene loci, or gene modules (Du *et al.* 2023). In addition, this leaf inoculation method can be used in the large-scale detection of pathogenicity, and the differentiation assays of poplar and tree stem canker or leaf pathogens.

**Author Contributions:** Conceptualization, J.Z. and B.Z.; methodology, J.Z. and B.Z.; software, Z.L. and Y.S.; validation, Z.L., Y.S., W.S. and Y.F.; formal analysis, Z.L., Y.S., W.S. and Y.F.; investigation, Z.L., Y.F. and Y.S.; resources, B.Z. and J.Z.; data curation, Z.L, Y.S. and J.F..; writing—original draft preparation, Z.L. and J.Z.; writing—review and editing, J.Z., Y.Z., Z.L., H.L. and L.P.; visualization, Z.L. and W.S.; supervision, X.S.; project administration, J.Z. and B.Z.; funding acquisition, J.Z. All authors have read and agreed to the published version of the manuscript.

**Declaration of Competing Interest**

**Data availability**

Data will be made available on request.

**Funding**

This research was jointly funded by the Central Public-interest Scientific Institution Basal Research Fund of State Key Laboratory of Tree Genetics and Breeding (grant number CAFYBB2020ZY001-2) and the National Natural Science Foundation of China (grant number 32171776) to Jiaping Zhao.

# References

Zhang X, Luo Y. Major forest diseases and insect pests in China [M]. Beijing: China Forestry Publishing House, 2003.

Zhang L, Li C, Huang M. A study on the resistance to *Botryosphaeria dothided* in callus tissues
of poplar clones [J]. Journal of Nanjing Forestry University, 1989, 32 (04): 9–15.

Chen H, Liu Z, Guo Y, *et al.* The pathogenicity assays of the main poplar cultivar clones in
China [J]. Journal of Liaoning Forestry Technology, 1985 (5): 2–3, 7.

Shi Y. Identification of canker resistance to selection poplar clones [D]. Northwest Agriculture
& Forestry University, 2014.

Zhang L, Zhang H, Ou D, *et al.* Investigation on the resistance to canker of some new leuce
hybrids inoculated with *Botryosphaeria dothidea* [J]. Journal of Northwest Forestry University,
2017, 32 (6): 210–213.

Marsberg A, Kemler M, Jami F, *et al.Botryosphaeria dothidea*: a latent pathogen of global
importance to woody plant health [J]. Molecular Plant Pathology, 2017, 18 (4): 477–488.

Zheng W, Jia X, Wang Y, *et al.* Construction and evaluation of the mutant library of
*Botryosphaeria dothidea* [J]. Journal of Shandong Agricultural Universit (Natural Science
Edition), 2017, 48 (01): 1–6.

Wei J, Huang L, Gao Z, *et al.* Laboratory evaluation methods of apple *Valsa* canker disease
caused by *Valsa keratoses* a sensu Kobayashi [J]. Acta Phytopathologica Sinica, 2010, 40 (1):
14–20.

Xie S. Comparative genomics of five pathogenic species of genus *Valsa* [D]. Northwest
Agriculture and Forestry University, 2019.

Feau N, Mottet M J, Périnet P, *et al.* Recent advances related to poplar leaf spot and canker
caused by *Septoria musiva*: Minireview/Minisynthèse [J]. Canadian Journal of Plant Pathology,
2010, 32 (2): 122–134.

Dunnell K L, LeBoldus J M. The correlation between *Septoria* leaf spot and stem canker
resistance in hybrid poplar [J]. Plant Disease, 2017, 101 (3): 464–469.

Zhang M, Sun X, Cui L, *et al.* The plant infection test: Spray and wound-mediated inoculation
with the plant pathogen *Magnaporthe Grisea* [J]. JoVE (Journal of Visualized Experiments),
2018 (138): e57675.

Huang Q, Su X, Wang S, *et al.* Integrative evaluation of *Populus × euramericana cl*．'Bofeng 1'
and *Populus × euramericana cl*．'Bofeng 2' [J]. Scientia Silvae Sinicae, 2014, 50 (05): 75–81.

Yang J, Li W, Fu Y, *et al.* A study on the disease resistances of *Eastern Cottonwood* Clones [J]. Shanxi Forest Science and Technology, 1989, (02): 7–14.

Kline R B. Principles and practice of structural equation modeling [M]. Guilford Publications, 2023.

LeBoldus J M, Blenis P V, Thomas B R. A method to induce stem cankers by inoculating nonwounded populus clones with *Septoria musiva* spore suspensions [J]. Plant Disease, 2010, 94 (10): 1238–1242.

Gusberti M, Gessler C, Broggini G A L. RNA-Seq analysis reveals candidate genes for ontogenic resistance in *Malus-Venturia pathosystem* [J]. PLoS One, 2013, 8 (11): e78457.

Asalf B, Gadoury D M, Tronsmo A M, *et al.* Ontogenic resistance of leaves and fruit, and how leaf folding influences the distribution of powdery mildew on strawberry plants colonized by *Podosphaera aphani*s [J]. Phytopathology, 2014, 104 (9): 954–963.

Xu Y, Lv L, Xu Y, *et al.* Leaf stage-associated resistance is correlated with phytohormones in a pathosystem-dependent manner [J]. Journal of Integrative Plant Biology, 2018, 60 (8): 703–722.

Zhao J, Yang F, Feng J, *et al.* Genome-wide constitutively expressed gene analysis and new reference gene selection based on transcriptome data: a case study from poplar/canker disease interaction [J]. Frontiers in Plant Science, 2017, 8: 1876.

Xing J, Li P, Zhang Y, *et al.* Fungal pathogens of canker disease trigger canopy dieback in poplar saplings by inducing functional failure of the phloem and cambium and carbon starvation in the xylem [J]. Physiological and Molecular Plant Pathology, 2020, 112: 101523.

Li P, Liu W, Zhang Y, *et al.* Fungal canker pathogens trigger carbon starvation by inhibiting carbon metabolism in poplar stems [J]. Scientific Reports, 2019, 9 (1): 10111.

Shen W, Fu Y, Wang L, *et al.* Transcriptomic analysis revealed that low-density aphid infestation temporarily changes photosynthesis and disease resistance but persistently promotes insect resistance in poplar leaves [J]. Forests, 2023, 14 (9): 1866.

Li M, Li J, Yao Y, *et al.* Pathological and genome assembly of transcriptome sequencing data implied multiple relationships between *Bean Common Mosaic Virus* (BCMV), *Batocera lineolata* and aphids in Populus *alba var. Pyramidalis* [J]. Physiological and Molecular Plant Pathology, 2024, 130: 102218.

Gessler C, Stumm D. Infection and stroma formation by *Venturia inaequalis* on apple leaves
with different degrees of susceptibility to scab [J]. Journal of Phytopathology, 1984, 110 (2):
119–126.

Gessler C, Stumm D. Infection and stroma formation by *Venturia inaequalis* on apple leaves
with different degrees of susceptibility to scab [J]. Journal of Phytopathology, 1984, 110 (2):
119–126.

MacHardy W E. Ontogenic resistance to scab in Mauls [J]. Apple Scab Biology, Epidemiology,
and Management: APS Press, St. Paul, Minnesota, 1996: 104–116.

Ballaré C L. Light regulation of plant defense [J]. Annual Review of Plant Biology, 2014, 65:
335–363.

Roberts M R, Paul N D. Seduced by the dark side: integrating molecular and ecological
perspectives on the influence of light on plant defense against pests and pathogens [J]. New
Phytologist, 2006, 170 (4): 677–699.

Augspurger C K, Kelly C K. Pathogen mortality of tropical tree seedlings: experimental studies
of the effects of dispersal distance, seedling density, and light conditions [J]. Oecologia, 1984, 61:
211–217.

Bell T, Freckleton R P, Lewis O T. Plant pathogens drive density-dependent seedling mortality
in a tropical tree [J]. Ecology Letters, 2006, 9 (5): 569–574.

Burdon J J, Chilvers G A. Host density as a factor in plant disease ecology [J]. Annual review of
phytopathology, 1982, 20 (1): 143–166.

Jurke C J, Fernando W G D. Effects of seeding rate and plant density on sclerotinia stem rot
incidence in canola [J]. Archives of phytopathology and plant protection, 2008, 41 (2): 142–155.

Shen, W.; Pan, L.; Fu, Y.; Suo, Y.; Zhang, Y.; Liu, H.; Su, X.; Zhao, J. Comparative study on the
effectiveness of three inoculation methods for *Valsa* sordida in populus alba var. pyramidalis.
preprints 2024, 2024030186. https://doi.org/10.20944/preprints202403.0186.v1

Kadir J, Hailmi M S, Charudattan R, *et al.* Effects of culture age on conidia production, viability
and pathogenicity of *Dactylaria higginsii* [J]. Journal of Agrobiotechnology, 2011, 2: 17–23.

Richard A. Hall, Dorothy D. Peterkin, Bibi Al, *et al.* Influence of culture age on rate of
conidiospore germination in four deuteromycetous entomogenous fungi [J]. Mycological
Research, 1994, 98 (7): 763–768.

Balendres M A, De Torres R, Dela Cueva F. Culture storage age and fungal re-isolation from

host–tissue influence *Colletotrichum* spp. virulence to pepper fruits [J]. Journal of

Phytopathology, 2019, 167 (9): 510–515.

Du J, Ge X, Wei H, *et al.* PsPRE1 is a basic helix-loop-helix transcription factor that confers

enhanced root growth and tolerance to salt stress in poplar [J]. Forestry Research, 2023 (FR–

2023–0016).

## Figures:

**Figure 1:** The schematic diagram of the leaf inoculation method provided in this study (A). This study provided a schematic diagram of the stem segments' inoculation method. (B).

**Figure 2:** Using a novel leaf inoculation method, the necrotic symptoms caused by pathogen *Valsa sordida* isolate CZC on the poplar leaves. In the WA medium inoculated poplar leaf, only unextended spots ($1.5 \pm 0.5$ mm$^2$) were observed (A); *V. sordida* inoculated the poplar leaf isolate CZC, some extended necrotic spots (more than 2.0 mm$^2$) were observed (B); the poplar leaves and the necrotic spots induced by *V. sordida* isolate CZC (A-B); the necrotic spots on the PDA inoculated leaves (C-F); the necrotic spots on the PDA inoculated leaves (G-I); the formation of fungal hyphen and pycnidium structure on the leaves of high susceptible hybrid clone B246 at five days after inoculation by *V. sordida* isolate CZC (J-K).

**Figure 3:** The effects of leaf position (A), leaf light condition (B), and culture age of the mycelium inoculants (C) on the leaf disease severity inoculated by poplar stem canker *Valsa sordida* isolate CZC, using the leaf inoculation method.

**Figure 4:** The disease incidence rate (A), necrotic area (B), and disease index (C) of the leaves of 48 poplar hybrid clones were inoculated by poplar stem canker *Valsa sordida* isolate CZC using the leaf inoculation method.

**Figure 5:** The leaf inoculation method revealed that the resistance of the 48 hybrid poplar clones is generally normally distributed. The standard distribution model (A) and the P-P plot of 48 hybrid poplar clones (B).

**Figure 6:** The disease incidence rate (A), disease index of onset time (B), disease index of onset area (C), and two-dimensional representation of DI$_{stem-time}$ and DI$_{leaf-area}$ (D) of the stems of 15 poplar hybrid clones were inoculated by poplar stem canker *Valsa sordida* isolate CZC using the stem inoculation method.

**Figure 7:** The necrotic area in the inoculated leaves of "Bofeng 3" poplar against 12 *Botryophaeria dothidea* isolates.

**Tables:**

Table 1: The hosts and locations of fungal isolates used in this study.

Table 2: Severity grading of necrotic symptoms induced by canker pathogen on leaves.

Table 3: The disease index of 15 poplar hybrid clones was detected using in vivo leaf inoculation and in vitro stem segments inoculation (the ranking values are based on the results of the leaf inoculation method).

Table 4: The necrotic areas in the inoculated leaves of "Bofeng 3" poplar against 12 *Botryophaeria dothidea* isolates.

Table S1: Severity grading criteria for disease onset time in poplar stems.

Table S2: Severity grading criteria for disease onset area in poplar stems.

Table S3: The leaf necrotic areas of the 48 hybrid poplar clones inoculated by *Valsa sordida* isolate CZC.

Table S4: The number of diseased poplars stem segments at the different time points (days after inoculation) during the experiments.