Predicting absolute protein folding stability using generative models

Matteo Cagiada[1*], **Sergey Ovchinnikov**[2], **Kresten Lindorff-Larsen**[1*]

*For correspondence:
matteo.cagiada@bio.ku.dk (MC);
lindorff@bio.ku.dk (KLL)

[1]Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen, DK-2200 Copenhagen, Denmark; [2]Massachusetts Institute of Technology, Cambridge, USA

Abstract

While there has been substantial progress in our ability to predict changes in protein stability due to amino acid substitutions, progress has been slow in methods to predict the absolute stability of a protein. Here we show how a generative model for protein sequence can be leveraged to predict absolute protein stability. We benchmark our predictions across a broad set of proteins and find a mean error of 1.5 kcal/mol and a correlation coefficient of 0.7 for the absolute stability across a range of small–medium sized proteins up to ca. 150 amino acid residues. We analyse current limitations and future directions including how such model may be useful for predicting conformational free energies. Our approach is simple to use and freely available via an online implementation.

Introduction

The stability of a protein is a critical property that affects its biological functions, as well as our ability engineer proteins for new purposes. This stability is mainly the result of a delicate balance between the atomic interactions in the folded state, the higher conformational entropy in the unfolded state, as well as a complex relationship between the protein structure and water properties (*Anfinsen, 1973*; *Fersht, 2017*). By protein stability, we here concern ourselves with the (thermodynamic) free energy of unfolding, ΔG, under a given set of conditions such as temperature and pH.

Experimentally, ΔG can be measured using a range of biophysical and spectroscopic techniques, though generally the value under native-like conditions requires extrapolations from for example high concentrations of denaturant or elevated temperatures (*Lindorff-Larsen and Teilum, 2021*). These measurements generally require purification of each variant and is relatively labour intensive, even though automated methods have been used (*Nisthal et al., 2019*). While the thermodynamic stability has been determined for a number of proteins and variants thereof using these approaches, the resulting data are relatively inhomogeneous in terms of methods used and experimental conditions (*Nikam et al., 2021*; *Xavier et al., 2021*). As an alternative, protein stability may in certain cases also be indirectly quantified using kinetic experiments (*Park and Marqusee, 2005*), and this has recently been combined with large-scale DNA synthesis and display technologies to quantify protein stability at a very large scale (*Tsuboyama et al., 2023*). While this type of experiments has provided a wealth of information, it is for technical reasons currently limited to small proteins with less than about 80 amino acid residues and which fold in a two-state manner.

As a supplement to experimental measurements, computational predictions can be used to assess the stability of a protein, there has been substantial interest in and development of computational predictions related to protein stability. With few exceptions, these methods have mostly focused on predictions of *changes* in thermodynamic stability ($\Delta\Delta G$), typically between a protein

and a variant that differs from it by one or very few amino acid substitutions. A number of methods to predict $\Delta\Delta G$ have been developed (*Pancotti et al., 2022*), and their accuracy in part originates from a number of important compensating effects. These include that while ΔG may be sensitive to the conditions (pH, temperature, etc), $\Delta\Delta G$ will be less sensitive. Similarly, while both ΔG and $\Delta\Delta G$ contain contributions from many interactions across the protein, many of these will cancel out when comparing the stability of a wild-type protein and a variant with a single amino acid substitution. Thus, in these senses $\Delta\Delta G$ is a more robust quantity than ΔG.

One additional complication in analysing and predicting protein stability is that many of the underlying thermodynamic parameters (ΔH, ΔS and ΔC_p) are strongly correlated with the number of residues (length) of the protein (*Robertson and Murphy, 1997*). These effects in turn suggest that it will be easier to predict the thermal stability (melting temperature) of a protein than the stability at a fixed temperature (*Pucci and Rooman, 2014*), and indeed several methods have been developed to predict melting temperatures. Previous work used these properties to predict, for example, stability from chain length (*Ghosh and Dill, 2009*) with more recent methods taking additional physical and statistical information into account (*Ruiz-Blanco et al., 2013*; *Pucci and Rooman, 2014*; *Yang et al., 2022*).

While progress in predicting stability (ΔG) has been relatively slow, developments in the field of machine learning have resulted in an increase in the number of prediction methods to estimate changes in protein stability ($\Delta\Delta G$) (*Pucci et al., 2022*). While these developments are encouraging, a recent example illustrates the difficulty of training models that learn both ΔG and $\Delta\Delta G$ (*Chu and Siegel, 2023*). Specifically, the authors trained a model directly on large-scale measurements of ΔG for small proteins, and showed that the model could be used to estimate both the global stability of similar folds and the effect of substitutions. However, the work also showed limited transferability, suggesting that training broadly transferable models for ΔG on a set of small proteins may be difficult.

Generative models for proteins are deep learning architectures that can be used in protein design to generate novel protein sequences or structures using representations learned from self-supervised training with large sequence or structure databases. One such model, the so-called 'Inverse Folding' approach of the Evolutionary Scale Model family (ESM-IF; *Hsu et al.* (*2022*)), uses the backbone structure of a target protein fold as input to sample novel sequences that fit the structural constraints imposed by the backbone. The new sequences are generated by designing one amino acid position at a time, sampling the amino acid type from position-specific likelihoods, given the input fold. The amino acid likelihoods generated by ESM-IF are learned during training on millions of sequence-structure pairs.

Recently, it has been shown that the ESM-IF likelihoods can also be used to assess $\Delta\Delta G$ by comparing the likelihoods of the wild-type and variant sequence, using the wild-type structure as input (*Hsu et al., 2022*; *Chen and Gong, 2022*; *Reeves and Kalyaanamoorthy, 2023*; *Notin et al., 2024*). Here, we explored whether models such as ESM-IF can also be used to solve the more difficult problem of predicting protein stability across different folds and sequences (ΔG). Our manuscript is structured as follows. First, we show how information from ESM-IF can be used to predict absolute stability for proteins with equilibrium characterised by a two-state folding mechanism. Second, we explore how well the model can assess the relative stability between two different configurations of a protein. Third, we show limitations for predicting stability for larger and more complex systems, where cooperativity between structural elements may also play a role. Finally, we briefly discuss issues that we believe need to be addressed by experiments and future models in order to advance the field.

Figure 1. Predicting absolute protein stability using a generative model. We show a comparison between predicted and experimental stability values for 265 proteins from two different data sets (represented by different colours). The precicted values are based on Eq. 1 and have been rescaled to units of kcal/mol (ΔG_{pred}). The Spearman's correlation coefficient (ρ_s) and the RMSE (in kcal/mol) are shown. The standard error for the correlation coefficient was obtained by bootstrapping.

Results and Discussion

Predicting global folding stability using generative models

It has been shown that the ESM-IF model can be used to predict changes in protein stability (*Hsu et al., 2022*; *Chen and Gong, 2022*; *Reeves and Kalyaanamoorthy, 2023*; *Notin et al., 2024*), suggesting that it might encode some level of protein biophysics that relates protein structure to thermodynamics. We therefore hypothesized that the ESM-IF model could also be used to predict the absolute stability of a given fold (ΔG) given its sequence. Because protein stability may vary substantially across conditions, we created a benchmark data set of experimental ΔG measurements by merging a curated data set of ΔG measurements of two-state folding proteins obtained under 'standard' experimental conditions (25 °C, pH 7.0) (*Maxwell et al., 2005*) with ΔG measurements from a large-scale proteolysis experiment on small single-domain proteins (*Tsuboyama et al., 2023*) (performed at ca. 25 °C, pH 7.4). In total, the benchmark data include 265 proteins ranging in size from 30 to 150 residues (Fig. S1A) and ΔG ranging from 1–12 kcal/mol (Fig. S1B). The dataset includes (i) 26 proteins whose kinetics and thermodynamics have been studied biophysically and that range in size from 61–151 residues (*Maxwell et al., 2005*) and (ii) 239 naturally occurring wild-type proteins (40–72 residues long) from the recently described 'megascale' experiment (*Tsuboyama et al., 2023*).

We used this data set to assess whether conditional likelihoods for amino acids calculated from ESM-IF for a given structure could be used to predict its global stability. Since it has been shown that protein stability correlates with protein size (*Ghosh and Dill, 2009*), we used as measure the sum of likelihoods for the target sequence $\sum_i^N \mathscr{L}_i^{\text{WT}}$ (Eq. 1). Here, the sum runs over the N residues in the protein, and $\mathscr{L}_i^{\text{WT}}$ is the likelihood from ESM-IF for having the reference (wild-type) amino acid at position i (see Methods). We calculated this measure for each of the 265 proteins and obtain a Pearson's correlation of 0.69 with the experimental ΔG values, demonstrating how ESM-IF can be used to predict absolute protein stabilities. To convert the values from Eq. 1 into ΔG_{pred} in physical units, we fitted a scaling constant and offset using the experimental data (see Methods). The resulting predictions have a root-mean-square-error between ΔG_{exp} and ΔG_{pred} of 1.4 kcal/mol (Fig. 1).

We then compared the accuracy achieved by Eq. 1 and ΔG_{pred} to other simpler measures such as the number of contacts in the fold or the sequence length (*Bastolla and Demetrius, 2005*; *Ghosh*

and Dill, 2009). We find that both of these measures are substantially less correlated with the experimental ΔG values than ΔG_{pred} calculated from ESM-IF (Fig. S2). We also assessed whether the summed model entropy could be used to predict protein stability and find that there was no correlation with experimental data S3).

Having assessed that the summed likelihoods correlate with global stability, we investigated whether absolute stability was correlated with the summed likelihoods of specific amino acid types or groups of residue types. We found that the summed likelihoods of two aliphatic residue types, valine and isoleucine, correlated the strongest with the global stability (Fig. S4A), and the sum of these two contributions correlate strongly with global stability (Fig. S4B). When we aggregated the amino acid likelihoods over all the hydrophobic amino acids, the results correlate with the global stability as much as the total sum of the likelihoods (Fig. S4C), in line with the assumption that protein stability is largely determined by the hydrophobic effect; the correlation using only buried residues was slightly lower (Fig. S4D). Together, these results show that the summed model likelihoods correlate with global protein stability, and that this correlation to a large extent is driven by the hydrophobic amino acids.

Having shown that our approach can be used to predict absolute folding stability at a level greater than simpler baseline models, we then asked whether other machine learning models than ESM-IF might achieve similar accuracy using the same approach. To investigate this we used Eq. 1 together with likelihoods obtained from (i) a sequence-based language model (ESM-1b, (*Rives et al., 2021*)) to assess the amount of information that could be extracted directly from the sequence or (ii) likelihoods from a different generative model (Chroma; *Ingraham et al.* (*2023*)) as a comparison between different design architectures. We also tested energy evaluations using the empirical force field FoldX (*Schymkowitz et al., 2005*), which can be used to predict changes in protein stability. Among the methods tested, we find that the ESM-IF likelihoods achieved the best correlation with experimental stability measurements (Fig. S5). We note that the sequence-based model, when used together with Eq. 1, achieves a near-zero correlation with the stability measures, highlighting that the stability of a given fold is a difficult task to learn from evolutionary conservation information alone. We also note that while FoldX has been shown to be useful for predicting $\Delta\Delta G$ from both experimental (*Schymkowitz et al., 2005*) and predicted (*Akdel et al., 2022*) structures, it was not developed to predict ΔG.

Prediction of stability differences between different protein conformations

Having shown that we can use the ESM-IF likelihoods to predict absolute protein stability from a given structure, we then asked whether we could use the same model to predict the difference in stability between two distinct conformations. By thermodynamic consistency, this corresponds to predicting the free energy difference between two distinct conformational states, each represented by a single structure.

To assess this ability, we selected two systems where the energetic difference between two states in solution has been measured experimentally using NMR relaxation dispersion techniques. The first system is the L99A variant of T4 lysozyme (T4L L99A, Fig. S6A), which is known to exchange between two conformations, termed G (ground) and E (excited), and with the E state being less stable than the G state by ca. 2 kcal/mol (*Mulder et al., 2001*; *Bouvignies et al., 2011*). Using structures of the G and E states (*Bouvignies et al., 2011*) as input to ESM-IF and Eq. 1 we predict that the E state is 1.2 kcal/mol less stable than G, in qualitative agreement with experiments (Fig. 2A).

As the second example, we selected Cyclophilin A (CYPA, Fig. S6B) where NMR experiments have shown a free energy difference between two states of 1.7 kcal/mol (*Eisenmesser et al., 2005*), and where models of the two states have been obtained by crystallography (*Fraser et al., 2009*). Again we used ESM-IF to predict the free energy difference between the two configurations. As for T4L L99A, we also predict a free energy difference whose sign is correct, but with a magnitude (0.3 kcal/mol) that is substantially smaller than found in experiments (1.7 kcal/mol; Fig. 2A). We here note that the ESM-IF model only uses the backbone structure as input, and that it is still not fully

Figure 2. Assessing the ability to predict the free energy difference between two distinct conformations of a protein. Comparison between predicted and experimental conformational free energies for (A) T4 lysozyme L99A (T4L) and (B) cyclophilin A (CYPA). In both panels the predicted ΔG from the two state are on the y-axis and the experimental $\Delta \Delta G$ is reported on the x-axis. The arrows indicate the $\Delta \Delta G$ between the two states.

known whether the free energy difference in CYPA measured by NMR is fully reflected in the small structural difference (C_α RMSD of 0.34 Å), between the two structures (*Eisenmesser et al., 2005*; *Fraser et al., 2009*; *Papaleo et al., 2014*; *Wapeesittipan et al., 2019*; *Thompson et al., 2019*).

While these results demonstrate that our approach cannot yet accurately predict free energy differences between states, they suggest that as prediction accuracy of the stability predictions increase, it might be possible to resolve conformational free energies of the magnitudes found in these two test systems.

Prediction of the absolute stability of more complex systems

Having shown that ESM-IF provides relatively accurate predictions of absolute stability for small–medium sized proteins that fold in a two-state manner, we asked whether the model could also be used to estimate stability for more complex systems. Specifically, we were interested in assessing whether our approach could also be used to predict the stability of larger, multi-domain proteins—potentially with cooperativity between the different domains—and for proteins where global unfolding does not follow an equilibrium two-state mechanism.

To assess this, we first study two different protein systems, each involving repeated domains: the G5[1]-E-G5[2] repeat of the Surface protein G (SasG, Fig. S6D; *Gruszka et al.* (*2015*)) and the Notch ankyrin domain (Nank, Fig. S6C; *Mello and Barrick* (*2004*)). In both cases, experimental measures of absolute stability are available for different combinations of the domains (*Gruszka et al., 2015*; *Mello and Barrick, 2004*). While our approach to predict stability captures the size-dependency across small proteins, we wondered whether it might overestimate stability for larger proteins with (semi-)independent domains. We therefore evaluated the stability for the different domain constructs using either (i) the full structure as input to ESM-IF or (ii) using the sum of stability predictions for the individual domains.

For the SasG system we find that the absolute stabilities predicted for the different combinations of domains (G5[1]-E, E-G5[2], and G5[1]-E-G5[2]) are substantially overestimated when we use the full structure as input to ESM-IF (Fig. 3A). When we instead calculate the stability as the sum of those of the individual domains we obtain results that are closer to experiments, though still overestimated relative to experiments (Fig. 3A). The imperfect agreement may in part originate from the complexity of the system; the E-domain located in between the G5[1] and G5[2] domains is disordered on its own and only folds when located between the folded G-domains (*Gruszka et al., 2012*). Thus, the calculations for the G5[1]-E and E-G5[2] constructs (using models with folded E domains) overestimate the stability. Also, the stability of the G5[1]-E-G5[2] construct contains a substantial contribution from the interfaces between the different domains (*Gruszka et al., 2015*) that is not captured by our

Figure 3. Predicting stability of multi-domain repeat proteins. For both (A) SasG and (B) Nank we calculated the stability of different domain constructs using either structural models with all domains in the construct (yellow) or from the sum of the predictions using the individual domains as input (green). The dashed coloured lines represent the best fit for the specific class of predictions.

---

simplified domain-additive approach.

We repeated the analysis for the different combinations of Nank repeats, again predicting the stability both from full structures and as the sum of the stability of the individual repeats. Like for SasG, we find that predictions using the full structures overestimate the absolute stability of the system, whereas stability predictions from summing the stability of the individual repeats were in better agreement with experiments (Fig. 2B). Nevertheless, while the domain-additive model gives results that are in overall better agreement with the magnitude of ΔG_{exp}, the increase in stability as additional domains are added is not captured accurately. Again, this might reflect that this model does not take into account important contributions from domain interfaces.

Finally, we tested the accuracy of our prediction against proteins with a folding mechanism more complex than two-state folding. We searched the ProTherm database (***Nikam et al., 2021***) for proteins with a multi-state folding process, measured with reversible denaturation and at standard experimental conditions (25 °C, pH 7.0). We found only few systems (less than 10) that met these criteria and selected the two monomeric systems for our analysis: creatine kinase (***Liu et al., 2009***) and exo-small β lactamase (***Santos et al., 2007***). Like for the systems described above, our model overestimates the global stability of the two proteins (Fig. S7).

Conclusions

Protein stability is a key property that underlies protein function, evolution, and our ability to engineer proteins. For example, while accurate prediction of $\Delta\Delta G$ (for one or a few substitutions) might be sufficient when optimizing the stability of a protein, accurate predictions of ΔG would enable comparing the stability of a series of orthologues or even completely different proteins. In some sense, prediction of ΔG corresponds to predicting $\Delta\Delta G$ between wildly different sequences and structures.

While the prediction of the change in stability due to one or a few substitutions ($\Delta\Delta G$) is a well-developed area of computational protein science, the prediction of the absolute stability of a protein fold has been a more difficult problem to tackle. One reason is that the global stability contains contributions from a large number of interactions by the protein and solvent in both the folded and unfolded state, making this a difficult quantity to calculate or learn. Also, since $\Delta\Delta G$ is less likely to depend on experimental conditions than ΔG, training and benchmarking predic-

tion methods for $\Delta\Delta G$ can use a wider set of experimental data that are collected under different conditions.

Given the relative scarcity of experimental ΔG values for distinct proteins under similar conditions, we have here taken the approach of using a pre-trained deep-learning model as basis for our predictions, rather than training or parameterizing a novel biophysical or machine learning model. Nevertheless, we show that there is enough data for benchmarking methods for prediction of absolute stability at physiologically relevant conditions. Here we use the pre-trained model in a relatively simple way (Eq. 1), and while the results are encouraging and useful, we recognize the limitations from both the use of the general ESM-IF model and the simple form of Eq. 1. We note also the limitation from examining only proteins with folded regions, and that future work should explore the contributions from disordered regions both to stability of folded domains and to the model predictions.

Despite these current limitations, our work shows how the amino acid likelihoods from an 'inverse folding model' (ESM-IF) can be used to predict, with useful accuracy, the absolute stability for a series of small–medium sized single-domain proteins with equilibrium two-state folding. As point of comparison, recently developed deep-learning methods for the prediction of $\Delta\Delta G$ achieve an accuracy of ca. 0.5–1.0 kcal/mol for small proteins with larger errors for proteins with more than 100 residues (*Blaabjerg et al., 2023*; *Dieckhaus et al., 2024*). While these numbers are smaller than the 1.5 kcal/mol that we report here for ΔG, we note that the spread of ΔG in our benchmark is greater than for $\Delta\Delta G$. Thus, while substantial work remains to be done, our results demonstrate that absolute stability predictions are possible at a level approaching that of $\Delta\Delta G$, at least for simpler systems. We expect that representations from deep learning models such as ESM-IF or other structure-aware model may be an excellent starting point for building more accurate and precise predictors in the future.

For larger proteins, including those with multiple domain, our results show more variability. In some cases we find that accuracy improves when predicting the stability via decomposing the protein into individual domains. However, when coupling between domains is substantial, such a divide-and-conquer approach may not work well. In some sense, this is analogous to the predictions of effects of multi-mutants as the sum of contributions of the individual variants, which depends on the additivity of the individual effects. While it may be possible to use our approach also to calculate interface energies between domains, such a decomposition would require automated assessment of how to divide the protein into the relevant domains.

Given the differences we observe between the results on small, simple proteins and larger more complex systems, we caution about making extrapolations about accuracy of ΔG predictions from results on small, single domain proteins. We suggest that modelling efforts will benefit both from an increase in the number of ΔG values that are measured under consistent conditions, and from making such measurements across proteins of different folds and sizes, and possibly for proteins whose equilibrium folding involves more than two states.

Another interesting avenue for future research is to use approaches such as that described here to examine other questions in protein biophysics. One example we have here explored is to predict the relative stability of different configurations of a protein. We have here tested our approach using two experimentally well-characterized systems where structural models exist for two distinct states. While the free energy differences in the two systems are only slightly larger than the average error of the stability predictions, NMR relaxation dispersion methods can be used to measure free energy differences between conformational states as large as ca. 3 kcal/mol (*Mittermaier and Kay, 2006*). Thus, as prediction accuracy increases we envision that such data will be particularly useful to benchmark methods to predict conformational free energies. We note here that ESM-IF uses only the backbone structure as input, and so may have difficulties in predicting effects that depend also on differences in side chain packing. We suggest that models that represent the atomic structure might instead be used in the future.

In summary, we have described an approach that uses a generative model to predict abso-

lute protein stability under ambient conditions, and benchmarked it across a number of data sets. While inaccuracies remain, we envision that the ability to predict whether a small–medium sized protein is stable, and by how much, will be useful across many areas of protein science. We also suggest that the data we have collected will be useful to benchmark other approaches to predict stability and other related properties.

Methods

Benchmark data

We applied a few filters to the data sets to help ensure the coherence on experimental conditions and types of folding. From the curated dataset (*Maxwell et al., 2005*) we collected 30 protein ΔG values and then removed proteins that form obligate homodimers. We also removed the protein VlsE, as it has been shown that despite being a two-state folder protein, it does not follow the usual contact order relationship (*Jones and Wittung-Stafshede, 2003*), reducing the total number of proteins to 26. We note for completeness that the predicted stability of VlsE (18 kcal/mol) is substantially greater than experiment (5.7 kcal/mol), suggesting again that our approach is limited for large proteins.

For the proteins from the previously described 'megascale' experiment (*Tsuboyama et al., 2023*), which included 478 unique proteins with estimated ΔG, we kept only natural proteins, reducing the size of the data set to 239 proteins. Where a protein ΔG was measured in multiple replicates, we used the average of the replicates as the final value.

Generation of the input structure

As ESM-IF was trained using structures predicted using AlphaFold 2 (AF2, *Jumper et al.* (*2021*)), we use LocalColabFold v.1.5.0 (AlphaFold 2.3.1) to generate a structural model for each target fold. We use LocalColabFold with MMseqs2 (*Steinegger and Söding, 2017*) and default parameters including three recycles, no relaxation of the predicted structure, and the maximum multiple sequence alignment size to generate an accurate fold. If an experimentally-derived structure is available for a target protein and conformation, we use it as a template for ColabFold to minimise possible structural deviations introduced by AF2. If the prediction by AF2 differed substantially from this template structure, we performed the prediction again while reducing the MSA size, and thus the influence of the co-evolutionary signal, until the predicted structure results similar to the input structure template (with an RMSD less than 0.5 Å). For the calculations T4 lysozyme we used PDB IDs 3DMV and 2LCB (*Bouvignies et al., 2011*) and for Cyclophilin A we based our calculations on PDB IDs 3K0N and 3K0O (*Fraser et al., 2009*).

Global stability predictions using ESM-IF

We predicted the global stability value from each structure using the Evolutionary Scale Model 'Inverse Folding' model (ESM-IF - model: esm_if1_gvp4_t16_142M_UR50, *Hsu et al.* (*2022*)). We used the output logits from the language-model decoder to evaluate the likelihoods for the amino acids at a specific position given the sequence position from the N-term to the preceding position. Specifically, we evaluated the likelihood ($\mathscr{L}_k$) for each amino acid k by applying the Softmax on the 20 amino acid logits (x) at each position:

$$\mathscr{L}_k = \text{Softmax}(x_k) = \frac{\exp(x_k)}{\sum_{j=1}^{20} \exp(x_j)})$$

Other methods for stability predictions

As comparison to the predictions based on ESM-IF we also predicted stabilities using other methods.

Absolute contact number

We evaluated the absolute contact number (ACN) of a structure using:

$$\text{ACN} = \sum_{i}^{N} \sum_{j>i}^{N} 1 \;\; if \; d_{ij} < d_{\text{cutoff}}$$

with N is the number of residues, d_{ij} is the distance between the C_α atoms of residues i and j, and $d_{\text{cutoff}} = 8$ Å.

FoldX

We predicted global stability with FoldX v.5 (*Schymkowitz et al., 2005*) using AF2 predicted structures as input. Specifically, we first used the command 'RepairPDB', followed by the command 'Stability' to evaluate the different energy contributions and used their sum (total-energy) as absolute stability value. We reverted the sign of this number to represent ΔG. We stress here that FoldX was not developed to predict ΔG.

Chroma

We predicted global stability with the Chroma model (*Ingraham et al., 2023*). Specifically, we used the Github version of Chroma (Dec. 2023) and the design protocol to collect the conditional likelihoods at diffusion time t=0.5. We then use the likelihoods for the wild-type residues and Eq. 1 to evaluate protein stability.

ESM-1b

We used ESM-1B (model: esm1b_t33_650M_UR50S, *Rives et al.* (*2021*)) to predict absolute stability using only the target protein sequence. For each protein, we collected the output logits of the language model decoder and then used the same approach as for ESM-IF to predict the absolute stability.

Predictions for Nank and SasG

For both Nank and SasG we first generated AF2 models of the individual repeats using the full structure. For G5^1-E-G5^2 we used PDB ID 3TIQ (*Gruszka et al., 2015*) and for Nank used PDB ID 1OT8 (*Mello and Barrick, 2004*). If a loop region was present between the domains/repeats, we included it in the N-terminal fragment. For the additive domain model we simply summed the ΔG values predicted for each domain in the construct.

Data and code availability

Code and data is available at https://github.com/KULL-Centre/_2024_cagiada_stability/

Acknowledgments

The research was supported by the PRISM (Protein Interactions and Stability in Medicine and Genomics) centre funded by the Novo Nordisk Foundation (NNF18OC0033950, to K.L.-L.). We acknowledge access to computational resources via a grant from the Carlsberg Foundation (CF21-0392).

References

Akdel M, Pires DE, Pardo EP, Jänes J, Zalevsky AO, Mészáros B, Bryant P, Good LL, Laskowski RA, Pozzati G, et al. A structural biology community assessment of AlphaFold2 applications. Nature Structural & Molecular Biology. 2022; 29(11):1056–1067.

Anfinsen CB. Principles that govern the folding of protein chains. Science. 1973 Jul; 181(4096):223–230.

Bastolla U, Demetrius L. Stability constraints and protein evolution: the role of chain length, composition and disulfide bonds. Protein Engineering Design and Selection. 2005; 18(9):405–415.

Blaabjerg LM, Kassem MM, Good LL, Jonsson N, Cagiada M, Johansson KE, Boomsma W, Stein A, Lindorff-Larsen
K. Rapid protein stability prediction using deep learning representations. Elife. 2023; 12:e82593.

Bouvignies G, Vallurupalli P, Hansen DF, Correia BE, Lange O, Bah A, Vernon RM, Dahlquist FW, Baker D, Kay
LE. Solution structure of a minor and transiently formed state of a T4 lysozyme mutant. Nature. 2011 Aug;
477(7362):111–114.

Chen T, Gong C. HotProtein: A novel framework for protein thermostability prediction and editing. NeurIPS
2022. 2022; .

Chu KS, Siegel JB. Protein Stability Prediction by Fine-tuning a Protein Language Model on a Mega-scale Dataset.
bioRxiv. 2023; p. 2023–11.

Dieckhaus H, Brocidiacono M, Randolph NZ, Kuhlman B. Transfer learning to leverage larger datasets for
improved prediction of protein stability changes. Proceedings of the National Academy of Sciences. 2024;
121(6):e2314853121.

Eisenmesser EZ, Millet O, Labeikovsky W, Korzhnev DM, Wolf-Watz M, Bosco DA, Skalicky JJ, Kay LE, Kern D.
Intrinsic dynamics of an enzyme underlies catalysis. Nature. 2005 Nov; 438(7064):117–121.

Fersht A. Structure and Mechanism in Protein Science: A Guide to Enzyme Catalysis and Protein Folding. World
Scientific; 2017.

Fraser JS, Clarkson MW, Degnan SC, Erion R, Kern D, Alber T. Hidden alternative structures of proline isomerase
essential for catalysis. Nature. 2009; 462(7273):669–673.

Ghosh K, Dill KA. Computing protein stabilities from their chain lengths. Proc Natl Acad Sci U S A. 2009 Jun;
106(26):10649–10654.

Gruszka DT, Whelan F, Farrance OE, Fung HKH, Paci E, Jeffries CM, Svergun DI, Baldock C, Baumann CG, Brock-
well DJ, Potts JR, Clarke J. Cooperative folding of intrinsically disordered domains drives assembly of a strong
elongated protein. Nat Commun. 2015 Jun; 6:7271.

Gruszka DT, Wojdyla JA, Bingham RJ, Turkenburg JP, Manfield IW, Steward A, Leech AP, Geoghegan JA, Foster
TJ, Clarke J, Potts JR. Staphylococcal biofilm-forming protein has a contiguous rod-like structure. Proc Natl
Acad Sci U S A. 2012 Apr; 109(17):E1011–8.

Hsu C, Verkuil R, Liu J, Lin Z, Hie B, Sercu T, Lerer A, Rives A. Learning inverse folding from millions of predicted
structures. In: *International conference on machine learning* PMLR; 2022. p. 8946–8970.

Ingraham JB, Baranov M, Costello Z, Barber KW, Wang W, Ismail A, Frappier V, Lord DM, Ng-Thow-Hing C,
Van Vlack ER, Tie S, Xue V, Cowles SC, Leung A, Rodrigues JV, Morales-Perez CL, Ayoub AM, Green R, Puentes
K, Oplinger F, et al. Illuminating protein space with a programmable generative model. Nature. 2023 Nov;
623(7989):1070–1078.

Jones K, Wittung-Stafshede P. The largest protein observed to fold by two-state kinetic mechanism does not
obey contact-order correlation. J Am Chem Soc. 2003 Aug; 125(32):9606–9607.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A,
Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J,
Back T, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug; 596(7873):583–
589.

Lindorff-Larsen K, Teilum K. Linking thermodynamics and measurements of protein stability. Protein Engi-
neering, Design and Selection. 2021; 34:gzab002.

Liu YM, Feng S, Ding XL, Kang CF, Yan YB. Mutation of the conserved Asp122 in the linker impedes creatine
kinase reactivation and refolding. Int J Biol Macromol. 2009 Apr; 44(3):271–277.

Maxwell KL, Wildes D, Zarrine-Afsar A, De Los Rios MA, Brown AG, Friel CT, Hedberg L, Horng JC, Bona D, Miller
EJ, Vallée-Bélisle A, Main ERG, Bemporad F, Qiu L, Teilum K, Vu ND, Edwards AM, Ruczinski I, Poulsen FM,
Kragelund BB, et al. Protein folding: defining a "standard" set of experimental conditions and a preliminary
kinetic data set of two-state proteins. Protein Sci. 2005 Mar; 14(3):602–616.

Mello CC, Barrick D. An experimentally determined protein folding energy landscape. Proc Natl Acad Sci U S A.
2004 Sep; 101(39):14102–14107.

Mittermaier A, Kay LE. New tools provide new insights in NMR studies of protein dynamics. science. 2006;
312(5771):224–228.

Mulder FA, Mittermaier A, Hon B, Dahlquist FW, Kay LE. Studying excited states of proteins by NMR spec-
troscopy. Nature structural biology. 2001; 8(11):932–935.

Nikam R, Kulandaisamy A, Harini K, Sharma D, Gromiha MM. ProThermDB: thermodynamic database for
proteins and mutants revisited after 15 years. Nucleic Acids Res. 2021 Jan; 49(D1):D420–D424.

Nisthal A, Wang CY, Ary ML, Mayo SL. Protein stability engineering insights revealed by domain-wide compre-
hensive mutagenesis. Proceedings of the National Academy of Sciences. 2019; 116(33):16367–16377.

Notin P, Kollasch A, Ritter D, Van Niekerk L, Paul S, Spinner H, Rollins N, Shaw A, Orenbuch R, Weitzman R, Frazer
J, Dias M, Franceschi D, Orenbuch R, Gal Y, Marks DS. ProteinGym: large-scale benchmarks for protein fitness
prediction and design. Advances in Neural Information Processing Systems. 2024; 36.

Pancotti C, Benevenuta S, Birolo G, Alberini V, Repetto V, Sanavia T, Capriotti E, Fariselli P. Predicting protein
stability changes upon single-point mutation: a thorough comparison of the available tools on a new dataset.
Brief Bioinform. 2022 Mar; 23(2).

Papaleo E, Sutto L, Gervasio FL, Lindorff-Larsen K. Conformational changes and free energies in a proline
isomerase. Journal of Chemical Theory and Computation. 2014; 10(9):4169–4174.

Park C, Marqusee S. Pulse proteolysis: a simple method for quantitative determination of protein stability and
ligand binding. Nature methods. 2005; 2(3):207–212.

Pucci F, Rooman M. Stability curve prediction of homologous proteins using temperature-dependent statistical
potentials. PLoS computational biology. 2014; 10(7):e1003689.

Pucci F, Schwersensky M, Rooman M. Artificial intelligence challenges for predicting the impact of mutations
on protein stability. Current opinion in structural biology. 2022; 72:161–168.

Reeves S, Kalyaanamoorthy S. Zero-Shot Transfer of Protein Sequence Likelihood Models to Thermostability
Prediction. bioRxiv. 2023; p. 2023–07.

Rives A, Meier J, Sercu T, Goyal S, Lin Z, Liu J, Guo D, Ott M, Zitnick CL, Ma J, Fergus R. Biological structure and
function emerge from scaling unsupervised learning to 250 million protein sequences. Proc Natl Acad Sci U
S A. 2021 Apr; 118(15).

Robertson AD, Murphy KP. Protein structure and the energetics of protein stability. Chemical reviews. 1997;
97(5):1251–1268.

Ruiz-Blanco YB, Marrero-Ponce Y, Paz W, García Y, Salgado J. Global stability of protein folding from an empir-
ical free energy function. J Theor Biol. 2013 Mar; 321:44–53.

Santos J, Risso VA, Sica MP, Ermácora MR. Effects of serine-to-cysteine mutations on beta-lactamase folding.
Biophys J. 2007 Sep; 93(5):1707–1718.

Schymkowitz J, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. The FoldX web server: an online force field.
Nucleic Acids Res. 2005 Jul; 33(Web Server issue):W382–8.

Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive
data sets. Nat Biotechnol. 2017 Nov; 35(11):1026–1028.

Thompson MC, Barad BA, Wolff AM, Sun Cho H, Schotte F, Schwarz DM, Anfinrud P, Fraser JS. Temperature-
jump solution X-ray scattering reveals distinct motions in a dynamic enzyme. Nature chemistry. 2019;
11(11):1058–1066.

Tsuboyama K, Dauparas J, Chen J, Laine E, Mohseni Behbahani Y, Weinstein JJ, Mangan NM, Ovchinnikov S,
Rocklin GJ. Mega-scale experimental analysis of protein folding stability in biology and design. Nature. 2023
Jul; p. 1–11.

Wapeesittipan P, Mey AS, Walkinshaw MD, Michel J. Allosteric effects in cyclophilin mutants may be explained
by changes in nano-microsecond time scale motions. Communications Chemistry. 2019; 2(1):41.

Xavier JS, Nguyen TB, Karmarkar M, Portelli S, Rezende PM, Velloso JP, Ascher DB, Pires DE. ThermoMutDB: a
thermodynamic database for missense mutations. Nucleic acids research. 2021; 49(D1):D475–D479.

Yang Y, Zhao J, Zeng L, Vihinen M. ProTstab2 for Prediction of Protein Thermal Stabilities. Int J Mol Sci. 2022
Sep; 23(18).

Supplementary Figures

Figure S1. Properties of the benchmark data set. Distribution of (A) ΔG_{exp} and (B) number of residues for the 265 proteins in the benchmark data set.

Figure S2. Predicting absolute protein stability using baseline models. We show a comparison between ΔG_{exp} values for 265 proteins from two different data sets (represented by different colours) and (A) the number of residues, (B) the total number of contacts formed in the structure and, for comparison, (C) the predictions using ESM-IF likelihoods. Panel C is similar to Fig. 1 (without converting to physical units) and is shown here for comparison. The Spearman's and Pearson's correlation coefficient are shown in each panel. The reported standard errors were obtained by bootstrapping.

Figure S3. Comparison between absolute protein stability and summed entropy. We show the correlation between the ΔG_{exp} (x-axis) values for the 265 proteins in our benchmark dataset and the stability prediction (y-axis) obtained through the sum of positional entropy from ESM-IF amino acid likelihoods. The reported standard error was obtained by bootstrapping.

Figure S4. Accuracy of absolute stability predictions using amino acid subgroups. (A) Box plots show the contribution of different amino acid types to the overall absolute stability for each of the 265 proteins in our benchmark set. The Spearman's and Pearson's correlation coefficients with the experimental measures are shown at the top of each amino class boxplot. We also show the correlation between the ΔG_{exp} values for the 265 proteins in our dataset and the absolute stability calculated from the summed likelihoods of (B) valines and isoleucines only, (C) hydrophobic residues only, and (D) buried residues only. In each panel we show Spearman's and Pearson's correlation coefficients and standard errors obtained by bootstrapping.

Figure S5. Predicting absolute protein stability using alternative models. We show a comparison between ΔG_{exp} values for 265 proteins from two different data sets (represented by different colours). (A) Using likelihoods ($\mathscr{L}$) from ESM-1b together with Eq. 1. (B) Total energies from FoldX. (C) Using likelihoods from Chroma together with Eq. 1. For comparison we show (D) Using likelihoods from ESM-IF together with Eq. 1. Panel D is similar to Fig. 1 (without converting to physical units) and is shown here for comparison The Spearman's and Pearson's correlation coefficient are shown in each panel (ρ_s, ρ_p). The reported standard errors were obtained by bootstrapping.

Figure S6. Benchmarking calculations for more complex problems and systems. (A) Structure of the G and E states of T4 lysozyme L99A variants. (B) Structure of the G and E states of Cyclophilin A. (C) Structure of Nank1–7 with each repeat indicated by a different colour. (D) Structure of SasG indicating the location of the G5[1], E and G5[2] domains.

Figure S7. Stability prediction for non-two-state folding proteins. We compared the predicted ΔG (green) with the experimental free energy differences going from the native to intermediate states (yellow) and from the intermediate state to the unfolded state (red). Results are shown for (left) creatine kinase and (right) the exo-small β lactamase.