Predicting absolute protein folding stability using generative models

Matteo Cagiada[1*]**, Sergey Ovchinnikov**[2]**, Kresten Lindorff-Larsen**[1*]

*For correspondence:
matteo.cagiada@bio.ku.dk (MC);
lindorff@bio.ku.dk (KLL)

[1]Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen, DK-2200 Copenhagen, Denmark; [2]Massachusetts Institute of Technology, Cambridge, USA

Abstract

While there has been substantial progress in our ability to predict changes in protein stability due to amino acid substitutions, progress has been slower in methods to predict the absolute stability of a protein. Here we show how a generative model for protein sequence can be leveraged to predict absolute protein stability. We benchmark our predictions across a broad set of proteins and find a mean error of 1.5 kcal/mol and a correlation coefficient of 0.7 for the absolute stability across a range of natural, small–medium sized proteins up to ca. 150 amino acid residues. We analyse current limitations and future directions including how such model may be useful for predicting conformational free energies. Our approach is simple to use and freely available via an online implementation.

Introduction

The stability of a protein is a critical property that affects its biological functions, as well as our ability to engineer proteins for new purposes. This stability is mainly the result of a delicate balance between the atomic interactions in the folded state, the higher conformational entropy in the unfolded state, as well as a complex relationship between the protein structure and water properties (*Anfinsen, 1973*; *Fersht, 2017*). By protein stability, we here concern ourselves with the (thermodynamic) free energy of unfolding, ΔG, under a given set of conditions such as temperature and pH.

Experimentally, ΔG can be measured using a range of biophysical and spectroscopic techniques, though generally the value under native-like conditions requires extrapolations from for example high concentrations of denaturant or elevated temperatures (*Lindorff-Larsen and Teilum, 2021*). These measurements generally require purification of each variant and is relatively labour intensive, even though automated methods have been used (*Nisthal et al., 2019*). While the thermodynamic stability has been determined for a number of proteins and their variants using these approaches, the resulting data are relatively inhomogeneous in terms of methods used and experimental conditions (*Nikam et al., 2021*; *Xavier et al., 2021*). As an alternative, protein stability may in certain cases also be indirectly quantified using kinetic experiments (*Park and Marqusee, 2005*), and this has recently been combined with large-scale DNA synthesis and display technologies to quantify protein stability at a very large scale (*Tsuboyama et al., 2023*). While this type of experiments has provided a wealth of information, it is for technical reasons currently limited to small proteins with less than about 80 amino acid residues and which fold in a two-state manner.

As a supplement to experimental measurements, computational predictions can be used to assess the stability of a protein. With few exceptions, these methods have mostly focused on predictions of *changes* in thermodynamic stability ($\Delta\Delta G$), typically between a protein and a variant that

differs from it by one or very few amino acid substitutions. A number of methods to predict $\Delta\Delta G$ have been developed (*Pancotti et al., 2022*), and their accuracy in part originates from a number of important compensating effects. These include that while ΔG may be sensitive to the conditions (pH, temperature, etc), $\Delta\Delta G$ will be less sensitive. Similarly, while both ΔG and $\Delta\Delta G$ contain contributions from many interactions across the protein, many of these will cancel out when comparing the stability of a wild-type protein and a variant with a single amino acid substitution. Thus, in these senses $\Delta\Delta G$ is a more robust quantity than ΔG.

One additional complication in analysing and predicting protein stability is that many of the underlying thermodynamic parameters (ΔH, ΔS and ΔC_p) are strongly correlated with the number of residues (length) of the protein (*Robertson and Murphy, 1997*). These effects in turn suggest that it will be easier to predict the thermal stability (melting temperature) of a protein than the stability at a fixed temperature (*Pucci and Rooman, 2014*), and indeed several methods have been developed to predict melting temperatures. Previous work used these properties to predict, for example, stability from chain length (*Ghosh and Dill, 2009*; *De Sancho et al., 2009*) with more recent methods taking additional physical and statistical information into account (*Ruiz-Blanco et al., 2013*; *Pucci and Rooman, 2014*; *Yang et al., 2022*).

While progress in predicting stability (ΔG) has been relatively slow, developments in the field of machine learning have resulted in an increase in the number of prediction methods to estimate changes in protein stability ($\Delta\Delta G$) (*Pucci et al., 2022*). While these developments are encouraging, a recent example illustrates the difficulty of training models that learn both ΔG and $\Delta\Delta G$ (*Chu and Siegel, 2023*). Specifically, the authors trained a model directly on large-scale measurements of ΔG for small proteins, and showed that the model could be used to estimate both the global stability of similar folds and the effect of substitutions. However, the work also showed limited transferability, suggesting that training broadly transferable models for ΔG on a set of small proteins may be difficult.

Our work here uses a generative model as input to predict protein stability. Generative models for proteins are deep learning architectures that can be used in protein design to generate novel protein sequences or structures. They are generally based on representations learned from self-supervised training on large sequence or structure databases. In this class of models, fixed-backbone design architectures use the backbone coordinates of a target fold to generate novel sequences that fit the given structure, implicitly maximizing the fitness of the amino acid side chains in the local protein environment. Among these models, the so-called 'inverse folding' approach of the Evolutionary Scale Model family (ESM-IF)((*Hsu et al., 2022*)) has been trained to recover and generate novel amino acid sequences given an input protein fold during training on millions of sequence-structure pairs. ESM-IF achieves this by using an invariant graph neural network followed by a generic transformer(*Vaswani, 2017*). First, the graph model uses rotation equivalence and invariant transformations to generate residue embeddings that represent the geometric features of a backbone for a target protein fold, and then the transformer module uses the generated embeddings to return amino acid likelihoods and sampled sequences. The model generates likelihoods and sequences autoregressively, decoding the amino acid from the N- to C-terminus and using the information from the previous sampled positions to condition the next predicted residue.

It has been shown that the ESM-IF likelihoods can be used to assess $\Delta\Delta G$ by comparing the likelihoods of the wild-type and variant sequence, using the wild-type structure as input (*Hsu et al., 2022*; *Chen and Gong, 2022*; *Reeves and Kalyaanamoorthy, 2023*; *Notin et al., 2024*). Here, we explored whether models such as ESM-IF can also be used to solve the more difficult problem of predicting protein stability across different folds and sequences (ΔG). Our manuscript is structured as follows. First, we show how information from ESM-IF can be used to predict absolute stability for proteins with an equilibrium two-state folding mechanism. Second, we explore how well the model can assess the relative stability between two different configurations of a protein. Third, we show limitations for predicting stability for larger and more complex systems, where cooperativity between structural elements may also play a role. Finally, we briefly discuss issues that we believe

94 need to be addressed by experiments and future models in order to advance the field.

Results and Discussion
95

Predicting global folding stability using generative models
96

97 It has been shown that the ESM-IF model can be used to predict changes in protein stability (*Hsu*
98 *et al., 2022*; *Chen and Gong, 2022*; *Reeves and Kalyaanamoorthy, 2023*; *Notin et al., 2024*), sug-
99 gesting that it might encode some level of protein biophysics that relates protein structure to ther-
100 modynamics. We therefore hypothesized that the ESM-IF model could also be used to predict the
101 absolute stability of a given fold (ΔG) given its sequence. Because protein stability may vary sub-
102 stantially across conditions, we created a benchmark data set of experimental ΔG measurements
103 by merging a curated data set of ΔG measurements of two-state folding proteins obtained under
104 'standard' experimental conditions (25 °C, pH 7.0) (*Maxwell et al., 2005*) with ΔG measurements
105 from a large-scale proteolysis experiment on small single-domain proteins (*Tsuboyama et al., 2023*)
106 (performed at ca. 25 °C, pH 7.4). In total, the benchmark data include 265 proteins ranging in size
107 from 30 to 150 residues (Fig. S1A) and ΔG ranging from 1–12 kcal/mol (Fig. S1B). The dataset in-
108 cludes (i) 26 proteins whose kinetics and thermodynamics have been studied biophysically and that
109 range in size from 61–151 residues (*Maxwell et al., 2005*) and (ii) 239 naturally occurring wild-type
110 proteins (40–72 residues long) from the recently described 'megascale' experiment (*Tsuboyama*
111 *et al., 2023*).

112 We used this data set to assess whether conditional likelihoods for amino acids calculated from
113 ESM-IF for a given structure could be used to predict its global stability. Since it has been shown
114 that protein stability correlates with protein size (*Ghosh and Dill, 2009*; *De Sancho et al., 2009*), we
115 used as measure the sum of likelihoods for the target sequence $\sum_i^N \mathscr{L}_i^{\mathrm{WT}}$ (Eq. 1). Here, the sum
116 runs over the N residues in the protein, and $\mathscr{L}_i^{\mathrm{WT}}$ is the likelihood from ESM-IF for having the refer-
117 ence (wild-type) amino acid at position i (see Methods). We calculated this measure for each of the
118 265 proteins and obtain a Pearson's correlation of 0.69 with the experimental ΔG values, demon-
119 strating how ESM-IF can be used to predict absolute protein stabilities (Fig. 1A). We then compared
120 the accuracy achieved by Eq. 1 to other simpler measures such as the number of contacts in the
121 fold or the sequence length (*Bastolla and Demetrius, 2005*; *Ghosh and Dill, 2009*; *De Sancho et al.,*
122 *2009*). We find that both of these measures are substantially less correlated with the experimental
123 ΔG values than ΔG_{pred} calculated from ESM-IF (Fig. 1B,C). We also assessed whether the summed
124 model entropy could be used to predict protein stability and find that there was no correlation with
125 experimental data (Fig. S2).

126 We then convert the values from Eq. 1 into ΔG_{pred} in physical units, by fitting a scaling constant
127 and offset using the experimental data (see Methods). The resulting predictions have a root-mean-
128 square-error between ΔG_{exp} and ΔG_{pred} of 1.4 kcal/mol (Fig. 2), mostly constant across proteins
129 of different stabilities (Fig. S3). This conversion from likelihoods to physical units involves fitting
130 two parameters, and cross-validation shows the errors to be robust (Fig. S4), and lower than the
131 baselines we examined (Table 1).

132 Having assessed that the summed likelihoods correlate with global stability, we investigated
133 whether absolute stability was correlated with the summed likelihoods of specific amino acid types
134 or groups of residue types. We found that the summed likelihoods of two aliphatic residue types,
135 valine and isoleucine, correlated the strongest with the global stability (Fig. 3A), and the sum of
136 these two contributions correlate strongly with global stability (ρ_P: 0.64, Fig. 3B). When we aggre-

Table 1. RMSE comparison for the null model and different models, including ESM-IF, for the predictions
fitted with experimental data on the experimental data, again.

Method	Null model	Sequence model	ACN model	ESM-IF model
RMSE	1.88 kcal/mol	1.55 kcal/mol	1.53 kcal/mol	1.39 kcal/mol

Figure 1. Predicting absolute protein stability. We show a comparison between ΔG_{exp} values for 265 proteins from two different data sets (represented by different colours) and (A) the sum of raw likelihoods predicted by ESM-IF, (B) the total number of contacts formed in the structure and, for comparison, (C) the number of residues. The Spearman's and Pearson's correlation coefficient are shown in each panel. The reported standard errors were obtained by bootstrapping.

Figure 2. Predicting absolute protein stability using a generative model. We show a comparison between predicted and experimental stability values for 265 proteins from two different data sets (represented by different colours). The predicted values are based on Eq. 1 and have been rescaled to units of kcal/mol (ΔG_{pred}). The Pearson's correlation coefficient (ρ_p) and the RMSE (in kcal/mol) are shown. The standard error for the correlation coefficient was obtained by bootstrapping.

Figure 3. Accuracy of absolute stability predictions using amino acid subgroups. (A) Box plots show the contribution of different amino acid types to the overall absolute stability for each of the 265 proteins in our benchmark set. The Spearman's and Pearson's correlation coefficients with the experimental measures are shown at the top of each amino class boxplot. We also show the correlation between the $\Delta G_{\exp}$ values for the 265 proteins in our dataset and the absolute stability calculated from the summed likelihoods of (B) valines and isoleucines only, (C) hydrophobic residues only, and (D) buried residues only. In each panel we show Spearman's and Pearson's correlation coefficients and standard errors obtained by bootstrapping.

gated the amino acid likelihoods over all the hydrophobic amino acids, the results correlate with the global stability as much as the total sum of the likelihoods (ρ_P: 0.67, Fig. 3C), in line with the assumption that protein stability is largely determined by the hydrophobic effect; the correlation using only buried residues was slightly lower (ρ_P: 0.63, Fig. 3D). Together, these results show that the summed model likelihoods correlate with global protein stability, and that this correlation to a large extent is driven by the hydrophobic amino acids. We note that these correlations are all obtained by representing the protein structure solely via the polypeptide backbone.

Having shown that our approach can be used to predict absolute folding stability at a level greater than simpler baseline models, we then asked whether other machine learning models than ESM-IF might achieve similar accuracy using the same approach. To investigate this we used Eq. 1 together with likelihoods obtained from (i) a sequence-based language model (ESM-1b) (*Rives et al., 2021*) to assess the amount of information that could be extracted directly from the sequence using a generic transformer architecture or (ii) likelihoods from a different generative model (Chroma; *Ingraham et al.* (*2023*)) as a comparison between different design architectures. We also tested energy evaluations using the empirical force field FoldX (*Schymkowitz et al., 2005*), which can be used to predict changes in protein stability. Among the methods tested, we find that the ESM-IF likelihoods achieved the highest correlation with experimental stability measurements (Table 2 and Fig. S5). We note that the sequence-based model, when used together with Eq. 1, achieves a near-zero correlation with the stability measures, highlighting that the stability of a given fold is a difficult task to learn from evolutionary conservation information alone. We also note that while FoldX has been shown to be useful for predicting $\Delta\Delta G$ from both experimental (*Schymkowitz et al., 2005*) and predicted (*Akdel et al., 2022*) structures, it was not developed to predict ΔG.

Table 2. Pearson's correlation coefficient between the different alternative model tested including ESM-IF, for the predictions of absolute stability on the experimental dataset

Method	ESM-1b	FoldX	Chroma	ESM-IF
ρ_P	0.25	0.04	0.62	0.69

Figure 4. Evaluation of the ability to predict the free energy difference between possible conformations for selected proteins. For proteins with experimentally determined structures of distinct conformational states and the free energy difference between these states, such as the G and E states of (A) T4 lysozyme (T4L) L99A variants and Cyclophilin A (CYPA), we calculate (C) their free energy difference as the difference between their predicted absolute stability. We have predicted and compared with the experimental conformational free energies for (D) T4L and (E) CYPA. In panels D and E, the left plots show the predicted absolute stability for the two states; the right plots shows a comparison of the experimental stability difference and the difference between the predicted absolute folding stabilities.

Prediction of stability differences between different protein conformations

Having shown that we can use the ESM-IF likelihoods to predict absolute protein stability from a given structure, we then asked whether we could use the same model to predict the difference in stability between two distinct conformations (Fig. 4C). By thermodynamic consistency, this corresponds to predicting the free energy difference between two distinct conformational states, each represented by a single backbone structure.

To assess this ability, we selected two systems where the energetic difference between two states in solution has been measured experimentally using NMR relaxation dispersion techniques. The first system is the L99A variant of T4 lysozyme (T4L L99A, Fig. 4A), which is known to exchange between two conformations, termed G (ground) and E (excited), and with the E state being less stable than the G state by ca. 2 kcal/mol (*Mulder et al., 2001*; *Bouvignies et al., 2011*). Using structures of the G and E states (*Bouvignies et al., 2011*) as input to ESM-IF and Eq. 1 we predict that the E state is 1.2 kcal/mol less stable than G, in qualitative agreement with experiments (Fig. 4D).

As the second example, we selected Cyclophilin A (CYPA, Fig. 4B) where NMR experiments have shown a free energy difference between two states of 1.7 kcal/mol (*Eisenmesser et al., 2005*), and where models of the two states have been obtained by crystallography (*Fraser et al., 2009*).

175 Again, we used ESM-IF to predict the free energy difference between the two configurations. As
176 for T4L L99A, we also predict a free energy difference whose sign is correct, but with a magnitude
177 (0.3 kcal/mol) that is substantially smaller than found in experiments (1.7 kcal/mol; Fig. 4E). We
178 here note that the ESM-IF model only uses the backbone structure as input, and that it is still not
179 fully known whether the free energy difference in CYPA measured by NMR is fully reflected in the
180 small structural difference (C_α RMSD of 0.34 Å), between the two structures (*Eisenmesser et al.,*
181 *2005*; *Fraser et al., 2009*; *Papaleo et al., 2014*; *Wapeesittipan et al., 2019*; *Thompson et al., 2019*).
182   While these results demonstrate that our approach cannot yet accurately predict free energy
183 differences between states, they suggest that as prediction accuracy of the stability predictions
184 increase, it might be possible to resolve conformational free energies of the magnitudes found in
185 these two test systems.

Prediction of the absolute stability of more complex systems
186
187 Having shown that ESM-IF provides relatively accurate predictions of absolute stability for small–
188 medium sized proteins that fold in a two-state manner, we asked whether the model could also
189 be used to estimate stability for more complex systems. Specifically, we were interested in as-
190 sessing whether our approach could also be used to predict the stability of larger, multi-domain
191 proteins—potentially with cooperativity between the different domains—and for proteins where
192 global unfolding does not follow an equilibrium two-state mechanism.
193   To assess this, we first study two different protein systems, each involving repeated domains:
194 the G5[1]-E-G5[2] repeat of the Surface protein G (SasG, Fig. 5A; *Gruszka et al.* (*2015*)) and the Notch
195 ankyrin domain (Nank, Fig. 5B; *Mello and Barrick* (*2004*)). In both cases, experimental measures
196 of absolute stability are available for different combinations of the domains (*Gruszka et al., 2015*;
197 *Mello and Barrick, 2004*). While our approach to predict stability captures the size-dependency
198 across small proteins, we wondered whether it might overestimate stability for larger proteins
199 with (semi-)independent domains. We therefore evaluated the stability for the different domain
200 constructs using either (i) the full structure as input to ESM-IF or (ii) using the sum of stability pre-
201 dictions for the individual domains.
202   For the SasG system we find that the absolute stabilities predicted for the different combina-
203 tions of domains (G5[1]-E, E-G5[2], and G5[1]-E-G5[2]) are substantially overestimated when we use the full
204 structure as input to ESM-IF (Fig. 5C). When we instead calculate the stability as the sum of those
205 of the individual domains we obtain results that are closer to experiments, though still overesti-
206 mated relative to experiments (Fig. 5D). The imperfect agreement may in part originate from the
207 complexity of the system; the E-domain located in between the G5[1] and G5[2] domains is disordered
208 on its own and only folds when located between the folded G-domains (*Gruszka et al., 2012*). Thus,
209 the calculations for the G5[1]-E and E-G5[2] constructs (using models with folded E domains) overesti-
210 mate the stability. Also, the stability of the G5[1]-E-G5[2] construct contains a substantial contribution
211 from the interfaces between the different domains (*Gruszka et al., 2015*) that is not captured by
212 our simplified domain-additive approach.
213   We repeated the analysis for the different combinations of Nank repeats, again predicting the
214 stability both from full structures and as the sum of the stability of the individual repeats. Like for
215 SasG, we find that predictions using the full structures overestimate the absolute stability of the
216 system, whereas stability predictions from summing the stability of the individual repeats were in
217 better agreement with experiments (Fig. 4B). Nevertheless, while the domain-additive model gives
218 results that are in overall better agreement with the magnitude of ΔG_{exp}, the increase in stability as
219 additional domains are added is not captured accurately. Again, this might reflect that this model
220 does not take into account important contributions from domain interfaces.
221   Finally, we tested the accuracy of our prediction against proteins with a folding mechanism
222 more complex than two-state folding. We searched the ProTherm database (*Nikam et al., 2021*) for
223 proteins with a multi-state folding process, measured with reversible denaturation and at standard
224 experimental conditions (25 °C, pH 7.0). We found only few systems (less than 10) that met these

Figure 5. Predicting stability of multi-domain repeat proteins. Starting from (A) SasG and (B) Nank structures we calculated (C,D) the stability of different domain constructs using either structural models with all domains in the construct (yellow) or from the sum of the predictions using the individual domains as input (green). The dashed coloured lines represent the best fit for the specific class of predictions.

225 criteria and selected the two monomeric systems for our analysis: creatine kinase (*Liu et al., 2009*)
226 and exo-small β lactamase (*Santos et al., 2007*). Like for the systems described above, our model
227 overestimates the global stability of the two proteins (Fig. S6).

Conclusions

229 Protein stability is a key property that underlies protein function, evolution, and our ability to en-
230 gineer proteins. For example, while accurate prediction of $\Delta\Delta G$ (for one or a few substitutions)
231 might be sufficient when optimizing the stability of a protein, accurate predictions of ΔG would
232 enable comparing the stability of a series of orthologues or even completely different proteins. In
233 some sense, prediction of ΔG corresponds to predicting $\Delta\Delta G$ between wildly different sequences
234 and structures.

235 While the prediction of the change in stability due to one or a few substitutions ($\Delta\Delta G$) is a
236 well-developed area of computational protein science, the prediction of the absolute stability of
237 a protein fold has been a more difficult problem to tackle. One reason is that the global stability
238 contains contributions from a large number of interactions by the protein and solvent in both the
239 folded and unfolded state, making this a difficult quantity to calculate or learn. Also, since $\Delta\Delta G$
240 is less likely to depend on experimental conditions than ΔG, training and benchmarking predic-
241 tion methods for $\Delta\Delta G$ can use a wider set of experimental data that are collected under different
242 conditions.

243 Given the relative scarcity of experimental ΔG values for distinct proteins under similar condi-
244 tions, we have here taken the approach of using a pre-trained deep-learning model as basis for
245 our predictions, rather than training or parameterizing a novel biophysical or machine learning
246 model. Nevertheless, we show that there is enough data for benchmarking methods for predic-
247 tion of absolute stability at physiologically relevant conditions. Here we use the pre-trained model
248 in a relatively simple way (Eq. 1), and while the results are encouraging and useful, we recognize the
249 limitations from the use of the general ESM-IF model and the simple form of Eq. 1. For example, the
250 ESM-IF model represents protein structures via the backbone coordinates providing limitations to
251 the kinds of interactions that are explicitly represented. Our results indicate that the stability of nat-
252 ural proteins is to a substantial extent determined by how likely the amino acid sequence is, with a
253 focus on the side chain packing in the protein chemical environment especially for buried regions
254 of the target fold. While it is not surprising that the environment of the side chains in the structure
255 influences stability, the finding that a simple sum of likelihoods can be used to predict stability of
256 the kinds of proteins we have examined suggests that less stable natural proteins have residues
257 that are suboptimal for their structures. We note that these predictions can be made even though
258 the model does not explicitly represent for example backbone strain or dynamics. While some of
259 these properties may have been learned indirectly during training of ESM-IF, it is also likely that
260 the imperfections in model predictions come from limitations in how the model represents pro-
261 tein biophysics. We also note that the ESM-IF model was trained on the structure and sequences
262 of natural proteins, and that our tests also involve natural (and mostly single-domain) proteins. It
263 thus remains to be tested how accuracy extends to a larger region of the protein sequence and
264 structure universe.

265 Despite these current limitations, our work shows how the amino acid likelihoods from an 'in-
266 verse folding model' (ESM-IF) can be used to predict, with useful accuracy, the absolute stability for
267 a series of small–medium sized single-domain proteins with equilibrium two-state folding. As point
268 of comparison, recently developed deep-learning methods for the prediction of $\Delta\Delta G$ achieve an
269 accuracy of ca. 0.5–1.0 kcal/mol for small proteins with larger errors for proteins with more than
270 100 residues (*Blaabjerg et al., 2023*; *Dieckhaus et al., 2024*). While these numbers are smaller than
271 the 1.5 kcal/mol that we report here for ΔG, we note that the spread of ΔG in our benchmark is
272 greater than for $\Delta\Delta G$. Thus, while substantial work remains to be done, our results demonstrate
273 that absolute stability predictions are possible at a level approaching that of $\Delta\Delta G$, at least for sim-
274 pler systems. We expect that representations from deep learning models such as ESM-IF or other

structure-aware models, ideally also representing amino acid side chains, may be an excellent starting point for building more accurate and precise predictors in the future.

For larger proteins, including those with multiple domain, our results show more variability. In some cases we find that accuracy improves when predicting the stability via decomposing the protein into individual domains. However, when coupling between domains is substantial, such a divide-and-conquer approach may not work well. In some sense, this is analogous to the predictions of effects of multi-mutants as the sum of contributions of the individual variants, which depends on the additivity of the individual effects. While it may be possible to use our approach also to calculate interface energies between domains, such a decomposition would require automated assessment of how to divide the protein into the relevant domains.

Given the differences we observe between the results on small, simple proteins and larger more complex systems, we caution about making extrapolations about accuracy of ΔG predictions from results on small, single domain proteins. We suggest that modelling efforts will benefit both from an increase in the number of ΔG values that are measured under consistent conditions, and from making such measurements across proteins of different folds and sizes, and possibly for proteins whose equilibrium folding involves more than two states.

Another interesting avenue for future research is to use approaches such as that described here to examine other questions in protein biophysics. One example we have here explored is to predict the relative stability of different configurations of a protein. We have here tested our approach using two experimentally well-characterized systems where structural models exist for two distinct states. While the free energy differences in the two systems are only slightly larger than the average error of the stability predictions, NMR relaxation dispersion methods can be used to measure free energy differences between conformational states as large as ca. 3 kcal/mol (*Mittermaier and Kay, 2006*). Thus, as prediction accuracy increases we envision that such data will be particularly useful to benchmark methods to predict conformational free energies. We again remind that ESM-IF uses only the backbone structure as input, and so may have difficulties in predicting effects that depend also on differences in side chain packing. Also here, models that represent the atomic structure including side chains might instead be used in the future.

In summary, we have described an approach that uses a generative model to predict absolute protein stability under ambient conditions, and benchmarked it across a number of data sets. While inaccuracies remain, we envision that the ability to predict whether a small–medium sized protein is stable, and by how much, will be useful across many areas of protein science. We also suggest that the data we have collected will be useful to benchmark other approaches to predict stability and other related properties.

Methods

Benchmark data

We applied a few filters to the data sets to help ensure the coherence on experimental conditions and types of folding. From the curated dataset (*Maxwell et al., 2005*) we collected 30 protein ΔG values and then removed proteins that form obligate homodimers. We also removed the protein VlsE, as it has been shown that despite being a two-state folder protein, it does not follow the usual contact order relationship (*Jones and Wittung-Stafshede, 2003*), reducing the total number of proteins to 26. We note for completeness that the predicted stability of VlsE (18 kcal/mol) is substantially greater than experiment (5.7 kcal/mol), suggesting again that our approach is limited for large proteins.

For the proteins from the previously described 'megascale' experiment (*Tsuboyama et al., 2023*), which included 478 unique proteins with estimated ΔG, we kept only natural proteins, reducing the size of the data set to 239 proteins. Where a protein ΔG was measured in multiple replicates, we used the average of the replicates as the final value.

323 ## Generation of the input structure

324 As ESM-IF was trained using structures predicted using AlphaFold 2 (AF2) (*Jumper et al., 2021*)), we
325 use LocalColabFold v.1.5.0 (AlphaFold 2.3.1) to generate a structural model for each target fold. We
326 use LocalColabFold with MMseqs2 (*Steinegger and Söding, 2017*) and default parameters includ-
327 ing three recycles, no relaxation of the predicted structure, and the maximum multiple sequence
328 alignment size to generate an accurate fold. If an experimentally-derived structure is available
329 for a target protein and conformation, we use it as a template for ColabFold to minimise possible
330 structural deviations introduced by AF2. If the prediction by AF2 differed substantially from this
331 template structure, we performed the prediction again while reducing the MSA size, and thus the
332 influence of the co-evolutionary signal, until the predicted structure results similar to the input
333 structure template (with an RMSD less than 0.5 Å). For the calculations T4 lysozyme we used PDB
334 IDs 3DMV and 2LCB (*Bouvignies et al., 2011*) and for Cyclophilin A we based our calculations on
335 PDB IDs 3K0N and 3K0O (*Fraser et al., 2009*).

336 ## Global stability predictions using ESM-IF

We predicted the global stability value from each structure using the Evolutionary Scale Model
'Inverse Folding' model (ESM-IF - model: esm_if1_gvp4_t16_142M_UR50) (*Hsu et al., 2022*)). We
used the output logits from the language-model decoder to evaluate the likelihoods for the amino
acids at a specific position given the sequence position from the N-term to the preceding position.
Specifically, we evaluated the likelihood ($\mathscr{L}_k$) for each amino acid k by applying the Softmax on the
20 amino acid logits (x) at each position:

$$\mathscr{L}_k = \text{Softmax}(x_k) = \frac{\exp(x_k)}{\sum_{j=1}^{20} \exp(x_j)})$$

337 ## Other methods for stability predictions

338 As comparison to the predictions based on ESM-IF we also predicted stabilities using other meth-
339 ods.

340 ### Absolute contact number

We evaluated the absolute contact number (ACN) of a structure using:

$$\text{ACN} = \sum_{i}^{N} \sum_{j>i}^{N} 1 \quad if \; d_{ij} < d_{\text{cutoff}}$$

341 with N is the number of residues, d_{ij} is the distance between the C_α atoms of residues i and j, and
342 $d_{\text{cutoff}} = 8$ Å.

343 ### FoldX

344 We predicted global stability with FoldX v.5 (*Schymkowitz et al., 2005*) using AF2 predicted struc-
345 tures as input. Specifically, we first used the command 'RepairPDB', followed by the command
346 'Stability' to evaluate the different energy contributions and used their sum (total-energy) as abso-
347 lute stability value. We reverted the sign of this number to represent ΔG. We stress here that FoldX
348 was not developed to predict ΔG.

349 ### Chroma

350 We predicted global stability with the Chroma model (*Ingraham et al., 2023*). Specifically, we used
351 the Github version of Chroma (Dec. 2023) and the design protocol to collect the conditional likeli-
352 hoods at diffusion time t=0.5. We then use the likelihoods for the wild-type residues and Eq. 1 to
353 evaluate protein stability.

ESM-1b

We used ESM-1B (model: esm1b_t33_650M_UR50S) (*Rives et al., 2021*)) to predict absolute stability using only the target protein sequence. For each protein, we collected the output logits of the language model decoder and then used the same approach as for ESM-IF to predict the absolute stability.

Predictions for Nank and SasG

For both Nank and SasG we first generated AF2 models of the individual repeats using the full structure. For G5^1-E-G5^2 we used PDB ID 3TIQ (*Gruszka et al., 2015*) and for Nank used PDB ID 1OT8 (*Mello and Barrick, 2004*). If a loop region was present between the domains/repeats, we included it in the N-terminal fragment. For the additive domain model we simply summed the ΔG values predicted for each domain in the construct.

Data and code availability

Code and data is available at https://github.com/KULL-Centre/_2024_cagiada_stability/

Acknowledgments

We thank Gabriel Rocklin for comments and suggestions on the manuscript. The research was supported by the PRISM (Protein Interactions and Stability in Medicine and Genomics) centre funded by the Novo Nordisk Foundation (NNF18OC0033950, to K.L.-L.). We acknowledge access to computational resources via a grant from the Carlsberg Foundation (CF21-0392).

References

Akdel M, Pires DE, Pardo EP, Jänes J, Zalevsky AO, Mészáros B, Bryant P, Good LL, Laskowski RA, Pozzati G, et al. A structural biology community assessment of AlphaFold2 applications. Nature Structural & Molecular Biology. 2022; 29(11):1056–1067.

Anfinsen CB. Principles that govern the folding of protein chains. Science. 1973 Jul; 181(4096):223–230.

Bastolla U, Demetrius L. Stability constraints and protein evolution: the role of chain length, composition and disulfide bonds. Protein Engineering Design and Selection. 2005; 18(9):405–415.

Blaabjerg LM, Kassem MM, Good LL, Jonsson N, Cagiada M, Johansson KE, Boomsma W, Stein A, Lindorff-Larsen K. Rapid protein stability prediction using deep learning representations. Elife. 2023; 12:e82593.

Bouvignies G, Vallurupalli P, Hansen DF, Correia BE, Lange O, Bah A, Vernon RM, Dahlquist FW, Baker D, Kay LE. Solution structure of a minor and transiently formed state of a T4 lysozyme mutant. Nature. 2011 Aug; 477(7362):111–114.

Chen T, Gong C. HotProtein: A novel framework for protein thermostability prediction and editing. NeurIPS 2022. 2022; .

Chu KS, Siegel JB. Protein Stability Prediction by Fine-tuning a Protein Language Model on a Mega-scale Dataset. bioRxiv. 2023; p. 2023–11.

De Sancho D, Doshi U, Munoz V. Protein folding rates and stability: how much is there beyond size? Journal of the American Chemical Society. 2009; 131(6):2074–2075.

Dieckhaus H, Brocidiacono M, Randolph NZ, Kuhlman B. Transfer learning to leverage larger datasets for improved prediction of protein stability changes. Proceedings of the National Academy of Sciences. 2024; 121(6):e2314853121.

Eisenmesser EZ, Millet O, Labeikovsky W, Korzhnev DM, Wolf-Watz M, Bosco DA, Skalicky JJ, Kay LE, Kern D. Intrinsic dynamics of an enzyme underlies catalysis. Nature. 2005 Nov; 438(7064):117–121.

Fersht A. Structure and Mechanism in Protein Science: A Guide to Enzyme Catalysis and Protein Folding. World Scientific; 2017.

397 **Fraser JS**, Clarkson MW, Degnan SC, Erion R, Kern D, Alber T. Hidden alternative structures of proline isomerase
398 essential for catalysis. Nature. 2009; 462(7273):669–673.

399 **Ghosh K**, Dill KA. Computing protein stabilities from their chain lengths. Proc Natl Acad Sci U S A. 2009 Jun;
400 106(26):10649–10654.

401 **Gruszka DT**, Whelan F, Farrance OE, Fung HKH, Paci E, Jeffries CM, Svergun DI, Baldock C, Baumann CG, Brock-
402 well DJ, Potts JR, Clarke J. Cooperative folding of intrinsically disordered domains drives assembly of a strong
403 elongated protein. Nat Commun. 2015 Jun; 6:7271.

404 **Gruszka DT**, Wojdyla JA, Bingham RJ, Turkenburg JP, Manfield IW, Steward A, Leech AP, Geoghegan JA, Foster
405 TJ, Clarke J, Potts JR. Staphylococcal biofilm-forming protein has a contiguous rod-like structure. Proc Natl
406 Acad Sci U S A. 2012 Apr; 109(17):E1011–8.

407 **Hsu C**, Verkuil R, Liu J, Lin Z, Hie B, Sercu T, Lerer A, Rives A. Learning inverse folding from millions of predicted
408 structures. In: *International conference on machine learning* PMLR; 2022. p. 8946–8970.

409 **Ingraham JB**, Baranov M, Costello Z, Barber KW, Wang W, Ismail A, Frappier V, Lord DM, Ng-Thow-Hing C,
410 Van Vlack ER, Tie S, Xue V, Cowles SC, Leung A, Rodrigues JV, Morales-Perez CL, Ayoub AM, Green R, Puentes
411 K, Oplinger F, et al. Illuminating protein space with a programmable generative model. Nature. 2023 Nov;
412 623(7989):1070–1078.

413 **Jones K**, Wittung-Stafshede P. The largest protein observed to fold by two-state kinetic mechanism does not
414 obey contact-order correlation. J Am Chem Soc. 2003 Aug; 125(32):9606–9607.

415 **Jumper J**, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A,
416 Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J,
417 Back T, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug; 596(7873):583–
418 589.

419 **Lindorff-Larsen K**, Teilum K. Linking thermodynamics and measurements of protein stability. Protein Engi-
420 neering, Design and Selection. 2021; 34:gzab002.

421 **Liu YM**, Feng S, Ding XL, Kang CF, Yan YB. Mutation of the conserved Asp122 in the linker impedes creatine
422 kinase reactivation and refolding. Int J Biol Macromol. 2009 Apr; 44(3):271–277.

423 **Maxwell KL**, Wildes D, Zarrine-Afsar A, De Los Rios MA, Brown AG, Friel CT, Hedberg L, Horng JC, Bona D, Miller
424 EJ, Vallée-Bélisle A, Main ERG, Bemporad F, Qiu L, Teilum K, Vu ND, Edwards AM, Ruczinski I, Poulsen FM,
425 Kragelund BB, et al. Protein folding: defining a "standard" set of experimental conditions and a preliminary
426 kinetic data set of two-state proteins. Protein Sci. 2005 Mar; 14(3):602–616.

427 **Mello CC**, Barrick D. An experimentally determined protein folding energy landscape. Proc Natl Acad Sci U S A.
428 2004 Sep; 101(39):14102–14107.

429 **Mittermaier A**, Kay LE. New tools provide new insights in NMR studies of protein dynamics. science. 2006;
430 312(5771):224–228.

431 **Mulder FA**, Mittermaier A, Hon B, Dahlquist FW, Kay LE. Studying excited states of proteins by NMR spec-
432 troscopy. Nature structural biology. 2001; 8(11):932–935.

433 **Nikam R**, Kulandaisamy A, Harini K, Sharma D, Gromiha MM. ProThermDB: thermodynamic database for
434 proteins and mutants revisited after 15 years. Nucleic Acids Res. 2021 Jan; 49(D1):D420–D424.

435 **Nisthal A**, Wang CY, Ary ML, Mayo SL. Protein stability engineering insights revealed by domain-wide compre-
436 hensive mutagenesis. Proceedings of the National Academy of Sciences. 2019; 116(33):16367–16377.

437 **Notin P**, Kollasch A, Ritter D, Van Niekerk L, Paul S, Spinner H, Rollins N, Shaw A, Orenbuch R, Weitzman R, Frazer
438 J, Dias M, Franceschi D, Orenbuch R, Gal Y, Marks DS. ProteinGym: large-scale benchmarks for protein fitness
439 prediction and design. Advances in Neural Information Processing Systems. 2024; 36.

440 **Pancotti C**, Benevenuta S, Birolo G, Alberini V, Repetto V, Sanavia T, Capriotti E, Fariselli P. Predicting protein
441 stability changes upon single-point mutation: a thorough comparison of the available tools on a new dataset.
442 Brief Bioinform. 2022 Mar; 23(2).

443 **Papaleo E**, Sutto L, Gervasio FL, Lindorff-Larsen K. Conformational changes and free energies in a proline
444 isomerase. Journal of Chemical Theory and Computation. 2014; 10(9):4169–4174.

445 **Park C**, Marqusee S. Pulse proteolysis: a simple method for quantitative determination of protein stability and
446   ligand binding. Nature methods. 2005; 2(3):207–212.

447 **Pucci F**, Rooman M. Stability curve prediction of homologous proteins using temperature-dependent statistical
448   potentials. PLoS computational biology. 2014; 10(7):e1003689.

449 **Pucci F**, Schwersensky M, Rooman M. Artificial intelligence challenges for predicting the impact of mutations
450   on protein stability. Current opinion in structural biology. 2022; 72:161–168.

451 **Reeves S**, Kalyaanamoorthy S. Zero-Shot Transfer of Protein Sequence Likelihood Models to Thermostability
452   Prediction. bioRxiv. 2023; p. 2023–07.

453 **Rives A**, Meier J, Sercu T, Goyal S, Lin Z, Liu J, Guo D, Ott M, Zitnick CL, Ma J, Fergus R. Biological structure and
454   function emerge from scaling unsupervised learning to 250 million protein sequences. Proc Natl Acad Sci U
455   S A. 2021 Apr; 118(15).

456 **Robertson AD**, Murphy KP. Protein structure and the energetics of protein stability. Chemical reviews. 1997;
457   97(5):1251–1268.

458 **Ruiz-Blanco YB**, Marrero-Ponce Y, Paz W, García Y, Salgado J. Global stability of protein folding from an empir-
459   ical free energy function. J Theor Biol. 2013 Mar; 321:44–53.

460 **Santos J**, Risso VA, Sica MP, Ermácora MR. Effects of serine-to-cysteine mutations on beta-lactamase folding.
461   Biophys J. 2007 Sep; 93(5):1707–1718.

462 **Schymkowitz J**, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. The FoldX web server: an online force field.
463   Nucleic Acids Res. 2005 Jul; 33(Web Server issue):W382–8.

464 **Steinegger M**, Söding J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive
465   data sets. Nat Biotechnol. 2017 Nov; 35(11):1026–1028.

466 **Thompson MC**, Barad BA, Wolff AM, Sun Cho H, Schotte F, Schwarz DM, Anfinrud P, Fraser JS. Temperature-
467   jump solution X-ray scattering reveals distinct motions in a dynamic enzyme. Nature chemistry. 2019;
468   11(11):1058–1066.

469 **Tsuboyama K**, Dauparas J, Chen J, Laine E, Mohseni Behbahani Y, Weinstein JJ, Mangan NM, Ovchinnikov S,
470   Rocklin GJ. Mega-scale experimental analysis of protein folding stability in biology and design. Nature. 2023
471   Jul; p. 1–11.

472 **Vaswani A**. Attention is all you need. Advances in Neural Information Processing Systems. 2017; .

473 **Wapeesittipan P**, Mey AS, Walkinshaw MD, Michel J. Allosteric effects in cyclophilin mutants may be explained
474   by changes in nano-microsecond time scale motions. Communications Chemistry. 2019; 2(1):41.

475 **Xavier JS**, Nguyen TB, Karmarkar M, Portelli S, Rezende PM, Velloso JP, Ascher DB, Pires DE. ThermoMutDB: a
476   thermodynamic database for missense mutations. Nucleic acids research. 2021; 49(D1):D475–D479.

477 **Yang Y**, Zhao J, Zeng L, Vihinen M. ProTstab2 for Prediction of Protein Thermal Stabilities. Int J Mol Sci. 2022
478   Sep; 23(18).

480 **Supplementary Figures**

Figure S1. Properties of the benchmark data set. Distribution of (A) ΔG_{exp} and (B) number of residues for the 265 proteins in the benchmark data set.

Figure S2. Comparison between absolute protein stability and summed entropy. We show the correlation between the ΔG_{exp} (x-axis) values for the 265 proteins in our benchmark dataset and the stability prediction (y-axis) obtained through the sum of positional entropy from ESM-IF amino acid likelihoods. The reported standard error was obtained by bootstrapping.

Figure S3. Statistical analysis of the quality of the fit. (A) Residual values for each point during the fit used to convert the ESM-iF score scale into kcal/mol. The residual values are reported on the y-axis for each fitted ΔG_pred. (B) QQPlot (quantile–quantile plot)) for the fitted residuals, suggesting that the errors are roughly normally distributed.

Figure S4. Results of a leave-one-out cross validation of the ΔG_{pred} predictions. For each of the 265 predictions we fit the conversion from summed likelihoods (Eq. 1) to physical units by fitting on 264 proteins and predicting the ΔG_{pred} for the left out protein. (A) The distribution of $\Delta G_{pred} - \Delta G_{exp}$ values calculated in this way. (B) Box plot of the absolute difference between the experimental and predicted values. The box plot shows the median as the orange line crossing the box, the lower quartile (Q1, 25%) and the upper quartile (Q3, 75%) as the edges of the box. The lines outside the box represent the one and a half interquantile range (IQR) and the wiskers represent the data points outside the IQR range.

Figure S5. Predicting absolute protein stability using alternative models. We show a comparison between ΔG_{exp} values for 265 proteins from two different data sets (represented by different colours). (A) Using likelihoods ($\mathscr{L}$) from ESM-1b together with Eq. 1. (B) Total energies from FoldX. (C) Using likelihoods from Chroma together with Eq. 1. For comparison we show (D) Using likelihoods from ESM-IF together with Eq. 1. Panel D is similar to Fig. 1 (without converting to physical units) and is shown here for comparison The Spearman's and Pearson's correlation coefficient are shown in each panel (ρ_s, ρ_p). The reported standard errors were obtained by bootstrapping.

Figure S6. Stability prediction for non-two-state folding proteins. We compared the predicted ΔG (green) with the experimental free energy differences going from the native to intermediate states (yellow) and from the intermediate state to the unfolded state (red). Results are shown for (left) creatine kinase and (right) the exo-small β-lactamase.