Revealing cancer driver genes through integrative transcriptomic and epigenomic analyses with Moonlight

Mona Nourbakhsh[1,2], Yuanning Zheng[3], Humaira Noor[3], Hongjin Chen[1], Subhayan Akhuli[1], Matteo Tiberti[2], Olivier Gevaert[3,4], Elena Papaleo*[1,2]

[1]Cancer Systems Biology, Section for Bioinformatics, Department of Health Technology, Technical University of Denmark, Lyngby, Denmark
[2]Cancer Structural Biology, Danish Cancer Institute, Copenhagen, Denmark
[3]Stanford Center for Biomedical Informatics Research, Department of Medicine, Stanford University, Stanford, CA, USA
[4]Department of Biomedical Data Science, Stanford University, Stanford, CA, USA

*Corresponding author
E-mail: elpap@dtu.dk, elenap@cancer.dk (EP)

Short title
Methylation-driven discovery of cancer drivers

Keywords
Cancer, driver, gene expression, methylation, oncogene, tumor suppressor

Abstract

Cancer involves dynamic changes caused by (epi)genetic alterations such as mutations or abnormal DNA methylation patterns which occur in cancer driver genes. These driver genes are divided into oncogenes and tumor suppressors depending on their function and mechanism of action. Discovering driver genes in different cancer (sub)types is important not only for increasing current understanding of carcinogenesis but also from prognostic and therapeutic perspectives. We have previously developed a framework called Moonlight which uses a systems biology multi-omics approach for prediction of driver genes. Here, we present an important development in Moonlight2 by incorporating a DNA methylation layer which provides epigenetic evidence for de-regulated expression profiles of driver genes. To this end, we present a novel functionality called Gene Methylation Analysis (GMA) which investigates abnormal DNA methylation patterns to predict driver genes. This is achieved by integrating the tool EpiMix which is designed to detect such aberrant DNA methylation patterns in a cohort of patients and further couples these patterns with gene expression changes. To showcase GMA, we applied it to three cancer (sub)types (basal-like breast cancer, lung adenocarcinoma, and thyroid carcinoma) where we discovered 33, 190, and 263 epigenetically driven genes, respectively. A subset of these driver genes had prognostic effects with expression levels significantly affecting survival of the patients. Moreover, a subset of the driver genes demonstrated therapeutic potential as drug targets. This study provides a framework for exploring the driving forces behind cancer and provides novel insights into the landscape of three cancer sub(types) by integrating gene expression and methylation data. Moonlight2R is available on GitHub (https://github.com/ELELAB/Moonlight2R) and BioCondcutor (https://bioconductor.org/packages/release/bioc/html/Moonlight2R.html). The associated case studies

52 presented here are available on GitHub (https://github.com/ELELAB/Moon-
53 light2_GMA_case_studies) and OSF (https://osf.io/j4n8q/).

Author summary

Cancer is a complex disease and a main cause of mortality worldwide. This heteroge-
neous disease arises due to accumulation of changes which occur in driver genes that
drive cancer progression when they are altered. These driver genes are commonly
divided into oncogenes, which promote cancer, and tumor suppressors, which prevent
it. A major goal of cancer research is identifying these driver genes, crucial for increas-
ing our current understanding of cancer biology and for developing novel treatment
approaches. A large number of cancer driver genes have already been identified.
However, the underlying mechanisms for the alterations in these genes is challenging
to predict given their context-dependent behavior and the complexity of cancer. Such
explanations are the focus of this study with the aim of providing evidence of why
certain genes do not function normally in cancer. Within this context, we present new
functionalities to our previously developed cancer driver predictive framework, Moon-
light. These new functionalities integrate multiple data types to predict oncogenes and
tumor suppressors in a systems-biology-oriented manner that is freely available as a
R package for the community.

Introduction

Cancer is a complex and heterogeneous disease and a leading cause of death globally
[1]. This widespread disease is categorized into multiple (sub)types and is character-
ized by stepwise accumulation of (epi)genetic alterations in cancer driver genes [2].

Driver genes are classified according to their function, i.e. oncogenes (OCGs) acti-

vated by gain-of-function mechanisms and tumor suppressor genes (TSGs) inacti-

vated by loss-of-function mechanisms [3]. Recently, dual role genes also emerged

which show context-dependent behavior and can act as both OCGs and TSGs in dif-

ferent biological contexts [4,5]. Driver genes participate in several cellular pathways

conceptualized in the Hallmarks of Cancer, a collection of functional capabilities that

cells gain during their transition from normal to tumor cells [6–8]. Distinct driver genes

can initiate cancer development in different cancer types and even within subtypes of

cancers originating from the same tissue. Thus, context-specific discovery of driver

genes in light of the cancer hallmarks is essential. Numerous tools have been devel-

oped for prediction of driver genes based on varying computational methods which we

recently reviewed [9]. Prediction of driver genes is essential for increasing current

knowledge of cancer development and for analyzing and interpreting the vast amount

of data in relation to the cancer phenotypes. This knowledge can be a step towards

reversing these phenotypes, discovering novel drug targets, facilitating new treatment

strategies, and designing precision medicine strategies [10–13]. We have contributed

to this field with Moonlight which uses a multi-omics systems biology approach for

prediction of driver genes [14,15].

The accumulated (epi)genetic alterations in driver genes include mutations, copy num-

ber variations, aberrant methylation levels, and histone modifications [3,16]. While ab-

normal methylation patterns are recognized as cancer-causing mechanisms, they

have been described to a lesser extent compared to mutations [9]. Hypomethylation

and hypermethylation, respectively representing loss and gain of methylation com-

pared to normal conditions, have been described as activating and inactivating mech-

anisms of OCGs and TSGs, respectively [17–19]. For instance, Søes et al. found pro-

moter hypomethylation and increased expression of putative OCG *ELMO3* to be as-

sociated with development of non-small cell lung cancer [20].

Here, we present novel functionalities to Moonlight2, expanding upon features pre-

sented in our previous work [15]. Specifically, we incorporate methylation evidence to

Moonlight2 predicted driver genes as a source of epigenetic explanation of the dereg-

ulated expression of these genes. Information about methylation state is provided by

EpiMix, an integrative tool for detecting aberrant DNA methylation patterns connected

with expression changes in patient cohorts [21]. To showcase this new feature, we

apply it to three cancer (sub)types (basal-like breast cancer, lung adenocarcinoma,

and thyroid carcinoma) and discover driver genes in the context of cell proliferation

and apoptosis, two well-established cancer hallmarks, and explore the prognostic and

therapeutic potentials of the predicted driver genes. We apply our new method on data

from The Cancer Genome Atlas (TCGA) [22,23].

115 Design and implementation

116 Design and implementation of new functionalities in Moon-
117 light

Here, we present new functionalities to Moonlight, our framework for driver gene pre-

diction [14,15]. In brief, Moonlight requires a set of differentially expressed genes

(DEGs) as input and is built up on two layers: a primary layer discovering putative

driver genes, termed oncogenic mediators, that uses gene expression changes and

information about cancer-related biological processes; and a secondary layer that cou-

ples mechanistic evidence to the oncogenic mediators by investigating (epi)genetic

alterations in the oncogenic mediators (namely, mechanistic indicators). From this sec-

ondary layer, the critical driver genes are predicted among the oncogenic mediators.

We recently presented Moonlight2 with the overall goal of implementing new function-

alities to provide standardized and automatized solutions to the analysis of the mech-

anistic indicators. At first, we developed a secondary layer for mechanistic indicators

based on mutational data [15].

In this contribution, we tackled the challenge of adding functionalities to Moonlight2 to

cover mechanistic indicators related to methylation changes. Each layer can be ap-

plied together or separately depending on the source of -omics data available for the

samples under analysis. This new functionality is termed Gene Methylation Analysis

(GMA) and should be applied following the Pattern Recognition Analysis (PRA) func-

tion which predicts the oncogenic mediators in the primary layer (Fig 1A). The biolog-

ical foundation for GMA lies within the observed roles of DNA methylation in both

physiological and cancer states. Under healthy conditions, DNA methylation serves

an essential regulatory role in cells by regulating expression of genes [24]. However,

in cancer, DNA methylation processes are altered, where hypo- and hypermethylation

can activate and inactivate OCGs and TSGs, respectively, leading to overexpression

of OCGs and silencing of TSGs [17–19] (Fig 1B).

GMA predicts methylation-driven driver genes by using EpiMix [21]. EpiMix models

DNA methylation in patient cohorts and predicts differential methylation associated

with gene expression and further allows for DNA methylation analysis of non-coding

regulatory regions [21], therefore being perfectly suitable to integrate with Moonlight's

primary layer. Moreover, EpiMix is available as a R BioConductor package, which al-

lows for easy integration with Moonlight. A key result of EpiMix is a table which in-

cludes functional CpG-gene pairs containing differentially methylated CpG sites whose

DNA methylation state is associated with the expression of the corresponding genes

they map to. Moreover, the methylation state (e.g. hypo- or hypermethylated) of each

CpG site is reported. This table is integrated with the main output table from Moon-

light's primary layer, specifically the output from PRA, which provides a list of onco-

genic mediators and their putative driver role (e.g. putative TSG or OCG) (Fig 1C).

This integration step involves the following: for each oncogenic mediator, the number

of associated CpG sites is summarized. EpiMix's predictions of methylation state and

Moonlight's predictions of driver gene role are then compared and used to assess

whether the gene's methylation status supports the putative role (OCG or TSG) of the

oncogenic mediator. These comparisons are subsequently used to define the driver

genes (Fig 1D). Those oncogenic mediators with correspondence between methyla-

tion state and putative driver role from EpiMix and Moonlight's primary layer are re-

tained as the final set of driver genes. See S1 Text for a detailed description of this

comparison.

As input, GMA requires i) a gene expression matrix with genes in rows and tumor and

normal samples in columns, ii) a methylation matrix with CpG sites in rows and tumor

and normal samples in columns which should be the same samples as in the expres-

sion data, iii) output of PRA from Moonlight's primary layer, i.e. the predicted onco-

genic mediators and their putative driver role, and finally, iv) output of a differential

expression analysis (DEA) which includes information about the DEGs. In return, GMA

outputs the following: i) a list of predicted driver genes categorized into TSGs and

OCGs, ii) a summary of the oncogenic mediators which includes the number of asso-

ciated CpG sites and evidence label, iii) a summary of various annotations found to all

DEGs input to Moonlight on the gene and methylation level, and iv) raw EpiMix results

corresponding to applying EpiMix on the input data independent of the GMA function.

We have also created three functions for visualizing genes and methylation states:

plotGMA which visualizes the number of differentially methylated hypo-, hyper- or

dual-methylated CpG sites, plotMoonlightMet which visualizes the effect of genes on

biological processes estimated in Moonlight's primary layer, and plotMetExp which

calls a visualization function from EpiMix, EpiMix_PlotModel, to display gene expres-

sion and methylation levels of a specific gene and CpG site [21].

181
Fig 1. Overview of the Moonlight framework with new methylation functionality. (A) Moonlight
consists of a primary layer requiring differentially expressed genes and gene expression data as input.
The primary layer predicts oncogenic mediators through a series of functions called functional enrich-
ment analysis (FEA), gene regulatory network analysis (GRN), upstream regulator analysis (URA), and
pattern recognition analysis (PRA). Moonlight's secondary mutation layer requires mutation data as
input and is carried out via the driver mutation analysis (DMA) function and similarly, Moonlight's sec-
ondary methylation layer implemented in the gene methylation analysis (GMA) function requires meth-
ylation data as input. The secondary layer results in the final prediction of driver genes. **(B)** DNA meth-
ylation is a mechanism occurring under physiological conditions in cells which functions to regulate
gene expression. However, in cancer, the DNA methylation process is altered. A loss of methylation
called hypomethylation can occur which can lead to increased expression of a gene and thus an in-
creased amount of the resulting protein. In contrast, gain of methylation called hypermethylation can
also occur which can silence gene expression and lead to decreased protein expression. These two

mechanisms can finally lead to cancer. Hypo- and hypermethylation can activate and inactivate onco-
genes and tumor suppressors, respectively, the biological principle that GMA is built on. **(C)** The outputs
of EpiMix and Moonlight are integrated to predict driver genes. EpiMix outputs a table of CpG-gene
pairs containing differentially methylated CpG sites whose DNA methylation state is associated with
gene expression. Moonlight outputs a list of oncogenic mediators and their putative driver role as tumor
suppressors or oncogenes. **(D)** Driver genes are defined in GMA by comparing EpiMix's predictions of
methylation state and Moonlight's predictions of driver role in "evidence" categories. Those oncogenic
mediators labeled with an "agreement" evidence are retained as the final set of predicted driver genes.
203
204

205 ## Application of new functionality to three cancer (sub)types
206
Following implementation of the new functionality, GMA, in Moonlight2, we conducted

a case study applying GMA to basal-like breast cancer, lung adenocarcinoma, and

thyroid carcinoma data from TCGA to discover methylation-driven driver genes. More-

over, we compared these predicted drivers with mutation-driven drivers by applying

our previously developed secondary mutational layer called Driver Mutation Analysis

(DMA) [15]. Detailed methods behind this case study are included in S2 Text.

Results

Case study: Prediction of driver genes with differential

methylation in three different cancer types using Moon-

light2

To showcase the new functionality in Moonlight2 and predict driver genes driven by

methylation changes, we applied Moonlight2 on three cancer (sub)types: basal-like

breast cancer, lung adenocarcinoma, and thyroid carcinoma. First, we performed DEA

between each of these cancer tissues and corresponding normal samples as this is

the input to Moonlight's primary layer (Table 1). Following DEA, Moonlight's primary

layer predicted 159, 1228, and 1598 oncogenic mediators in these three cancer

(sub)types, respectively (Table 1). Additionally, EpiMix alone identified 9483, 10018,

and 6142 functional gene-CpG pairs in these three cancer (sub)types, respectively.

These functional gene-CpG pairs represent differentially methylated CpG sites whose

DNA methylation state is associated with the expression of the corresponding genes

they map to. The number of hits discovered individually from EpiMix and Moonlight's

primary layer indicate a substantial volume of significant associations. Consequently,

integrating the results from EpiMix with the oncogenic mediators identified in Moon-

light's primary layer, as implemented in GMA presented here, help to narrow down the

most critical findings and yield the benefits of both approaches. From GMA, we found

that those oncogenic mediators in basal-like breast cancer that are associated with

differentially methylated CpGs include 38 hypomethylated CpGs, 165 hypermethyl-

ated CpGs, and 22 methylated CpGs with a dual status, meaning the CpG site was

found hypomethylated in cancer tissues from some patients, while hypermethylated in

other patients. Similarly, oncogenic mediators in lung adenocarcinoma that are asso-

ciated with differentially methylated CpGs include in total 218 hypomethylated CpGs,

625 hypermethylated CpGs, and 48 dual-methylated CpGs. Finally, oncogenic medi-

ators in thyroid carcinoma associated with differentially methylated CpGs contain in

total 945 hypomethylated CpGs, 305 hypermethylated CpGs, and 230 dual-methyl-

ated CpGs (Fig 2A).

Across all three cancer (sub)types, the number of differentially methylated CpG sites

mapped to the oncogenic mediators ranges between 0 and 28. The classifications of

methylation status in the oncogenic mediators in basal-like breast cancer are shown

in Fig 2B, generated with the plotGMA function. Next, we compared Moonlight's onco-

genic mediators with EpiMix' functional genes. For this, we included only those func-

tional genes that contained the same methylation state in all of its associated CpGs

and moreover, the dual states were excluded. In basal-like breast cancer, this com-

parison revealed 109 oncogenic mediators not associated with differentially methyl-

ated CpGs, 2754 functional genes not predicted as oncogenic mediators, 17 onco-

genic mediators with a "conflicting" evidence label, and 33 oncogenic mediators with

an "agreement" evidence label (Fig 2C). Consequently, these 33 oncogenic mediators

are retained as the final set of driver genes divided into 32 TSGs and 1 OCG (Table

1). Next, we visualized the effect of these predicted driver genes in basal-like breast

cancer on two well-known cancer hallmarks, apoptosis and proliferation of cells, using

the function plotMoonlightMet. These effects define the basis upon which the onco-

genic mediators are predicted from the PRA step in Moonlight's primary layer, demon-

strating that the predicted OCGs have a positive effect on proliferation of cells and a

negative effect on apoptosis and vice versa for the predicted TSGs (Fig 2D). Similar

overviews for lung adenocarcinoma and thyroid carcinoma are shown in S1 Fig, which

resulted in a final prediction of 190 driver genes divided into 110 TSGs and 80 OCGs

in lung adenocarcinoma and 263 driver genes categorized into 5 TSGs and 258 OCGs

in thyroid carcinoma (Table 1). We did not discover any dual role genes across the

three cancer (sub)types, i.e. genes predicted as OCGs in one of the three cancer

(sub)types and as TSGs in another cancer (sub)type and vice versa.

Table 1. Number of predicted DEGs, oncogenic mediators, and driver genes in three cancer (sub)types: basal-like breast cancer, lung adenocarcinoma, and thyroid carcinoma. The oncogenic mediators and driver genes predicted by Moonlight's primary and secondary methylation layer, respectively, are divided into (putative) TSGs and OCGs.

Cancer (sub)type	DEGs	Oncogenic media-tors [putative TSGs/putative OCGs]	Driver genes [TSGs/OCGs]
Basal-like breast cancer	4292	159 [125/34]	33 [32/1]

Lung adenocarci-noma	4468	1228 [521/707]	190 [110/80]
Thyroid carcinoma	2972	1598 [118/1480]	263 [5/258]

Abbreviations: DEGs, differentially expressed genes; OCGs, oncogenes; TSGs, tumor suppressor genes.

We then compared the predicted driver genes with the predicted oncogenic mediators

in each cancer (sub)type. We quantified these comparisons in terms of overlaps with

genes reported in the COSMIC Cancer Gene Census (CGC) [25]. Specifically, we

computed the precision as (TP/(TP+FP))*100 and sensitivity as (TP/(TP+FN))*100.

We defined the true positives (TP) as the overlap between the gene set (either the

driver genes or the oncogenic mediators) and the CGC, whereas the false positives

(FP) are those genes found in the gene set but are not included in CGC. In contrast,

the false negatives (FN) comprise those genes reported in CGC but are not predicted

in our gene set. For all three cancer (sub)types, we found that GMA had a greater

precision and lower sensitivity compared to using only Moonlight's primary layer (Fig

2E). A higher precision of GMA is desirable as it indicates that the predicted driver

gene sets have a higher fraction of genes from the CGC compared to the oncogenic

mediator sets. On the other hand, the higher sensitivity of using only Moonlight's pri-

mary layer compared to also using GMA might be attributed to the larger numbers of

oncogenic mediators. A larger number of oncogenic mediators results in a larger over-

lap between the CGC and the oncogenic mediators, thereby lowering the number of

FNs and increasing the sensitivity. In this case, prioritizing higher precision over sen-

sitivity is preferable since our aim is to find the most crucial driver genes among the

oncogenic mediators. Thus, a higher precision indicates a greater proportion of TPs,

corresponding with our objective. Next, we also evaluated the significance of associa-

tion between the gene sets and the CGC using a Fisher's exact test (Table 2). We only

found the oncogenic mediator and driver gene sets from basal-like breast cancer to

have a significant association with genes in the CGC (p-value = 0.000392 for the on-

cogenic mediators predicted using Moonlight's primary layer and p-value = 0.00228

for the driver genes predicted using GMA). However, in all three cancer (sub)types,

we found the driver genes to have a higher odds ratio than the oncogenic mediators,

demonstrating a greater association between the driver gene sets and the CGC com-

pared to the oncogenic mediators (Table 2).

297
298

Table 2. Significance of association between Moonlight's gene sets and genes from the Cancer Gene Census (CGC) evaluated using Fisher's exact test in three cancer (sub)types: basal-like breast cancer, lung adenocarcinoma, and thyroid carcinoma. The gene sets from Moonlight were found using Moonlight's primary layer and Moonlight's secondary layer through the Gene Methylation Analysis (GMA) functionality. p-values and odds ratios from Fisher's exact test are included.

Cancer (sub)type	Method	*p*-value	Odds ratio
Basal-like breast cancer	Moonlight's primary layer	0.000392*	2.77
	GMA	0.00228*	5.09
Lung adenocarci-noma	Moonlight's primary layer	0.0764	1.28
	GMA	0.272	1.40
Thyroid carcinoma	Moonlight's primary layer	0.472	0.895
	GMA	0.215	1.39

*p-value < 0.05.
Abbreviations: CGC, Cancer Gene Census; GMA, Gene Methylation Analysis.

299
300
While these results together demonstrate the added value of GMA, it is worth high-

lighting certain limitations. Notably, the driver genes reported in CGC are mainly based

on mutation evidence. In this study, we have used abnormal DNA methylation levels

as evidence for deregulated expression of the driver genes. Hence, these methylation

patterns may not be fully captured in the CGC, challenging our comparison with the

CGC. However, to date, no golden standard of cancer drivers exists, and the CGC

stands as the most robust and comprehensive resource available. Thus, it serves as

the main reference point that the majority of studies use to evaluate their predicted

driver genes and method [26–37]. To our knowledge, a similar well-curated resource

of cancer driver genes driven by methylation changes does not exist. Moreover, per-

forming cancer type-specific comparisons would be more desirable. While the CGC

reports which cancer types the driver genes are associated with, these annotations

are limited in scope. Therefore, while ideal, performing such cancer type-specific com-

parisons do not contain enough power. Finally, the quantitative statistical measures

are not taking into account that some of our predicted driver genes may be novel.

Consequently, some FPs may in fact be TPs but are not included in CGC, and some

FNs may not necessarily be FNs; rather, they may not represent drivers in the specific

cancer (sub)type.

Fig 2. Integration of Moonlight and EpiMix for prediction of cancer driver genes. (A) Number of differentially methylated CpGs as found from EpiMix in oncogenic mediators predicted from Moonlight's primary layer. The differentially methylated CpGs are categorized into methylation status and stratified by cancer (sub)type. **(B)** Heatmap showing number of differentially methylated CpGs and classifications of methylation status in the oncogenic mediators in basal-like breast cancer. The heatmap was generated using the plotGMA function. **(C)** Venn diagram comparing oncogenic mediators predicted from Moonlight's primary layer with functional genes predicted from EpiMix in basal-like breast cancer. The functional genes are genes containing differentially methylated CpG pairs whose DNA methylation state is associated with expression of the gene. Only those functional genes that contained the same methylation state in all of its associated CpGs were included in this comparison, and moreover, the dual methylation states were excluded. **(D)** Heatmap showing the effect of the predicted driver genes in basal-like breast cancer on apoptosis and proliferation of cells. This heatmap was generated using the function plotMoonlightMet. These effects define the basis upon which the oncogenic mediators are predicted from the PRA step in Moonlight's primary layer. **(E)** Comparison between the predicted driver genes with the predicted oncogenic mediators in all three cancer (sub)types where the driver genes

were predicted with the new functionality GMA in Moonlight's secondary layer, and the oncogenic me-
diators were predicted with Moonlight's primary layer. The comparisons were quantified in terms of
overlaps with genes reported in the COSMIC Cancer Gene Census (CGC) by computing the precision
and sensitivity. The precision was calculated as $(TP/(TP+FP))*100$ and sensitivity as
$(TP/(TP+FN))*100$. The true positives (TP) are the overlap between the gene set (either the driver genes
or the oncogenic mediators) and the CGC. The false positives (FP) are those genes found in the gene
set but are not included in CGC. The false negatives (FN) comprise those genes reported in CGC but
are not predicted in our gene set.

344
345

To investigate biological roles of the predicted driver genes, we performed enrichment

analyses (Fig 3). The predicted driver genes are involved in various signaling path-

ways such as KRAS signaling in basal-like breast cancer and thyroid carcinoma,

mTORC1 signaling in lung adenocarcinoma, and TNF−alpha signaling via NF−kB and

p53 pathway in thyroid carcinoma. Previously, TP53 and TNF signaling have been

associated with the onset of cancer among epigenetically modified pathways [38]. Fur-

thermore, IL−6/JAK/STAT3 signaling was significantly enriched among the predicted

driver genes in basal-like breast cancer (Fig 3). Basal-like breast cancers overexpress

Interleukin 6 (IL-6), a pro-inflammatory cytokine, and it has been reported that p53

absence triggers an IL-6 dependent epigenetic reprogramming driving breast cancer

cells towards a basal-like/stem cell-like gene expression profile [39]. Additionally, epi-

thelial-mesenchymal transition (EMT) is a recurring enriched term, observed in both

lung adenocarcinoma and thyroid carcinoma. Epigenetic regulation of EMT has previ-

ously been described, and DNA methylation and demethylation plays a key role in this

regulation [40–43].

Fig 3. Enrichment analyses of predicted driver genes. Enrichment analysis of predicted driver genes in **(A)** basal-like breast cancer, **(B)** lung adenocarcinoma, and **(C)** thyroid carcinoma using the "MSigDB Hallmark 2020" database. The top 10 most significantly enriched terms (adjusted *p*-value < 0.05) are included. The gene ratio on the *x* axis is the ratio between the number of predicted driver genes that intersect with genes annotated in the given hallmark gene set and the total number of genes annotated in the respective hallmark gene set. The point sizes reflect the number of driver genes playing a role in the respective hallmark gene set.

Association between expression of predicted driver genes and survival of cancer patients

We performed survival analysis to evaluate the prognostic potential of the predicted driver genes. We first used Cox proportional hazards regression and found that the expression level of 20 of the predicted OCGs in lung adenocarcinoma had a significant effect on survival at the multivariate level when accounting for tumor stage, age of patients, and sex of patients. Similarly, expression of two of the predicted OCGs in thyroid carcinoma had a significant effect on survival. Thus, we deemed these 22 OCGs as prognostic (Fig 4A). Next, we examined whether high or low expression of these prognostic genes were associated with survival of the patients. For this, we divided the patients into high and low expression groups and assessed differences in survival through Kaplan-Meier survival analyses and log-rank tests. These analyses

revealed a significant difference in survival between patients with high and low expres-

sion of 18 of the 20 prognostic OCGs in lung adenocarcinoma. The two OCGs that did

not show a significant difference were *RPL39L* and *GINS2.* On the other hand, we did

not observe a significant difference in survival between patients with high and low ex-

pression of the two predicted OCGs in thyroid carcinoma. These results together indi-

cate a greater prognostic potential of OCGs compared to TSGs and additionally, a

greater presence of prognostic OCGs in lung adenocarcinoma compared to basal-like

breast cancer and thyroid carcinoma. It is, however, worth mentioning that a smaller

subset of driver genes was predicted in basal-like breast cancer with only one pre-

dicted OCG, indicating a more limited search pool for prognostic OCGs.

To highlight a few examples, multivariate Cox regression analysis identified

GNPNAT1, *RRM2*, and *SLC2A1* as prognostic OCGs in lung adenocarcinoma with

hazard ratios of 1.4, 1.3, and 1.3, respectively. In all three cases, patients with high

expression of the OCG had a significantly lower survival probability compared to pa-

tients with low expression of these OCGs (Figs 4B-D) (*p*-values: < 0.0001, 0.0015 and

0.00055 for *GNPNAT1*, *RRM2* and *SLC2A1*, respectively). This aligns with the antici-

pated role of OCGs which are typically upregulated in cancer, indicating a worse prog-

nosis.

Fig 4. Survival analysis of predicted driver genes. **(A)** Hazard ratios from multivariate Cox proportional hazards regression of 20 of the predicted OCGs in lung adenocarcinoma and of two of the predicted OCGs in thyroid carcinoma. **(B-D)** Kaplan-Meier survival plots of three of the predicted OCGs in lung adenocarcinoma which were deemed prognostic from multivariate Cox regression analysis: **(B)** *GNPNAT1*, **(C)** *RRM2*, and **(D)** *SLC2A1*. Patients with expression values above and below the median expression level of the respective gene were divided into a high and low expression group, respectively. The *p*-values represent the significance of difference in survival between the two groups for each gene.

Predicted driver genes have therapeutic potential as drug targets

The potential of cancer driver genes as drug targets have previously been highlighted [44–46] and targeted therapies have been developed towards these genes. Thus, we next investigated the therapeutic potential of the predicted driver genes as drug targets by querying the Drug-Gene Interaction Database (DGIdb) [47] for driver gene-drug interactions using only cancer-specific data sources. In basal-like breast cancer, we

identified seven TSGs documented to interact with drugs in DGIdb. In lung adenocar-

cinoma, both OCGs and TSGs, numbering 12 each, were reported as drug targets.

Finally, in thyroid carcinoma, 23 OCGs were reported as interacting with drugs (S2

Fig). Across all three cancer (sub)types, the number of driver gene-drug interactions

varied between one and 55. Roughly half of all predicted driver genes interacted with

one drug while the other half interacted with two or more drugs (Fig 5A).

Next, we examined those driver gene-drug interactions for which the interaction type

was known. In basal-like breast cancer, lung adenocarcinoma, and thyroid carcinoma,

we found three (all TSGs), six (three OCGs and three TSGs), and five (all OCGs)

driver genes, respectively, for which the interaction type was known (Figs 5B-D). The

majority of the drugs were inhibitors. The two driver genes with the most interactions

were *PDGFRB* in basal-like breast cancer and *MET* in thyroid cancer. We predicted

PDGFRB as a TSG in basal-like breast cancer which is annotated to interact with 16

inhibitors and three drugs with antagonist or inhibitor interactions. These drugs exert

inhibitory mechanisms for targeting an OCG role of *PDGFRB*. As the gene-drug target

interactions are not specific for a certain cancer type, these results might suggest a

potential dual role of *PDGFRB*. On the other hand, *MET* predicted as an OCG in thy-

roid cancer interacted with 19 inhibitors, in accordance with the OCG role of *MET*.

Moreover, in lung adenocarcinoma, the predicted OCG *RRM2*, which we also identi-

fied as a prognostic gene above, interacted with one inhibitor, gemcitabine. Previously,

one study investigated the mRNA expression of RRM1 and RRM2 in tumors from pa-

tients with lung adenocarcinoma treated with docetaxel/gemcitabine. They found low

RRM2 mRNA expression to be associated with a higher response rate to treatment

compared to patients with high expression [48]. Similarly, in thyroid carcinoma, we

observed an interaction between *ERBB3*, a member of the epidermal growth factor

receptor (EGFR) family of receptor tyrosine kinases, and four inhibitors (sapitinib, pozi-

otinib, gefitinib, and dacomitinib). These inhibitors, all classified as tyrosine kinase in-

hibitors [49–56], align with *ERBB3*'s predicted role as an OCG. Another example is

the interaction between *EpCAM* and solitomab in lung adenocarcinoma. *EpCAM* is an

epithelial cell adhesion molecule which plays a role in cell proliferation, migration, and

signaling and is frequently overexpressed on the cell surface of several human carci-

nomas [57–59]. For instance, *EpCAM* was recently found to be upregulated in primary

lung cancer compared to normal lung tissues caused by gene amplification and pro-

moter hypomethylation [60]. Solitomab is a bispecific antibody binding to *EpCAM* and

CD3 [57] which previously has shown preliminary signs of antitumor activity [61].

454
Fig 5. Exploration of predicted driver genes as drug targets. (A) Distribution of driver gene-drug
interactions stratified by cancer type with the number of drug interactions on the *x* axis and number of
driver genes on the *y* axis. **(B-D)** Heatmaps visualizing driver gene-drug interactions in **(B)** lung adeno-
carcinoma, **(C)** basal-like breast cancer, and **(D)** thyroid carcinoma. Only those driver gene-drug inter-
actions where the interaction type was known are included in the heatmaps. The type of interaction is
shown in different colors. The driver genes are divided into OCGs and TSGs.

461
462

Integrating the results from DMA and GMA functions of Moonlight2

Next, we also applied the Moonlight2 DMA functionality [15] to the data used for the

case studies above to show the potential of integrating different mechanistic indicators.

For basal-like breast cancer, DMA predicted 46 driver genes (10 OCGs and 36 TSGs),

while GMA predicted 33 driver genes (32 OCGs and 1 TSG) (Figs 6A-C). For lung

adenocarcinoma, DMA predicted 842 driver genes (490 OCGs and 352 TSGs), while

GMA predicted 190 (80 OCGs and 110 TSGs) (Figs 6D-F). Both secondary layers

predicted a larger number of driver genes in lung adenocarcinoma than basal-like

breast cancer (Table 1, Fig 6). This is likely a direct consequence of Moonight's pri-

mary layer, which identified a larger number of oncogenic mediators in lung adenocar-

cinoma than basal-like breast cancer. At the same time, DMA predicted a larger num-

ber of driver genes for both datasets than GMA, with a larger proportion in lung ade-

nocarcinoma than basal-like breast cancer (~4.5 times against ~1.4 times, respec-

tively). This observation aligns with previous reports suggesting that lung adenocarci-

noma exhibits a high mutation burden [62,63], suggesting that DMA was able to iden-

tify a larger number of driver mutations overall. In most cases, we found an overlap

between driver genes identified by DMA and GMA, which suggests multiple mecha-

nisms at play. In basal-like breast cancer, 13 driver genes were predicted by both DMA

and GMA, which were all TSGs (Figs 6A-C). In lung adenocarcinoma, 141 driver genes

(63 OCGs and 78 TSGs) were identified by both methods (Figs 6D-F). In the case of

lung adenocarcinoma, and more so than in basal-like breast cancer, the driver genes

predicted by GMA were in good part also predicted by DMA.

Next, we performed enrichment analysis of the DMA predicted driver genes in basal-

like breast cancer and lung adenocarcinoma to understand whether DMA and GMA

can identify distinct or overlapping biological mechanisms. The significantly enriched

terms (adjusted *p*-value < 0.05) among the DMA predicted driver genes in basal-like

breast cancer were angiogenesis, KRAS signaling up, epithelial mesenchymal transi-

tion, and IL-2/STAT5 signaling (Fig 7A) while among the GMA predicted drivers they

were IL-6/JAK/STAT3 signaling, UV response dn, KRAS signaling up and adipogene-

sis (Fig 3A). Thus, results from both GMA and DMA were enriched in the KRAS sig-

naling term only.

Both GMA and DMA identified *NRP1* as a driver gene involved in KRAS signaling.

NRP1 has been shown to be highly expressed in different cancer types [64] and to-

gether with *FSTL1* is predicted to be driver for basal-like breast cancer by DMA. These

two genes are involved in angiogenesis, one of the cancer hallmarks [65–67], which

is also prognostic indicators of survival in breast cancer [68,69]. Additionally, for lung

adenocarcinoma, key enriched terms for both DMA and GMA predicted driver genes

included G2-M checkpoint, E2F targets and mTORC1 signaling (Figs 3B, 7B), sug-

gesting that the two mechanistic indicators identify at least partially overlapping bio-

logical processes. These processes are all important in cancer progression or metas-

tasis [70–73].

Finally, we also performed gene enrichment analysis of the driver genes identified by

both DMA and GMA. An enrichment analysis of the 141 overlapping driver genes be-

tween GMA and DMA in lung adenocarcinoma revealed E2F targets, G2-M checkpoint

and mTORC1 signaling to again be the most significant (Fig 7C), covering a vast ma-

jority of the overlapping genes. A similar enrichment analysis of the 13 overlapping

driver genes between GMA and DMA in basal-like breast cancer revealed no signifi-

cantly enriched terms.

Basal-like breast cancer

Lung adenocarcinoma

512
513 **Fig 6. Comparison of number of mutation- and methylation-driven driver genes.** Venn diagram
514 comparing **(A, D)** the number of driver genes, **(B, E)** TSGs, and **(C, F)** OCGs predicted by the Driver
515 Mutation Analysis (DMA) and Gene Methylation Analysis (GMA) functions of Moonlight2 for **(A-C)** basal-
516 like breast cancer and **(D-F)** lung adenocarcinoma.
517

Fig 7. Enrichment analysis of predicted mutation-driven driver genes. Enrichment analysis of **(A)** mutation-driven driver genes predicted by Driver Mutation Analysis (DMA) in basal-like breast cancer, **(B)** mutation-driven driver genes predicted by Driver Mutation Analysis (DMA) in lung adenocarcinoma, and **(C)** driver genes predicted by both DMA and GMA in lung adenocarcinoma. The "MSigDB Hallmark 2020" database was used for the enrichment analyses. The top 10 most significantly enriched terms (adjusted p-value < 0.05) are included. The gene ratio on the x axis is the ratio between the number of predicted driver genes that intersect with genes annotated in the given hallmark gene set and the total number of genes annotated in the respective hallmark gene set. The point sizes reflect the number of driver genes playing a role in the respective hallmark gene set.

Availability and future directions

The data that support the findings of this study are openly available in The Cancer Genome Atlas. The data used for this analysis are available at the Genomic Data Commons (https://portal.gdc.cancer.gov). GitHub and OSF repositories associated with this study are available at https://github.com/ELELAB/Moonlight2R, https://github.com/ELELAB/Moonlight2_GMA_case_studies, and https://osf.io/j4n8q/. Example data and vignette are available in the Moonlight2 R package. In the future, we envision incorporation of additional secondary -omics layers such as chromatin accessibility and copy number variation. Moreover, we would like to implement prote-omics and single-cell RNA sequencing data as additional input data types.

Acknowledgements

The results published here are in whole or in part based upon data generated by The Cancer Genome Atlas (TCGA) Research Network: https://www.cancer.gov/tcga.

Author contributions

Conceptualization: Mona Nourbakhsh, Yuanning Zheng, Matteo Tiberti, Olivier Gevaert, Elena Papaleo

Data curation: Mona Nourbakhsh, Hongjin Chen, Subhayan Akhuli, Matteo Tiberti

Formal analysis: Mona Nourbakhsh, Yuanning Zheng, Humaira Noor, Hongjin Chen, Subhayan Akhuli, Matteo Tiberti, Olivier Gevaert, Elena Papaleo

Funding acquisition: Olivier Gevaert, Elena Papaleo

Investigation: Mona Nourbakhsh, Hongjin Chen, Subhayan Akhuli, Elena Papaleo

Methodology: Mona Nourbakhsh, Yuanning Zheng, Humaira Noor, Matteo Tiberti, Olivier Gevaert, Elena Papaleo

Project administration: Elena Papaleo

Resources: Elena Papaleo

Software: Mona Nourbakhsh, Hongjin Chen, Subhayan Akhuli

Supervision: Mona Nourbakhsh, Yuanning Zheng, Humaira Noor, Matteo Tiberti, Olivier Gevaert, Elena Papaleo

Validation: Mona Nourbakhsh, Matteo Tiberti

Visualization: Mona Nourbakhsh, Hongjin Chen, Subhayan Akhuli

Writing - original draft preparation: Mona Nourbakhsh, Hongjin Chen, Subhayan Akhuli, Elena Papaleo

Writing - review & editing: all the authors

References

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. A Cancer Journal for Clinicians. 2021;71: 209–249. doi:10.3322/caac.21660

2. Stratton MR, Campbell PJ, Futreal PA. The cancer genome. Nature. 2009;458: 719–724. doi:10.1038/nature07943

3. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW. Cancer genome landscapes. Science. 2013;340: 1546–1558. doi:10.1126/science.1235122

4. Shen L, Shi Q, Wang W. Double agents: Genes with both oncogenic and tumor-suppressor functions. Oncogenesis. 2018;7: 1–14. doi:10.1038/s41389-018-0034-x

5. Datta N, Chakraborty S, Basu M, Ghosh MK. Tumor suppressors having oncogenic functions: The double agents. Cells. 2021;10: 1–26. doi:10.3390/cells10010046

6. Hanahan D, Weinberg RA. The Hallmarks of Cancer. Cell. 2000;100: 57–70.

7. Hanahan D, Weinberg RA. Hallmarks of cancer: The next generation. Cell. 2011;144: 646–674. doi:10.1016/j.cell.2011.02.013

8. Hanahan D. Hallmarks of Cancer: New Dimensions. Cancer Discov. 2022;12: 31–46. doi:10.1158/2159-8290.CD-21-1059

9.   Nourbakhsh M, Degn K, Saksager AB, Tiberti M, Papaleo E. Prediction of cancer
driver genes and mutations: the potential of integrative computational frameworks.
Brief Bioinform. 2024;25: 1–16. doi:10.1093/bib/bbad519
10.   Konda P, Garinet S, Van Allen EM, Viswanathan SR. Genome-guided discovery of
cancer therapeutic targets. Cell Rep. 2023;42: 1–18.
doi:10.1016/j.celrep.2023.112978
11.   Liu Y, Hu X, Han C, Wang L, Zhang X, He X, et al. Targeting tumor suppressor genes
for cancer therapy. BioEssays. 2015;37: 1277–1286. doi:10.1002/bies.201500093
12.   Yu X, Zhao H, Wang R, Chen Y, Ouyang X, Li W, et al. Cancer epigenetics: from la-
boratory studies and clinical trials to precision medicine. Cell Death Discov. 2024;10:
1–12. doi:10.1038/s41420-024-01803-z
13.   Iannuccelli M, Micarelli E, Surdo P Lo, Palma A, Perfetto L, Rozzo I, et al. Can-
cerGeneNet: Linking driver genes to cancer hallmarks. Nucleic Acids Res. 2020;48:
D416–D421. doi:10.1093/nar/gkz871
14.   Colaprico A, Olsen C, Bailey MH, Odom GJ, Terkelsen T, Silva TC, et al. Interpreting
pathways to discover cancer driver genes with Moonlight. Nat Commun. 2020;11: 1–
17. doi:10.1038/s41467-019-13803-0
15.   Nourbakhsh M, Saksager A, Tom N, Chen XS, Colaprico A, Olsen C, et al. A workflow
to study mechanistic indicators for driver gene prediction with Moonlight. Brief Bioin-
form. 2023;24: 1–13. doi:10.1093/bib/bbad274
16.   Ciriello G, Miller ML, Aksoy BA, Senbabaoglu Y, Schultz N, Sander C. Emerging land-
scape of oncogenic signatures across human cancers. Nat Genet. 2013;45: 1127–
1133. doi:10.1038/ng.2762
17.   Ehrlich M. DNA hypomethylation in cancer cells. Epigenomics. 2009;1: 239–259.
doi:10.2217/EPI.09.33
18.   Esteller M. Epigenetics in Cancer. N Engl J Med. 2008;358: 1148–1159.
doi:10.1056/NEJMra072067
19.   Lakshminarasimhan R, Liang G. The role of DNA methylation in cancer. Adv Exp Med
Biol. 2016;945: 151–172. doi:10.1007/978-3-319-43624-1_7
20.   Søes S, Daugaard IL, Singers Sørensen B, Carus A, Mattheisen M, Alsner J, et al.
Hypomethylation and increased expression of the putative oncogene ELMO3 are as-
sociated with lung cancer development and metastases formation. Oncoscience.
2014;1: 367–374. doi:10.18632/oncoscience.42
21.   Zheng Y, Jun J, Brennan K, Gevaert O. EpiMix is an integrative tool for epigenomic
subtyping using DNA methylation. Cell Reports Methods. 2023;3: 1–23.
doi:10.1016/j.crmeth.2023.100515
22.   Tomczak K, Czerwińska P, Wiznerowicz M. The Cancer Genome Atlas (TCGA): An
immeasurable source of knowledge. Contemp Oncol. 2015;19: A68–A77.
doi:10.5114/wo.2014.47136
23.   Hutter C, Zenklusen JC. The Cancer Genome Atlas: Creating Lasting Value beyond
Its Data. Cell. 2018;173: 283–285. doi:10.1016/j.cell.2018.03.042
24.   Moore LD, Le T, Fan G. DNA methylation and its basic function. Neuropsychopharma-
cology. 2013;38: 23–38. doi:10.1038/npp.2012.112
25.   Sondka Z, Bamford S, Cole CG, Ward SA, Dunham I, Forbes SA. The COSMIC Can-
cer Gene Census: describing genetic dysfunction across all human cancers. Nat Rev
Cancer. 2018;18: 696–705. doi:10.1038/s41568-018-0060-1

26. Rahimi M, Teimourpour B, Marashi SA. Cancer driver gene discovery in transcrip-
tional regulatory networks using influence maximization approach. Comput Biol Med.
2019;114: 1–11. doi:10.1016/j.compbiomed.2019.103362
27. Akhavan-Safar M, Teimourpour B. KatzDriver: A network based method to cancer
causal genes discovery in gene regulatory network. BioSystems. 2021;201: 1–13.
doi:10.1016/j.biosystems.2020.104326
28. Wei PJ, Zhang D, Li HT, Xia J, Zheng CH. DriverFinder: A gene length-based network
method to identify cancer driver genes. Complexity. 2017;2017: 1–11.
doi:10.1155/2017/4826206
29. Pham VVH, Liu L, Bracken CP, Goodall GJ, Long Q, Li J, et al. CBNA: A control the-
ory based method for identifying coding and non-coding cancer drivers. PLoS Comput
Biol. 2019;15: 23. doi:10.1371/journal.pcbi.1007538
30. Dinstag G, Shamir R. PRODIGY: Personalized prioritization of driver genes. Bioinfor-
matics. 2020;36: 1831–1839. doi:10.1093/bioinformatics/btz815
31. Wei PJ, Zhang D, Xia J, Zheng CH. LNDriver: Identifying driver genes by integrating
mutation and expression data based on gene-gene interaction network. BMC Bioinfor-
matics. 2016;17: 1–10. doi:10.1186/s12859-016-1332-y
32. Zhang D, Bin Y. DriverSubNet: A Novel Algorithm for Identifying Cancer Driver Genes
by Subnetwork Enrichment Analysis. Front Genet. 2021;11: 1–10.
doi:10.3389/fgene.2020.607798
33. Li A, Chapuy B, Varelas X, Sebastiani P, Monti S. Identification of candidate cancer
drivers by integrative Epi-DNA and Gene Expression (iEDGE) data analysis. Sci Rep.
2019;9: 1–12. doi:10.1038/s41598-019-52886-z
34. Han Y, Yang J, Qian X, Cheng WC, Liu SH, Hua X, et al. DriverML: A machine learn-
ing algorithm for identifying driver genes in cancer sequencing studies. Nucleic Acids
Res. 2019;47: 1–12. doi:10.1093/nar/gkz096
35. Gu H, Xu X, Qin P, Wang J. FI-Net: Identification of Cancer Driver Genes by Using
Functional Impact Prediction Neural Network. Front Genet. 2020;11: 1–15.
doi:10.3389/fgene.2020.564839
36. Hou Y, Gao B, Li G, Su Z. MaxMIF: A New Method for Identifying Cancer Driver
Genes through Effective Data Integration. Advanced Science. 2018;5: 1–9.
doi:10.1002/advs.201800640
37. Zapata L, Susak H, Drechsel O, Friedländer MR, Estivill X, Ossowski S. Signatures of
positive selection reveal a universal role of chromatin modifiers as cancer driver
genes. Sci Rep. 2017;7: 1–15. doi:10.1038/s41598-017-12888-1
38. Terekhanova N V., Karpova A, Liang WW, Strzalkowski A, Chen S, Li Y, et al. Epige-
netic regulation during cancer transitions across 11 tumour types. Nature. 2023;623:
432–441. doi:10.1038/s41586-023-06682-5
39. D'Anello L, Sansone P, Storci G, Mitrugno V, D'Uva G, Chieco P, et al. Epigenetic
control of the basal-like gene expression profile via Interleukin-6 in breast cancer
cells. Mol Cancer. 2010;9: 1–13. doi:10.1186/1476-4598-9-300
40. Lin Y Te, Wu KJ. Epigenetic regulation of epithelial-mesenchymal transition: Focusing
on hypoxia and TGF-β signaling. J Biomed Sci. 2020;27: 1–10. doi:10.1186/s12929-
020-00632-3
41. Skrypek N, Goossens S, De Smedt E, Vandamme N, Berx G. Epithelial-to-Mesenchy-
mal Transition: Epigenetic Reprogramming Driving Cellular Plasticity. Trends in Ge-
netics. 2017;33: 943–959. doi:10.1016/j.tig.2017.08.004

42. Liu QL, Luo M, Huang C, Chen HN, Zhou ZG. Epigenetic Regulation of Epithelial to
Mesenchymal Transition in the Cancer Metastatic Cascade: Implications for Cancer
Therapy. Front Oncol. 2021;11: 1–13. doi:10.3389/fonc.2021.657546

43. Lu W, Kang Y. Epithelial-Mesenchymal Plasticity in Cancer Progression and Metasta-
sis. Dev Cell. 2019;49: 361–374. doi:10.1016/j.devcel.2019.04.010

44. Yang H, Gan L, Chen R, Li D, Zhang J, Wang Z. From multi-omics data to the cancer
druggable gene discovery: A novel machine learning-based approach. Brief Bioin-
form. 2023;24: 1–11. doi:10.1093/bib/bbac528

45. Yoshimaru T, Nakamura Y, Katagiri T. Functional genomics for breast cancer drug
target discovery. J Hum Genet. 2021;66: 927–935. doi:10.1038/s10038-021-00962-6

46. Zsákai L, Sipos A, Dobos J, Erős D, Szántai-Kis C, Bánhegyi P, et al. Targeted drug
combination therapy design based on driver genes. Oncotarget. 2019;10: 5255–5266.
doi:10.18632/oncotarget.26985

47. Cannon M, Stevenson J, Stahl K, Basu R, Coffman A, Kiwala S, et al. DGIdb 5.0: re-
building the drug-gene interaction database for precision medicine and drug discovery
platforms. Nucleic Acids Res. 2024;52: D1227–D1235. doi:10.1093/nar/gkad1040

48. Souglakos J, Boukovinas I, Taron M, Mendez P, Mavroudis D, Tripaki M, et al. Ribo-
nucleotide reductase subunits M1 and M2 mRNA expression levels and clinical out-
come of lung adenocarcinoma patients treated with docetaxel/gemcitabine. Br J Can-
cer. 2008;98: 1710–1715. doi:10.1038/sj.bjc.6604344

49. Atwell B, Chen CY, Christofferson M, Montfort WR, Schroeder J. Sorting nexin-de-
pendent therapeutic targeting of oncogenic epidermal growth factor receptor. Cancer
Gene Ther. 2023;30: 267–276. doi:10.1038/s41417-022-00541-7

50. Robichaux JP, Le X, Vijayan RSK, Hicks JK, Heeke S, Elamin YY, et al. Structure-
based classification predicts drug response in EGFR-mutant NSCLC. Nature.
2021;597: 732–737. doi:10.1038/s41586-021-03898-1

51. Robichaux JP, Elamin YY, Tan Z, Carter BW, Zhang S, Liu S, et al. Mechanisms and
clinical activity of an EGFR and HER2 exon 20-selective kinase inhibitor in non-small
cell lung cancer. Nat Med. 2018;24: 638–646. doi:10.1038/s41591-018-0007-9

52. Cohen P, Cross D, Jänne PA. Kinase drug discovery 20 years after imatinib: progress
and future directions. Nat Rev Drug Discov. 2021;20: 551–569. doi:10.1038/s41573-
021-00195-4

53. Poels KE, Schoenfeld AJ, Makhnin A, Tobi Y, Wang Y, Frisco-Cabanos H, et al. Iden-
tification of optimal dosing schedules of dacomitinib and osimertinib for a phase I/II
trial in advanced EGFR-mutant non-small cell lung cancer. Nat Commun. 2021;12: 1–
12. doi:10.1038/s41467-021-23912-4

54. van Alderwerelt van Rosenburgh IK, Lu DM, Grant MJ, Stayrook SE, Phadke M, Wal-
ther Z, et al. Biochemical and structural basis for differential inhibitor sensitivity of
EGFR with distinct exon 19 mutations. Nat Commun. 2022;13: 1–16.
doi:10.1038/s41467-022-34398-z

55. Pu X, Zhou Y, Kong Y, Chen B, Yang A, Li J, et al. Efficacy and safety of dacomitinib
in treatment-naïve patients with advanced NSCLC harboring uncommon EGFR muta-
tion: an ambispective cohort study. BMC Cancer. 2023;23: 1–10. doi:10.1186/s12885-
023-11465-2

56. Momeny M, Zarrinrad G, Moghaddaskho F, Poursheikhani A, Sankanian G, Zaghal A,
et al. Dacomitinib, a pan-inhibitor of ErbB receptors, suppresses growth and invasive
capacity of chemoresistant ovarian carcinoma cells. Sci Rep. 2017;7: 1–12.
doi:10.1038/s41598-017-04147-0

57. Brischwein K, Schlereth B, Guller B, Steiger C, Wolf A, Lutterbuese R, et al. MT110: A
novel bispecific single-chain antibody construct with high efficacy in eradicating estab-
lished tumors. Mol Immunol. 2006;43: 1129–1143. doi:10.1016/j.molimm.2005.07.034

58. Ferrari F, Bellone S, Black J, Schwab CL, Lopez S, Cocco E, et al. Solitomab, an Ep-
CAM/CD3 bispecific antibody construct (BiTE®), is highly active against primary uter-
ine and ovarian carcinosarcoma cell lines in vitro. Journal of Experimental and Clinical
Cancer Research. 2015;34: 1–8. doi:10.1186/s13046-015-0241-7

59. Keller L, Werner S, Pantel K. Biology and clinical relevance of EpCAM. Cell Stress.
2019;3: 165–180. doi:10.15698/cst2019.06.188

60. Cui Y, Li J, Liu X, Gu L, Lyu M, Zhou J, et al. Dynamic Expression of EpCAM in Pri-
mary and Metastatic Lung Cancer Is Controlled by Both Genetic and Epigenetic
Mechanisms. Cancers (Basel). 2022;14: 1–12. doi:10.3390/cancers14174121

61. Kebenko M, Goebeler ME, Wolf M, Hasenburg A, Seggewiss-Bernhardt R, Ritter B, et
al. A multicenter phase 1 study of solitomab (MT110, AMG 110), a bispecific Ep-
CAM/CD3 T-cell engager (BiTE®) antibody construct, in patients with refractory solid
tumors. Oncoimmunology. 2018;7: 1–12. doi:10.1080/2162402X.2018.1450710

62. Greenman C, Stephens P, Smith R, Dalgliesh GL, Hunter C, Bignell G, et al. Patterns
of somatic mutation in human cancer genomes. Nature. 2007;446: 153–158.
doi:10.1038/nature05610

63. Chalmers ZR, Connelly CF, Fabrizio D, Gay L, Ali SM, Ennis R, et al. Analysis of
100,000 human cancer genomes reveals the landscape of tumor mutational burden.
Genome Med. 2017;9: 1–14. doi:10.1186/s13073-017-0424-2

64. Chen ZY, Gao H, Dong Z, Shen YN, Wang ZC, Wei W, et al. NRP1 regulates radia-
tion-induced EMT via TGF-β/Smad signaling in lung adenocarcinoma cells. Int J Ra-
diat Biol. 2020;96: 1281–1295. doi:10.1080/09553002.2020.1793015

65. Madu CO, Wang S, Madu CO, Lu Y. Angiogenesis in breast cancer progression, diag-
nosis, and treatment. J Cancer. 2020;11: 4474–4494. doi:10.7150/jca.44313

66. Teleanu RI, Chircov C, Grumezescu AM, Teleanu DM. Tumor angiogenesis and anti-
angiogenic strategies for cancer treatment. J Clin Med. 2020;9: 1–21.
doi:10.3390/jcm9010084

67. Badodekar N, Sharma A, Patil V, Telang G, Sharma R, Patil S, et al. Angiogenesis in-
duction in breast cancer: A paracrine paradigm. Cell Biochem Funct. 2021;39: 860–
873. doi:10.1002/cbf.3663

68. Vartanian RK, Weidner N. Correlation of Intratumoral Endothelial Cell Proliferation
with Microvessel Density (Tumor Angiogenesis) and Tumor Cell Proliferation in Breast
Carcinoma. Breast CA Endothelial Cell Proliferation. 1994;144: 1188–1194.

69. Weidner N, Semple JP, Welch WR, Folkman J. Tumor angiogenesis and metastasis -
correlation in invasive breast carcinoma. N Engl J Med. 1991;324: 1–8.
doi:10.1056/NEJM199101033240101

70. Gargalionis AN, Papavassiliou KA, Papavassiliou AG. Implication of mTOR Signaling
in NSCLC: Mechanisms and Therapeutic Perspectives. Cells. 2023;12: 1–10.
doi:10.3390/cells12152014

71. Nevins JR. The Rb/E2F pathway and cancer. Hum Mol Genet. 2001;10: 699–703.

72. Kent LN, Leone G. The broken cycle: E2F dysfunction in cancer. Nat Rev Cancer.
2019;19: 326–338. doi:10.1038/s41568-019-0143-7

73. Stark GR, Taylor WR. Analyzing the G2/M Checkpoint. Checkpoint Controls and Can-
cer. New Jersey: Humana Press; 2004. pp. 051–082. doi:10.1385/1-59259-788-2:051

Supporting information

S1 Fig. Integration of Moonlight and EpiMix for prediction of cancer driver genes. (A) Heatmap showing number of differentially methylated CpGs and classifications of methylation status in the oncogenic mediators in lung adenocarcinoma. The heatmap was generated using the plotGMA function. **(B)** Venn diagram comparing oncogenic mediators predicted from Moonlight's primary layer with functional genes predicted from EpiMix in lung adenocarcinoma. The functional genes are genes containing differentially methylated CpG pairs whose DNA methylation state is associated with expression of the gene. Only those functional genes that contained the same methylation state in all of its associated CpGs were included in this comparison, and moreover, the dual methylation states were excluded. **(C)** Heatmap showing the effect of the predicted driver genes in lung adenocarcinoma on apoptosis and proliferation of cells. This heatmap was generated using the function plotMoonlightMet. These effects define the basis upon which the oncogenic mediators are predicted from the PRA step in Moonlight's primary layer. **(D)** Heatmap showing number of differentially methylated CpGs and classifications of methylation status in the oncogenic mediators in thyroid carcinoma. The heatmap was generated using the plotGMA function. **(E)** Venn diagram comparing oncogenic mediators predicted from Moonlight's primary layer with functional genes predicted from EpiMix in thyroid carcinoma. The functional genes are genes containing differentially methylated CpG pairs whose DNA methylation state is associated with expression of the gene. Only those functional genes that contained the same methylation state in all of its associated CpGs were included in this comparison, and moreover, the dual methylation states were excluded. **(F)** Heatmap showing the effect of the predicted driver genes in thyroid carcinoma on apoptosis and proliferation of cells. This heatmap was generated using the function plotMoonlightMet. These effects define the basis upon which the oncogenic mediators are predicted from the PRA step in Moonlight's primary layer.
(PDF)

S2 Fig. Number of driver gene-drug interactions. Number of driver gene-drug interactions in **(A)** basal-like breast cancer, **(B)** lung adenocarcinoma, and **(C)** thyroid carcinoma found by querying DGIdb. The driver genes are stratified into OCGs and TSGs. The number of drug interactions is shown on the *x* axis and the driver genes are shown on the *y* axis.
(PDF)

S1 Text. Evidence integration for defining driver genes.
(PDF)

S2 Text. Methods of case study: Prediction of driver genes with differential methylation in basal-like breast cancer, lung adenocarcinoma, and thyroid carcinoma using Moonlight.
(PDF)