THE HISTONE CHAPERONE SPN1 PRESERVES CHROMATIN PROTECTIONS AT PROMOTERS AND NUCLEOSOME POSITIONING IN OPEN READING FRAMES

AUTHORS

Andrew J. Tonsager[1], Alexis Zukowski[2], Catherine A. Radebaugh[1], Abigail Weirich[2], Laurie A. Stargell[1,*], Srinivas Ramachandran[2,3*]

[1]Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins, CO, 80523-1870, USA

[2]Department of Biochemistry and Molecular Genetics, [3]RNA Bioscience Initiative, University of Colorado School of Medicine, Aurora, CO, 80045, USA

*Correspondence: srinivas.ramachandran@cuanschutz.edu, laurie.stargell@colostate.edu

ABSTRACT

Spn1 is a multifunctional histone chaperone that associates with RNA polymerase II during elongation and is essential for life in eukaryotes. While previous work has elucidated regions of the protein important for its many interactions, it is unknown how these domains contribute to the maintenance of chromatin structure. Here, we employ digestion by micrococcal nuclease followed by single-stranded library preparation and sequencing (MNase-SSP) to characterize chromatin structure in *Saccharomyces cerevisiae* expressing wild-type or mutants of Spn1 (*spn1^{K192N} or spn1$^{141-305}$*). We mapped protections of all sizes genome-wide. Surprisingly, we observed a widespread loss of short fragments over nucleosome-depleted regions (NDRs) at promoters in the *spn1^{K192N}*-containing strain, indicating critical functions of Spn1 in maintaining normal chromatin architecture outside open reading frames. Additionally, there are shifts in DNA protections in both Spn1 mutant expressing strains over open reading frames, which indicate changes in nucleosome and subnucleosome positioning. This was observed in markedly different Spn1 mutant strains, demonstrating that multiple functions of Spn1 are required to maintain proper chromatin structure in open reading frames. Changes in chromatin structure correlate positively with changes in gene expression as shown by RNA-seq analysis in the Spn1 mutant strains. Taken together, our results reveal a previously unknown role of Spn1 in the maintenance of NDR architecture and deepen our understanding of Spn1-dependent chromatin maintenance over transcribed regions.

INTRODUCTION

Eukaryotes package their genomes by wrapping ~147 bp of DNA in about 1.7 superhelical turns around an octamer of histone proteins to form nucleosomes (Davey et al. 2002; Flaus et al. 1996; Luger et al. 1997). This process occurs in a stepwise manner, as DNA initially wraps around a tetramer of histones H3–H4 to form a tetrasome, followed by sequential additions of two H2A-H2B dimers to form a hexasome then a nucleosome (Polo and Almouzni 2006). Tetrasome and hexasome subnucleosomal structures have been observed both *in vitro* (Brower-Toland et al. 2002) and *in vivo* (Ramachandran et al. 2017) and represent stable subnucleosomal intermediates in nucleosome assembly and disassembly. Nucleosomes regulate DNA access to many proteins that perform essential processes such as DNA replication (Eaton et al. 2010; Groth et al. 2007), repair (Adkins et al. 2013; Groth et al. 2007), recombination (Bevington and Boyes 2013), and transcription. Nucleosomes regulate transcription in a variety of ways. Positioning of nucleosomes at promoters can hinder transcription initiation by preventing the binding of transcription factors (Henikoff et al. 2011; Kubik et al. 2015) and nucleosome-depleted regions (NDRs) are important for regulation of gene expression (Kubik et al. 2015). Chromatin remodelers, such as the RSC complex, actively maintain NDRs after nucleosome deposition (Lorch et al. 2014). This region is bound by many other factors, such as activators that bind and compete with nucleosome formation within NDRs, contributing to the low nucleosome density in these regions (Rando and Winston 2012). NDRs upstream of transcription start sites contain promoter sequences bound by transcription initiation factors. Positioning of nucleosomes within coding regions can play an important role in preventing intragenic transcription initiation by reducing the accessibility of factors to intragenic initiation sites (Kaplan et al. 2003). Nucleosomes inhibit transcription elongation as they pose a significant barrier to RNA polymerase II (RNAPII) and reduce its transcription rate (Brahma and Henikoff 2020; Selth et al. 2010). Hence, chromatin structure *in vivo* must be dynamic to allow for selective gene expression. An important class of proteins responsible for modifying chromatin structure are histone chaperones, which bind histones and facilitate histone deposition, exchange, or removal from chromatin, independent of ATP (Bai and Morozov 2010).

Spn1 is a multifunctional and essential histone chaperone that is highly conserved across

eukaryotes. *S. cerevisiae* Spn1 preferentially binds H3-H4 histones (Li et al. 2018) via its N-

terminal region, with a high binding affinity ($K_D \sim 10nM$). Spn1 affinity to H3-H4 is at least 50-

fold higher than that to H2A-H2B dimers (Li et al. 2022). Spn1 can deposit H3-H4 onto DNA and

facilitate the assembly of H3-H4 tetrasomes *in vitro* (Li et al. 2018). Spn1 also has a well-

characterized interaction with Spt6, another essential histone chaperone; they bind each other

with high affinity ($K_D = 10\text{-}100nM$) to form a heterodimer (Gavin et al. 2002; Krogan et al. 2002;

Lindstrom et al. 2003; McDonald et al. 2010). The central domain of Spn1 is a highly conserved

eight-helix bundle and is primarily responsible for binding Spt6 (Li et al. 2022). Depletion of

either protein through the auxin-inducible degron system results in a reduction in recruitment of

the other to chromatin (Reim et al. 2020). Spn1 binds nucleosomes with its C-terminal region (Li

et al. 2018; Li et al. 2022).

A mutant of Spn1, *spn1$^{141\text{-}305}$*, lacks the N- and C-terminal domains, which are responsible for

histone and nucleosome binding, respectively. When *spn1$^{141\text{-}305}$* is the sole source of Spn1, yeast

cell growth is relatively healthy (Fischbeck et al. 2002; Li et al. 2018). Though *spn1$^{141\text{-}305}$* cannot

bind histones and nucleosomes, it retains the ability to bind Spt6 and RNAPII (Li et al. 2018; Li et

al. 2022). A different allele of Spn1, *spn1^{K192N}*, is similarly viable, but we have previously

demonstrated that these cells have reduced Spn1 recruitment to the *CYC1* locus and fail to

recruit Spt6 to *CYC1* (Zhang et al. 2008). The *spn1^{K192N}* mutant also has reduced Spn1 binding to

RNAPII by co-immunoprecipitation (Zhang et al. 2008). While both alleles cover a *SPN1*

knockout, we have previously shown that they are partially defective or involved in redundant

functions with other cellular proteins. For example, yeast expressing the Spn1 mutants in a

background containing knockouts in other elongation and transcription factors experience

significant defects in growth (Li et al. 2018; Zhang et al. 2008). Recent work demonstrates that

Iws1, the human ortholog of Spn1, serves as a hub that maintains association of numerous

additional transcription factors (Cermakova et al. 2021).

While we and others have previously demonstrated that Spn1 possesses multiple interactions

pertinent to its chromatin functions, the role of Spn1 *in vivo* is not well understood. Here,

micrococcal nuclease digestion followed by single-stranded library preparation and sequencing

(MNase-SSP) (Ramani et al. 2019) is used to characterize chromatin structure in yeast expressing

wild-type or mutant Spn1 alleles. Compared to traditional MNase-seq (with or without selection

of ~147 bp fragments), MNase-SSP allows for the analysis of a wider range of protections by

better retaining MNase-protected fragments of a variety of lengths (Ramani et al. 2019). This

includes both nucleosomal protections and subnucleosomal protections. Subnucleosome

protections are broadly defined as DNA protections with footprints smaller than the protection

by a nucleosome (~147bp) (Henikoff et al. 2011; Kent et al. 2011). Specific subnucleosome

protections can arise due to histone-containing subnucleosomal intermediates at open reading

frames (Ramachandran et al. 2017; Rhee et al. 2014). The MNase-SSP protections also include

non-nucleosomal protections, defined as DNA fragments obtained through non-histone

protections, such as protection by transcription factors, pre-initiation complex, and other DNA

binding factors. In this study, using MNase-SSP, we outline the effects of $spn1^{K192N}$ and $spn1^{141-305}$

on chromatin structure *in vivo*. Furthermore, to determine whether changes in chromatin

structure identified through MNase-SSP correlate with changes in gene expression, we also

performed RNA-seq to identify differentially expressed genes in cells expressing $spn1^{K192N}$ or

$spn1^{141-305}$.

We find that despite distinct defects with known binding partners, both $spn1^{K192N}$ and $spn1^{141-305}$

expression show similar chromatin defects across genes, namely a progressive 3' downstream

shift in chromatin protections over gene bodies. Intriguingly, MNase-SSP also revealed a

molecular mutant phenotype specific to cells expressing $spn1^{K192N}$: the loss of protections over

nucleosome-depleted regions (NDRs) in promoters, most dramatic for fragments <60bp in

length. Additionally, expression of $spn1^{141-305}$, but not $spn1^{K192N}$, produces a unique molecular

mutant phenotype of a shift in the +1 and +2 nucleosome positions. Mapping the midpoints of

the sequenced fragments indicates that shifts correspond to a downstream redistribution of

nucleosome and subnucleosomal occupancy at alternative rotational positions; alternative

rotational positions have been previously observed in the yeast genome (Brogaard et al. 2012).

This molecular mutant phenotype of the shift downstream is largest at longer genes. Our RNA-

seq results also revealed that expression of $spn1^{K192N}$ or $spn1^{141-305}$ produces global reduction in

gene expression. Intriguingly, genes differentially expressed in the Spn1 mutant strains are

enriched in longer genes, revealing a correlative relationship between instances of nucleosome

positioning shift and changes in gene expression. Taken together, these studies reveal the

importance of Spn1 in preserving chromatin structure across the yeast genome.

MATERIAL AND METHODS

Yeast Culturing and Strains

All *Saccharomyces cerevisiae* strains were grown in yeast extract, peptone, and dextrose (YPD;

2% dextrose) media at 30°C in biological duplicate independently. A description of yeast strains

and plasmids is provided in Supplemental Table S1. The strains used in this study were

previously created as described (Li et al. 2018; Thurston et al. 2018; Zhang et al. 2008). In

summary, strains were transformed with a covering plasmid containing the *URA3* gene and the

wild-type *SPN1* open reading frame flanked by the *TOA1* promoter and terminator; the *TOA1*

sequences were used to prevent integration within the covering plasmid of the *LEU2* derivative

described next. Endogenous *SPN1* was replaced by a *LEU2* fragment flanked by the *SPN1*

promoter and terminator through homologous recombination. Plasmids containing the *HIS3*

gene and wild-type *SPN1*, *spn1^{K192N}*, or *spn1^{141-305}* under the control of their native promoter and

terminator were introduced into strains by plasmid shuffling, which results in removal of the

covering plasmid.

RNA preparation for sequencing

RNA was isolated from yeast expressing either wild-type Spn1, *spn1^{K192N}*, or *spn1^{141-305}* in

triplicate 10-ml cultures by hot-acid phenol extraction as previously described (Iyer and Struhl

1996). Cells were grown in YPD at 30°C to mid-exponential phase (OD600 ~ 1.0), harvested, and

each sample was spiked to obtain a final concentration of 1 *S. pombe* to 10 *S. cerevisae* cells.

RNA libraries were prepared with poly-A enrichment and sequenced using the NovaSeq X Plus

Series (PE150) platform (Novogene).

RNA-seq analysis

A concatenated fasta file was created using i) cDNA sequences of *S. cerevisiae* (release 113 from

Ensembl), ii) cDNA sequences of *S. pombe* (from PomBase), iii) whole genome sequence of *S.*

cerevisiae, and iv) whole genome sequence of *S. pombe*. A Salmon index was created using the

concatenated fasta, setting the whole genome sequences as decoys. Transcript quantification

was performed using Salmon v1.9 with libType set as automatic (Patro et al. 2017). Differential

gene expression analysis was performed using DeSeq2 (Love et al. 2014) (v1.30.1) in R (v4.0.3).

For spike-in normalization, size factors were estimated by setting *S. pombe* genes as control

genes, resulting in the distribution of log2FC for S. pombe genes centered at 0 (**Figure S18A-C,**

File S1). The expression matrix across genotypes was transformed using the "rlog" function in

DESeq2 and the transformed matrix was used for performing principal component analysis

(PCA) using the "prcomp" function in R and plotted using ggplot2. Gene set enrichment analysis

was performed using g:Profiler (Kolberg et al. 2023).

Isolation of nuclei and Micrococcal nuclease digestion

Yeast nuclei isolation and MNase digestion were performed as described in (Zhang and Reese

2005) with modifications. Yeast strains were grown in 400 ml of YP +2% glucose until an OD_{600}

of ~1.0 was reached. The cells were collected by centrifugation in a JA-10 rotor at 5,000 RPM

for 10 minutes at 4^{o}C. The supernatant was decanted from above the cell pellets, and the cells

were suspended in 30 ml of Sorbitol buffer (50 mM Tris-Cl pH 7.5, 1.0 M Sorbitol, 10 mM $MgCl_2$,

10 mM 2-Mercaptoethanol and 1mM Phenylmethylsulfonyl fluoride (PMSF)). The cell

suspensions were transferred to chilled 50 ml conical tubes and collected by centrifugation in a

GPKR centrifuge at 3,200 RPM for 5 minutes at 4^{o}C. The supernatants were decanted from above

the cell pellets and the cells were washed two additional times with 30 mL of Sorbitol buffer. The

mass of the cell pellets was determined, and the cells were suspended in 0.66 ml of Sorbitol

buffer per gram of cells.

Spheroplasting was completed by adding 0.34 mL of Zymolase (10 $^{mg}/_{mL}$) per gram of cells and

incubating at 30^{o}C with gentle agitation until ~90% of the cells were converted to spheroplasts.

The spheroplasts were collected by centrifugation in a GPKR centrifuge at 3,200 RPM for 5

minutes at 4°C and then gently washed two times with 5 ml of Sorbitol buffer as described

above. The washed spheroplasts were suspended in 25 mL of Ficoll buffer (18% Ficoll 400, 20

mM Tris-Cl pH 7.5, and 0.5 mM $MgCl_2$), transferred to a 50 ml Potter-Elvehjem homogenizer and

lysed on ice with six up-down even strokes with a handheld drill at max rpm. The lysates were

transferred to chilled 25 ml centrifuge tubes, and the remaining cells were pelleted in a JA-20

rotor at 6400 x rpm for 5 minutes at 4°C. The supernatants containing yeast nuclei were

transferred to new centrifuge tubes and pelleted at 15,000 x rpm at 4°C. The nuclei were

suspended in 10 ml of MNase digestion buffer (20mM Tris-Cl pH 8.0, 1.5mM $CaCl_2$, 150mM

NaCl, and 1mM PMSF). To estimate the yield of nuclei, 100 µl aliquots of the nuclei suspensions

were diluted to 1 ml with MNase digestion buffer and the OD_{600} was determined. The nuclei

were again pelleted at 15,000 x rpm for 30 minutes at 4°C and suspended in MNase digestion

buffer at a ratio of 2.4 ml per 0.2 OD_{600}. Five aliquots (400 µl) of the nuclei suspensions were

transferred to microcentrifuge tubes and incubated at 37°C for 10 minutes. Digestion buffer or

MNase was added to samples to yield a final concentration of 0, 32, 64, or 128 U/ml, and

incubation at 37°C continued for 10 minutes. 100 µl of stop solution (50mM EDTA, 2.5% SDS,

0.25mg/ml Proteinase K) was added to each reaction, and the samples were mixed and

incubated at 37°C overnight. The MNase-digested DNA samples were extracted twice with

phenol:chloroform, once with chloroform, and precipitated with ethanol. The DNA was washed

with 70% ethanol, dried at room temperature, then suspended in 100 µl of TE buffer (pH 7.5)

and 3 µl of RNaseA (10 mg/ml) added. The RNase reactions were incubated overnight at 37°C,

followed by ethanol precipitation, washing with 70% ethanol, and drying at room temperature.

The precipitated DNA samples were suspended in 100µl of TE buffer (10mM Tris-Cl, pH 8.0 and

1mM EDTA, pH 8.0), and 10µl of each sample was subjected to agarose gel electrophoresis.

Single-stranded DNA library preparation (SSP) and sequencing

MNase-protected DNA fragments were made into sequencing libraries using the single-

stranded library protocol as previously described (Gansauge and Meyer 2013; Ramani et al.

2019; Snyder et al. 2016). In brief, 10ng DNA was dephosphorylated using FastAP

Thermosensitive Alkaline Phosphatase (Thermo Scientific cat. EF0651), denatured, and incubated

overnight with CircLigaseII (Lucigen cat. CL9025K) and 0.093-0.125 µM biotinylated CL78 primer

(35) at 60ºC with shaking every 5 minutes. DNA fragments were then denatured and bound to

magnetic streptavidin M-280 beads (Invitrogen cat. 11205D) for 30 minutes at room

temperature with nutation. Beads were washed, and second-strand synthesis was performed

using Bst 2.0 DNA polymerase (NEB cat. M0537) with an increasing temperature gradient of 15-

31ºC with shaking at 1750 rpm. Beads were washed, and a 3' gap fill was performed using T4

DNA polymerase (Thermo Scientific cat. EL0011) for 30 minutes at room temperature. Beads

were washed, and a double-stranded adapter was ligated using T4 DNA ligase (Thermo

Scientific cat. EP0062) for 2 hours at room temperature with shaking at 1750 rpm. Beads were

washed and resuspended in 30 µL 10 mM TET buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA pH

8.0, 0.05% Tween-20). DNA was denatured at 95ºC for 3 min, and DNA libraries were collected

after immediate magnetic separation.

Quantitative real-time PCR was performed on DNA libraries using iTAQ Supermix (Bio-Rad cat.

1725124), and Ct values were used to determine the number of PCR cycles needed to amplify

each library. PCR was performed with KAPA HiFi DNA polymerase (Kapa Biosystems cat. KK2502)

using barcoded indexing primers for Illumina. Primer dimers were removed from the libraries

using AMPure beads (Beckman Coulter cat. A63881). Libraries were eluted in 0.1X TE and

concentrations were determined using Qubit. The length distribution of each library was

assessed by the Agilent Bioanalyzer using the D1000 or HSD1000 cassette. Libraries were

sequenced for 150 cycles in paired-end mode on the NovaSeq 6000 system at the University of

Colorado Cancer Center Genomics Shared Resource.

MNase-SSP analysis

Sequencing reads were trimmed using Cutadapt (Martin 2011): Illumina adapter sequences were

removed, and reads were trimmed to 140 bp. Reads less than 35 bp were discarded. Trimmed

reads were aligned to the sacCer3 version of *Saccharomyces cerevisiae* genome using bowtie2

(Martin 2011). Samtools (Li et al. 2009) and bedtools (Quinlan and Hall 2010) were used for

processing aligned reads from sam to bed files. Coverage at 1 bp windows genome-wide for

specific fragment lengths was calculated as the number of reads that mapped at each window,

normalized by the factor N: N = 139712364/(Total number of mapped reads). 139712364 was a

number chosen arbitrarily. Coverage of fragment centers for specific fragment lengths was also

calculated similarly at 1 bp windows. V-plots were generated using custom scripts that are

publicly available (https://github.com/srinivasramachandran/VPLOTS/). Nucleosome positions

were obtained from (Brogaard et al. 2012). Transcription start sites were obtained from (Park et

al. 2014). Genes were stratified by expression using published NET-seq data (Churchman and

Weissman 2011). NET-seq normalized read counts were summed between TSS+100 and

TSS+200. Z-scores across rotational positions spanning -50 to +50 bp surrounding each

nucleosome dyad position were calculated for each mapped dyad for each gene. Positive Z-

scores denote preferred rotational positions within a nucleosome position of one gene. The Z-

scores of wild-type Spn1 were subtracted from the Z-scores of Spn1 mutants for each gene to

obtain changes in rotation position preference due to Spn1 mutation.

RESULTS

MNase-SSP interrogates chromatin protections across the yeast genome

To study the role of Spn1 in maintaining chromatin structure, wild-type Spn1, $spn1^{K192N}$, and

$spn1^{141-305}$ containing strains were subjected to MNase-SSP. MNase-SSP was performed with

increasing levels of MNase, which digests linker DNA at 25 times the rate of nucleosomal DNA

and produces protected genomic fragments of varied sizes (Chereji et al. 2017). Samples were

subjected to agarose gel electrophoresis to qualitatively assess the population of

polynucleosomal, nucleosomal, and smaller fragment sizes after MNase digestion (**Figure S1A,**

File S1). Fragments of all lengths were sequenced and mapped to the *S. cerevisiae* genome. A

metagene analysis was performed to generate plots of fragment midpoint versus fragment

length ("V-plots") (Henikoff et al. 2011) centered at the dyad of each nucleosome from the +1 to

+10 positions. V-plots centered at +1, +2, and +5 exemplify that plots at each nucleosome

position show a checkered pattern that appears similar across both wild-type (**Figure 1A**) and

mutant samples (**Figure 1B, C**). To construct V-plots, we randomly sampled fragments genome-

wide to yield similar fragment length distributions across WT and Spn1 mutant datasets (**Figure**

S1B, File S1). Dyad definitions from chemical cleavage mapping were used to define

nucleosome dyad positions spanning 4688 yeast genes (Brogaard et al. 2012) (**Figure S2, File**

S1). The density at position 0 and fragment length 147 bp corresponds to the strongest dyad

signal (**Figure 1**). For fragments ~150 bp in length, this arises from peaks in density separated

by a 10 bp periodicity on either side of the dyad, suggesting nucleosome occupancy at

previously mapped rotational positions surrounding yeast nucleosomes (Brogaard et al. 2012;

Chereji et al. 2018). Additionally, the V-plots indicate the existence of rotational positioning of

subnucleosomal protections in all the samples, as this pattern persists at shorter fragment

lengths. Notably, at ~50-150 bp upstream of the +1 dyad, the $spn1^{K192N}$ mutation, but not

$spn1^{141-305}$, produces a loss in fragment density for short protections, corresponding to a

decrease in protection over nucleosome-depleted regions (NDR) (**Figure 1B**). This finding

suggests that Spn1 may contribute towards maintenance of the architecture of NDRs, a newly

described function for Spn1.

The $spn1^{K192N}$ mutation reduces stability of chromatin protections over nucleosome-

depleted regions

The V-plot analyses indicate that MNase-protected fragments are reduced in the nucleosome-

depleted regions (NDRs) specifically in the $spn1^{K192N}$ mutant yeast strain (**Figure 1**). NDRs are

flanked by well-positioned -1 and +1 nucleosomes, which reside directly upstream and

downstream of this region respectively. There are many contributors to the architecture of

NDRs, including chromatin remodelers, activators, transcription initiation factors, etc. (Lorch et

al. 2014; Rando and Winston 2012). Many of these non-nucleosomal factors have small

footprints (<60 bp) (Henikoff et al. 2011; Kent et al. 2011). The V-plot analyses suggested that

short protections of these lengths are reduced over the NDR in $spn1^{K192N}$ cells. To further explore

this, we mapped the enrichment of short (<60 bp) fragments relative to the +1 nucleosome

position of genes. This range of fragments produces a significant peak between the locations of

the +1 and -1 nucleosomes in wild-type cells, and the abundance of fragments within this peak

is dramatically reduced in $spn1^{K192N}$, but not $spn1^{141-305}$ cells (**Figure 2A**). The enrichment of <60

bp fragments shows that short non-nucleosomal protections are lost over NDRs in *spn1^{K192N}*

cells.

NDRs are maintained and protected by a wide range of proteins, including those sensitive to

MNase (Chereji et al. 2017; Henikoff et al. 2011; Kubik et al. 2015). By varying MNase

concentration, less stable structures can be revealed that are more sensitive to higher levels of

MNase. To probe the MNase sensitivity of the NDR in Spn1 mutants, datasets generated from

treatment with a lower amount of MNase were analyzed. We mapped smaller fragments from

these samples and found that the NDR is largely unaffected in these samples (**Figure 2B**). This

demonstrates that *spn1^{K192N}* cells exhibit a defect in protein protections in the NDRs, which

renders the protections significantly less stable than wild-type cells at higher MNase

concentrations. Taken together, this indicates that the stability of protein protections in the NDR

is compromised in *spn1^{K192N}* cells.

The loss of short protections in the NDR prompted the question of whether longer protections

are also affected in *spn1^{K192N}*. To assess which lengths of protected fragments are reduced in this

region, bins of fragment lengths were generated to selectively analyze their enrichment in wild-

type versus Spn1 mutant-expressing yeast. Fragments approximately 147bp in length are

captured in a bin of fragments between 142-152 bp in length, and can be interpreted as

nucleosomal protections, as the DNA is protected from MNase by its histone contacts. These

fragments represent a significant portion of the sample, as the MNase-SSP libraries exhibit a

local peak in fragments of this length (**Figure S1B, File S1**). The abundance of protected

fragments both 83-93 bp and 102-112 bp can be interpreted as subnucleosomal protections

with loss of histone-DNA contacts due to removal of an H2A-H2B dimer (Ramachandran et al.

2017; Rhee et al. 2014). Additionally, mapping fragments of 61-71 bp can be interpreted as

subnucleosomal protections with loss of histone-DNA contacts for both H2A-H2B dimers

(Ramachandran et al. 2017). All of these ranges in fragment lengths also correspond to peaks in

the fragment length distribution of our samples (**Figure S1B, File S1**). To determine if these

protections over NDRs change upon Spn1 mutation, fragments from each of these bins were

mapped over the yeast genome. The loss of fragments observed in *spn1^{K192N}* diminishes as

fragment length increases (**Figure S3, File S1**). This suggests that Spn1 maintains mainly short
non-nucleosomal protections in the NDR.

The prior metagene analyses demonstrated that *spn1^{K192N}* reduces protections over the NDR,
with this defect becoming increasingly dramatic as the DNA fragment lengths shorten. To
investigate the fraction of NDRs driving this effect in the metaplots, the normalized counts of 0-
60 bp fragments were plotted as a heatmap, with genes aligned by their +1 nucleosome dyad
position and ranked by length (**Figure S4, File S1**). To visualize the difference in fragment
occupancy between wild-type and mutant strains, the wild-type occupancy scores were
subtracted from the *spn1^{K192N}* or *spn1$^{141-305}$* samples to generate a difference heatmap (**Figure**
2C). In *spn1^{K192N}* cells but not *spn1$^{141-305}$* cells, reduced 0-60 bp fragment occupancy was
observed directly upstream of the +1 dyad locations across genes ranked by length. This
indicates that the defect in short protections in NDRs is consistent across all yeast genes in
spn1^{K192N} cells.

While nucleosomes are generally depleted in NDRs, recent work has elucidated that a subset of
yeast NDR regions do contain fragile nucleosomes which are highly sensitive to MNase (Chereji
et al. 2017; Kubik et al. 2015). Our analyses above demonstrate that Spn1 maintains primarily
short non-nucleosomal protections in NDRs across the yeast genome. However, it is possible
that the subset of NDRs containing fragile nucleosomes may exhibit larger defects in these non-
nucleosomal protections compared to NDRs without fragile nucleosomes in *spn1^{K192N}* cells. To
address this possibility, we mapped the enrichment of short (<60 bp) fragments relative to the
+1 nucleosome position of genes with either a fragile –1 nucleosome or stable –1 nucleosome
as defined by Kubik *et al.* (Kubik et al. 2015). To assess the MNase sensitivity of these
protections, we mapped fragments from both conditions treated with a higher or lower amount
of MNase as previously described. This analysis demonstrates that both NDRs with a fragile –1
nucleosome and those with a stable –1 nucleosome experience a similar loss of 0-60 bp
protections in *spn1^{K192N}* cells but display no differences in nucleosomal protections from wild-
type in the NDR (**Figure S5, File S1**). Taken together, this indicates that Spn1 maintenance of

short non-nucleosomal protections in NDRs is critical at both of these classes of NDRs found in

the yeast genome.

358   Spn1 influences nucleosome positioning over genes

Spn1 is a transcription elongation factor physically associated with the RNAPII elongation

complex (Ehara et al. 2022), and its genome-wide occupancy is highest over gene bodies (Mayer

et al. 2010; Rossi et al. 2021). To further investigate the role of Spn1 on preserving chromatin

structure over gene bodies, the normalized counts of 80-170 bp fragments were first plotted as

a heatmap (**Figure S6A, File S1**). To visualize differences in pattern between wild-type and

mutant strains, the wild-type occupancy scores were subtracted from the *spn1^{K192N}* or *spn1$^{141-305}$*

samples to generate a difference heatmap (**Figure 3A**). As this range of fragments is sufficiently

wide to include subnucleosomal protections, this analysis also shows the loss of protection

across NDRs in the *spn1^{K192N}* mutant strain (**Figure 3A,** left panel). The difference heatmap also

reveals a striated pattern across the open reading frames of genes, which is strikingly similar in

both mutants. This striated pattern of blue and red stripes seen in the difference heatmap

highlight regions of lost fragment occupancy over wild-type dyad positions followed by an

increase in fragment occupancy directly downstream of these positions. To specifically assess

changes in nucleosome protections in the Spn1 mutant expressing strains, we repeated our

analysis by exclusively mapping 142-152 bp fragments over our gene list (**Figure S6B, File S1**).

We observe a striated pattern in the difference heatmap over open reading frames similar to the

pattern observed in the difference heatmap generated with 80-170 bp fragments. Consistent

with our prior profile analysis (**Figure S3, File S1**), we observe no loss of protection in 142-152

bp fragments over the NDR in the *spn1^{K192N}* strain (**Figure S6C, File S1**).

To investigate nucleosome positioning across gene bodies further, an additional metagene

analysis was performed by averaging the normalized read count over genes and plotting the

result as a profile. The wild-type strain produces the expected profile of regularly phased and

well-positioned peaks, interpreted as peaks in nucleosomes occupancy (**Figure 3B**). Nucleosome

positions become "fuzzier" with each downstream nucleosome from the TSS; increases in the

variability of distance of downstream nucleosome positions from the TSS result in a wider and

flatter peak within the population. To quantify the differences between the wild-type and

mutant profiles, a difference plot between wild-type and mutant enrichment was generated

(**Figure S6D, File S1**). This difference plot indicates that nucleosome positions +1 and +2 shift

specifically in cells expressing the $spn1^{141-305}$ allele, and further downstream nucleosome

positions shift in both mutant strains. This result suggests that Spn1 maintains nucleosome

positioning across genes, but the contributions of its functional domains differ in the

maintenance of nucleosomes at the 5′ ends of genes versus across gene bodies.

The profiles of wild-type and mutant normalized read count indicate that that expression of

either $spn1^{K192N}$ or $spn1^{141-305}$ mutant alleles produces a progressive 3′ downstream shift in

nucleosome positioning. Rescaling this profile into individual profiles at different distances from

the +1 dyad to feature the +1 to +6 nucleosome positions illustrates that expression of either

mutant produces the trend of a progressive 3′ downstream shift in nucleosome positioning

(**Figure 3C; Figure S7, File S1**). These findings align with recently published work that employed

conventional MNase-seq to demonstrate Spn1 maintains nucleosome positioning over gene

bodies, as expression of $spn1^{K192N}$ produced a progressive downstream shift in nucleosome

positioning (Mayer et al. 2010; Rossi et al. 2021). The findings from our MNase-SSP approach are

similar and support this notion, as well as provide the new finding that expression of the

functionally distinct mutant allele $spn1^{141-305}$ also exhibits a defect in nucleosome positioning.

Quantification of nucleosome positioning shifts reveals that rotational position preference

is redistributed in Spn1-mutant cells particularly at longer genes

Our analysis of the MNase-SSP data from wild-type and mutant Spn1 expressing yeast thus far

has revealed global shifts in chromatin protections over open reading frames. Genes across the

yeast genome vary in length, and this prompted an investigation of gene length as a parameter

for nucleosome shift. To begin examining this relationship, the gene list was ranked by length

into quartiles of equal gene numbers (**Figure S8A, File S1**). Mapping fragments 80-170 bp and

generating a difference plot as performed previously shows that the furthest downstream

nucleosomes present in the longest quartile of genes exhibit the largest shift in nucleosome

position (**Figure S8B, File S1**). Strikingly, the set of shortest genes experienced, on average,

minimal defects in nucleosome positioning, even at nucleosome positions that were shifted in

the longer genes.

To statistically validate shifts in nucleosome positioning, we developed a method to quantify the

occupancy of nucleosomes at each rotational position. These positions correspond to DNA

sequences with favorable histone-DNA interactions (Satchwell et al. 1986; Segal et al. 2006).

Midpoints from sequenced fragments 142-152 bp in length were first mapped to the genome

over each quartile ranked by gene length. This analysis reveals that peaks in 142-152 bp

fragments in the mutant strains occupy the same rotational positions as wild-type cells, but with

changes in occupancy at specific alternative rotational positions. Occupancy is gained

downstream of the dyad and lost upstream of the dyad location in the mutant strains, indicative

of a shift in nucleosome positioning (**Figure 4A; Figure S9, File S1**). As this shift manifests as a

change in occupancy at discrete positions, this can be quantified by calculating the change in

occupancy at each rotational position for each nucleosome. To accomplish this, Z-scores across

rotational positions spanning -50 to +50 bp surrounding each nucleosome dyad position were

calculated for each mapped dyad for each gene. The Z-scores of wild-type Spn1 were subtracted

from the Z-scores of Spn1 mutants for each gene to obtain changes in rotation position

preference due to Spn1 mutation. The median difference in Z-scores was plotted across all

genes at each rotational position of each nucleosome position was plotted. The previous

analysis indicates that the shifts in nucleosome positioning correlated with gene length. To

evaluate the extent to which these correlations are significant, Z-scores and p-values (using the

Wilcoxon Rank Sum test) were calculated for the changes in fragment midpoint occupancy at

each rotational position around each nucleosome dyad for the genes in our previously

determined quartiles of gene length. This analysis demonstrates that shifts in nucleosome

positioning occur most dramatically at downstream nucleosome positions in longer genes, as

there were more rotational positions with significant changes in Z-scores over these genes.

(**Figure 4B**). Additionally, *spn1$^{141-305}$* mutant cells exclusively have rotational positions with

statistically significant changes in nucleosome occupancy at the +1 and +2 nucleosome

positions. This demonstrates that the disordered N- and C-terminal regions of Spn1 specifically

play a role in the proper maintenance of nucleosomes at the 5' end of open reading frames.

While both mutant strains experience shifts in nucleosome positions, $spn1^{141-305}$ mutant cells

exhibit more positions with significant changes in 142-152 bp protected fragment occupancy

over open reading frames. These results support the model that Spn1 preserves nucleosome

positioning over genes, and this function becomes increasingly critical at longer genes.

Subnucleosomal protections are shifted in Spn1 mutant strains

An advantage to the MNase-SSP approach is the retention of small fragments in MNase-

digested samples as previously described (**Figure S1B, File S1**), which can be mapped to the

genome to determine the occupancy and positioning of subnucleosomal protections (Henikoff

et al. 2011; Ramachandran et al. 2017). The abundance of protected fragments 102-112 bp can

be interpreted as subnucleosomal protections with loss of histone-DNA contacts to an H2A-H2B

dimer and the abundance of protected fragments 61-71 bp can be interpreted as

subnucleosomal protections with loss of histone-DNA contacts to both H2A-H2B dimers

(Ramachandran et al. 2017). To test whether Spn1 mutation leads to a shift in the location of

these subnucleosomal protections, the midpoints of protected fragments 102-112 bp and 61-71

bp in length were mapped over nucleosome dyad positions at genes ranked by gene length. As

observed in the V-plots, plotting the midpoints of these fragments demonstrates the existence

of alternative rotational positions around mapped nucleosome dyad positions with 10 bp

periodicity on either side of the dyad (**Figure 4C; Figures S10, S11, File S1**). Occupancy at these

positions can be interpreted as discrete locations of subnucleosome protections with loss of

contacts to an H2A-H2B dimer. In contrast to nucleosome-protected fragments, the relative

occupancy of 102-112 bp and 61-71 bp protected fragments is distributed more equally across

the rotational positions upstream and downstream of the dyad position. For 102-112 bp

fragments, this most probably occurs since loss of contacts to a single H2A-H2B dimer shifts the

location of the midpoint on the DNA protected by a hexasome (Ramachandran et al. 2017). As

transcription through nucleosomes generates hexasomes with transient loss of contacts to both

proximal and distal H2A-H2B dimers, the protected fragments have midpoints shifted on either

side of the dyad that result in a wider distribution of hexasome-occupied rotational positions.

Intriguingly, mapping 61-71 bp protected fragments revealed that rotational positions were

even more equally distributed compared to the positions observed in the 102-112 bp bin of

protected fragments due to the larger shifts in their midpoints.

To quantify and statistically validate the shifts observed in $spn1^{K192N}$ and $spn1^{141-305}$ cells, Z-

scores and p-values were calculated at rotational positions around each mapped dyad location

as described before with gene lists ranked by gene length (**Figure 4D; Figure S12, File S1**). This

demonstrated that protected 102-112 bp fragments are redistributed from upstream rotational

positions to downstream positions in both $spn1^{K192N}$ and $spn1^{141-305}$ cells. As seen before, more

positions had significant changes in occupancy in $spn1^{141-305}$ cells than $spn1^{K192N}$, including shifts

at the +1 and +2 dyad locations seen exclusively in $spn1^{141-305}$ cells, with more positions

experiencing significant changes in 102-112 bp fragment occupancy at longer genes (**Figure**

4D). In contrast to the 102-112 bp fragments, protections 61-71 bp in length in $spn1^{K192N}$ have

minimal changes compared to wild-type, even at the longest genes (**Figure S12, File S1**).

However, 61-71 bp protections in $spn1^{141-305}$ show significant shifts to downstream rotational

positions as a function of gene length. Taken together, while both Spn1 mutants exhibited a

length-dependent downstream shift in rotational positioning preference of subnucleosomal

protections, more positions had significant changes in occupancy in $spn1^{141-305}$ cells than

$spn1^{K192N}$.

Shifts in nucleosome and subnucleosome positioning do not exhibit a clear correlation

with gene expression as assayed by RNAPII occupancy in the Spn1 mutant strains

Spn1 is part of the RNAPII elongation complex, and its occupancy over coding genes is highly

correlated to RNAPII occupancy over those genes (Mayer et al. 2010; Reim et al. 2020;

Viktorovskaya et al. 2021). We next assessed if a relationship exists between gene expression

and the extent of nucleosome positioning defect over gene bodies in the mutant strains. Genes

were ranked by Rpb3 occupancy (a subunit of RNAPII) using published NET-seq data to

generate gene quartiles ranked by gene expression (Churchman and Weissman 2011). Across all

quartiles, the number of genes with nucleosomes positioned at +1 through +10 was relatively

consistent (**Figure S13A, File S1**), indicating gene length will not be a confounding factor in this

analysis. Mapping the sequenced fragments 80-170 bp over genes ranked with the NET-seq

data and generating difference plots shows a slight trend in nucleosome positioning defects

between the lowest and highest quartiles, especially at the promoter-proximal nucleosome

positions (**Figure S13B, File S1**). Additionally, it appears that this effect might be greater for

spn1^{141-305}.

We utilized our quantitative and statistically validated approach to determine whether

nucleosome positioning defects correlate with gene expression as assayed by RNAPII

occupancy. To test this possibility, the midpoints from sequenced fragments 142-152 bp were

mapped to the genome over each quartile ranked by gene expression, using the quartiles

ranked with NET-seq data (**Figure 5A, Figure S14, File S1**). As expected, peaks in 142-152 bp

fragments in the mutant strains occupy the same rotational positions as wild-type cells, but with

changes in occupancy at specific rotational positions. To quantify the shift in nucleosome

positioning over gene bodies, Z-scores across rotational positions spanning -50 to +50 bp

surrounding each nucleosome dyad position were calculated for each mapped dyad for each

gene as described previously. Z-scores and p-values were calculated for the changes in fragment

midpoint occupancy as described previously. This analysis demonstrated that shifts in

nucleosome positioning do not exhibit a linear correlation with expression in the Spn1 mutant

strains (**Figure 5B**). Intriguingly, for *spn1^{141-305}*, the nucleosome positioning shift is more

dramatic than *spn1^{K192N}* with the second, third, and fourth quartiles ranked by expression are

similarly impacted. Given that Spn1 is an elongation factor, we found this result surprising. For

completeness, the midpoints of protected fragments 102-112 bp and 61-71 bp in length were

mapped over nucleosome dyad positions at genes ranked by gene expression (**Figure 5C;**

Figures S15, S16, File S1). Shifts in subnucleosomal protections were quantified with change in

Z-score as described previously for genes in quartiles ranked by expression (**Figure 5D; Figure**

S17, File S1). This analysis demonstrated that shifts in subnucleosome positioning do not

exhibit a correlation with gene expression as assayed by RNAPII occupancy in the Spn1 mutant

strains, similar to nucleosome sized fragments.

Spn1 mutant strains display global reduction in gene expression with an increased effect

at longer genes

To ask if the nucleosome and subnucleosome positioning shifts correlate with changes in gene

expression within the mutant strains, we performed RNA-seq on wild-type and Spn1

mutants (**File S2**). Others have shown that rapid depletion of Spn1 results in global reduction in

RNA levels (Reim et al. 2020), hence we included *S. pombe* cells as a spike-in to query for the

same. We performed principal component analysis of normalized transcript counts from

triplicate samples of wild-type, *spn1^{K192N}*, and *spn1$^{141-305}$* expressing yeast and observed the

replicates clustered together and away from replicates of the other genotypes when PC1 and

PC2 loadings were plotted (**Figure 6A**). Spike-in normalization resulted in *S. pombe* genes

having log$_2$ fold changes (log$_2$FC) distributed tightly around zero, whereas the distribution of

log$_2$FC of *S. cerevisiae* genes were left shifted (**Figures S18A-C, File S1**), indicating a global

reduction in RNA levels. Volcano plots showed that most genes that changed significantly in

spn1^{K192N} and *spn1$^{141-305}$* expressing yeast compared to wild-type were downregulated (**Figures**

6B-D). We combined all genes that significantly changed in at least one of the three

comparisons and grouped them based on the change in patterns of expression (**Figure 6E, File**

S2). Group 1 (n=821) features genes that are significantly downregulated only in *spn1$^{141-305}$*

expressing yeast compared to wild-type. Group 2 (n=280) features genes that are significantly

downregulated in both *spn1^{K192N}* and *spn1$^{141-305}$* expressing yeast compared to wild-type. Group

3 (n=102) consists of genes significantly downregulated only in *spn1^{K192N}* expressing yeast

compared to wild-type. Group 4 (n=61) consists of genes significantly downregulated in *spn1^{141-}*

^{305} expressing yeast compared to both wild-type and *spn1^{K192N}* expressing yeast. The rest of the

genes belong to other combinations of changes that are <20 genes per group.

Our MNase-seq analysis had shown downstream nucleosome shifts that positively correlated

with increasing gene length and not with genes ranked by RNAPII occupancy. We asked if the

gene expression changes measured by RNA-seq have a similar correlation. Hence, we

intersected lists of genes in the four groups we defined with quartiles of genes based on length

and RNAPII occupancy. Strikingly, we observed a significant depletion of short genes and an

enrichment of long genes in groups 1, 2, and 4 (**Figure 6E**). Intriguingly, group 3, which consists

of genes that are downregulated only in *spn1^{K192N}* expressing yeast compared to wild-type did

not show an enrichment for long genes. Instead, group 3 genes show a clear enrichment for the

highest quartile ranked by RNAPII occupancy (**Figure 6F**). Group 2 genes also show an

enrichment for the highest quartile of RNAPII occupancy, indicating that $spn1^{K192N}$ expressing

yeast have increased defects in genes with the highest gene expression quartile. We next

performed gene set enrichment analysis of genes belonging to groups 1-4 and found highly

significant overlap with terms related to metal ion transport, glycolysis, and ribosome biogenesis

among others (**Figure S18D, File S1**).

Taken together, most genes significantly downregulated in $spn1^{141-305}$ or both $spn1^{141-305}$ and

$spn1^{K192N}$ expressing yeast are enriched for long genes, which also correlates with the observed

downstream shifts in nucleosome positioning observed for long genes. Furthermore, genes

significantly downregulated in $spn1^{K192N}$ expressing yeast are enriched in the highest quartile of

expression, which also exhibited increased downstream shifts in nucleosome positioning.

DISCUSSION

In this study, an extensive mapping of chromatin protections in yeast expressing mutant alleles

with distinct defects for known binding partners was performed to assess the role of this

essential histone chaperone in maintaining chromatin structure. By retaining fragments of a

wide range of sizes, the MNase-SSP approach demonstrates that Spn1 preserves nucleosomal,

subnucleosomal, and non-nucleosomal protections across the yeast genome. Our analyses

revealed that this function of Spn1 maintains chromatin over NDRs, the 5' ends of genes, and

within gene bodies.

The role of Spn1 in the maintenance of NDR architecture was previously unknown. While

chromatin defects in Spn1 mutants have been primarily characterized over gene bodies during

transcription elongation (Reim et al. 2020; Viktorovskaya et al. 2021), the $spn1^{K192N}$ mutation

clearly reduces subnucleosomal protections in the NDR, most notably for very short protections

(<60bp). How this mutation leads to this defect is unclear, although, the $spn1^{K192N}$ mutant is

known to be defective for binding to RNAPII, likely in part through disruption of interactions

with Spt5 and Spt6 (Ehara et al. 2022; Zhang et al. 2008). Also, previous work with mutants that

disrupt the Spn1-Spt6 complex has implicated the complex in monitoring nucleosomes and

weakening the Spn1-Spt6 interaction has impacts on the assembly and maintenance of

chromatin (McCullough et al. 2015; Viktorovskaya et al. 2021). Additionally, compromise of Spt6

function has been shown to produce aberrations in chromatin over gene bodies, namely the

misincorporation of histone variant H2A.Z (McCullough et al. 2015; Viktorovskaya et al. 2021).

Our findings support the notion that surveillance of proper chromatin structure by the Spn1-

Spt6 complex may extend to the NDRs.

Both $spn1^{K192N}$ and $spn1^{141-305}$ produced a downstream shift in nucleosome positioning over

gene bodies. However, $spn1^{141-305}$ exclusively exhibited defects in the positioning of the +1 and

+2 nucleosomes. This finding reveals that the mechanism of chromatin maintenance at the 5'

ends of genes may be functionally distinct from that over the rest of the gene and that the

histone- and nucleosome-binding functions of Spn1 are part of this mechanism. As $spn1^{K192N}$ did

not produce significant shifts in nucleosome or subnucleosome positioning at the 5' ends of

genes, one possible interpretation is that the mechanism of chromatin maintenance at the

promoter-proximal region of genes requires Spn1 to directly bind chromatin, whose binding

regions are unaffected in $spn1^{K192N}$. In contrast, maintenance of chromatin over the rest of the

body of the gene requires both Spn1 recruitment to the RNAPII complex and its histone- and

nucleosome-binding capabilities, as evidenced by defects in nucleosome and subnucleosome

positioning at nucleosomes distal from the promoter in both mutants.

By mapping the fragment midpoints, we utilized our dataset to demonstrate that shifts in

nucleosome positioning in Spn1 mutants occur in discrete steps at alternate favored rotational

positions. By quantifying the occupancy of nucleosomes at each rotational position, we present

a method for statistically validating shifts in nucleosome positioning. Mapping the midpoints of

smaller subnucleosomal protections also provided *in vivo* evidence suggesting the existence of

alternate rotational positions occupied by subnucleosomal intermediates. These positions also

occurred in discrete steps of 10 bp on either side of the dyad, and occupancy of these shorter

protected fragments were distributed more evenly across the positions surrounding the mapped

nucleosome dyad position. This approach of quantifying rotational positioning may be useful for

other datasets of sufficiently high sequencing depth in organisms with small genomes like *S.*
cerevisiae.

Quantification and statistical validation of change in preference of rotational positions at each
nucleosome position of each gene allowed us to assess the relationship between the shifts in
nucleosomal and subnucleosomal protections in Spn1 mutant expressing yeast and genes
ranked by expression or length. As Spn1 is part of the RNAPII elongation complex, we assessed
whether defects in chromatin structure observed in Spn1 were more dramatic at highly
expressed genes as ranked by RNAPII occupancy. Surprisingly, our findings indicate that the
relationship between transcription and changes in chromatin structure induced by Spn1
mutation were not correlated. However, Spn1 maintenance of chromatin structure became
increasingly critical with gene length, as demonstrated by an increase in rotational positions with
statistically significant changes in occupancy from the lowest quartile to the highest quartile of
genes ranked by length. This finding provides novel insights into how chromatin structure is
maintained at genes with varied length. In Spn1 mutant expressing yeast, the longest genes
exhibited significant shifts in nucleosomal and subnucleosomal protections even at the most
distal nucleosome positions assessed. For shorter genes, there were few significant changes in
nucleosomal and subnucleosomal rotational position occupancies, even at nucleosome
positions with significant changes in the longer quartiles of genes, such as the +3 and +4
positions.

To ascertain whether changes in chromatin overlap with the set of genes differentially expressed
in the Spn1 mutant strains, we performed RNA-seq on wild-type and Spn1 mutant strains. This
experiment revealed that yeast expressing $spn1^{K192N}$ or $spn1^{141-305}$ produce a global reduction in
gene expression, with over 400 genes significantly downregulated in $spn1^{K192N}$ and nearly 1200
genes significantly downregulated in $spn1^{141-305}$. Remarkably, we observed that these
differentially expressed genes were highly enriched for long genes, the same class of genes
which exhibited the largest shifts in nucleosome positioning. Additionally, we observed that
genes downregulated in $spn1^{K192N}$ expressing yeast are enriched in highly expressed genes,
which also exhibited increased shifts in nucleosome positioning.

The requirement for Spn1 in preserving nucleosome positioning shares features with the

functional requirements for Spt4 and Spt6 in maintaining chromatin structure (Uzun et al. 2021;

Viktorovskaya et al. 2021). Each of these factors are physically associated with RNAPII during

transcription elongation, with Spn1 and Spt4 positioned at the tip of the RNAPII clamp facing

the downstream DNA (Uzun et al. 2021; Viktorovskaya et al. 2021). In contrast, the main body of

Spt6 is docked near the upstream DNA exit channel, although, the Spt6 N-terminal helices that

bind Spn1 are proximal to the downstream DNA (Ehara et al. 2022). Spt6 binds histones and

nucleosomes when the HMG-box-like protein Nhp6 is present (Ehara et al. 2022). Spt6 is

commonly classified as a histone chaperone necessary for nucleosome reassembly in the wake

of RNAPII, supported by its proximity to upstream DNA and its ascribed ability to suppress

intragenic transcription (Doris et al. 2018; Kaplan et al. 2003; Van Bakel et al. 2013). While Spt4

has not been shown to bind chromatin, it oscillates on and off RNAPII along with its binding

partner Spt5 based on nucleosome positions, and its deletion produces a downstream

nucleosome positioning shift. Moreover, Spt5 deletion produces increased stalling and

backtracking during early transcription elongation (Uzun et al. 2021). Despite differences in their

transcription-associated functions, Spn1, Spt4, and Spt5 are all critical for maintaining proper

chromatin structure and their relationships are further supported in that defects in each

produced a similar pattern of downstream shifts in nucleosome positioning. One explanation for

this commonality is that all participate in this process of co-transcriptional nucleosome

disassembly and reassembly, and a defect in any step of this process produces a similar defect in

chromatin. Others have posed that this defect in passage through nucleosomes may produce an

elongation complex that transcribes through chromatin more slowly than in wild-type cells,

which might produce mispositioned nucleosomes (Uzun et al. 2021). Indeed, there is evidence

that loss of RNAPII transcription results in shifts in nucleosome positions, as depletion of Rpb1

(Tramantano et al. 2016) and growth of the temperature-sensitive *rpb1-1* allele at a non-

permissive temperature (Tramantano et al. 2016), produces a downstream nucleosome shift.

Maintenance of nucleosome positioning is also associated with ATP-dependent nucleosome

remodelers, which position nucleosomes and evict or exchange histones in an ATP-dependent

manner (reviewed in (Clapier et al. 2017)). Recent work demonstrated that the ISW1a

nucleosome remodeler complex subunit *Ioc4* mutant or depletion produces a downstream shift

in nucleosome positioning comparable to the shift we observed in the Spn1 mutants (Bhardwaj

et al. 2020). However, other MNase-seq experiments demonstrate that depletion or deletion of

nucleosome remodeling factors, notably Isw1, Chd1, Ino80, and Sth1 produce upstream shifts in

nucleosome positioning over gene bodies (Klein-Brill et al. 2019; Ocampo et al. 2016). Spn1

does collaborate with the SWI/SNF nucleosome remodeler complex, as we've previously

observed through a significant genetic interaction with *SNF2*, which encodes the catalytic

subunit of SWI/SNF (Zhang et al. 2008).Intriguingly, degradation of Snf2 has little effect on

nucleosome positioning (Klein-Brill et al. 2019). Taken together, the shifts observed in the Spn1

mutants more closely resemble the shifts produced by Spt4, Spt6, and Rpb1 rather than most

nucleosome remodelers.

A puzzling question that remains is how *spn1^{K192N}* and *spn1$^{141-305}$* produce such similar changes

in chromatin protections over nucleosome positions distal to promoters despite their distinct

defects for known binding partners. Both Spn1 recruitment to the elongating RNAPII complex

and its ability to bind histones and/or nucleosomes are critical for preserving chromatin

structure, and likely occur co-transcriptionally. Wild-type Spn1 is poised near the downstream

nucleosome in the RNAPII elongation complex (Ehara et al. 2022). Spn1 directly binds histones

and nucleosomes, a function that may be important for the disassembly of downstream

nucleosomes during transcription. If the recruitment of Spn1 to sites of transcription is

diminished, the process of disassembling the downstream nucleosome may be impacted.

Conversely, the *spn1$^{141-305}$* mutant is likely still recruited to sites of transcription. However, its loss

of histone- and nucleosome-binding regions renders it unable to participate in the process of

downstream nucleosome disassembly. In either mutant, the loss of Spn1 function results in

defects in chromatin preservation during transcription elongation, resulting in a shift in

nucleosome and subnucleosome protections. Taken together, these findings deepen our

understanding of Spn1-dependent chromatin maintenance over the yeast genome and provide

insight into mechanisms underlying this process.

AUTHOR CONTRIBUTIONS

Andrew J. Tonsager: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing—original draft. **Alexis Zukowski**: Formal analysis, Methodology, Investigation. **Catherine A. Radebaugh**: Conceptualization, Formal analysis, Investigation, Methodology, Visualization. **Abigail Weirich**: Investigation. **Laurie A. Stargell**: Conceptualization, Funding Acquisition, Project administration, Supervision, Writing—review & editing. **Srinivas Ramachandran**: Conceptualization, Formal analysis, Funding Acquisition, Investigation, Methodology, Project administration, Supervision, Visualization, Writing—review & editing.

FUNDING

This work was supported by the RNA Bioscience Initiative, the University of Colorado School of Medicine, and NIH grant R35GM133434. S. R. is a Pew-Stewart Scholar for Cancer Research, supported by the Pew Charitable Trusts and the Alexander and Margaret Stewart Trust.

CONFLICT OF INTEREST

The authors declare that they have no known competing financial interests or connections, direct or indirect, that could have appeared to influence the work reported in this paper.

ACKNOWLEDGEMENTS

We thank Sarah Swygert for her expert technical assistance.

REFERENCES

Adkins NL, Niu H, Sung P, Peterson CL. 2013. Nucleosome dynamics regulates DNA processing. Nature Structural & Molecular Biology. 20(7):836-842.

Bai L, Morozov AV. 2010. Gene regulation by nucleosome positioning. Trends in Genetics. 26(11):476-483.

Bevington S, Boyes J. 2013. Transcription-coupled eviction of histones h2a/h2b governs v(d)j recombination. The EMBO Journal. 32(10):1381-1392.

Bhardwaj SK, Hailu SG, Olufemi L, Brahma S, Kundu S, Hota SK, Persinger J, Bartholomew B. 2020. Dinucleosome specificity and allosteric switch of the isw1a atp-dependent chromatin remodeler in transcription regulation. Nat Commun. 11(1):5913.

Brahma S, Henikoff S. 2020. Epigenome regulation by dynamic nucleosome unwrapping. Trends in Biochemical Sciences. 45(1):13-26.

Brogaard K, Xi L, Wang J-P, Widom J. 2012. A map of nucleosome positions in yeast at base-pair resolution. Nature. 486(7404):496-501.

Brower-Toland BD, Smith CL, Yeh RC, Lis JT, Peterson CL, Wang MD. 2002. Mechanical disruption of individual nucleosomes reveals a reversible multistage release of DNA. Proceedings of the National Academy of Sciences. 99(4):1960-1965.

Cermakova K, Demeulemeester J, Lux V, Nedomova M, Goldman SR, Smith EA, Srb P, Hexnerova R, Fabry M, Madlikova M et al. 2021. A ubiquitous disordered protein interaction module orchestrates transcription elongation. Science. 374(6571):1113-1121.

Chereji RV, Ocampo J, Clark DJ. 2017. Mnase-sensitive complexes in yeast: Nucleosomes and non-histone barriers. Molecular Cell. 65(3):565-577.e563.

Chereji RV, Ramachandran S, Bryson TD, Henikoff S. 2018. Precise genome-wide mapping of single nucleosomes and linkers in vivo. Genome Biology. 19(1):19.

Churchman LS, Weissman JS. 2011. Nascent transcript sequencing visualizes transcription at nucleotide resolution. Nature. 469(7330):368-373.

Clapier CR, Iwasa J, Cairns BR, Peterson CL. 2017. Mechanisms of action and regulation of atp-dependent chromatin-remodelling complexes. Nat Rev Mol Cell Biol. 18(7):407-422.

Davey CA, Sargent DF, Luger K, Maeder AW, Richmond TJ. 2002. Solvent mediated interactions in the structure of the nucleosome core particle at 1.9å resolution. Journal of Molecular Biology. 319(5):1097-1113.

Doris SM, Chuang J, Viktorovskaya O, Murawska M, Spatt D, Churchman LS, Winston F. 2018. Spt6 is required for the fidelity of promoter selection. Molecular Cell. 72(4):687-699.e686.

Eaton ML, Galani K, Kang S, Bell SP, MacAlpine DM. 2010. Conserved nucleosome positioning defines replication origins. Genes & Development. 24(8):748-753.

Ehara H, Kujirai T, Shirouzu M, Kurumizaka H, Sekine S-i. 2022. Structural basis of nucleosome disassembly and reassembly by rnapii elongation complex with fact. Science. 377(6611):eabp9466.

Fischbeck JA, Kraemer SM, Stargell LA. 2002. *Spn1* , a conserved gene identified by suppression of a postrecruitment-defective yeast tata-binding protein mutant. Genetics. 162(4):1605-1616.

Flaus A, Luger K, Tan S, Richmond TJ. 1996. Mapping nucleosome position at single base-pair resolution by using site-directed hydroxyl radicals. Proceedings of the National Academy of Sciences. 93(4):1370-1375.

Gansauge M-T, Meyer M. 2013. Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. Nature Protocols. 8(4):737-748.

Gavin A-C, Bösche M, Krause R, Grandi P, Marzioch M, Bauer A, Schultz J, Rick JM, Michon A-M, Cruciat
C-M et al. 2002. Functional organization of the yeast proteome by systematic analysis of protein
complexes. Nature. 415(6868):141-147.
Groth A, Rocha W, Verreault A, Almouzni G. 2007. Chromatin challenges during DNA replication and
repair. Cell. 128(4):721-733.
Henikoff JG, Belsky JA, Krassovsky K, MacAlpine DM, Henikoff S. 2011. Epigenome characterization at
single base-pair resolution. Proceedings of the National Academy of Sciences. 108(45):18318-
18323.
Iyer V, Struhl K. 1996. Absolute mrna levels and transcriptional initiation rates in saccharomyces
cerevisiae. Proc Natl Acad Sci U S A. 93(11):5208-5212.
Kaplan CD, Laprade L, Winston F. 2003. Transcription elongation factors repress transcription initiation
from cryptic sites. Science. 301(5636):1096-1099.
Kent NA, Adams S, Moorhouse A, Paszkiewicz K. 2011. Chromatin particle spectrum analysis: A method
for comparative chromatin structure analysis using paired-end mode next-generation DNA
sequencing. Nucleic Acids Research. 39(5):e26-e26.
Klein-Brill A, Joseph-Strauss D, Appleboim A, Friedman N. 2019. Dynamics of chromatin and transcription
during transient depletion of the rsc chromatin remodeling complex. Cell Rep. 26(1):279-292
e275.
Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. 2023. G:Profiler-interoperable web service
for functional enrichment analysis and gene identifier mapping (2023 update). Nucleic Acids Res.
51(W1):W207-W212.
Krogan NJ, Kim M, Ahn SH, Zhong G, Kobor MS, Cagney G, Emili A, Shilatifard A, Buratowski S, Greenblatt
JF. 2002. Rna polymerase ii elongation factors of *saccharomyces cerevisiae* : A targeted
proteomics approach. Molecular and Cellular Biology. 22(20):6979-6992.
Kubik S, Bruzzone Maria J, Jacquet P, Falcone J-L, Rougemont J, Shore D. 2015. Nucleosome stability
distinguishes two different promoter types at all protein-coding genes in yeast. Molecular Cell.
60(3):422-434.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Subgroup
GPDP. 2009. The sequence alignment/map format and samtools. Bioinformatics. 25(16):2078-
2079.
Li S, Almeida AR, Radebaugh CA, Zhang L, Chen X, Huang L, Thurston AK, Kalashnikova AA, Hansen JC,
Luger K et al. 2018. The elongation factor spn1 is a multi-functional chromatin binding protein.
Nucleic Acids Research. 46(5):2321-2334.
Li S, Edwards G, Radebaugh CA, Luger K, Stargell LA. 2022. Spn1 and its dynamic interactions with spt6,
histones and nucleosomes. Journal of Molecular Biology. 434(13):167630.
Lindstrom DL, Squazzo SL, Muster N, Burckin TA, Wachter KC, Emigh CA, McCleery JA, Yates JR, Hartzog
GA. 2003. Dual roles for spt5 in pre-mrna processing and transcription elongation revealed by
identification of spt5-associated proteins. Molecular and Cellular Biology. 23(4):1368-1378.
Lorch Y, Maier-Davis B, Kornberg RD. 2014. Role of DNA sequence in chromatin remodeling and the
formation of nucleosome-free regions. Genes & Development. 28(22):2492-2497.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for rna-seq data
with deseq2. Genome Biol. 15(12):550.
Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. 1997. Crystal structure of the nucleosome
core particle at 2.8 å resolution. Nature. 389(6648):251-260.
Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnetjournal. 17(1):10.

Mayer A, Lidschreiber M, Siebert M, Leike K, Söding J, Cramer P. 2010. Uniform transitions of the general
812   rna polymerase ii transcription complex. Nature Structural & Molecular Biology. 17(10):1272-
813   1278.

814   McCullough L, Connell Z, Petersen C, Formosa T. 2015. The abundant histone chaperones spt6 and fact
815   collaborate to assemble, inspect, and maintain chromatin structure in *saccharomyces cerevisiae*.
816   Genetics. 201(3):1031-1045.

817   McDonald SM, Close D, Xin H, Formosa T, Hill CP. 2010. Structure and biological importance of the spn1-
818   spt6 interaction, and its regulatory role in nucleosome binding. Molecular Cell. 40(5):725-735.

819   Ocampo J, Chereji RV, Eriksson PR, Clark DJ. 2016. The isw1 and chd1 atp-dependent chromatin
820   remodelers compete to set nucleosome spacing in vivo. Nucleic Acids Res. 44(10):4625-4635.

821   Park D, Morris AR, Battenhouse A, Iyer VR. 2014. Simultaneous mapping of transcript ends at single-
822   nucleotide resolution and identification of widespread promoter-associated non-coding rna
823   governed by tata elements. Nucleic Acids Research. 42(6):3736-3749.

824   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-aware
825   quantification of transcript expression. Nat Methods. 14(4):417-419.

826   Polo SE, Almouzni G. 2006. Chromatin assembly: A basic recipe with various flavours. Current Opinion in
827   Genetics & Development. 16(2):104-111.

828   Quinlan AR, Hall IM. 2010. Bedtools: A flexible suite of utilities for comparing genomic features.
829   Bioinformatics. 26(6):841-842.

830   Ramachandran S, Ahmad K, Henikoff S. 2017. Transcription and remodeling produce asymmetrically
831   unwrapped nucleosomal intermediates. Molecular Cell. 68(6):1038-1053.e1034.

832   Ramani V, Qiu R, Shendure J. 2019. High sensitivity profiling of chromatin structure by mnase-ssp. Cell
833   Reports. 26(9):2465-2476.e2464.

834   Rando OJ, Winston F. 2012. Chromatin and transcription in yeast. Genetics. 190(2):351-387.

835   Reim NI, Chuang J, Jain D, Alver BH, Park PJ, Winston F. 2020. The conserved elongation factor spn1 is
836   required for normal transcription, histone modifications, and splicing in saccharomyces
837   cerevisiae. Nucleic Acids Research. 48(18):10241-10258.

838   Rhee Ho S, Bataille Alain R, Zhang L, Pugh BF. 2014. Subnucleosomal structures and nucleosome
839   asymmetry across a genome. Cell. 159(6):1377-1388.

840   Rossi MJ, Kuntala PK, Lai WKM, Yamada N, Badjatia N, Mittal C, Kuzu G, Bocklund K, Farrell NP, Blanda TR
841   et al. 2021. A high-resolution protein architecture of the budding yeast genome. Nature.
842   592(7853):309-314.

843   Satchwell SC, Drew HR, Travers AA. 1986. Sequence periodicities in chicken nucleosome core DNA.
844   Journal of Molecular Biology. 191(4):659-675.

845   Segal E, Fondufe-Mittendorf Y, Chen L, Thåström A, Field Y, Moore IK, Wang J-PZ, Widom J. 2006. A
846   genomic code for nucleosome positioning. Nature. 442(7104):772-778.

847   Selth LA, Sigurdsson S, Svejstrup JQ. 2010. Transcript elongation by rna polymerase ii. Annual Review of
848   Biochemistry. 79(1):271-293.

849   Snyder Matthew W, Kircher M, Hill Andrew J, Daza Riza M, Shendure J. 2016. Cell-free DNA comprises an
850   in vivo nucleosome footprint that informs its tissues-of-origin. Cell. 164(1-2):57-68.

851   Thurston AK, Radebaugh CA, Almeida AR, Argueso JL, Stargell LA. 2018. Genome instability is promoted
852   by the chromatin-binding protein spn1 in *saccharomyces cerevisiae*. Genetics. 210(4):1227-1237.

853   Tramantano M, Sun L, Au C, Labuz D, Liu Z, Chou M, Shen C, Luk E. 2016. Constitutive turnover of histone
854   h2a.Z at yeast promoters requires the preinitiation complex. eLife. 5:e14243.

855   Uzun Ü, Brown T, Fischl H, Angel A, Mellor J. 2021. Spt4 facilitates the movement of rna polymerase ii
856   through the +2 nucleosomal barrier. Cell Reports. 36(13):109755.

Van Bakel H, Tsui K, Gebbia M, Mnaimneh S, Hughes TR, Nislow C. 2013. A compendium of nucleosome
and transcript profiles reveals determinants of chromatin architecture and transcription. PLoS
Genetics. 9(5):e1003479.
Viktorovskaya O, Chuang J, Jain D, Reim NI, López-Rivera F, Murawska M, Spatt D, Churchman LS, Park PJ,
Winston F. 2021. Essential histone chaperones collaborate to regulate transcription and
chromatin integrity. Genes & Development. 35(9-10):698-712.
Zhang L, Fletcher AGL, Cheung V, Winston F, Stargell LA. 2008. Spn1 regulates the recruitment of spt6
and the swi/snf complex during transcriptional activation by rna polymerase ii. Molecular and
Cellular Biology. 28(4):1393-1403.
Zhang Z, Reese JC. 2005. Isolation of yeast nuclei and micrococcal nuclease mapping of nucleosome
positioning. Yeast protocols. New Jersey: Humana Press. p. 245-256.

FIGURES LEGENDS

Figure 1. An extensive characterization of fragments protected from MNase digestion

suggests shifts in chromatin protections in Spn1 mutant yeast across fragment lengths. (A)

Nuclei harvested from two biological replicates of wild-type, (B) $spn1^{K192N}$, and (C) $spn1^{141-305}$

expressing yeast grown to log phase were digested with 128U MNase and protected fragments

were isolated and sequenced. V-plots were produced by mapping fragment midpoints to the

yeast genome anchored at +1, +2, and +5 nucleosome dyad positions from 4688 genes.

Figure 2. The $spn1^{K192N}$ mutation reduces the stability of nucleosome-depleted regions. (A)

Profiles of MNase protected fragments (0-60 bp) mapped over 4688 genes aligned to the +1

nucleosome dyad in wild-type, $spn1^{K192N}$, and $spn1^{141-305}$ expressing yeast. (B) Profiles of MNase

protected fragments (0-60 bp) from low MNase-digested samples (32U) mapped over 4688

genes aligned to the +1 nucleosome dyad in wild-type, $spn1^{K192N}$, and $spn1^{141-305}$ expressing

yeast. (C) Difference heatmap between wild-type and mutant Spn1 expressing yeast strains

scored by changes in occupancy of 0-60 bp fragments. Genes are aligned by their +1

nucleosome dyad position and ranked by length. Scores are calculated by subtracting wild-type

occupancy data from mutant occupancy. Regions in red are enriched for fragments at that

position while blue regions are depleted for fragments.

Figure 3. Mutations in Spn1 produce a downstream shift in nucleosome positions over

open reading frames genome wide. (A) Difference heatmap between wild-type and mutant

Spn1 expressing yeast strains scored by changes in nucleosome occupancy. Genes are aligned

by their +1 nucleosome dyad position and ranked by length. Scores are calculated by

subtracting wild-type nucleosome occupancy data from mutant nucleosome occupancy.

Regions in red are enriched for nucleosomes at that position while blue regions are depleted for

nucleosomes. (B) Profiles of MNase protected fragments (80-170 bp) mapped over 4688 genes

aligned to the +1 nucleosome dyad in wild-type, $spn1^{K192N}$, and $spn1^{141-305}$ expressing yeast. (C)

Profiles of MNase protected fragments (80-170 bp) aligned to the +1 nucleosome dyad in wild-

type, $spn1^{K192N}$, and $spn1^{141-305}$ expressing yeast highlighting the +5 nucleosome position.

Figure 4. Quantification of nucleosome and subnucleosome positioning shifts in Spn1-

mutant cells reveals increasing defects with increasing gene length. (A) Metagene plots of

fragment midpoints 142-152 bp in length mapped to the +5 nucleosome dyad at gene quartiles

ranked by length. Asterisks indicate rotational positions with statistically significant change in Z-

score. (B) Z-score changes in 142-152 bp fragment occupancy at rotational positions

surrounding each nucleosome dyad position at genes ranked by length in mutant Spn1

expressing yeast strains compared to wild-type plotted as a heatmap. (C) Metagene plots of

fragment midpoints 102-112 bp in length mapped to the +5 nucleosome dyad at genes quartile

ranked by length. Asterisks indicate rotational positions with statistically significant change in Z-

score. (D) Z-score changes in 102-112 bp fragment occupancy at rotational positions

surrounding each nucleosome dyad position at genes ranked by length in mutant Spn1

expressing yeast strains compared to wild-type plotted as a heatmap.

Figure 5. Spn1 maintains nucleosomal and subnucleosomal positioning at actively

transcribed genes. (A) Metagene plots of fragment midpoints 142-152 bp in length mapped to

the +5 nucleosome dyad at gene quartiles ranked by expression using published NET-seq data

(38). Asterisks indicate rotational positions with statistically significant change in Z-score. (B) Z-

score changes in 142-152 bp fragment occupancy at rotational positions surrounding each

nucleosome dyad position at genes ranked by expression in mutant Spn1 expressing yeast

strains compared to wild-type plotted as a heatmap. (C) Metagene plots of fragment midpoints

102-112 bp in length mapped to the +5 nucleosome dyad at genes quartile ranked by

expression. Asterisks indicate rotational positions with statistically significant change in Z-score.

(D) Z-score changes in 102-112 bp fragment occupancy at rotational positions surrounding each

nucleosome dyad position at genes ranked by expression in mutant Spn1 expressing yeast

strains compared to wild-type plotted as a heatmap.

Figure 6. Spn1 mutants have global defects in gene expression with increased effects on

longer and highly expressed genes. (**A**) Loadings of the first two principal components from

the principal component analysis of normalized gene count matrix shows concordance between

replicates and a clear separation between *wild-type*, *spn1^{K192N}*, and *spn1$^{141-305}$* expressing yeast.

(**B**) Volcano plot where the Log$_2$(Fold Change) for *spn1^{K192N}* expressing yeast compared to wild-

type is plotted against -log$_{10}$(FDR). Genes with Log$_2$(Fold Change) >0 and FDR ≤ 0.05 are

marked in red and genes with Log$_2$(Fold Change) <0 and FDR ≤ 0.05 are marked in blue. (**C**)

Same as (**B**) for *spn1$^{141-305}$* expressing yeast compared to wild-type. (**D**) Same as (**B**) for *spn1$^{141-305}$*

expressing yeast compared to *spn1^{K192N}* expressing yeast. **E**) Heatmap of log$_2$(Fold Change)

across three possible comparisons (*spn1^{K192N}* vs. wild-type, *spn1$^{141-305}$* vs. wild-type, and *spn1^{K192N}*

vs. *spn1$^{141-305}$* for all genes that have adjusted p-value<0.05 for at least one of the three

comparisons. The genes are grouped (G1-G4) based on which comparisons yield significant

down- or up-regulation. The rest of the genes (n=90) belong to other combinations of changes

that are <20 genes per group. (**F**) Log$_2$ enrichment of the overlap of gene groups defined in (D)

with gene quartiles ranked by length compared to background enrichment. The statistical

significance of the enrichment for each overlap was determined using hypergeometric test in R

followed by multiple testing correction using the Benjamini and Hochberg method and overlaps

with adjusted p-value<0.05 are marked with an asterisk. (**G**) Same as (E) for gene quartiles

ranked by expression using published NET-seq data.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6