A random mutagenesis screen enriched for missense mutations in bacterial effector

proteins.

Malene L. Urbanus[1], Thomas M. Zheng[1], Anna N. Khusnutdinova[2], Doreen Banh[1] Harley

O'Connor Mount[3], Alind Gupta[3], Peter J. Stoigos[2], Alexei Savchenko[2,4], Ralph R. Isberg[5],

Alexander F. Yakunin[2,6], and Alexander W. Ensminger[1,3]*

Affiliations

[1]Department of Biochemistry, University of Toronto, Toronto, ON, Canada.

[2]Department of Chemical Engineering and Applied Chemistry, University of Toronto, ON,

Canada.

[3]Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada.

[4]Department of Microbiology, Immunology & Infectious Diseases, Health Research Innovation

Centre, University of Calgary, AB, Canada.

[5]Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

Boston, MA, USA

[6]Centre for Environmental Biotechnology, School of Natural Sciences, Bangor University,

Bangor, UK.

*Corresponding author:

Email: alex.ensminger@utoronto.ca

Abstract

To remodel their hosts and escape immune defenses, many pathogens rely on large arsenals of

proteins (effectors) that are delivered to the host cell using dedicated translocation machinery.

Effectors hold significant insight into the biology of both the pathogens that encode for them and

the host pathways that they manipulate. One of the most powerful systems biology tools for

studying effectors is the model organism, *Saccharomyces cerevisiae*. For many pathogens, the

heterologous expression of effectors in yeast is growth inhibitory at a frequency much higher

than housekeeping genes, an observation ascribed to targeting conserved eukaryotic proteins.

Abrogation of yeast growth inhibition has been used to identify bacterial suppressors of effector

activity, host targets, and functional residues and domains within effector proteins. We present

here a yeast-based method for enriching for informative, in-frame, missense mutations in a pool

of random effector mutants. We benchmark this approach against three effectors from *Legionella*

pneumophila, an intracellular bacterial pathogen that injects a staggering >330 effectors into the

host cell. For each protein, we show how *in silico* protein modeling (AlphaFold2) and missense-

directed mutagenesis can be combined to reveal important structural features within effectors.

We identify known active site residues within the metalloprotease RavK, highly conserved

residues in SdbB, and previously unidentified functional motifs within the C-terminal domain of

SdbA. We show that this domain has structural similarity with glycosyltransferases and exhibits

in vitro activity consistent with this predicted function.

Introduction

For many bacterial pathogens, host manipulation derives from the collective activity of large

numbers of translocated proteins ("effectors") that are injected into the host cell using dedicated

secretion machinery. A striking example is the gram-negative bacterium, *Legionella*

pneumophila, the causative agent of Legionnaires' disease, an often-fatal severe pneumonia that

naturally replicates in freshwater protozoa. *L. pneumophila* encodes for the largest effector

arsenal described to date (>330 effectors per isolate, or roughly 10% of the proteome) (Burstein

et al. 2009; Huang *et al.* 2011; Zhu *et al.* 2011), which is injected into the host cell using the

Dot/Icm type IVB secretion system (Segal *et al.* 1998; Vogel *et al.* 1998). *L. pneumophila*

effectors modulate host vesicle trafficking, post-translational modification, protein translation,

autophagy, vacuolar function, and the cytoskeleton to avoid lysosomal fusion and to establish a

replicative, neutral pH vacuole (Isberg *et al.* 2009; Escoll *et al.* 2013; Sherwood and Roy 2016;

Qiu and Luo 2017). While over 50 effectors have been studied, the majority of these remain

uncharacterized (Finsel and Hilbi 2015).

Determining the function of the >330 effectors and the role they play in establishing the

Legionella-containing vacuole is complicated by extensive genetic redundancy within the

effector arsenal (O'Connor *et al.* 2011) and the lack of predicted conserved domains or functions

for many substrates (Gomez-Valero *et al.* 2011, 2014; Burstein *et al.* 2016). In a study

comparing genomes from 38 *Legionella* species, only half of the predicted effectors contained

conserved domains and many of these are of uncharacterized function (Burstein *et al.* 2016).

Although the amino acid sequence of many effectors may not yield obvious clues to their

function, some effectors have structural homology to characterized proteins or domains, along

with conserved active site motifs or other signature motifs (Toulabi *et al.* 2013; Morar *et al.*

2015; Wong *et al.* 2015; Urbanus *et al.* 2016; Pinotsis and Waksman 2017; Kozlov *et al.* 2018).

Looking beyond *L. pneumophila*, over 18,000 effector genes have been predicted across the

entire *Legionella* genus (Burstein *et al.* 2016; Gomez-Valero *et al.* 2019). This number suggests

both a wealth of novel effector activities and host biology waiting to be discovered.

We set out to develop a method to efficiently identify important motifs or amino acid

residues in uncharacterized *Legionella* effectors by random mutagenesis and selection for loss-

of-function mutations to facilitate the prediction of mechanism and function. Approximately

10% of the effectors severely inhibit yeast growth when overexpressed (Campodonico *et al.*

2005; Felipe *et al.* 2008; Heidtman *et al.* 2009; Shen *et al.* 2009; Guo *et al.* 2014; Urbanus *et al.*

2016), such that loss-of-function by random mutagenesis can be selected for as an alleviation of

the yeast growth defect. However, a random mutant pool contains a multitude of mutations that

can potentially cause a loss-of-function phenotype, such as frameshift, nonsense and missense

mutations in the effector or regulatory elements such as the promoter region. Most of these

mutations are uninformative as only missense loss-of-function mutations in the effector are

relevant. To enrich for full-length missense clones, we used a C-terminal in-frame His3 fusion

and selected for histidine prototrophic yeast requiring the presence of a full-length protein. A

similar strategy (C-terminal His3 fusions) was previously shown to enrich for in-frame human

open reading frames amongst a randomly primed pool of cDNAs cloned into a yeast expression

vector (Holz *et al.* 2001). Here, we benchmark this in-frame mutagenesis approach against three

L. pneumophila effectors previously shown to inhibit yeast growth: SdbA and SdbB, whose

functions remain uncharacterized and RavK, a previously described metalloprotease. We show

our approach identifies active site residues within RavK (Liu *et al.* 2017) highly conserved

residues in SdbB, and previously unidentified functional motifs in the C-terminal domain of

SdbA. These motifs are part of the donor and acceptor binding regions of glycosyltransferases,

which we show the C-terminal domain of SdbA shares homology with. Finally, we show that a

C-terminal fragment exhibits *in vitro* activity consistent with this predicted function.

Results

A random mutagenesis screen to identify regions important for bacterial effector function

To efficiently screen for randomly generated mutations in *Legionella* effectors that cause a loss-

of-function phenotype and represent full-length protein rather than frameshift or nonsense

mutations, we applied the method of Holz and colleagues to select for expression of full-length

proteins in yeast (Holz *et al.* 2001). We cloned the yeast *HIS3* gene in frame behind yeast-

cytotoxic *Legionella* effectors on a high-copy galactose-inducible yeast expression plasmid.

After confirming that the His3 fusion does not interfere with the yeast-cytotoxic phenotype, and

therefore likely does not interfere with effector function, we generated a pool of random mutants

using the *E. coli* mutator strain XL1-Red. We then transformed this mutant plasmid pool to yeast

and monitored growth on different media types to assess the number of transformable plasmids (-

ura/glucose) and the efficiency of the random mutagenesis step (-ura/galactose) selecting all

mutations that caused a loss of function (Fig 1). To specifically select for full-length missense

loss-of-function mutants, we grew the transformed pool on medium with galactose and lacking

uracil and histidine (-ura-his/galactose), which requires the production of full-length effector-

His3 fusion protein and mutations in the effector that disrupt its activity.

Missense loss-of-function screen identifies conserved SdbB amino acid residues

As a proof of principle, we looked at SdbB which severely inhibits yeast growth (Fig 2A)

(Heidtman *et al.* 2009) and is part of the SidB paralog family, whose members are predicted to

be lipases from the α/β hydrolase enzyme family (Luo and Isberg 2004). After verifying that the

SdbB-His3 fusion was still capable of inhibiting yeast growth (Fig 2A) we created a random

mutagenesis pool of SdbB-His3 and quantified the number of colony-forming units (CFUs) on

the different selection media. While 1.89% of the transformable plasmids carried a mutation that

allowed for growth on SD-ura/galactose medium indicating some type of loss-of-function

mutation (Fig 2B), only 0.03% of the transformable plasmids carried a mutation allowing growth

on SD-ura/-his/galactose medium which requires the expression of full-length fusion protein.

The efficiency of the histidine selection step was verified by sequencing 20 clones from each

condition. The loss-of-function clones selected on –ura/galactose consisted of 16 frameshift

mutations, 2 nonsense mutations, 1 combination of a missense and frameshift mutation and 1

missense mutation in the *sdbB* gene (Fig 2C). In contrast, the loss-of-function clones selected on

–ura/-his/galactose only contained missense mutations (Fig 2D,F). For both conditions a number

of mutations were recovered several times suggesting that the mutational screen is reaching

saturation. To confirm that the identified mutations indeed rescued SdbB toxicity in yeast, we

compared the fitness of mutants to an empty vector control in a liquid growth curve assay (see

Experimental methods). The wild-type and His3-fused SdbB almost completely inhibited yeast

growth, while the SdbB mutants showed a fitness of 60-90% compared to an empty vector

control (Fig 2E and 2G).

The positions of the frameshift and nonsense mutations in SdbB (Fig 2D) indicate that a

large part of the protein is required for function, as even a nonsense mutation at S365, 84 amino

acid residues from the C-terminus, almost completely rescued activity. The missense loss-of-

function mutations (Fig 2D and 2F) target four amino acid residues (G116, G189, D273 and

H351) that are invariant in SdbB orthologs from *Legionella pneumophila* and other *Legionella*

species (Burstein *et al.* 2016) (Fig S1), suggesting they are essential for function or structure.

The G198E mutation is part of the GXS/CXG motif that is invariant across the SdbB orthologs

and is predicted to align with the so-called nucleophile elbow of the nucleophile-acid-base triad

of the α/β hydrolase active site (Brenner 1988; Ollis *et al.* 1992; Schrag and Cygler 1997;

Marchler-Bauer *et al.* 2017) (Fig S2). When the mutations are mapped onto the SdbB

AlphaFold2 model (Jumper *et al.* 2021; Varadi *et al.* 2021) the majority localize in the pocket

with the putative catalytic cysteine (C187), including the invariant D273 and H351 residues

captured in the screen suggesting they are the remaining residues of the catalytic triad (Fig 2H).

Thus, the SdbB example demonstrates that functionally important amino acid residues

can be identified using the random mutagenesis method in conjunction with the histidine

selection for full-length protein. Importantly, this approach significantly reduces the number of

sequenced clones required to identify amino acid residues or regions of interest, by

approximately 60-fold in the case of SdbB.

Missense loss-of-function screen identifies the active site of the characterized effector RavK

To benchmark the missense loss-of-function screen on a characterized effector, we looked at

RavK which also severely inhibits yeast growth (Heidtman *et al.* 2009). RavK was recently

shown to be a small, soluble metalloprotease that specifically cleaves host actin, and mutation of

the active site motif HExxH abolishes all activity and yeast toxicity (Liu *et al.* 2017). The RavK-

His3 fusion was still able to inhibit yeast growth (Fig 3B) and subjected to random mutagenesis

and selection for loss-of-function mutants on medium lacking histidine. Of the 14 loss-of-

function clones we sequenced, one clone contained a large, in-frame deletion from amino acid

residue 70 to residue 166, which was unexpected but confirms the strength of the histidine

selection for full-length protein. All other loss-of-function clones were caused by single point

mutations resulting in missense mutations (Fig 3A). In the fitness assay, wild-type and His3-

fused RavK almost completely inhibited yeast growth, while the RavK mutants displayed a

fitness of 70-90% compared to the empty vector control. The loss-of-function mutations all map

to the first half of RavK, suggesting that the N-terminal half of RavK is essential for RavK

function. This is in agreement with Liu and colleagues (Liu *et al.* 2017), who identified the

active site motif ($_{95}$HExxH$_{99}$) in the N-terminal half of the protein and found that the 50 C-

terminal residues of RavK can be deleted without any effect on its activity on actin.

Four of the loss-of-function mutations targeted the active site motif H$_{95}$ExxH$_{99}$, three of

which are mutations to E$_{96}$. Our screen also identified several residues outside of this motif that

are critical for RavK function. To investigate why these might be functionally important, we

performed an HHpred analysis which looks for structural homologs of proteins (Zimmermann *et*

al. 2018). HHpred identified many hits with homology to the HExxH metalloprotease motif.

Among the top five HHpred hits are three small soluble metalloproteases or minigluzincins,

anthrax lethal factor and a zinc dependent peptidase from the M48 family (Dalkas *et al.* 2010;

López-Pelegrín *et al.* 2013; López-Pelegrín *et al.* 2014) (Fig S3). Notably, some of the other

loss-of-function mutations occur in areas that have homology with structural elements in the

minigluzincins that contribute to the active site cleft (López-Pelegrín *et al.* 2013) (Fig S3).

Indeed, when the missense mutations are mapped on the RavK AlphaFold2 model (Jumper *et al.*

2021; Varadi *et al.* 2021) (Fig 3C) they cluster around the active site HExxH including the top

rim of the active site cleft.

The C-terminal domain of SdbA is a putative glycosyltransferase

SdbA is a member of the SidB paralog family (Luo and Isberg 2004). While the function of

SdbA remains undefined, experimental evolution of *Legionella* in mouse macrophages selected

for parallel SdbA nonsense and frameshift mutations in three out of four independent lineages

(Ensminger *et al.* 2012), suggesting that its normal function partially restricts growth in this

accidental host. While the N-terminal domain of SdbA has homology with SidB (Luo and Isberg

2004), the additional C-terminal domain does not have significant sequence homology to other

known proteins (data not shown). SdbA completely inhibits yeast growth when heterologously

expressed (Heidtman *et al.* 2009), making the missense loss-of-function screen an informative

tool by which to identify functional residues that might suggest a specific activity inside the

eukaryotic cell.

The missense loss-of-function screen identified 19 mutations in 24 sequenced clones

targeting 17 amino acid residues in SdbA (Fig 4A). In contrast to the smaller proteins SdbB and

RavK, the SdbA results included several double mutants. In some cases, it is not clear how much

each mutation contributes to the phenotype, considering that the fitness of the double mutants is

not dramatically different from the single loss-of-function mutations (Fig 4B). However, all the

single mutations that lead to a loss-of-function phenotype fall in the C-terminal domain and

mainly concentrate in two regions: $_{541}$GGTGHIS$_{547}$ and $_{957}$GGLSVME$_{963}$. An HHpred homology

search (Zimmermann *et al.* 2018) predicted with high confidence that the C-terminal domain is a

glycosyltransferase of the GT-B fold. When comparing the SdbA C-terminal domain with the

sequence of *E. coli* MurG, a well-studied member of the GT-B fold glycosyltransferase family,

the two mutated regions align with the G-loop 1 and a consensus region in GT-B fold

superfamily involved in binding the donor molecule (Ha *et al.* 2000; Hu *et al.* 2003; Crouvoisier

et al. 2007) (Fig 4C). Glycosyltranferases hydrolyze UDP-sugar donor molecules and transfer

the sugar to the acceptor molecule – which can be a variety of molecules such as small

molecules, lipid or proteins (Lairson *et al.* 2008). In MurG amino acid residues A263, L264,

L265, E268, Q287 and Q288 contact the donor molecule UDP-GlcNAc (Hu *et al.* 2003) (Fig4C)

while the G-loop 1 is thought to be involved in acceptor molecule binding (Ha *et al.* 2000).

Notably, mutations in these motifs abrogate MurG enzymatic activity, including mutation of the

residues H18 and E268 (Hu *et al.* 2003; Crouvoisier *et al.* 2007), whose corresponding residues

in SdbA (H545 and E963) were found to be mutated in our screen. Fig 4D shows the single

missense mutants abrogating SdbA activity mapped onto the AlphaFold2 model (Jumper *et al.*

2021; Varadi *et al.* 2021)of a C-terminal fragment (amino acid residues 510-1050) with the

residues H545 and E963 highlighted in yellow (43)

To test whether the C-terminal domain of SdbA is indeed a glycosyltransferase, we

purified the C-terminal fragment 510-1050 and the equivalent of the MurG E268A inactive

mutant in SdbA (E963A) and tested several UDP-sugars as substrate using the UDP-Glo assay

(Fig 4E). Glycosyltransferases can hydrolyze UDP-sugars in the absence of an acceptor molecule

(with water acting as an acceptor in the reaction)(Sheikh *et al.* 2017; Vicente *et al.* 2023). Indeed

wild-type SdbA 510-1050 hydrolyzes the UDP-GlcNAc donor while the E963A mutant cannot

(Fig 4E), confirming that SdbA is a glycosyltransferase with specificity for UDP-GlcNAc and

that a mutation in the E963 residue identified by the missense loss-of-function screen abrogates

that activity. Using the same assay, we determined the kinetic parameters of UDP-GlcNAc

hydrolysis by SdbA (Fig 4F), which revealed high affinity (low micromolar Km) of this enzyme

to UDP-GlcNAc suggesting efficient glycosylation activity *in vivo.*

Taken together, the missense loss-of-function screen identified critical residues in the C-

terminal domain of SdbA, which together with HHpred analysis predict that SdbA contains a

glycosyltransferase domain. The discovery and confirmation of the C-terminal

glycosyltransferase domain allows for targeted follow-up experiments to elucidate the action and

target of SdbA on eukaryotic host cells.

Discussion

Our observations demonstrate that the combination of a random mutagenesis loss-of-function

screen with a selection for full-length protein is very effective in specifically selecting loss-of-

function missense clones. In fact, all but one of the clones recovered in our assay contained

missense mutations, while the remaining one contained an in-frame deletion which included the

active site of RavK. The screens correctly identified the previously described active site of

RavK, and the predicted active site nucleophile motif and the remaining residues of the catalytic

triad of SdbB. Surprisingly, the N-terminal α/β hydrolase domain of SdbA did not accumulate

loss-of-function mutations, but the C-terminal domain appears to be crucial for the function of

SdbA. The prediction that the C-terminal domain of SdbA is a glycosyltransferase was supported

by *in vitro* activity towards UDP-GlcNAc and provides a way forward to study this *Legionella*

effector.

To effectively apply the random mutagenesis missense enrichment selection or to extend

this approach to bacteria and mammalian cells, a number of considerations need to be taken into

account. First, the protein of interest must have a selectable loss-of-function phenotype such as

alleviation of growth inhibition. Growth fitness is a universal phenotype that is easy to measure

in bacteria, yeasts and mammalian cell lines. Second, the C-terminal fusion of a selection marker

must not interfere with protein function. If the function of protein of interest is inhibited by the

C-terminal fusion, it could potentially be overcome by introducing linker regions of varying

length and flexibility (Chen *et al.* 2013) or by using a cleavable linker such as the ubiquitin K0

mutant that is processed by cytosolic deubiquitinases in eukaryotic cells (Bachran *et al.* 2013).

Similarly, the C-terminal selection marker must be able to function as a fusion protein or be

liberated by an *in vivo* cleavable linker.

To extend the random mutagenesis missense enrichment selection to bacteria, the

chloramphenicol acetyltransferase (CAT) gene which confers resistance to chloramphenicol

could be a good candidate as a C-terminal fusion partner. CAT has been successfully used in

protein fusions where it conferred chloramphenicol resistance during colony selection as a C-

terminal fusion partner, with increased selection efficiency when mutant fusion proteins are

soluble (Maxwell *et al.* 1999). In mammalian cell lines, a positive selection marker such as

blasticidin S deamidase could be used as a C-terminal fusion partner. Blasticidin S deamidase is

functional as a C-terminally fused protein (Suarez and McElwain 2009) and confers resistance

against blasticidin, which rapidly inhibits mammalian cell growth at a low dose (Kimura *et al.*

1994). An alternative, if no positive selection marker is available, is GFP, which has been used

extensively as a fused localization marker for various cellular compartments and organisms

(Margolin 2000; Roessel and Brand 2002; Huh *et al.* 2003). After a standard number of

generation doublings, GFP-positive cells, indicative of the presence of full-length protein, can be

selected by flow cytometry and cell sorting or by screening for GFP positive colonies in bacteria

and yeast. However, these approaches would reduce the screening throughput and/or require

specialized equipment.

Our initial results suggest that missense-directed mutagenesis will be a useful tool to help

identify potential functions for other bacterial effector proteins, many of which have low

sequence homology to characterized proteins (Gomez-Valero *et al.* 2011; Burstein *et al.* 2016).

Rather than being replaced by *in silico* protein modeling, we show how the two methodologies

complement one another and can be used to identify structural features essential for activity

against the eukaryotic cell. In *Legionella pneumophila* alone 10% of the translocated effectors

(Campodonico *et al.* 2005; Felipe *et al.* 2008; Heidtman *et al.* 2009; Shen *et al.* 2009; Guo *et al.*

2014; Urbanus *et al.* 2016), severely inhibit yeast growth and are possible candidates for the

random mutagenesis missense enrichment screen. Identifying functional residues within

uncharacterized effectors is a logical first step towards validating *in silico* protein models,

predicting effector activity, and designing protein-protein interaction studies.

Experimental methods

In frame effector-His3 fusion by yeast recombinational cloning

The *S. cerevisiae* BY4742 (*MATa, his3Δ1, leu2Δ0, met15Δ0, ura3Δ0*; (Brachmann *et al.* 1998))

strains overexpressing *lpg0275*, *lpg0969* and *lpg2482* (*sdbA, ravK,* and *sdbB* respectively) from

the high-copy vector pYES2 NT/A (Life Technologies, GAL 1 promoter, N-terminal 6X

HIS/Xpress tag and URA3 selectable marker) (Heidtman *et al.* 2009) were used to create the

effector-His3 fusion mutants by yeast recombinational cloning. The *S. cerevisiae HIS3* gene was

PCR amplified from pAG423GAL-ccdB (Alberti *et al.* 2007) using an effector specific forward

primer containing the last 50-60 nucleotides of the effector (minus the stop codon) followed by

the first 20-30 nucleotides of the *HIS3* sequence and the pYES-HIS3 reverse primer (Table S1).

The resulting PCR products were transformed together with XbaI/PmeI digested pYES2 NT/A

lpg0275, lpg0969 or lpg2482 to BY4742 using the high-efficiency LiOAC/PEG method (Gietz

and Schiestl 2007) and plated onto SD-uracil with 2% glucose. The resulting transformants were

screened by PCR and sequence verified. To confirm that the His3 fusion does not interfere with

effector function, the ability of the effector-HIS3 fusion to inhibit yeast growth was tested by

comparing the growth BY4742 with empty vector control, the wild-type effector and the

effector-HIS3 fusion in a yeast spot dilution assay as described previously (Urbanus *et al.* 2016).

Selection of loss-of-function mutations

The effector-HIS3 fusion vectors were mutagenized in XL1-red (Agilent) as per manufacturer's

instructions. XL1-red transformants were washed off the transformation plate, grown overnight

in 50 ml LB with ampicillin and the resulting mutant plasmid pool was purified by midiprep

(Promega). The mutant plasmid pool was transformed to BY4742 using the high-efficiency

LiOAC/PEG method (Gietz and Schiestl 2007). Four transformation reactions were performed

per screen, each using 1 µg of plasmid pool per reaction. One reaction was split in three parts

and plated onto different media types to quantify the transformable plasmids (SD- uracil + 2%

glucose), loss-of-function mutants (SD- uracil +2% galactose), and missense loss-of-function

mutants (SD-uracil/histidine + 2% galactose) and incubated for 2-4 days at 30°C. The remaining

transformation reactions were plated onto 150 mm SD-uracil/histidine + 2% galactose plates and

allowed to grow until colonies appeared (3-4 days). Plasmids were rescued from missense loss-

of-function mutants and transformed to the *E. coli* Top10 strain before sequencing using primers

in the vector and effector, if required (Table S1). Details of the sequenced mutants are shown in

Table S2.

Analysis of mutant fitness

Liquid growth assays were used to assess the effect of missense loss-of-function mutations on

yeast fitness as described (Urbanus *et al.* 2016) with the following modifications. Overnight

cultures of freshly transformed BY4742 with empty vector control, pYES2 NT/A effector-HIS3

wild-type and mutants were diluted 100-fold into 100 µl of SD-ura/2% galactose and grown with

Breathe-Easy adhesive seals (EK scientific) in a CellGrower robot (S&P robotics) at 30ºC with

intermittent shaking. Yeast growth was monitored for 30 h by measuring the OD_{620} every 15

min. Growth fitness was calculated as the ratio of the area under the curve (AUC) of a effector-

expressing strain over an empty vector control after 30 h using the R package GrowthCurver

(Sprouffske and Wagner 2016). The average AUC ratio and standard deviation was calculated

from three technical replicates.

HHpred analysis and sequence alignments

The amino acid sequence of RavK and SdbA (amino acid residues 528-1116) were submitted to

the HHpred server (https://toolkit.tuebingen.mpg.de/#/) (Zimmermann *et al.* 2018) analyzed

using MSA generation HHblits Uniclust20_2017_07 and Uniprot20_2016_02, respectively and

otherwise default parameters. The resulting alignments were visualized using Boxshade

(https://embnet.vital-it.ch/software/BOX_form.html). Amino acid sequence alignments of SdbB

with its orthologs or with SidB were generated using T-coffee (Tommaso *et al.* 2011) and

visualized using Jalview (Waterhouse *et al.* 2009) or Boxshade.

Nucleotide sugar donor specificity of the SdbA C-terminal domain

The gene fragment corresponding to SdbA (Lpg0275) residues 510-1050 was PCR amplified

from *Legionella pneumophila* str Philadelphia-1 genomic DNA and inserted into the pMCSG53

plasmid (Eschenfeldt *et al.* 2013) by ligation independent cloning, providing an N-terminal

6xHIS-TEV tag. The point mutant E963A was prepared by site-directed mutagenesis using

QuikChange™ site-directed mutagenesis kit (Stratagene) according to the manufacturer's

protocol. Plasmids were sequenced and transformed to the *E. coli* BL21 DE Gold strain for

purification. Recombinant proteins were purified to near homogeneity (>95%) using Ni-chelate

affinity chromatography on Ni-NTA Superflow resin (Qiagen) using standard protocols. Cultures

were grown in TB and expression was induced at an OD_{595} of 0.8 with 0.4 mM IPTG overnight

at 16 °C. Cells were harvested by centrifugation at 9,300× *g*, resuspended in 50 mM HEPES pH

7.5, 400 mM NaCl, 5% glycerol, 5 mM imidazole, and lysed by sonication. Lysates were

clarified by centrifugation at 21,000× *g* at 4°C and loaded onto gravity flow Ni-NTA agarose

columns (Qiagen), followed by washing with 50 mM HEPES pH 7.5, 400 mM NaCl, 5%

glycerol, 30 mM imidazole. Proteins were eluted using 50 mM HEPES pH 7.5, 400 mM NaCl,

5% glycerol, 250 mM imidazole and flash-frozen in liquid nitrogen for storage at −80°C. The

purity of the protein samples was assessed by SDS-PAGE and visualized by Coomassie Brilliant

Blue R.

The nucleotide sugar donor specificity of SdbA 510-1050 was assayed using the UDP-

Glo Glycosyltransferase Assay (Promega) according to manufacturer's protocol. Briefly, 0.09

μM of purified wild-type and E963A mutant SdbA 510-1050 protein was incubated with 100 μM

UDP-glucose, UDP-glcNAc, UDP-glucuron, UDP-galactose or UDP-galNAc for 1h at 30°C in

50 mM HEPES, pH 7.5, 100 mM KCl, 2 mM $MgCl_2$, 1 mM $MnCl_2$. The hydrolase activity of the

UDP-substrate was detected as the release of UDP by the UDP-Glo assay (Promega) after 20

min of incubation with UDP-Glo detection reagent. Luminescence was measured using a

SpectraMax M2 plate reader. Three technical repeats were performed per reaction.

The V_{max}, K_m and k_{cat} for wild-type SdbA 510-1050 with UDP-glcNAc was determined

by incubating 0.16 μM SdbA 510-1050 with UDP-glcNAc concentration range of 0.0039 – 2

mM for 1h at 30°C in 50 mM HEPES, pH 7.5, 100 mM KCl, 2 mM $MgCl_2$, 1 mM $MnCl_2$. Three

technical repeats were performed per reaction. Kinetic parameters were determined by non-linear

curve fitting from the Michaelis Menten plot using GraphPad Prism (version 5.00 for Windows,

GraphPad Software).

Acknowledgments

We thank Beth Nicholson, Morgan Petersen and John McPherson, for their suggestions and

careful reading of the manuscript and Kamran Rizzolo for help with cloning. This work was

supported by a Project Grant (AWE) from the Canadian Institutes of Health Research (PJT-

162256).

 Figure legends

 Figure 1: A method to enrich full-length missense mutants in a random mutagenesis screen.

Legionella effectors with a severe growth phenotype in yeast are fused in frame with the yeast

HIS3 gene in a high-copy yeast expression vector with galactose-inducible expression and an

uracil selection marker. The plasmid is randomly mutated in a *E. coli* mutator strain, such as

XL1-red. The resulting mutant pool contains a variety of mutants; sense, nonsense, missense and

frameshifts, of which the latter three can cause loss-of-function phenotypes. The plasmid pool is

transformed to *S. cerevisiae* strain BY4742 and grown under conditions that select all

transformable plasmids (–ura/glucose), all loss-of-function mutations (-ura with galactose

induction) and only full-length proteins containing missense loss-of-function mutations (-ura-his

with galactose induction).

Figure 2: The full-length ORF enriched random mutagenesis screen identifies missense

mutations in the putative active site pocket. A SdbB (lpg2482) and SdbB in-frame fusion with

His3 show a severe growth defect in a yeast spot dilution assay. **B** The percentage of loss-of-

function mutation in the pool of transformable plasmids selected on medium lacking uracil (-ura,

grey) and lacking uracil and histidine (-ura-his, black) with galactose induction. The average and

standard deviation of three independent replicates are shown. **C** Occurrence of frameshift,

nonsense and missense mutations in 20 sequenced clones selected on -ura and -ura-his with

galactose induction. **D** A schematic representation of loss-of-function clones selected on –ura

with galactose induction. Mutations recovered from the same clone are shown in black,

connected by a line while single mutations are shown in red. Mutation type is indicated as a

closed triangle (frameshift), open hexagon (nonsense) or closed circle (missense), and the

number of symbols reflects the occurrence of the mutation in the dataset. **E** The fitness of wild-

type SdbB, His3 fusion and loss-of-function mutations (selected on –ura/galactose) compared to

empty vector controls in liquid growth assays confirms the loss-of-function phenotype for the

SdbB random mutagenesis clones. The average and standard deviation of three technical

replicates are shown. **F** A schematic representation of SdbB loss-of-function clones selected on -

ura-his/galactose. Missense loss-of-function mutations are shown in red with a closed circle, the

number of symbols reflects the occurrence of the mutation in the dataset. Amino acid residues

shown for presentation purposes are shown in grey. **G** The fitness of wild-type SdbB, His3

fusion and loss-of-function mutations (selected on -ura,-his/galactose) compared to empty vector

controls in liquid growth assays confirms the loss-of-function phenotype for the SdbB random

mutagenesis clones. The average and standard deviation of three technical replicates are shown.

H The missense mutations identified by the random mutagenesis screen (D,F) are shown in

orange on an Alphafold2 model of SdbB (AF-Q5ZSN5-F1-model_v4), and residues from the

putative active site 185-GxCxG189 not captured by the screen are shown in magenta. Putative

catalytic triad C187-D273-H351 residies are shown with sticks and the box shows the

enlargement of the putative catalytic site. The AlphaFold2 model was visualized using Pymol 2.2

(Schrödinger).

Figure 3: RavK random mutagenesis captures residues lining the active site cleft. A A

schematic representation of the mutations causing a RavK loss-of-function phenotype when

expressed in yeast. Mutated residues are shown in red, amino acid residues shown for

presentation purposes are in grey and the number of symbols reflects the occurrence of the

mutation in the dataset. **B** The growth fitness of wild-type RavK, RavK-His3 and loss-of-

function mutants normalized to empty vector controls (see Methods) confirms the loss-of-

function phenotype for the RavK random mutagenesis clones. The average and standard

deviation of three technical replicates are shown. **C** The Alphafold2 model of RavK (AF-

Q5ZWW5-F1-model_v4.pdb) with missense mutations shown in orange. The histidine residues

of the active site motive $_{95}$HExxH$_{99}$ (Liu *et al.* 2017)not captured by the screen are shown in red.

The residues in the active site cleft are shown as sticks. The AlphaFold2 model was visualized

using Pymol 2.2 (Schrödinger).

Figure 4: The C-terminal domain of SdbA is a putative glycosyltransferase domain.

A A schematic representation of the mutations causing a SdbA loss-of-function phenotype when

expressed in yeast. Mutations recovered from the same clone are shown in black, connected by a

line and single mutations alleviating a SdbA induced growth defect are shown in red. Amino acid

residues shown for presentation purposes are shown in grey. Black and red closed circles

indicate the number of times the mutation was identified. **B** The growth fitness of wild-type

SdbA and loss-of-function mutants normalized to empty vector controls (see Methods for

details). SdbA and SdbA-His3 display a severe growth defect, while the loss-of-function

mutations rescue growth up to 50% of the empty vector control. The average and standard

deviation of three technical replicates are shown. **C** HHpred alignment of SdbA with 1F0K (Ha

et al. 2000) (*E. coli* MurG). The G-loop 1 and consensus sequence are shown with identical

residues (black) and similar residues (grey) highlighted. SdbA amino acid residues that when

mutated abrogate activity are indicated by a red closed circle. Amino acid residues in the MurG

consensus sequence contacting UDP-GlcNac are indicated by a black star and a red start if

mutations abrogate MurG activity. **D** The Alphafold2 model of SdbA (AF-Q5ZYT6-F1-

model_v4.pdb) with missense mutations shown in orange. Residues H545 and E963,

corresponding to residues H18 and E268 in MurG, are shown in yellow (43). The AlphaFold2

model was visualized using Pymol 2.2 (Schrödinger). **E** The SdbA glycosyltransferase domain

uses GlcNac as a donor substrate. Donor substrate specificity was tested by incubating 0.09 µM

of purified fragment (residues 510-1050) of wild-type SdbA or an inactive mutant (E963A) with

100 µM UDP-glucose, UDP-glcNAc, UDP-glucuron, UDP-galactose or UDP-galNAc for 1h at

30°C. The hydrolase activity of the UDP-substrate was detected as the release of UDP by the

UDP-Glo assay (Promega) after 20 min of incubation with UDP-Glo detection reagent.

Luminescence was measured using a SpectraMax M2 platereader. Three technical repeats were

performed per reaction. **F** Determination of the kinetic parameters for GlcNac hydrolysis by

SdbA. Reactions with 0.16 µg purified wild-type C-terminal domain of SdbA (residues 510-

1050) and a range of 0.0039 – 2 mM GlcNac and were incubated for 1 h at 30°C. GlcNac

hydrolysis was measured using the UDP-Glo glycosyltransferase assay as described above; three

technical replicates were performed per reaction. Kinetic parameters were determined by non-

linear curve fitting from the Michaelis-Menten plot.

References

Alberti, S., A. D. Gitler, and S. Lindquist, 2007 A suite of Gateway® cloning vectors for high-
throughput genetic analysis in Saccharomyces cerevisiae. Yeast 24: 913–919.

Bachran, C., T. Morley, S. Abdelazim, R. J. Fattah, S. Liu *et al.*, 2013 Anthrax Toxin-Mediated
Delivery of the Pseudomonas Exotoxin A Enzymatic Domain to the Cytosol of Tumor Cells
via Cleavable Ubiquitin Fusions. Mbio 4: e00201-13.

Brachmann, C. B., A. Davies, G. J. Cost, E. Caputo, J. Li *et al.*, 1998 Designer deletion strains
derived from Saccharomyces cerevisiae S288C: A useful set of strains and plasmids for PCR-
mediated gene disruption and other applications. Yeast 14: 115–132.

Brenner, S., 1988 The molecular evolution of genes and proteins: a tale of two serines. Nature
334: 528–530.

Burstein, D., F. Amaro, T. Zusman, Z. Lifshitz, O. Cohen *et al.*, 2016 Genomic analysis of 38
Legionella species identifies large and diverse effector repertoires. Nat Genet 48: 167–175.

Burstein, D., T. Zusman, E. Degtyar, R. Viner, G. Segal *et al.*, 2009 Genome-Scale Identification
of Legionella pneumophila Effectors Using a Machine Learning Approach. Plos Pathog 5:
e1000508.

Campodonico, E. M., L. Chesnel, and C. R. Roy, 2005 A yeast genetic system for the
identification and characterization of substrate proteins transferred into host cells by the
Legionella pneumophila Dot/Icm system. Mol Microbiol 56: 918–933.

Chen, X., J. L. Zaro, and W.-C. Shen, 2013 Fusion protein linkers: Property, design and
functionality. Adv Drug Deliver Rev 65: 1357–1369.

Crouvoisier, M., G. Auger, D. Blanot, and D. Mengin-Lecreulx, 2007 Role of the amino acid
invariants in the active site of MurG as evaluated by site-directed mutagenesis. Biochimie 89:
1498–1508.

Dalkas, G. A., C. T. Chasapis, P. V. Gkazonis, D. Bentrop, and G. A. Spyroulias, 2010
Conformational Dynamics of the Anthrax Lethal Factor Catalytic Center. Biochemistry 49:
10767–10769.

Ensminger, A. W., Y. Yassin, A. Miron, and R. R. Isberg, 2012 Experimental Evolution of
Legionella pneumophila in Mouse Macrophages Leads to Strains with Altered Determinants
of Environmental Survival. Plos Pathog 8: e1002731.

Eschenfeldt, W. H., M. Makowska-Grzyska, L. Stols, M. I. Donnelly, R. Jedrzejczak *et al.*, 2013
New LIC vectors for production of proteins from genes containing rare codons. J. Struct.
Funct. Genom. 14: 135–144.

Escoll, P., M. Rolando, L. Gomez-Valero, and C. Buchrieser, 2013 Molecular Mechanisms in
Legionella Pathogenesis. Curr Top Microbiol 376: 1–34.

Felipe, K. S. de, R. T. Glover, X. Charpentier, O. R. Anderson, M. Reyes *et al.*, 2008 Legionella
Eukaryotic-Like Type IV Substrates Interfere with Organelle Trafficking. Plos Pathog 4:
e1000117.

Finsel, I., and H. Hilbi, 2015 Formation of a pathogen vacuole according to Legionella
pneumophila: how to kill one bird with many stones. Cell Microbiol 17: 935–950.

Gietz, R. D., and R. H. Schiestl, 2007 High-efficiency yeast transformation using the LiAc/SS
carrier DNA/PEG method. Nat Protoc 2: 31–34.

Gomez-Valero, L., C. Rusniok, D. Carson, S. Mondino, A. E. Pérez-Cobas *et al.*, 2019 More
than 18,000 effectors in the Legionella genus genome provide multiple, independent
combinations for replication in human cells. Proc National Acad Sci 116: 201808016.

Gomez-Valero, L., C. Rusniok, C. Cazalet, and C. Buchrieser, 2011 Comparative and Functional
Genomics of Legionella Identified Eukaryotic Like Proteins as Key Players in Host–Pathogen
Interactions. Front Microbiol 2: 208.

Gomez-Valero, L., C. Rusniok, M. Rolando, M. Neou, D. Dervins-Ravault *et al.*, 2014
Comparative analyses of Legionella species identifies genetic features of strains causing
Legionnaires' disease. Genome Biol 15: 505.

Guo, Z., R. Stephenson, J. Qiu, S. Zheng, and Z.-Q. Luo, 2014 A Legionella effector modulates
host cytoskeletal structure by inhibiting actin polymerization. Microbes Infect 16: 225–236.

Ha, S., D. Walker, Y. Shi, and S. Walker, 2000 The 1.9 Å crystal structure of Escherichia coli
MurG, a membrane-associated glycosyltransferase involved in peptidoglycan biosynthesis.
Protein Sci 9: 1045–1052.

Heidtman, M., E. J. Chen, M. Moy, and R. R. Isberg, 2009 Large-scale identification of
Legionella pneumophila Dot/Icm substrates that modulate host cell vesicle trafficking
pathways. Cell Microbiol 11: 230–248.

Holz, C., A. Lueking, L. Bovekamp, C. Gutjahr, N. Bolotina *et al.*, 2001 A Human cDNA
Expression Library in Yeast Enriched for Open Reading Frames. Genome Res 11: 1730–
1735.

Hu, Y., L. Chen, S. Ha, B. Gross, B. Falcone *et al.*, 2003 Crystal structure of the MurG:UDP-
GlcNAc complex reveals common structural principles of a superfamily of
glycosyltransferases. Proc National Acad Sci 100: 845–849.

Huang, L., D. Boyd, W. M. Amyot, A. D. Hempstead, Z. Luo *et al.*, 2011 The E Block motif is
associated with Legionella pneumophila translocated substrates. Cell Microbiol 13: 227–245.

Huh, W.-K., J. V. Falvo, L. C. Gerke, A. S. Carroll, R. W. Howson *et al.*, 2003 Global analysis
of protein localization in budding yeast. Nature 425: 686–691.

Isberg, R. R., T. J. O'Connor, and M. Heidtman, 2009 The Legionella pneumophila replication
vacuole: making a cosy niche inside host cells. Nat Rev Microbiol 7: 13–24.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov *et al.*, 2021 Highly accurate protein
structure prediction with AlphaFold. Nature 1–11.

Kimura, M., A. Takatsuki, and I. Yamaguchi, 1994 Blasticidin S deaminase gene from
Aspergillus terreus (BSD): a new drug resistance gene for transfection of mammalian cells.
Biochim. Biophys. Acta (BBA) - Gene Struct. Expr. 1219: 653–659.

Kozlov, G., K. Wong, and K. Gehring, 2018 Crystal structure of the Legionella effector Lem22.
Proteins Struct Funct Bioinform 86: 263–267.

Lairson, L. L., B. Henrissat, G. J. Davies, and S. G. Withers, 2008 Glycosyltransferases:
Structures, Functions, and Mechanisms. Annu Rev Biochem 77: 521–555.

Liu, Y., W. Zhu, Y. Tan, E. S. Nakayasu, C. J. Staiger *et al.*, 2017 A Legionella Effector
Disrupts Host Cytoskeletal Structure by Cleaving Actin. Plos Pathog 13: e1006186.

López-Pelegrín, M., N. Cerdà-Costa, A. Cintas-Pedrola, F. Herranz-Trillo, P. Bernadó *et al.*,
2014 Multiple Stable Conformations Account for Reversible Concentration-Dependent
Oligomerization and Autoinhibition of a Metamorphic Metallopeptidase. Angewandte
Chemie Int Ed 53: 10624–10630.

López-Pelegrín, M., N. Cerdà-Costa, F. Martínez-Jiménez, A. Cintas-Pedrola, A. Canals *et al.*,
2013 A Novel Family of Soluble Minimal Scaffolds Provides Structural Insight into the
Catalytic Domains of Integral Membrane Metallopeptidases*. J Biol Chem 288: 21279–
21294.

Luo, Z.-Q., and R. R. Isberg, 2004 Multiple substrates of the Legionella pneumophila Dot/Icm
system identified by interbacterial protein transfer. P Natl Acad Sci Usa 101: 841–846.

Marchler-Bauer, A., Y. Bo, L. Han, J. He, C. J. Lanczycki *et al.*, 2017 CDD/SPARCLE:
functional classification of proteins via subfamily domain architectures. Nucleic Acids Res
45: D200–D203.

Margolin, W., 2000 Green Fluorescent Protein as a Reporter for Macromolecular Localization in
Bacterial Cells. Methods 20: 62–72.

Maxwell, K. L., A. K. Mittermaier, J. D. Forman-Kay, and A. R. Davidson, 1999 A simple in
vivo assay for increased protein solubility. Protein Sci 8: 1908–1911.

Morar, M., E. Evdokimova, C. Chang, A. W. Ensminger, and A. Savchenko, 2015 Crystal
structure of the Legionella pneumophila lem10 effector reveals a new member of the HD
protein superfamily. Proteins Struct Funct Bioinform 83: 2319–2325.

O'Connor, T. J., Y. Adepoju, D. Boyd, and R. R. Isberg, 2011 Minimization of the Legionella
pneumophila genome reveals chromosomal regions involved in host range expansion. Proc
National Acad Sci 108: 14733–14740.

Ollis, D. L., E. Cheah, M. Cygler, B. Dijkstra, F. Frolow *et al.*, 1992 The α / β hydrolase fold.
Protein Eng Des Sel 5: 197–211.

Pinotsis, N., and G. Waksman, 2017 Structure of the WipA protein reveals a novel tyrosine
protein phosphatase effector from Legionella pneumophila. J Biol Chem 292: 9240–9251.

Qiu, J., and Z.-Q. Luo, 2017 Legionella and Coxiella effectors: strength in diversity and activity.
Nat Rev Microbiol 15: 591–605.

Roessel, P. van, and A. H. Brand, 2002 Imaging into the future: visualizing gene expression and
protein interactions with fluorescent proteins. Nat. Cell Biol. 4: E15–E20.

Schrag, J. D., and M. Cygler, 1997 [4] Lipases and αβ hydrolase fold. Methods Enzymol 284:
85–107.

Schrödinger, L. The PyMOL Molecular Graphics System, Version~2.2.

Segal, G., M. Purcell, and H. A. Shuman, 1998 Host cell killing and bacterial conjugation require
overlapping sets of genes within a 22-kb region of the Legionella pneumophila genome. Proc
National Acad Sci 95: 1669–1674.

Sheikh, M. O., S. M. Halmo, S. Patel, D. Middleton, H. Takeuchi *et al.*, 2017 Rapid screening of
sugar-nucleotide donor specificities of putative glycosyltransferases. Glycobiology 27: 206–
212.

Shen, X., S. Banga, Y. Liu, L. Xu, P. Gao *et al.*, 2009 Targeting eEF1A by a Legionella
pneumophila effector leads to inhibition of protein synthesis and induction of host stress
response. Cell Microbiol 11: 911–926.

Sherwood, R. K., and C. R. Roy, 2016 Autophagy Evasion and Endoplasmic Reticulum
Subversion: The Yin and Yang of Legionella Intracellular Infection. Annu Rev Microbiol 70:
413–433.

Sprouffske, K., and A. Wagner, 2016 Growthcurver: an R package for obtaining interpretable
metrics from microbial growth curves. Bmc Bioinformatics 17: 172.

Suarez, C. E., and T. F. McElwain, 2009 Stable expression of a GFP-BSD fusion protein in
Babesia bovis merozoites. Int J Parasitol 39: 289–297.

Tommaso, D. P., S. Moretti, and X.-I. acids …, 2011 T-Coffee: a web server for the multiple
sequence alignment of protein and RNA sequences using structural information and
homology extension.

Toulabi, L., X. Wu, Y. Cheng, and Y. Mao, 2013 Identification and Structural Characterization
of a Legionella Phosphoinositide Phosphatase*. J Biol Chem 288: 24518–24527.

Urbanus, M. L., A. T. Quaile, P. J. Stogios, M. Morar, C. Rao *et al.*, 2016 Diverse mechanisms
of metaeffector activity in an intracellular bacterial pathogen, Legionella pneumophila. Mol
Syst Biol 12: 893.

Varadi, M., S. Anyango, M. Deshpande, S. Nair, C. Natassia *et al.*, 2021 AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-sequence space
with high-accuracy models. Nucleic Acids Res. 50: D439–D444.

Vicente, J. B., A. C. L. Guerreiro, B. Felgueiras, D. Chapla, D. Tehrani *et al.*, 2023
Glycosyltransferase 8 domain-containing protein 1 (GLT8D1) is a UDP-dependent
galactosyltransferase. Sci. Rep. 13: 21684.

Vogel, Joseph. P., H. L. Andrews, S. K. Wong, and R. R. Isberg, 1998 Conjugative Transfer by
the Virulence System of Legionella pneumophila. Science 279: 873–876.

Waterhouse, A. M., J. B. Procter, D. M. A. Martin, M. Clamp, and G. J. Barton, 2009 Jalview
Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics 25:
1189–1191.

Wong, K., G. Kozlov, Y. Zhang, and K. Gehring, 2015 Structure of the Legionella Effector,
lpg1496, Suggests a Role in Nucleotide Metabolism*. J Biol Chem 290: 24727–24737.

Zhu, W., S. Banga, Y. Tan, C. Zheng, R. Stephenson *et al.*, 2011 Comprehensive Identification
of Protein Substrates of the Dot/Icm Type IV Transporter of Legionella pneumophila. Plos
One 6: e17638.

Zimmermann, L., A. Stephens, S.-Z. Nam, D. Rau, J. Kübler *et al.*, 2018 A Completely
Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at its Core. J Mol
Biol 430: 2237–2243.

Figure 1: A method to enrich full-length missense mutants in a random mutagenesis screen

Figure 2: Random mutagenesis screen identifies missense mutations in amino acid residues of the putative active site pocket

Figure 3 - RavK random mutagenesis captures residues lining the active site cleft

Figure 4: The C-terminal domain of SdbA is a putative glycosyltransferase domain

Figure S1: Random mutagenesis targets conserved residues in SdbB orthologs from seven *Legionella* species. The sequences of SdbB orthologs identified by Burstein et al (Nat Genet 48(2): 167-175, 2016) were aligned using T-coffee, visualized with Jalview and coloured by % identity. The missense mutations identified by the random mutagenesis screen are indicated with a red closed circle above the SdbB (lpg2482) sequence. Four of the seven mutations target invariant residues; G116, G189, D273 and H351 shown in red boxes.

```
SidB    1  MAKIYNAPKPKYSGWEWFKFIAIRTVFPPVLLWDLIKIGANKLLGEWVSGLVLPAQNENF
SdbB    1  MAKTITKDSFFHSRLQ--QFVA-NSLFPAG-------------AGDWYKNKGYGKKGE-V

SidB   61  DDLAISDDTVSNYNED-------------D--LICEKHDVITHDGAHLDTFEVRHRSQESI
SdbB   44  DDEAPFRDFVEKQKTDKKSHYYKEFQGLDLSLKKVKTKLVSGESCQLEVMKCQPESNEPK

SidB  107  DPKYQKYIINLVGNGMCYEHIIDDIKEDSKALKANVIGFNLRGVGQSTGKAKSSEDLVAD
SdbB  104  KPGTGKHIVYFPGANTYYQACFRDISTACKETGATVHAFNFPGTGLSSGKVREANDLINA

                                            GxS/CxG
SidB  167  GIAQVQRLLDQGVSPQNITLKGHSLGAGVASLVAQHFHQLGQ-PINLFNSRSFSTITNFL
SdbB  164  GISVVSSLIKQGVHPDDIILQGDCYGASIALEVKKQLEDQADIKVRAIMNNVFKSFKAAV

SidB  226  VGHMRLERDEIGRAIGHKDSTVGTILGWLAKPFIKFGVALAKWEINAGSAFKSVPEAYKD
SdbB  224  C----------------DMITQSPWLPNILKSIVKRLLEFTGWHVTPGKKYKHA-DPYQC

SidB  286  YIVVRSRKEIRGERIDDAVIPHYASIHKELASERHKKKAEIDEEIANLDDIIRKADPLAK
SdbB  267  HIQHLGDQTLESSTLSGKVSKYHHEIQT--GQTKSQKRAPITDTC--PEEYRKDRDELDR

SidB  346  PGLANARDALVQAREKIKSDRKMETDVQYANGHNSDWNALHNRSGKSA-QTFFREFVQRT
SdbB  323  KHYVRVKE---SAKERLASKFGVD-KFGRVNAHFADLCELEMLDGQSVYQGFVNDYIARS

SidB  405  EA---DHAVKSIPEIN--------------------------------------------
SdbB  379  NAYIEKHPQKGIKEVQDDLRKLHYLQPADSIEITEDEAQDFNTVVDLITEEQQLRHDRFN

SidB       ----------
SdbB  439  DNTIGKSISM
```

Figure S2: SidB and SdbB amino acid sequence alignment. The SidB (lpg1642) and SdbB (lpg2482) were aligned with T-coffee and visualized with Boxshade, where identical residues are shown in black and similar residues in grey. The active site motif GxS/CxG predicted by NCBI conserved domain search (Marchler-Bauer A et al.,2017, Nucleic Acids Res.45(D)200-3.) for SidB is indicated, suggesting that SdbB C187 is part of the active site catalytic triad.

```
                                                                                    β2
                                                                                 ________
ravK      1   MVSLEHIQKLISECRKLGKDGLDNGTNGLIPELEIDVVPPSAFLG-VGNNPAIFVNSKTY
4QHJ_B    1   MKDRKILNEILSNTINE------LNLNDKKANIKIKIKPLKRKIA-SISL----------
4JIU_A    1   ------------------------------------------------LG-YQRPVKVRIRPLK-
4JIX_B    1   ------------------------------------------------LN-INESINIEIKPMK-
2L0R_A    1   ----------------------------------------------------------
3C37_A    1   -VNDPEVQRYVDKVGKRLLSGARAV----EFDYVFKVVKDDSVNAFAIPGGRVYVHTGLL

          rim of active site cleft
          _________________             β3
                                    ___________
ravK     60   KLMRTTHEKWVENKTIVFKSYLL-S-QPAIKIIGAIVHETGHAFN------------V--
4QHJ_B   44   -----------TNKTIYINKNILPY-LSDEEIRFILAHELLHLKY------------GKY
4JIU_A   16   ---MSIARVSFKYGTITLDPAVL-N-LEEEMFYILIHELAHLKAETSYHSSSFWREV--
4JIX_B   16   ---QKIASFSFKTKTLRLNKYVV-ENFDEELLHYIILHELIHFKI--------------
2L0R_A    1   ---------------------------NDSEGFIHEFGHAVD------------D--
3C37_A   56   KAAD---------------------NETELAGVLAHEINHAVA--------------

ravK    104   -----AAKIPNT-EANACIFEIEVLMRLFQVKSPLLLGCTELDMQSYFKSRLTDYNKC--
4QHJ_B   80   HINEFEEELLFL-FPNKEAILFNLINKLFQKK--------------------------
4JIU_A   69   -----EKVFPGE-R--AKEIEDRIMTKLQR---------------------------
4JIX_B        ---------------------------------------------------------
2L0R_A   17   -----YAGYLLDKNQSDLVTNSKKFIDIFKEEGSNLTSYGRTNEAEFFAEAFRLMHSTDH
3C37_A        ---------------------------------------------------------

ravK    156   ------VKDCQCLAEMVEFITHQ
4QHJ_B        ----------------------
4JIU_A        ----------------------
4JIX_B        ----------------------
2L0R_A   72   AERLKVQKNAPKTFQFINDQIKF
3C37_A        ----------------------
```

Supplemental figure 3 - HHpred alignment of RavK with metalloproteases.
Alignment of the HHpred top hits; MJ1213 (4QHJ), proabylysin (4JIU), projannalysin (4JIX) anthrax lethal factor (2L0R) and a zinc-dependent endopeptidase of the M84 family (3C37) with RavK is shown where grey and black background indicate the level of amino acid residue conservation as 50% or more similarity or 50% or more identity, respectively. Structural elements of proabylysin (4JIU) are indicated on top of the alignment and mutations that abolish RavK activity are indicated by a red closed circle.

Table S1: Primers used for yeast recombinational cloning and sequencing. In the yeast recombinational cloning primers the effector sequence is underlined, the HIS3 sequence in bold and the pYES2 vector sequence unchanged.

Primer	Sequence (5'-3')	Use
lpg0275-HIS3 FW	AAGGAAGATTTACATCATAATGAGCTTTCCAGTCAAAGTTTGGTTAGTGTCAGTGCTGGT**atgacagagcagaaagccctagtaaagcgt**	yeast recombinational cloning
lpg0969-HIS3 FW	TGGTTAAAGAGCATTTTCATTCTCCTGAAACAGAGATAAAGCTTGATATA**atgacagagcagaaagccctagtaaagcgt**	yeast recombinational cloning
lpg2482-HIS3 FW	GAACAACAACTAAGGCATGATCGATTCAACGATAATACAATAGGTAAAAGCATCTCTATG**atgacagagcagaaagccctagtaaagcgt**	yeast recombinational cloning
pYES2-HIS3 rv	CGTGAATGTAAGCGTGACATAACTAATTACATGATGCGGCCCTCTAGGATCAGCGGGTTT**ctacataagaacacctttggtggagggaac**	yeast recombinational cloning
scHIS3_seqF	GCTCTGGCCAAGCATTCC	verification of his3 fusion
scHIS3_seqR	TTCAGTGGTGTGATGGTCGT	verification of his3 fusion , sdbB and ravK random mutant sequencing
Gal1 fw	AATATACCTCTATACTTTAACGTC	sdbB and ravK random mutant sequencing
lpg0969seqF	GCTGCCAAAATTCCCAATAC	ravK random mutant sequencing
lpg0969seqR	CCAGTTCCGTACACCCTAGC	ravK random mutant sequencing
lpg0275_467F	AGGGATTGGAGTTCCTGGAC	sdbA random mutant sequencing
lpg0275_637R	CGTCTTCACCTGTAACTTGCAC	sdbA random mutant sequencing
lpg0275_942F	TGAGAAGGGCATGAAAGTCA	sdbA random mutant sequencing
lpg0275_1417F	TCCCGACAAGATTTGGTAGC	sdbA random mutant sequencing
lpg0275_1920F	CAGAACCCGTCCTTATGTGG	sdbA random mutant sequencing
lpg0275_2435F	ACCCACGCAATAACCCTATG	sdbA random mutant sequencing
lpg0275_2825F	TGGCACCAATAATGACCAGA	sdbA random mutant sequencing

Table S2. Random mutagenesis mutants for RavK, SdbA and SdbB. Mutants suppressing the growth defect of RavK,SdbA and SdbB in yeast were selected on SD-gal-ura (induced expression and plasmid selection) or SD-gal-ura-his (induced expression, plasmid selection and full-length fusion protein selection respectively). This table summarizes the identified mutations, the type of mutation and the consequences of the mutation.

Mutant no	Locus no	Gene name	Selection	Nucleotide mutation	Type of mutation	Amino acid mutation(s)	1st mutation	2nd mutation
1	lpg0275	sdbA	+ gal, -ura-his	367G>A, 1624G>A	amino acid substitution	V123M, G542R	123	542
2	lpg0275	sdbA	+ gal, -ura-his	1622G>A, 452C>T	amino acid substitution	S151L, G541E	151	541
3	lpg0275	sdbA	+ gal, -ura-his	1222G>A, 1622G>A	amino acid substitution	V408I, G541E	408	541
4	lpg0275	sdbA	+ gal, -ura-his	1622G>A	amino acid substitution	G541E	541	
5	lpg0275	sdbA	+ gal, -ura-his	1621G>A	amino acid substitution	G541E	541	
6	lpg0275	sdbA	+ gal, -ura-his	1622G>A	amino acid substitution	G541E	541	
7	lpg0275	sdbA	+ gal, -ura-his	1622G>A	amino acid substitution	G541E	541	
8	lpg0275	sdbA	+ gal, -ura-his	1622G>A	amino acid substitution	G541E	541	
9	lpg0275	sdbA	+ gal, -ura-his	1625G>A	amino acid substitution	G542E	542	
10	lpg0275	sdbA	+ gal, -ura-his	1630G>A, 2378C>T	amino acid substitution	G544R, P793L	544	793
11	lpg0275	sdbA	+ gal, -ura-his	1630G>A, 2826G>A	amino acid substitution	G544R, M942I	544	942
12	lpg0275	sdbA	+ gal, -ura-his	1634A>G	amino acid substitution	H545R	545	
13	lpg0275	sdbA	+ gal, -ura-his	1633C>T	amino acid substitution	H545Y	545	
14	lpg0275	sdbA	+ gal, -ura-his	1639T>C	amino acid substitution	S547P	547	
15	lpg0275	sdbA	+ gal, -ura-his	2275G>A	amino acid substitution	D759N	759	
16	lpg0275	sdbA	+ gal, -ura-his	2275G>A	amino acid substitution	D759N	759	
17	lpg0275	sdbA	+ gal, -ura-his	2731C>A, 2866G>A	amino acid substitution	G911R, G956R	911	956
18	lpg0275	sdbA	+ gal, -ura-his	2870G>A	amino acid substitution	G957D	957	
19	lpg0275	sdbA	+ gal, -ura-his	2873G>A	amino acid substitution	G958D	958	
20	lpg0275	sdbA	+ gal, -ura-his	2887G>A	amino acid substitution	E963K	963	
21	lpg0275	sdbA	+ gal, -ura-his	2887G>A	amino acid substitution	E963K	963	
22	lpg0275	sdbA	+ gal, -ura-his	2887G>A	amino acid substitution	E963K	963	
23	lpg0275	sdbA	+ gal, -ura-his	2980G>A	amino acid substitution	E994K	994	
24	lpg0275	sdbA	+ gal, -ura-his	2980G>A	amino acid substitution	E994K	994	
25	lpg0969	ravK	+ gal, -ura-his	41G>A	amino acid substitution	C14Y	14	
26	lpg0969	ravK	+ gal, -ura-his	98T>C	amino acid substitution	L33S	33	
27	lpg0969	ravK	+ gal, -ura-his	152C>T	amino acid substitution	A51V	51	
28	lpg0969	ravK	+ gal, -ura-his	158T>C	amino acid substitution	F53S	53	
29	lpg0969	ravK	+ gal, -ura-his	203A>C	amino acid substitution	K68T	68	
30	lpg0969	ravK	+ gal, -ura-his	205T>A	amino acid substitution	W69R	69	
31	lpg0969	ravK	+ gal, -ura-his	220A>G	amino acid substitution	T74A	74	
32	lpg0969	ravK	+ gal, -ura-his	286G>A	amino acid substitution	E96K	96	
33	lpg0969	ravK	+ gal, -ura-his	286G>A	amino acid substitution	E96K	96	
34	lpg0969	ravK	+ gal, -ura-his	287A>G	amino acid substitution	E96G	96	
35	lpg0969	ravK	+ gal, -ura-his	293G>T	amino acid substitution	G98V	98	
36	lpg0969	ravK	+ gal, -ura-his	331G>A	amino acid substitution	E111K	111	
37	lpg0969	ravK	+ gal, -ura-his	354A>C	amino acid substitution	E118D	118	
38	lpg2482	sdbB	+gal,-ura	33_34insT	frameshift	H12SfsX1	12	
39	lpg2482	sdbB	+gal,-ura	33_34insT	frameshift	H12SfsX1	12	
40	lpg2482	sdbB	+gal,-ura	33_34insT	frameshift	H12SfsX1	12	
41	lpg2482	sdbB	+gal,-ura	119delA	frameshift	K40RfsX33	39	
42	lpg2482	sdbB	+gal,-ura	119delA	frameshift	K40RfsX33	39	

43	lpg2482	sdbB	+gal,-ura	166_167insA	frameshift	Q56TfsX4	56	
44	lpg2482	sdbB	+gal,-ura	172_173insA	frameshift	T58NfsX2	58	
45	lpg2482	sdbB	+gal,-ura	172_173insA	frameshift	T58NfsX2	58	
46	lpg2482	sdbB	+gal,-ura	172_173insA	frameshift	T58NfsX2	58	
47	lpg2482	sdbB	+gal,-ura	172delA	frameshift	T58PfsX15	58	
48	lpg2482	sdbB	+gal,-ura	265C>T	premature stop codon	Q89AMBER	89	
49	lpg2482	sdbB	+gal,-ura	312delA	frameshift	K104NfsX56	104	
50	lpg2482	sdbB	+gal,-ura	407G>A, 469_470insG	amino acid substitution and frameshift	G136E, G157AfsX2	136	157
51	lpg2482	sdbB	+gal,-ura	555_556insG	frameshift	G185=fsX1	185	
52	lpg2482	sdbB	+gal,-ura	594_595insA	frameshift	Q199AfsX21	199	
53	lpg2482	sdbB	+gal,-ura	818A>G	amino acid substitution	D273G	273	
54	lpg2482	sdbB	+gal,-ura	905_906insA	frameshift	R302KfsX17	302	
55	lpg2482	sdbB	+gal,-ura	905_906insA	frameshift	R302KfsX17	302	
56	lpg2482	sdbB	+gal,-ura	905delA	frameshift	R302GfsX40	302	
57	lpg2482	sdbB	+gal,-ura	1094C>A	premature stop codon	S365AMBER	365	
58	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
59	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
60	lpg2482	sdbB	+ gal, -ura-his	346G>A	amino acid substitution	G116R	116	
61	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
62	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
63	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
64	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
65	lpg2482	sdbB	+ gal, -ura-his	347G>A	amino acid substitution	G116E	116	
66	lpg2482	sdbB	+ gal, -ura-his	566G>A	amino acid substitution	G189E	189	
67	lpg2482	sdbB	+ gal, -ura-his	566G>A	amino acid substitution	G189E	189	
68	lpg2482	sdbB	+ gal, -ura-his	836C>T	amino acid substitution	S279L	279	
69	lpg2482	sdbB	+ gal, -ura-his	836C>T	amino acid substitution	S279L	279	
70	lpg2482	sdbB	+ gal, -ura-his	836C>T	amino acid substitution	S279L	279	
71	lpg2482	sdbB	+ gal, -ura-his	842T>C	amino acid substitution	L281S	281	
72	lpg2482	sdbB	+ gal, -ura-his	842T>C	amino acid substitution	L281S	281	
73	lpg2482	sdbB	+ gal, -ura-his	842T>C	amino acid substitution	L281S	281	
74	lpg2482	sdbB	+ gal, -ura-his	926G>A	amino acid substitution	C309Y	309	
75	lpg2482	sdbB	+ gal, -ura-his	926G>A	amino acid substitution	C309Y	309	
76	lpg2482	sdbB	+ gal, -ura-his	926G>A	amino acid substitution	C309Y	309	
77	lpg2482	sdbB	+ gal, -ura-his	1051C>T	amino acid substitution	H351Y	351	