7 # When visual attention is divided in the flash-lag effect
8

Jane Yook[1,2,3], Hinze Hogendoorn[1,4], Gereon R. Fink[2,5], Simone Vossel[2,3]*, Ralph Weidner[2]*

*These authors contributed equally

[1] Melbourne School of Psychological Sciences, The University of Melbourne, Melbourne VIC
3010, Australia

[2] Cognitive Neuroscience, Institute of Neuroscience and Medicine (INM-3), Forschungszentrum
Jülich, 52428 Jülich, Germany

[3] Department of Psychology, Faculty of Human Sciences, University of Cologne, 50923 Cologne,
Germany

[4] School of Psychology and Counselling, Queensland University of Technology, Brisbane QLD
4059, Australia

[5] Department of Neurology, Faculty of Medicine, University Hospital Cologne, University of
Cologne, 50923 Cologne, Germany

Abstract

The flash-lag effect (FLE) is the perceived lagging behind of a flash physically aligned to a continuously moving object. The illusion is attributed to a compensatory mechanism of motion extrapolation, which ensures that the perceived position of a moving object aligns with its real-time position despite neural processing delays. While several studies have demonstrated that attention can modulate the FLE, the precise role of attention in the extrapolation process remains elusive. Thus, in the current study, we sought to disentangle the influence of visual attention by manipulating the amount of attention allocated to moving stimuli in different locations. By directing attention to one, two, three, or four stimuli presented in different quadrants across trials, we measured the trial-wise FLE to investigate potential FLE magnitude or variability modulations under different attention conditions. Our results showed that FLE magnitudes were significantly larger when attention was divided among two, three, or four stimuli than when attention was focused on one stimulus. Surprisingly, FLE variability showed no difference across attention conditions. These findings show that the compensatory mechanisms for neural processing delays in dynamic environments are modulated by attention to enable efficient processing of multiple objects simultaneously.

Keywords

Flash-lag effect, attention, divided attention, motion extrapolation, neural processing delays, multiple objects

1. Introduction

The flash-lag effect (FLE) is a classic demonstration of a motion-induced position shift. As initially

shown by Mackay (1958), the illusion occurs when a static object flashes next to a continuously

moving object, leading to a perceived spatial offset between their positions despite their physical

alignment at the moment of the flash. Specifically, the flash is misperceived as lagging behind the

moving object.

While the FLE has garnered many explanations including theories of attention shifts, differential

processing latency, temporal position averaging, postdiction, and discrete perception (Baldo &

Klein, 2010; Brenner & Smeets, 2000; Eagleman & Sejnowski, 2000; Khurana et al., 2000;

Krekelberg & Lappe, 1999, 2001; Purushothaman et al., 1998; Schneider, 2018; Whitney &

Murakami, 1998), motion extrapolation provides a parsimonious explanation. This mechanism

compensates for inherent delays in perceptual processing (Nijhawan, 1994, 2002, 2008), already

operating on early stages of visual processing (Johnson et al., 2023). It ensures that the perceived

position of a moving object aligns more closely with its veridical position. In the context of the

FLE, the predictable motion trajectory of the moving object allows for extrapolation. In contrast,

the static flash, having no motion trajectory, cannot compensate for neural delays through

extrapolation, resulting in the misperception of the flash as lagging (Murakami, 2001).

Our limited processing capacity poses a challenge for motion extrapolation in dynamic

environments with multiple moving objects (cf. multiple object tracking; Pylyshyn & Storm, 1988;

see also Adamian & Andersen, 2022; Howe & Holcombe, 2012; Luu & Howe, 2015). Our

attentional capacity may be insufficient to track the motion of all objects simultaneously (Norman

& Bobrow, 1975), necessitating a strategic allocation of cognitive resources to focus on

information most relevant to behavior. For example, spatial attention enhances processing at

specific locations in anticipation of a task-relevant stimulus (Posner, 1980; for review, see

Reynolds & Chelazzi, 2004; Summerfield & Egner, 2009). This prompts the intriguing question:

how does the allocation of attention influence the ability to predict motion, particularly in the FLE?

Several studies have demonstrated that variations in attentional resources can modulate the FLE.

For instance, Sarich et al. (2007) employed a dual-task paradigm to compare the magnitude of the

FLE in a flash-lag task performed alone or concurrently with a target detection task. They found

that the FLE, measured as a point of subjective equality, was smallest when the flash-lag task was

performed separately or with a slight interval from the detection task. Notably, when another task

necessitated attention, the magnitude of the FLE increased (and detection worsened). Similarly,

Shioiri et al. (2010) manipulated attention by varying the number of stimuli moving and presenting

the flash next to one of these stimuli. They observed an FLE increase when attention was divided

among two or six stimuli but an FLE decrease when focused solely on one stimulus. Furthermore,

when participants were pre-cued to upcoming flash locations, the FLE reduced following valid

cues compared to invalid cues (Baldo & Klein, 2010; Namba & Baldo, 2004; Vreven & Verghese,

2005; but see Khurana et al., 2000) or no cues (Shioiri et al., 2010). However, the specific

implications of these findings for the role of attention in motion extrapolation supposed to underlie

the FLE remain unclear.

While attentional mechanisms prioritize and allocate resources to relevant features or stimuli, they

could play a role in facilitating or modulating the extrapolation process. Extrapolation may become

prioritized in the context of divided attention, where attentional resources are distributed across

multiple objects. When detailed processing resources are limited, the brain may rely more heavily

on predictive mechanisms to maintain coherence in perception. In this sense, the brain might

compensate for the reduced attentional focus by generating predictions of each object's future

position based on generalized motion information. This could lead to a trade-off in the accuracy

and precision of predictions, potentially increasing the magnitude and variability of the FLE.

Furthermore, divided attention could lead to delays in processing motion information. Studies have

demonstrated that when attention is divided among multiple tasks or stimuli, processing efficiency

is compromised, leading to slower reaction times and delays in accessing relevant information

(Carrasco & McElree, 2001; Giordano et al., 2009). This time-consuming process could be

prolonged (cf. temporal integration; Lappe & Krekelberg, 1998), resulting in slower recognition

of motion and delayed availability of position information for each object. The brain may attempt

to extrapolate over larger distances to compensate for these delays, potentially increasing the FLE

magnitude. These possibilities suggest that the brain may prioritize efficiency over reliability of

motion processing.

Alternatively, divided attention may disrupt or attenuate the extrapolation process altogether.

When attention is divided, cognitive resources may be insufficient to accurately anticipate multiple

objects' trajectories simultaneously. This disruption could be limited by the quality of motion

information, such as spatial resolution (Anton-Erxleben & Carrasco, 2013; Barbot & Carrasco,

2017; Yeshurun & Carrasco, 1998, 1999). The brain may fail to generate reliable predictions about

each object's future positions, leading to errors or inconsistencies in perception. The perceived

position of the flash relative to the moving object may vary widely from trial to trial, potentially

increasing FLE variability.

The present study explored the effect of divided attention on the FLE using trial-wise spatial cues

to disentangle these possibilities. Previous studies have often used separate stimuli grouped as a

single moving object (dots arranged in a line; Baldo & Klein, 1995; Krekelberg & Lappe, 1999)

or multiple objects not grouped but having the same motion (dots revolving in a circular path;

Khurana et al., 2000; Shioiri et al., 2010; for an extensive review, see Hubbard, 2014), also

potentially causing grouping effects. In contrast, we employed several objects characterized by

independent motion trajectories. In this novel paradigm, an array of identically looking bars was

presented in different locations, and a cue preceding each trial indicated the bar to be tracked. All

bars rotated, but participants were required to covertly track the moving bar in the cued quadrant(s)

of the display. We varied the amount of attention available to the bar by dividing the participants'

attention between other bars in one, two, three, or across four quadrants, while a target flashed

next to one of the bars simultaneously. Subsequently, participants were asked to indicate the

position of that bar at the time of the flash. Assuming the FLE would be present in all attention

conditions, we hypothesized that the effects of divided attention would be expressed in three

scenarios as increased FLE magnitude and variability ("prediction model"), increased FLE

magnitude without changing the variability ("processing speed model"), or increased FLE

variability without changing the magnitude ("processing quality model").

2. Materials and Methods

2.1 Participants

Twenty-six healthy participants were recruited from the Forschungszentrum Jülich. Two participants were excluded due to insufficient data quality (see **section 2.5** for details), so data from 24 participants (15 female, 9 male, age [M: 28.8 years, SD: 4 years], range 21–40 years) were included in the final analyses. This sample size was motivated by a preliminary power calculation for a desired medium effect size (Cohen's $f = 0.25$) with a power of 80% and an alpha level of 0.05 for a repeated-measures analysis of variance (ANOVA). All participants were fluent in spoken and written English (in order to understand task instructions), had normal or corrected-to-normal visual acuity, and had no neurological or psychiatric disorders. Handedness was not a selection criterion, and participants self-reported their handedness preferences (19 right-handed and 5 left-handed). All participants provided informed written consent and received compensation of 10 Euros per hour for their participation. This study was conducted following the Declaration of Helsinki and approved by the ethics committee of the German Psychological Society (DGPs) (Ethics ID: 2022-02-03VA).

2.2 Apparatus

Visual stimuli were generated using Psychopy 3.0 (Peirce, 2007, 2009) and presented on a 22-inch Samsung SyncMaster 2233RZ LCD monitor with a 1680 x 1050 pixel resolution and a refresh rate of 120 Hz (as outlined in Wang & Nikolic, 2011). A chinrest was used to stabilize the head and maintain a viewing distance of 70 cm. The center of the screen was approximately at eye level. A standard QWERTY keyboard was positioned below the chinrest and not visible from the participants' field of view, with the left hand on the space bar and the right hand on the arrow keys. The experiment was conducted in a soundproof, light-attenuated room.

2.3 Stimuli

A white fixation cross subtending 0.5° x 0.5° of visual angle (dva) was continuously displayed at

the center of a black background. The spatial cue consisted of one, two, three, or four yellow dots

(radius = 0.15 dva) presented at the corners of the central fixation cross. The dots indicated the

quadrant(s) of the display in which the flash could occur. Every possible combination of dots was

realized (e.g., for two quadrants, two dots on the left, right, upper, lower, and diagonal quadrants).

The moving stimuli consisted of four gray bars, which rotated smoothly around the center of each

quadrant at an angular velocity of 240°/s. Therefore, the placement of each bar was 0.13 x 6.04

dva, with its inner end at 9.98 dva eccentricity from fixation. On one end of each bar was a small

dot (radius = 0.13 dva) used to indicate the bar's position, namely, relative to the position of the

static stimulus. The static stimulus (target) was a red dot of the same size as the dot end (radius =

0.13 dva) and was always positioned at 0.8 dva from the bar (dot end).

2.4 Procedure

In each trial, participants were instructed to fixate on the central fixation cross for the duration of

the stimulation sequence (**Figure 1A**). Attention cues were displayed for 600 ms, pointing to the

quadrant(s) to be attended. The cues signaled the quadrant but not the position within the quadrant,

in which the target likely appeared in the upcoming display (see below). After a 500-ms interval

allowing participants to covertly orient their attention to the cued quadrant(s), an array of four

rotating bars was presented surrounding the central fixation cross. The bars rotated in the same

clockwise or counter-clockwise direction, randomly interwoven across trials, with the restriction

that each bar had a distinct starting orientation: 0°, 22.5°, 45°, 67.5°, 90°, 112.5°, 135°, 157.5°,

180°, 202.5°, 225°, 247.5°, 270°, 292.5°, 315°, or 337.5° from the vertical. This arrangement

allowed the bars to rotate along independent motion trajectories despite their shared direction of

motion. Periodically after 850, 1000, or 1150 ms chosen at random, the target briefly flashed (25

ms) next to one of the bars. After the target offset, the bars continued to rotate for an additional

1150, 1000, or 850 ms. The participants' task was to make a delayed response about the

instantaneous position of the cued bar at the time of the flash, using a response probe.

Figure 1. Experimental paradigm. (**A**) Trial sequence. Spatial cues were presented at the central fixation cross to indicate that the participants should orient their attention to the upcoming display, featuring an array of four rotating bars (not drawn to scale) in corresponding quadrants. The target flashed next to a bar at a randomly selected moment during the sequence. Here, a target is presented at a cued quadrant, but sometimes the target appeared at a non-cued quadrant (see text for details). At the end of the sequence, participants were asked to respond using a response probe. Once the participants responded, the subsequent trial began after an inter-trial interval of 500 ms. Each trial lasted up to 8 s. (**B**) Procedure timeline. Participants performed at least 30 practice trials for task familiarization, followed by four blocks of 105 trials, totaling 420 trials over approximately one hour, including eye-tracking. Each block comprised trials from all four attention conditions and every combination of cue-target configurations with equal probability. The order of trials was randomized within the block.

The response probe comprised a wheel of 60 small grey dots (each dot radius = 0.13 dva) – whose

diameter was equal to the length of a bar. To perform the task, participants navigated the wheel

using left and right arrow keys to select their perceived position of the bar (dot end). Once satisfied

with their selection, they pressed the enter key to record the response and prompt the subsequent

trial. Because the starting position on the wheel randomly varied from trial to trial, participants

were instructed to respond as accurately as possible within 5000 ms. Because the target position

and the corresponding bar position were always physically aligned, the participants' perceived

displacement of the stimuli in the direction of motion, or lack thereof, allowed us to determine the

FLE of that trial directly. This approach contrasts with the conventional method of deriving an

average offset through the point of subjective equality across a block for a given condition (cf.

Yook et al., 2022). The subsequent trial began after an inter-trial interval (ITI) of 500 ms.

The probe location corresponded to the quadrant in which the target was presented in the previous

display, so participants did not additionally report in which quadrant they saw the target in each

trial. However, the cue-target validity differed between trials. Notably, the target was presented in

a cued quadrant 90% of the time. The target was presented in a non-cued quadrant on the remaining

10% of the time, and participants were asked to press the space bar and refrain from indicating the

bar's position with the response probe. We refer to these trials as "catch trials", which served to

ensure that participants would pay attention to the cues. If participants incorrectly responded to

catch trials by responding using the probe or to a non-catch trial by pressing space, they were given

feedback regarding their focus of attention: "Please pay attention to the cues as they are helpful".

Otherwise, no feedback on behavioral performance was provided.

The task was divided into four blocks of 105 trials (96 non-catch trials and 9 catch trials; **Figure**

1B). This yielded a total of 420 trials over approximately one hour, with each trial lasting up to 8

s. Every combination of cue-target configurations could appear within each block with equal

probability. Participants performed four attention conditions with the order of trials randomized

within each block. Participants were instructed to take regular self-paced breaks between each of

the four blocks and after every ~5–10 mins within each block (indicated by a break screen) to

minimize fatigue.

At the beginning of the experiment, participants completed at least one block of 30 practice trials

to familiarize themselves with the task. At the end of the experiment, participants were debriefed

about the purpose of the study involving the FLE. None of the participants indicated that they were

aware that the positions of the to-be-attended bar and target were aligned.

2.5 Data quality assessment

Data from participants who failed to identify at least 75% of catch trials were excluded from the

analyses to minimize the potential impact of non-compliance with task instructions and task

artifacts. This decision ensured that participants attended to the cued bar(s), as instructed. Data

from two participants who detected an average of 2.5 (SD: 3.53) out of 36 catch trials throughout

the experiment were excluded. Besides the catch trials, incorrectly responded non-catch trials were

also excluded from further analyses. Among the remaining 24 participants who detected an average

of 33.29 catch trials (SD: 2.35), an average of 379 correct non-catch trials (min: 371, max: 384,

SD: 3.75) were available for analyses.

2.6 Behavioral analysis

Behavioral data were pre-processed in Anaconda 4.2.0 (Anaconda Inc., 2016) with Python 3.5

(Van Rossum & Drake, 2009) and analyzed in R Statistical Software v4.3.1 (R Core Team, 2023)

with custom scripts.

Across trials, participants were required to focus on a single moving bar (focused attention

condition) or simultaneously divide their attention among two, three, or four bars (divided attention

conditions: attend-to-two, -three, or -four conditions). We obtained the FLE of each trial by

estimating the difference between the target position and the reported bar position. Given that a

value of zero indicated no difference in the positions of the two stimuli, a positive value indicated

a flash-lag effect (i.e., moving bar perceived ahead of the flash). Conversely, a negative value

indicated a flash-lead effect (i.e., flash perceived ahead of the moving bar).

To assess whether the magnitude of the FLE was influenced when attention was divided, we
calculated the median FLE values for each condition of each participant. We then compared the
means of medians (assessed per individual participant) between conditions using a one-way
repeated-measures ANOVA using the ez package (Lawrence, 2016) in R. The pairwise differences
between two conditions were evaluated with paired-samples t-tests. To correct for multiple
comparisons, statistically significant differences are reported based on a threshold of $p = (0.05/n -$
rank $+ 1$) (one-tailed), adjusted for familywise error rate (FWE) using Holm-Bonferroni correction
(Holm, 1979). Greenhouse-Geisser corrections were performed when the results of Mauchly's
sphericity test results were significant. The same procedure was repeated using the standard
deviation to assess whether the variability of the FLE across trials was influenced by divided
attention.

2.7 Eye-tracking recording and analysis

We monitored eye movements using an infrared EyeLink 1000 Plus (SR Research Ltd.,
Mississauga, Canada) system at a sampling rate of 1000 Hz to verify that participants maintained
fixation during the stimulation sequence of each trial. The eye tracker was calibrated with a 5-
point calibration procedure to establish an accurate gaze position at the beginning of the
experiment. Acceptable calibration values had to meet the validation criterion of < 0.5 dva and
maximum error of < 1.5 dva. The left eye's gaze positions were successfully recorded in 12 (of 24)
participants due to technical problems.

Eye-tracking data were pre-processed and analyzed in R using the eyelinker package (Barthelme,
2021) and custom scripts. Fixations and saccades were determined from the raw gaze position data
using Eyelink's default event parser. A fixation was defined as an event lasting at least 100 ms
within a 1-dva threshold from central fixation. Saccades were identified using a velocity threshold
of 30°/s and acceleration threshold of 8000°/s^2. We applied a drift correction by subtracting the
mean gaze position measured during the 500-ms ITI before the cue onset of each trial from all
gaze positions of that trial to account for potential gaze position drifts. This approach ensured
sensitivity to changes in head/eye position. Eye-blink events were excluded.

For each saccade, we examined the distance from a pre-determined region considered as
acceptable fixation within a 2-dva threshold at any point while the stimulus was on screen (i.e.
closer to stimulus position than central fixation). This analysis was conducted during three distinct
time intervals of interest: during the cue presentation lasting 600 ms, immediately after the cue
offset for a duration of 500 ms, and before and after the target presentation, with a combined
duration of 2000 ms. During these time windows, participants were instructed to maintain fixation.
By contrast, participants were allowed to make eye movements freely during the response interval.
Therefore, we assessed whether participants' perceptions of the bar position were systematically
influenced by voluntary eye movements, if any, during the critical time windows. The mean
deviations from fixation were compared between conditions using a two-way repeated-measures
ANOVA.

3. Results

3.1 Fixation controlled under all attention conditions

We compared gaze positions between conditions to ensure that overt attentional shifts did not

influence the participants' report of the bar position. Across the three windows of interest (during

the cue, after the cue, and before and after the target onset of each trial), there was no detectable

influence of our attention manipulations in deviations from fixation (two-way ANOVA,

*attention*timeWindow*: $F(6, 66) = 0.48$, $p = 0.82$; *attention*: $F(3, 33) = 0.38$, $p = 0.8$; *timeWindow*:

$F(2, 22) = 36.33$, $p < 0.001$).

3.2 Attentional modulation of FLE magnitude, but not FLE variability

Figure 2A shows the FLE magnitude, expressed as the average of individual medians, in relation

to the number of to-be-attended moving bars. The illusion manifested prominently across all

attention conditions, ranging between 15-20° (see **Appendix A**). Notably, the mean FLE

magnitude was significantly larger in divided attention conditions than in the condition where

attention was focused on a single bar (one-way ANOVA, *attention*: $F(3, 69) = 9.21$, $p < 0.001$).

Figure 2. FLE results. (**A**) The mean FLE magnitude is plotted as a function of number of attended

moving bars. All values are reported as means ± within-subject standard error of the mean (S.E.M.,

error bars). ns (not significant) denotes $p > 0.05$, ** denotes $p < 0.01$, *** denotes $p < 0.001$. (**B**)

The same described in **A** but for FLE variability.

A qualitatively similar pattern was observed for the FLE variability shown in **Figure 2B** (see also

Appendix B) but with no significant effect of attention (one-way ANOVA, *attention*: $F(3, 69) =$

$1.059, p = 0.3$) (**Table 1**).

Table 1.

Results of paired-sample *t*-tests (one-tailed) with correction for multiple comparisons for mean

FLE magnitude and variability.

Pair	FLE Magnitude		FLE Variability	
	t_{23}	p	t_{23}	p
1 vs 2	-3.86	< 0.01	-1.44	0.825
1 vs 3	-4.31	0.001	-1.57	0.774
1 vs 4	-5.19	< 0.001	-1.15	1
2 vs 3	-0.38	1	-0.43	1
2 vs 4	-0.27	1	-0.35	1
3 vs 4	0.26	1	-0.01	1

4. Discussion

In this study, we measured the perceived position of the moving object during the FLE across different attention conditions. When attention was distributed across multiple quadrants of the display, we found that FLE magnitude was more significant than when attention was directed to a single quadrant, which is in line with the prediction of the processing speed model. While the prediction model and the processing quality model predicted changes in variability, the observed variability of the FLE did not differ across conditions.

In each trial, participants viewed an array of four independently moving objects in corresponding quadrants of the display. Using spatial cues, we manipulated the allocation of attention to a single object, comparing the FLE under conditions where attention was either focused or divided. This approach was accompanied by catch trials, ensuring that participants accurately directed their attention according to the provided cues. The FLE was stronger when attention was divided among multiple objects, which is consistent with previous studies (Namba & Baldo, 2004; Sarich et al., 2007; Shioiri et al., 2010; but see Khurana et al., 2000), with our study benefiting from a larger sample size (N = 24) and a trial-wise measure of the FLE.

Subtle differences emerged within conditions of divided attention – specifically, when attention was distributed among two, three, or four moving objects. While there was a notable increase in FLE magnitude when attention shifted from one to two objects, subsequent increases were qualitatively smaller and did not reach statistical significance when attention was divided among three or four objects. These findings align with Shioiri et al. (2010), who observed a similar pattern in cued conditions where up to six dots formed a circle rather than being neatly arranged in single quadrants. Interestingly, Hogendoorn et al. (2010) presented a comparable set-up where attention was divided among six clocks in a circle; however, they observed an opposite pattern, with a gradual increase in magnitude (measured as time) from one to six clocks and a step-wise increase in variability from one to two clocks, but not beyond.

Our attentional manipulation may have peaked in the attend-to-two condition, and further dividing attention was less impactful in the FLE, indicating a saturation effect under highly divided-

attention conditions (Howe & Holcombe, 2012; Luu & Howe, 2015). According to Shioiri et al.,

this saturation effect could be related to the spatial spread of attention, independent of the number

of attended objects. For example, Khurana et al.'s (2000) study involved two moving objects above

and below the fixation point. Therefore, their cues indicated one of two possible locations where

the flash could appear, resulting in no difference in FLE magnitude (although in response time;

see **Appendix D**). Given that our study was similarly limited to four moving objects in four

quadrants, this pattern of results is coherent. Under these conditions, attention effects may not

necessarily follow a linear pattern but may critically depend on task demands (Khurana et al., 2000;

Shioiri et al., 2010). However, there is some evidence that attentional resources may be hemifield-

limited (Alvarez & Cavanagh, 2005; Luck et al., 1989) or even quadrant-limited (Carlson et al.,

2007, 2011). If each quadrant has discrete attentional capacity, then dividing attention would not

impact tracking additional moving objects in extra quadrants. This explanation could account for

the deviation observed in our supplementary analysis exploring FLE anisotropy (see **Appendix C**)

and the discrepancy in Khurana et al. (2000), showing no attention modulation compared to other

studies.

A promising avenue for future research involves increasing the demand for top-down attention to

address the observed saturation effect. It is possible that the current experiment set-up was too easy.

This draw-back could be addressed by manipulating attentional load, such as varying the number

of stimuli (> 1) in each quadrant across the four quadrants or varying eccentricities from each other

(Baldo et al., 2002). Additionally, motion speed has been reported to influence the attention

modulation in the FLE (Shioiri et al., 2010). Selectively manipulating the speed of moving stimuli

within this paradigm could further reveal whether more attention is required to track faster motion,

potentially influencing the FLE to a greater extent than observed here.

The results of the current study show a similar pattern as those reported by Namba and Baldo

(2004), Sarich et al. (2007), and Shioiri et al. (2010). However, since these studies did not employ

catch trials as done here, their results could be explained by anticipatory processing of the flash

(Baldo & Klein, 1995; Khurana et al., 2000; Namba & Baldo, 2004; Shioiri et al., 2010). According

to this interpretation, attention facilitates the processing of the flash (Baldo et al., 2002; Vreven &

Verghese, 2005), reducing the latency difference between the static flash and the moving object

under focused attention conditions (Shioiri et al., 2010). Conversely, under divided attention

conditions, the latency difference for each object may increase, potentially leading to a larger

perceived offset between the two (Öğmen et al., 2004; Purushothaman et al., 1998; Whitney &

Cavanagh, 2000; Whitney & Murakami, 1998). Although it is difficult to definitively rule out

attention's influence on the flash, it seems implausible that the flash's impact can solely explain

our results. While this interpretation would also hypothesize larger FLE magnitudes when attention

is divided among multiple objects, it implies that participants might anticipate the occurrence of

the flash even without explicit cues. The inclusion of catch trials in our analysis reduces the

likelihood that participants were just waiting to detect the flash. Instead, the catch trials added a

control measure to validate the effectiveness of the cues. If participants were waiting to detect the

flash, then they would be expected to report the FLE on these trials, rendering the cues ineffective.

However, our data indicate that participants did notice when the flash was presented next to a non-

cued bar, suggesting that they did not merely wait for the flash on each trial.

Besides, as our analysis focused on non-catch trials, another valuable avenue for future research

could involve measuring the FLE on catch trials. Increasing the frequency of catch trials as a

control condition would be advantageous for a more comprehensive investigation. A lack of

discernable difference in FLE magnitudes between catch and non-catch trials would strongly

indicate that observed FLE changes were solely driven by the flash. This would provide further

insights into how attention influences the perception of motion rather than the detection of the

flash. Additionally, our investigation of the FLE led us to consider another motion-induced position

shift known as the representational momentum (RM; Freyd & Finke, 1984; Hubbard, 1995).

Unlike the FLE, RM illustrates a perceived forward shift of a moving object's stopping position,

reflecting prediction about its motion trajectory without comparing it with another stimulus.

However, like the FLE, RM has been reported to observe similar modulations due to attention

(Hayes & Freyd, 2002; Hubbard et al., 2009; for reviews, see Hubbard, 2005, 2013, 2018),

suggesting that a common extrapolation mechanism might underlie the attentional modulation in

these illusions (but see White, 2018).

Our current results and previous findings consistently demonstrate larger FLE magnitudes under

divided attention conditions and smaller FLE magnitudes with focused attention. If we interpret

the FLE as a consequence of motion extrapolation, this pattern could be explained by two potential

mechanisms: attention modulates the impact of neural processing latencies or influences reliance

on predictive processing. Here, we speculated that divided attention may give rise to existing

predictions rather than wait for slow sensory input. However, the data do not seem to support the

possibility that divided attention enhances prediction itself since FLE variability did not modulate

with the presence of multiple simultaneously moving objects. Instead, our results are more in

accord with the view of an attentional effect on processing latencies, thereby modulating the degree

of compensatory motion extrapolation.

Conclusions

In conclusion, the experiment reported here is the first to manipulate attention in the flash-lag

paradigm using a combination of conventional attentional cueing and divided attention procedures

with trial-wise FLE readouts. Our results contribute to the growing body of evidence

demonstrating that for the brain to partly overcome its delays and allow us to interact with dynamic

stimuli without significant delays, it must always predict the future. More specifically, our results

reveal a dependency of these compensatory mechanisms on attention in dynamic environments.

Attention is crucial in efficiently processing multiple moving objects and facilitates motion

extrapolation.

Acknowledgements

We thank our colleagues of the INM-3 for their valuable discussions and support.

Commercial relationships: none

Corresponding author: Simone Vossel

Email: s.vossel@fz-juelich.de

Address: Cognitive Neuroscience, Institute of Neuroscience and Medicine (INM-3),

Forschungszentrum Jülich, 52428 Jülich, Germany

Appendices for

When visual attention is divided in the flash-lag effect

Jane Yook *et al.*

Appendix A: Above-zero flash-lag effect (FLE) magnitude

Table A1.

Results of one-sample *t*-tests (one-tailed) for mean FLE magnitude.

Bars	*M*	*SD*	t_{23}	*p*
1	15.82°	6.4°	12.11	< 0.001
2	18.99°	9.01°	10.33	< 0.001
3	19.36°	6.5°	14.61	< 0.001
4	19.18°	7.22°	13.01	< 0.001

488   **Appendix B: Descriptive statistics for FLE variability**

490   **Table B1.**

491   Descriptive statistics for mean FLE variability across attention conditions.

Bars	*M*	*SD*
1	23.22°	14.65°
2	24.44°	14.72°
3	24.79°	12.78°
4	24.81°	14.96°

Appendix C: Hemifield effects in the FLE

A

  B

Figure C1. Hemifield results. We investigated the attentional modulation of the FLE, focusing on where attention was allocated. Using linear motion, previous studies have demonstrated an FLE asymmetry in the left and right sides of the visual field (Kanai et al., 2004; Shi & Nijhawan, 2008; Suzuki et al., 2023). Notably, the FLE tends to be more substantial when stimuli are presented in the left hemifield. Therefore, we aimed to test whether the effects of divided attention were more pronounced with FLE anisotropy, particularly within the attend-to-two condition. To this end, we examined the magnitude and variability of the FLE when participants were cued to two quadrants within their left hemifield, right hemifield, or diagonally across the upper and lower visual hemifields. (**A**) The mean FLE magnitude is plotted as a function of the attended hemifields. All values are reported as means ± within-subject standard error of the mean (S.E.M., error bars). ns (not significant) denotes $p > 0.05$. Although the FLE magnitude in the left hemifield was marginally higher than in the right or diagonal hemifields (**Table S3**), no significant modulation of attention was detected (one-way ANOVA, *hemiField: $F(2,46) = 0.16$, $p = 0.741$*). (**B**) The same as described in **A** but for mean FLE variability based on the oriented attention across hemifields (one-way ANOVA, *hemiField: $F(2,46) = 0.32$, $p = 0.73$*).

Table C2.

Results of the paired-sample *t*-tests (two-tailed) with Bonferroni correction for multiple comparisons for mean FLE magnitude and variability.

Pair	FLE Magnitude		FLE Variability	
	t_{23}	p	t_{23}	p
left vs right	0.19	1	0.67	1
diagonal vs left	-0.79	1	-0.11	1
diagonal vs right	-0.38	1	0.86	1

Appendix D: Response times

While response time (RT) is commonly utilized as an indicator of faster processing in the literature (see also Khurana et al., 2000), our study primarily focused on assessing the magnitude and variability of the FLE. There were two key reasons for not analyzing RTs in this paradigm. Firstly, our experimental task deliberately emphasized response accuracy (i.e., the bar position) over speed in determining the FLE. Unlike the emphasis on speed in previous studies such as Khurana et al. (2000), our goal was to ensure accuracy in the FLE task. Secondly, the starting position on the response probe introduced inherent variability. Longer response times were needed when the distance from the starting position to the desired position was greater. The focus of our study was the magnitude and variability of the FLE, and the RT analysis was not aligned with our primary research questions.

The RTs from the onset of the response probe averaged approximately ~3400 ms across all conditions (**Table D1**).

Table D1.

Descriptive statistics for mean RTs across attention conditions.

Bars	*M*	*SD*
1	3438 ms	133 ms
2	3450 ms	135 ms
3	3435 ms	135 ms
4	3439 ms	138 ms

540   **References**

542   Adamian, N., & Andersen, S. K. (2022). Attentional Enhancement of Tracked Stimuli in Early

543   Visual Cortex Has Limited Capacity. *The Journal of Neuroscience*, *42*(46), 8709–8715.

544   https://doi.org/10.1523/JNEUROSCI.0605-22.2022

545   Alvarez, G. A., & Cavanagh, P. (2005). Independent Resources for Attentional Tracking in the

546   Left and Right Visual Hemifields. *Psychological Science*, *16*(8), 637–643.

547   https://doi.org/10.1111/j.1467-9280.2005.01587.x

548   Anaconda Inc. (2016). *Anaconda Software Distribution* (Version 2-2.4.0). Anaconda Inc. Austin,

549   TX, USA. https://www.anaconda.com/

550   Anton-Erxleben, K., & Carrasco, M. (2013). Attentional enhancement of spatial resolution:

551   Linking behavioural and neurophysiological evidence. *Nature Reviews. Neuroscience*,

552   *14*(3), 188–200. https://doi.org/10.1038/nrn3443

553   Baldo, M. V. C., Kihara, A. H., Namba, J., & Klein, S. A. (2002). Evidence for an Attentional

554   Component of the Perceptual Misalignment between Moving and Flashing Stimuli.

555   *Perception*, *31*(1), 17–30. https://doi.org/10.1068/p3302

556   Baldo, M. V. C., & Klein, S. A. (1995). Extrapolation or attention shift? *Nature*, *378*(6557),

557   Article 6557. https://doi.org/10.1038/378565a0

558   Baldo, M. V. C., & Klein, S. A. (2010). Paying attention to the flash-lag effect. In R. Nijhawan

559   & B. Khurana (Eds.), *Space and Time in Perception and Action* (1st ed., pp. 396–407).

560   Cambridge University Press. https://doi.org/10.1017/CBO9780511750540.023

561   Barbot, A., & Carrasco, M. (2017). Attention Modifies Spatial Resolution According to Task

562   Demands. *Psychological Science*, *28*(3), 285–296.

563   https://doi.org/10.1177/0956797616679634

564   Barthelme, S. (2021). *eyelinker: Import ASC Files from EyeLink Eye Trackers.* R package

565   version 0.2.1. https://cran.r-project.org/web/packages/eyelinker/index.html

566   Brenner, E., & Smeets, J. B. J. (2000). Motion extrapolation is not responsible for the flash–lag
567   effect. *Vision Research*, *40*(13), 1645–1648. https://doi.org/10.1016/S0042-
568   6989(00)00067-5

569   Carlson, T. A., Alvarez, G. A., & Cavanagh, P. (2007). Quadrantic deficit reveals anatomical
570   constraints on selection. *Proceedings of the National Academy of Sciences*, *104*(33),
571   13496–13500. https://doi.org/10.1073/pnas.0702685104

572   Carlson, T. A., Cho, H., Turret, J., & Dakin, S. (2011). Psychoanatomy of visual attention: A
573   unified account of quadrant and hemifield effects. *Journal of Vision*, *11*(11), 101–101.
574   https://doi.org/10.1167/11.11.101

575   Carrasco, M., & McElree, B. (2001). Covert attention accelerates the rate of visual information
576   processing. *Proceedings of the National Academy of Sciences*, *98*(9), 5363–5367.
577   https://doi.org/10.1073/pnas.081074098

578   Eagleman, D. M., & Sejnowski, T. J. (2000). Motion Integration and Postdiction in Visual
579   Awareness. *Science*, *287*(5460), 2036–2038.
580   https://doi.org/10.1126/science.287.5460.2036

581   Freyd, J. J., & Finke, R. A. (1984). Representational momentum. *Journal of Experimental
582   Psychology: Learning, Memory, and Cognition*, *10*(1), 126–132.
583   https://doi.org/10.1037/0278-7393.10.1.126

584   Giordano, A. M., McElree, B., & Carrasco, M. (2009). On the automaticity and flexibility of
585   covert attention: A speed-accuracy trade-off analysis. *Journal of Vision*, *9*(3), 30.
586   https://doi.org/10.1167/9.3.30

587   Hayes, A. E., & Freyd, J. J. (2002). Representational momentum when attention is divided.
588   *Visual Cognition*, *9*(1–2), 8–27. https://doi.org/10.1080/13506280143000296

589   Hogendoorn, H., Carlson, T. A., Vanrullen, R., & Verstraten, F. A. J. (2010). Timing divided
590   attention. *Attention, Perception, & Psychophysics*, *72*(8), 2059–2068.
591   https://doi.org/10.3758/BF03196683

Holm, S. (1979). A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian*
Journal of Statistics, *6*(2), 65–70.

Howe, P. D. L., & Holcombe, A. O. (2012). Motion information is sometimes used as an aid to
the visual tracking of objects. *Journal of Vision*, *12*(13), 10–10.
https://doi.org/10.1167/12.13.10

Hubbard, T. L. (1995). Environmental invariants in the representation of motion: Implied
dynamics and representational momentum, gravity, friction, and centripetal force.
Psychonomic Bulletin & Review, *2*(3), 322–338. https://doi.org/10.3758/BF03210971

Hubbard, T. L. (2005). Representational momentum and related displacements in spatial
memory: A review of the findings. *Psychonomic Bulletin & Review*, *12*(5), 822–851.
https://doi.org/10.3758/BF03196775

Hubbard, T. L. (2013). Do the flash-lag effect and representational momentum involve similar
extrapolations? *Frontiers in Psychology*, *4*. https://doi.org/10.3389/fpsyg.2013.00290

Hubbard, T. L. (2014). The flash-lag effect and related mislocalizations: Findings, properties,
and theories. *Psychological Bulletin*, *140*(1), 308–338. https://doi.org/10.1037/a0032899

Hubbard, T. L. (2018). Influences on representational momentum. In *Spatial biases in perception*
and cognition (pp. 121–138). Cambridge University Press.
https://doi.org/10.1017/9781316651247.009

Hubbard, T. L., Kumar, A. M., & Carp, C. L. (2009). Effects of spatial cueing on
representational momentum. *Journal of Experimental Psychology: Learning, Memory,*
and Cognition, *35*(3), 666–677. https://doi.org/10.1037/a0014870

Johnson, P. A., Blom, T., Van Gaal, S., Feuerriegel, D., Bode, S., & Hogendoorn, H. (2023).
Position representations of moving objects align with real-time position in the early
visual response. *eLife*, *12*, e82424. https://doi.org/10.7554/eLife.82424

Kanai, R., Sheth, B. R., & Shimojo, S. (2004). Stopping the motion and sleuthing the flash-lag
effect: Spatial uncertainty is the key to perceptual mislocalization. *Vision Research*,
44(22), 2605–2619. https://doi.org/10.1016/j.visres.2003.10.028

Khurana, B., Watanabe, K., & Nijhawan, R. (2000). The Role of Attention in Motion
Extrapolation: Are Moving Objects 'Corrected' or Flashed Objects Attentionally
Delayed? *Perception*, *29*(6), 675–692. https://doi.org/10.1068/p3066

Krekelberg, B., & Lappe, M. (1999). Temporal recruitment along the trajectory of moving
objects and the perception of position. *Vision Research*, *39*(16), 2669–2679.
https://doi.org/10.1016/S0042-6989(98)00287-9

Krekelberg, B., & Lappe, M. (2001). Neuronal latencies and the position of moving objects.
Trends in Neurosciences, *24*(6), 335–339. https://doi.org/10.1016/S0166-2236(00)01795-
1

Lappe, M., & Krekelberg, B. (1998). The Position of Moving Objects. *Perception*, *27*(12), 1437–
1449. https://doi.org/10.1068/p271437

Lawrence, M. (2016). *ez: Easy Analysis and Visualization of Factorial Experiments*. R package
version 4.4-0. https://CRAN.R-project.org/package=ez

Luck, S. J., Hillyard, S. A., Mangun, G. R., & Gazzaniga, M. S. (1989). Independent
hemispheric attentional systems mediate visual search in split-brain patients. *Nature*,
342(6249), 543–545. https://doi.org/10.1038/342543a0

Luu, T., & Howe, P. D. L. (2015). Extrapolation occurs in multiple object tracking when eye
movements are controlled. *Attention, Perception, & Psychophysics*, *77*(6), 1919–1929.
https://doi.org/10.3758/s13414-015-0891-8

Mackay, D. M. (1958). Perceptual Stability of a Stroboscopically Lit Visual Field containing
Self-Luminous Objects. *Nature*, *181*(4607), 507–508. https://doi.org/10.1038/181507a0

Murakami, I. (2001). The flash-lag effect as a spatiotemporal correlation structure. *Journal of
Vision*, *1*(2), 6. https://doi.org/10.1167/1.2.6

Namba, J., & Baldo, M. V. C. (2004). The Modulation of the Flash-Lag Effect by Voluntary
Attention. *Perception*, *33*(5), 621–631. https://doi.org/10.1068/p5212

Nijhawan, R. (2002). Neural delays, visual motion and the flash-lag effect. *Trends in Cognitive*
Sciences, *6*(9), 387–393. https://doi.org/10.1016/S1364-6613(02)01963-0

Nijhawan, R. (2008). Visual prediction: Psychophysics and neurophysiology of compensation
for time delays. *Behavioral and Brain Sciences*, *31*(2), 179–198.
https://doi.org/10.1017/S0140525X08003804

Norman, D. A., & Bobrow, D. G. (1975). On data-limited and resource-limited processes.
Cognitive Psychology, *7*(1), 44–64. https://doi.org/10.1016/0010-0285(75)90004-3

Öğmen, H., Patel, S. S., Bedell, H. E., & Camuz, K. (2004). Differential latencies and the
dynamics of the position computation process for moving targets, assessed with the flash-
lag effect. *Vision Research*, *44*(18), 2109–2128.
https://doi.org/10.1016/j.visres.2004.04.003

Peirce, J. W. (2007). PsychoPy—Psychophysics software in Python. *Journal of Neuroscience*
Methods, *162*(1), 8–13. https://doi.org/10.1016/j.jneumeth.2006.11.017

Peirce, J. W. (2009). Generating stimuli for neuroscience using PsychoPy. *Frontiers in*
Neuroinformatics, *2*. https://www.frontiersin.org/articles/10.3389/neuro.11.010.2008

Posner, M. I. (1980). Orienting of attention. *The Quarterly Journal of Experimental Psychology*,
32(1), 3–25. https://doi.org/10.1080/00335558008248231

Purushothaman, G., Patel, S. S., Bedell, H. E., & Ogmen, H. (1998). Moving ahead through
differential visual latency. *Nature*, *396*(6710), 424–424. https://doi.org/10.1038/24766

Pylyshyn, Z. W., & Storm, R. W. (1988). Tracking multiple independent targets: Evidence for a
parallel tracking mechanism. *Spatial Vision*, *3*(3), 179–197.
https://doi.org/10.1163/156856888x00122

R Core Team. (2023). *R: A Language and Environment for Statistical Computing.* Vienna,
667   Austra. R Foundation for Statistical Computing. https://www.r-project.org/

668   Reynolds, J. H., & Chelazzi, L. (2004). Attentional modulation of visual processing. *Annual*
669   *Review of Neuroscience*, *27*(1), 611–647.
670   https://doi.org/10.1146/annurev.neuro.26.041002.131039

671   Sarich, D., Chappell, M., & Burgess, C. (2007). Dividing attention in the flash-lag illusion.
672   *Vision Research*, *47*(4), 544–547. https://doi.org/10.1016/j.visres.2006.09.029

673   Schneider, K. A. (2018). The Flash-Lag, Fröhlich and Related Motion Illusions Are Natural
674   Consequences of Discrete Sampling in the Visual System. *Frontiers in Psychology*, *9*,
675   1227. https://doi.org/10.3389/fpsyg.2018.01227

676   Shi, Z., & Nijhawan, R. (2008). Behavioral significance of motion direction causes anisotropic
677   flash-lag, flash-drag, flash-repulsion, and movement-mislocalization effects. *Journal of*
678   *Vision*, *8*(7), 24. https://doi.org/10.1167/8.7.24

679   Shioiri, S., Yamamoto, K., Oshida, H., Matsubara, K., & Yaguchi, H. (2010). Measuring
680   attention using flash-lag effect. *Journal of Vision*, *10*(10), 10–10.
681   https://doi.org/10.1167/10.10.10

682   Summerfield, C., & Egner, T. (2009). Expectation (and attention) in visual cognition. *Trends in*
683   *Cognitive Sciences*, *13*(9), 403–409. https://doi.org/10.1016/j.tics.2009.06.003

684   Suzuki, Y., Atmaca, S., & Laeng, B. (2023). The lateralized flash-lag illusion: A psychophysical
685   and pupillometry study. *Brain and Cognition*, *166*, 105956.
686   https://doi.org/10.1016/j.bandc.2023.105956

687   Van Rossum, G., & Drake, F. L. (2009). *Python 3 Reference Manual*. Scotts Valley, CA:
688   CreateSpace.

689   Vreven, D., & Verghese, P. (2005). Predictability and the Dynamics of Position Processing in the
690   Flash-Lag Effect. *Perception*, *34*(1), 31–44. https://doi.org/10.1068/p5371

691   Wang, P., & Nikolic, D. (2011). An LCD Monitor with Sufficiently Precise Timing for Research
692   in Vision. *Frontiers in Human Neuroscience*, *5*.
693   https://www.frontiersin.org/articles/10.3389/fnhum.2011.00085

694   Whitney, D., & Cavanagh, P. (2000). Motion distorts visual space: Shifting the perceived
695   position of remote stationary objects. *Nature Neuroscience*, *3*(9), 954–959.
696   https://doi.org/10.1038/78878

697   Whitney, D., & Murakami, I. (1998). Latency difference, not spatial extrapolation. *Nature
698   Neuroscience*, *1*(8), 656–657. https://doi.org/10.1038/3659

699   Yeshurun, Y., & Carrasco, M. (1998). Attention improves or impairs visual performance by
700   enhancing spatial resolution. *Nature*, *396*(6706), Article 6706.
701   https://doi.org/10.1038/23936

702   Yeshurun, Y., & Carrasco, M. (1999). Spatial attention improves performance in spatial
703   resolution tasks1Parts of this study were presented at the Annual Meeting of the
704   Association for Research in Vision and Ophthalmology (May 1997) and at the Annual
705   Meeting of the Psychonomics Society (November 1997) and published in Abstract format
706   (Yeshurun and Carrasco, 1997and Carrasco and Yeshurun, 1997, respectively).1. *Vision
707   Research*, *39*(2), 293–306. https://doi.org/10.1016/S0042-6989(98)00114-X

708   Yook, J., Lee, L., Vossel, S., Weidner, R., & Hogendoorn, H. (2022). Motion extrapolation in the
709   flash-lag effect depends on perceived, rather than physical speed. *Vision Research*, *193*,
710   107978. https://doi.org/10.1016/j.visres.2021.107978

711